Identifying the drivers of computationally detected correlated evolution among sites under antibiotic selection

Jonathan Dench[1] ⋆, Aaron Hinz[1], Stéphane Aris-Brosou[1,2], and Rees Kassen[1]

[1]Department of Biology, University of Ottawa, Ottawa, ON K1N 6N5, Canada

[2]Department of Mathematics and Statistics, University of Ottawa, Ottawa, ON K1N 6N5, Canada

⋆Author for correspondence: Jonathan Dench (`Jdenc017@gmail.com`).

ABSTRACT

The ultimate causes of correlated evolution among sites in a genome remain difficult to tease apart. To address this problem directly, we performed a high-throughput search for correlated evolution among sites associated with resistance to a fluoroquinolone antibiotic using whole genome data from clinical strains of *Pseudomonas aeruginosa*, before validating our computational predictions experimentally. We show that for at least two sites, this correlation is underlain by epistasis. Our analysis also revealed eight additional pairs of synonymous substitutions displaying correlated evolution underlain by physical linkage, rather than selection associated with antibiotic resistance. Our results provide direct evidence that both epistasis and physical linkage among sites can drive the correlated evolution identified by high throughput computational tools. In other words, the observation of correlated evolution is not by itself sufficient evidence to guarantee that the sites in question are epistatic; such a claim requires additional evidence, ideally coming from direct estimates of epistasis, based on experimental evidence.

Keywords: Correlated evolution, epistasis, antibiotic resistance, *Pseudomonas aeruginosa*, computational methods, experimental validation

1 INTRODUCTION

Adaptive evolution often involves changes to multiple characters (or traits) in concert, a process called correlated evolution. Such coordinated changes arise either because of physical linkage in the genome, or from strong selection that generates an association between distinct genomic sites. The analysis of correlated evolution among traits has a long history in quantitative genetics (Falconer, 1960; Cheverud, 1984; Lynch and Walsh, 1998), molecular biology (Goh *et al.*, 2000; Callahan *et al.*, 2011), life history evolution (Baer and Lynch, 2003; Sexton *et al.*, 2009; Kelley *et al.*, 2013), and comparative biology (Pagel, 1994; Shimizu *et al.*, 2014). A comparable

effort at the genomic level remains a daunting task because the number of potentially interacting sites (*i.e.* nucleotides) can be overwhelmingly large, making it difficult to distinguish genuine instances of correlated evolution arising from linkage or selection from spurious correlations arising due to chance.

In an effort to fill this gap, we have extended a computational approach, implemented in a software called AEGIS (Analysis of Epistasis and Genomic Interacting Sites), designed to detect both positively and negatively correlated pairs of mutations with very high specificity (Nshogozabahizi *et al.*, 2017). Although the original approach focused on identifying correlated evolution among sites within genes (Ibeh *et al.*, 2016; Nshogozabahizi *et al.*, 2017; Aris-Brosou *et al.*, 2017), nothing prevents it from being used to perform the same task across entire genomes. AEGIS makes use of Pagel's phylogenetically informed maximum likelihood model for predicting correlated evolution between pairs of traits based on the co-distribution of their values (Pagel, 1994), and was extensively tested through both simulations and analyses of real data in the context of detecting correlated evolution at the molecular level, between pairs of nucleotide positions in a particular genome alignment (Nshogozabahizi *et al.*, 2017). The output of AEGIS is a list of pairs of sites and their associated probabilities of co-evolving by chance. While this approach uses phylogenetic information to reduce the detection of spurious associations, it is agnostic to the underlying mechanism responsible for correlation among sites, and thus cannot distinguish between linkage and selection leading to nonadditive fitness effects, or epistasis, as the cause of correlated evolution. This approach, like most statistical comparative methods, are now known to detect correlated evolution even in the case of a single unreplicated co-distribution of traits (Maddison and FitzJohn, 2014; Uyeda *et al.*, 2018).

Consequently, identifying the mechanisms driving correlated evolution still requires direct experimental measurements of epistasis between sites, preferably under conditions similar to the selective setting in which the pairs of sites evolved. To do this, we focused on the evolution

of resistance to fluoroquinolone antibiotics in the opportunistic pathogen *Pseudomonas aeruginosa*. This pathogen is a ubiquitous, Gram-negative bacterium that causes both acute opportunistic infections and chronic respiratory tract infections in cystic fibrosis patients. Treatment with fluoroquinolone antibiotics imposes strong selection that regularly leads to the evolution of resistance at a number of well-known target sites such as efflux pump regulators (e.g. *nfxB*) and DNA topoisomerases (i.e. *gyrA*, *parC*) (Akasaka *et al.*, 2001; Wong *et al.*, 2012; Melnyk *et al.*, 2015; Kos *et al.*, 2015). As *P. aeruginosa* is amenable to genetic manipulation, we can introduce putative correlated mutations, either one at a time or in combination, into a range of genetic backgrounds. The level of epistasis can then be measured by comparing the effects of the two substitutions against the expected combined effects of each single substitution on traits relevant to selection, such as antibiotic resistance or fitness in the absence of antibiotics.

Here, we use a unique data set of 393 *P. aeruginosa* exomes to predict correlated evolution among sites and evaluate the mechanistic causes of correlations that evolve during selection by fluoroquinolone antibiotics. We investigate the role of epistasis and linkage, both physical and through hitchhiking, as causes of correlated evolution using a combination of tools including site-directed mutagenesis to reconstruct single and double mutants. For nonsynonymous substitutions that co-evolved in response to antibiotic selection, we test for epistasis with direct estimates of the minimum inhibitory concentration (MIC) of antibiotics and competitive fitness in the absence of drugs. We employ additional computational analyses to infer the mechanism leading to the correlated evolution of synonymous substitutions. Our results demonstrate that epistasis can drive correlated evolution among nonsynonymous sites tied to antibiotic resistance, whereas correlated evolution among synonymous sites is best explained by idiosyncratic processes.

2 MATERIALS AND METHODS

2.1 Alignment and phylogeny

We assembled a multiple sequence alignment which included the complete genomes of *P. aeruginosa* strains PA01 (PA01), *P. aeruginosa* UCBPP PA14 (PA14), and *P. aeruginosa* PA7 (PA7) downloaded from www.pseudomonas.com (Winsor *et al.*, 2016, accessed Dec 4, 2014), as well as 390 draft *P. aeruginosa* genomes (Kos *et al.*, 2015), used here as they were published alongside information detailing which of the strains were resistant to the fluoroquinolone antibiotic levofloxacin. The genomes used in our study represent a collection of clinical strains isolated from around the world (Table S1). Due to dissimilarities among contigs across the draft genomes, we were unable to construct a global alignment using whole genome alignment algorithms as implemented in either MAUVE (Darling *et al.*, 2004) or MUGSY (Angiuoli and Salzberg, 2011). Using PA14 reference gene sequences detailed in a database downloaded from www.pseudomonas.com (Winsor *et al.*, 2016, accessed Dec 4, 2014), we assembled an exome alignment using a custom algorithm to concatenate individual gene alignments.

The species tree of these bacterial genomes was reconstructed based on 'highly networked' genes that, according to the complexity hypothesis, are unlikely to be horizontally transferred. We identified these highly networked genes from the Cluster of Orthologous Groups database's (Tatusov *et al.*, 1997; Galperin *et al.*, 2015) information class genes (those with COG terms A, B, J, K, and L). From an alignment of 1,290 highly networked *P. aeruginosa* genes, we estimated the species tree by maximum likelihood with FastTree (Price *et al.*, 2010) (GTR $+ \Gamma$ model of evolution (Aris-Brosou and Rodrigue, 2012)) compiled with *DUSE_DOUBLE* as recommended for large sequence alignments. This tree was rooted with the subclade containing the taxonomic outliers PA7 (Roy *et al.*, 2010), AZPAE14941, AZPAE14901 and AZPAE15042 (Kos *et al.*, 2015). After rooting the tree, we removed the subclade from both the phylogeny and the above

103 exome alignment, so that downstream analyses were based on 389 genomes. For additional
104 information please see *Alignment algorithm* and *Complexity Hypothesis* in the supplementary
105 information.

2.2 Analysis of correlated evolution

We used AEGIS (Nshogozabahizi *et al.*, 2017) to detect pairs of nucleotide positions (sites)
that evolved in a correlated manner. For this, AEGIS performed pairwise comparisons be-
tween sites, testing if a model of dependent evolution was significantly better than an inde-
pendent model at explaining the observed phylogenetic distribution of nucleotide states. This
maximum likelihood analysis relied on the algorithm (Pagel, 1994), computationally validated
with AEGIS for use with amino acid data (Nshogozabahizi *et al.*, 2017), as implemented in
BayesTraits for binary states (Pagel and Meade, 2006). Nucleotide states in our alignment were
converted into binary characters where '0' was assigned where there was a consensus of nu-
cleotide character at 80% of the levofloxacin sensitive strains and '1' otherwise. We chose this
method of recoding so that signals of correlated evolution would more likely reflect adapta-
tion to fluoroquinolone antibiotic selection. All R scripts and complete results can be found
at `github.com/JDench/sigResults_AEGIS_inVivo`. Due to computational limitations, we
limited our analysis of correlated evolution to a 12 focal gene by exome analysis. Six of the focal
genes were previously shown to evolve during fluoroquinolone selection (Akasaka *et al.*, 2001;
Wong *et al.*, 2012) (*gyrA*, *gyrB*, *nfxB*, *parC*, *parE*) or are linked to biofilm formation (Choy *et al.*,
2004), which can contribute to antibiotic resistance (Starkey *et al.*, 2009) (*morA*). The six other
genes (*dnaA*, *dnaN*, *lpd3*, *ribD*, *rpoB*, *serC*) were randomly selected from among the other 5,971
genes in our alignment. For additional information please see *Analysis of correlated evolution*
in the supplementary information.

2.3 Direct experimental assessment of epistasis

All growth and incubation conditions were performed in lysogeny broth (LB) (Bertani, 1951) containing half the normal salt (i.e., 5g/L NaCl), at 37°C, in an orbital shaker set at 150 rpm.

Generating mutants: To measure the *in vitro* fitness and antibiotic resistance effects of substitutions predicted with AEGIS, we created mutant constructs using a modified selection counter-selection allelic replacement protocol (Melnyk *et al.*, 2017). Here, replacement alleles were constructed from the wild-type (WT) template and contained only the mutations of interest rather than coming from an evolved strain which may carry non-focal mutations. As the effect of a mutation may be contingent upon the genetic background (i.e., WT) in which it arose (Sorrells *et al.*, 2015; Kryazhimskiy *et al.*, 2014; Blount *et al.*, 2008), constructs were created using two distinct genetic backgrounds: PA01, and PA14 (Dettman *et al.*, 2013). For additional information please see *Modified selection counter-selection protocol* in the supplementary information.

Minimum inhibitory concentration assays: We performed minimum inhibitory concentration assays to identify if our mutant constructs had increased resistance to the fluoroquinolone antibiotic ciprofloxacin. Using a 96-well plate, we carried out serial 2-fold dilutions of ciprofloxacin, in a liquid LB broth, such that columns represented a concentration gradient from 32 to 0.015625 (*i.e.* $2^{(5,4,\ldots,-6)}$) μg/mL. We inoculated each well with 5 μL of overnight mutant culture such that each row represented a single type of inoculum. After 24 hours of growth, we read the absorbance values (at 600 nm) using a plate reader (Table S7). Each plate contained at least one blank row, used as reference for reads of the absorbance. We performed four replicates of each genotype, while ensuring that no genotype appeared on a single plate more than once.

Competitive fitness assays: We measured relative fitness of WT and mutant constructs via competitive fitness assays. Competitions began when we mixed an equal volume of overnight culture, diluted 100 fold, of a focal strain and a marked competitor (Lenski *et al.*, 1991). Similar to previous work, the marked competitor was a WT strain bearing the neutral *lacZ* marker

gene. The frequency of focal and marked strains were estimated using direct counts of diluted culture plated immediately after mixing and again after competition during overnight growth to stationary phase. We plated six replicates of each competition, and time point, on minimal media agar plates containing X-Gal. Counting was done after overnight growth at 37°C, followed by 2-4 days of growth at room temperature. For additional information please see *Calculating competitive fitness* in the supplementary information.

2.4 Quantifying importance of sites in predicting resistance

We quantified the importance of polymorphic sites in our alignment for predicting resistance to levofloxacin based on the machine learning classification algorithm 'adaptive boosting', implemented in the *adaBoost* (Alfaro *et al.*, 2013) R package. Measurements of importance provided a relative measure of the strength of association between resistance and substitution at a site in our alignment. We compared the machine learning results to substitutions known to be within the six, previously identified, focal *P. aeruginosa* genes of which five evolve in response to fluoroquinolone selection (Akasaka *et al.*, 2001; Wong *et al.*, 2012; Kos *et al.*, 2015) and one is involved in biofilm formation (Choy *et al.*, 2004) which can contribute to antibiotic resistance (Starkey *et al.*, 2009). For additional information please see *adaptive boosting algorithm* in the supplementary information.

2.5 Assessing horizontal gene transfer

Horizontal gene transfer (HGT) was assessed with a phylogenetic approach (Ravenhall *et al.*, 2015), where we reconstructed each gene tree as above (maximum likelihood under GTR + Γ), and tested for tree concordance with the Approximately Unbiased (AU) test (Shimodaira, 2002). This was done with CONSEL (Shimodaira and Hasegawa, 2001), on the *baseml* (GTR + Γ_5, no

173 clock) output from PAML (Yang, 2007).

174 2.6 Testing for biological effects of synonymous substitutions

To test if pairs of synonymous substitutions were correlated due to translational effects, we es-
timated the free energy (ΔG) of folded mRNA transcripts for both WT and mutant transcripts,
which we normalized as relative values through division of the WT estimate (ΔG_{rel}). We also es-
timated if synonymous mutations affected translation elongation rates via estimates of the index
of translation elongation (I_{TE}: Xia, 2014). For additional information please see *Calculating*
ΔG and I_{TE} in the supplementary information.

181 3 RESULTS AND DISCUSSION

182 3.1 Predicting correlated evolution among sites

We identified pairs of sites that evolve in a correlated manner in the context of fluoroquinolone
resistance by first assembling a multiple sequence alignment using the entire set of 5,977 protein-
coding genes from 393 previously sequenced *P. aeruginosa* strains. This multiple sequence
alignment included three reference strains (PA01, PA14, and PA7) and 390 clinical strains for
which we had information on the level of levofloxacin (a fluoroquinolone antibiotic) resistance
(Kos *et al.*, 2015). We then used this alignment to construct a phylogenetic tree rooted by PA7,
an outlier strain (Roy *et al.*, 2010) (Fig. S2). The subclade containing this strain was removed for
downstream analyses. We acknowledge that the comparative approach implemented in AEGIS
may not be ideally suited to analyze highly promiscuous bacteria; however, we constructed our
phylogeny from genes unlikely to undergo horizontal transfer so that signals of correlated evo-
lution would represent independent evolutionary events.

We subsequently employed AEGIS to identify pairs of nucleotide positions that show ev-

idence of correlated evolution. As this approach can be computationally expensive, requiring

on the order of n^2 comparisons in a genome of length n ($\sim 84 \times 10^9$ comparisons), we focused

our analyses on a subset of twelve genes against all other positions in the exome. AEGIS calcu-

lated the probability that each position within these twelve genes evolved in a correlated manner

with any other nucleotide position in the entire *P. aeruginosa* exome. Among the more than

144 million pairwise comparisons, we found that $\sim 127,000$ pairs of sites showed evidence of

significant correlation with $P \leq 0.01$ after a Benjamini-Hochberg correction for false discovery

rate (FDR). At medium ($P \leq 10^{-7}$) or high ($P \leq 10^{-11}$) significance levels, as previously deter-

mined based on extensive simulations (Nshogozabahizi *et al.*, 2017), only 178 and nine pairs,

respectively, showed evidence for correlated evolution (Fig. 1). Among the nine pairs of sites

showing the strongest evidence for correlated evolution, we found substitutions in the nucleotide

sequence for each of the two DNA gyrases (*gyrA* and *gyrB*) targeted by fluoroquinolones, within

the canonical fluoroquinolone resistance determining gene *parC*, in the biofilm-associated gene

morA, but also in *dnaN* which was among the set of randomly chosen genes. Notably, all highly

significant pairs of substitutions were synonymous, and hence unlikely to affect drug resistance

or fitness (but see Plotkin and Kudla, 2011; Bailey *et al.*, 2014; Agashe *et al.*, 2016), save for one

pair of nonsynonymous sites, *gyrA* (c248t)-*parC* (c260g/t), that presumably impacts the level of

resistance (Fig. 1). The next most strongly correlated pair of nonsynonymous substitutions in-

volved *parC* and an uncharacterized gene at locus tag *PA14_34000*, but its significance was sev-

eral orders of magnitude lower than any of the nine strongest correlated pairs ($P = 8.018 \times 10^{-7}$

here *vs.* $P \leq 10^{-11}$ above). In total, we found evidence of correlated evolution among 96 paired

nonsynonymous substitutions with $P \leq 10^{-4}$, and many more with $P \leq 10^{-2}$. However, most

involved uncharacterized genes while the rest involved genes without sufficient biological ra-

tionale to justify investigation within our study. It should be noted that the algorithm we em-

ployed to detect correlated evolution (*i.e.* Pagel's algorithm Pagel, 1994), just like pretty much

any alternative phylogeny-aware methods used in comparative studies (*e.g.*, Maddison, 1990;

Huelsenbeck *et al.*, 2003; Maddison *et al.*, 2007), can detect a significant statistical association

even when there is a single case of co-distribution of traits (Maddison and FitzJohn, 2014; Uyeda

et al., 2018). This is why, despite the care taken in our approach, we performed additional exper-

imental analyses to validate the statistical signal detected between the most strongly correlated

pairs.

3.2 Experimental validation of the nonsynonymous pair

To test if epistasis was driving the correlated evolution of these fluoroquinolone resistance sub-

stitutions in *gyrA-parC*, we used a modified allelic replacement protocol (Melnyk *et al.*, 2017) to

construct mutants bearing all possible combinations of *gyrA-parC* single and double mutations

in two genetic backgrounds (PA01 and PA14) that lack the mutations of interest (Methods).

We then quantified the MIC of a fluoroquinolone antibiotic (ciprofloxacin) for all genotypes

and their competitive fitness against the ancestral (wild-type [WT]) genotypes in the absence

of drug. We used the PA01 and PA14 genetic backgrounds because they are phylogenetically

distinct (Dettman *et al.*, 2013), representing two major clades (Kos *et al.*, 2015).

Our results reveal that the pattern of changes in resistance for single and double mutants was

similar for both genotypes (Fig. 2a,b). On its own, the *gyrA* mutation increases resistance, by

$\sim$ 8-fold in PA14 and at least twice that in PA01, whereas the *parC* mutations on their own have

no effect. In combination, however, the *gyrA* mutation together with either of the *parC* mutations

confers between 128-256 fold increases in resistance, depending on genetic background. Support

for this interpretation comes from a likelihood ratio test for significant fixed effects in a mixed

effects model using biological replicates as a random effect. Results of the tests show that MIC

depends on both the genetic background ($X^2 = 46.52$, $df = 1$, $P = 9.059 \times 10^{-12}$) and mutation

$(X^2 = 813.95, df = 6, P < 2.2 \times 10^{-16})$, but also that the magnitude of the fold-increase in MIC was similar for PA01 and PA14 constructs (i.e. interaction term, $X^2 = 7.47, df = 5$, $P = 0.1878$). This analysis confirms the presence of strong positive epistasis for resistance between these *gyrA-parC* mutants, the magnitude of which is independent of the two genetic backgrounds tested.

Notably, mutants carrying the *gyrA c248t* substitution had the expected 8-32 fold increase in ciprofloxacin resistance (Akasaka *et al.*, 2001). However, the *parC c260g/t* substitutions did not affect resistance on their own but did show a $\sim$ 16 fold additional increase for double mutants (Fig. 2 a,b; Table 1). This strong epistasis for resistance could explain why *gyrA c248t* and *parC c260g/t* double mutants are common among fluoroquinolone resistant clinical isolates of *P. aeruginosa* (Akasaka *et al.*, 2001; Kos *et al.*, 2015) and is consistent with previous work showing that the fluoroquinolone resistance conferred by substitutions in *parC* depended upon mutations in *gyrA* (Moon *et al.*, 2010). We note that Akasaka *et al.* (2001) quantified the effect on MIC of both the *gyrA* and *parC* mutations by measuring decatenation activity of purified protein and found that both mutations would increase resistance. This result suggests a possible Bateson-like molecular mechanism of the observed epistasis for resistance (Bateson, 1911). As ciprofloxacin acts by inhibiting both DNA gyrase and topoisomerase IV (the protein products of *gyrA* and *parC*, respectively), the changed decatenation activity of topoisomerase IV caused by the *parC* substitution may be masked if DNA replication is prevented by non-functional DNA gyrase. Since *gyrA* single mutants can grow under elevated fluoroquinolones levels, but *parC* single mutants cannot, it is reasonable to posit that the *gyrA* mutation compensates for the inhibition of *parC*, or that fluoroquinolones do not entirely inhibit *parC* protein activity. A study of the molecular basis for this epistasis for resistance may provide valuable insight in the development of next generation fluoroquinolones.

The results for fitness costs associated with resistance mutations tell quite a different story

(Fig. 2c,d). We found no evidence for a cost of resistance associated with the *gyrA* and *parC* mu-

tations, either singly or in combination, in the PA14 background, a result that has been observed

previously (Melnyk *et al.*, 2015). By contrast, all the single and double mutations in PA01 lead

to significant fitness costs, and the double mutant including the *parC c260g* mutation, exhibits

significant positive epistasis (Table 2). In other words, the resistance mutations compensate for

each other, leading to costs that are lower than expected from their additive effects. Our results

support the idea that the cost of antibiotic resistance in bacteria can vary substantially across

genetic backgrounds (Melnyk *et al.*, 2015). Combinations of mutations in *gyrA* and *parC* other

than those tested here have been shown to be cost-free in *Streptococcus pneumoniae* (Gillespie

et al., 2002), reinforcing the idea that the effect of a mutation, or combination of mutations,

depends intimately on genetic background. Moreover, our observation of positive epistasis for

resistance in both genetic backgrounds, but not consistently for fitness costs, indicates that the

correlated evolution detected in *gyrA* and *parC* is likely driven by epistatic effects on antibiotic

resistance, rather than competitive fitness in antibiotic-free environments.

Six of the genes in our twelve gene-by-exome analysis were expected to evolve in response

to fluoroquinolone selection. Yet, we only had sufficient evidence to support experimental val-

idation of epistasis between a single pair of sites canonically associated with fluoroquinolone

resistance mutations. Two questions remained: (i) how come AEGIS did not identify correlated

evolution involving additional nonsynonymous resistance mutations, and (ii) what was/were the

mechanism/-s underlying the highly significant correlation identified for the eight pairs of syn-

onymous substitutions?

3.3 Resistance is strongly associated with only two substitutions in our alignment

The computational approach employed above identified only two nonsynonymous substitutions likely to be correlated during the evolution of resistance to fluoroquinolone antibiotics (Wibowo, 2013; Kos *et al.*, 2015). Our lack of success in identifying more pairs of nonsynonymous substitutions could be because no other resistance mutations in our alignment evolved in a correlated manner, or because AEGIS is only able to detect instances of strong correlated evolution (Nshogozabahizi *et al.*, 2017), or perhaps that only a few substitutions in our dataset are strongly associated with drug resistance. We therefore performed three additional analyses to uncover further substitutions associated with drug resistance in our dataset.

First, we quantified the importance of all polymorphic positions in the *P. aeruginosa* exome for predicting fluoroquinolone resistance (Long *et al.*, 2019). We did this by training a supervised machine-learning algorithm, based on adaptive boosting, which measured the strength of correlation between substitutions in our alignment and the resistance phenotype. This approach detected that the *gyrA 248* and *parC 260* nucleotide positions were the only two sites in our alignment that strongly predicted resistance (Fig. S4). These two positions were also the only ones among the 100 most important associations that were located in genes expected to evolve in response to fluoroquinolone selection (Fig. S4).

Second, we tested the association between fluoroquinolone resistance and nonsynonymous mutations at known fluoroquinolone resistance determining sites (Akasaka *et al.*, 2001; Kos *et al.*, 2015). From our alignment, the *gyrA 248* (*T83I*) and *parC 260* (*S87L* and *S87W*) mutations were present in 41.4% and 32.9% of all strains, respectively, and these mutations were correlated to the resistance phenotype (χ^2 tests: *gyrA 248*, $X^2 = 231.941$, $df = 1$, $P = 2.250 \times 10^{-52}$; *parC 260*, $X^2 = 177.21$, $df = 1$, $P = 1.98 \times 10^{-40}$, Table S2). We note that another study has

shown evidence for epistasis in the context of drug resistance when mutations in *parC* (*S87L* and *S87W*) co-occur with a mutation in *gyrA*, but at a different position (*S91F* instead of *T83I*; Schubert *et al.*, 2019). In contrast, nonsynonymous substitutions at additional amino acid sites known to confer fluoroquinolone resistance were identified in our alignment, but either did not correlate with resistance (at $P \leq 0.05$), or represented less than 10% of all strains included in our analysis (max. 8.74%, mean (sd) 3.54 (1.81)%; Table S2). Previous work showed that the signal of correlated evolution (i.e. the sensitivity of the AEGIS algorithm) is maximized when roughly half the strains are mutants, and when these mutations are evenly distributed throughout the phylogeny (Nshogozabahizi *et al.*, 2017), which is not the case here.

Third, we searched for loss of function mutations that are known to occur under fluoroquinolone selection, such as those affecting the *nfxB* or *orfN* genes (Wong *et al.*, 2012). There were no examples of premature stop codons within the sequences of these two genes in our alignment, although the presence of nonsynonymous loss of function mutations cannot be ruled out. Although it is in principle possible to resort to branch-site codon models to identify positions under selection in branches of interest (Yang and Nielsen, 2002; Zhang *et al.*, 2005), and hence detect sites potentially involved in drug resistance, our alignment contains too few taxa and too little evolutionary depth for proper statistical analysis (Arenas, 2015). Altogether, these three lines of evidence suggest that among the subset of known fluoroquinolone resistance mutations, only the *gyrA c248t* and *parC c260g/t* substitutions are present in enough strains to have been detected as correlated and show a significant association with fluoroquinolone resistance in our alignment.

3.4 Synonymous substitutions: the role of multiple drivers

AEGIS provided strong evidence for correlated evolution among synonymous mutations. This result is surprising because synonymous mutations are often assumed to evolve neutrally, and

so we would not *a priori* expect them to exhibit strong correlated evolution unless they were

under strong selection or were physically linked. This result could be due to strong phylogenetic

uncertainty, which can affect comparative methods (Shoji *et al.*, 2016; Guigueno *et al.*, 2019).

To account for this we could rerun the analyses on bootstrapped trees (Nshogozabahizi *et al.*,

2017) but this is a taxing solution as it would be computationally quite demanding, especially

here, possessing statistical properties that would deserve scrutiny at a level beyond the scope of

the present work. However, the phylogenetic tree that we reconstructed has internal nodes that

are fairly well supported (Fig. S2), making it unlikely that the signal of correlated evolution that

we detect between synonymous mutations be due to phylogenetic uncertainty.

On the other hand, it has been known for some time that synonymous sites can experience

purifying selection (Zhou *et al.*, 2010; Lawrie *et al.*, 2013), and a number of recent reports show

that they can sometimes contribute to adaptation as well (Bailey *et al.*, 2014; Agashe *et al.*, 2016).

However, we find no signal of epistasis-driven selection acting on the most strongly correlated

pairs of synonymous sites identified in our study (Fig. 1). Estimates of the change in stability

of mRNA transcripts and index of translation efficiency, which are proxies for translation rate

(Kudla *et al.*, 2009) and efficiency (Xia, 2014), respectively, relative to the wild type are shown

in Fig. 3 and Tables S2,3. While two of the synonymous mutations in *parC* (*c*1581*t* and *c*1587*t*)

could produce more stable and abundant mRNA transcripts, the only evidence for epistasis in

either metric we could identify was for translation efficiency in the mutations in *morA*, and the

direction of this effect is opposite to what would be expected if it were to result in positive

selection.

By contrast, two lines of evidence point to physical linkage between sites as the proximate

cause of correlation between synonymous mutations. First, the proportion of synonymous pairs

occurring within the same gene was higher than expected by chance among all observed (syn-

onymous and nonsynonymous) pairs ($X^2 = 7246.037$, $df = 2$, $P \leq 2.16 \times 10^{-16}$; see Table

S6). Second, the physical distance between pairs of correlated synonymous substitutions, as mapped on the circular genome of PA14, was negatively associated with the strength of correlation (for pairs with $P \leq 10^{-4}$; $t = -5.1784$, $df = 7,528$, $P = 2.29 \times 10^{-7}$ Fig. S5) , though the correlation was quite poor ($R^2 = 0.03$). Physical distance remains significantly associated to strength of correlation even when the 8 most significantly correlated pairs are removed substitutions ($t = -6.922$, $df = 7,520$, $P \approx 4.8 \times 10^{-12}$, $R^2 = 0.006$).

The mechanism responsible for the close physical linkage between synonymous mutations is less obvious. One possibility is that both hitchhiked to high frequency alongside a third, nonsynonymous mutation in the same gene that was under strong positive selection. We found a nonsynonymous mutation at nucleotide position 1374 of *gyrB*, which is near the gene's quinolone-resistance determining region, that was always present alongside the synonymous mutations. Moreover 97% of the strains that have substituted the pair of synonymous mutations in *morA* also fixed nonsynonymous mutations at other positions along the gene that could have functional effects on regulating or sensing internal oxygen levels (positions 292, 293, 306) (Taylor and Zhulin, 1999) or signal transduction (position 1356) (The UniProt Consortium, 2017). While these results point to hitchhiking as a plausible explanation for the correlation between sites, we failed to detect additional functional links with nonsynonymous mutations in the other genes. Moreover, AEGIS did not detect correlations between any of the correlated synonymous mutations and additional nonsynonymous mutations, even at $P \leq 10^{-4}$ (Fig. S1), suggesting that hitchhiking is not a general explanation for the correlated evolution of synonymous mutations in this data set. It should further be noted for the correlated pairs of substitutions found in *gyrB* and *morA*, that unless the pair of mutations repeatedly arose in the context of a third substitutions under selection, hitchhiking would not have produced a signal of correlated evolution.

A second possible mechanism we considered is horizontal gene transfer (HGT) and/or incomplete lineage sorting. Comparing the similarity of the species tree and the gene trees for

each of the four genes with a strongly correlated pair of synonymous mutations reveals little correspondence between the two for all but *parC* (Table 3), suggesting widespread recombination in our strain collection. To further assess the prevalence of recombination, we included tRNA genes located in close proximity of each of our four focal genes in the PA14 assembly. The rationale for this analysis was that both tRNA genes and three of our four focal genes, being involved in information processing, should interact with a large number of other gene products (transcripts and/or proteins), so that a successful HGT of such genes should involve the co-transfer of most of their interacting partners, an unlikely event (Jain *et al.*, 1999; Aris-Brosou, 2005). However, we found that all tRNA gene trees differed from each other and the species tree (Table 3) suggesting that recombination (and thus HGT and/or incomplete lineage sorting) may be widespread in our strain collection. Widespread HGT is consistent with the findings of Kos *et al.* (2015), who found that the *parE* and *β*-lactamase genes were horizontally transferred in a number of the strains included in our exome alignment.

Importantly, both hitchhiking and HGT represent population-level processes that can lead to a pattern of physical linkage between synonymous sites. However, neither mechanism explains how such strong intra-gene correlations arose in the first place. Functional constraints, for example via intra-molecular pairing that forms strong hairpins in mRNA, and locally-biased mutation that depends on nucleotide context, have been proposed as potential mechanisms (Tsunoyama *et al.*, 2001). A full analysis of the contribution of these processes to the patterns we observe here, while interesting, is beyond the scope of this work, however.

Altogether, these results suggest that the close physical linkage between pairs of synonymous mutations is due in part to widespread HGT, except in the *parC* gene. Without any evidence that either HGT or hitchhiking with a nonsynonymous mutation are driving the correlated evolution of the four synonymous mutations in *parC*, which form a network of five correlated pairs (Fig. 1), these mutations could in principle have evolved repeatedly as a driverless cohort - that is, muta-

₄₁₂ tions that self-assemble by chance, and drift to fixation via linkage (Buskirk *et al.*, 2017). How-

₄₁₃ ever, both the species tree (results of AEGIS analysis) and the gene trees (Fig. S3) show that these

₄₁₄ four synonymous *parC* substitutions have evolved independently numerous times. It is therefore

₄₁₅ unlikely that a cohort of four substitutions repeatedly self-assembled by chance. Whether these

₄₁₆ mutations co-occur for selective reasons related to the ecology of the CF lung (Sriramulu *et al.*,

₄₁₇ 2005; Smith *et al.*, 2006; Poole, 2005) and antibiotic treatment (Kos *et al.*, 2015) or other pro-

₄₁₈ cesses such as functional constraints associated with transcription and translation or mutational

₄₁₉ biases remains unclear.

₄₂₀ 4 CONCLUSIONS

₄₂₁ Our results show many instances of correlated evolution ($\sim 127,000$ interactions at $P \leq 0.01$

₄₂₂ from $\sim 10,220,000$ comparisons tested - *i.e.* : $\sim 1.2\%$). In the context of antibiotic resistance

₄₂₃ in *P. aeruginosa*, moreover, some instances of correlated evolution can be underlain by, among

₄₂₄ other processes, epistasis. This result is consistent with experimental work showing that multiple

₄₂₅ mutations can contribute to the evolution of antibiotic resistance (Trindade *et al.*, 2009; MacLean

₄₂₆ *et al.*, 2010; Hall and MacLean, 2011), both within and between genes (Lehner, 2011), although

₄₂₇ typically the number of mutations identified remains quite small. Our results suggest, unsurpris-

₄₂₈ ingly perhaps, that a wide range of mutations can evolve in a correlated manner in the context of

₄₂₉ antibiotic selection – but also that only a limited number of these mutations are easily interpreted

₄₃₀ in the context of drug resistance (here, those found in *gyrA* and *parC*). Furthermore, our results

₄₃₁ lend support to the idea that high-throughput techniques for identifying correlated evolution *can*

₄₃₂ lead to identifying epistasis. Similar interpretations have been made previously, for instance in

₄₃₃ the case of the evolution of RNA genes where base pairing is critical (Dutheil *et al.*, 2010), or

₄₃₄ in the context of genome-wide association studies, where the detection of interacting residues

is used as a first screen for epistasis, before fitting the linear models traditionally employed to identify genetic determinants of drug resistance (Schubert *et al.*, 2019). This latter approach, which notably also revealed epistatic interactions between *gyrA* and *parC*, requires that MIC values of all the genomes included in the analysis be known – data that was not available to us. In spite of this, both approaches should be highly complementary.

However correlated evolution is necessary but not sufficient evidence for epistasis (Nshogozabahizi *et al.*, 2017), as it can also arise through other mechanisms such as physical linkage. Computational approaches like ours detect relationships between phenotype and genomic substitutions though these associations are not evidence alone for a causal relationship. Indeed, we have observed a strong signal that physical linkage has driven the correlated evolution between pairs of synonymous substitutions in our data set, perhaps linked to the wholesale transfer of genes during recombination. Despite recent reports that synonymous mutations may not, in fact, be silent (Duan *et al.*, 2003; Chamary and Hurst, 2005; Plotkin and Kudla, 2011; Cuevas *et al.*, 2012; Bailey *et al.*, 2014; Agashe *et al.*, 2016; Fragata *et al.*, 2018), we have little evidence that synonymous substitutions, either on their own or in combination, impact fitness in our data. The fact that many of the most strongly correlated synonymous substitutions we identified here are GC→AT mutations (Fig. 3) reflects the pronounced GC bias of *P. aeruginosa* and reinforces the notion that these pairs of synonymous mutations have not been selected. It is notable that another computational study found evidence of physical linkage underlying correlated evolution between nonsynonymous but not synonymous substitutions and so attributed their co-occurrence to functional constraints (Callahan *et al.*, 2011). While this does not seem to be the case for the synonymous mutations in our analysis, together these results suggest that physical linkage may often be important in generating signals of correlated evolution. If so, this result suggests using caution when interpreting the results of computational studies that identify strong correlated evolution among traits or sites within a genome.

While computational methods provide us with a high-throughput means of determining candidates for epistasis, we often lack a sufficient understanding of epistasis at the molecular level to have confidence in this interpretation in the absence of additional evidence. Our work represents a first step in this direction: by using a computational approach to predict candidates for correlated evolution, with follow-up analyses suggesting that only a subset of these candidates are actually under epistasis, we were able to provide direct experimental evidence that epistasis underlies at least some instances of correlated evolution. However, this interpretation relies on our ability to conduct experimental validations under selective conditions that we presume to closely resemble those in which the sites evolved. This may not always be possible. Indeed, most of the strongest instances of correlated evolution in our data set appear to be unconnected to fitness or epistasis in an antibiotic selective context. As densely sampled population genomic data are not always available (such as the > 3000 genomes in Skwark *et al.* (2017)), we urge caution in interpreting instances of correlation between sites from high throughput computational analyses as solely due to epistasis.

ACKNOWLEDGEMENTS

We would like to thank Felipe Dargent and Matti Ruuskanen, as well as three anonymous reviewers, for their constructive feedbacks on the manuscript. We are grateful for the computational resources of the Centre for Advanced Computing at Queens University. This work was funded by the Natural Sciences and Engineering Research Council of Canada (S.A.B. and R.K.).

CONFLICT OF INTEREST

None declared.

DATA ACCESSIBILITY

All R scripts and complete results of the computational analysis of correlated evolution can be found at `github.com/JDench/sigResults_AEGIS_inVivo`.

ORCID

Jonathan Dench https://orcid.org/0000-0003-2193-8965

Stéphane Aris-Brosou https://orcid.org/0000-0003-4987-0296

References

Agashe, D., Sane, M., Phalnikar, K., Diwan, G. D., Habibullah, A., Martinez-Gomez, N. C., Sahasrabuddhe, V., Polachek, W., Wang, J., Chubiz, L. M., and Marx, C. J. 2016. Large-effect beneficial synonymous mutations mediate rapid and parallel adaptation in a bacterium. *Molecular Biology and Evolution*, 33(6): 1542–1553.

Akasaka, T., Tanaka, M., Yamaguchi, A., and Sato, K. 2001. Type II topoisomerase mutations in fluoroquinolone-resistant clinical strains of *Pseudomonas aeruginosa* isolated in 1998 and 1999: role of target enzyme in mechanism of fluoroquinolone resistance. *Antimicrob Agents Chemother*, 45(8): 2263–8.

Alfaro, E., Gámez, M., and García, N. 2013. adabag: An R package for classification with boosting and bagging. *Journal of Statistical Software*, 54(2): 1–35.

Angiuoli, S. V. and Salzberg, S. L. 2011. Mugsy: fast multiple alignment of closely related whole genomes. *Bioinformatics*, 27(3): 334–342.

Arenas, M. 2015. Trends in substitution models of molecular evolution. *Frontiers in Genetics*, 319(6).

Aris-Brosou, S. 2005. Determinants of adaptive evolution at the molecular level: the extended complexity hypothesis. *Mol Biol Evol*, 22(2): 200–9.

Aris-Brosou, S. and Rodrigue, N. 2012. The essentials of computational molecular evolution. In M. Anisimova, editor, *Evolutionary Genomics*, volume 855 of *Methods in Molecular Biology*, pages 111–152. Humana Press.

Aris-Brosou, S., Ibeh, N., and Noël, J. 2017. Viral outbreaks involve destabilized evolutionary networks: evidence from ebola, influenza and zika. *Sci Rep*, 7(1): 11881.

Baer, C. F. and Lynch, M. 2003. Correlated evolution of life-history with size at maturity in *Daphnia pulicaria*: patterns within and between populations. *Genetics Research*, 81(2): 123–132.

Bailey, S. F., Hinz, A., and Kassen, R. 2014. Adaptive synonymous mutations in an experimentally evolved *Pseudomonas fluorescens* population. *Nature Communications*, 5: 4076–4076.

Bateson, W. 1911. Review from cambridge: Mendel's principles of heredity. *The Edinburgh Review*, 213(435): 77.

Bertani, G. 1951. Studies on lysogenesis I. : The mode of phage liberation by lysogenic *Escherichia coli*. *Journal of Bacteriology*, 62(3): 293–300.

Blount, Z. D., Borland, C. Z., and Lenski, R. E. 2008. Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*. *Proceedings of the National Academy of Sciences*, 105(23): 7899–7906.

Buskirk, S. W., Peace, R. E., and Lang, G. I. 2017. Hitchhiking and epistasis give rise to cohort dynamics in adapting populations. *Proceedings of the National Academy of Sciences*, 114(31): 8330–8335.

Callahan, B., Neher, R. A., Bachtrog, D., Andolfatto, P., and Shraiman, B. I. 2011. Correlated evolution of nearby residues in drosophilid proteins. *PLoS Genetics*, 7(2): e1001315–e1001315.

Chamary, J. and Hurst, L. D. 2005. Evidence for selection on synonymous mutations affecting stability of mRNA secondary structure in mammals. *Genome Biology*, 6(9): R75.

Cheverud, J. M. 1984. Quantitative genetics and developmental constraints on evolution by selection. *Journal of Theoretical Biology*, 110(2): 155 – 171.

Choy, W.-K., Zhou, L., Syn, C. K.-C., Zhang, L.-H., and Swarup, S. 2004. MorA defines a new class of regulators affecting flagellar development and biofilm formation in diverse *Pseudomonas* species. *Journal of Bacteriology*, 186(21): 7221–7228.

Cuevas, J. M., Domingo-Calap, P., and Sanjuán, R. 2012. The fitness effects of synonymous mutations in DNA and RNA viruses. *Molecular Biology and Evolution*, 29(1): 17–20.

Darling, A. C. E., Mau, B., Blattner, F. R., and Perna, N. T. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res*, 14(7): 1394–403.

Dettman, J. R., Rodrigue, N., Aaron, S. D., and Kassen, R. 2013. Evolutionary genomics of epidemic and nonepidemic strains of *Pseudomonas aeruginosa*. *Proc. of the Nat. Acad. of Sci. of the U.S.A.*, 110(52): 21065–21070.

Duan, J., Wainwright, M. S., Comeron, J. M., Saitou, N., Sanders, A. R., Gelernter, J., and Gejman, P. V. 2003. Synonymous mutations in the human dopamine receptor D2 (DRD2) affect mRNA stability and synthesis of the receptor. *Human Molecular Genetics*, 12(3): 205–216.

Dutheil, J. Y., Jossinet, F., and Westhof, E. 2010. Base pairing constraints drive structural epistasis in ribosomal RNA sequences. *Molecular Biology and Evolution*, 27(8): 1868–1876.

Falconer, D. S. 1960. *Introduction to quantitative genetics*. Oliver And Boyd; Edinburgh; London.

Fragata, I., Matuszewski, S., Schmitz, M. A., Bataillon, T., Jensen, J. D., and Bank, C. 2018. The fitness landscape of the codon space across environments. *Heredity*, 121(5): 422–437.

Galperin, M. Y., Makarova, K. S., Wolf, Y. I., and Koonin, E. V. 2015. Expanded microbial genome coverage and improved protein family annotation in the COG database. *Nucleic Acids Research*, 43(D1): D261–D269.

Gillespie, S. H., Voelker, L. L., and Dickens, A. 2002. Evolutionary barriers to quinolone resistance in *Streptococcus pneumoniae*. *Microbial Drug Resistance*, 8(2): 79–84.

Goh, C.-S., Bogan, A. A., Joachimiak, M., Walther, D., and Cohen, F. E. 2000. Co-evolution of proteins with their interaction partners. *Journal of Molecular Biology*, 299(2): 283 – 293.

Guigueno, M. F., Shoji, A., Elliott, K. H., and Aris-Brosou, S. 2019. Flight costs in volant vertebrates: A phylogenetically-controlled meta-analysis of birds and bats. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, 235: 193 – 201.

Hall, A. R. and MacLean, R. C. 2011. Epistasis buffers the fitness effects of rifampicin-resistance mutations in *Pseudomonas aeruginosa*. *Evolution*, 65(8): 2370–9.

Huelsenbeck, J. P., Nielsen, R., and Bollback, J. P. 2003. Stochastic Mapping of Morphological Characters. *Systematic Biology*, 52(2): 131–158.

Ibeh, N., Nshogozabahizi, J. C., and Aris-Brosou, S. 2016. Both epistasis and diversifying selection drive the structural evolution of the Ebola virus glycoprotein mucin-like domain. *J Virol*, 90(11): 5475–84.

Jain, R., Rivera, M. C., and Lake, J. A. 1999. Horizontal gene transfer among genomes: The complexity hypothesis. *Proceedings of the National Academy of Sciences of the United States of America*, 96(7): 3801–3806.

Kelley, J. L., Fitzpatrick, J. L., and Merilaita, S. 2013. Spots and stripes: ecology and colour pattern evolution in butterflyfishes. *Proceedings of the Royal Society of London B: Biological Sciences*, 280(1757).

Kos, V. N., Déraspe, M., McLaughlin, R. E., Whiteaker, J. D., Roy, P. H., Alm, R. A., Corbeil, J., and Gardner, H. 2015. The resistome of *Pseudomonas aeruginosa* in relationship to phenotypic susceptibility. *Antimicrobial Agents and Chemotherapy*, 59(1): 427–436.

Kryazhimskiy, S., Rice, D. P., Jerison, E. R., and Desai, M. M. 2014. Global epistasis makes adaptation predictable despite sequence-level stochasticity. *Science*, 344(6191): 1519–1522.

Kudla, G., Murray, A. W., Tollervey, D., and Plotkin, J. B. 2009. Coding-sequence determinants of gene expression in escherichia coli. *Science*, 324.

Lawrie, D. S., Messer, P. W., Hershberg, R., and Petrov, D. A. 2013. Strong purifying selection at synonymous sites in d. melanogaster. *PLoS Genetics*, 9(5): 1–18.

Lehner, B. 2011. Molecular mechanisms of epistasis within and between genes. *Trends in Genetics*, 27(8): 323–331.

Lenski, R. E., Rose, M. R., Simpson, S. C., and Tadler, S. C. 1991. Long-term experimental evolution in *Escherichia coli*. i. adaptation and divergence during 2,000 generations. *The American Naturalist*, 138(6): 1315–1341.

Long, G. S., Hussen, M., Dench, J., and Aris-Brosou, S. 2019. Identifying genetic determinants of complex phenotypes from whole genome sequence data. *BMC Genomics*, 20(1): 470.

Lynch, M. and Walsh, B. 1998. *Genetics and analysis of quantitative traits*, volume 1. Sinauer Sunderland, MA.

MacLean, R. C., Perron, G. G., and Gardner, A. 2010. Diminishing returns from beneficial mutations and pervasive epistasis shape the fitness landscape for rifampicin resistance in pseudomonas aeruginosa. *Genetics*, 186(4): 1345–54.

Maddison, W. P. 1990. A method for testing the correlated evolution of two binary characters: Are gains or losses concentrated on certain branches of a phylogenetic tree? *Evolution*, 44(3): 539–557.

Maddison, W. P. and FitzJohn, R. G. 2014. The Unsolved Challenge to Phylogenetic Correlation Tests for Categorical Characters. *Systematic Biology*, 64(1): 127–136.

Maddison, W. P., Midford, P. E., and Otto, S. P. 2007. Estimating a Binary Character's Effect on Speciation and Extinction. *Systematic Biology*, 56(5): 701–710.

Melnyk, A. H., Wong, A., and Kassen, R. 2015. The fitness costs of antibiotic resistance mutations. *Evol Appl*, 8(3): 273–83.

Melnyk, A. H., McCloskey, N., Hinz, A. J., Dettman, J., and Kassen, R. 2017. Evolution of cost-free resistance under fluctuating drug selection in *Pseudomonas aeruginosa*. *mSphere*, 2(4).

Moon, D. C., Seol, S. Y., Gurung, M., Jin, J. S., Choi, C. H., Kim, J., Lee, Y. C., Cho, D. T., and Lee, J. C. 2010. Emergence of a new mutation and its accumulation in the topoisomerase iv gene confers high levels of resistance to fluoroquinolones in *Escherichia coli* isolates. *International Journal of Antimicrobial Agents*, 35(1): 76–79.

Nshogozabahizi, J. C., Dench, J., and Aris-Brosou, S. 2017. Widespread historical contingency in influenza viruses. *Genetics*, 205(1): 409–420.

Pagel, M. 1994. Detecting correlated evolution on phylogenies: A general method for the comparative analysis of discrete characters. *Proceedings: Biological Sciences*, 255(1342): 37–45.

Pagel, M. and Meade, A. 2006. Bayesian analysis of correlated evolution of discrete characters by reversible-jump Markov chain Monte Carlo. *American Naturalist*, 167(6): 808–825.

Plotkin, J. B. and Kudla, G. 2011. Synonymous but not the same: the causes and consequences of codon bias. *Nat Rev Genet*, 12.

Poole, K. 2005. Efflux-mediated antimicrobial resistance. *Journal of Antimicrobial Chemotherapy*, 56(1): 20–51.

Price, M. N., Dehal, P. S., and Arkin, A. P. 2010. FastTree 2 - approximately maximum-likelihood trees for large alignments. *PLoS ONE*, 5(3).

Ravenhall, M., Škunca, N., Lassalle, F., and Dessimoz, C. 2015. Inferring horizontal gene transfer. *PLOS Computational Biology*, 11(5): 1–16.

Roy, P. H., Tetu, S. G., Larouche, A., Elbourne, L., Tremblay, S., Ren, Q., Dodson, R., Harkins, D., Shay, R., Watkins, K., Mahamoud, Y., and Paulsen, I. T. 2010. Complete genome sequence of the multiresistant taxonomic outlier *Pseudomonas aeruginosa* PA7. *PLoS ONE*, 5(1): e8842.

Schubert, B., Maddamsetti, R., Nyman, J., Farhat, M. R., and Marks, D. S. 2019. Genome-wide discovery of epistatic loci affecting antibiotic resistance in neisseria gonorrhoeae using evolutionary couplings. *Nature Microbiology*, 4(2): 328–338.

Sexton, J. P., McIntyre, P. J., Angert, A. L., and Rice, K. J. 2009. Evolution and ecology of species range limits. *Annual Review of Ecology Evolution and Systematics*, 40: 415–436.

Shimizu, A., Dohzono, I., Nakaji, M., Roff, D. A., Miller III, D. G., Osato, S., Yajima, T., Niitsu, S., Utsugi, N., Sugawara, T., and et al. 2014. Fine-tuned bee-flower coevolutionary state hidden within multiple pollination interactions. *Scientific Reports*, 4(1).

Shimodaira, H. 2002. An approximately unbiased test of phylogenetic tree selection. *Syst Biol*, 51(3): 492–508.

Shimodaira, H. and Hasegawa, M. 2001. CONSEL: for assessing the confidence of phylogenetic tree selection. *Bioinfo*, 17(12): 1246–1247.

Shoji, A., Aris-Brosou, S., and Elliott, K. 2016. Physiological constraints and dive behavior scale in tandem with body mass in auks: A comparative analysis. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, 196: 54 – 60.

Skwark, M. J., Croucher, N. J., Puranen, S., Chewapreecha, C., Pesonen, M., Xu, Y. Y., Turner, P., Harris, S. R., Beres, S. B., Musser, J. M., Parkhill, J., Bentley, S. D., Aurell, E., and Corander, J. 2017. Interacting networks of resistance, virulence and core machinery genes identified by genome-wide epistasis analysis. *PLOS Genetics*, 13(2): 1–24.

Smith, E. E., Buckley, D. G., Wu, Z., Saenphimmachak, C., Hoffman, L. R., D'Argenio, D. A., Miller, S. I., Ramsey, B. W., Speert, D. P., Moskowitz, S. M., Burns, J. L., Kaul, R., and Olson, M. V. 2006. Genetic adaptation by pseudomonas aeruginosa to the airways of cystic fibrosis patients. *Proceedings of the National Academy of Sciences of the United States of America*, 103(22): 8487–8492.

Sorrells, T. R., Booth, L. N., Tuch, B. B., and Johnson, A. D. 2015. Intersecting transcription networks constrain gene regulatory evolution. *Nature*.

Sriramulu, D. D., Lünsdorf, H., Lam, J. S., and Römling, U. 2005. Microcolony formation: a novel biofilm model of pseudomonas aeruginosa for the cystic fibrosis lung. *Journal of Medical Microbiology*, 54(7): 667–676.

Starkey, M., Hickman, J. H., Ma, L., Zhang, N., De Long, S., Hinz, A., Palacios, S., Manoil, C., Kirisits, M. J., Starner, T. D., Wozniak, D. J., Harwood, C. S., and Parsek, M. R. 2009. *Pseudomonas aeruginosa* rugose small-colony variants have adaptations that likely promote persistence in the cystic fibrosis lung. *Journal of Bacteriology*, 191(11): 3492–3503.

Tatusov, R. L., Koonin, E. V., and Lipman, D. J. 1997. A genomic perspective on protein families. *Science*, 278(5338): 631–637.

Taylor, B. L. and Zhulin, I. B. 1999. Pas domains: Internal sensors of oxygen, redox potential, and light. *Microbiology and Molecular Biology Reviews*, 63(2): 479–506.

The UniProt Consortium 2017. UniProt: the universal protein knowledgebase. *Nucleic Acids Research*, 45(D1): D158–D169.

Trindade, S., Sousa, A., Xavier, K. B., Dionisio, F., Ferreira, M. G., and Gordo, I. 2009. Positive epistasis drives the acquisition of multidrug resistance. *PLoS Genetics*, 5(7): e1000578–e1000578.

Tsunoyama, K., Bellgard, M. I., and Gojobori, T. 2001. Intragenic variation of synonymous substitution rates is caused by nonrandom mutations at methylated cpg. *Journal of Molecular Evolution*, 53(4): 456–464.

Uyeda, J. C., Zenil-Ferguson, R., and Pennell, M. W. 2018. Rethinking phylogenetic comparative methods. *Systematic Biology*, 67(6): 1091–1109.

Wibowo, A. 2013. Performance standards for antimicrobial susceptibility testing: 23rd informational supplement m100-s23. In *Performance standards for antimicrobial susceptibility testing: 23rd informational supplement M100-S23*, volume 33 of *Developments in Environmental Modelling*, pages 62–65. Clinical and Laboratory Standards Institute.

Winsor, G. L., Griffiths, E. J., Lo, R., Dhillon, B. K., Shay, J. A., and Brinkman, F. S. L. 2016. Enhanced annotations and features for comparing thousands of pseudomonas genomes in the pseudomonas genome database. *Nucleic Acids Res*, 44(D1): D646–53.

Wong, A., Rodrigue, N., and Kassen, R. 2012. Genomics of adaptation during experimental evolution of the opportunistic pathogen *Pseudomonas aeruginosa*. *PLoS Genet*, 8(9): e1002928.

Xia, X. 2014. A major controversy in codon-anticodon adaptation resolved by a new codon usage index. *Genetics*, 199(2): 573–579.

Yang, Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol*, 24(8): 1586–91.

Yang, Z. and Nielsen, R. 2002. Codon-substitution models for detecting molecular adaptation at individual sites along specific lineages. *Molecular biology and evolution*, 19(6): 908—917.

Zhang, J., Nielsen, R., and Yang, Z. 2005. Evaluation of an improved branch-site likelihood method for detecting positive selection at the molecular level. *Molecular biology and evolution*, 22(12): 2472–2479.

Zhou, T., Gu, W., and Wilke, C. O. 2010. Detecting positive and purifying selection at synonymous sites in yeast and worm. *Molecular biology and evolution*, 27(8): 1912–1922.

 # Tables

Table 1. Results from MIC assays of mutant constructs under ciprofloxacin. We measured MIC from four technical replicates for each of two independent constructs (biological replicates), except for *gyrA c*248*t* that had three independent constructs. The significance of differences in $\log_2$ MIC between wild-type and each mutant construct was assessed with the Dunn test and a Bonferonni correction; $P \leq 0.05$ shown in bold. Epistasis was measured with a multiplicative model, using the MIC determined from reads of the optical density (600 nm) after 24 hours growth, and measurement error was calculated using error propagation (Trindade *et al.*, 2009). There is evidence for epistasis when the absolute value of ϵ is greater than the error of our measures (in bold).

Background	Mutant		X^2	Z	P	ϵ	Error
PA14	*gyrA c*248*t*		6.480	-2.546	**0.005**		
	*parC c*260*t*		0.000	0.000	0.500		
	*parC c*260*g*		0.000	0.000	0.500		
	*gyrA c*248*t*	*parC c*260*t*	5.478	-2.341	**0.010**	**0.547**	0.382
	*gyrA c*248*t*	*parC c*260*g*	5.478	-2.341	**0.010**	**0.547**	0.382
PA01	*gyrA c*248*t*		4.253	-2.062	0.020		
	*parC c*260*t*		0.000	0.000	0.500		
	*parC c*260*g*		0.000	0.000	0.500		
	*gyrA c*248*t*	*parC c*260*t*	4.000	-2.000	**0.023**	**4.461**	2.051
	*gyrA c*248*t*	*parC c*260*g*	3.529	-1.879	**0.030**	**3.271**	3.138

Table 2. Results from competitive fitness assays of mutant constructs in permissive LB media. We measured fitness from six technical replicates for each of two independent constructs (biological replicates), except for $gyrA\ c248t$ which had three independent constructs. Relative fitness was calculated by dividing the fitness of each mutant construct by that of its wild-type (not shown here). The significance of differences in competitive fitness of wild-type and each mutant construct was assessed with the Dunn test and a Bonferonni correction; $P \leq 0.05$ shown in bold. Epistasis was measured with a multiplicative model and error was calculated using error propagation (Trindade *et al.*, 2009). There is evidence for epistasis when the absolute value of ϵ is greater than the error of our measures (in bold).

Background	Mutant		X^2	Z	P	ϵ	Error
PA14	*gyrA c248t*		1.778	1.333	0.456		
	parC c260t		3.509	1.873	0.153		
	parC c260g		0.561	0.749	1.000		
	gyrA c248t	*parC c260t*	1.010	1.005	0.787	0.016	0.024
	gyrA c248t	*parC c260g*	3.509	1.873	0.153	-0.003	0.021
PA01	*gyrA c248t*		8.266	2.875	**0.010**		
	parC c260t		7.364	2.714	**0.017**		
	parC c260g		8.576	2.929	**0.009**		
	gyrA c248t	*parC c260t*	11.000	3.317	**0.002**	0.019	0.021
	gyrA c248t	*parC c260g*	6.224	2.495	**0.032**	**0.022**	0.021

Table 3. Approximately unbiased (AU) tests of significant differences among phylogenetic trees. We built each tree using the multiple sequence alignment for one of the four focal genes that contain the most significantly correlated synonymous substitutions (*dnaN, gyrB, morA, parC*) and four tRNA genes unlikely, according to the complexity hypothesis (Jain *et al.*, 1999; Aris-Brosou, 2005) to undergo HGT (*arginyl tRNA synthetase, glycyl tRNA synthetase subunit beta, isoleucyl tRNA synthetase, methionyl tRNA formyltransferase*). We tested for significant differences among all the gene trees and the species tree used in the analysis of correlated evolution. Values in each cell represent the probability, calculated via the AU test, that a tree (column names) describes evolution observed in an alignment (row names). In bold are the "best" tree(s) (*i.e.*: insignificantly different trees).

Focal Gene	*dnaN*	*gyrB*	*morA*	*parC*	arginyl tRNA synthetase	glycyl tRNA synthetase subunit beta	isoleucyl tRNA synthetase	methionyl tRNA formyltransferase	Species Tree
dnaN	**1.000**	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
gyrB	0.000	**1.000**	0.000	0.000	0.000	0.000	0.000	0.000	0.000
morA	0.000	0.001	**1.000**	0.002	0.001	0.000	0.000	0.000	0.001
parC	0.000	0.000	0.000	**0.423**	0.000	0.000	0.000	0.000	**0.577**
arginyl tRNA synthetase	0.000	0.000	0.000	0.000	**1.000**	0.000	0.000	0.000	0.000
glycyl tRNA synthetase subunit beta	0.000	0.000	0.000	0.000	0.000	**1.000**	0.000	0.000	0.000
isoleucyl tRNA synthetase	0.000	0.000	0.000	0.000	0.000	0.000	**1.000**	0.000	0.000
methionyl tRNA formyltransferase	0.000	0.000	0.000	0.000	0.000	0.000	0.000	**1.000**	0.000

 # Figures

Figure 1. Correlated pairs of mutations with strongest evidence. (a) a histogram showing the distribution of significance for all correlated pairs with strong ($P < 10^{-7}$) evidence for correlated evolution. The purple bar highlights the strength of evidence for the pairs presented in B. (b) substitutions with the strongest evidence for correlated evolution ($P < 10^{-11}$). Circles represent a substitution in a gene (top text), at a particular nucleotide position (bottom text) on the coding strand. Colours show whether or not the substitution was synonymous, and whether or not it was found in one of the six genes expected to evolve in response to fluoroquinolone selection. Edges connect the predicted pairs of correlated substitutions.

Figure 2. Empirical tests of epistasis for resistance and fitness. The mean and 95% confidence interval of measurements for each genotype is represented by the dark black lines and colored rectangles respectively. The color used reflects the genetic background of genotypes while the hue identifies the biological replicate. (a,b) results of the ciprofloxacin MIC assay of *P. aeruginosa* WT and mutant constructs. Dashed horizontal lines highlight the MIC value for WT strains. Numbers to the left of mean (black) lines represent the MIC fold increase, compared to WT, of mutants. (c,d) results of the competitive fitness assays for WT and mutant construct genotypes. The relative fitness of WT strains is highlighted by the horizontal dashed red line. Mutants with relative fitness significantly different from wild-type are denoted with an asterisk (see Table 2).

Figure 3. Computationally predicted biological effects of the strongly correlated pairs of synonymous substitutions. Each set of bars is for a pair of substitutions, where the first bar (grey) is the wild-type value (inferred from PA14 reference genome), the second bar (red) is a mutant carrying the first substitution (A) listed below the bars while the third bar (blue) represents a mutant carrying the second listed substitution (B), and the fourth bar (purple) represents a double mutant. (a) estimates of the relative change of mRNA folding free energy (ΔG) are relative to the WT value (gold dashed line) and error bars represent the 95% confidence interval. (b) measures of the mean index of translation elongation (I_{TE}, B) with error bars presenting standard deviation among measures estimated from double mutant strains present in our data set.