Strengths and potential pitfalls of hay-transfer for ecological restoration revealed by RAD-seq analysis in floodplain *Arabis* species

Hannes Dittberner[1], Christian Becker[2], Wen-Biao Jiao[3], Korbinian Schneeberger[3], Norbert Hölzel[4], Aurélien Tellier[5], Juliette de Meaux*[1]

Affiliations: [1]Institute of Botany, University of Cologne, 50674 Cologne, Germany; Cologne Center for Genomics, University of Cologne, 50931 Cologne, Germany; [3]Max-Planck-Institute for Plant Breeding Research, 50829 Cologne, Germany; [4]Institute of Landscape Ecology, University of Münster, 48149 Münster, Germany;[5]Center of Life and Food Sciences Weihenstephan, Technical University of Munich, 85354, Freising, Germany

*Corresponding Author E-mail: jdemeaux@uni-koeln.de

Abstract

Achieving high intraspecific genetic diversity is a critical goal in ecological restoration as it increases the adaptive potential and long-term resilience of populations. Thus, we investigated genetic diversity within and between pristine sites in a fossil floodplain and compared it to sites restored by hay-transfer between 1997 and 2014. RAD-seq genotyping revealed that the stenoecious flood-plain species *Arabis nemorensis* is co-occurring with individuals that, based on ploidy, ITS-sequencing and morphology, probably belong to the close relative *Arabis sagittata*, which has a documented preference for dry calcareous grasslands but has not been reported in

floodplain meadows. We show that hay-transfer maintains genetic diversity for both species.

Additionally, in *A. sagittata*, transfer from multiple genetically isolated pristine sites resulted in

restored sites with increased diversity and admixed local genotypes. In *A. nemorensis,* transfer

did not create novel admixture dynamics because genetic diversity between pristine sites was

less differentiated. Thus, the effects of hay-transfer on genetic diversity also depend on the

genetic makeup of the donor communities of each species, especially when material is mixed.

Our results demonstrate the efficiency of hay-transfer for habitat restoration and emphasize the

importance of pre-restoration characterization of micro-geographic patterns of intraspecific

diversity of the community to guarantee that restoration practices reach their goal, i.e.

maximize the adaptive potential of the entire restored plant community. Overseeing these

patterns may differentially alter the adaptive potential of species present, thereby altering the

balance between species in the community. Additionally, our comparison of summary statistics

obtained from *de novo* and reference-based RAD-seq pipelines shows that the genomic impact

of restoration can be reliably monitored in species lacking prior genomic knowledge.

Keywords

Introduction

Habitat degradation is an ever growing problem in our modern world, causing unprecedented loss of biodiversity and essential ecosystem services (Baillie, Hilton-Taylor, & Stuart, 2004). Thus, there is a growing demand for ecological restoration, i.e. measures assisting the recovery of ecosystems that have been degraded, damaged or destroyed (Society for Ecological Restoration International Science & Policy Working Group, 2004). However, ecological restoration is a difficult process rarely leading to full ecosystem recovery (Benayas, Newton, Diaz, & Bullock, 2009). Thus, the young field of restoration ecology, which studies ecological processes in the light of restoration, is vital to improve restoration practices (Bullock, Aronson, Newton, Pywell, & Rey-Benayas, 2011; Roberts, Stone, & Sugden, 2009; Suding, 2011).

One of the main goals of ecological restoration is the recovery of biodiversity, including both species richness and intraspecific genetic diversity (henceforth called genetic diversity). Genetic diversity has generally positive effects on ecosystems (reviewed in Hughes, Inouye, Johnson, Underwood, & Vellend, 2008). For example, experimentally increasing the genetic diversity of *Solidago altissima* increased primary above-ground biomass productivity and arthropod diversity (Crutsinger et al., 2006). Genetic diversity also boosts resistance of populations to invasion and environmental fluctuations, presumably because it enhances the adaptive potential of populations (Reed & Frankham, 2003; Vrijenhoek, 1994). For example, high diversity experimental *Arabidopsis thaliana* populations showed higher resistance against invasion by *Senecio vulgaris* than low diversity populations (Scheepens, Rauschkolb, Ziegler, Schroth, & Bossdorf, 2017). Moreover, *Zoestra marina* populations with higher genetic diversity

showed increased biomass production, plant density and faunal abundance during an extremely

warm period (Reusch, Ehlers, Hämmerli, & Worm, 2005). Concordantly, restored populations of

Z. marina with increased genetic diversity showed longer plant survival, grew more rapidly and

provided enhanced ecosystem services, measured by increased primary productivity,

invertebrate density and nitrogen retention. This effect was stable in a range of environmental

conditions along a water-depth gradient (Reynolds, McGlathery, & Waycott, 2012). These

examples demonstrate the importance of genetic diversity for ecosystem function and stability

and hence the need to consider population genetics in the design and evaluation of restoration

practices, which is the focus of restoration genetics.

Restoration genetics can not only inform the planning of restoration efforts, e.g. by identifying

suitable source populations, but also help evaluating the success of restoration projects, e.g. by

monitoring genetic diversity in restored populations (Mijangos, Pacioni, Spencer, & Craig, 2015;

Williams, Nevill, & Krauss, 2014). In fact, studies comparing the level of genetic diversity in

pristine and restored populations frequently report limited success, with a reduction of genetic

diversity in restored populations. This decline in genetic diversity may be caused by genetic

bottlenecks in plant nurseries, biases introduced by seed harvesting strategies, founder effects

during recolonization and/or unreliable commercial seeds (reviewed in Mijangos, Pacioni,

Spencer, & Craig, 2015). By contrast, the transfer of seed-containing hay from pristine (donor)

to restoration (donee) sites, termed hay-transfer, is expected to limit the loss of genetic

diversity and maintain site-specific local adaption (Hufford & Mazer, 2003; Kiehl, Kirmer, &

Shaw, 2014). In addition, this method has the unique feature that it can, theoretically, restore

an entire community without altering the genetic composition of populations and thus is the

best method available for restoring entire ecosystems (Hölzel & Otte, 2003; Kiehl, Kirmer,

Donath, Rasran, & Hölzel, 2010). So far, however, there is no empirical support for the efficiency

of this practice (Bucharova et al., 2017), especially since many species maintain seed banks in

the soil. Indeed, the genetic diversity specific to the seed bank will not be sampled with the hay,

although it is known that it can contribute significantly to the maintenance of diversity (Tellier,

Laurent, Lainer, Pavlidis, & Stephan, 2011).

The field of restoration genetics has witnessed a major technological shift over recent years.

Restoration genetics studies have initially relied on microsatellites and AFLP markers and thus

provided a limited overview on patterns of genetic variation within and between restored or

pristine populations (reviewed in Mijangos et al., 2015). Now, genotyping-by-sequencing (GBS)

methods are beginning to be more broadly adopted (Gruenthal et al., 2014; Massatti, Doherty,

& Wood, 2018; O'Leary, Hollenbeck, Vega, Gold, & Portnoy, 2018; Torres-Martinez & Emery,

2016). These methods drastically reduce sequencing costs through strategies to sequence a

reduced portion of the genome, e.g. Restriction site Associated DNA-sequencing (RAD-seq)

(Elshire et al., 2011; Etter, Bassham, Hohenlohe, Johnson, & Cresko, 2011; Peterson, Weber,

Kay, Fisher, & Hoekstra, 2012). In contrast to previous methods (AFLP, microsatellites), GBS

approaches sample proportions of the genome that are sufficiently large to allow resolving

patterns of genetic diversity and spatial structure even at very local scale where overall levels of

genetic diversity are low (Bradbury et al., 2015; Jeffries et al., 2016; Reitzel, Herrera, Layden,

Martindale, & Shank, 2013). In principle, GBS approaches have a third major advantage: they

are well suited to unravel genetic diversity in non-model species without prior genomic

information. Yet, the accuracy of genotyping in the absence of a reliable reference genome has

been questioned (Shafer et al., 2016). Since target species in restoration projects rarely coincide

with species or genera with advanced prior genomic knowledge, it is important to assess

whenever possible, whether conclusions from RAD-seq-based restoration genetics studies

depend on the availability of a reference genome.

Flood-plain meadows are species-rich ecosystems, accommodating many endangered and

stenoecious species adapted to the variable moisture regime. Due to increased agricultural

land-use and river regulation a large proportion of flood-plain ecosystems are degraded and

have become a target for ecological restoration. Here, we use RAD-seq to evaluate whether

genetic diversity is maintained in flood-plain meadows restored by hay-transfer in the Upper

Rhine valley in Germany (Hölzel & Otte, 2003, Donath, Bissels, Hölzel, & Otte, 2007). We

focused on *Arabis nemorensis*, a stenoecious species typically restricted to flood-plain

meadows, and thus strongly endangered in Central Europe (Schnittler & Günther, 1999). *A.*

nemorensis is a short-lived, mostly biennial hemicryptophyte that is known to maintain a long-

lived soil seed bank (Hölzel & Otte, 2004). It is part of the *Arabis hirsuta* species aggregate,

which comprises several morphologically similar but ecologically diverse species, which diverged

about 1.2 million years ago (Karl & Koch, 2014).

We performed RAD-seq for over 130 plants collected across pristine and restored sites. This

dataset allows us to ask the following questions: i) what is the level of genetic diversity and

structuration of the pristine sites that served as source populations for restoration? ii) do

restored sites show a lower level of genetic diversity than the pristine sites? iii) how did

restoration affect the distribution of diversity within and among restored sites? iv) is the use of

a reference genome necessary to reliably characterize the impact of restoration on genetic

diversity? This work demonstrates that a thorough genetic analysis of source and restored sites

reveals the complex dynamics at stake in the restoration process. Through genetic analysis, we

reveal the unanticipated co-occurrence of *A. nemorensis* with *A. sagittata,* a morphologically

similar and closely related species from the *A. hirsuta* aggregate, which normally exhibits a

preference for drier habitats such as calcareous grasslands (Hand & Gregor, 2006). We further

show that restoration by hay-transfer has maintained and can even enhance local diversity. Our

analysis further shows that the use of a reference genome yields higher estimates of genetic

diversity, but does not affect the resolution of patterns of genetic variation within and between

sites, indicating that this approach can find broad applications in the field.

Methods

Plant material and DNA extraction

The sampling area comprises the fossil, dyke-protected flood plain of the River Rhine near

Riedstadt in Hessen, Germany. The area is dominated by arable fields, but also contains

remnants of pristine flood meadow communities in low lying depressions that are submerged

by ascending groundwater during high floods of the River Rhine. Since ca. 20 years, new flood

meadow communities have been restored on ex-arable land using the transfer of green hay

from the pristine sites as donors to overcome significant dispersal limitation (Hölzel & Otte,

2003). During this process, hay from different donor sites was placed in distinct yet adjacent

patches, making admixture possible. Since restoration is still ongoing, restored sites differ in age

(Table S1).

From previous monitoring of the sites we knew that populations that *Arabis* plants were not

present every year in all sites. Thus, we sampled in two consecutive years to maximize the

number of study sites. In the few sites sampled in both years, the genetic composition of the

samples was not markedly different across years, so samples from both years were bulked for

these sites (see Table S1, Figure S1). We harvested seeds from a total of 134 plants of *Arabis*

nemorensis/sagittata in nine sites named from A to I, in order of collection. The presence of *A.*

sagitata was not previously reported in these sites and was thus unexpected. Although some

individuals showed the reduced stem leaf density and shorter siliques typical of *A. sagittata*,

these phenotypic criteria were not always clearly distinguishable in the field, especially at the

end of the season when siliques matured, so that the presence of the two species has remained

overlooked in previous studies (Burmeier, Eckstein, Donath, & Otte, 2011). Species identity was

thus determined by post-hoc analysis, after the RAD-seq analysis revealed the presence of two

taxonomic units.

Four sites were sampled in pristine habitat (B, C, D and I) and five in restored habitat (A, E, F, G

and H, Figure 1). In site A two distinct stands of plants separated by about 100 m were sampled

and were treated as sub-sites A-1 and A-2 throughout this study. To produce material for DNA

extraction, we stratified seeds on wet filter paper for 6 days at 4°C in darkness. Afterwards we

sowed seeds in soil (33% VM, 33% ED-73, 33% Seramis (clay granules)). After 4 to 6 weeks of

growth in the greenhouse, we harvested about 200mg of leaf material from one offspring of

each wild parent (genotype). We homogenized freshly harvested leaf material using a Precellys

Evolution homogenizer (Bertin technologies) for 2x20 seconds at 6800 rpm. We extracted DNA

using the NucleoSpin Plant II Mini kit (Macherey-Nagel) following the manufacturer's

instructions. We verified DNA quality using gel-electrophoresis with a 0.8% agarose gel. We

measured DNA quantity using Qubit (broad-range kit) following manufacturer's instructions.

Figure 1: Overview map of studied sites
Each point on the map represents one site. The color of the points represents the type of the site, i.e.
whether it is pristine or restored. The purple square represents the area shown in the zoomed inset. All
sites, except site F are protected by a dyke.

Draft genome assembly

To facilitate genotype calling we assembled a draft genome for one *A. nemorensis* accession (ID

29). Library preparation and sequencing was done at the Cologne Center for Genomics. Three

libraries were created: one paired-end library with approximately 280 bp insert size creating an

overlap of 20 bp with 150 bp reads, and two mate-pair libraries with 3 kbp and 6 kbp inserts,

respectively. The paired-end library was created using the Illumina TruSeq® Nano DNA Library

Prep Kit, with 2µg of input DNA, without PCR. The mate-pair libraries were created using the

Illumina Nextera® Mate Pair Library Prep Kit, with 4 µg of input DNA and 10 cycles of PCR for the

3 kbp library and 15 cycles of PCR for the 6 kbp library. All libraries were sequenced together as

150bp paired-end reads as part of an Illumina HiSeq 4000 lane for a total of 66 Gbp.

We used *FastQC* (Andrews, 2010) to quality-check the resulting reads. We filtered the resulting

reads to remove reads shorter than 100bp and trimmed Illumina adapters using *Cutadapt*

(Martin, 2011). We assembled reads using the *ALLPATHS-LG* assembler (Gnerre et al., 2011)

with default settings, running it on the CHEOPS high-performance computing cluster of the

University of Cologne. The resulting contig assembly had a size of 199 Mbp and an N50 of 47

kbp. The scaffold assembly had a size of 206 Mbp and an N50 102 kbp. To further scaffold the

genome, we generated 2.9 Gbp of PacBio sequence data. Library preparation and sequencing

was done at the Max-Planck-Institute for Plant Breeding Research (Cologne, Germany). We

scaffolded the genome using *OPERA-LG* with default settings (Gao, Bertrand, Chia, & Nagarajan,

2016). This increased the N50 to 150kbp. To achieve chromosome-level assembly, we made use

of the available reference genome of *Arabis alpina* (Jiao et al., 2017; Willing et al., 2015). To

facilitate whole-genome alignment, repetitive regions in the genome were masked using

RepeatMasker (Smit, Hubley, & Green, 2013). We used 'brassicaceae' as the search term ('-

species' option) for the RepeatMasker repeat-database. We created the pseudo-chromosome

assembly using the *CoGe* website (Lyons & Freeling, 2008). We aligned our draft genome with

the *Arabis alpina* genome using *SynMap2* (Haug-Baltzell, Stephens, Davey, Scheidegger, &

Lyons, 2017) and afterwards performed synthenic path assembly (Lyons, Freeling, Kustu, &

Inwood, 2011) to assemble the chromosomes. The pseudo-chromosomes had a total size of 192

Mbp and were used for all following analyses.

The size of the genome was estimated by flow-cytometry, which was performed commercially

at Plant Cytometry Services. The estimated genome size was 274 Mbp. Thus, the assembly size

was 70% of the genome size.

Annotation

To detect and annotate Transposable elements (TE) specific to *A. nemorensis* (not in the

previously used database) we used the softwares RepeatModeler (Smit & Hubley, 2008) and

RepeatMasker (Smit et al., 2013). The consensus repeat library was first constructed by

RepeatModeler and then used by RepeatMasker to search TEs. Protein-coding genes were

annotated by integrating predictions of *ab initio* gene annotation tools and alignments of

homologous proteins. Three different tools including Augustus v3.2.3 (Stanke & Waack, 2003),

GlimmerHMM v3.0 (Majoros, Pertea, & Salzberg, 2004) and SNAP v2013 (Korf, 2004) were used

to predict the initial gene models. Protein sequences from *A. thaliana*, *A. lyrata* and *A. alpina*

(Arabidopsis Genome Initiative, 2000; Hu et al., 2011; Willing et al., 2015) were aligned to the

assembly by the tool Exonerate v2.2.0 (Slater & Birney, 2005). Then, the *ab initio* predictions

and protein alignment hits were further combined to build the consensus gene models by the

tool EVidenceModeler (EVM) v2012 (Haas et al., 2008). Finally, TE related genes in these models

were annotated by checking the TE annotation, blastp (Altschul, Gish, Miller, Myers, & Lipman,

1990) alignments with Plant TE related proteins and blastp alignments with *A. thaliana* proteins.

If a gene's protein sequence had blastp alignment identity and coverage both larger than 50%

with a TE related protein, or at least 30% of its exon regions overlapped with TEs but had no

good blastp hit (identity >50% and coverage >50%) from *A. thaliana* protein sequences, this

gene would be marked as a TE related gene.

RAD-sequencing and SNP calling

We genotyped 134 samples using the original RAD-sequencing (RADseq) protocol (Etter et al.,

2011), with the following modifications (Document S1): i) we used the enzyme KpnI-HF (New

England Biolabs) for DNA digestion, ii) we ligated digested DNA with complementary adapters

containing one of ten different barcodes and a stretch of five random nucleotides, used for

post-hoc removal of PCR duplicates (Table S2), iii) we created 14 pools of 10 barcoded samples

each in equal amounts, iv) we used indexed reverse primers for amplification, described in

(Peterson, Weber, Kay, Fisher, & Hoekstra, 2012), to allow multiplexing of pools. Libraries were

sequenced on two Illumina HiSeq 4000 lanes with 2x150bp.

We used *FastQC* (Andrews, 2010) to quality-check the resulting reads. We trimmed adapters

and removed reads shorter than 100bp using *Cutadapt* (Martin, 2011). We removed PCR

duplicates based on a 5 bp stretch of random nucleotides at the end of the adapter, using the

clone_filter module of *Stacks* version 1.37 (Catchen, Hohenlohe, Bassham, Amores, & Cresko,

2013). We de-multiplexed samples using the *process_radtags* module from *Stacks*. We filtered

reads with ambiguous barcodes (allowed distance 2) and cut-sites, reads with uncalled bases

and low-quality reads (default threshold).

For reference-based genotyping, we mapped reads using *BWA* (Li & Durbin, 2009) with default

settings. We filtered mapped reads using *SAMtools* (Li et al., 2009) and custom python scripts

using the following criteria to remove reads: mapping quality < 30, number soft-clipped bases >

30; reads were unpaired; mates mapped on different chromosomes; mate mapping distance >

700. We called genotypes using *SAMtools mpileup* and *VarScan2* (Koboldt et al., 2012) with the

following options: base quality > 20; re-calculation of base quality on the fly (-E option); read

depth > 14; strand filter de-activated; SNP calling p-value < 0.01. We filtered genotyped loci

using *VCFtools* (Danecek et al., 2011) and custom python scripts removing loci with missing data

in more than 5% of individuals and loci in masked (repetitive) regions. We clustered contiguous

loci spaced less than 100 bp apart into RAD-regions. Regions with excessively low or high

coverage are likely results of allele dropout or paralogous mapping, respectively. Thus, we

removed regions fulfilling one of the following criteria: mean coverage of the region greater

than twice the overall mean coverage; mean coverage of the region smaller than a third of the

overall mean coverage; region maximum coverage greater than twice the mean maximal

coverage over all regions; region shorter than 250bp; region longer than 1300bp.

After depth filtering we still found loci with a high frequency of heterozygotes (up to 100%),

which is unlikely in highly selfing species. Notably, in over 90% of these loci only one of the two

homozygous genotypes was observed. Moreover, these highly heterozygous loci clustered in

high density on single RAD-fragments. Thus, these loci likely resulted from paralogous mapping,

which passed the depth filter due to sequencing depth variation among RAD-regions. Therefore,

we removed RAD-regions containing loci with a frequency of heterozygotes greater than 20%.

This threshold was picked to limit the impact of noise from mapping artifacts while still allowing

for reasonable levels of heterozygosity, since some low- level outcrossing likely occurs. From the

resulting genotype dataset, we extracted single nucleotide polymorphisms (SNPs) using

VCFtools (Danecek et al., 2011).

For *de-novo*-genotyping we used the *Stacks* 2.2 *denovo_map.pl* pipeline (Catchen et al., 2013).

The aim of the de-novo analysis was to test whether one would reach similar results and

conclusions without the use of a reference genome. Thus, we performed this analysis without

knowledge about the two species (since we gained this from the reference-based analysis) and

ran *Stacks* for all samples combined. As recommended by the authors of the tool (Rochette &

Catchen, 2017), we first used a subset of 15 representative genotypes to tune the parameters (-

M and -n) of the algorithm, which control the number of mismatches between stacks within (M)

and between (n) individuals. We varied M and n from 1 to 9. For each set of parameters, we

analysed the number of loci shared between 80% of the samples. This measure peaked at the

value six for M and n. Thus, we used this value for both parameters for the full analysis. We ran

the *denovo_map.pl* pipeline using 0.01 as the p-value threshold for calling genotypes and SNPs,

and otherwise default options. We used the *populations* program to create a VCF-file for further

analysis using the following filters: 5% maximum missing data per locus; 20% maximum

observed heterozygosity per locus; locus must be present in all sites.

Population genetics statistics

We did all statistical analysis using R version 3.4.4 (R Development Core Team, 2008) and

provide a supplemental R Markdown file (Document S2). The following packages were used for

plotting: *ggplot2* (Wickham, 2009), *ggmap* (Kahle & Wickham, 2013), *ggthemes* (Arnold et al.,

2017), *ggsn* (Baquero, 2017) and *heatmap3* (Zhao, Guo, Sheng, & Shyr, 2015). We performed all

analysis for the reference-based and the *de-novo*-based dataset and compared the results. We

used the *vcfR* package (Knaus & Grünwald, 2017) to load VCF-files into R and make the SNP data

available for processing with other libraries. Based on our annotation we determined whether

SNPs are in coding regions and whether they are synonymous or non-synonymous using the

PopGenome package (Pfeifer, Wittelsbuerger, Li, & Handsaker, 2018). We performed principal

component analysis (PCA) of SNP data for all samples using the *adegenet* package (Jombart et

al., 2016). Missing data was scaled to the mean for PCA.

Based on the first principal component most individuals (83%) could be assigned to two distinct

taxonomic groups. We used molecular methods to assign species labels to the two taxonomic

groups. We sequenced the internal transcribed spacer (ITS) sequence of nine individuals, three

from the first (left) and five from the second (right) cluster and one located between the

clusters (Table S3). Primers for amplification were taken from (Mummenhoff, Franzke, & Koch,

1997). We used the sequences as input to the taxonomy tool of the *Brassibase* website (Kiefer

et al., 2014). Interpretation of the output required ploidy information. Leaf samples were

therefore collected from twelve individuals (seven putative *A. sagittata*, two putative *A.*

nemorensis, three putative hybrids; Table S4) and their ploidy determined by Plant Cytometry

Services (Didam, Netherlands).

We conducted all population genetic analysis separately for the two species. Individuals, which

were not assigned to any species, were likely interspecific hybrids and excluded from further

analysis. We used the *pegas* library to calculate within-site genetic diversity (Nei's π; average

pairwise nucleotide differences) for each site (Paradis, Jombart, Schliep, Potts, & Winter, 2016),

excluding sites with less than two individuals per respective species. To scale the estimates of π,

we divided the average number of pairwise nucleotide differences among samples by the total

number of successfully genotyped bases, excluding all bases which failed any of our previously

described genotype filters (missing data, region heterozygosity, region depth and length). For

the de-novo pipeline we extracted the total number of genotyped bases from *Stacks* output. We

calculated correlation coefficients between reference-based and *de-novo*-based π estimates

using Pearson's method. We calculated pairwise F_{ST} (Nei, 1987) and genetic distance (Cavalli-

Sforza & Edwards, 1967) between all pairs of sites using the *hierfstat* package (Goudet &

Jombart, 2015). Negative F_{ST} values were set to zero. Differences of genetic distance and F_{ST}

among pristine and restored sites were tested using a Wilcoxon-rank-sum-test on pairwise

distance matrices. We tested for correlation between the distance matrices of the reference

and *de novo* datasets using a Mantel test with 10000 permutations.

Admixture analysis

For ADMIXTURE analysis (Alexander, Novembre, & Lange, 2009), we converted VCF-files to bed-

files using PLINK (Purcell, 2009; Purcell et al., 2007). First, we conducted ADMIXTURE analysis

for all samples combined for K=1 to K=10 (reference-pipeline only). Then for each of the species

and pipeline (reference/*de novo*), we ran ADMIXTURE analysis for K=2 to K=10, with 10

iterations of cross-validation each. Before plotting, we normalized clusters across runs using

CLUMPAK (Kopelman, Mayzel, Jakobsson, Rosenberg, & Mayrose, 2015). We created plots using

a custom R-script.

Results

RAD-sequencing uncovers two hybridizing species

Unless otherwise stated, all described results were obtained using the reference-based pipeline.

We genotyped 134 individuals from 10 sites – 4 pristine and 6 restored (Figure 1) – yielding 3.6

Mb of sequence of which 32,880 single nucleotide positions were polymorphic (SNPs). Only 20%

of SNPs were in coding regions, 40% and 56% of which were synonymous and non-synonymous,

respectively (4% unassigned). To visualize patterns of genetic diversity across sites, we

conducted principal component analysis (PCA) for all individuals (Figure 2, top). Almost 80% of

the total genetic variation was explained by the first principal component, which separated

most individuals into two clearly defined clusters, likely representing taxonomic units. Based on

phylogenetic analysis of the ITS region of 10 individuals (Table S3), we identified one taxonomic

unit as *Arabis nemorensis* (Figure S2). The ITS region of individuals from the other taxonomic

unit was highly similar to *A. sagittata* and *Arabis hirsuta* (Figure S3). Both are sibling species

from the *A. hirsuta* complex, but *A. hirsuta* is tetraploid and *A. sagittata* diploid. Since all twelve

tested samples were diploid (Table S4), we conclude that the second cluster most likely

corresponds to *A. sagittata*. Average genetic distance (d_{XY}) between *A. nemorensis* and *A.*

sagittata was 8.1e-03, i.e. 8 fixed SNP differences for 1000 bp.

Twenty-three individuals showed a positioning along the first PC that was intermediate between

the two clusters, suggesting they were interspecific hybrids. Most of these hybrids were closer

to *A. sagittata* on the first PC. Since sterile F1-hybrids should be located exactly in the middle

between the two species, this suggests that they are somewhat fertile and preferentially back-

cross with *A. sagittata.* Laboratory observations have shown that hybrids are indeed fertile

(Dittberner, personal observation). Additionally, in an ADMIXTURE analysis of all samples with

K=2, most hybrids showed ancestry over 50% from *A. sagittata*, as expected from preferential

back-crossing to this parent (Figure S4).

Overall our sample was composed of 31% *A. nemorensis*, 52% *A. sagittata* and 17% hybrids. The

species composition differed among sites (Figure 2, bottom). *A. nemorensis* was present in 7

sites (3 pristine, 4 restored), *A. sagittata* in 9 sites (3 pristine, 6 restored) and hybrids in 6 sites

(3 pristine, 3 restored). Notably, the pristine site D was dominated by hybrids with over 56%.

Figure 2: Presence of two hybridizing *Arabis* species in study sites.
Top) Plot of the first two principal components (PC) of genetic variation based on the reference-based
pipeline. Text labels and colors distinguish the different sites. Text labels are approximately in the
centroid of the respective site (with offset to avoid overlaps). Values in brackets in the axis labels show
the amount of variance explained by each PC. Note that the first PC explains about 80% of the total
variation and splits most individuals into two clearly defined clusters. The clusters on the left and on the
right end of PC1 correspond to two distinct species, identified here as *Arabis nemorensis* and *Arabis*
sagittata. Species identity is indicated by text labels below the plot. Points in between the two clusters
are likely natural interspecific hybrids. The dashed lines show the thresholds used for distinction
between species and hybrids throughout this study. Middle) Plot showing the same analysis as above but
based on the de-novo-pipeline. Bottom) Stacked barplot showing the distribution of species among the

371   study sites. Population labels are colored by type: darkgreen = pristine, lightgreen = restored. Species are
372   distributed heterogeneously among sites.

No reduction of genetic diversity in restored sites

We computed species-specific estimates of genetic diversity within each site, excluding hybrid

genotypes. The *A. nemorensis* dataset consisted of 2746 SNPs. Levels of genetic diversity (π)

varied up to two-fold among sites, ranging from 6.6e-05 in A-2 to 1.4e-04 in A-1 (Figure 3**Fehler!**

Verweisquelle konnte nicht gefunden werden., left; Table S5). Total diversity was 1.5e-04.

However, pristine and restored sites did not differ significantly in their level of diversity (mean

difference= +10% in restored; W = 4, p = 1).

The *A. sagittata* dataset consisted of 6366 SNPs. Total genetic diversity in *A. sagittata* was about

three times as high as in *A. nemorensis*. Yet, in contrast to *A. nemorensis*, genetic diversity

differed strongly among sites, ranging from 1.03e-06 in site I to 4.26e-04 in site E (Figure

3**Fehler! Verweisquelle konnte nicht gefunden werden.**, right; Table S5). Notably, genetic

diversity was low in two of three pristine sites. In contrast, we found high levels of diversity in all

restored sites, except site F. However, the overall difference between pristine and restored sites

was not significant (mean difference= +163% in restored sites; W=13, p=0.14).

Since hybridization potentially enables gene flow between the two species we also compared

levels of genetic diversity for both species combined, including the hybrids. Overall genetic

diversity increased by an order of magnitude and mixed sites were more diverse than mostly

pure sites, as would be expected (Figure S5). Again, restored sites did not show significantly

different levels of genetic diversity from pristine sites (mean difference= +22%; W=13, p=0.91).

Figure 3: No reduction in genetic diversity in restored sites
Barplot of average pairwise genetic diversity (π) within each site of each species. Bar-color indicates the
type of site.

Restoration reduces population structure and facilitates recombination

To quantify the degree of population structure, we estimated genetic distance and

differentiation (F_{ST}) among all pairs of sites (Table S6). Genetic distance among *A. nemorensis*

sites ranged from 0.03 to 0.31 and F_{ST} estimates from 0 to 0.5 (Figure 4, A+C). Population

structure was slightly more pronounced among pristine sites than restored sites: Mean genetic

distance was 0.26 among pristine sites and 0.13 among restored sites (W=17, p=0.047); mean

F_{ST} was 0.37 among pristine sites and 0.25 among restored sites (W=14, p=0.26).

In *A. sagittata*, genetic distance ranged from 0.32 to 0.34 and F_{ST} estimates from 0 to 0.91

(Figure 4, B+D). The difference in population structure between pristine and restored sites was

stronger than in *A. nemorensis*: mean genetic distance was 0.33 among pristine sites and 0.12

among restored sites (W=45, p=0.002); mean F_{ST} was 0.81 among pristine sites and 0.2 among

restored sites (W=43, p=0.015).

Figure 4: Overall strong population structure with signatures of admixture in restored sites
This panel shows different measurements of population structure of each species. A-D) Heatmaps
showing pairwise comparisons between all sites of genetic distance (Cavalli-Sforza chord distance) and
FST, respectively. Values of the respective variable are indicated by tile color, corresponding to the color
scale in the top-left corner of each heatmap. Sites were clustered based on the respective variable, as
indicated by the dendrograms. Colors of text labels and bars next to the grid indicate the type of the site
(same color scheme as Figure 3).

Reduced differentiation among restored sites suggests that genetic material of pristine sites was

mixed in restored sites. To visualize this in more detail, we conducted ADMIXTURE analysis for

each species. This analysis assumes a given number (K) of genetic clusters (ancestral

populations) and assigns cluster ancestry proportions to each individual, allowing for mixed

ancestry. For *A. sagittata*, K=6 was estimated as the optimal value (Figure 5). Up to K=4, the

three pristine sites consisted of three distinct genetic clusters, which were mixed in all but one

of the restored sites. For higher values of K, pristine population B consisted in two clusters, with

pure and admixed individuals in equal proportions, in agreement with the higher genetic

diversity observed at this site. All individuals from other pristine sites had pure ancestry from a

single cluster per population.

For *A. nemorensis*, K=7 was estimated as the optimal value for clustering (Figure 5). In contrast

to *A. sagittata*, we found individuals with different or mixed ancestry within populations of *A.*

nemorensis, even at low values of K, indicating more genetic mixture than in *A. sagittata*. Since

genetic structure was less pronounced for *A. nemorensis*, the restoration procedure did not

impact the genetic distribution of this species, by contrast with *A. sagittata*.

Figure 5: Increased impact of genetic admixture of pristine sites in *A. sagittata*
The figures shows a comparison of ADMIXTURE results for both species. The top panels show the cross-validation (CV) error for all tested values of K. The following panels show the ancestry proportions of each cluster for each individual for all values of K until the first minimum of the CV-error curve. The numbers on the right represent the value of K. Site labels are colored according to the type of the site: light-green = restored, dark-green = pristine.

De-novo- and reference-based summary statistics reach the same conclusion

Finally, we tested whether a reference genome is required to determine the impact of

restoration on genetic diversity by comparing results from a reference-based and a *de novo*

pipeline. We found that estimates of genetic diversity, genetic distance or F_{ST} yielded by the two

methods were highly correlated, with all correlation coefficients being greater than 0.95

(maximum p < 0.001, Figure 6). However, we observed that estimates of genetic diversity

generated without a reference genome were deflated, especially for high diversity sites (Figure

6), by a median factor of 0.83 for *A. nemorensis* and 0.36 for *A. sagittata*. Estimates of genetic

distance were underestimated by a median factor of 0.61 in both species. Interestingly,

however, both pipelines yielded almost identical estimates of F_{ST} (median factor of 0.93) and

revealed a very similar extent of admixture between sites (Figures S6-7). Moreover, the

presence of the two species and their hybrids was detected with both methods (Figure 2

middle). We concluded that, in this study, the use of a reference genome was not required to

determine the impact of restoration on genetic diversity. Since the two pipelines coincide, we

could further conclude that the distribution of variation in two species and across pristine and

restored sites reported above is not the result of possible mapping biases to the reference

genome.

Figure 6: De-novo- and reference-based summary statistics are highly correlated.
Plots show the correlation of summary statistics based on the reference-based and the de-novo-based
pipeline. Line and dot colors represent the species. Each dot represents one individual. Lines represent a
linear fit through the dots and the gray shadows indicate the error of the fit. The dashed line represents
a hypothetical 1:1 relationship of the variables. Correlation-coefficients (Pearson's product-moment
correlation) are indicated as colored text corresponding to the species. Stars indicate the level of the
coefficient: **p<0.01, ***p<0.001.

Discussion

A reference-genome is not required to characterize the impact of restoration

While RAD-seq and related methods are a cheap tool to acquire genotype information across

the genome without need for a reference genome (Elshire et al., 2011; Etter et al., 2011;

Peterson et al., 2012), the reliability of *de novo* assembly pipelines has been questioned (Shafer

et al., 2016). The availability of a reference genome allowed us to ask whether reference-based

read mapping pipelines yielded distinct conclusions from a pipeline based on *de novo* read

assembly. Conversely, this also allowed verifying that the use of an *A. nemorensis* reference

genome to map A. sagittata samples did not bias the outcome of our study. The results from

both pipelines were highly correlated. Thus, comparative analysis of sites was reliable with both

pipelines. However, the *de-novo*-pipeline (*Stacks*, Catchen et al., 2013) underestimated the

amount of genetic diversity compared to the reference-based pipeline. This is in contrast to a

previous study comparing different RAD-seq pipelines (Shafer et al., 2016), where *Stacks*

produced slightly inflated estimates compared to reference-based pipelines. However, in the

same study, other de novo pipelines produced substantially lower estimates of genetic diversity

(Shafer et al., 2016). Thus, the magnitude of genetic diversity estimates might depend on the

study system and pipelines/parameters used, and caution is advised when comparing these

estimates across studies.

In contrast, both pipelines agreed for analyses comparing diversity between species or sites (e.g.

F_{ST}, ADMIXTURE). Thus, we conclude that RAD-seq is an efficient tool to characterize the

distribution of diversity, even in the absence of a reference genome. We therefore hope that

this study will pave the way for exploring how species, with diverse life-history and

uncharacterized genomes, will be maintained after hay transfer or how modalities of hay

transfer affects not only single species but the balance between multiple species in the

community.

First documented presence of *A. sagittata* in floodplain meadow habitat

We detected two independent genetic clusters. Based on ITS sequences, ecological preference

and morphological descriptors, the first one appears to represent *bona fide A. nemorensis*

individuals. Taxonomic boundaries in the *Arabis hirsuta* clade can be tenuous, yet ITS sequence

differences between the two clusters strongly suggests that they belong to distinct species (Karl

& Koch, 2014). ITS sequences, ploidy level and morphological evidence collectively indicated

that individuals in the second genetic cluster are likely to belong to *Arabis sagittata*. Species

barriers are often incomplete, and patterns of divergence between closely related species of the

same genus can be strongly intertwined. Population genomics approaches and wider sampling

will be needed to consolidate our understanding of the *A. sagittata* taxonomic group and its

historical relationship with the *A. nemorensis* taxon. Population genomics can indeed yield

detailed view on the process and history of speciation in plant genera (see e.g. Novikova et al.,

2016).

We did not anticipate the presence of the *A. sagittata* in flood-plain habitats because it is

predominantly found in warm and dry habitats (Hand & Gregor, 2006). However, agricultural

land-use and flood regulation have considerably modified the flood-plain ecosystem: flooding is

contained by a dyke and the ground water level has decreased (Hölzel & Otte, 2001). Man-made

modifications of the environment can change selection regimes and facilitate the establishment

of non-native species (Byers, 2002; Crooks, Chang, & Ruiz, 2011; Fukasawa, Miyashita,

Hashimoto, Tatara, & Abe, 2013; Tyrrell & Byers, 2007). Thus, it is possible that the migration of

A. sagittata into the flood-plain ecosystem was facilitated by the decreased frequency and

severity of flooding caused by human river regulation. We want to stress that the species co-

occurred both in pristine and restored sites. Thus, the contact between the species was not

caused by the restoration efforts. In fact, we observed that *A. sagittata* is more frequent than *A.*

nemorensis in our sample, which raises the concern that this species may be in the process of

displacing *A. nemorensis*. This trend could be enhanced by climate change, which may lead to

conditions that further favor the xero-thermophylic *A. sagittata*. Additionally, if *A. sagittata* is

often mistaken for *A. nemorensis* in flora reports on flood-plain environments, the remaining *A.*

nemorensis population in Central Europe might be even smaller than is currently assumed.

The parallel transfer of two sympatric relatives shows that hay transfer maintains genetic

diversity

Hay-transfer has proven to be a particularly successful method for establishing new populations

of target species in ecological restoration across a variety of herbaceous vegetation types

(Coiffait-Gombault, Buisson, & Dutoit, 2011; Hölzel & Otte, 2003; Kiehl et al., 2010; Kiehl &

Wagner, 2006; Török et al., 2012). Furthermore, hay-transfer is often seen as the gold standard

to preserve local levels of genetic diversity and adaptation (Vander Mijnsbrugge, Bischoff, &

Smith, 2010). The latter is probably the main reason, why it is increasingly used in in ecological

restoration (Kiehl et al., 2014). The aim of this study was to characterize the level of diversity in

the pristine source sites and document the impact of hay-transfer on the genetic diversity in

restored sites. Although our initial plan was to focus on *A. nemorensis*, a typical representative

of species-rich floodplain meadows, the unanticipated presence of *A. sagittata* in our sample

allowed us to compare the genetic effects of restoration by hay-transfer on the two species.

Several years after restoration, we did not find a significant difference in genetic diversity

between pristine and restored sites for either of the two species. Thus, the hay-transfer method

can restore populations with levels of diversity indistinguishable from the source populations in

the long-term. Our findings are in agreement with studies on population life-stage structure and

dynamics comparing pristine and restored sites of *A. nemorensis/A. sagittata* in the same region

(Burmeier et al., 2011). This outcome is particularly remarkable given that *A. nemorensis* has a

long-term seed bank, with up to 25,000 germinable seeds*m^{-2}, which was not transferred to

restored sites (Burmeier et al., 2011). Although species for which the genetic diversity present in

the seed bank tends to differ more strongly from the above ground diversity may fare

differently after hay-transfer, we note that populations restored with alternative methods, e.g.

spontaneous recolonization (Vandepitte et al., 2012) or propagated seed mixtures (Espeland et

al., 2017; Fant, Holmstrom, Sirkin, Etterson, & Masi, 2008), both excluded the seed bank and

revealed a reduction of diversity (Mijangos et al., 2015). Thus, hay-transfer might be superior to

other restoration methods not only in restoration success (Hölzel & Otte, 2003; Kiehl et al.,

2010) but also in transferring genetic diversity.

The modalities of habitat restoration can modify the relative adaptive potential of species

in the ecosystem

Genetic diversity was low in *A. nemorensis* and *A. sagittata,* compared to other outcrossing or

selfing Brassicaceae species (Mattila, Tyrmi, Pyhäjärvi, & Savolainen, 2017; Onge, Källman,

Slotte, Lascoux, & Palmé, 2011). In fact, genetic distance between species was only on the order

of magnitude as diversity found within populations of *Arabidopsis lyrata* (Mattila et al., 2017).

Genetic diversity within our species was similar to that reported for *Arabis alpina* populations in

Scandinavia (Laenen et al., 2018). As in our study, these populations are selfing and located on

the margin of the species' range (Jalas & Suominen, 1994), two factors often coinciding with

lower levels of genetic diversity. Since populations with low genetic diversity may suffer from

increased genetic load and decreased adaptive potential, the transfer of non-local material is

often envisaged to preserve endangered species (Breed, Stead, Ottewell, Gardner, & Lowe,

2013; Weeks et al., 2011). This approach is however controversial. Strategies that introduce

non-local seeds can decrease population fitness either by introducing maladapted genotypes

(Crémieux, Bischoff, Müller-Schärer, & Steinger, 2010; McKay, Christian, Harrison, & Rice, 2005)

or by causing outbreeding depression (Frankham et al., 2011). As a compromise, a strategy of

mixing regional seeds was recently proposed (Bucharova et al., 2018). Our results show that this

strategy was incidentally implemented in the examined restoration effort: ADMXITURE analysis

showed that pristine *A. sagittata* sites are dominated by one or two ancestral groups, with very

low genetic diversity within each group. Yet, in some of the restored sites, admixture of low

diversity *A. sagittata* groups took place, leading to a strong increase in genetic diversity and

decrease in population structure. ADMIXTURE analysis also revealed that this led to genetic

recombination in *A. sagittata,* possibly increasing the adaptive potential of restored sites in this

species. Increased genetic connectivity between populations has indeed been shown to help

maintain or even increase genetic diversity in the long term (DiLeo, Rico, Boehmer, & Wagner,

2017). Fitness assays of recombined *A. sagittata* genotypes are needed to verify whether this

local admixture has reinforced the establishment of this species in restored floodplain

meadows.

Interestingly, we find less pronounced population structure in pristine sites of *A. nemorensis*

than in *A. sagittata*, suggesting that a higher level of gene flow helps maintain diversity within

pristine sites. The restoration of this species will therefore not directly benefit from post-

transfer admixture. More so, our results suggest that it is possible that *A. nemorensis* is at

increased disadvantage in restored habitats, if admixture and resulting recombination were to

favor the emergence of more competitive genotypes were to evolve in *A. sagittata* thanks to

admixture.

The coexistence of these species in the Rhine floodplain ecosystem will be further impacted by

the ongoing hybridization dynamic that our analysis uncovers for the first time in this system

and which occurred independently of the restoration effort. *A. nemorensis* could receive alleles

conferring drought-adaptation from the xero-thermophilic *A. sagittata* that may enhance its

ability to cope with increased drought exposure in its habitat. The genomic composition of the

hybrids, however, also indicates that hybrids back-cross preferentially with *A. sagittata*. Gene-

flow from *A. nemorensis* could facilitate adaptation of *A. sagittata* to the flood-plain

environment. Such adaptive introgressions could also potentially accelerate the extinction of *A.*

nemorensis. While hybridization is common in plants (Mallet, Besansky, & Hahn, 2016), well

documented cases of adaptive introgression are rare and require elaborate experiments

(Goulet, Roda, & Hopkins, 2017; Suarez-Gonzalez, Lexer, & Cronk, 2018). Such experiments are

now warranted to determine the impact of hybridization in this sympatric species complex.

Conclusions

Clearly, genetic analysis helps with species identification as sibling species can be difficult to

distinguish morphologically even for specialists. A unique feature of hay-transfer restoration

approaches is that the whole plant community can be transplanted. It is therefore particularly

important to determine the composition of the source populations to limit the spread of non-

target species (Bickford et al., 2007). Our study demonstrates that hay-transfer has maintained

genetic diversity in restored populations. However, we also note that it may have inadvertently

contributed to increase the genetic diversity and adaptive potential of only one of the two

species, due to differences in the genetic makeup of the donor populations. This might lead to a

competitive advantage of one over the other species in the long term, potentially disturbing the

balance in the community. On the one hand this shows that restoration by hay-transfer can

indeed enhance the adaptive potential, which is especially important in the face of a rapid

climate change. On the other hand, this also highlights that understanding the underlying

genetics of the community to be transferred is a pre-requisite for the design of restoration

strategies that promote the maintenance of both endangered ecosystems and endangered

species.

Acknowledgements

We thank Matthias Harnisch for providing information about the populations, Markus Koch for

insightful discussions about the *A. hirsut*a tribe, Eric Schranz for his assistance with the pseudo-

chromosome assembly and Gregor Schmitz for helpful feedback. Further, we thank Janine

Altmüller and the team of the Cologne Center for Genomics for their assistance in RAD-seq

optimization, library preparation and sequencing. This work was partly funded by the German

Research Foundation 'Deutsche Forschungsgemeinschaft DFG [DFG priority program 1529

'ADAPTOMICS'].

References

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in
unrelated individuals. *Genome Research*. https://doi.org/10.1101/gr.094052.109

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment
search tool. *Journal of Molecular Biology*, *215*(3), 403–410.
https://doi.org/10.1016/S0022-2836(05)80360-2

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data (Version
0.11.3). Retrieved from http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Arabidopsis Genome Initiative. (2000). Analysis of the genome sequence of the flowering plant
Arabidopsis thaliana. *Nature*, *408*(6814), 796–815. https://doi.org/10.1038/35048692

Arnold, J. B., Daroczi, G., Werth, B., Weitzner, B., Kunst, J., Auguie, B., … London, J. (2017).
ggthemes: Extra Themes, Scales and Geoms for "ggplot2" (Version 3.4.0). Retrieved from
https://cran.r-project.org/web/packages/ggthemes/index.html

Baillie, J., Hilton-Taylor, C., & Stuart, S. N. (2004). *2004 IUCN Red List of threatened species : a*
global species assessment. Retrieved from https://portals.iucn.org/library/node/9830

Baquero, O. S. (2017). ggsn: North Symbols and Scale Bars for Maps Created with "ggplot2" or
"ggmap" (Version 0.4.0). Retrieved from https://CRAN.R-project.org/package=ggsn

Benayas, J. M. R., Newton, A. C., Diaz, A., & Bullock, J. M. (2009). Enhancement of Biodiversity
and Ecosystem Services by Ecological Restoration: A Meta-Analysis. *Science*, *325*(5944),
1121–1124. https://doi.org/10.1126/science.1172460

Bickford, D., Lohman, D. J., Sodhi, N. S., Ng, P. K. L., Meier, R., Winker, K., … Das, I. (2007).
Cryptic species as a window on diversity and conservation. *Trends in Ecology &*
Evolution, *22*(3), 148–155. https://doi.org/10.1016/j.tree.2006.11.004

Bradbury, I. R., Hamilton, L. C., Dempson, B., Robertson, M. J., Bourret, V., Bernatchez, L., &
Verspoor, E. (2015). Transatlantic secondary contact in Atlantic Salmon, comparing
microsatellites, a single nucleotide polymorphism array and restriction-site associated
DNA sequencing for the resolution of complex spatial structure. *Molecular Ecology*,
24(20), 5130–5144. https://doi.org/10.1111/mec.13395

Breed, M. F., Stead, M. G., Ottewell, K. M., Gardner, M. G., & Lowe, A. J. (2013). Which
provenance and where? Seed sourcing strategies for revegetation in a changing

environment. *Conservation Genetics*, *14*(1), 1–10. https://doi.org/10.1007/s10592-012-0425-z

Bucharova, A., Bossdorf, O., Hölzel, N., Kollmann, J., Prasse, R., & Durka, W. (2018). Mix and match: regional admixture provenancing strikes a balance among different seed-sourcing strategies for ecological restoration. *Conservation Genetics*. https://doi.org/10.1007/s10592-018-1067-6

Bucharova, A., Michalski, S., Hermann, J.-M., Heveling, K., Durka, W., Hölzel, N., … Bossdorf, O. (2017). Genetic differentiation and regional adaptation among seed origins used for grassland restoration: lessons from a multispecies transplant experiment. *Journal of Applied Ecology*, *54*(1), 127–136. https://doi.org/10.1111/1365-2664.12645

Bullock, J. M., Aronson, J., Newton, A. C., Pywell, R. F., & Rey-Benayas, J. M. (2011). Restoration of ecosystem services and biodiversity: conflicts and opportunities. *Trends in Ecology & Evolution*, *26*(10), 541–549. https://doi.org/10.1016/j.tree.2011.06.011

Burmeier, S., Eckstein, R. L., Donath, T. W., & Otte, A. (2011). Plant Pattern Development during Early Post-Restoration Succession in Grasslands—A Case Study of Arabis nemorensis. *Restoration Ecology*, *19*(5), 648–659. https://doi.org/10.1111/j.1526-100X.2010.00668.x

Byers, J. E. (2002). Impact of non-indigenous species on natives enhanced by anthropogenic alteration of selection regimes. *Oikos*, *97*(3), 449–458. https://doi.org/10.1034/j.1600-0706.2002.970316.x

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool set for population genomics. *Molecular Ecology*, *22*(11), 3124–3140. https://doi.org/10.1111/mec.12354

Cavalli-Sforza, L. L., & Edwards, A. W. F. (1967). Phylogenetic analysis. Models and estimation procedures. *American Journal of Human Genetics*, *19*(3 Pt 1), 233–257.

Coiffait-Gombault, C., Buisson, E., & Dutoit, T. (2011). Hay Transfer Promotes Establishment of Mediterranean Steppe Vegetation on Soil Disturbed by Pipeline Construction. *Restoration Ecology*, *19*(201), 214–222. https://doi.org/10.1111/j.1526-100X.2010.00706.x

Crémieux, L., Bischoff, A., Müller-Schärer, H., & Steinger, T. (2010). Gene flow from foreign provenances into local plant populations: Fitness consequences and implications for biodiversity restoration. *American Journal of Botany*, *97*(1), 94–100. https://doi.org/10.3732/ajb.0900103

Crooks, J. A., Chang, A. L., & Ruiz, G. M. (2011). Aquatic pollution increases the relative success of invasive species. *Biological Invasions*, *13*(1), 165–176. https://doi.org/10.1007/s10530-010-9799-3

Crutsinger, G. M., Collins, M. D., Fordyce, J. A., Gompert, Z., Nice, C. C., & Sanders, N. J. (2006). Plant Genotypic Diversity Predicts Community Structure and Governs an Ecosystem Process. *Science*, *313*(5789), 966–968. https://doi.org/10.1126/science.1128326

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

DiLeo, M. F., Rico, Y., Boehmer, H. J., & Wagner, H. H. (2017). An ecological connectivity network
maintains genetic diversity of a flagship wildflower, Pulsatilla vulgaris. *Biological*
Conservation, *212*, 12–21. https://doi.org/10.1016/j.biocon.2017.05.026

Donath, T. W., Bissels, S., Hölzel, N., & Otte, A. (2007). Large scale application of diaspore
transfer with plant material in restoration practice – Impact of seed and microsite
limitation. *Biological Conservation*, *138*(1), 224–234.
https://doi.org/10.1016/j.biocon.2007.04.020

Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., & Mitchell, S. E.
(2011). A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity
Species. *PLOS ONE*, *6*(5), e19379. https://doi.org/10.1371/journal.pone.0019379

Espeland, E. K., Emery, N. C., Mercer, K. L., Woolbright, S. A., Kettenring, K. M., Gepts, P., &
Etterson, J. R. (2017). Evolution of plant materials for ecological restoration: insights
from the applied and basic literature. *Journal of Applied Ecology*, *54*(1), 102–115.
https://doi.org/10.1111/1365-2664.12739

Etter, P. D., Bassham, S., Hohenlohe, P. A., Johnson, E. A., & Cresko, W. A. (2011). SNP discovery
and genotyping for evolutionary genetics using RAD sequencing. *Methods in Molecular*
Biology (Clifton, N.J.), *772*, 157–178. https://doi.org/10.1007/978-1-61779-228-1_9

Fant, J. B., Holmstrom, R. M., Sirkin, E., Etterson, J. R., & Masi, S. (2008). Genetic Structure of
Threatened Native Populations and Propagules Used for Restoration in a Clonal Species,
American Beachgrass (Ammophila breviligulata Fern.). *Restoration Ecology*, *16*(4), 594–
603. https://doi.org/10.1111/j.1526-100X.2007.00348.x

Frankham, R., Ballou, J. D., Eldridge, M. D. B., Lacy, R. C., Ralls, K., Dudash, M. R., & Fenster, C. B.
(2011). Predicting the probability of outbreeding depression. *Conservation Biology: The*
Journal of the Society for Conservation Biology, *25*(3), 465–475.
https://doi.org/10.1111/j.1523-1739.2011.01662.x

Fukasawa, K., Miyashita, T., Hashimoto, T., Tatara, M., & Abe, S. (2013). Differential population
responses of native and alien rodents to an invasive predator, habitat alteration and
plant masting. *Proceedings of the Royal Society of London B: Biological Sciences*,
280(1773), 20132075. https://doi.org/10.1098/rspb.2013.2075

Gao, S., Bertrand, D., Chia, B. K. H., & Nagarajan, N. (2016). OPERA-LG: efficient and exact
scaffolding of large, repeat-rich eukaryotic genomes with performance guarantees.
Genome Biology, *17*, 102. https://doi.org/10.1186/s13059-016-0951-y

Gnerre, S., MacCallum, I., Przybylski, D., Ribeiro, F. J., Burton, J. N., Walker, B. J., … Jaffe, D. B.
(2011). High-quality draft assemblies of mammalian genomes from massively parallel
sequence data. *Proceedings of the National Academy of Sciences*, *108*(4), 1513–1518.
https://doi.org/10.1073/pnas.1017351108

Goudet, J., & Jombart, T. (2015). hierfstat: Estimation and Tests of Hierarchical F-Statistics
(Version 0.04-22). Retrieved from https://cran.r-
project.org/web/packages/hierfstat/index.html

Goulet, B. E., Roda, F., & Hopkins, R. (2017). Hybridization in Plants: Old Ideas, New Techniques.
Plant Physiology, *173*(1), 65–78. https://doi.org/10.1104/pp.16.01340

Gruenthal, K. M., Witting, D. A., Ford, T., Neuman, M. J., Williams, J. P., Pondella, D. J., … Larson, W. A. (2014). Development and application of genomic tools to the restoration of green abalone in southern California. *Conservation Genetics*, *15*(1), 109–121. https://doi.org/10.1007/s10592-013-0524-5

Haas, B. J., Salzberg, S. L., Zhu, W., Pertea, M., Allen, J. E., Orvis, J., … Wortman, J. R. (2008). Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biology*, *9*(1), R7. https://doi.org/10.1186/gb-2008-9-1-r7

Hand, R., & Gregor, T. (2006). Die Verbreitung von Arabis sagittata In Deutschland. Ergebnisse einer Herbarstudie. *Kochia*, *1*, 21–31.

Haug-Baltzell, A., Stephens, S. A., Davey, S., Scheidegger, C. E., & Lyons, E. (2017). SynMap2 and SynMap3D: web-based whole-genome synteny browsers. *Bioinformatics*, *33*(14), 2197–2198. https://doi.org/10.1093/bioinformatics/btx144

Hölzel, N., & Otte, A. (2001). The impact of flooding regime on the soil seed bank of flood-meadows. *Journal of Vegetation Science*, *12*(2), 209–218. https://doi.org/10.2307/3236605

Hölzel, N., & Otte, A. (2003). Restoration of a species-rich flood meadow by topsoil removal and diaspore transfer with plant material. *Applied Vegetation Science*, *6*(2), 131–140. https://doi.org/10.1111/j.1654-109X.2003.tb00573.x

Hölzel, N., & Otte, A. (2004). Assessing soil seed bank persistence in flood-meadows: The search for reliable traits. *Journal of Vegetation Science*, *15*(1), 93–100. https://doi.org/10.1111/j.1654-1103.2004.tb02241.x

Hu, T. T., Pattyn, P., Bakker, E. G., Cao, J., Cheng, J.-F., Clark, R. M., … Guo, Y.-L. (2011). The *Arabidopsis lyrata* genome sequence and the basis of rapid genome size change. *Nature Genetics*, *43*(5), 476–481. https://doi.org/10.1038/ng.807

Hufford, K. M., & Mazer, S. J. (2003). Plant ecotypes: genetic differentiation in the age of ecological restoration. *Trends in Ecology & Evolution*, *18*(3), 147–155. https://doi.org/10.1016/S0169-5347(03)00002-8

Hughes, A. R., Inouye, B. D., Johnson, M. T. J., Underwood, N., & Vellend, M. (2008). Ecological consequences of genetic diversity. *Ecology Letters*, *11*(6), 609–623. https://doi.org/10.1111/j.1461-0248.2008.01179.x

Jalas, J., & Suominen, J. (1994). *Atlas Florae Europaeae. Distribution of Vascular Plants in Europe 10: Cruciferae (Sisymbrium to Aubrieta)* (Vol. 10). The Committee for Mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.

Jeffries, D. L., Copp, G. H., Handley, L. L., Olsén, K. H., Sayer, C. D., & Hänfling, B. (2016). Comparing RADseq and microsatellites to infer complex phylogeographic patterns, an empirical perspective in the Crucian carp, Carassius carassius, L. *Molecular Ecology*, *25*(13), 2997–3018. https://doi.org/10.1111/mec.13613

Jiao, W.-B., Accinelli, G. G., Hartwig, B., Kiefer, C., Baker, D., Severing, E., … Schneeberger, K. (2017). Improving and correcting the contiguity of long-read genome assemblies of three plant species using optical mapping and chromosome conformation capture data. *Genome Research*, gr.213652.116. https://doi.org/10.1101/gr.213652.116

Jombart, T., Kamvar, Z. N., Lustrik, R., Collins, C., Beugin, M.-P., Knaus, B., … Calboli, F. (2016).
adegenet: Exploratory Analysis of Genetic and Genomic Data (Version 2.0.1). Retrieved
from https://cran.r-project.org/web/packages/adegenet/index.html

Kahle, D., & Wickham, H. (2013). ggmap: Spatial Visualization with ggplot2. *The R Journal*, *5*(1),
144–161.

Karl, R., & Koch, M. A. (2014). Phylogenetic signatures of adaptation: The Arabis hirsuta species
aggregate (Brassicaceae) revisited. *Perspectives in Plant Ecology, Evolution and*
Systematics, *16*(5), 247–264. https://doi.org/10.1016/j.ppees.2014.06.001

Kiefer, M., Schmickl, R., German, D. A., Mandáková, T., Lysak, M. A., Al-Shehbaz, I. A., … Koch,
M. A. (2014). BrassiBase: introduction to a novel knowledge database on Brassicaceae
evolution. *Plant & Cell Physiology*, *55*(1), e3. https://doi.org/10.1093/pcp/pct158

Kiehl, K., Kirmer, A., Donath, T. W., Rasran, L., & Hölzel, N. (2010). Species introduction in
restoration projects - evaluation of different techniques for the establishment of semi-
natural grasslands in Central and Northwestern Europe. *Basic and Applied Ecology*, *11*(4),
285–299.

Kiehl, K., Kirmer, A., & Shaw, N. (2014). *Guidelines for Native Seed Production and Grassland*
Restoration. Cambridge Scholars Publishing.

Kiehl, K., & Wagner, C. (2006). Effect of Hay Transfer on Long-Term Establishment of Vegetation
and Grasshoppers on Former Arable Fields. *Restoration Ecology*, *14*(1), 157–166.
https://doi.org/10.1111/j.1526-100X.2006.00116.x

Knaus, B. J., & Grünwald, N. J. (2017). VCFR: a package to manipulate and visualize variant call
format data in R. *Molecular Ecology Resources*, *17*(1), 44–53.

Koboldt, D. C., Zhang, Q., Larson, D. E., Shen, D., McLellan, M. D., Lin, L., … Wilson, R. K. (2012).
VarScan 2: Somatic mutation and copy number alteration discovery in cancer by exome
sequencing. *Genome Research*, *22*(3), 568–576. https://doi.org/10.1101/gr.129684.111

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak: a
program for identifying clustering modes and packaging population structure inferences
across K. *Molecular Ecology Resources*, *15*(5), 1179–1191. https://doi.org/10.1111/1755-
0998.12387

Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics*, *5*, 59.
https://doi.org/10.1186/1471-2105-5-59

Laenen, B., Tedder, A., Nowak, M. D., Toräng, P., Wunder, J., Wötzel, S., … Slotte, T. (2018).
Demography and mating system shape the genome-wide impact of purifying selection in
Arabis alpina. *Proceedings of the National Academy of Sciences*, 201707492.
https://doi.org/10.1073/pnas.1707492115

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics*, *25*(14), 1754–1760.
https://doi.org/10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … 1000 Genome Project Data
Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools.
Bioinformatics (Oxford, England), *25*(16), 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352

Lyons, E., & Freeling, M. (2008). How to usefully compare homologous plant genes and chromosomes as DNA sequences. *The Plant Journal*, *53*(4), 661–673. https://doi.org/10.1111/j.1365-313X.2007.03326.x

Lyons, E., Freeling, M., Kustu, S., & Inwood, W. (2011). Using Genomic Sequencing for Classical Genetics in E. coli K12. *PLOS ONE*, *6*(2), e16717. https://doi.org/10.1371/journal.pone.0016717

Majoros, W. H., Pertea, M., & Salzberg, S. L. (2004). TigrScan and GlimmerHMM: two open source ab initio eukaryotic gene-finders. *Bioinformatics (Oxford, England)*, *20*(16), 2878–2879. https://doi.org/10.1093/bioinformatics/bth315

Mallet, J., Besansky, N., & Hahn, M. W. (2016). How reticulated are species? *BioEssays*, *38*(2), 140–149. https://doi.org/10.1002/bies.201500149

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal*, *17*(1), 10–12. https://doi.org/10.14806/ej.17.1.200

Massatti, R., Doherty, K. D., & Wood, T. E. (2018). Resolving neutral and deterministic contributions to genomic structure in <Emphasis Type="Italic">Syntrichia ruralis</Emphasis> (Bryophyta, Pottiaceae) informs propagule sourcing for dryland restoration. *Conservation Genetics*, *19*(1), 85–97. https://doi.org/10.1007/s10592-017-1026-7

Mattila, T. M., Tyrmi, J., Pyhäjärvi, T., & Savolainen, O. (2017). Genome-Wide Analysis of Colonization History and Concomitant Selection in Arabidopsis lyrata. *Molecular Biology and Evolution*, *34*(10), 2665–2677. https://doi.org/10.1093/molbev/msx193

McKay, J. K., Christian, C. E., Harrison, S., & Rice, K. J. (2005). "How Local Is Local?"—A Review of Practical and Conceptual Issues in the Genetics of Restoration. *Restoration Ecology*, *13*(3), 432–440. https://doi.org/10.1111/j.1526-100X.2005.00058.x

Mijangos, J. L., Pacioni, C., Spencer, P. B. S., & Craig, M. D. (2015). Contribution of genetics to ecological restoration. *Molecular Ecology*, *24*(1), 22–37. https://doi.org/10.1111/mec.12995

Mummenhoff, K., Franzke, A., & Koch, M. (1997). Molecular phylogenetics of Thlaspi s.l. (Brassicaceae) based on chloroplast DNA restriction site variation and sequences of the internal transcribed spacers of nuclear ribosomal DNA. *Canadian Journal of Botany*, *75*(3), 469–482. https://doi.org/10.1139/b97-051

Nei, M. (1987). *Molecular evolutionary genetics*. Columbia university press.

Novikova, P. Y., Hohmann, N., Nizhynska, V., Tsuchimatsu, T., Ali, J., Muir, G., … Nordborg, M. (2016). Sequencing of the genus *Arabidopsis* identifies a complex history of nonbifurcating speciation and abundant trans-specific polymorphism. *Nature Genetics*, *48*(9), 1077–1082. https://doi.org/10.1038/ng.3617

O'Leary, S. J., Hollenbeck, C. M., Vega, R. R., Gold, J. R., & Portnoy, D. S. (2018). Genetic mapping and comparative genomics to inform restoration enhancement and culture of southern flounder, Paralichthys lethostigma. *BMC Genomics*, *19*(1), 163. https://doi.org/10.1186/s12864-018-4541-0

Onge, K. R. S., Källman, T., Slotte, T., Lascoux, M., & Palmé, A. E. (2011). Contrasting demographic history and population structure in Capsella rubella and Capsella

grandiflora, two closely related species with different mating systems. *Molecular Ecology*, *20*(16), 3306–3320. https://doi.org/10.1111/j.1365-294X.2011.05189.x

Paradis, E., Jombart, T., Schliep, K., Potts, A., & Winter, D. (2016). pegas: Population and Evolutionary Genetics Analysis System (Version 0.9). Retrieved from https://cran.r-project.org/web/packages/pegas/index.html

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double Digest RADseq: An Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-Model Species. *PLoS ONE*, *7*(5), e37135. https://doi.org/10.1371/journal.pone.0037135

Pfeifer, B., Wittelsbuerger, U., Li, H., & Handsaker, B. (2018). PopGenome: An Efficient Swiss Army Knife for Population Genomic Analyses (Version 2.6.1). Retrieved from https://CRAN.R-project.org/package=PopGenome

Purcell, S. (2009). PLINK (Version 1.07).

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., … Sham, P. C. (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *American Journal of Human Genetics*, *81*(3), 559–575. https://doi.org/10.1086/519795

R Development Core Team. (2008). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from http://www.R-project.org

Reed, D. H., & Frankham, R. (2003). Correlation between Fitness and Genetic Diversity. *Conservation Biology*, *17*(1), 230–237. https://doi.org/10.1046/j.1523-1739.2003.01236.x

Reitzel, A. M., Herrera, S., Layden, M. J., Martindale, M. Q., & Shank, T. M. (2013). Going where traditional markers have not gone before: utility of and promise for RAD sequencing in marine invertebrate phylogeography and population genomics. *Molecular Ecology*, *22*(11), 2953–2970. https://doi.org/10.1111/mec.12228

Reusch, T. B. H., Ehlers, A., Hämmerli, A., & Worm, B. (2005). Ecosystem recovery after climatic extremes enhanced by genotypic diversity. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(8), 2826–2831. https://doi.org/10.1073/pnas.0500008102

Reynolds, L. K., McGlathery, K. J., & Waycott, M. (2012). Genetic Diversity Enhances Restoration Success by Augmenting Ecosystem Services. *PLOS ONE*, *7*(6), e38397. https://doi.org/10.1371/journal.pone.0038397

Roberts, L., Stone, R., & Sugden, A. (2009). The Rise of Restoration Ecology. *Science*, *325*(5940), 555–555. https://doi.org/10.1126/science.325_555

Rochette, N. C., & Catchen, J. M. (2017). Deriving genotypes from RAD-seq short-read data using Stacks. *Nature Protocols*, *12*(12), 2640–2659. https://doi.org/10.1038/nprot.2017.123

Scheepens, J. F., Rauschkolb, R., Ziegler, R., Schroth, V., & Bossdorf, O. (2017). Genotypic diversity and environmental variability affect the invasibility of experimental plant populations. *Oikos*, *127*(4), 570–578. https://doi.org/10.1111/oik.04818

Schnittler, M., & Günther, K.-F. (1999). Central European vascular plants requiring priority
conservation measures – an analysis from national Red Lists and distribution maps.
Biodiversity & Conservation, *8*(7), 891–925. https://doi.org/10.1023/A:1008828704456
Shafer, A. B. A., Peart, C. R., Tusso, S., Maayan, I., Brelsford, A., Wheat, C. W., & Wolf, J. B. W.
(2016). Bioinformatic processing of RAD-seq data dramatically impacts downstream
population genetic inference. *Methods in Ecology and Evolution*, *8*(8), 907–917.
https://doi.org/10.1111/2041-210X.12700
Slater, G. S. C., & Birney, E. (2005). Automated generation of heuristics for biological sequence
comparison. *BMC Bioinformatics*, *6*, 31. https://doi.org/10.1186/1471-2105-6-31
Smit, A., & Hubley, R. (2008). RepeatModeler Open-1.0 (Version 1.0). Retrieved from
http://www.repeatmasker.org
Smit, A., Hubley, R., & Green, P. (2013). *RepeatMasker Open-4.0*. Retrieved from
http://www.repeatmasker.org
Society for Ecological Restoration International Science & Policy Working Group. (2004). *The SER*
International Primer on Ecological Restoration. www.ser.org & Tucson: Society for
Ecological Restoration Internationa.
Stanke, M., & Waack, S. (2003). Gene prediction with a hidden Markov model and a new intron
submodel. *Bioinformatics (Oxford, England)*, *19 Suppl 2*, ii215-225.
Suarez-Gonzalez, A., Lexer, C., & Cronk, Q. C. B. (2018). Adaptive introgression: a plant
perspective. *Biology Letters*, *14*(3), 20170688. https://doi.org/10.1098/rsbl.2017.0688
Suding, K. N. (2011). Toward an Era of Restoration in Ecology: Successes, Failures, and
Opportunities Ahead. *Annual Review of Ecology, Evolution, and Systematics*, *42*(1), 465–
487. https://doi.org/10.1146/annurev-ecolsys-102710-145115
Tellier, A., Laurent, S. J. Y., Lainer, H., Pavlidis, P., & Stephan, W. (2011). Inference of seed bank
parameters in two wild tomato species using ecological and genetic data. *Proceedings of*
the National Academy of Sciences, *108*(41), 17052–17057.
https://doi.org/10.1073/pnas.1111266108
Török, P., Miglécz, T., Valkó, O., Kelemen, A., Tóth, K., Lengyel, S., & Tóthmérész, B. (2012). Fast
restoration of grassland vegetation by a combination of seed mixture sowing and low-
diversity hay transfer. *Ecological Engineering*, *44*, 133–138.
https://doi.org/10.1016/j.ecoleng.2012.03.010
Torres-Martinez, L., & Emery, N. C. (2016). Genome-wide SNP discovery in the annual herb,
Lasthenia fremontii (Asteraceae): genetic resources for the conservation and restoration
of a California vernal pool endemic. *Conservation Genetics Resources*, *8*(2), 145–158.
https://doi.org/10.1007/s12686-016-0524-0
Tyrrell, M. C., & Byers, J. E. (2007). Do artificial substrates favor nonindigenous fouling species
over native species? *Journal of Experimental Marine Biology and Ecology*, *342*(1), 54–60.
https://doi.org/10.1016/j.jembe.2006.10.014
Vandepitte, K., Gristina, A. S., Hert, K. D., Meekers, T., Roldán-Ruiz, I., & Honnay, O. (2012).
Recolonization after habitat restoration leads to decreased genetic variation in
populations of a terrestrial orchid. *Molecular Ecology*, *21*(17), 4206–4215.
https://doi.org/10.1111/j.1365-294X.2012.05698.x

Vander Mijnsbrugge, K., Bischoff, A., & Smith, B. (2010). A question of origin: Where and how to
collect seed for ecological restoration. *Basic and Applied Ecology*, *11*(4), 300–311.
https://doi.org/10.1016/j.baae.2009.09.002
Vrijenhoek, R. C. (1994). Genetic diversity and fitness in small populations. In V. Loeschcke, S. K.
Jain, & J. Tomiuk (Eds.), *Conservation Genetics* (pp. 37–53). Basel: Birkhäuser Basel.
https://doi.org/10.1007/978-3-0348-8510-2_5
Weeks, A. R., Sgro, C. M., Young, A. G., Frankham, R., Mitchell, N. J., Miller, K. A., … Hoffmann, A.
A. (2011). Assessing the benefits and risks of translocations in changing environments: a
genetic perspective. *Evolutionary Applications*, *4*(6), 709–725.
https://doi.org/10.1111/j.1752-4571.2011.00192.x
Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
Retrieved from http://ggplot2.org
Williams, A. V., Nevill, P. G., & Krauss, S. L. (2014). Next generation restoration genetics:
applications and opportunities. *Trends in Plant Science*, *19*(8), 529–537.
https://doi.org/10.1016/j.tplants.2014.03.011
Willing, E.-M., Rawat, V., Mandáková, T., Maumus, F., James, G. V., Nordström, K. J. V., …
Schneeberger, K. (2015). Genome expansion of Arabis alpina linked with
retrotransposition and reduced symmetric DNA methylation. *Nature Plants*, *1*(2), 14023.
https://doi.org/10.1038/nplants.2014.23
Zhao, S., Guo, Y., Sheng, Q., & Shyr, Y. (2015). heatmap3: An Improved Heatmap Package
(Version 1.1.1). Retrieved from https://cran.r-
project.org/web/packages/heatmap3/index.html

Data availability

968 The genome assembly, annotation, raw reads and RAD-seq samples will be uploaded to EMBL

969 ENA upon acceptance (available upon request for review). VCF files and custom script will be

970 uploaded to Dryad repository upon acceptance (available upon request for review). An R

971 Markdown file describing the population genetic analysis is provided in the supplemental

972 material.

Author contributions

974 JdM, HD, AT and NH conceived the study. HD and NH collected plant material; HD and CB

975 prepared material for sequencing; WBJ, KS and HD were responsible for bioinformatic

976   processing. JdM, HD and NH analyzed data and wrote the manuscript with significant

977   contributions from CB, WBJ, KS and AT.