Whole plant community transplants across climates reveal structural community

stability due to large shifts in species assemblage

Article type: original research

Authors: Sara Tomiolo[a,b*], Mark C. Bilton[a], Katja Tielbörger[a]

Affiliation:

a- Department of Ecology and Evolution, Plant Ecology Group, University of

Tübingen, Auf der Morgenstelle 5, 72076 Tübingen, Germany

b- Department of Bioscience, Aarhus University, Vejlsøvej 25, 8600 Silkeborg,

Denmark

Contacts:

*Corresponding author: sara.tomiolo@gmail.com; **Orcid ID:** orcid.org/0000-0002-

5071-5851

Mark C. Bilton: mcbilton.ecology@hotmail.com

Katja Tielbörger: katja.tielboerger@uni-tuebingen.de

Running headline: Community stability under climate change

Keywords: annual plant communities, climate change, Climatic Niche Groups,

community ecology, reciprocal transplants, determinants of plant community

diversity and assemblage, distribution range, community stability, field experiments

Summary

(1) Climate change will decrease precipitation and increase rainfall variability in

Eastern Mediterranean regions, with responses of plant communities largely

uncertain. Here, we tested short-term responses of dryland plant communities to

contrasting rainfall regimes using a novel experimental approach.

(2) We exposed three annual plant communities to sharp changes in climatic

conditions using whole community reciprocal transplants of soil and seed banks. We

tested for the role of climate *vs.* community origin on community response and

resistance. In parallel, we asked whether origin-specific climatic adaptations predict

compositional shifts across climates.

(3) For both community origins, the most dry-adapted species in each community

increased in dry climate and the wet-adapted species increased in wet climate. Dry

community origins showed large compositional shifts while maintaining stable plant

density, biomass and species richness across climates. Conversely, wet communities

showed smaller compositional shifts, but larger variation in biomass and richness.

(4) Asynchrony in species abundances in response to rainfall variability could

maintain structural community stability. This, in combination with seed dormancy,

has the ability to delay extinction in response to climate change. However, increasing

occurrence of extreme droughts may, in the long-term, lead to loss of wet-adapted

species.

Introduction

Understanding how climate may alter overall available plant resources (Sardans *et al.*, 2008; Garcia *et al.*, 2014) and impact upon community structure (Tilman & Downing, 1994; Gilman *et al.*, 2010) is a major challenge in current ecological research (Maestre *et al.*, 2012; Parmesan & Hanley, 2015). In cold regions for example, warming is likely to improve growing conditions and thus increase plant community biomass by increasing nutrient mobilization and expanding the length of the growing season (Garcia *et al.*, 2014). In contrast, decreased rainfall in drier regions will likely have negative impacts on primary productivity, community composition, and their corresponding ecosystem services (Sala & Lauenroth, 1982; Peñuelas *et al.*, 2007; Miranda *et al.*, 2011). These effects may be particularly strong in those dryland ecosystems for which climate predictions indicate increasing incidents of droughts (Cubasch *et al.*, 1996; Smiatek *et al.*, 2011). Here, droughts and increasing temperatures will increase evapotranspiration, shorten the growing season and limit access to nutrients, thereby decreasing total community biomass (Peñuelas *et al.*, 2007; Doblas−Miranda *et al.*, 2015; Harrison *et al.*, 2015). In extreme cases, this may lead to the collapse of entire ecological communities (Forey *et al.*, 2010).

One of the predicted impacts of climate change is a re-assembly of plant communities (Hobbs *et al.*, 2006; Williams & Jackson, 2007; Alexander *et al.*, 2016) due to the differential ability of single species to either track their climatic niche or to survive under changed conditions, by means of adaptation or plasticity (Fernandez-Going *et al.*, 2013; Shi *et al.*, 2015). Such community reshuffling may be expressed in a loss or gain of certain species, a shift in species relative abundance or both. In plant communities already exposed to large inter-annual variations in climate, immediate rearrangement of community assembly in response to climate extremes may be an inherent property of plant communities, and may promote community stability in the long-term. This effect is particularly pronounced when species numbers are large and population sizes vary asynchronously (Doak *et al.*, 1998; Schindler *et al.*, 2015). Indeed, ecological theory and models support the idea that high inter-annual variation in species response to climate can lead to community-level stability (Anderson *et al.*, 1982; Tilman *et al.*, 1998; Thompson *et al.*, 2015; Abbott *et al.*, 2017). This may be an important mechanism for maintaining dryland communities' stability in response to large year-to-year variation in rainfall, and for slowing down ongoing selection

processes due to climate change (Bonebrake & Mastrandrea, 2010; Bilton *et al.*,

2016).

Although long-term climate manipulations are the gold standard in ecological

climate impact research and are fundamental for understanding long-term community

shifts (Brown *et al.*, 2001; Rinnan *et al.*, 2007; Blume-Werry *et al.*, 2016), they are

very costly to set up and maintain, often outliving funding cycles and scientific

research positions (Lindenmayer *et al.*, 2012). The monitoring of communities for

short-term responses may be a useful complement to long-term experiments, as

besides being less costly, it can be vitally important for parsing mechanistic

information about plant responses to large inter-annual variation, as well as extreme

events (De Dato *et al.*, 2006; Barbosa *et al.*, 2014; Blume-Werry *et al.*, 2016).

Reciprocal transplants represent a promising approach for indirectly studying plant

responses to climate change on a short temporal scale. These manipulative

experiments have been widely adopted in single species (e.g. Link *et al.*, 2003; Casper

& Castelli, 2007; Macel *et al.*, 2007; Alexander *et al.*, 2015; Tomiolo *et al.*, 2015) for

studying local adaptation and, more recently, for studying their responses to a climate

that matches conditions predicted by climate change scenarios (the so called "space-

for-time approach"). Reciprocal transplants have also been applied to entire

communities in studies of soil microbiomes (Waldrop & Firestone, 2006; Lazzaro *et*

al., 2011), leaf litter (Ayres *et al.*, 2009; Allison *et al.*, 2013), and occasionally to

whole plant communities in different habitats ranging from wetlands to alpine

grasslands (Maranon & Bartolome, 1993; Wetzel *et al.*, 2004; Wu *et al.*, 2012;

Alexander *et al.*, 2015). However, the potential for using whole community reciprocal

transplants to study plant community response to climate change has not been fully

exploited, particularly in dryland systems, which often provide ideal conditions.

Dryland ecosystems are often dominated by annual plants that survive the dry

season as a permanent seed bank (Cohen, 1966). Therefore, the community (i.e. the

seed bank) can be conveniently transplanted as a whole during the dry season without

any damage to the plants. In addition, by transplanting seed banks with their

associated soil, it is possible to evaluate plant communities' responses to climate

while preserving soil abiotic and biotic interactions. To test the response of dryland

annual plant communities characterized by very different climates, we transplanted

home soil with seed bank among three sites situated along a steep aridity gradient in

the Eastern Mediterranean region, ranging from arid to Mediterranean climate. In this

region, rainfall is the main limiting factor for plant growth (Ziv *et al.*, 2014) and

differs up to eight-fold between the driest and wettest site (Holzapfel *et al.*, 2006). We

classified species based on their climatic requirements, adopting the Climatic Niche

Group approach (CNG; *sensu* Bilton et al. 2016) that has been successfully employed

for the species in our study region (Bilton *et al.*, 2016) and in other dryland

ecosystems (Liu *et al.*, 2018). By identifying those species responsive to drier or

wetter conditions, this approach provided us with testable predictions about directions

of shifts in community assembly across climates within the reciprocal transplants.

Finally, the study sites used for our reciprocal transplant also hosted a long-term

climate manipulation experiment (Tielbörger *et al.*, 2014). This allowed for a

qualitative comparison between long-term dynamics, resulting from consistently

imposed climate change, and the short-term responses observed in our transplant

experiment.

We predicted that the community emerging from the reciprocal soil transplants

would be greatly determined by community origin, with fewer individuals emerging

from drier origins than wetter origins. Secondly, we hypothesized that, regardless of

their origin, communities emerging at the drier transplant site (i.e. lower rainfall

availability) would experience a reduction in total biomass and plant density. We also

predicted that climate would select the emerging community from the species pool of

each origin in a predictable manner, with more wet adapted species emerging when

communities were exposed to wetter climates, and more dry-adapted species in drier

climates.

Methods

Study area

This study was conducted in Israel at three fenced sites (area approximately 100 m

x 400 m) with respectively Mediterranean (M), semi-arid (SA) and arid (A) climate.

The three study sites share the same calcareous bedrock, southern aspect, altitude and

mean annual temperatures, so that they differ chiefly in mean and variance of annual

rainfall, and vegetation. The M site is located southwest of Jerusalem (N 31° 42' E

35° 3') at 620 masl, on Terra Rossa soil. The climate is characterized by 550 mm

average annual rainfall with 20% inter-annual variation. The SA site (N 31° 23' E 34°

54') is located in the northern portion of the Negev Desert near the city of Beersheba,

at 590 masl, on Light Brown Rendzina. Average annual rainfall is 270 mm with approximately 30% inter-annual variation. The A site is located in the central Negev near Sde Boqer (N 30º52' E 34º46') at 470 masl, on desert Lithosol. Average annual rainfall amounts to 90 mm with 43% inter-annual variation (Holzapfel *et al.*, 2006). The plant communities at the three sites are semi-natural shrublands dominated by *Sarcopoterium spinosum* (L.) Spach, and winter annuals (approx. 85% of all species) that persist during summer in the form of dormant seed banks stored in the soil (Noy-Meir, 1973; Alon & Steinberger, 1999). The species pool is overlapping among sites, and annual plant cover amounts to 25% at the M site, 10% at the SA site, and < 1% at the A site (Tielbörger *et al.*, 2014).

Experimental set up

During summer of 2010, we collected soil with seed bank from forty square plots (20cm x 20cm, depth: 5cm) at the M and A sites and sixty plots at the SA site. Within each site, plots were situated at least 20 cm apart from each other, and away from rocks and shrubs. Following Tomiolo et al. (2015) soil collected from each site was pooled to produce a baseline community. Previous studies showed that small-scale heterogeneity in the soil seed bank may be very large, with some patches having almost no seeds and others very many (Siewert & Tielbörger, 2010). Therefore, we pooled the soil samples per site prior to the transplant, following the procedure adopted in many previous studies using field soil (Maranon & Bartolome, 1993; Macel *et al.*, 2007; Burns & Strauss, 2011; Lazzaro *et al.*, 2011).

The soil was stored in a net-house at the University of Rehovot, Israel, where it experienced summer temperatures necessary for breaking seed dormancy (Baskin *et al.*, 1993). In September 2010, twenty of the previously excavated plots at each site were randomly selected and filled with home soil, while the remaining plots were filled with soil from the closest away-from-home site (i.e. M site received M and SA soil; SA site received M, SA and A soil; A site received SA and A soil, Supplementary Material Appendix 1 Fig. A1). Transplanted soil was separated from the surrounding soil by a layer of absorbent paper that provided initial isolation between soils, while not impeding water percolation. After transplanting, we placed patches of organza (a thin transparent fabric) over the surface of each plot to avoid contamination from seed dispersal or seed predation (Petrů & Tielbörger, 2008), and we removed them at the time of germination.

Because the transplants were carried out during the dormant season we could

relocate the community of dormant seeds and soil biota with minimum damage. By

transplanting communities with their maternal soil we could test direct effects of

climate (e.g. decreasing rainfall) while preserving biotic interactions with

neighbouring plants and soil biota, which are also affected by the novel climate

(Emmett *et al.*, 2004). At peak development (spring 2011), we recorded the identity

and number of individuals of the emerging species in each plot. In order to minimize

edge effects, we excluded plants growing in the outer 1 cm margin of each plot. After

species identification, aboveground biomass was collected, oven-dried at 70°C for 48

hours and weighed.

Unfortunately, the season of recording was very dry and the arid site received

only 30% of the average annual rainfall. Therefore, only a handful of seedlings of two

desert species (*Stipa capensis*, *Erodium touchyanum*) emerged at the arid site. As a

result, there was no home arid community to be compared to the transplants, and we

had to restrict our subsequent analyses to the reciprocal transplants between the SA

and M community origins.

Climatic Niche Groups (CNG)

Each species within the target communities was assigned to a Climatic Niche

Group (Bilton *et al.*, 2016) classified by their distribution range in relation to rainfall.

This approach has proven powerful for predicting species-specific response to climate

change (Bilton *et al.*, 2016; Liu *et al.*, 2018). A similar method has been employed for

defining thermal niches of species in high elevation and tundra habitats (Gottfried *et*

al., 2012; Elmendorf *et al.*, 2015), and it is conceptually similar to Ellenberg values,

which determine species habitat requirements based on several abiotic parameters

(Ellenberg, 1974).

The rationale for the CNG grouping is that rainfall is the main driver of

community composition in the region, therefore species sharing similar climate

adaptations (approximated by the realized climatic rainfall niche) are likely to co-

occur in the same community (García-Camacho *et al.*, 2017). Species realized

climatic niche values were derived as in Bilton *et al.* (2016). For each single species

the observed occurrences within Israel (distribution range) were overlaid with mean

annual rainfall climate data, and the mean value was taken (obtained from BioGIS,

2012, available at http://www.biogis.huji.ac.il/). Boundaries between climatic niche

groups spanned similar ranges of average annual rainfall (approximately 130 mm) and

resulted in four groups that ranked species with respect to their hypothesized response

to climate. Climatic Niche Group 1 (CNG1) represented species associated with the

lowest rainfall extremes of the gradient, conversely CNG4 gathered species

distributed in areas with high rainfall. Species from all four CNGs were present in

both communities (Supplementary Material, Appendix 2 Table A1), but varied in

their proportional representation at each site, and could therefore be compared across

sites and climates (Bilton *et al.*, 2016).

Statistical analyses

We first analyzed how total density (number of individuals per plot), total

biomass, species richness (number of species per plot) and diversity (Shannon-Wiener

Index) varied in response to climate, community origin and their interaction. In

addition, we analyzed how the number of individuals belonging to each climatic niche

group per plot (i.e. CNG density) varied in response to climate, community origin,

with respect to the four-level categorical explanatory variable CNG identity (i.e.

CNG1 – CNG4), including all two-way and three-way interactions. We applied

generalized linear models with negative binomial distribution to total individual, CNG

density and species richness using the MASS package (Venables *et al.*, 2002) within

the R software version 3.3.3 (R Development Core Team, 2014). Biomass and species

diversity were analyzed using linear models. To meet model requirements biomass

square root transformed. The significance of the models was assessed with a Type 3

ANOVA, using the "car" package (Fox & Weisberg, 2011).

Visual representation of the CNG density interactions was done using log-ratios

calculated from the overall mean abundance of each group in each climate or origin.

Showing relative change in overall abundances was also extremely helpful for

visualizing the significant interactions we found with our models.

For testing how species composition varied with community origin and climate

we used Redundancy Analysis (RDA, (Legendre *et al.*, 2011)) in the R package

'vegan' (Oksanen *et al.*, 2015). The interaction term was included in a full model and

confirmed using a step-wise approach. The data were Hellinger transformed

(Legendre & Gallagher, 2001) and scaled within plots. Significance of the model was

tested using 999 permutations. To test if species composition could be explained by

rainfall distribution range we regressed the resulting RDA 'species mean scores'

against the 'climatic niche value' of each species, both for individual species and for

the CNG classifications. Furthermore, we performed an RDA on the community-

weighted means (Garnier *et al.*, 2007) using the species 'climatic niche value' as a

pseudo-trait.

Results

Overall, 97 species were recorded in our transplant plots, among which 12.3%

were grasses, 23.7% legumes and 64% belonged to other families (Supplementary

Material, Appendix 2 Table A1). In total, 68 species emerged from the semi-arid soil

seed bank, 81 from the Mediterranean origin, and 53 species were shared between the

two origins. Fourteen of these appeared in all four combinations of community origin

and climate.

Total plant density, diversity, richness and biomass

Number of species, species diversity, total plant density and total biomass (Fig. 1,

Table 1) were all significantly higher for M community origins rather than SA

origins. Additionally, M community origins attained significantly lower species

richness and diversity and biomass (Fig. 1) when exposed to the drier SA climate

compared to their home M climate. For the SA community origins, climate had no

significant effect on total plant density, biomass, species richness or diversity (Fig.1,

Table 1).

Species composition

The RDA indicated four distinct communities emerging from the respective

treatments, with a significant effect of community origin and climate on species

assembly, as well as a significant interaction between these terms (Fig. 2a, b). Using

simple correlations we assessed which plots/species scores changed and had most

impact on each axis. We obtained three main RDA axes describing the species

composition. For plot mean scores, RDA1 (9.2% explained) was correlated to overall

differences between community origins, whereas the constrained RDA2 (3.3%;

explained) and RDA3 (1.7%; explained) distinguished the climate x community

origin interaction term. For species mean scores, RDA1 was positively correlated to

species Climatic Niche values, and the correlation was positive but less strong for

RDA2 and RDA3 (Fig. 2c). Results were further validated by an RDA on the

community weighted mean traits using species Climatic Niche values as a trait, and

showed significant community origin and climate effects (p<0.05). In combination,

these results suggest that rainfall niche partially explained variation in species

composition across treatments.

CNG density across sites and community origins

Overall, densities of individuals in each CNG group significantly differed

(CNG identity effect: Fig. 3a, b, Table 2), and all group densities were higher in M

origin than SA origin (Origin and Origin x CNG effect; Fig. 3c; Table 2). The

representation of CNGs in the communities also changed significantly across

climates, dry groups were more abundant in SA climate and wet groups were more

abundant in M climate (Climate x CNG effect: Fig. 3c, Table 2, Supplementary

Material Appendix 2 Table A2). The magnitude in CNG shifts across climates was

different among community origins as indicated by a significant 3-way interaction

(CNG identity x community origin x climate, Table 2, Fig. 3e, f). In SA community

origins, the mean abundance of individuals belonging to dry CNGs (CNG 1 and 2)

was halved in the wet (M) compared to the dry (SA) climate; on the other hand

individuals belonging to CNG 4, the wettest adapted group, were 6.5 times more

abundant in SA communities origins emerging in the wet climate (Fig. 3a, e). In M

community origins, the shift in CNG densities across climates was less strong

compared to SA origins, but the hierarchical response of the CNGs was in the same

order (i.e. densities of dry CNGs were higher in the dry climate and densities of wet

CNGs were higher in the wet climate). The largest shift in density was seen in the

wettest CNG that counted twice as many individuals in wet *vs*. dry climate (Fig. 3b,

f).

Discussion

Our results revealed that climate played a large role in determining the species

assemblages in our whole community transplant experiment, and that such changes

were predicted by species-specific climatic adaptations. One of our most notable

findings was that the communities showed different qualitative and quantitative

responses to climate change depending on their origin. Mediterranean community

origins from a wetter and more predictable climate, responded with changes in species

richness, diversity and total biomass, but showed small shifts in community

composition. Conversely, semi-arid community origins, from a drier and more

unpredictable climate, showed little variation in species richness, diversity and

biomass, and large shifts in species and CNG composition.

Perhaps unsurprisingly, and as predicted from previous observations across the

rainfall gradient (Tielbörger *et al.*, 2014), communities establishing from the

Mediterranean (M) origins had higher individual density and total biomass than semi-

arid (SA) origins. This is also consistent with other dryland systems (Guo & Brown,

1997; Cleland *et al.*, 2013). The higher plant density in Mediterranean community

origins corresponded with higher densities of individuals for all CNG groups,

Interestingly, the species composition and CNG abundances in the experimental

communities rapidly matched the concomitant climate. Namely, plants establishing

from both community origins showed a relative increase of dry adapted species (CNG

1 and 2) when exposed to the drier (SA) climate. Similarly, wet adapted species

(CNG 3 and 4) were more abundant in both communities when exposed to the wetter

(M) climate. This response was particularly large for semi-arid community origins

where the reshuffling in community composition was strikingly well explained by

hierarchical switches in CNG abundances. Therefore, CNGs revealed species

responses to short-term climate effects in a predictable way, suggesting potential

short-term selection mechanisms (e.g. environmental filtering) that act on the

communities in response to yearly differences in rainfall. This result is consistent with

the findings of Bilton *et al.* (2016), and suggests that CNGs are representative of

species rainfall requirements and possibly climatic adaptations at the different sites.

Our most intriguing finding was that climate filtered for predictable species

groups, but the magnitude of the structural community shifts was largely different

among community origins. In addition, the magnitude of the compositional shifts

between transplanted community origins was inversely related to changes in total

community parameters across climates. Mediterranean origin transplants, had smaller

compositional shifts, but larger shifts in biomass and richness. Conversely, semi-arid

community origins, with higher between- and within-year rainfall variability

(Tielbörger *et al.*, 2014), experienced the greatest shifts in community composition,

while showing only marginal response in community parameters. Consistent with our

results, large interannual compositional shifts were observed also in other

communities from drier and highly variable climates (Guo & Brown, 1997; Cleland *et*

al., 2013). Interestingly, we did not record an increase in total biomass or total density

in semi-arid community origins when exposed to wetter climates. This is possibly the

result of an increase in wet CNGs in the communities, which have a core distribution

in wetter regions where there are larger plant densities, productivity, and a higher

intensity of competition (Schiffers & Tielbörger, 2006; Liancourt & Tielbörger,

2009). Wet CNGs may possess a better competitive ability and might have curbed the

growth and density of dry CNGs, thus leading to little changes in total biomass and

total density.

Large inter-annual variation in species abundances in the short-term may lead to

higher community stability in the long-term (Bai *et al.*, 2004; Grime *et al.*, 2008).

Similar patterns have been previously explained in plant community studies, albeit in

a different context, by the portfolio effect (Doak *et al.*, 1998; Schindler *et al.*, 2015).

The portfolio effect predicts that greater numbers of species in a community lead both

mathematically and ecologically to a greater chance of asynchronous relationships

forming year-to-year. Here we show, consistent with previous studies (Cleland *et al.*,

2013; Hallett *et al.*, 2014), that in the community originating in a more unpredictable

climate (in our case the semi-arid community origin), greater asynchrony and greater

species turnover led to greater stability across climates. Plant species in more variable

climates have been found to exhibit a larger phenotypic plasticity (Sultan, 1987; Pratt

& Mooney, 2013; Lazaro-Nogal *et al.*, 2015; Spence *et al.*, 2016), which results in

fitness homeostasis (Richards *et al.*, 2006; Nicotra *et al.*, 2010). This could be an

explanation for the higher resistance of species in SA communities but also reveals an

interesting analogy with the community level, where 'homeostasis' may be associated

with a larger compositional change. Overall, we suggest that asynchronous shifts in

abundance of species according to their climate adaptations may allow for fast

responses to year-to-year climatic variation in dryland annual communities (Abbott *et*

al., 2017).

In the short-term, high species turnover may assure community stability in

biomass and density, but in the long-term, such processes may also lead to greater

resistance and stability of dry communities to rainfall fluctuations by favoring species

adapted to arid conditions. This high turnover is possible without immediate loss of

species because in dryland environments, plants often display bet-hedging strategies

such as long-lived seed banks and seed dormancy that can buffer against inter-annual

fluctuations (Petrů & Tielbörger, 2008; Tielbörger *et al.*, 2012). Delayed germination

of dormant seeds during unfavorable years decreases the risk of extinction over time

and also promotes coexistence of species with different climatic requirements via

storage effects (Chesson & Grubb, 1990; Pake & Venable, 1995). Interestingly, the

findings from this short-term community transplant study are consistent with those of

a parallel long-term experiment conducted at the same study sites (Tielbörger *et al.*,

2014), where community composition was monitored for 10 years in permanent plots

receiving respectively ambient rainfall, experimental drought (-30% rainfall) or

increased rainfall (+30% rainfall). Plant communities exposed to the long-term

climate manipulation treatments showed no detectable long-term effect on total

density, species richness and community biomass (Tielbörger *et al.*, 2014). However,

at both time-scales, shifts in species abundance in relation to their CNG was observed

(Bilton *et al.*, 2016). While in our short-term study we found these effects to be more

pronounced for semi-arid community origins, the long-term study found stronger

patterns in Mediterranean communities (Bilton *et al.*, 2016). This suggests that high

inter-annual community fluctuations may contribute to stability in community

composition in the long run, whereas low species turnover across years may result in

long-term loss of wet-adapted species.

Our overall findings allow some careful conclusions about the potential

response of these communities to climate change. It should be noted that the variation

in rainfall experienced by the community origins in our study approximated the

extremes of climatic variability at each site, but exceeded the decrease in rainfall

predicted by climate change scenarios for the next 50-80 years (Smiatek *et al.*, 2011;

Tomiolo *et al.*, 2015). These results suggest that, as long as inter-annual climatic

fluctuations keep within the limits of climatic variability commonly experienced by

these communities, and rainy years that replenish the seed bank periodically occur,

wet adapted species will persist within the communities. However, with increasing

drought and unpredictability, communities are likely to experience species loss that

will affect primarily species with high rainfall requirements (Tielbörger *et al.*, 2014;

Bilton *et al.*, 2016). The similarity of results between the long-term experiment and

our reciprocal transplant indicate that the latter may be a powerful complement to

long-term field experiments. However, annual communities are particularly suited for

our experimental approach and the same may not hold for long-lived communities.

For example, in temperate systems a lag between shifts in climatic conditions and

subsequent changes in community structure is often observed (Adler & Levine, 2007; Jones *et al.*, 2016). The unexpected community resistance and resilience of dryland ecosystems to extreme events compared with temperate ecosystems (Ruppert *et al.*, 2015), may be attributable to the large variability in rainfall to which plant species are pre-adapted via bet hedging strategies (Tielbörger *et al.*, 2012; Gremer & Venable, 2014) or enhanced phenotypic plasticity (Petrů *et al.*, 2006). Such differences among rainfall variability and plant life history should be taken into account when drawing comparisons among habitats.

Declarations: KT and ST developed the experimental design. ST set up the experiments and collected the data. MB and ST performed statistical analyses. ST wrote the first draft of the manuscript and all authors contributed substantially to the following versions. We declare that we do not have conflicts of interest.

Acknowledgements: Jaime Kigel and Marcelo Sternberg provided logistic support. The Hebrew University of Jerusalem (Rehovot) kindly provided material for fieldwork. We thank Jake Alexander for providing comments on a previous version of the manuscript. This study is part of the GLOWA Jordan River Project and was funded by the German Ministry of Education and Research (BMBF). Further support for MB and ST was obtained by the German Research Foundation (TI338_12-1; TI338_11-1; and TI338_11-2; TI 338/15-1).

Data Accessibility: upon publication data will be made accessible on Dryad

References

Abbott RE, Doak DF, Peterson ML. 2017. Portfolio effects, climate change, and the persistence of small populations: analyses on the rare plant *Saussurea weberi*. *Ecology* **98**(4): 1071-1081.

Adler PB, Levine JM. 2007. Contrasting relationships between precipitation and

species richness in space and time. *Oikos* **116**(2): 221-232.

Alexander JM, Diez JM, Hart SP, Levine JM. 2016. When climate reshuffles

competitors: a call for experimental macroecology. *Trends in Ecology and*

Evolution **31**(11): 831-841.

Alexander JM, Diez JM, Levine JM. 2015. Novel competitors shape species'

responses to climate change. *Nature* **525**(7570): 515-518.

Allison SD, Lu Y, Weihe C, Goulden ML, Martiny AC, Treseder KK, Martiny

JB. 2013. Microbial abundance and composition influence litter

decomposition response to environmental change. *Ecology* **94**(3): 714-725.

Alon A, Steinberger Y. 1999. Response of the soil microbial biomass and nematode

population to a wetting event in nitrogen-amended Negev desert plots. *Biology*

and Fertility of Soils **30**(1-2): 147-152.

Anderson R, Gordon D, Crawley M, Hassell M. 1982. Variability in the abundance

of animal and plant species. *Nature* **296**(5854): 245-248.

Ayres E, Steltzer H, Simmons BL, Simpson RT, Steinweg JM, Wallenstein MD,

Mellor N, Parton WJ, Moore JC, Wall DH. 2009. Home-field advantage

accelerates leaf litter decomposition in forests. *Soil Biology and Biochemistry*

41(3): 606-610.

Bai Y, Han X, Wu J, Chen Z, Li L. 2004. Ecosystem stability and compensatory

effects in the Inner Mongolia grassland. *Nature* **431**(7005): 181.

Barbosa ERM, Tomlinson KW, Carvalheiro LG, Kirkman K, de Bie S, Prins

HHT, van Langevelde F. 2014. Short-term effect of nutrient availability and

rainfall distribution on biomass production and leaf nutrient content of

savanna tree species. *PloS ONE* **9**(3).

Baskin CC, Chesson PL, Baskin JM. 1993. Annual seed dormancy cycles in two

desert winter annuals. *Journal of Ecology* **81**: 551-556.

Bilton MC, Metz J, Tielbörger K. 2016. Climatic niche groups: A novel application

of a common assumption predicting plant community response to climate

change. *Perspectives in Plant Ecology, Evolution and Systematics* **19**: 61-69.

BioGIS 2012. Israel Biodiversity Information System [

http://www.biogis.huji.ac.il/]. available at

http://www.biogis.huji.ac.il/.

Blume-Werry G, Kreyling J, Laudon H, Milbau A. 2016. Short-term climate

change manipulation effects do not scale up to long-term legacies: effects of

an absent snow cover on boreal forest plants. *Journal of Ecology* **104**(6):

1638-1648.

Bonebrake TC, Mastrandrea MD. 2010. Tolerance adaptation and precipitation

changes complicate latitudinal patterns of climate change impacts.

Proceedings of the National Academy of Sciences of the United States of

America **107**(28): 12581-12586.

Brown JH, Morgan Ernest SK, Parody JM, Haskell JP. 2001. Regulation of

diversity: maintenance of species richness in changing environments.

Oecologia **126**: 321-332.

Burns JH, Strauss SY. 2011. More closely related species are more ecologically

similar in an experimental test. *Proceedings of the National Academy of*

Sciences **108**(13): 5302-5307.

Casper BB, Castelli JP. 2007. Evaluating plant-soil feedback together with

competition in a serpentine grassland. *Ecology Letters* **10**(5): 394-400.

Chesson PL, Grubb P. 1990. Geometry, heterogeneity and competition in variable

environments. *Philosophical Transactions of the Royal Society of London B:*

Biological Sciences **330**(1257): 165-173.

Cleland EE, Collins SL, Dickson TL, Farrer EC, Gross KL, Gherardi LA,

Hallett LM, Hobbs RJ, Hsu JS, Turnbull L, et al. 2013. Sensitivity of

grassland plant community composition to spatial vs. temporal variation in

precipitation. *Ecology* **94**(8): 1687-1696.

Cohen D. 1966. Optimizing reproduction in a randomly varying environment.

Journal of Theoretical Biology **12**: 119-129.

Cubasch U, vonStorch H, Waszkewitz J, Zorita E. 1996. Estimates of climate

change in Southern Europe derived from dynamical climate model output.

Climate Research **7**(2): 129-149.

De Dato GD, Pellizzaro G, Cesaraccio C, Sirca CB, De Angelis P, Duce P, Spano

DEI, Scarascia Mugnozza G. 2006. Effects of warmer and drier climate

conditions on plant composition and biomass production in a Mediterranean

shrubland community. *Forest@* **3**(4): 511-526.

Doak DF, Bigger D, Harding EK, Marvier MA, O'Malley RE, Thomson D. 1998.

The statistical inevitability of stability-diversity relationships in community

ecology. *American Naturalist* **151**(3): 264-276.

Doblas−Miranda E, Martínez−Vilalta J, Lloret F, Álvarez A, Ávila A, Bonet F,

Brotons L, Castro J, Curiel Yuste J, Díaz M. 2015. Reassessing global

change research priorities in mediterranean terrestrial ecosystems: how far

have we come and where do we go from here? *Global Ecology and*

Biogeography **24**(1): 25-43.

Ellenberg H. 1974. Indicator values of vascular plants in central Europe. *Scripta*

Geobotanica **9**: 97.

Elmendorf SC, Henry GHR, Hollister RD, Fosaa AM, Gould WA, Hermanutz L,

Hofgaard A, Jónsdóttir IS, Jorgenson JC, Lévesque E, et al. 2015.

Experiment, monitoring, and gradient methods used to infer climate change

effects on plant communities yield consistent patterns. *Proceedings of the*

National Academy of Sciences **112**(2): 448-452.

Emmett BA, Beier C, Estiarte M, Tietema A, Kristensen HL, Williams D,

Penuelas J, Schmidt I, Sowerby A. 2004. The response of soil processes to

climate change: results from manipulation studies of shrublands across an

environmental gradient. *Ecosystems* **7**(6): 625-637.

Fernandez-Going BM, Harrison SP, Anacker BL, Safford HD. 2013. Climate

interacts with soil to produce beta diversity in Californian plant communities.

Ecology **94**(9): 2007-2018.

Forey E, Touzard B, Michalet R. 2010. Does disturbance drive the collapse of biotic

interactions at the severe end of a diversity-biomass gradient? *Plant Ecology*

206(2): 287-295.

Fox J, Weisberg S. 2011. *An R Companion to Applied Regression*. Thousand Oaks

CA: Sage Publications.

Garcia RA, Cabeza M, Rahbek C, Araújo MB. 2014. Multiple Dimensions of

Climate Change and Their Implications for Biodiversity. *Science* **344**(6183).

García-Camacho R, Metz J, Bilton MC, Tielbörger K. 2017. Phylogenetic

structure of annual plant communities along an aridity gradient. Interacting

effects of habitat filtering and shifting plant–plant interactions. *Israel Journal*

of Plant Sciences: 1-13.

Garnier E, Lavorel S, Ansquer P, Castro H, Cruz P, Dolezal J, Eriksson O,

Fortunel C, Freitas H, Golodets C, et al. 2007. Assessing the effects of land-

use change on plant traits, communities and ecosystem functioning in

grasslands: a standardized methodology and lessons from an application to 11

European sites. *Annals of Botany* **99**(5): 967-985.

Gilman SE, Urban MC, Tewksbury J, Gilchrist GW, Holt RD. 2010. A

framework for community interactions under climate change. *Trends in*

Ecology & Evolution **25**(6): 325-331.

Gottfried M, Pauli H, Futschik A, Akhalkatsi M, Barancok P, Benito Alonso JL,

Coldea G, Dick J, Erschbamer B, Fernandez Calzado MR, et al. 2012.

Continent-wide response of mountain vegetation to climate change. *Nature*

Clim. Change **2**(2): 111-115.

Gremer JR, Venable DL. 2014. Bet hedging in desert winter annual plants: optimal

germination strategies in a variable environment. *Ecology Letters* **17**(3): 380-

387.

Grime JP, Fridley JD, Askew AP, Thompson K, Hodgson JG, Bennett CR. 2008.

Long-term resistance to simulated climate change in an infertile grassland.

Proceedings of the National Academy of Sciences of the United States of

America **105**(29): 10028-10032.

Guo Q, Brown JH. 1997. Interactions between winter and summer annuals in the

Chihuahuan Desert. *Oecologia* **111**(1): 123-128.

Hallett LM, Hsu JS, Cleland EE, Collins SL, Dickson TL, Farrer EC, Gherardi

LA, Gross KL, Hobbs RJ, Turnbull L, et al. 2014. Biotic mechanisms of

community stability shift along a precipitation gradient. *Ecology* **95**(6): 1693-

1700.

Harrison SP, Gornish ES, Copeland S. 2015. Climate-driven diversity loss in a

grassland community. *Proceedings of the National Academy of Sciences of the*

United States of America **112**(28): 8672-8677.

Hobbs RJ, Arico S, Aronson J, Baron JS, Bridgewater P, Cramer VA, Epstein

PR, Ewel JJ, Klink CA, Lugo AE, et al. 2006. Novel ecosystems: theoretical

and management aspects of the new ecological world order. *Global Ecology*

and Biogeography **15**(1): 1-7.

Holzapfel C, Tielbörger K, Parag HA, Kigel J, Sternberg M. 2006. Annual plant-

shrub interactions along an aridity gradient. *Basic and Applied Ecology* **7**:

268-279.

Jones SK, Collins SL, Blair JM, Smith MD, Knapp AK. 2016. Altered rainfall

patterns increase forb abundance and richness in native tallgrass prairie.

Scientific Reports **6**: 20120.

Lazaro-Nogal A, Matesanz S, Godoy A, Perez-Trautman F, Gianoli E,

Valladares F. 2015. Environmental heterogeneity leads to higher plasticity in

dry-edge populations of a semi-arid Chilean shrub: insights into climate

change responses. *Journal of Ecology* **103**(2): 338-350.

Lazzaro A, Gauer A, Zeyer J. 2011. Field-scale transplantation experiment to

investigate structures of soil bacterial communities at pioneering sites. *Applied*

and Environmental Microbiology **77**(23): 8241-8248.

Legendre P, Gallagher ED. 2001. Ecologically meaningful transformations for

ordination of species data. *Oecologia* **129**(2): 271-280.

Legendre P, Oksanen J, ter Braak CJF. 2011. Testing the significance of canonical

axes in redundancy analysis. *Methods in Ecology and Evolution* **2**(3): 269-

277.

Liancourt P, Tielbörger K. 2009. Competition and a short growing season lead to

ecotypic differentiation at the two extremes of the ecological range.

Functional Ecology **23**(2): 397-404.

Lindenmayer DB, Likens GE, Andersen A, Bowman D, Bull CM, Burns E,

Dickman CR, Hoffmann AA, Keith DA, Liddell MJ. 2012. Value of long−

term ecological studies. *Austral Ecology* **37**(7): 745-757.

Link SO, Smith JL, Halvorson JJ, Bolton H. 2003. A reciprocal transplant

experiment within a climatic gradient in a semiarid shrub-steppe ecosystem:

effects on bunchgrass growth and reproduction, soil carbon, and soil nitrogen.

Global Change Biology **9**(7): 1097-1105.

Liu D, Peñuelas J, Ogaya R, Estiarte M, Tielbörger K, Slowik F, Yang X, Bilton

MC. 2018. Species selection under long−term experimental warming and

604 drought explained by climatic distributions. *New Phytologist* **217**(4): 1494-

605 1506.

606 **Macel M, Lawson CS, Mortimer SR, Smilauerova M, Bischoff A, Cremieux L,**

607 **Dolezal J, Edwards AR, Lanta V, Bezemer TM, et al. 2007.** Climate vs.

608 soil factors in local adaptation of two common plant species. *Ecology* **88**(2):

609 424-433.

610 **Maestre FT, Quero JL, Gotelli NJ, Escudero A, Ochoa V, Delgado-Baquerizo M,**

611 **Garcia-Gomez M, Bowker MA, Soliveres S, Escolar C, et al. 2012.** Plant

612 species richness and ecosystem multifunctionality in global drylands. *Science*

613 **335**: 214-218.

614 **Maranon T, Bartolome JW. 1993.** Reciprocal transplants of herbaceous

615 communities between *Quercus agrifolia* woodland and adjacent grassland.

616 *Journal of Ecology* **81**(4): 673-682.

617 **Miranda JD, Armas C, Padilla FM, Pugnaire FI. 2011.** Climatic change and

618 rainfall patterns: effects on semi-arid plant communities of the Iberian

619 Southeast. *Journal of Arid Environments* **75**(12): 1302-1309.

620 **Nicotra AB, Atkin OK, Bonser SP, Davidson AM, Finnegan EJ, Mathesius U,**

621 **Poot P, Purugganan MD, Richards CL, Valladares F, et al. 2010.** Plant

622 phenotypic plasticity in a changing climate. *Trends in Plant Science* **15**(12):

623 684-692.

Noy-Meir I. 1973. Desert ecosystems: Environment and producers. *Annual Review of*

Ecology and Systematics **4**: 25–51.

Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB,

Simpson GL, Solymos P, Henry M, Stevens H, et al. 2015. vegan:

Community ecology package.

Pake CE, Venable DL. 1995. Is coexistence of Sonoran desert annuals mediated by

temporal variablilty in reproductive success? *Ecology* **76**: 246-261.

Parmesan C, Hanley ME. 2015. Plants and climate change: complexities and

surprises. *Annals of Botany* **116**(6): 849-864.

Peñuelas J, Prieto P, Beier C, Cesaraccio C, De Angelis P, De Dato G, Emmett

BA, Estiarte M, Garadnai J, Gorissen A. 2007. Response of plant species

richness and primary productivity in shrublands along a north–south gradient

in Europe to seven years of experimental warming and drought: reductions in

primary productivity in the heat and drought year of 2003. *Global Change*

Biology **13**(12): 2563-2581.

Petrů M, Tielbörger K. 2008. Germination strategies of annual plants under

changing climatic conditions: teasing apart local and regional effects.

Oecologia **155**: 717-728.

Petrů M, Tielbörger K, Belkin R, Sternberg M, Jeltsch F. 2006. Life history

variation in an annual plant under two opposing environmental constraints

along an aridity gradient. *Ecography* **29**(1): 66-74.

Pratt JD, Mooney KA. 2013. Clinal adaptation and adaptive plasticity in *Artemisia*

californica: implications for the response of a foundation species to predicted

climate change. *Global Change Biology* **19**(8): 2454-2466.

R Development Core Team 2014. R: A language and environment for statistical

computing. .In Computing RFfS. Vienna, Austria.

Richards CL, Bossdorf O, Muth NZ, Gurevitch J, Pigliucci M. 2006. Jack of all

trades, master of some? On the role of phenotypic plasticity in plant invasions.

Ecology Letters **9**(8): 981-993.

Rinnan R, Michelsen A, Baath E, Jonasson S. 2007. Fifteen years of climate

change manipulations alter soil microbial communities in a subarctic heath

ecosystem. *Global Change Biology* **13**(1): 28-39.

Ruppert JC, Harmoney K, Henkin Z, Snyman HA, Sternberg M, Willms W,

Linstädter A. 2015. Quantifying drylands' drought resistance and recovery:

the importance of drought intensity, dominant life history and grazing regime.

Global Change Biology **21**(3): 1258-1270.

Sala OE, Lauenroth WK. 1982. Small rainfall events: an ecological role in semiarid

regions. *Oecologia* **53**: 301-304.

Sardans J, Peñuelas J, Estiarte M, Prieto P. 2008. Warming and drought alter C

and N concentration, allocation and accumulation in a Mediterranean

shrubland. *Global Change Biology* **14**(10): 2304-2316.

Schiffers K, Tielbörger K. 2006. Ontogenetic shifts in interactions among annual

plants. *Journal of Ecology* **94**(2): 336-341.

Schindler DE, Armstrong JB, Reed TE. 2015. The portfolio concept in ecology and

evolution. *Frontiers in Ecology and the Environment* **13**(5): 257-263.

Shi Z, Sherry R, Xu X, Hararuk O, Souza L, Jiang LF, Xia JY, Liang JY, Luo

YQ. 2015. Evidence for long-term shift in plant community composition

under decadal experimental warming. *Journal of Ecology* **103**(5): 1131-1140.

Siewert W, Tielbörger K. 2010. Dispersal-dormancy relationships in annual plants:

putting model predictions to the test. *American Naturalist* **176**(4): 490-500.

Smiatek G, Kunstmann H, Heckl A. 2011. High-resolution climate change

simulations for the Jordan River area. *Journal of Geophysical Research-*

Atmospheres **116**.

Spence LA, Liancourt P, Boldgiv B, Petraitis PS, Casper BB. 2016. Short-term

manipulation of precipitation in Mongolian steppe shows vegetation

influenced more by timing than amount of rainfall. *Journal of Vegetation*

Science **27**(2): 249-258.

681 **Sultan SE 1987.** Evolutionary Implications of Phenotypic Plasticity in Plants. In:

682   Hecht MK, Wallace B, Prance GT eds. *Evolutionary Biology: Volume 21*.

683   Boston, MA: Springer US, 127-178.

684 **Thompson PL, Beisner BE, Gonzalez A. 2015.** Warming induces synchrony and

685   destabilizes experimental pond zooplankton metacommunities. *Oikos* **124**(9):

686   1171-1180.

687 **Tielbörger K, Bilton MC, Metz J, Kigel J, Holzapfel C, Lebrija-Trejos E,**

688   **Konsens I, Parag HA, Sternberg M. 2014.** Middle-Eastern plant

689   communities tolerate 9 years of drought in a multi-site climate manipulation

690   experiment. *Nature Communications* **5**.

691 **Tielbörger K, Petrů M, Lampei C. 2012.** Bet-hedging germination in annual plants:

692   a sound empirical test of the theoretical foundations. *Oikos* **121**(11): 1860-

693   1868.

694 **Tilman D, Downing JA. 1994.** Biodiversity and stability in grasslands. *Nature*

695   **367**(6461): 363-365.

696 **Tilman D, Lehman CL, Bristow CE. 1998.** Diversity-stability relationships:

697   statistical inevitability or ecological consequence? *The American Naturalist*

698   **151**(3): 277-282.

Tomiolo S, van der Putten WH, Tielbörger K. 2015. Separating the role of biotic

interactions and climate in determining adaptive response of plants to climate

change. *Ecology* **96**(5): 1298-1308.

Venables WN, Ripley BD, . 2002. *Modern Applied Statistics with S.* . New York,

N.Y.: Springer.

Waldrop MP, Firestone MK. 2006. Response of microbial community composition

and function to soil climate change. *Microbial Ecology* **52**(4): 716-724.

Wetzel PR, Kitchens WM, Brush JM, Dusek ML. 2004. Use a reciprocal transplant

study to measure the rate of plant community change in a tidal marsh along a

salinity gradient. *Wetlands* **24**(4): 879-890.

Williams JW, Jackson ST. 2007. Novel climates, no-analog communities, and

ecological surprises. *Frontiers in Ecology and the Environment* **5**(9): 475-482.

Wu ZT, Dijkstra P, Koch GW, Hungate BA. 2012. Biogeochemical and ecological

feedbacks in grassland responses to warming. *Nature Climate Change* **2**(6):

458-461.

Ziv B, Saaroni H, Pargament R, Harpaz T, Alpert P. 2014. Trends in rainfall

regime over Israel, 1975–2010, and their relationship to large-scale variability.

Regional Environmental Change **14**(5): 1751-1764.

Figures and tables

Table 1: Type III ANOVA table results for the models applied to individual density, species
richness, species diversity, and total biomass. Lines correspond to response variables and
columns to explanatory variables of each model. In each column the first value represent Chi-
square test values and the second the p-value. Probability values for significant terms are
reported in bold.

	Origin	**Climate**	**Origin x Climate**
Density	312.86; **<0.001**	0.12; 0.72	0.59; 0.44
Species richness	281.29; **<0.001**	12.99; **<0.001**	0.085; 0.77
Species diversity	130.41; **<0.001**	10.40; **0.001**	1.36; 0.24
Total biomass	103.86; **0.001**	2.34; 0.13	4.66; **0.03**

726
727

728 **Table 2: Type III ANOVA table of results for Generalized Linear Models determining the effect**
729 **of community origin, climate and Climatic Niche Group (CNG) on plant community densities**
730 **establishing from seed bank.**

Effect	DF	LR Chi-sq.	P-value
		236.9	
Origin	1	3	**<0.001**
Climate	1	2.35	0.12540
		343.2	
CNG	3	1	**<0.001**
Origin x Climate	1	0.07	0.78714
Origin x CNG	3	83.55	**<0.001**
Climate x CNG	3	50.09	**<0.001**
Origin x Climate x CNG	3	16.85	**<0.001**

731
732
733

Figure Captions:

Figure 1: Mean ±1SE A) species richness (i.e. number of species per plot), B) diversity C) total individual density, D) total biomass of annual plant communities established from two community origins ('M' Mediterranean; 'SA' Semi-Arid) grown in two sites (i.e. climates: 'M' Mediterranean; 'SA' Semi-Arid).

Figure 2: Graphical representation of Redundancy Analysis (RDA) showing species composition in community origins (M, SA) exposed to two different climates (M, SA). The ellipses represent the 95% confidence intervals for the groupings/categories. Fig. 2 a, b represents the plot centroids of each community origin-climate combination. Lines are vectors from the centre of a category to each site score (points). RDA-axis 1: correlated to distance between origins, RDA-axis 3: the effect of climate on SA community origins, and RDA-axis 2 the effect of climate on M community origins. In red: SA community origins - SA climate; yellow: SA origins - M climate; blue: M origins- M climate; green: M origins -SA climate. Fig. 2c represents the species centroids for each Climatic Niche Group (CNG). Lines are vectors connecting the centre of each group with species scores. In red: CNG1, yellow: CNG2, green: CNG3, blue: CNG4.

Figure 3: Impact of community origin, climate and Climatic Niche Group (CNG) on plant densities establishing from seed banks in home *vs*. away-from-home climate. Each Climatic Niche Group (CNG) aggregates species with similar climatic adaptation, ranging from dry climates (CNG 1) to wetter climates (CNG 4). Fig. 3 a, b: Total mean individuals' densities across climates and origins are broken down according to CNG abundances. Fig. 3 c-f: Shifts in the abundance of CNG densities across community origins (3c), climates (3d) and the combination of the two (3 e, f). Shifts in densities of CNG groups are expressed as log ratios, thus visualizing changes in plant density on a relative scale. Positive values indicate higher CNG abundances in SA community origins or climate, whereas negative values indicate higher CNG abundances in M community origins or climate.

Figure A1: Dots on the map represent the three experimental sites. On the right hand, the the scheme of the community transplants at each sites. Communities (soil and relative seed bank) were transplanted between the home and the closest away-from-home site.

790
791 **Figure 1**
792

793
Figure 2
794
795
796

Raw Density

2-way interactions

3-way interaction

Figure 3