Whole plant community transplants across climates reveal total community

stability due to large shifts in species assemblage

Authors: Sara Tomiolo[a,b*], Mark C. Bilton[a], Katja Tielbörger[a]

Affiliation:

a- Department of Ecology and Evolution, Plant Ecology Group, University of

Tübingen, Auf der Morgenstelle 5, 72076 Tübingen, Germany

b- Department of Bioscience, Aarhus University, Vejlsøvej 25, 8600 Silkeborg,

Denmark

Contacts:

*Corresponding author: sara.tomiolo@gmail.com; **Orcid ID:** orcid.org/0000-0002-

5071-5851

Mark C. Bilton: mark.bilton@uni-tuebingen.de

Katja Tielbörger: katja.tielboerger@uni-tuebingen.de

Running headline: Community stability under climate change

Keywords: annual plant communities, climate change, Climatic Niche Groups,

community ecology, reciprocal transplants, determinants of plant community

diversity and assemblage, distribution range, community stability, field experiments

Abstract

Climate change will decrease average precipitation and increase rainfall

variability in Eastern Mediterranean regions. This may affect the performance of

many plant species either directly or via altered biotic interactions in ways that are

hard to predict. In such highly fluctuating climates, year-to-year community variation

may override long-term selection processes, thus masking directional community

responses to climate change.

Here we tested short-term responses of dryland plant communities to sharp

changes in climatic conditions, by means of whole community reciprocal transplants

of soil and seed banks. We exposed annual plant communities to two very different

climates (Mediterranean and semi-arid) and measured changes in community

composition, total biomass, plant density, and species diversity. In addition, we

grouped species into dry-adapted and wet-adapted species in each community.

Our results revealed that climate played a large role in determining community

assembly. For both community origins, we observed a relative increase of dry-adapted

species when exposed to the drier climate, and a relative increase in wet-adapted

species in the wetter climate. However, the compositional shifts were much larger in

semi-arid origin communities. At the same time community density, biomass and

species richness remained remarkably stable across climates, indicating that

community stability was maintained between climates by predictable asynchronous

shifts of species. Similar but smaller shifts were present in the Mediterranean origin

communities, where exposure to the drier climate reduced biomass and species

richness.

Our study suggests that large asynchronous variation in species abundances,

matching high year-to-year rainfall variability, may provide a mechanism for

community homeostasis, and slow down selection processes in response to climate

change. However, increased occurrence of extreme droughts exceeding the climatic

fluctuations to which species are adapted may, in the long-term, lead to loss of wet-

adapted species.

Introduction

Understanding how climate may alter overall available resources (Sardans et al. 2008, Garcia et al. 2014) and impact upon community structure (Tilman and Downing 1994, Gilman et al. 2010) is a major challenge in current ecological research (Maestre et al. 2012, Parmesan and Hanley 2015). In cold regions for example, warming is likely to improve growing conditions and thus increase plant community biomass by increasing nutrient mobilization and expanding the length of the growing season (Garcia et al. 2014). In contrast, decreased rainfall in drier regions will likely have negative impacts on primary productivity, community composition, and their corresponding ecosystem services (Sala and Lauenroth 1982, Peñuelas et al. 2007, Miranda et al. 2011). These effects may be particularly strong in those dryland ecosystems for which climate predictions indicate increasing incidents of droughts (Cubasch et al. 1996, Smiatek et al. 2011). Here, droughts and increasing temperatures will increase evapotranspiration, shorten the growing season and limit access to nutrients, thereby decreasing total community biomass (Peñuelas et al. 2007, Doblas−Miranda et al. 2015, Harrison et al. 2015). In extreme cases, this may lead to the collapse of entire ecological communities (Forey et al. 2010).

One of the predicted impacts of climate change is a re-assembly of plant communities (Hobbs et al. 2006, Williams and Jackson 2007, Alexander et al. 2016) due to the differential ability of single species to either track their climatic niche or to survive under changed conditions, by means of adaptation or plasticity (Fernandez-Going et al. 2013, Shi et al. 2015). Such community reshuffling may be expressed in a loss or gain of certain species, a shift in species relative abundance or both. In plant communities already exposed to large inter-annual variations in climate, immediate rearrangement of community assembly in response to climate extremes may be an

inherent property of plant communities, and may promote community stability in the

long-term. This effect is particularly pronounced when species numbers are large and

population sizes vary asynchronously (Doak et al. 1998, Schindler et al. 2015).

Indeed, ecological theory and models support the idea that high inter-annual variation

in species response to climate can lead to community-level stability (Anderson et al.

1982, Tilman et al. 1998, Thompson et al. 2015, Abbott et al. 2017). This may be an

important mechanism for maintaining dryland communities' stability in response to

large year-to-year variation in rainfall, and for slowing down ongoing selection

processes due to climate change (Bonebrake and Mastrandrea 2010, Bilton et al.

2016).

Although long-term climate manipulations are the gold standard in ecological

climate impact research and are fundamental to understanding long-term community

shifts (Brown et al. 2001, Rinnan et al. 2007, Blume-Werry et al. 2016), they are very

costly to set up and maintain, often outliving funding cycles and scientific research

positions (Lindenmayer et al. 2012). The monitoring of communities for short-term

responses may be a useful complement to long-term experiments, as besides being

less costly, can be vitally important for parsing mechanistic information about plant

responses to large inter-annual variation, as well as extreme events (De Dato et al.

2006, Barbosa et al. 2014, Blume-Werry et al. 2016). Reciprocal transplants represent

a promising approach for indirectly studying plant responses to climate change on a

short temporal scale. These manipulative experiments have been widely adopted in

single species (e.g. Link et al. 2003, Casper and Castelli 2007, Macel et al. 2007,

Alexander et al. 2015, Tomiolo et al. 2015) for studying local adaptation and, more

recently, for studying their responses to a climate that matches conditions predicted

by climate change scenarios (the so called "space-for-time approach"). Reciprocal

transplants have also been applied to entire communities in studies of soil

microbiomes (Waldrop and Firestone 2006, Lazzaro et al. 2011), leaf litter (Ayres et

al. 2009, Allison et al. 2013), and occasionally to whole plant communities in

different habitats ranging from wetlands to alpine grasslands (Maranon and

Bartolome 1993, Wetzel et al. 2004, Wu et al. 2012, Alexander et al. 2015). However,

the potential for using whole community reciprocal transplants to study plant

community response to climate change has not been fully exploited, particularly in

dryland systems, which often provide ideal conditions.

Dryland ecosystems are often dominated by annual plants that survive the dry

season as a permanent seed bank (Cohen 1966). Therefore, the community (i.e. the

seed bank) can be conveniently transplanted as a whole during the dry season without

any harm to the plants. In addition, by transplanting seed banks with their associated

soil, it is possible to evaluate plant communities' response to climate while preserving

soil abiotic and biotic interactions. To test the response of dryland annual plant

communities characterized by very different climates, we transplanted home soil with

seed bank among three sites situated along a steep aridity gradient in the Eastern

Mediterranean region, ranging from an arid to Mediterranean climate. In this region,

rainfall is the main limiting factor to plant growth (Ziv et al. 2014) and differs up to 5-

fold between the driest and wettest site(Holzapfel et al. 2006). The study sites used

for our reciprocal transplant also hosted a long-term climate manipulation experiment

(Tielbörger et al. 2014). This allowed for a comparison between long-term dynamics,

resulting from consistently imposed climate change, and the short-term responses

observed in our transplant experiment. Finally, we classified species based on their

climatic requirements, adopting the Climatic Niche Group approach (CNG; *sensu*

Bilton et al. 2016) that has been successfully employed for the species in our study

region (Bilton et al. 2016) and in other dryland ecosystems (Liu et al. 2018). By

identifying those species responsive to drier or wetter conditions, the CNG approach

provided us with testable predictions about shifts in community assembly across

climates within the reciprocal transplants.

We predicted that the community emerging from the reciprocal soil transplants

would be greatly determined by community origin, with less individuals emerging

from drier origins than wetter origins. Secondly, we hypothesized that the climate (i.e.

rainfall availability) at the transplant site would affect the emerging communities,

resulting in less biomass and less individuals emerging from community origins

exposed to drier climates. We also predicted that climate would select the emerging

community from the species pool of each origin in a predictable manner, with more

wet adapted species emerging in communities exposed to wetter climates, and more

dry-adapted species in drier climates.

Methods

Study area

This study was conducted in Israel at three fenced sites (area approximately 100 m

x 400 m) with respectively Mediterranean (M) semi-arid (SA), and arid (A) climate.

The three study sites share the same calcareous bedrock, southern aspect, altitude and

mean annual temperatures, so that they differ chiefly in mean and variance of annual

rainfall, and vegetation. The M site is located southwest of Jerusalem (N 31° 42' E

35° 3') at 620 masl, on Terra Rossa soil. The climate is characterized by 550 mm

average annual rainfall with 20% inter-annual variation. The SA site (N 31° 23' E 34°

54') is located in the northern portion of the Negev Desert near the city of Beersheba,

at 590 masl, on Light Brown Rendzina. Average annual rainfall is 270 mm with

approximately 30% inter-annual variation. The A site is located in the central Negev

near Sde Boqer (N 30º52' E 34º46') at 470 masl, on desert Lithosol. Average annual

rainfall amounts to 90 mm with 43% inter-annual variation (Holzapfel et al. 2006).

The plant communities at the three sites are semi-natural shrublands dominated by

Sarcopoterium spinosum (L.) Spach, and winter annuals (approx. 85% of all species)

that persist during summer in the form of dormant seed banks stored in the soil (Noy-

Meir 1973, Alon and Steinberger 1999). The species pool is overlapping among sites,

and annual plant cover amounts to 25% at the M site, 10% at the SA site, and < 1% at

the A site (Tielbörger et al. 2014).

Experimental set up

During the summer of 2010, we collected soil with seed bank from forty square

plots (20cm x 20cm, depth: 5cm) at the M and A sites and sixty plots at the SA site.

Within each site, plots were situated at least 20 cm apart from each other and away

from rocks and shrubs. Following Tomiolo et al. (2015) soil collected from each site

was pooled to produce a baseline community as there is substantial small-scale

heterogeneity in the seed bank (Siewert and Tielbörger 2010). The soil was stored in a

net-house at the University of Rehovot, Israel, where it experienced summer

temperatures necessary for breaking seed dormancy (Baskin et al. 1993). In

September 2010, twenty of the previously excavated plots at each site were randomly

selected and filled with home soil, while the remaining plots were filled with soil

from the closest away-from-home site (i.e. M site received M and SA soil; SA site

received M, SA and A soil; A site received SA and A soil, Supplementary Material

Appendix 1 Fig. A1). Transplanted soil was separated from the surrounding soil by a

layer of absorbent paper that provided initial isolation between soils, while not

impeding water percolation. After transplanting, we placed patches of organza (a thin

transparent fabric) over the surface of each plot to avoid contamination from seed

dispersal or seed predation (Petrů and Tielbörger 2008), and we removed them at the

time of germination.

Because the transplants were carried out during the dormant season we could

relocate the community of dormant seeds and soil biota with minimum damage. By

transplanting communities with their maternal soil we could test direct effects of

climate (e.g. decreasing rainfall) while preserving biotic interactions with

neighbouring plants and soil biota, which are also affected by the novel climate

(Emmett et al. 2004). At peak development (spring 2011), we recorded the identity

and number of individuals of the emerging species in each plot. In order to minimize

edge effects, we excluded plants growing in the outer 1 cm margin of each plot. After

species identification, aboveground biomass was collected, oven-dried at 70°C for 48

hours and weighed.

Unfortunately, the season of recording was very dry and the arid site received

only 30% of the average annual rainfall. Therefore, only a handful of seedlings of two

desert species (*Stipa capensis*, *Erodium touchyanum*) emerged at the arid site. As a

result, there was no home arid community to be compared to the transplants, and we

had to restrict our subsequent analyses to the reciprocal transplants between the SA

and M community origins.

Climatic Niche Groups (CNG)

Each species within the target communities was assigned to a Climatic Niche

Group (Bilton et al. 2016) classified by their distribution range in relation to rainfall.

A similar method has been employed for defining thermal niches of species in high

elevation and tundra habitats (Gottfried et al. 2012, Elmendorf et al. 2015), and

ideologically similar for Ellenberg values, which determine species habitat

requirements based on several abiotic parameters (Ellenberg 1974).

The rationale for the CNG grouping is that rainfall is the main driver of

community composition in the region, therefore species sharing similar climate

adaptations (approximated by the realized climatic rainfall niche) are likely to co-

occur in the same community by virtue of habitat filtering (García-Camacho et al.

2017). Species realized climatic niche values were derived as in Bilton et al. (2016).

For each single species the observed occurrences within Israel (distribution range)

were overlaid with mean annual rainfall climate data, and the mean value was taken

(obtained from BioGIS 2012, available at http://www.biogis.huji.ac.il/). Boundaries

between climatic niche groups spanned similar ranges of average annual rainfall

(approximately 130 mm) and resulted in four groups that ranked species with respect

to their hypothesized response to climate. Climatic Niche Group 1 (CNG1)

represented species associated with the lowest rainfall extremes of the gradient,

conversely CNG4 gathered species distributed in areas with high rainfall. Species

from all four CNGs were present in both communities (Supplementary Material,

Appendix 2 Table A1), but varied in their proportional representation at each site, and

could therefore be compared across sites and climates (Bilton et al. 2016).

Statistical analyses

We first analyzed how total density (number of individuals per plot), biomass,

species richness (number of species per plot), evenness (Simpson's evenness) and

diversity (Shannon-Wiener Index) varied in response to climate, community origin

and their interaction. In addition, we analyzed how the number of individuals

belonging to each climatic niche group per plot (i.e. CNG density) varied in response

to climate, community origin, with respect to the four-level categorical explanatory

variable CNG identity (i.e. CNG1 – CNG4), including all two-way and three-way

interactions. We applied generalized linear models with negative binomial

distribution to total individual, CNG density and species richness using the MASS

package (Venables et al. 2002) within the R software version 2.14 (R Development

Core Team 2014). Biomass, species diversity and evenness were analyzed using

linear models. To meet model requirements, evenness was log transformed and

biomass square root transformed. The significance of the models was assessed with a

Type 3 ANOVA, using the "car" package (Fox and Weisberg 2011). Post-hoc tests

were conducted using Tukey HSD test. Visual representation of the CNG density

interactions was done using log-ratios (log(SA density + 1) - log(M density +1)).

For testing how species composition varied with community origin and climate

we used Redundancy Analysis (RDA, (Legendre et al. 2011)) in the R package

'vegan' (Oksanen et al. 2015). The interaction term was included in a full model and

confirmed using a step-wise approach. The data were Hellinger transformed

(Legendre and Gallagher 2001) and scaled within plots. Significance of the model

was tested using 999 permutations. To test if species composition could be explained

by rainfall distribution range we regressed the resulting RDA 'species mean scores'

against the 'climatic niche value' of each species, both for individual species and for

the CNG classifications. Furthermore, we performed an RDA on the community-

weighted means (Garnier et al. 2007) using the species 'climatic niche value' as a

pseudo-trait.

Results

Overall, 97 species were recorded, among which 12.3% were grasses, 23.7%

legumes and 64% belonged to other families (Supplementary Material, Appendix 2

Table A1). In total, 68 species emerged from the semi-arid soil seed bank, 81 from the

Mediterranean origin, and 53 species were shared between the two origins. Fourteen

of these appeared in all four combinations of community origin and climate.

Diversity, richness and biomass

Total biomass, plant density, number of species and species diversity (Fig. 1b, 2a,

b, Table 1) were all significantly higher for communities of M origin rather than SA

origin. Additionally, for the M community origins, plant biomass, species richness,

diversity (Fig.1a, b, d) and total density (Fig 2.a, b) were significantly lower when

communities were exposed to the drier SA climates compared to their home climate,

whereas evenness remained the same across climates (Fig. 1c). For the SA

community origins, climate had no significant effect on any of the whole community

parameters total density (Fig. 2a, b, Table 2), biomass, species richness or diversity

(Fig.1, Table 1). Higher evenness was found in SA community origins exposed to M

climate compared to M community origins exposed to SA climate (Fig. 1c, Table 1).

Relative abundance of CNG across sites and community origins

Relative density of individuals changed considerably across community

origins and CNG groups (Table 2). In particular, the relative abundance of CNGs

shifted significantly across climates and community origins as indicated by the

significant two-way interactions (Table 2, Fig. 2a-d, Supplementary Material

Appendix 2 Table A2). In SA community origins, the mean abundance of individuals

belonging to dry CNGs (CNG 1and 2) was halved in M climate compared to SA

climate; on the other hand individuals belonging to CNG 4, the wettest adapted group,

were 6.5 times more abundant in M climate (Fig. 2a, c). In M communities, the shift

in CNG relative abundances was less strong but the relative hierarchical response of

the CNGs was in the same order (densities of CNG1 and 2 higher in SA climate;

CNG3 and 4 higher in M climate). The largest shift was seen in CNG4 that counted

twice as many individuals in home *vs.* away from home climate (Fig. 2b, d). The

different magnitude in CNG shifts across community origins and climates was

confirmed by a significant 3-way interaction (CNG x community origin x climate,

Table 2, Fig. 2e, f).

The RDA indicated four distinct communities emerging from the respective

treatments, with a significant effect of community origin and climate on species

assembly, as well as a significant interaction between these terms (Fig. 3a, b). Using

simple correlations we assessed which plots/species scores changed and had most

impact on each axis. We obtained three main RDA axes describing the species

composition. For plot mean scores: RDA1 (9.2% explained) was highly correlated to

overall differences between community origins, whereas the constrained RDA2

(3.3%; explained) and RDA3 (1.7%; explained) distinguished the climate x

community origin interaction term. For species mean scores, RDA1 was positively

correlated to species Climatic Niche values, and the correlation was positive but less

strong for RDA2 and RDA3 (Fig. 3c). Results were further validated by an RDA on

the community weighted mean traits using species Climatic Niche values as a trait,

and showed significant community origin and climate effects ($p < 0.05$). In

combination, these results suggest that rainfall niche strongly explained variation in

species composition across treatments.

Discussion

Our results revealed that climate played a large role in determining the species assemblages in our whole community transplant experiment, and that such changes were predictably related to possible species-specific climatic adaptations. One of our most notable findings was that large shifts in species composition occurred when transplants were exposed to vastly different climates, whereas total community parameters such as biomass, density, and diversity remained remarkably stable.

Perhaps unsurprisingly, and as predicted from previous observations across the rainfall gradient (Tielbörger et al. 2014), communities establishing from the Mediterranean (M) origin soil had higher individual density and total biomass than semi-arid (SA) origins. This is also similar to other dryland systems, where rainfall acts as a limiting resource (Guo and Brown 1997, Cleland et al. 2013). Consistently, all four CNGs had higher densities of individuals establishing from the M origin compared to SA origin. However, the wetter species groups (CNG 3 & 4) were proportionally more represented in M community origins, and the drier species groups (CNG 1 & 2) were more abundant in SA community origins. This, consistently with the findings of Bilton et al. (2016), suggests that CNGs are representative of species rainfall requirements and possibly climatic adaptations at the different sites. All communities rapidly matched the concomitant climate with shifts in community composition across climates that were strikingly well explained by hierarchical switches in relative CNG abundances. Plants establishing from both community origins showed a relative increase of species assumed to be most dry adapted when exposed to the drier (SA) climate. Accordingly, wet adapted species were more abundant in both communities when exposed to the wetter (M) climate. Therefore, CNGs revealed species responses to short-term climate effects in a predictable way,

suggesting potential short-term selection mechanisms that act on the communities in

response to yearly or extreme differences in rainfall.

Our most intriguing finding was that climate not only selected for predictable

species groups, but that the community origins differed greatly in the magnitude of

the community shifts. The SA origin, with the more variable rainfall regime – both

between and within years (Tielbörger et al. 2014) – also experienced the greater shifts

in community composition. This result matches other studies showing that

communities from drier climates often have higher inter-annual turnover of species

composition (Guo and Brown 1997, Cleland et al. 2013). Fascinatingly, the

magnitude of the compositional shifts between transplanted community origins was

inversely related to changes in total community parameters across climates. Namely,

in SA community origins total density, biomass and species richness remained

constant. Conversely, for the M origin transplants, which had smaller compositional

shifts, total density, biomass and richness decreased when exposed to the drier SA

climate. Large inter-annual variation in species abundances often leads to higher

community stability across time (Bai et al. 2004, Grime et al. 2008). Similar patterns

have been previously explained in plant community studies, albeit in a different

context, by the portfolio effect (Doak et al. 1998, Schindler et al. 2015). The portfolio

effect predicts that greater numbers of species in a community lead both

mathematically and ecologically to a greater chance of asynchronous relationships

forming year-to-year. Here we show, consistent with a previous study (Cleland et al.

2013), that in the community with the lower species richness (SA origin), greater

asynchrony and greater species turnover led to greater stability across climates. This

asynchrony is not random, but directly linked to species-specific climatic adaptations.

Thus, asynchronous shifts in relative abundance of species according to their climate

adaptations may allow for fast responses to year-to-year climatic variation in dryland

annual communities (Abbott 2017).

In the short-term, high species turnover may assure community stability, by

maintaining stable biomass and density, but in the long-term, such processes may also

lead to greater resistance and stability of dry (SA) communities to rainfall fluctuations

by favoring species adapted to more arid conditions. The high turnover is possible

without immediate loss of species because in dryland environments subject to large

year-to-year climatic variability, plants often display bet-hedging strategies such as

long-lived seed banks and seed dormancy that can buffer against inter-annual

fluctuations (Petrů and Tielbörger 2008, Tielbörger et al. 2012). Delayed germination

of dormant seeds during unfavorable years decreases the risk of extinction over time

and also promotes coexistence of species with different climatic requirements via

storage effects (Chesson and Grubb 1990, Pake and Venable 1995). Interestingly, the

findings from this short-term community transplant study are consistent with those of

a parallel long-term experiment conducted at the M and SA sites (Tielbörger et al.

2014), where community composition was monitored for 10 years in permanent plots

receiving respectively ambient rainfall, experimental drought (-30% rainfall) or

increased rainfall (+30% rainfall). Plant communities exposed to the long-term

climate manipulation treatments showed no detectable long-term effect on total

density, species richness and community biomass (Tielbörger et al. 2014). This

unexpected community resistance may be attributable to the large variability in

rainfall to which plant species in both community origins are pre-adapted (Tielbörger

et al. 2014, Tomiolo et al. 2015). Also at the two time-scales, selection of species in

relation to their CNG was observed (Bilton et al. 2016). However, while in the current

study we found stronger short-term selection for SA origin communities, long-term

selection patterns were found to be weaker in SA and stronger for M communities

(Bilton et al. 2016), suggesting that high inter-annual fluctuations result in stable

composition in the long run, whereas low species turnover across years results in long

term loss of wet-adapted species.

Finally, the observation of the community-level responses among climates

prompts an interesting parallel with well-known concepts of population ecology. High

environmental variability may select for a high degree of phenotypic plasticity (Sultan

1987, Pratt and Mooney 2013, Lazaro-Nogal et al. 2015, Spence et al. 2016). Thus,

when exposed to novel climates, individuals will display a large plasticity in adaptive

traits (Reed et al. 2011, Liancourt et al. 2015, Mathiasen and Premoli 2016). A better

match between the new trait value and the novel climate (Valladares et al. 2014)

confers an adaptive advantage and may result in stable fitness across home and away

environments, i.e. fitness homeostasis (Richards et al. 2006, Nicotra et al. 2010) and

may eventually assure the persistence of a species in changing climates. We propose

an analogy to that concept on a community level, where relative species abundances

(rather than trait values) shift across years (and sites) in order to better match the

current climate. Here, we show that the SA community origin, which is historically

exposed to larger year-to-year climatic variability (Holzapfel et al. 2006, Tielbörger et

al. 2014), experienced a greater compositional fluctuation (or "compositional

plasticity"), and this led to greater "community homeostasis" (i.e. maintenance of

stable density and biomass) across climates.

Our overall findings allow some careful conclusions about the potential

response of these communities to climate change. It should be noted that the variation

in rainfall experienced by the community origins in our study approximated the

extremes of climatic variability at each site, but exceeded the decrease in rainfall

predicted by climate change scenarios for the next 50-80 years (Smiatek et al. 2011,

Tomiolo et al. 2015). These results suggest that, as long as inter-annual climatic

fluctuations keep within the limits of climatic variability commonly experienced by

these communities, and rainy years that replenish the seed bank periodically occur,

wet adapted species will persist within the communities. However, with increasing

drought and unpredictability, communities are likely to experience species loss that

will affect primarily species with high rainfall requirements (Tielbörger et al. 2014,

Bilton et al. 2016). The similar results from the long-term experiment with our

reciprocal transplant indicate that the latter may be a powerful complement to long-

term field experiments. However, it should be noted that annual communities are

particularly suited for this approach and the same may not hold for long-lived

communities. Moreover, the fast response observed in our communities is uncommon

compared to temperate systems, where a lag between shifts in climatic conditions and

subsequent changes in community structure is often observed (Adler and Levine

2007, Jones et al. 2016). This suggests a higher resistance and resilience of dryland

ecosystems to extreme events compared to temperate ecosystems (Ruppert et al.

2015).

Declarations: KT and ST developed the experimental design. ST set up the

experiments and collected the data. MB and ST performed statistical analyses. ST

wrote the first draft of the manuscript and all authors contributed substantially to

revisions. We declare that we do not have conflicts of interest.

Acknowledgements: Jaime Kigel and Marcelo Sternberg provided logistic support.

The Hebrew University of Jerusalem (Rehovot) kindly provided material for

fieldwork. We thank Jake Alexander for providing comments on a previous version of

the manuscript. This study is part of the GLOWA Jordan River Project and was

funded by the German Ministry of Education and Research (BMBF). Further support

for MB and ST was obtained by the German Research Foundation (TI338_12-1;

TI338_11-1; and TI338_11-2; TI 338/15-1).

References

Abbott, R. E. et al. 2017. Portfolio effects, climate change, and the persistence of

small populations: analyses on the rare plant *Saussurea weberi*. — Ecology

98: 1071-1081.

Adler, P. B. and Levine, J. M. 2007. Contrasting relationships between precipitation

and species richness in space and time. — Oikos 116: 221-232.

Alexander, J. M. et al. 2015. Novel competitors shape species' responses to climate

change. — Nature 525: 515-518.

Alexander, J. M. et al. 2016. When climate reshuffles competitors: a call for

experimental macroecology. — Trends in Ecology and Evolution 31: 831-841.

Allison, S. D. et al. 2013. Microbial abundance and composition influence litter

decomposition response to environmental change. — Ecology 94: 714-725.

Alon, A. and Steinberger, Y. 1999. Response of the soil microbial biomass and

nematode population to a wetting event in nitrogen-amended Negev desert

plots. — Biol. Fertility Soils 30: 147-152.

Anderson, R. et al. 1982. Variability in the abundance of animal and plant species. —

Nature 296: 245-248.

Ayres, E. et al. 2009. Home-field advantage accelerates leaf litter decomposition in

forests. — Soil Biol. Biochem. 41: 606-610.

Bai, Y. et al. 2004. Ecosystem stability and compensatory effects in the Inner

Mongolia grassland. — Nature 431: 181.

Barbosa, E. R. M. et al. 2014. Short-term effect of nutrient availability and rainfall

distribution on biomass production and leaf nutrient content of savanna tree

species. — Plos One 9:

Baskin, C. C. et al. 1993. Annual seed dormancy cycles in two desert winter annuals.

— J. Ecol. 81: 551-556.

Bilton, M. C. et al. 2016. Climatic niche groups: A novel application of a common

assumption predicting plant community response to climate change. —

Perspect. Plant Ecol. Evol. Syst. 19: 61-69.

BioGIS 2012. Israel Biodiversity Information System [http://www.biogis.huji.ac.il/

]. p available at http://www.biogis.huji.ac.il/.

Blume-Werry, G. et al. 2016. Short-term climate change manipulation effects do not

scale up to long-term legacies: effects of an absent snow cover on boreal forest

plants. — J. Ecol. 104: 1638-1648.

Bonebrake, T. C. and Mastrandrea, M. D. 2010. Tolerance adaptation and

precipitation changes complicate latitudinal patterns of climate change

impacts. — Proceedings of the National Academy of Sciences of the United

States of America 107: 12581-12586.

Brown, J. H. et al. 2001. Regulation of diversity: maintenance of species richness in

changing environments. — Oecologia 126: 321-332.

Casper, B. B. and Castelli, J. P. 2007. Evaluating plant-soil feedback together with

competition in a serpentine grassland. — Ecol. Lett. 10: 394-400.

Chesson, P. L. and Grubb, P. 1990. Geometry, heterogeneity and competition in

variable environments. — Philosophical Transactions of the Royal Society of

London B: Biological Sciences 330: 165-173.

Cleland, E. E. et al. 2013. Sensitivity of grassland plant community composition to

spatial vs. temporal variation in precipitation. — Ecology 94: 1687-1696.

Cohen, D. 1966. Optimizing reproduction in a randomly varying environment. — J.

Theor. Biol. 12: 119-129.

Cubasch, U. et al. 1996. Estimates of climate change in Southern Europe derived from

dynamical climate model output. — Clim. Res. 7: 129-149.

De Dato, G. D. et al. 2006. Effects of warmer and drier climate conditions on plant

composition and biomass production in a Mediterranean shrubland

community. — Forest@ 3: 511-526.

491 Doak, D. F. et al. 1998. The statistical inevitability of stability-diversity relationships

492   in community ecology. — Am. Nat. 151: 264-276.

493 Doblas−Miranda, E. et al. 2015. Reassessing global change research priorities in

494   mediterranean terrestrial ecosystems: how far have we come and where do we

495   go from here? — Global Ecol. Biogeogr. 24: 25-43.

496 Ellenberg, H. 1974. Indicator values of vascular plants in central Europe. — Scripta

497   Geobotanica 9: 97.

498 Elmendorf, S. C. et al. 2015. Experiment, monitoring, and gradient methods used to

499   infer climate change effects on plant communities yield consistent patterns. —

500   Proc. Natl. Acad. Sci. U.S.A. 112: 448-452.

501 Emmett, B. A. et al. 2004. The response of soil processes to climate change: results

502   from manipulation studies of shrublands across an environmental gradient. —

503   Ecosystems 7: 625-637.

504 Fernandez-Going, B. M. et al. 2013. Climate interacts with soil to produce beta

505   diversity in Californian plant communities. — Ecology 94: 2007-2018.

506 Forey, E. et al. 2010. Does disturbance drive the collapse of biotic interactions at the

507   severe end of a diversity-biomass gradient? — Plant Ecol. 206: 287-295.

508 Fox, J. and Weisberg, S. 2011. An R Companion to Applied Regression. — Sage

509   Publications.

Garcia, R. A. et al. 2014. Multiple Dimensions of Climate Change and Their

Implications for Biodiversity. — Science 344:

García-Camacho, R. et al. 2017. Phylogenetic structure of annual plant communities

along an aridity gradient. Interacting effects of habitat filtering and shifting

plant–plant interactions. — Isr. J. Plant Sci. 1-13.

Garnier, E. et al. 2007. Assessing the effects of land-use change on plant traits,

communities and ecosystem functioning in grasslands: a standardized

methodology and lessons from an application to 11 European sites. — Ann.

Bot. 99: 967-985.

Gilman, S. E. et al. 2010. A framework for community interactions under climate

change. — Trends Ecol. Evol. 25: 325-331.

Gottfried, M. et al. 2012. Continent-wide response of mountain vegetation to climate

change. — Nature Clim. Change 2: 111-115.

Grime, J. P. et al. 2008. Long-term resistance to simulated climate change in an

infertile grassland. — Proceedings of the National Academy of Sciences of the

United States of America 105: 10028-10032.

Guo, Q. and Brown, J. H. 1997. Interactions between winter and summer annuals in

the Chihuahuan Desert. — Oecologia 111: 123-128.

Harrison, S. P. et al. 2015. Climate-driven diversity loss in a grassland community. —

Proceedings of the National Academy of Sciences of the United States of

America 112: 8672-8677.

Hobbs, R. J. et al. 2006. Novel ecosystems: theoretical and management aspects of

the new ecological world order. — Global Ecol. Biogeogr. 15: 1-7.

Holzapfel, C. et al. 2006. Annual plant-shrub interactions along an aridity gradient. —

Basic Appl. Ecol. 7: 268-279.

Jones, S. K. et al. 2016. Altered rainfall patterns increase forb abundance and richness

in native tallgrass prairie. — Scientific Reports 6: 20120.

Lazaro-Nogal, A. et al. 2015. Environmental heterogeneity leads to higher plasticity

in dry-edge populations of a semi-arid Chilean shrub: insights into climate

change responses. — J. Ecol. 103: 338-350.

Lazzaro, A. et al. 2011. Field-scale transplantation experiment to investigate

structures of soil bacterial communities at pioneering sites. — Appl. Environ.

Microbiol. 77: 8241-8248.

Legendre, P. and Gallagher, E. D. 2001. Ecologically meaningful transformations for

ordination of species data. — Oecologia 129: 271-280.

Legendre, P. et al. 2011. Testing the significance of canonical axes in redundancy

analysis. — Methods Ecol. Evol. 2: 269-277.

Liancourt, P. et al. 2015. Leaf-trait plasticity and species vulnerability to climate

change in a Mongolian steppe. — Global Change Biol. 21: 3489-3498.

Lindenmayer, D. B. et al. 2012. Value of long−term ecological studies. — Austral

Ecol. 37: 745-757.

Link, S. O. et al. 2003. A reciprocal transplant experiment within a climatic gradient

in a semiarid shrub-steppe ecosystem: effects on bunchgrass growth and

reproduction, soil carbon, and soil nitrogen. — Global Change Biol. 9: 1097-

1105.

Liu, D. et al. 2018. Species selection under long−term experimental warming and

drought explained by climatic distributions. — New Phytol. 217: 1494-1506.

Macel, M. et al. 2007. Climate vs. soil factors in local adaptation of two common

plant species. — Ecology 88: 424-433.

Maestre, F. T. et al. 2012. Plant species richness and ecosystem multifunctionality in

global drylands. — Science 335: 214-218.

Maranon, T. and Bartolome, J. W. 1993. Reciprocal transplants of herbaceous

communities between *Quercus agrifolia* woodland and adjacent grassland. —

J. Ecol. 81: 673-682.

Mathiasen, P. and Premoli, A. C. 2016. Living on the edge: adaptive and plastic

responses of the tree *Nothofagus pumilio* to a long-term transplant experiment

predict rear-edge upward expansion. — Oecologia 181: 607-619.

Miranda, J. D. et al. 2011. Climatic change and rainfall patterns: effects on semi-arid

plant communities of the Iberian Southeast. — J. Arid Environ. 75: 1302-

1309.

Nicotra, A. B. et al. 2010. Plant phenotypic plasticity in a changing climate. —

Trends Plant Sci. 15: 684-692.

Noy-Meir, I. 1973. Desert ecosystems: Environment and producers. — Annu. Rev.

Ecol. Syst. 4: 25–51.

Oksanen, J. et al. 2015. vegan: Community ecology package.

Pake, C. E. and Venable, D. L. 1995. Is coexistence of Sonoran desert annuals

mediated by temporal variablilty in reproductive success? — Ecology 76: 246-

261.

Parmesan, C. and Hanley, M. E. 2015. Plants and climate change: complexities and

surprises. — Ann. Bot. 116: 849-864.

Peñuelas, J. et al. 2007. Response of plant species richness and primary productivity

in shrublands along a north–south gradient in Europe to seven years of

582 experimental warming and drought: reductions in primary productivity in the

583 heat and drought year of 2003. — Global Change Biol. 13: 2563-2581.

584 Petrů, M. and Tielbörger, K. 2008. Germination strategies of annual plants under

585 changing climatic conditions: teasing apart local and regional effects. —

586 Oecologia 155: 717-728.

587 Pratt, J. D. and Mooney, K. A. 2013. Clinal adaptation and adaptive plasticity in

588 *Artemisia californica*: implications for the response of a foundation species to

589 predicted climate change. — Global Change Biol. 19: 2454-2466.

590 R Development Core Team 2014. R: A language and environment for statistical

591 computing. . — In: Computing, R. F. f. S. (ed).

592 Reed, T. E. et al. 2011. Interacting effects of phenotypic plasticity and evolution on

593 population persistence in a changing climate. — Conserv. Biol. 25: 56-63.

594 Richards, C. L. et al. 2006. Jack of all trades, master of some? On the role of

595 phenotypic plasticity in plant invasions. — Ecol. Lett. 9: 981-993.

596 Rinnan, R. et al. 2007. Fifteen years of climate change manipulations alter soil

597 microbial communities in a subarctic heath ecosystem. — Global Change

598 Biol. 13: 28-39.

599   Ruppert, J. C. et al. 2015. Quantifying drylands' drought resistance and recovery: the

600   importance of drought intensity, dominant life history and grazing regime. —

601   Global Change Biol. 21: 1258-1270.

602   Sala, O. E. and Lauenroth, W. K. 1982. Small rainfall events: an ecological role in

603   semiarid regions. — Oecologia 53: 301-304.

604   Sardans, J. et al. 2008. Warming and drought alter C and N concentration, allocation

605   and accumulation in a Mediterranean shrubland. — Global Change Biol. 14:

606   2304-2316.

607   Schindler, D. E. et al. 2015. The portfolio concept in ecology and evolution. — Front.

608   Ecol. Environ. 13: 257-263.

609   Shi, Z. et al. 2015. Evidence for long-term shift in plant community composition

610   under decadal experimental warming. — J. Ecol. 103: 1131-1140.

611   Siewert, W. and Tielbörger, K. 2010. Dispersal-dormancy relationships in annual

612   plants: putting model predictions to the test. — Am. Nat. 176: 490-500.

613   Smiatek, G. et al. 2011. High-resolution climate change simulations for the Jordan

614   River area. — J. Geophys. Res.-Atmos. 116:

615   Spence, L. A. et al. 2016. Short-term manipulation of precipitation in Mongolian

616   steppe shows vegetation influenced more by timing than amount of rainfall. —

617   J. Veg. Sci. 27: 249-258.

Sultan, S. E. 1987. Evolutionary Implications of Phenotypic Plasticity in Plants. — In:

Hecht, M. K. et al. (eds), Evolutionary Biology: Volume 21. Springer US, pp.

127-178.

Thompson, P. L. et al. 2015. Warming induces synchrony and destabilizes

experimental pond zooplankton metacommunities. — Oikos 124: 1171-1180.

Tielbörger, K. et al. 2012. Bet-hedging germination in annual plants: a sound

empirical test of the theoretical foundations. — Oikos 121: 1860-1868.

Tielbörger, K. et al. 2014. Middle-Eastern plant communities tolerate 9 years of

drought in a multi-site climate manipulation experiment. — Nature

Communications 5:

Tilman, D. and Downing, J. A. 1994. Biodiversity and stability in grasslands. —

Nature 367: 363-365.

Tilman, D. et al. 1998. Diversity-stability relationships: statistical inevitability or

ecological consequence? — The American Naturalist 151: 277-282.

Tomiolo, S. et al. 2015. Separating the role of biotic interactions and climate in

determining adaptive response of plants to climate change. — Ecology 96:

1298-1308.

Valladares, F. et al. 2014. The effects of phenotypic plasticity and local adaptation on

forecasts of species range shifts under climate change. — Ecol. Lett. 17: 1351-

1364.

Venables, W. N. et al. 2002. Modern Applied Statistics with S. . — Springer.

Waldrop, M. P. and Firestone, M. K. 2006. Response of microbial community

composition and function to soil climate change. — Microb. Ecol. 52: 716-

724.

Wetzel, P. R. et al. 2004. Use a reciprocal transplant study to measure the rate of plant

community change in a tidal marsh along a salinity gradient. — Wetlands 24:

879-890.

Williams, J. W. and Jackson, S. T. 2007. Novel climates, no-analog communities, and

ecological surprises. — Front. Ecol. Environ. 5: 475-482.

Wu, Z. T. et al. 2012. Biogeochemical and ecological feedbacks in grassland

responses to warming. — Nat. Clim. Chang. 2: 458-461.

Ziv, B. et al. 2014. Trends in rainfall regime over Israel, 1975–2010, and their

relationship to large-scale variability. — Regional Environmental Change 14:

1751-1764.

Figures and tables

656
657   **Table 1: Type III ANOVA table results for the models applied to individual density, species**

658   **richness, species diversity, evenness and total biomass. Lines correspond to response variables**

659   **and columns to explanatory variables of each model. In each column the first value represent**

660   **Chi-square test values and the second the p-value. Probability values for significant terms are**

661   **reported in bold.**

	Origin	**Climate**	**Origin x Climate**
Density	312.86; **<0.001**	0.12; 0.72	0.59; 0.44
Species richness	281.29; **<0.001**	12.99; **<0.001**	0.085; 0.77
Evenness	11.96; **<0.001**	7.18; **0.009**	0.028; 0.86
Species diversity	130.41; **<0.001**	10.40; **0.001**	1.36; 0.24
Total biomass	103.86; **0.001**	2.34; 0.13	4.66; **0.03**

Table 2: Type III ANOVA table of results for Generalized Linear Models determining the effect

of community origin, climate and Climatic Niche Group (CNG) on plant community densities

establishing from seed banks. N=320.

Effect	DF	LR Chi-sq.	P-value
Origin	1	236.93	**<0.00001**
Climate	1	2.35	0.12540
CNG	3	343.21	**<0.00001**
Origin x Climate	1	0.07	0.78714
Origin x CNG	3	83.55	**<0.00001**
Climate x CNG	3	50.09	**<0.00001**
Origin x Climate x CNG	3	16.85	**0.00076**

 Figure Captions:

Figure 1: Mean ±1SE A) species richness (i.e. number of species per plot), B) total biomass, C)

evenness, D) diversity, of annual plant communities established from two community origins

('M' Mediterranean; 'SA' Semi-Arid) grown in two sites (i.e. climates: 'M' Mediterranean; 'SA'

Semi-Arid).

Figure 2: Impact of community origin, climate and Climatic Niche Group (CNG) on plant

densities establishing from seed banks in home *vs*. away-from-home climate. Each Climatic Niche

Group (CNG) aggregates species with similar climatic adaptation, ranging from dry climates

(CNG 1) to wetter climates (CNG 4). Fig. 2 a, b: Total mean individuals density across climates is

broken down according to CNG relative abundance. Fig. 2 c-f: Differences in relative densities of

CNG groups are expressed as log ratios, where positive values indicate higher CNG relative

abundances in SA community origins or sites, whereas negative values indicate higher CNG

relative abundances in M community origins and climate. Community composition shifts are

represented across community origins (2c), climates (2d) and the combination of the two (2 e,f).

Figure 3: Redundancy Analysis (RDA) of species compositional change in community origins

emerging in home *vs*. away-from-home climates. Indicated are the 95% confidence intervals for

the groupings/categories (ellipses). Fig. 3 a, b represents the plot centroids of each community

origin-climate combination. Lines are vectors from the centre of a category to each site score

(points). RDA-axis 1: correlated to distance between origins, RDA-axis 3: the effect of climate on

SA community origins, and RDA-axis 2 the effect of climate on M community origins. In red: SA

community origins - SA climate; yellow: SA origins - M climate; blue: M origins- M climate;

green: M origins -SA climate. Fig. 3c represents the species centroids for each species group.

Lines are vectors connecting the centre of each group with species scores. In red: CNG1, yellow:

CNG2, green: CNG3, blue: CNG4.

716
717   **Figure 1**
718

Raw Density

2-way interactions

3-way interaction

Figure 2

Figure 3

748
749
750
751