Functional connectivity with short-term dynamics explains diverse patterns of excitatory spike transmission *in vivo*

Abed Ghanbari[1], Naixin Ren[2], Christian Keine[3], Carl Stoelzel[2], Bernhard Englitz[4], Harvey A. Swadlow[2], and Ian H. Stevenson[1,2]

[1] *University of Connecticut, Department of Biomedical Engineering*
[2] *University of Connecticut, Department of Psychological Sciences*
[3] *Carver College of Medicine, Iowa Neuroscience Institute, Department of Anatomy and Cell Biology, University of Iowa*
[4] *Radboud University, Donders Institute for Brain, Cognition and Behavior, Department of Neurophysiology*

Abstract

Fast information transmission in neural networks is heavily influenced by short-term synaptic plasticity (STP), and the type and timescale of STP varies by cell-type and brain region. Although STP has been widely characterized *in vitro* from recordings of postsynaptic potentials or currents, characterizing STP in *in vivo* in behaving animals is difficult due to the lack of large-scale intracellular recordings. Here, we use paired extracellular observations to estimate the short-term dynamics of synaptic transmission from spikes alone. We introduce an augmented generalized linear model (GLM) that includes a dynamic functional connection as well as several, non-synaptic factors that alter spike transmission probability. Our model captures the diverse short-term dynamics of *in vivo* spike transmission at identified synapses and accurately captures the effects of local pre- and postsynaptic spike patterns. We applied this model to large-scale multi-electrode recordings to describe stimulus-dependent shifts in spike transmission and cell-type specific differences in STP.

Introduction

Neural information processing is largely governed by synapses and their dynamics [1,2]. Short-term synaptic plasticity (STP) alters synaptic transmission on timescales from a few milliseconds to several seconds depending on the sequence of presynaptic spiking. Presynaptic STP arises from a mixture of two main processes: depletion of neurotransmitter-containing vesicles, which causes depression, and the elevation of residual Ca^{2+} in the presynaptic terminal, which causes facilitation by increasing vesicle release probability [3]. This can be observed in intracellular recordings where, following repetitive stimulation of the presynaptic terminal, the amplitudes of postsynaptic potentials (PSPs) or currents (PSCs) will either decrease (depression) or increase (facilitation) [3,4]. The degree of STP differs depending on the pre- and postsynaptic cell type [5] and brain region [6,7]. Functionally, STP can act as a temporal filter [8], can allow neural circuits to specialize for specific tasks [9,10], and may also underlie gain control [11], network stability [12], and long-term synaptic plasticity [13]. Here we focus on understanding how STP-induced changes

in PSP/PSC amplitudes shape postsynaptic spiking. *In vivo* studies have shown that postsynaptic
spiking probability, similar to the amplitude of PSP/PSCs, depends on the recent history of
presynaptic spiking [14,15]. Just as PSP/PSCs show diverse patterns of depression and facilitation,
the postsynaptic spiking probability also appears to have complex patterns depending on the brain
region and cell-types [16]. However, postsynaptic spiking probability is modified by many
additional variables besides STP at a single synaptic input. Here we aim to understand how the
pattern of presynaptic spiking activity and short-term synaptic plasticity shape postsynaptic
spiking probability.

To do so, we use simultaneously recorded pre- and postsynaptic spiking activity to detect and study
putative monosynaptic connections. When an excitatory, monosynaptic connection is present,
cross-correlations between the spiking of a pre- and postsynaptic neuron often show a rapid,
transient increase in postsynaptic spikes following the presynaptic spike. This occurs at an interval
reflecting the presynaptic axonal conduction time plus the synaptic delay (usually < 4 ms) [17,18].
However, this cross-correlation is not static. Previous studies have found that the cross-correlation
often differs for presynaptic spikes that are part of a burst compared to isolated spikes [14]. Spike
transmission probability appears to depend on the timing of previous presynaptic spikes, and one
factor influencing spike probability may be STP [19,20]. For example, depressing synapses would
have more reliable synaptic transmission in response to isolated presynaptic spikes following long
inter-spike intervals (ISIs) compared to shorter intervals (in bursts) [14,16]. On the other hand,
facilitating synapses would show a stronger response to presynaptic spikes following shorter ISIs
(bursts) compared to the presynaptic spikes following longer ISIs (isolated) [21]. By looking at
the corresponding cross-correlograms from a subset of presynaptic spikes with specific ISIs,
previous studies have found highly diverse, non-monotonic spike transmission patterns for
different synapses [16]. This diversity in patterns of spike transmission probability, however, is
not solely attributable to STP. When two presynaptic spikes occur in close succession, the
membrane time-constant may cause postsynaptic potentials (PSP) to sum and increase the spike
probability even if the individual PSPs were sub-threshold [22]. Moreover, the history of
postsynaptic spiking also affects spike probability such that even if the PSP is strong, it may not
trigger a spike if it falls within the refractory period or during an after-hyperpolarization (I_{AHP})
current [23]. Finally, slow fluctuations in the overall excitability of the postsynaptic neuron, due
to neuro-modulation, for instance, could also change synaptic transmission probability [24]. In
different synapses the degree that each factor contributes varies and leads to diverse patterns of
postsynaptic spike transmission probability.

The overall correlation structure in spiking data can often be estimated by generalized linear
models (GLMs) [25,26]. However, previous models have treated these interactions as static, and,
thus, cannot capture dynamic changes in spike transmission probability. Here we extend these
GLMs to describe dynamic interactions between neurons and account for the diversity of spike
transmission patterns [26–28]. For each individual presynaptic spike, our model aims to predict
postsynaptic spikes accounting for the postsynaptic neuron's baseline firing rate, slow fluctuations

of the postsynaptic firing rate, the effect of postsynaptic spiking history, and a coupling term
affected by synaptic summation and short-term synaptic plasticity. The conditional intensity of our
model provides estimates of postsynaptic spiking probability following every single presynaptic
event based on the previously observed sequence of pre- and postsynaptic spiking. The split cross-
correlogram only describes the average response conditioned on the ISI preceding the most recent
presynaptic spike. By using a model-based approach we can incorporate the full sequence of
presynaptic spikes beyond the most recent one, explicitly account for factors such as postsynaptic
history, and link the observed patterns of spike transmission to the underlying dynamics of vesicle
depletion and release probability.

To evaluate the model, we first examined its ability to capture the observed patterns of spike
transmission probability for three well-studied, strong putative synapses using pre- and
postsynaptic spike observations from: 1) a pair of neurons in the mouse thalamus, 2) an auditory
nerve projection onto a spherical bushy cell (ANF-SBC) in the gerbil, and 3) a neuron in
ventrobasal (VB) thalamus of the rabbit projecting to a putative fast-spike inhibitory neuron in
primary somatosensory (S1) barrel cortex (VB – Barrel). Short-term synaptic dynamics of this
latter system have been extensively characterized *in vivo* [14,29,30]. Similarly, ANF-SBC
synapses have been extensively studied in previous experiments and are well-characterized *in vitro*
[31–33]. Using the auditory brainstem connection, we explore how synaptic transmission
probability changes depending on the stimulus. Our result suggests that a simplified model,
without considering short-term plasticity, is insufficient to explain how patterns of spike
transmission change as the pattern of presynaptic input changes. Finally, we apply our model to
spiking data from a large-scale, multi-electrode array recorded from multiple areas in an awake
mouse. Here we investigate the STP dynamics in putative synapses from excitatory neurons onto
two putative inhibitory subtypes (e.g. FS: fast-spiking, RS: regular-spiking). We find that these
two types of connections have distinct patterns of spike transmission, where excitatory-FS
connections appear to be slightly more depressing than excitatory-RS connections. Together, these
results illustrate the diversity of spike transmission patterns *in vivo* and present one potential
approach to studying short-term synaptic dynamics in behaving animals.

Most previous approaches to describing interactions between neurons using spiking activity have
focused on static functional connectivity. These models improve both encoding and decoding
accuracy and have been shown to capture physiological network structure in some cases [34]. Here
we model dynamic functional connectivity where the effect of each presynaptic spike on the
probability of postsynaptic spiking depends on the previously observed sequence of presynaptic
spiking. This augmented GLM can be directly compared with the observed spike transmission
probability and also allows us to disentangle the contributions of short-term synaptic plasticity,
synaptic summation, presynaptic firing rate fluctuations, and spike history. Moreover, we find that
modeling dynamic functional connections allows us to better predict postsynaptic responses
compared to the static models. Since modeling static functional connectivity can improve decoding
[25,26,35], modeling dynamic functional connectivity may improve decoding further as well. As

multi-electrode arrays improve, and the number of simultaneously recorded neurons increases,
models of dynamic functional connectivity may provide insight into not just network structure, but
also the extent of depression or facilitation in these networks, as well as differences in network
dynamics across multiple brain areas and under different behavioral conditions. Here, our findings
suggest that, at least in some cases, *in vivo* spike transmission dynamics differ substantially for
different stimuli and different cell-types.

Results

Short-term synaptic plasticity directly affects synaptic information processing by altering the
amplitude of presynaptic currents [2], but in most neural systems it remains unclear how these
presynaptic effects translate to altered postsynaptic spike probability. Postsynaptic spiking is
affected by many factors including short-term plasticity, postsynaptic spike history, summation of
PSPs, and slow fluctuations in presynaptic firing rate. Here we developed a statistical model that
includes each of these factors and allows their effects to be estimated solely using pre- and
postsynaptic spiking activity. In this approach, working with spikes rather than PSC/PSPs enables
us to understand the short-term changes in synaptic transmission probability *in vivo* where large-
scale intracellular recordings have not been achieved.

Spike transmission probability varies strongly as a function of presynaptic ISIs

Here we define spike transmission probability as the probability of postsynaptic spiking in a
window shortly after each presynaptic spike. One conventional approach to study spike
transmission and changes in transmission probability are cross-correlograms. Cross-correlograms
of excitatory monosynaptic connections show a rapid, transient increase in the postsynaptic spiking
probability shortly (usually < 4ms, although this depends on the presynaptic axonal conduction
delay) after the presynaptic spike [17]. The timing and shape of the cross-correlogram depends on
the synaptic delay, the strength of the connection and varies between synapses. However, in the
overall cross-correlogram since all presynaptic ISIs are averaged, the dependence of spike
transmission probability on the presynaptic ISIs remains hidden (Fig. 1A). To determine the effect
of presynaptic ISI on spike transmission probability we can calculate the cross-correlogram for a
subset of presynaptic spikes with a specific ISI. and previous studies showed that transmission
probabilities can vary for different ISIs within the same synapse [14,16]. Moreover, the short-term
dynamics of spike transmission probability can differ for different synapses as a function of
presynaptic ISIs. To illustrate this diversity, we examined three strong synapses from three distinct
neural systems: (i) a pair of neurons in thalamus in a mouse, (ii) a projection from ventrobasal to
somatosensory barrel cortex (VB-Barrel) in a rabbit, and (iii) the auditory nerve fiber to spherical
bushy cell projection in a gerbil (ANF-SBC). Although the presynaptic neurons have diverse ISI
distributions (Fig. 1B), splitting the spikes into ISI quantiles and calculating the correlogram for
each quantile, demonstrates how postsynaptic responses differ following short and long
presynaptic ISIs. For the pair of neurons in thalamus, spike transmission has a higher probability
at short and long intervals and a lower probability for mid-range ISIs. For VB-Barrel transmission

probability is higher for longer ISIs, while for ANF-SBC the highest transmission probability
occurs at intermediate intervals (Fig. 1C).

Fig. 1: Spike transmission probability depends on the presynaptic ISI and differs between synapses.
A) Cross-correlograms between pre- and postsynaptic spiking show an increase in the postsynaptic spike
count (or probability) after a short latency, indicative of a monosynaptic connection. Efficacy (Eff.) for
each synapse denotes the ratio between numbers of postsynaptic spikes in the synaptic peak (horizontal
bars) corrected for the baseline number of expected postsynaptic spikes to number of presynaptic spikes.
B) Inter-spike interval distributions (log-scale) for the presynaptic neuron for three different synapses. The
distributions are color-coded into 5 quantiles with equal number of presynaptic spikes. **C)** For each ISI
quantile, we calculate a separate cross-correlogram. Colors correspond to (B) going from shorter
presynaptic ISIs (left) to longer ISIs (right). Note that both the baseline firing rate and the synaptic peak for
each connection change as a function of presynaptic ISI. Solid lines overlaying the cross-correlograms
illustrate model fits used to estimate the synaptic effect and the smooth baseline.

The shape of spike transmission patterns depends on multiple factors

In synapses exhibiting short-term synaptic plasticity the postsynaptic response after each
presynaptic spike changes according to the recent history of presynaptic spiking [28,36]. However,
besides synaptic dynamics there are additional factors that alter spike timing. At short presynaptic
ISIs, membrane potential summation can lead to larger PSPs and increased spike probability, even
in absence of short-term synaptic plasticity [22]. The spiking nonlinearity and the history of
postsynaptic spiking can also alter how a given pattern of presynaptic input is transformed into
postsynaptic spiking [26,37].

To illustrate how STP, synaptic summation, and postsynaptic history interact to create the observed
spike transmission pattern we simulated from a simplified rate model with linear voltage
summation, short-term plasticity, a soft spiking nonlinearity, and an after-hyperpolarization (Fig.

2). Similar to experimental data, the spike transmission probability in this simplified model
depends on the presynaptic ISI as well as the type of STP. For depressing synapses, the spike
transmission probability increases for longer presynaptic ISIs while for facilitating synapses it
increases for mid-range ISIs [28,36]. Independent of STP type, PSPs sum at short ISIs (Fig. 2A).
However, the exact shape of transmission probabilities also depends on the strength of the synapse
and, possibly, the history of postsynaptic spiking. An after-hyperpolarization current following
each postsynaptic spike, for instance, can briefly decreases the probability of subsequent spikes.
In our simulation, we find that "spike interference" from previous postsynaptic activity can
counteract membrane potential summation (Fig. 2B). This type of postsynaptic spike interference
generally decreases the spike probability for shorter presynaptic ISIs, but the magnitude of this
decrease depends on the synaptic strength and type of STP (Fig. 2C). These simulations illustrate
how the pattern of spike transmission probability results from the complex interaction between the
membrane potential, the spike nonlinearity, the post-spike history, and short-term synaptic
plasticity.

Fig. 2: A simulation of a simplified rate model shows how spike transmission probability
depends on multiple factors. A: For different types of short-term synaptic plasticity, postsynaptic
summation increases the amplitudes of the postsynaptic potentials (PSP) at shorter ISIs. Lines
denote the membrane potential of a postsynaptic neuron in a simplified model as it responds to
short (dark traces) and long (light) paired presynaptic pulses. Relative amplitudes of excitatory
postsynaptic potentials increase under the simplified model depending on the type of STP (right
panel). **B:** Spike generation changes with synaptic strength. In this paired-pulse stimulation
paradigm, stronger synapses are more likely to generate a spike following the first presynaptic
impulse which can then decrease the spiking probability following the second impulse if there are
post-spike history effects. As in (B) traces denote postsynaptic membrane potential responses to
short (dark) and long (light) presynaptic ISIs. Dashes denote example postsynaptic spiking, with
"spike interference" occurring for strong synapses and short ISIs. **C:** The pattern of spike
transmission probability under the simplified model changes depending on the type of STP, the
coupling strength, and presence of post-spike interference. Dashed lines show transmission
probability without interference from previous postsynaptic spikes, while solid lines show how
post-spike history effects can decrease the spike transmission probability.

Spike transmission patterns are diverse across regions and species

210
The combination of these factors could be one explanation for to the diversity of spike transmission
patterns in experimental data. To account for STP, postsynaptic history effects, and slow changes
of firing rate we extend a previously developed GLM framework for static functional connections
[26] to include short-term dynamics. In the previous, static GLM the probability of postsynaptic
spiking is modeled as a linear combination of a baseline firing rate parameter, a post-spike history
filter to capture the postsynaptic spike dynamics, such as refractoriness and burstiness, and a
coupling filter describing the fixed influence of presynaptic spikes. The sum of these effects is
then passed through a spiking non-linearity [26]. In our extended model we added a linear term
that allows changes in excitability of the postsynaptic neuron as a function of the presynaptic firing
rate (timescale >1min) and allow the coupling term to change for each presynaptic spike according
to the Tsodyks and Markram (TM) model of STP [36]. We fit the parameters of this TM-GLM
using only the pre- and postsynaptic spike observations and obtain parameters for each effect using
approximate maximum likelihood estimation (see Methods). This provides estimates of the history
and coupling filters, as in a static GLM, as well as additional parameters for the dynamical synapse
(TM model) including facilitation, depression, membrane time constants, and release probability.
Given these parameters, the model estimates the postsynaptic spiking probabilities following each
observed presynaptic spike and predicts spike transmission probabilities in response to arbitrary
patterns of presynaptic inputs.

After fitting the model to real pre- and postsynaptic spike-trains, we compared its behavior to
experimentally observed patterns of spike transmission probability. In particular, we compare
peaks in the split cross-correlograms to the average model prediction for the same sets of
presynaptic spikes (see Methods). We find that our model is flexible enough to explain the changes
in synaptic transmission probability observed in spiking statistics for all three synapses above
(Fig. 3A). Moreover, using the model-based approach, the contributions of each model component
can be disentangled. Our results suggest that the pattern of spike transmission probability for the
thalamus connection is dominated by a combination of membrane potential summation and short-
term depression. Although depression decreases spike transmission probability at shorter ISIs,
membrane summation acts to increase postsynaptic spiking. The ANF-SBC synapse, in contrast,
shows an increase in spike transmission probability for a medium range of ISIs that is explained
by a model dominated by short-term facilitation. Lastly, the VB-Barrel connection shows a higher
postsynaptic response for spikes following longer ISIs (isolated) that is explained by the model as
an effect of short-term synaptic depression.

In addition to separating the factors affecting spike transmission, the model also improves the
prediction of postsynaptic spike timing. To evaluate how spike prediction accuracy is influenced
by STP, we compare the prediction of postsynaptic spiking activity after each presynaptic spike
for our model with a static model containing all components except STP. In all three datasets, a
model with short-term synaptic plasticity provides substantially better predictions of the
postsynaptic spiking activity, assessed by Receiver Operating Characteristics (ROC) curves. For

the model with short-term synaptic plasticity accuracies were AUC=0.76, 0.70, and 0.79 for the
Thalamus pair, VB-Barrel, and ANF-SBC connections, respectively; compared to a model without
STP where the model accuracies were AUC=0.54, 0.48, and 0.56.

In our model, STP is described by two coupled differential equations with five parameters: $\theta_{stp} =$
$\{\tau_d, \tau_f, U, f, \tau_s\}$ (see Methods). Here we estimate values for depression, facilitation, and
membrane time-constants along with release probability, U, and magnitude of facilitation, f, (Fig.
3B). Our result for the thalamus pair shows a higher release probability and depression time-
constant with a larger membrane time constant. The VB-Barrel connection shows a higher
depression compared to facilitation time constant with a lower membrane time constant. The ANF-
SBC synapse shows a lower release probability compared to the other two connections and a lower
depression and membrane time constant. Although here we estimate these parameters from pre-
and postsynaptic spiking alone, they could also be estimated from intracellular measurements [38].
We are not aware of any *in vivo* experiments that measured depression or facilitation time-
constants for these systems. However, previous *in vitro* studies found a wide range of paired-pulse
ratios (0.3 to 0.9) in thalamocortical projections [39], consistent with the depressing VB-Barrel
synapse here. Additionally, *in vitro* observations of ANF-SBC connections report depression time-
constants on the order of 2-25ms in response to a 100 Hz stimulus train [40,41]. These previous
estimates are substantially faster than the time-constants estimated by the TM-GLM for the ANF-
SBC connection here. However, as mentioned in [40], different patterns of presynaptic input (e.g.
regular, Poisson, natural) can result in different time constants, which makes it difficult to compare
in vitro and *in vivo* STP parameters directly. One parameter that may be more readily comparable
across preparations is the membrane time-constant. We find that the estimated membrane time-
constant from the TM-GLM for the thalamus pair is consistent with thalamus relay cells observed
intracellularly (12.2 ± 1.1 ms (n=8)) [42], and the estimated membrane time-constant for ANF-
SBC is approximately consistent with *in vitro* measurements (1.05 ± 0.09 ms) [40], as well.

Previous work modeling intracellular recordings suggests that the full TM model may not be
necessary to explain STP at some, purely depressing synapses [38]. Therefore, we explored how
simplified TM models of STP, with fewer parameters, compare with the full model using the
Akaike information criterion (AIC; see Methods). AIC evaluates model accuracy (log-likelihood)
penalized by the number of parameters and determines if a simplified model with fewer parameters
is preferred over a more complex model. We compare the full model to five reduced models: 1) a
model with only integration, without dynamic release probability and resources ($\tau_d, \tau_f = 0$ and
$f, U = 1$), 2) a facilitation only model ($\tau_d = 0$), 3) a depression only model ($\tau_f = 0$), 4) a 3-
parameter TM model where the magnitude of facilitation is fixed ($f = U$), and 5) a full TM model
without resetting integration when a postsynaptic spike occurs ($\pi_i = 1$). For the thalamus pair and
VB-Barrel, a model with fixed magnitude of facilitation ($f = U$) performs better while for the
ANF-SBC connection the full model gives a better prediction. The full TM model performs well

in all cases, but, for some synapses, as previous results suggested [38], there may be ambiguity with parameter identifiability where many parameter settings explain the data.

Fig. 3: Model predictions of spike transmission dynamics. A: Spike transmission patterns are diverse across different connections. For three different connections (a pair in thalamus, ventrobasal thalamus to somatosensory cortex, auditory nerve fiber to spherical bushy cell) transmission patterns are explained by a combination of different factors. For each synapse, top panels show the presynaptic ISI distributions (log-spaced). In the second row, the observed spike transmission probability (red data points) and model predictions (blue with 95% confidence bands). We then used the estimated TM parameters for each synapse and simulated responses to for paired presynaptic pulses. Blue curves denote the PPRs of the full model, and gray lines denote PPRs for a model without synaptic summation. In the fourth row, we evaluate how accurately the TM-GLM can prediction individual postsynaptic transmission events. For each individual presynaptic spike, we compare the model transmission probability with the observed binary outcome. ROC curves show the prediction accuracy for the TM-GLM (blue) and a standard GLM without STP (orange). **B:** Estimates for the four STP parameters of the model for each synapse. Each dot represents estimation from a distinct bootstrap sample. **C:** Model comparison for 6 different models (Akaike information criteria relative to a model without plasticity). Models: 1) Integration only ($\tau_d, \tau_f = 0$ and $f, U = 1$), 2) Facilitation only ($\tau_d = 0$), 3) Depression only ($\tau_f = 0$), 4) 3-parameter TM ($f = U$), 5) 4-parameter TM without resetting integration ($\pi_i = 1$), 6) 4-parameter TM.

Recent patterns of pre- and postsynaptic spiking shape the synaptic transmission probability

Although previous studies have focused largely on how spike transmission probability varies as a function of the single preceding presynaptic ISI, synaptic dynamics depend on the full sequence of presynaptic spiking. Unlike *in vitro* experiments where the state of the synapse can, to some

extent, be controlled before studying responses to a specific presynaptic pattern, *in vivo*
measurements of spike transmission can be heavily influenced by higher-order correlations
between successive ISIs [29]. Additionally, it is difficult to assess the effects of multi-spike
patterns empirically by splitting the correlograms, since the number of observations for any given
presynaptic spike pattern rapidly decreases with the number of spikes in the pattern. Here we
examine how spike transmission depends, not just on the preceding presynaptic ISI, but on triplets
of spikes. We compare the empirically observed spike transmission probability following triplets
to the estimated spike transmission probability from the TM-GLM. Then, after fitting the TM-
GLM, we simulate postsynaptic responses to isolated, local patterns of spikes and determine to
what extent the observed spike transmission patterns are influenced by higher-order correlations
between successive ISIs.

First, in addition to the timing of the two preceding presynaptic spikes (ISI_1), we split correlograms
based on the timing of the three preceding presynaptic spikes using both ISI_1 and ISI_2. Since the
TM-GLM provides estimates of the post-synaptic spike probability following every presynaptic
spike, we can split both the data and model fits the same way (Fig. 4A). We find that the spike
transmission patterns clearly depend on the triplet patterns of presynaptic spikes. That is, the spike
transmission probability is influenced by both ISI_1 and ISI_2, and their interaction differed between
synapses, as expected from the TM-GLM model. However, similar to the descriptions of spike
transmission as a function of ISI_1, the TM-GLM accurately captures the patterns of spike
transmission for triplets of presynaptic spikes at those three synapses. In the thalamus pair, spike
transmission was dominated by ISI_1, and the effect of ISI_2 appears to be weak or, at least, doesn't
appear to be monotonic. Spike transmission at the VB-Barrel connection depends strongly on both
ISI_1 and ISI_2, with higher spike transmission probability for longer ISI_2, consistent with recovery
from depression. Lastly, for the ANF-SBC connection, transmission probabilities decrease for
shorter ISI_2, but there also appears to be a strong interaction between ISI_1 and ISI_2, where
transmission probability is high for multiple combinations of these two intervals (e.g. intervals of
10ms then 100ms and intervals of 100ms then 10ms both result in high probability transmission).

To examine to what extent the empirical observations of spike transmission are affected by higher-
order correlations between successive ISIs, we again use the estimated parameters in the TM-GLM
to simulate postsynaptic responses to hypothetical, isolated triplets of presynaptic spikes. In these
simulations we fix the post-spike history effect and the excitability in the model to their average
values from model fits, and we fix the initial STP state (initial values of R and u in TM model) for
the first spike in triplets to the average R and u values from the model fits. In experimental data,
the initial state of the pre- and postsynaptic neurons before the triplets occur can wildly differ
between different values of ISI_1 and ISI_2. By simulating, we can compare the influence of different
triplets (ISI_1 and ISI_2) when the pre- and postsynaptic neurons start at the same state. Here we find
that for the thalamus pair, although the empirical data showed no clear effect for ISI_2, the simulated
spike transmission probability increases with short ISI_2, consistent with strong synaptic
summation. One reason that this effect may be masked in the empirical transmission probabilities

is that post-spike history effects could act to decrease the probability of future postsynaptic spikes.
For the VB-Barrel simulations, we find that short ISI_2 decreases transmission probability,
consistent with the empirical transmission patterns, although less pronounced. Serial correlations
in the sequence of presynaptic spikes (such as long bursts) could act to accentuate the depression
in the empirical observations beyond what we see with the simulated responses to isolated triplets.
Finally, for the ANF-SBC, although the empirical transmission probability showed decreased
transmission for short ISI_2, the simulated responses to isolated patterns have increasing
transmission at short ISI_2 (due to synaptic summation). This difference is likely due to the post-
spike history effects, which has been fixed for the simulations, but can have a large effect in the
experimental data. Since the overall efficacy of this synapse is quite high (>0.7), is likely that a
postsynaptic spike follows the first or second presynaptic spike which then influences the response
to the third spike.

To better understand the effects of post-spike history, we examined how the postsynaptic spiking
history changes the spike transmission patterns with a similar approach. In addition to splitting the
correlograms based on ISI_1, we also split based on the previous postsynaptic ISI, ISI_{post} (Fig. 4B).
Here, as with the triplets of presynaptic spikes, we find that the spike transmission patterns depend
on the triplet patterns of 2 pre- and 1 postsynaptic spike, and the TM-GLM accurately captures the
patterns of spike transmission at our three synapses (Fig. 4B). Here, for both thalamus and VB-
Barrel pairs, synaptic transmission probability decreases after a long postsynaptic ISI for all values
of ISI_1. In contrast, the ANF-SBC connection shows decreased transmission probability at short
postsynaptic ISIs.

As with the triplets of presynaptic spikes, we then simulate how these local patterns of 2 pre- and
1 postsynaptic spike change spike transmission probability when the neurons start from the same
initial conditions (average values of excitability, post-spike history, R and u). For the thalamus
and VB-Barrel pairs, the simulations of isolated, local patterns match the general trends of
empirical spike transmission. However, for the VB-Barrel synapse, the effect of ISI_{post} in the
empirical transmission patterns is stronger than in the simulations, suggesting that serial
correlations in ISIs could again play a role and does decrease transmission probability for isolated
patterns. However, as with the responses to triplets of presynaptic spikes, these local patterns alone
are insufficient to explain the empirically observed patterns of spike transmission.

Fig. 4: Effects of pre- and postsynaptic spiking patterns. A: Synaptic transmission patterns
change based on the full sequence of presynaptic spiking. Top panel shows a schematic of 4
different patterns of presynaptic spike triplets with a fixed ISI between the two most recent
presynaptic spikes (black dashed lines). We split the presynaptic ISI distribution into 8 quantiles.
Each data point shows the observed spike transmission probability corresponding to the ISI_2
quantile with the same color. Solid lines are the average estimated probability for each pattern
under the model (based on the full sequence of observed spikes). To examine the influence of
serial correlations, we then stimulate model responses to the isolated triplet pattern, assuming the
synapse is initially in an average state. **B:** Synaptic transmission patterns change depending on the
history of postsynaptic spiking, as well. Here each data point in the scatter plots shows the spike
transmission probability of the corresponding to the postsynaptic ISIs of the same color in the ISI
distribution. Colors represent the corresponding time difference between presynaptic and previous
postsynaptic spike. Solid lines are the average predicted probability for quantiles with
corresponding colors. Last row shows simulations of the model using estimated STP parameters
and fixing the excitability from the model fits to their average values. Here the history effect for
each ISI interval is set to the post-spike history filter value on that interval.

Spike transmission patterns change depending on stimulus type

These results suggest that the presynaptic spike pattern has a complex effect on spike transmission
probability. In sensory systems, one variable that affects the presynaptic spike pattern is the
external stimulus. To examine how differences in stimulus statistics might alter spike transmission,
we fitted our model to a dataset recorded juxtacellularly from an ANF-SBC synapse, presented
with *natural sounds*, a range of randomized frequency-level pure-tones (*tuning stimuli*), and

spontaneous activity in the absence of acoustic stimulation. We merged these three datasets and
fitted the model to the merged dataset. As with the previous fits of the ANF-SBC connection (based
on a different set of *tuning stimuli*), the transmission probability under all three conditions exhibits
a bandpass-like pattern suggesting facilitation and little to no synaptic summation. However, spike
transmission during natural stimuli was markedly different from that during pure tone stimulation.
During natural sounds, transmission probability is maximized at 100 ms rather than 10 ms in the
tuning stimuli and during *spontaneous activity*. Further, *natural stimuli* have much lower
transmission probability at short ISIs. Interestingly, the TM-GLM captures the overall facilitation,
but also captures differences due to the different stimuli. In contrast, a static GLM captures almost
none of the variations in spike transmission probability suggesting that a fixed coupling term,
postsynaptic history, and, slow fluctuations of presynaptic spiking are not sufficient to capture
patterns of spike transmission probabilities (Fig. 5A). Together, these results suggest that the
combination of STP, synaptic summation, history, and excitability is sufficient to explain the
observed differences between stimuli, without requiring any additional adaptation or plasticity.

Since these recordings were performed juxtacellularly, we also have access to the slope of
individual (extracellular) PSPs, which correlates with the intracellular PSP amplitude. We
compared patterns of individual PSP slopes for each stimulus type and how they correlate with the
estimated coupling amplitude following individual presynaptic spikes in our model (Fig. 5B, 5C).
Note that patterns of PSP slopes do not have the same pattern as spike transmission probability,
since there are other factors (e.g. postsynaptic spiking history) contributing to postsynaptic spiking.
These results show the stimulus-dependence of PSP amplitudes and a static GLM without STP
cannot account for these variations. Although the correlation is not perfect, the model does
correlate with the measured PSP slope, even though the model only has access to spikes. By
modeling dynamic functional connectivity, we can approximately reconstruct the amplitude of
individual synaptic events.

Fig. 5: The TM-GLM captures stimulus-dependent changes in spike transmission probability at the ANF-SBC synapse. A) The TM-GLM captures stimulus-dependent spike transmission probability patterns better than a static model without short-term synaptic plasticity. Asterisks show spike transmission probability for (log-spaced) presynaptic ISIs during two types of auditory stimuli and during spontaneous activity: Natural Sounds (yellow), Spontaneous Activity (red), and Tuning Stimuli (blue). Solid lines and 95% confidence bands show model predictions for each stimulus type. Corresponding inter-spike interval distributions are shown on the right. **B)** The TM-GLM captures changes in extracellularly recorded PSPs. Here the observed PSP slope (dashed lines) approximately matches and coupling term in the TM-GLM (solid lines) for each three stimuli. Although the spike transmission probability of the static GLM can vary as a function of presynaptic ISI due to non-synaptic factors, the coupling term is fixed. **C)** Estimates of individual PSP amplitudes predicted by the model and their PSP slopes in the juxtacellular recording. Black lines denote linear fits and the bar plot shows the corresponding Spearman correlations.

Postsynaptic cell-type specific changes in spike transmission patterns

We applied our model to spiking data from a large-scale multi-electrode array recording to investigate the spike transmission dynamics in synapses from putative excitatory neurons to two different putative inhibitory subtypes. We detected putative synapses using the log-likelihood ratio (LLR < -6, ~200 synapses) between a full model of the correlogram that includes the synaptic effect and smooth model of the correlogram that only captures the slow structure (see Methods). We then found excitatory-inhibitory microcircuits where putative excitatory neurons (based on the cross-correlogram and spike waveform) give inputs to putative inhibitory neurons (41 excitatory synapses onto 9 inhibitory neurons in total). To identify inhibitory neurons as inhibitory, we required the neuron to have an outgoing connection to a third neuron with a fast, transient decrease

in the cross-correlogram. Each of the 9 putative inhibitory neurons here had at least one outgoing
connection where the spiking probability of a downstream neuron decreases >18% relative to
baseline following its spiking (Fig. 6A and supplementary figures for individual cross-
correlograms). We then categorized each neuron as a putative fast-spiking (FS, n=5) or regular-
spiking (RS, n=4) unit based on the spike waveform and firing rate (Fig. 6B). Putative FS units
had narrow-width spike waveforms (half-width of the trough = .08±.02 ms) and higher firing rates
(26.07±9.6 Hz) compared to putative RS neurons (n=4) with broader waveforms (half-width =
.14±.02 ms) and lower firing rate (10.18±10.01 Hz).

We identified these microcircuits in different regions with 4 putative excitatory-inhibitory
microcircuits recorded in hippocampus (depth differences: 77.2±49.4 μm), 3 in thalamus
(49.4±26.2 μm), and 2 in motor cortex (36.4±23.5 μm). Putative excitatory neurons showed a
wide spike waveform (half-width = .18±.04 ms) similar to the putative regular-spiking inhibitory
neurons, but these two classes can be distinguished by their outgoing connection types (e.g.
inhibitory/excitatory) [43] (Fig. 6B). Average efficacies from putative excitatory-FS connections
(.22±.12, n=22) were larger, on average, compared to putative excitatory-RS efficacies (.13±.13,
n=19). We then fit the TM-GLM to data from these 41 putative synapses, similar to the three
identified synapses above. We find that the STP parameters for these two types of synapses largely
overlap, except for the membrane time-constant. Interestingly, the membrane time-constants
measured for these inhibitory subtypes *in vitro* overlap with our estimates here (Fig. 6C) [44].
Although *in vitro* studies have not explored the same TM model used here, there is evidence of
postsynaptic cell-type specific STP where putative excitatory-RS connections show facilitation
and putative excitatory-FS connections show depression [5]. Here we find that both connection
types are somewhat facilitating but excitatory-FS connections having a slightly shorter facilitation
time constant. However, unlike what would be expected if excitatory-FS connections were
depressing, the release probability of excitatory-FS connections is lower than for excitatory-RS
connections (Fig. 6C).

To better understand synaptic transmission *in vivo* it is important to consider not just the
parameters of the synapse but the full history of presynaptic spiking in the individual presynaptic
neurons. We use the estimated model parameters to simulate responses to a train of regular
presynaptic spikes with the frequency matched to the average firing rate of the corresponding
excitatory input. In simulating postsynaptic responses to the spike train, we fix the excitability and
postsynaptic history to their average values from model fits and set the initial STP state of the first
spike in the train to the average R and u values from model fits. With these input-matched
simulations, excitatory-RS connections show a higher postsynaptic potential compared to
excitatory-FS connections (Fig. 6D). Similarly, we simulated the paired-pulse ratio (PPR) at
different inter-stimulus intervals in our TM model following the average state. On average,
connections to regular-spiking inhibitory neurons show a higher PPR (Fig. 6E). For all
connections, we then evaluated the spike prediction accuracy of a model without STP (e.g. static
GLM) with our TM-GLM using the Area Under the ROC Curve (Fig. 6F). The model with STP

(TM-GLM) gives more accurate predictions when the postsynaptic neuron spikes following a
presynaptic spike for our population of 41 putative excitatory-inhibitory connections
(AUC=.69±.05) in comparison with the static GLM (AUC=.50±.03). Altogether, these results
illustrate how a dynamic model of functional connectivity, such as the TM-GLM, may allow us to
investigate cell-type-specific differences in short-term synaptic dynamics in behaving animals
using only pre- and postsynaptic spiking.

Fig. 6: Distinctive short-term synaptic plasticity dynamics in connections between excitatory
neurons to putative Regular-Spiking (RS) and Fast-Spiking (FS) inhibitory neurons. A) Here
we examine putative synapses between excitatory neurons and inhibitory neurons (identified by
their cross-correlations) and separate the putative inhibitory neurons into two classes: fast-spiking,
which have narrow spike waveforms and high rates (left), and regular-spiking (right), which have
wide waveforms and lower rates. Identifying these synapses requires both finding both a putative
excitatory input and a putative inhibitory output for the same neuron. B) Half-widths (of the
trough) of the spike waveforms and firing rates for the FS (orange) and RS (blue) inhibitory

neurons, as well as, their excitatory inputs (grey). Individual blue and orange waveforms
(maximum amplitude across the MEA) are shown for all 9 putative inhibitory neurons. C)
Estimated depression, facilitation, and membrane time-constants for excitatory-RS and excitatory-
FS connections, along with the release probability (right). The purple error-bar next to the
membrane time-constant estimations show the median and standard deviations from *in vitro*
experiments [44]. D) Simulated postsynaptic potential amplitudes estimated from Tsodyks-
Markram model of short-term synaptic plasticity using estimated parameters. For each synapse,
PSPs are estimated in response to a pulse train with inter-pulse intervals set to their corresponding
average presynaptic inter-spike intervals. Dots and error bars denote the median and inter-quartile
range for excitatory-RS (blue) and excitatory-FS (red) connections. E) Simulated Paired-Pulse
Ratio for individual synapses of excitatory-RS (blue) and excitatory-FS (red) connections as a
function of the presynaptic ISI. F) Area Under the Curve (AUC) of postsynaptic spiking prediction
using the static GLM without short-term synaptic plasticity (green) and the TM-GLM with short-
term synaptic plasticity (blue).

Discussion

Short-term synaptic plasticity (STP) has been extensively studied *in vitro* and with intracellular
recordings where the amplitudes of individual postsynaptic potential/currents (PSP/PSCs) can be
directly measured. By using controlled experiments with specific, structured presynaptic spike
patterns these studies established how short-term synaptic dynamics can be described by the
interactions between release probability and vesicles/resource dynamics [36]. These alterations in
PSP/PSP amplitudes can affect the statistics of postsynaptic spiking. Thus, STP could, explain
why the probability of postsynaptic spiking depends not just on the presence of a presynaptic spike,
but on the timing of the most recent presynaptic inter-spike interval [14,16]. However, the
relationship between STP and *in vivo* spike transmission patterns is complex. Patterns of
postsynaptic spike transmission are highly diverse and multiple factors beyond STP and the most
recent presynaptic ISI shape these patterns. Here we aimed to disentangle the different
contributions to spike transmission by developing an augmented generalized linear model, the TM-
GLM that explicitly includes STP dynamics as slow changes in postsynaptic excitability and the
history of postsynaptic spiking.

Synapses with different types of STP can allow the same sequence of presynaptic spikes to
generate different patterns of postsynaptic spiking and thereby control the information flow in the
brain. Here we tracked the observed spike transmission probability of three strong synapses from
different species and brain areas. The dynamical spike transmission model enables us to
disentangle different factors (e.g. slow firing rate changes, postsynaptic spiking history, synaptic
summation, and STP) that shape these diverse patterns. First, we investigate the role of STP and
the full sequence of presynaptic spiking activity in shaping the spike transmission patterns. In three
strong synapses (an intra-thalamic synapse, a thalamocortical synapse, and an auditory brainstem
synapse) we show how models of functional connectivity with short-term synaptic plasticity can

1) capture diverse pattern of spike transmission probability, 2) disentangle these transmission
patterns to the multiple factors that shape postsynaptic response, 3) extract biologically plausible
synaptic dynamics, and 4) improve prediction of postsynaptic spiking.

Estimating static functional connectivity using spike times has revealed network structure in the
retina [26] and hippocampus [45], can reconstruct true physiological circuitry [34], and improves
encoding and decoding [25,26,35]. However, synaptic weights change over a wide range of
timescales depend on external stimuli and behavior [20]. Additionally, synaptic dynamics can
shape information transmission in different ways for different pattern of presynaptic spiking, e.g.
different behavioral tasks. Although, standard GLMs can partially capture the first-order effects of
recent presynaptic spikes on postsynaptic spiking probability, they fail to capture the nonlinear
dynamics of synaptic transmission affected by the whole sequence. Here, in a recording from the
endbulb of Held (ANF-SBC) we found that spike transmission patterns differed for different
stimuli (natural sound stimuli, varying pure tones and without stimulation - e.g. spontaneous
activity), and these differences were well-described by the TM-GLM. Although the STP-
parameters were the same for all stimuli, the different presynaptic spike patterns yield different
synaptic dynamics and different patterns of spike transmission. Since spike transmission
probability in the TM-GLM depends on the full history of presynaptic spiking, this model can
account for changes on behavioral timescales even in the absence of adaptation or other forms of
plasticity (e.g. STDP, LTP).

Cell-type specific interactions in layers and regions of the brain perform different computational
tasks. Previous *in vitro* studies have shown that STP dynamics depend on both presynaptic and
postsynaptic cell-types [5]. Here in a large multi-electrode array recording of a freely behaving
mouse we investigated STP dynamics of synaptic connections from putative excitatory neurons to
two different subtypes of putative inhibitory neurons: fast-spiking and regular-spiking. Using
inferred short-term dynamics, predicted responses to train of spikes with the same input
frequencies as the presynaptic neuron in those connections show facilitation in excitatory-RS and
depression in excitatory-FS connections which are in line with previous *in vitro* findings [5].
Moreover, the model with short-term dynamics significantly improves the prediction of
postsynaptic activity. As large-scale extracellular recordings advance, models such as the TM-
GLM are promising to characterize and compare the short-term dynamics of spike transmission of
many different cell types, brain regions, and species.

Although our model provides a tool to characterize the dynamics of spike transmission, there are
limitations on how well TM-GLM can capture true synaptic dynamics. Firstly, functional
connections inferred from spikes do not necessarily guarantee anatomical connections. A peak in
the cross-correlogram does not uniquely indicate the presence of a monosynaptic connection
[46,47]. Here we assume that the transient, short-latency increase in postsynaptic spiking activity
following a presynaptic spike indicates the presence of an excitatory monosynaptic connection
[17]. Nevertheless, verifying connections using optogenetics, juxtacellular recordings [48],

imaging [49] or anatomical reconstruction provide a more accurate estimate of true anatomical
connections. Secondly, to model short-term dynamics in spiking neurons we employ a rate model
that does not explicitly account for the detailed membrane potential of the postsynaptic neuron.
Other approaches to modeling synaptic transmission with realistic spike-generation mechanisms,
currents, and even dendritic morphology do exist, but are typically less computationally tractable
[50]. Here we employed an augmented GLM with a logistic spike nonlinearity. We chose the
logistic nonlinearity over the conventional exponential function as it appears to better describe
strong connections, such as the ANF-SBC, but other nonlinearities may be better for other neurons
[51]. There are also alternatives to the Tsodyks-Markram model for modeling synaptic dynamics.
The TM model is biologically plausible, but, since it is deterministic, it ignores the stochasticity
of synaptic release and only tracks the dynamics of average postsynaptic potentials. Finally, there
are many covariates that could be added to improve model performance, including local field
potentials, connections to other simultaneously observed presynaptic neurons [35], higher-order
history or coupling terms [52,53], and covariates related to other types of plasticity [52,54–57].
Despite these simplifying assumptions and the fact that we only observe a fraction of inputs to the
neuron, the TM-GLM captures a wide diversity of *in vivo*, excitatory spike transmission patterns.

Short-term synaptic plasticity alters information transmission from presynaptic to postsynaptic
neurons by dynamically changing the synaptic efficacy [14,16,36]. Intracellular studies *in vitro* or
with artificial stimulation patterns have shown that short-term synaptic dynamics depend on cell-
types and brain regions [5,58]. However, there is evidence that, in addition to these anatomical
dependencies, short-term synaptic dynamics also depend on stimulus type and the larger
computational function of the neural circuit [59]. To understand how these synaptic dynamics alter
neural computations we will need to study them during natural patterns of presynaptic spiking [60]
and, ultimately, during natural behavior. Since large-scale intracellular recordings are currently
not feasible *in vivo*, here we examined the possibility of using existing large-scale extracellular
recordings to quantify the dynamics of spike transmission and infer the short-term dynamics of
synaptic responses. We find that including STP in models of spiking neurons can capture diverse
patterns of spike transmission, including patterns that are stimulus-dependent and cell-type-
specific. Additionally, these models substantially improve prediction of postsynaptic spiking
following presynaptic spikes and, at least in some cases, can approximately reconstruct individual
PSP amplitudes.

Material and methods

Neural Data

To illustrate how synaptic dynamics can be estimated from spikes, we first examined a set of three
strong putative or identified synapses with diverse spike transmission probability patterns: (i) a
dual-electrode recording of a thalamocortical projection in the barrel system, (ii) an *in vivo* loose-
patch (juxtacellular) recording at the calyceal endbulb of Held synapse in the auditory brainstem,
and (iii) a recording from a pair of neurons in the thalamus detected from a larger multi-electrode

array (MEA) recording. Next, we applied our model more generally to analyze a large sample of
putative synaptic connections recorded from the MEA dataset. The data from these three identified
strong synapses and the MEA data have been collected from different species, regions, cell-types,
under different stimulation and show a diverse pattern of postsynaptic spiking probability. In all
cases we deduce short-term synaptic dynamics on the basis of only pre- and postsynaptic spike
observations.

For the first putative synapse, we use *in vivo* data from simultaneous extracellular recordings in
ventrobasal (VB) thalamic barreloids and topographically aligned, somatosensory cortical barrel
columns (VB-Barrel) in awake, unanesthetized, adult rabbits. Detailed surgical and physiological
methods have been described previously [61]. Spike-triggered averages of the cortical spikes
following spiking of the VB neuron was used to identify connected S1 neurons. Based on the
presence of high frequency discharge (3+ spikes, > 600 Hz) following electrical stimulation of the
thalamus, and narrow spike waveforms, the S1 neuron in this recording was identified as a putative
inhibitory neuron [62]. These recordings identified several putative thalamocortical projections.
The putative synapse that we model here is particularly clear, with 68,345 pre- and 128,096
postsynaptic spikes recorded over the course of 92 minutes of spontaneous activity and has been
previously studied in [14,63].

For the second synapse, we examined *in vivo* loose-patch recordings at the Endbulb of Held in
young adult gerbils. Detailed surgical and physiological methods have been previously described
[64]. Briefly, the glass electrode was positioned in the anterior portion of the ventral cochlear
nucleus (AVCN) and single-units were recorded during varying acoustic stimulation. Single units
were classified when recording a positive action potential amplitude of at least 2 mV and showing
the characteristic complex waveform identifying them as large spherical bushy cells (SBC) of the
rostral AVCN. This recording included a mixture of juxtacellular waveforms: an isolated
excitatory PSP (EPSP) or an EPSP followed by a postsynaptic action potential. For both cases the
timing of EPSPs and spikes and rising slope of the EPSPs were extracted. The timing and slope of
the EPSPs were identified using a slope threshold for the rising part of EPSPs as previously
described [65]. We then modeled spike transmission probability patterns for two recordings: (i)
during randomized pure tone acoustic stimulation and (ii) during multiple stimuli, i.e. randomized
frequency-level pure tone stimulation interspaced with spontaneous activity, natural sounds, and
also during spontaneous activity. Using this second dataset, we characterized how variable
presynaptic spike patterns evoked by different stimuli affected the patterns of spike transmission
at the same synapse.

We also use MEA spiking data to study the factors shaping spike transmission probability patterns
in a large-scale recording with multiple cell-types. Here we use a previously collected, publicly
available recording from the Cortex Lab at UCL [66,67] with data from two Neuropixels electrode
arrays recorded simultaneously, each with 960 sites (384 active) with lengths of 10-mm and
spacing of 70 × 20-μm (http://data.cortexlab.net/dualPhase3/). The two electrode arrays span

multiple brain areas and ~90 min of data was collected in an awake, head-fixed mouse on a rotating
rubber wheel during visual stimulus presentations. Spikes were automatically detected and sorted
using Kilosort [68] on the broadband (0.3–10 kHz) signal and then manually curated. If two
clusters of spikes had similar waveforms, cross-correlogram features, and spike amplitudes, they
were merged into a single cluster and assigned to a single neuron. In total, 831 well-isolated single
neurons where identified from the two probes in several different brain areas: visual cortex (n=74),
hippocampus (n=64), thalamus (n=244), motor cortex (n=243), and striatum (n=200). Due to the
large number of simultaneously recorded neurons in this dataset, there are many potential synapses
($\sim 831^2$).

Synapse Detection:

To identify putative monosynaptic connections between well-isolated single neurons, we looked
for specific patterns in the cross-correlograms [46]. If two neurons are monosynaptically
connected, the probability of postsynaptic spiking increases/decreases rapidly following a
presynaptic spike. In spiking data, this rapid, transient change can be seen in cross-correlograms
as an asymmetric bump/dip in the number of postsynaptic spikes following presynaptic spikes
[18]. For each connection we calculated the cross-correlogram in a 5 ms window before and after
presynaptic spikes with bin-size of 0.1 ms. To avoid aliasing in the cross-correlograms, we added
a small, random shift to each postsynaptic spike drawn uniformly between $-\Delta t/2$ and $\Delta t/2$ where
Δt is the spike time resolution (0.01 ms in most cases). Here we used a model-based approach
using the cross-correlograms to decide whether two synapses are monosynaptically connected. To
fit the cross-correlogram we used a baseline rate μ, a linear combination of B-spline bases $\boldsymbol{B}(t)$,
and a weighted alpha function to model the synapse, $w\,\boldsymbol{\alpha}(t)$, all passed through an output
nonlinearity; $\lambda(t) = \exp(\mu + \boldsymbol{r}\boldsymbol{B}(t) + w\,\boldsymbol{\alpha}(t))$. The alpha function, $\boldsymbol{\alpha}(t) = (t - t_d)/$
$\tau_\alpha \exp(1 - (t - t_d)/\tau_\alpha)$, describes the shape of the synaptic potential where t_d is the synaptic
delay and τ_α is the synaptic time-constant [22]. For individual connections, we estimate these
parameters by maximizing the penalized Poisson log-likelihood $l(\mu, \boldsymbol{r}, w, t_d, \tau_\alpha) = \Sigma y_i log\lambda_i -$
$\Sigma\lambda_i + \epsilon\|\boldsymbol{r}\|_2$ where y_i is the number of postsynaptic spikes observed in the i-th bin of the
correlogram and $\|\boldsymbol{r}\|_2$ regularizes the model to penalize B-spline bases for capturing sharp
increases in the cross-correlogram. ϵ is a regularization hyper-parameter which we set to 1 based
on manual search. Due to the parameterization of $\boldsymbol{\alpha}(t)$, the log-likelihood is not concave.
However, since the gradient of the log-likelihood can be calculated analytically, we efficiently
optimize the likelihood using LBFGS. During the optimization, the delay and time-constant
parameters are log-transformed, allowing us to use unconstrained optimization, even though they
are strictly positive. We used random restarts to avoid local maxima. To identify putative
monosynaptic connections in the large-scale multi-electrode array data, we compared this model
with a smooth model with slow changes in cross-correlogram and without the synapse, $\lambda_0(t) =$
$\exp(\mu' + \boldsymbol{r}'\boldsymbol{B}(t))$, using the log-likelihood ratio (LLR) test between our full model with synapse
and the nested smooth model. Since low values of the likelihood ratio mean that the observed result
was better explained with full model as compared to the smooth model, we then visually screened

pair-wise connections with lowest ratios (LLR < -6) compared to the null model to find putative
synapses. Out of ~831^2 possible connections in this dataset we find ~200 putative synapses
(0.03%). We handpicked a strong putative synapse between two thalamic neurons to study its
efficacy pattern in detail alongside the VB-Barrel and ANF-SBC synapses.

In addition to this single strong synapse, we also categorize putative pre- and postsynaptic cell
types for the connections detected in the MEA dataset. For this purpose, we assessed single units
based on their cross-correlograms, firing rates, and spike waveforms. We categorized units as
excitatory or inhibitory if, in accordance with Dale's law [69], all outgoing cross-correlograms
showed transient, short-latency (<4ms) increase/decrease in spiking probability [35]. We then
looked into identified inhibitory neurons and categorized them into to putative fast-spiking (FS)
and regular-spiking (RS) inhibitory neurons. Using these putative Excitatory-FS and Excitatory-
RS synapses, we then examine how the spike transmission patterns differ for these two subtypes
of inhibitory neurons.

Extending a Generalized Linear Model to Account for Short-term Plasticity (TM-GLM)

Short-term synaptic plasticity causes the amplitude of postsynaptic potentials (PSP) to vary over
time depending on the dynamics of synaptic resources and utilization and can be modeled using
the pattern of presynaptic spiking [36,70]. However, changes in the overall postsynaptic spiking
probability cannot be uniquely attributed to changes in amplitudes of postsynaptic potentials. To
accurately describe the dynamics of spike transmission, we also need to account for the membrane
potential summation, the excitability of the postsynaptic neuron (e.g. slow changes in the
presynaptic firing rate) and the dynamics of postsynaptic spiking (e.g. refractory period, after
hyperpolarization current). We developed an extension of a generalized linear model, which we
call a TM-GLM to describe each of these effects. Concretely, the probability of a postsynaptic
spike shortly after each presynaptic spike accounts for the full sequence of previous presynaptic
spiking and the recent history of postsynaptic spiking. We define the conditional intensity of the
postsynaptic neuron after the i-th presynaptic spike, $t_s^{(i)}$, so that the probability of observing a
postsynaptic spike in the j-th time bin after the i-th presynaptic spike is given as:

$$\lambda_i(t_j) = \sigma\left(\beta_0 + X_c\left(t_s^{(i)}\right)\beta_c + \sum_{t_r^{(l)} < t_s^{(i)}} X_h\left(t_s^{(i)} - t_r^{(l)}\right)\beta_h + A_s\, w_i\, \alpha(t_j)\right)$$

where $t_r^{(l)}$ are the postsynaptic spike times preceding $t_s^{(i)}$. For each presynaptic spike, our model
decomposes the firing rate of the postsynaptic neuron into four effects: a baseline firing rate, β_0,
slow fluctuations in presynaptic firing rate $X_c\beta_c$, history effects from the recent postsynaptic spikes
(prior to $t_s^{(i)}$), $X_h\beta_h$, and a time-varying coupling effect from the presynaptic input, $A_s w\, \alpha(t)$ (Fig.
7).

Fig. 7: TM-GLM. Postsynaptic spiking probability before passing the spiking nonlinearity (yellow) changes as a linear combination of presynaptic coupling term with STP dynamics (blue), postsynaptic spiking history (green), the postsynaptic excitability (red). Transparent red curves show the bases of slow changes in postsynaptic probability at presynaptic spike times (X_c).

Here we model slow fluctuations in the postsynaptic rate $X_c\beta_c$ with a linear combination of B-splines with equally spaced knots every 50 seconds of recording time. In the history term, splines (X_h) span a period of 10 ms prior to each presynaptic spike with 4 logarithmically-spaced knots. By scaling $\alpha(t_j)$ with a multiplicative factor, w_i, the strength of a synapse can vary over time and, in this case, depends on the detailed sequence of presynaptic spiking and their corresponding inter-spike intervals. A_s is the magnitude of the synaptic strength. In this case we use a model for short-term synaptic plasticity that allows both depression (where the w_i decreases for shorter presynaptic ISIs) and facilitation (where the w_i increases for shorter presynaptic ISIs), and incorporates membrane summation. To model these effects, w_i is determined by a nonlinear dynamical system based on the Tsodyks and Markram (TM) model [36,71] where: $w_i = w_{i-1} \exp\left(-\frac{t_s^{(i)}-t_s^{(i-1)}}{\tau_s}\right)\pi_i +$ $R_i u_i$, where τ_s is the membrane time-constant and the first term of the equation describes how postsynaptic membrane potential summation increases the probability of postsynaptic spiking. This membrane summation will be ignored if there is a postsynaptic spike: $\pi_i = \left\{0 \text{ if } t_s^{(i-1)} < t_r^{(i-1)} < t_s^{(i)}; 1 \text{ otherwise}\right\}$. In the second term of this equation, R represents the dynamics of resources and u describes their utilization.

$$R_i = 1 - [1 - R_{i-1}(1 - u_{i-1})] \exp\left(-\frac{t_s^{(i)} - t_s^{(i-1)}}{\tau_d}\right)$$

$$u_i = U + [u_{i-1} + f(1 - u_{i-1}) + U] \exp\left(-\frac{t_s^{(i)} - t_s^{(i-1)}}{\tau_f}\right)$$

where τ_d and τ_f are the depression and facilitation time-constants. U is the release probability and f is the magnitude of facilitation. To make the estimation more tractable, we approximate the full optimization problem and estimate synaptic delay, t_d, and time-constant, τ_α, by fitting $\alpha(t)$ using the full cross-correlogram, as above. We fix these parameters for the rest of the optimization process. We then maximize a penalized, Bernoulli log-likelihood $l(\theta) = \Sigma\Sigma \left[y_{ij}\lambda_i(t_j) - (1 - y_{ij})(1 - \lambda_i(t_j)) \right] + \gamma\|\theta'_{stp}\|_2$ where $\gamma = 1$ is the regularization hyperparameter to estimate the parameters: $\theta = \{\beta_0, \beta_{c=1:C}, \beta_{h=1:H}, A_s, \theta_{stp}\}$, $\theta_{stp} = \{\tau_d, \tau_f, U, f, \tau_s\}$.

As with previous applications of GLMs, we assume that bins are conditionally independent given the covariates, but unlike many other GLMs, here we only calculate the log-likelihood during short intervals (5ms) after presynaptic spikes. With y_{ij} being a binary value representing the presence of a postsynaptic spike in the j-th time bin after the i-th presynaptic spike. We again used a logarithmic transformation for the time-constants to avoid negative values and logit transformation for U and f to bound their values in the interval $[0,1]$; $\theta'_{stp} = \{\log(\tau_d), \log(\tau_f), \text{logit}(U), \text{logit}(f), \log(\tau_s)\}$. By modeling STP this model is no longer a strict GLM, and the log-likelihood may have local maxima. Here we use random restarts to avoid local maxima in our optimization process. The parameters of each restart $\{\beta_0, \beta_{c=1:C}, \beta_{h=1:H}, A_s\}$ are initialized by adding noise ($\sim N(0,1)$) to the corresponding parameters in a standard GLM. We initialize the plasticity parameters with $\tau'^{(0)}_d \sim N(-1,5)$, $\tau'^{(0)}_f \sim N(-1,5)$, $U'^{(0)} \sim N(0,5)$, $f'^{(0)} \sim N(0,5)$, $\tau'^{(0)}_s \sim N(-3,5)$. We then use an LBFGS algorithm to optimize the log-likelihood where we calculate all derivatives analytically except for derivatives of θ_{stp} which we calculate numerically. To estimate the uncertainty of the parameters, we bootstrap the data from each of the strong synapses by chunking the whole recording time into samples of 50 seconds then resampling the chunks to generate a new spike train with the same original length.

Calculating spike transmission probability

To demonstrate how the probability of postsynaptic spiking changes according to the corresponding presynaptic inter-spike intervals, we estimated spike transmission probabilities from the cross-correlograms directly instead of using a model. To calculate this probability, we focused on a transmission interval after the presynaptic spike where the conditional intensity (when corrected for the baseline rate) goes above 10% of the maximum of $\boldsymbol{\alpha}(t)$. We split the presynaptic inter-spike interval distribution into log-spaced intervals, and, for each interval, we calculate the ratio between numbers of postsynaptic spikes in the transmission interval to the number of presynaptic spikes. Unlike previous studies [14,61] we do not correct this probability for the baseline postsynaptic rate. The uncorrected probability allows us to more directly compare the model predictions to the empirical spike transmission probabilities. Since our model gives an estimate of the postsynaptic probability after each individual presynaptic spike, we can average over the same transmission interval. However, we know if there is a postsynaptic spike in the

transmission interval, probability of a postsynaptic spike goes to ~0 for all consecutive bins due to
the post-spike dynamics (e.g. refractory period). Therefore, we measure the predicted probability
of a postsynaptic spike in a 5ms window after i-th presynaptic spike from binned $\lambda_i(t_j)$ as follows:
$z_i = \sum_{j=1}^{L} \lambda_i(t_j) \prod_{m=1}^{j-1}(1 - \lambda_i(t_m))$. Here we assume conditional independence of the j-th bin
after a presynaptic spike, but we enforce a refractory period for all bins after a postsynaptic spike
in our generative model. Here L is the first bin that y_{ij} is nonzero. z_i represents the probability of
postsynaptic spiking after each presynaptic spike and we fit a smooth curve over the distribution
of z_i's and their corresponding inter-spike intervals to compare with the empirical spike probability
patterns.

Modeling the effect of local patterns of pre- and postsynaptic spiking

The observed and modeled spike transmission patterns, as calculated above, reflect the expected
postsynaptic spike probability given a specific presynaptic ISI. However, since the presynaptic
ISIs are not independent and there are serial correlations in ISIs, the detailed sequence of the pre-
and postsynaptic spiking likely affects the shapes of these curves. To quantify the effects of serial
ISI correlations on the model of spike transmission probability we demonstrate how local patterns
of presynaptic spiking modifies spike transmission patterns in the data and the model. For each of
the three strong identified synapses we measure postsynaptic spiking probability in response to
presynaptic spike triplets. Due to the limited number of spikes in our data, we divide the
presynaptic ISI distribution into few log-spaced intervals and measure the postsynaptic spiking
probability for triplets with the two ISIs that fall in those intervals. Similarly, we measure the
predicted postsynaptic probability in response to the presynaptic triplets. After measuring
postsynaptic responses to presynaptic spike triplets in the data and the model, we simulate the
contribution of STP in shaping the transmission pattern in response to these triplets. To factor out
contributions of the postsynaptic history and slow changes in presynaptic firing rate, we fix the
corresponding values in the model to their average values within the model. In these simulations,
we also fix the initial values of the STP dynamics in the TM model for the first spike of the triplets
to the average R and u within the model. This approach enables us to illustrate how short-term
synaptic plasticity in triplets of presynaptic spikes changes spike transmission probability and how
serial correlations in presynaptic spiking affect spike transmission probability.

The postsynaptic spike history and the serial correlations between the pre- and postsynaptic spiking
also modify spike transmission probability patterns. To investigate history effects in the local
pattern of pre- and postsynaptic spikes, we measured the postsynaptic spiking probability in
response to two presynaptic spikes and a postsynaptic spike preceding the most recent presynaptic
spike. Due to the limited number of spikes and sparseness of the split cross-correlograms, we again
divided the presynaptic and postsynaptic ISI distributions into a few log-spaced intervals. We then
measure the spike transmission probability for a group of presynaptic spikes that their preceding
presynaptic ISIs and postsynaptic spike ISIs fall into different combinations of pre- and
postsynaptic log-spaced intervals. After measuring postsynaptic responses to any possible

combination of the two most recent presynaptic spikes and their postsynaptic spikes in the data
and the model, we simulate the contribution of the history and STP together in shaping the
transmission. In our simulation the excitability was set to the model estimates. To measure the
effects of postsynaptic spiking history, for each postsynaptic ISI, we fix the history contribution
to estimated post-spike history filter value at that postsynaptic ISI. We then use the predicted STP
parameters from the data to simulate the STP contribution in response to paired pulses of
presynaptic ISIs where we again fix the initial values of the TM model for the first presynaptic
spike to the average R and u within the model. This approach enables us to illustrate how short-
term synaptic plasticity in local patterns of two presynaptic spikes and a postsynaptic spike
changes spike transmission probability and quantifies how serial correlations between pre- and
postsynaptic spiking affect spike transmission probability.

Evaluating prediction accuracy

In addition to evaluating the estimated parameters and comparing the model to empirical spike
transmission probabilities, we also assess how accurately the model can predict postsynaptic
spiking. Not only can we predict the probability of a spike given specific presynaptic ISIs, but we
can also predict whether there will be a postsynaptic spike following each individual presynaptic
spike. To quantify how well the predicted postsynaptic spike probability, z_i, predicts the
postsynaptic spiking activity, we use Receiver Operating Characteristic (ROC) curves. To compute
the ROC curve, we first create a threshold version of z_i which operates as our prediction: $\{(\hat{r}_i =$
$1)$ if $(z_i > \text{thr})$; 0 otherwise$\}$. Changing the threshold from 0 to 1 traces out a relationship between
the true positive rate (TPR) and false positive rate (FPR). The area under the ROC curve (AUC)
reflects the performance of each model, where a perfect classifier has AUC=1 and a random
classifier has AUC=0.5. Effectively, the AUC is the probability of a randomly chosen spike having
a higher model probability than a randomly chosen non-spike [72]. Here we calculate the AUC for
short intervals (~5ms) after presynaptic spikes and check whether we detect a postsynaptic spike
in the transmission interval where $\alpha(t)$ is above 10% of its maximum. Here we compare the AUC
for the static model of connectivity without short-term synaptic plasticity with our dynamical
model.

A simplified rate model to simulate effects of synaptic summation and post-spike history

Our TM-GLM's prediction of the spike transmission pattern is data-driven and depends on the full
history of pre- and postsynaptic spiking. To better understand and illustrate how STP, synaptic
summation, and post-spike history interact to create the observed patterns of spike transmission,
we simulated postsynaptic responses in a simplified voltage model. Namely, we consider PSP
summation in response to a pattern of two presynaptic spikes. We assume that the synapse is
initially fully recovered, and the PSC amplitudes are determined by the 4-paramter TM model with
$U = 0.7, \tau_d = 1.7, \tau_f = 0.02, f = 0.05$ for the depressing synapse and $U = 0.1, \tau_d = 0.02, \tau_f =$
$1, f = 0.11$ for the facilitating synapse [28]. We then convolve the PSCs (delta function kernel)
with a PSP kernel, $\exp(-t/\tau_v) - \exp(-t/\tau_r)$, with $\tau_v = .01$ and $\tau_r = .001$ ms to describe synaptic

summation. We assume that the instantaneous postsynaptic spike probability is simply a nonlinear function of the distance to a threshold voltage $\sigma\big(5(V(t) - V_{th})\big)$ where $\sigma(x) = 1/(1 + e^{-x})$ and $V_{th} = .5, .75,$ and 1 correspond to strong, moderate, and weak inputs respectively. The spike transmission probability sums this instantaneous probability over a window of 20ms after each presynaptic spike. Finally, we adjust the spike transmission probability for the second PSP to account for potential post-spike history effects. Namely, we assume that the adjusted spike transmission probability for the second spike is $p_2^* = (1 - p_1)p_2 + p_1 p_2 f_{ahp}$ where p_1 is the transmission probability for the first spike, p_2 is the unadjusted probability for the second spike, and f_{ahp} is the effect of the after-hyperpolarization. Here we use $f_{ahp}(\Delta t) = (\sigma\big(150(\Delta t - 0.02)\big) - c)/d$ where Δt is the presynaptic ISI, and c and d are constants ensuring that $f_{ahp}(0) = 0$ and $f_{ahp}(\infty) = 1$. Although this simulation is highly simplified, it demonstrates how the observed spike transmission pattern depends, not just on the type and timescale of STP, but on the interaction between STP, synaptic summation, after-hyperpolarization effects, and the spike nonlinearity.

Acknowledgements

Thanks to Nick Steinmetz for providing MEA dataset. AG, NR, and IHS were supported by NSF CAREER 1651396 to IHS.

References

1.   Abbott LF, Nelson SB. Synaptic plasticity: taming the beast. Nat Neurosci. 2000;3: 1178–1183. doi:10.1038/81453

2.   Abbott LF, Regehr WG. Synaptic computation. Nature. 2004. doi:10.1038/nature03010

3.   Zucker RS, Regehr WG. Short-term synaptic plasticity. Annu Rev Physiol. 2002;64: 355–405. doi:DOI 10.1146/annurev.physiol.64.092501.114547

4.   Regehr WG. Short-term presynaptic plasticity. Cold Spring Harb Perspect Biol. 2012; doi:10.1101/cshperspect.a005702

5.   Thomson AM, Lamy C. Functional maps of neocortical local circuitry. Front Neurosci. Frontiers; 2007;1: 19–42. doi:10.3389/neuro.01.1.1.002.2007

6.   Dittman JS, Kreitzer AC, Regehr WG. Interplay between facilitation, depression, and residual calcium at three presynaptic terminals. J Neurosci. 2000; doi:10.1523/JNEUROSCI.20-04-01374.2000

7.   Wang Y, Markram H, Goodman PH, Berger TK, Ma J, Goldman-Rakic PS. Heterogeneity in the pyramidal network of the medial prefrontal cortex. Nat Neurosci. 2006;9: 534–542. doi:10.1038/nn1670

8.   Fortune ES, Rose GJ. Short-term synaptic plasticity as a temporal filter. Trends Neurosci. 2001;24: 381–385. doi:10.1016/S0166-2236(00)01835-X

9.   Cook DL, Schwindt PC, Grande LA, Spain WJ. Synaptic depression in the localization of
sound. Nature. 2003;421: 66–70. doi:10.1038/nature01248

10.   Mongillo G, Barak O, Tsodyks M. Synaptic theory of working memory. Science.
2008;319: 1543–1546. doi:10.1126/science.1150769

11.   Abbott LF. Synaptic Depression and Cortical Gain Control. Science (80-). 1997;275: 221–
224. doi:10.1126/science.275.5297.221

12.   Lindner B, Gangloff D, Longtin A, Lewis JE. Broadband Coding with Dynamic Synapses.
J Neurosci. 2009;29: 2076–2087. doi:10.1523/JNEUROSCI.3702-08.2009

13.   Drew PJ, Abbott LF. Extending the effects of spike-timing-dependent plasticity to
behavioral timescales. Proc Natl Acad Sci. 2006;103: 8876–8881. doi:10.1073/pnas.0600676103

14.   Swadlow HA, Gusev AG. The impact of "bursting" thalamic impulses at a neocortical
synapse. Nat Neurosci. 2001;4: 402–8. doi:10.1038/86054

15.   Usrey WM, Reppas JB, Reid RC. Paired-spike interactions and synaptic efficacy of retinal
inputs to the thalamus. Nature. 1998; doi:10.1038/26487

16.   English DF, McKenzie S, Evans T, Kim K, Yoon E, Buzsáki G. Pyramidal Cell-
Interneuron Circuit Architecture and Dynamics in Hippocampal Networks. Neuron. 2017;96: 505–
520. doi:10.1016/j.neuron.2017.09.033

17.   Perkel DH, Gerstein GL, Moore GP. Neuronal spike trains and stochastic point processes.
II.   Simultaneous   spike   trains.   Biophys   J.   1967;7:   419–440.   Available:
http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1368069/

18.   Barthó P, Hirase H, Monconduit L, Zugaro M, Harris KD, Buzsáki G. Characterization of
Neocortical Principal Cells and Interneurons by Network Interactions and Extracellular Features.
J Neurophysiol. 2004;92: 600–608. doi:10.1152/jn.01170.2003

19.   Csicsvari J, Hirase H, Czurko A, Buzsáki G. Reliability and state dependence of pyramidal
cell-interneuron synapses in the hippocampus: An ensemble approach in the behaving rat. Neuron.
1998;21: 179–189. doi:10.1016/S0896-6273(00)80525-5

20.   Fujisawa S, Amarasingham A, Harrison MT, Buzsáki G. Behavior-dependent short-term
assembly   dynamics   in   the   medial   prefrontal   cortex.   Nat   Neurosci.   2008;11:   823–833.
doi:10.1038/nn.2134

21.   Usrey WM, Alonso JM, Reid RC. Synaptic interactions between thalamic inputs to simple
cells in cat visual cortex. J Neurosci. 2000;20: 5461–5467. doi:20/14/5461 [pii]

22.   Carandini M, Horton JC, Sincich LC. Thalamic filtering of retinal spike trains by
postsynaptic summation. J Vis. 2007;7: 20. doi:10.1167/7.14.20

23.   Trussell LO, Zhang S, Ramant IM. Desensitization of AMPA receptors upon multiquantal
neurotransmitter release. Neuron. 1993;10: 1185–1196. doi:10.1016/0896-6273(93)90066-Z

24.   Henze DA, Buzsáki G. Action potential threshold of hippocampal pyramidal cells in vivo
is increased by recent spiking activity. Neuroscience. 2001;105: 121–130. doi:10.1016/S0306-
4522(01)00167-1

25.   Truccolo W, Eden UT, Fellows MR, Donoghue JP, Brown EN, John P. A Point Process
Framework for Relating Neural Spiking Activity to Spiking History, Neural Ensemble, and
Extrinsic Covariate Effects. J Neurophysiol. 2005;93: 1074–1089. doi:10.1152/jn.00697.2004

26.   Pillow JW, Shlens J, Paninski L, Sher A, Litke AM, Chichilnisky EJ, et al. Spatio-temporal
correlations and visual signalling in a complete neuronal population. Nature. Macmillan Publishers
Limited. All rights reserved; 2008;454: 995–9. doi:10.1038/nature07140

27.   Harris K, Csicsvari J, Hirase H, Dragoi G, Buzsáki G. Organization of cell assemblies in
the hippocampus. Nature. 2003;424: 552–556. doi:10.1038/nature01765.1.

28.   Ghanbari A, Malyshev A, Volgushev M, Stevenson IH. Estimating short-term synaptic
plasticity from pre- and postsynaptic spiking. PLoS Comput Biol. 2017;
doi:10.1371/journal.pcbi.1005738

29.   Stoelzel CR, Bereshpolova Y, Gusev AG, Swadlow HA. The Impact of an LGNd Impulse
on the Awake Visual Cortex: Synaptic Dynamics and the Sustained/Transient Distinction. J
Neurosci. 2008; doi:10.1523/JNEUROSCI.4726-07.2008

30.   Stoelzel CR, Bereshpolova Y, Swadlow HA. Stability of Thalamocortical Synaptic
Transmission across Awake Brain States. J Neurosci. 2009; doi:10.1523/JNEUROSCI.5983-
08.2009

31.   Thomson AM, West DC, Wang Y, Bannister AP. Synaptic Connections and Small Circuits
Involving Excitatory and Inhibitory Neurons in Layers 2-5 of Adult Rat and Cat Neocortex: Triple
Intracellular Recordings and Biocytin Labelling In Vitro. Cereb Cortex. Oxford University Press;
2002;12: 936–953. doi:10.1093/cercor/12.9.936

32.   Yang H, Xu-Friedman MA. Impact of Synaptic Depression on Spike Timing at the Endbulb
of Held. J Neurophysiol. 2009; doi:10.1152/jn.00072.2009

33.   Yang H, Xu-Friedman MA. Relative Roles of Different Mechanisms of Depression at the
Mouse Endbulb of Held. J Neurophysiol. 2008; doi:10.1152/jn.01293.2007

34.   Gerhard F, Kispersky T, Gutierrez GJ, Marder E, Kramer M, Eden U. Successful
Reconstruction of a Physiological Circuit with Known Connectivity from Spiking Activity Alone.
PLoS Comput Biol. 2013;9. doi:10.1371/journal.pcbi.1003138

35.   Stevenson IH, London BM, Oby ER, Sachs NA, Reimer J, Englitz B, et al. Functional
Connectivity and Tuning Curves in Populations of Simultaneously Recorded Neurons. PLoS
Comput Biol. Public Library of Science; 2012;8: e1002775. doi:10.1371/journal.pcbi.1002775

36.   Markram H, Wang Y, Tsodyks M. Differential signaling via the same axon of neocortical
pyramidal neurons. Proc Natl Acad Sci U S A. 1998;95: 5323–8. doi:10.1073/pnas.95.9.5323

37.   Huang C, Resnik A, Celikel T, Englitz B. Adaptive Spike Threshold Enables Robust and
Temporally Precise Neuronal Encoding. PLoS Comput Biol. 2016;
doi:10.1371/journal.pcbi.1004984

38.   Costa RP, Sjöström PJ, van Rossum MCW. Probabilistic inference of short-term synaptic
plasticity in neocortical microcircuits. Front Comput Neurosci. 2013;7: 75.
doi:10.3389/fncom.2013.00075

39.   Gil Z, Connors BW, Amitai Y. Differential regulation of neocortical synapses by
neuromodulators and activity. Neuron. 1997; doi:10.1016/S0896-6273(00)80380-3

40.   Wang Y, Manis PB. Synaptic Transmission at the Cochlear Nucleus Endbulb Synapse
During Age-Related Hearing Loss in Mice. J Neurophysiol. 2005; doi:10.1152/jn.00374.2005

41.   Wang Y, Manis PB. Short-term synaptic depression and recovery at the mature mammalian
endbulb of Held synapse in mice. J Neurophysiol. 2008; doi:10.1152/jn.90715.2008

42.   Paz JT, Chavez M, Saillet S, Deniau J-M, Charpier S. Activity of Ventral Medial Thalamic
Neurons during Absence Seizures and Modulation of Cortical Paroxysms by the Nigrothalamic
Pathway. J Neurosci. 2007; doi:10.1523/JNEUROSCI.4677-06.2007

43.   Moore AK, Wehr M. Parvalbumin-Expressing Inhibitory Interneurons in Auditory Cortex
Are Well-Tuned for Frequency. J Neurosci. 2013;33: 13713–13723.
doi:10.1523/JNEUROSCI.0663-13.2013

44.   Perrenoud Q, Rossier J, Geoffroy H, Vitalis T, Gallopin T. Diversity of gabaergic
interneurons in layer VIa and VIb of mouse barrel cortex. Cereb Cortex. 2013;
doi:10.1093/cercor/bhs032

45.   Harris K, Csicsvari J, Hirase H, Dragoi G, Buzsáki G. Organization of cell assemblies in
the hippocampus. Nature. 2003;424: 552–556. doi:10.1038/nature01765.1.

46.   Moore GP, Segundo JP, Perkel DH, Levitan H. Statistical Signs of Synaptic Interaction in
Neurons. Biophys J. 1970;10: 876–900. doi:10.1016/S0006-3495(70)86341-X

47.   London M, Schreibman A, Haä;usser M, Larkum ME, Segev I. The information efficacy
of a synapse. Nat Neurosci. 2002;5: 332–340. doi:10.1038/nn826

48.   Pinault D. The juxtacellular recording-labeling technique. Neuromethods. 2011.
doi:10.1007/978-1-60327-202-5_3

49.   Weiler N, Wood L, Yu J, Solla SA, Shepherd GMG. Top-down laminar organization of
the excitatory network in motor cortex. Nat Neurosci. 2008; doi:10.1038/nn2049

50.   Ladenbauer J, McKenzie S, English DF, Hagens O, Ostojic S. Inferring and validating
mechanistic models of neural microcircuits based on spike-train data. bioRxiv Prepr. 2018;
doi:10.1101/261016

51.   McFarland JM, Cui Y, Butts DA. Inferring Nonlinear Neuronal Computation Based on
Physiologically Plausible Inputs. PLoS Comput Biol. 2013;9. doi:10.1371/journal.pcbi.1003143

52.   Robinson BS, Berger TW, Song D. Identification of stable spike-timing-dependent
plasticity from spiking activity with generalized multilinear modeling. Neural Comput. 2016;
doi:10.1162/NECO_a_00883

53.   Song D, Robinson BS, Berger TW. Identification of Short-Term and Long-Term
Functional Synaptic Plasticity From Spiking Activities. Adaptive Learning Methods for Nonlinear
System Modeling. Elsevier; 2018. pp. 289–312. doi:10.1016/B978-0-12-812976-0.00017-8

54.   Stevenson I, Koerding K. Inferring spike-timing-dependent plasticity from spike train data.
Adv Neural Inf Process Syst. 2011; 1–9. Available:
http://machinelearning.wustl.edu/mlpapers/paper_files/NIPS2011_1399.pdf

55.   Linderman S, Stock C, Adams R. A framework for studying synaptic plasticity with neural
spike train data. Adv Neural Inf. 2014; Available: http://arxiv.org/abs/1411.4077

56.   Amidi Y, Nazari B, Sadri S, … UE-2018 40th A, 2018 U. Parameter Estimation in Synaptic
Coupling Model Using a Point Process Modeling Framework*. ieeexplore.ieee.org. 2018;
doi:10.1109/EMBC.2018.8512815

57.   Bayat Mokhtari E, Lawrence JJ, Stone EF. Data Driven Models of Short-Term Synaptic
Plasticity. Front Comput Neurosci. 2018; doi:10.3389/fncom.2018.00032

58.   Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, et al.
Reconstruction and Simulation of Neocortical Microcircuitry. Cell. 2015;163: 456–492.
doi:10.1016/j.cell.2015.09.029

59.   Karmarkar UR, Buonomano D V. Timing in the Absence of Clocks: Encoding Time in
Neural Network States. Neuron. 2007; doi:10.1016/j.neuron.2007.01.006

60.   Klyachko VA, Stevens CF. Excitatory and feed-forward inhibitory hippocampal synapses
work synergistically as an adaptive filter of natural spike trains. PLoS Biol. 2006;4: 1187–1200.
doi:10.1371/journal.pbio.0040207

61.   Swadlow HA, Gusev AG. Receptive-field construction in cortical inhibitory interneurons.
Nature Neuroscience. 2002. doi:10.1038/nn847

62.   Kawaguchi Y. Distinct firing patterns of neuronal subtypes in cortical synchronized
activities. J Neurosci. 2001; doi:21/18/7261 [pii]

63.   Swadlow HA. Thalamocortical control of feed-forward inhibition in awake somatosensory
"barrel" cortex. Philos Trans R Soc Lond B Biol Sci. 2002;357: 1717–27.
doi:10.1098/rstb.2002.1156

64.   Keine C, Rübsamen R, Englitz B. Signal integration at spherical bushy cells enhances
representation of temporal structure but limits its range. Elife. 2017;6. doi:10.7554/eLife.29639

65.   Keine C, Rubsamen R, Englitz B. Inhibition in the auditory brainstem enhances signal
representation and regulates gain in complex acoustic environments. Elife. 2016;5.
doi:10.7554/eLife.19295

66.   Mora Lopez C, Putzeys J, Raducanu BC, Ballini M, Wang S, Andrei A, et al. A Neural
Probe with Up to 966 Electrodes and Up to 384 Configurable Channels in 0.13 µm SOI CMOS.
IEEE Trans Biomed Circuits Syst. 2017;11: 510–522. doi:10.1109/TBCAS.2016.2646901

67.   Jun JJ, Steinmetz NA, Siegle JH, Denman DJ, Bauza M, Barbarits B, et al. Fully integrated
silicon probes for high-density recording of neural activity. Nature. 2017;551: 232–236.
doi:10.1038/nature24636

68.   Pachitariu M, Steinmetz N, Kadir S, Carandini M, Harris KD. Kilosort: realtime spike-
sorting for extracellular electrophysiology with hundreds of channels [Internet]. bioRxiv. 2016.
doi:10.1101/061481

69.   Dale H. Pharmacology and Nerve-endings. J R Soc Med. 1935;28: 319–332.
doi:10.1177/003591573502800330

70.   Tsodyks M, Pawelzik K, Markram H. Neural networks with dynamic synapses. Neural
Comput 1998 May;10(4)821-835. 1998;835: 821–835. doi:10.1162/089976698300017502

71.   Tsodyks M V., Markram H. The neural code between neocortical pyramidal neurons
depends on neurotransmitter release probability. Proc Natl Acad Sci. 1997;94: 719–723.
doi:10.1073/pnas.94.2.719

72.   Hatsopoulos NG, Xu Q, Amit Y. Encoding of Movement Fragments in the Motor Cortex.
J Neurosci. 2007;27: 5105–5114. doi:10.1523/JNEUROSCI.3570-06.2007

1061

1062

Figure 6— figure supplement 1: Spike transmission probabilities patterns and cross-
correlograms for each of the microcircuits identified from the multi-electrode array recording.
Excitatory inputs (cross-correlograms at left with corresponding transmission probability (Prob.)
numbers) to the putative inhibitory neurons are shown along with selected outgoing cross-
correlograms used to identify putative inhibitory neurons (right). Spike wave-shapes are shown in
each circle. Top panel shows the disentangled paired-pulse ratios using simulations following
model fits – membrane integration (yellow); STP (blue); integration and STP (orange). Numbers
on each line connecting putative excitatory neurons and the inhibitory neuron corresponds to depth
differences between recording electrodes.

1072

1073

1074

1075

1076

1077

1078

1079

1080

37