Modeling the short-term dynamics of *in vivo* excitatory spike transmission

Abed Ghanbari[1], Naixin Ren[2], Christian Keine[3], Carl Stoelzel[2], Bernhard Englitz[4], Harvey A. Swadlow[2], and Ian H. Stevenson[1,2]

[1] University of Connecticut, Department of Biomedical Engineering

[2] University of Connecticut, Department of Psychological Sciences

[3] Carver College of Medicine, Iowa Neuroscience Institute, Department of Anatomy and Cell Biology, University of Iowa

[4] Radboud University, Donders Institute for Brain, Cognition and Behavior, Department of Neurophysiology

Corresponding author: ian.stevenson@uconn.edu

Abstract (250 words)

Information transmission in neural networks is influenced by both short-term synaptic plasticity (STP) as well as non-synaptic factors, such as after-hyperpolarization currents and changes in excitability. Although these effects have been widely characterized *in vitro* using intracellular recordings, how they interact *in vivo* is unclear. Here we develop a statistical model of the short-term dynamics of spike transmission that aims to disentangle the contributions of synaptic and non-synaptic effects based only on observed pre- and postsynaptic spiking. The model includes a dynamic functional connection with short-term plasticity as well as effects due to the recent history of postsynaptic spiking and slow changes in postsynaptic excitability. Using paired spike recordings, we find that the model accurately describes the short-term dynamics of *in vivo* spike transmission at a diverse set of identified and putative excitatory synapses, including a thalamothalamic connection in mouse, a thalamocortical connection in a female rabbit, and an auditory brainstem synapse in a female gerbil. We illustrate the utility of this modeling approach by showing how the spike transmission patterns captured by the model may be sufficient to account for stimulus-dependent differences in spike transmission in the auditory brainstem (endbulb of Held). Finally, we apply this model to large-scale multi-electrode recordings to illustrate how such an approach has the potential to reveal cell-type specific differences in spike transmission *in vivo*. Although short-term synaptic plasticity parameters estimated from ongoing pre- and postsynaptic spiking are highly uncertain, our results are partially consistent with previous intracellular observations in these synapses.

Significance Statement (120 words)

Although synaptic dynamics have been extensively studied and modeled using intracellular recordings of post-synaptic currents and potentials, inferring synaptic effects from extracellular spiking is challenging. Whether or not a synaptic current contributes to postsynaptic spiking depends not only on the amplitude of the current, but also on many other factors, including the activity of other, typically unobserved, synapses, the overall excitability of the postsynaptic neuron, and how recently the postsynaptic neuron has spiked. Here we developed a model that, using only observations of pre- and postsynaptic spiking, aims to describe the dynamics of *in vivo* spike transmission by modeling both short-term synaptic plasticity and non-synaptic effects. This approach may provide a novel description of fast, structured changes in spike transmission.

Introduction (650 words)

In response to a presynaptic input, the amplitudes of elicited postsynaptic potentials (PSPs) can increase or decrease dramatically due to short-term synaptic plasticity (Zucker and Regehr, 2002; Regehr, 2012). The probability that a postsynaptic neuron spikes in response to a presynaptic spike can also increase or decrease depending on the recent history of pre- and postsynaptic activity

(Usrey et al., 2000; Swadlow and Gusev, 2001). Although many models exist to describe
intracellular observations of short-term synaptic plasticity (Costa et al., 2013; Hennig, 2013; Barri
et al., 2016; Bird et al., 2016), most models of *functional* connections between neurons based on
extracellular spike observations assume that connections are fixed over time (Truccolo et al., 2005;
Pillow et al., 2008). Unlike intracellular PSP observations, where the amplitude of each individual
presynaptic spike can be measured (subject to noise), extracellular spike observations are sparse,
typically all-or-none binary events. Modeling dynamic, functional connections from spike
observations, especially in the presence of uncontrolled, ongoing neural activity, presents a major
statistical challenge (Ghanbari et al., 2017). Here we further develop a model-based approach that,
given only pre- and postsynaptic spike observations, estimates the contributions of short-term
synaptic plasticity and several non-synaptic factors to the probability of spike transmission.

Traditionally, the influence of presynaptic spikes on postsynaptic spiking is measured using cross-
correlation (Perkel et al., 1967; Fetz et al., 1991; Csicsvari et al., 1998; Barthó et al., 2004). If two
neurons are monosynaptically connected, the probability of the postsynaptic neuron spiking will
briefly increase or decrease following a presynaptic spike, which appears as a fast-onset, short-
latency peak or trough in the cross-correlation, depending on whether the synapse is excitatory or
inhibitory (Perkel et al., 1967; Barthó et al., 2004). Just as synaptic potentials depress or facilitate
due to short-term synaptic plasticity, this spike transmission probability might also depend on the
recent history of presynaptic activity. By subdividing cross-correlograms to characterize the
specific effects of different presynaptic spike patterns, previous studies have found that certain,
putative synaptic connections show reduced spike transmission probability following recent
presynaptic spikes (Swadlow and Gusev, 2001; English et al., 2017), while others show increased
probability (Usrey et al., 2000), as might be expected of depressing or facilitating synapses,
respectively.

Here, rather than subdividing correlograms, we use a likelihood-based modeling approach that
extends previous static models of functional connectivity (Harris et al., 2003; Pillow et al., 2008;
Stevenson et al., 2008). This dynamic model describes not only the sign and strength of synaptic
connections, but also whether the dynamics are depressing or facilitating. In addition to describing
differences in responses to specific presynaptic spike patterns, the model-based approach also
allows us to predict how the postsynaptic neuron will respond to arbitrary patterns of presynaptic
activity. In previous work, we evaluated this type of dynamical functional connectivity model on
simulated and *in vitro* experiments where the ground-truth dynamics were known (Ghanbari et al.,
2017). These results demonstrated that, at least in a controlled setting, short-term synaptic
plasticity can be inferred from spike observations, even in the presence of sources of error, such
as spike sorting errors, stochastic vesicle release, and common input from unobserved neurons.
Here we build on this model and examine how well it can account for excitatory spike transmission
dynamics observed *in vivo* where the true synaptic currents are unknown.

A key element of our dynamical functional connectivity model is the inclusion of both synaptic
and non-synaptic contributions to spike transmission. For each individual presynaptic spike, our
model predicts postsynaptic spiking by taking into account synaptic coupling with STP, synaptic
summation, post-spike history effects, and slow fluctuations of excitability. Although these effects
do not include all factors that may influence spiking statistics (Herz et al., 2006), together they can
account for wide variety of phenomena, including subthreshold membrane integration (Carandini
et al., 2007) and slower fluctuations in the overall excitability of the postsynaptic neuron, such as
observed during neuromodulation (Henze and Buzsáki, 2001). The interaction between synaptic
and non-synaptic effects, as well as the degree to which each factor contributes is likely to lead to
diverse patterns of spike transmission. Here we show how models of dynamical functional
connectivity with short-term synaptic plasticity can capture these patterns of spike transmission
and disentangle the multiple factors that shape postsynaptic response.

Material and methods

Neural Data

All data analyzed here were obtained from previous studies (see below). Animal use procedures
were approved by the institutional animal care and use committees at University of Connecticut
(VB-Barrel), University of Leipzig (ANF-SBC), or University College London (MEA),
respectively, and conform to the principles outlined in the Guide for the Care and Use of
Laboratory Animals (National Institutes of Health publication no. 86-23, revised 1985).

To illustrate how synaptic dynamics can be estimated from spikes, we first examined a set of three
strong putative or identified synapses with diverse spike transmission probability patterns: (i) a
local, excitatory connection from one neuron in mouse thalamus to another detected from a larger
multi-electrode array (MEA) recording, (ii) a ventrobasal thalamus projection to primary
somatosensory cortex (VB – Barrel) in a rabbit, and (iii) an *in vivo* loose-patch (juxtacellular)
recording of an auditory nerve projection onto a spherical bushy cell (ANF-SBC) in the auditory
brainstem of a gerbil. We then use this auditory brainstem connection to explore how synaptic
transmission probability depends on the stimulus and compare the results with a model without
short-term synaptic plasticity. Next, we applied our model more generally to analyze a large
sample of putative synaptic connections recorded from the MEA dataset. The data from these three
identified strong synapses and the MEA data have been collected from different species, regions,
cell-types, under different stimulation and show a diverse pattern of postsynaptic spiking
probability. In all cases we deduce short-term synaptic dynamics on the basis of only pre- and
postsynaptic spike observations.

For the first putative synapse, we use *in vivo* data from simultaneous extracellular recordings in
ventrobasal (VB) thalamic barreloids and topographically aligned, somatosensory cortical barrel
columns (VB-Barrel) in awake, unanesthetized, adult rabbits. Detailed surgical and physiological

methods have been described previously (Swadlow and Gusev, 2002). Spike-triggered averages
of the cortical spikes following spiking of the VB neuron was used to identify connected S1
neurons. Based on the presence of high frequency discharge (3+ spikes, > 600 Hz) following
electrical stimulation of the thalamus, and narrow spike waveforms, the S1 neuron in this recording
was identified as a putative inhibitory neuron (Kawaguchi, 2001). These recordings identified
several putative thalamocortical projections. The putative synapse that we model here is
particularly clear, with 68,345 pre- and 128,096 postsynaptic spikes recorded over the course of
92 minutes of spontaneous activity and has been previously studied in (Swadlow and Gusev, 2001;
Swadlow, 2002).

For the second synapse, we examined *in vivo* loose-patch recordings at the Endbulb of Held in
young adult female gerbils. Detailed surgical and physiological methods have been previously
described (Keine et al., 2017). Briefly, the glass electrode was positioned in the anterior portion of
the ventral cochlear nucleus (AVCN) and single-units were recorded during varying acoustic
stimulation. Single units were classified when recording a positive action potential amplitude of at
least 2 mV and showing the characteristic complex waveform identifying them as large spherical
bushy cells (SBC) of the rostral AVCN. This recording included a mixture of juxtacellular
waveforms: an isolated excitatory PSP (EPSP) or an EPSP followed by a postsynaptic action
potential. For both cases the timing of EPSPs and spikes and rising slope of the EPSPs were
extracted. The timing and slope of the EPSPs were identified using a slope threshold for the rising
part of EPSPs as previously described (Keine et al., 2016). We then modeled spike transmission
probability patterns for two recordings: (i) during randomized pure tone acoustic stimulation and
(ii) during multiple stimuli, i.e. randomized frequency-level pure tone stimulation interspaced with
spontaneous activity, natural sounds, and also during spontaneous activity. Using this second
dataset, we characterized how variable presynaptic spike patterns evoked by different stimuli
affected the patterns of spike transmission at the same synapse.

We also use MEA spiking data to study the factors shaping spike transmission probability patterns
in a large-scale recording with multiple cell-types. Here we use a previously collected, publicly
available recording from the Cortex Lab at UCL (Jun et al., 2017; Mora Lopez et al., 2017) with
data from two Neuropixels electrode arrays recorded simultaneously, each with 960 sites (384
active) with lengths of 10-mm and spacing of 70 × 20-μm (http://data.cortexlab.net/dualPhase3/).
The two electrode arrays span multiple brain areas and ~90 min of data was collected in an awake,
head-fixed mouse on a rotating rubber wheel during visual stimulus presentations. Spikes were
automatically detected and sorted using Kilosort (Pachitariu et al., 2016) on the broadband (0.3–
10 kHz) signal and then manually curated. If two clusters of spikes had similar waveforms, cross-
correlogram features, and spike amplitudes, they were merged into a single cluster and assigned
to a single neuron. In total, 831 well-isolated single neurons where identified from the two probes
in several different brain areas: visual cortex (n=74), hippocampus (n=64), thalamus (n=244),

motor cortex (n=243), and striatum (n=200). Due to the large number of simultaneously recorded
neurons in this dataset, there are many potential synapses (~831^2).

Synapse Detection

To identify putative monosynaptic connections between well-isolated single neurons, we looked
for specific patterns in the cross-correlograms (Moore et al., 1970). If two neurons are
monosynaptically connected, the probability of postsynaptic spiking increases/decreases rapidly
following a presynaptic spike. In spiking data, this rapid, transient change can be seen in cross-
correlograms as an asymmetric bump/dip in the number of postsynaptic spikes following
presynaptic spikes (Barthó et al., 2004). For each connection we calculated the cross-correlogram
in a 5 ms window before and after presynaptic spikes with bin-size of 0.1 ms. To avoid aliasing in
the cross-correlograms, we added a small, random shift to each postsynaptic spike drawn
uniformly between $-\Delta t/2$ and $\Delta t/2$ where Δt is the spike time resolution (0.01 ms in most cases).
Here we used a model-based approach using the cross-correlograms to decide whether two
synapses are monosynaptically connected. To fit the cross-correlogram we used a baseline rate μ,
a linear combination of B-spline bases $\mathbf{B}(t)$, and a weighted alpha function to model the synapse,
$w\,\alpha(t)$, all passed through an output nonlinearity; $\lambda(t) = \exp\big(\mu + \mathbf{r}\mathbf{B}(t) + w\,\alpha(t)\big)$. The alpha
function, $\alpha(t) = (t - t_d)/\tau_\alpha \exp(1 - (t - t_d)/\tau_\alpha)$, describes the shape of the synaptic potential
where t_d is the synaptic delay and τ_α is the synaptic time-constant (Carandini et al., 2007). For
individual connections, we estimate these parameters by maximizing the penalized Poisson log-
likelihood $l(\mu, \mathbf{r}, w, t_d, \tau_\alpha) = \Sigma y_i log \lambda_i - \Sigma \lambda_i + \epsilon\|\mathbf{r}\|_2$ where y_i is the number of postsynaptic
spikes observed in the i-th bin of the correlogram and $\|\mathbf{r}\|_2$ regularizes the model to penalize B-
spline bases for capturing sharp increases in the cross-correlogram. ϵ is a regularization hyper-
parameter which we set to 1 based on manual search. Due to the parameterization of $\alpha(t)$, the log-
likelihood is not concave. However, since the gradient of the log-likelihood can be calculated
analytically, we efficiently optimize the likelihood using a gradient-based pseudo-Newton method
(LBFGS) (Boyd and Vandenberghe, 2004). During the optimization, the delay and time-constant
parameters are log-transformed, allowing us to use unconstrained optimization, even though they
are strictly positive. We used random restarts to avoid local maxima. To identify putative
monosynaptic connections in the large-scale multi-electrode array data, we compared this model
with a smooth model with slow changes in cross-correlogram and without the synapse, $\lambda_0(t) =$
$\exp(\mu' + \mathbf{r}'\mathbf{B}(t))$, using the log-likelihood ratio (LLR) test between our full model with synapse
and the nested smooth model. Since low values of the likelihood ratio mean that the observed result
was better explained with full model as compared to the smooth model, we then visually screened
pair-wise connections with lowest ratios (LLR <-6) compared to the null model to find putative
synapses. Out of ~831^2 possible connections in this dataset we find ~200 putative synapses
(0.03%). We handpicked a strong putative synapse between two thalamic neurons to study its
efficacy pattern in detail alongside the VB-Barrel and ANF-SBC synapses.

In addition to this single strong synapse, we also categorize putative pre- and postsynaptic cell
types for the connections detected in the MEA dataset. For this purpose, we assessed single units
based on their cross-correlograms, firing rates, and spike waveforms. We categorized units as
excitatory or inhibitory if, in accordance with Dale's law, all outgoing cross-correlograms showed
transient, short-latency (<4ms) increase/decrease in spiking probability. We then looked into
identified inhibitory neurons and categorized them into to putative fast-spiking (FS) and regular-
spiking (RS) inhibitory neurons. Using these putative Excitatory-FS and Excitatory-RS synapses,
we then examine how the spike transmission patterns differ for these two subtypes of inhibitory
neurons.

Extending a Generalized Linear Model to Account for Short-term Plasticity (TM-GLM)

Short-term synaptic plasticity causes the amplitude of postsynaptic potentials (PSP) to vary over
time depending on the dynamics of synaptic resources and utilization and can be modeled using
the pattern of presynaptic spiking (Markram et al., 1998; Tsodyks et al., 1998). However, changes
in the overall postsynaptic spiking probability cannot be uniquely attributed to changes in
amplitudes of postsynaptic potentials. To accurately describe the dynamics of spike transmission,
we also need to account for the membrane potential summation, the excitability of the postsynaptic
neuron (e.g. slow changes in the presynaptic firing rate) and the dynamics of postsynaptic spiking
(e.g. refractory period, after hyperpolarization current). We developed an extension of a
generalized linear model, which we call a TM-GLM to describe each of these effects. Concretely,
the probability of a postsynaptic spike shortly after each presynaptic spike accounts for the full
sequence of previous presynaptic spiking and the recent history of postsynaptic spiking. We define
the conditional intensity of the postsynaptic neuron after the i-th presynaptic spike, $t_s^{(i)}$, so that the
probability of observing a postsynaptic spike in the j-th time bin after the i-th presynaptic spike is
given as:

$$220 \quad \lambda_i(t_j) = \sigma\left(\beta_0 + \mathbf{X}_c\left(t_s^{(i)}\right)\boldsymbol{\beta}_c + \sum_{t_r^{(l)} < t_s^{(i)}} \mathbf{X}_h\left(t_s^{(i)} - t_r^{(l)}\right)\boldsymbol{\beta}_h + A_s\, w_i\, \alpha(t_j) \right)$$

221 where $t_r^{(l)}$ are the postsynaptic spike times preceding $t_s^{(i)}$. For each presynaptic spike, our model
222 decomposes the firing rate of the postsynaptic neuron into four effects: a baseline firing rate, β_0,
223 slow fluctuations in postsynaptic firing rate $\mathbf{X}_c\boldsymbol{\beta}_c$, history effects from the recent postsynaptic
224 spikes (prior to $t_s^{(i)}$), $\mathbf{X}_h\boldsymbol{\beta}_h$, and a time-varying coupling effect from the presynaptic input,
225 $A_s w\, \alpha(t)$ (Fig. 1).

Fig. 1: TM-GLM. Postsynaptic spiking probability before passing the spiking nonlinearity (yellow) changes as a linear combination of presynaptic coupling term with STP dynamics (blue), postsynaptic spiking history (green), the postsynaptic excitability (red). Transparent red curves show the bases of slow changes in postsynaptic probability at presynaptic spike times (X_c).

Here we model slow fluctuations in the postsynaptic rate $\mathbf{X_c}\beta_c$ with a linear combination of B-splines with equally spaced knots every 50 seconds of recording time. In the history term, splines ($\mathbf{X_h}$) span a period of 10 ms prior to each presynaptic spike with 4 logarithmically-spaced knots. By scaling $\alpha(t_j)$ with a multiplicative factor, w_i, the strength of a synapse can vary over time and, in this case, depends on the detailed sequence of presynaptic spiking and their corresponding inter-spike intervals. A_s is the magnitude of the synaptic strength. In this case we use a model for short-term synaptic plasticity that allows both depression (where the w_i decreases for shorter presynaptic ISIs) and facilitation (where the w_i increases for shorter presynaptic ISIs), and incorporates membrane summation. To model these effects, w_i is determined by a nonlinear dynamical system based on the Tsodyks and Markram (TM) model (Tsodyks and Markram, 1997; Markram et al., 1998) where: $w_i = w_{i-1} \exp\left(-\frac{t_s^{(i)} - t_s^{(i-1)}}{\tau_s}\right)\pi_i + R_i u_i$, where τ_s is the membrane time-constant and the first term of the equation describes how postsynaptic membrane potential summation increases the probability of postsynaptic spiking. This membrane summation will be ignored if there is a postsynaptic spike: $\pi_i = \left\{0 \text{ if } t_s^{(i-1)} < t_r^{(i-1)} < t_s^{(i)}; 1 \text{ otherwise}\right\}$. In the second term of this equation, R represents the dynamics of resources and u describes their utilization.

$$R_i = 1 - [1 - R_{i-1}(1 - u_{i-1})] \exp\left(-\frac{t_s^{(i)} - t_s^{(i-1)}}{\tau_d}\right)$$

$$u_i = U + [u_{i-1} + f(1 - u_{i-1}) + U] \exp\left(-\frac{t_s^{(i)} - t_s^{(i-1)}}{\tau_f}\right)$$

where τ_d and τ_f are the depression and facilitation time-constants. U is the release probability, and
f is the magnitude of facilitation. To make the estimation more tractable, we approximate the full
optimization problem and estimate synaptic delay, t_d, and time-constant, τ_α, by fitting $\alpha(t)$ using
the full cross-correlogram, as above. We fix these parameters for the rest of the optimization
process. We then maximize a penalized, Bernoulli log-likelihood $\log(l(\boldsymbol{\theta})) = \Sigma\Sigma\left[y_{ij}\lambda_i(t_j) - \right.$
$\left.(1 - y_{ij})(1 - \lambda_i(t_j))\right] + \gamma\|\boldsymbol{\theta}'_{stp}\|_2$ where $\gamma = 1$ is the regularization hyperparameter to estimate
the parameters: $\boldsymbol{\theta} = \{\beta_0, \boldsymbol{\beta}_{c=1:C}, \boldsymbol{\beta}_{h=1:H}, A_s, \boldsymbol{\theta}_{stp}\}, \boldsymbol{\theta}_{stp} = \{\tau_d, \tau_f, U, f, \tau_s\}$.

As with previous applications of GLMs, we assume that bins are conditionally independent given
the covariates, but unlike many other GLMs, here we only calculate the log-likelihood during short
intervals (5ms) after presynaptic spikes. With y_{ij} being a binary value representing the presence of
a postsynaptic spike in the j-th time bin after the i-th presynaptic spike. We again used a
logarithmic transformation for the time-constants to avoid negative values and logit transformation
for U and f to bound their values in the interval $[0, 1]$; $\boldsymbol{\theta}'_{stp} =$
$\{\log(\tau_d), \log(\tau_f), \text{logit}(U), \text{logit}(f), \log(\tau_s)\}$. By modeling STP this model is no longer a strict
GLM, and the log-likelihood may have local maxima. Here we use random restarts to avoid local
maxima in our optimization process. The parameters of each restart $\{\beta_0, \boldsymbol{\beta}_{c=1:C}, \boldsymbol{\beta}_{h=1:H}, A_s\}$ are
initialized by adding noise ($\sim N(0,1)$) to the corresponding parameters in a standard GLM. We
initialize the log-transformed plasticity parameters with $\tau'^{(0)}_d \sim N(-1,5)$, $\tau'^{(0)}_f \sim N(-1,5)$,
$U'^{(0)} \sim N(0,5), f'^{(0)} \sim N(0,5), \tau'^{(0)}_s \sim N(-3,5)$. We then use an LBFGS algorithm to optimize the
log-likelihood where we calculate all derivatives analytically except for derivatives of θ_{stp} which
we calculate numerically. To estimate the uncertainty of the parameters, we bootstrap the data
from each of the strong synapses by chunking the whole recording time into samples of 50 seconds
then resampling the chunks to generate a new spike train with the same original length.

Calculating spike transmission probability

To demonstrate how the probability of postsynaptic spiking changes according to the
corresponding presynaptic inter-spike intervals, we estimated spike transmission probabilities
from the cross-correlograms directly instead of using a model. To calculate this probability, we
focused on a transmission interval after the presynaptic spike where the conditional intensity (when
corrected for the baseline rate) goes above 10% of the maximum of $\boldsymbol{\alpha}(t)$ (horizontal bars in Fig
2A). We split the presynaptic inter-spike interval distribution into log-spaced intervals, and, for
each interval, we calculate the ratio between numbers of postsynaptic spikes in the transmission
interval to the number of presynaptic spikes. Unlike previous studies (Swadlow and Gusev, 2001,
2002) we do not correct this probability for the baseline postsynaptic rate. The uncorrected
probability allows us to more directly compare the model predictions to the empirical spike
transmission probabilities. Since our model gives an estimate of the postsynaptic probability after

each individual presynaptic spike, we can average over the same transmission interval. However,
we know if there is a postsynaptic spike in the transmission interval, probability of a postsynaptic
spike goes to ~0 for all consecutive bins due to the post-spike dynamics (e.g. refractory period).
Therefore, we measure the predicted probability of a postsynaptic spike in a 5ms window after i-
th presynaptic spike from binned $\lambda_i(t_j)$ as follows: $z_i = \sum_{j=1}^{L} \lambda_i(t_j) \prod_{m=1}^{j-1}(1 - \lambda_i(t_m))$. Here we
assume conditional independence of the j-th bin after a presynaptic spike, but we enforce a
refractory period for all bins after a postsynaptic spike in our generative model. Here L is the first
bin that y_{ij} is nonzero. z_i represents the probability of postsynaptic spiking after each presynaptic
spike and we fit a smooth curve over the distribution of z_i's and their corresponding inter-spike
intervals to compare with the empirical spike probability patterns.

Modeling the effect of local patterns of pre- and postsynaptic spiking

The observed and modeled spike transmission patterns, as calculated above, reflect the expected
postsynaptic spike probability given a specific presynaptic ISI. However, since the presynaptic
ISIs are not independent and there are serial correlations in ISIs, the detailed sequence of the pre-
and postsynaptic spiking likely affects the shapes of these curves. To quantify the effects of serial
ISI correlations on the model of spike transmission probability we demonstrate how local patterns
of presynaptic spiking modifies spike transmission patterns in the data and the model. For each of
the three strong identified synapses we measure postsynaptic spiking probability in response to
presynaptic spike triplets. Due to the limited number of spikes in our data, we divide the
presynaptic ISI distribution into few log-spaced intervals and measure the postsynaptic spiking
probability for triplets with the two ISIs that fall in those intervals. Similarly, we measure the
predicted postsynaptic probability in response to the presynaptic triplets. After measuring
postsynaptic responses to presynaptic spike triplets in the data and the model, we simulate the
contribution of STP in shaping the transmission pattern in response to these triplets. To factor out
contributions of the postsynaptic history and slow changes in presynaptic firing rate, we fix the
corresponding values in the model to their average values within the model. In these simulations,
we also fix the initial values of the STP dynamics in the TM model for the first spike of the triplets
to the average R and u within the model. This approach enables us to illustrate how short-term
synaptic plasticity in triplets of presynaptic spikes changes spike transmission probability and how
serial correlations in presynaptic spiking affect spike transmission probability.

The postsynaptic spike history and the serial correlations between the pre- and postsynaptic spiking
also modify spike transmission probability patterns. To investigate history effects in the local
pattern of pre- and postsynaptic spikes, we measured the postsynaptic spiking probability in
response to two presynaptic spikes and a postsynaptic spike preceding the most recent presynaptic
spike. Due to the limited number of spikes and sparseness of the split cross-correlograms, we again
divided the presynaptic and postsynaptic ISI distributions into a few log-spaced intervals. We then
measure the spike transmission probability for a group of presynaptic spikes that their preceding

presynaptic ISIs and postsynaptic spike ISIs fall into different combinations of pre- and
postsynaptic log-spaced intervals. After measuring postsynaptic responses to any possible
combination of the two most recent presynaptic spikes and their postsynaptic spikes in the data
and the model, we simulate the contribution of the history and STP together in shaping the
transmission. In our simulation the excitability was set to the model estimates. To measure the
effects of postsynaptic spiking history, for each postsynaptic ISI, we fix the history contribution
to estimated post-spike history filter value at that postsynaptic ISI. We then use the predicted STP
parameters from the data to simulate the STP contribution in response to paired pulses of
presynaptic ISIs where we again fix the initial values of the TM model for the first presynaptic
spike to the average R and u within the model. This approach enables us to illustrate how short-
term synaptic plasticity in local patterns of two presynaptic spikes and a postsynaptic spike
changes spike transmission probability and quantifies how serial correlations between pre- and
postsynaptic spiking affect spike transmission probability.

Evaluating prediction accuracy

In addition to evaluating the estimated parameters and comparing the model to empirical spike
transmission probabilities, we also assess how accurately the model can predict postsynaptic
spiking. Not only can we predict the probability of a spike given specific presynaptic ISIs, but we
can also predict whether there will be a postsynaptic spike following each individual presynaptic
spike. To quantify how well the predicted postsynaptic spike probability, z_i, predicts the
postsynaptic spiking activity, we use Receiver Operating Characteristic (ROC) curves. To compute
the ROC curve, we first create a threshold version of z_i which operates as our prediction: $\{(\hat{r}_i =$
$1)$ if $(z_i > \text{thr}); 0 \text{ otherwise}\}$. Changing the threshold from 0 to 1 traces out a relationship between
the true positive rate (TPR) and false positive rate (FPR). The area under the ROC curve (AUC)
reflects the performance of each model, where a perfect classifier has AUC=1 and a random
classifier has AUC=0.5. Effectively, the AUC is the probability of a randomly chosen spike having
a higher model probability than a randomly chosen non-spike (Hatsopoulos et al., 2007). Here we
calculate the AUC for short intervals (~5ms) after presynaptic spikes and check whether we detect
a postsynaptic spike in the transmission interval where $\alpha(t)$ is above 10% of its maximum. Here
we compare the AUC for the static model of connectivity without short-term synaptic plasticity
with our dynamical model.

A simplified rate model to simulate effects of synaptic summation and post-spike history

Our TM-GLM's prediction of the spike transmission pattern is data-driven and depends on the full
history of pre- and postsynaptic spiking. To better understand and illustrate how STP, synaptic
summation, and post-spike history interact to create the observed patterns of spike transmission,
we simulated postsynaptic responses in a simplified voltage model. Namely, we consider PSP
summation in response to a pattern of two presynaptic spikes. We assume that the synapse is
initially fully recovered, and the PSC amplitudes are determined by the 4-paramter TM model with

$U = 0.7, \tau_d = 1.7, \tau_f = 0.02, f = 0.05$ for the depressing synapse and $U = 0.1, \tau_d = 0.02, \tau_f =$
$1, f = 0.11$ for the facilitating synapse (Ghanbari et al., 2017). We then convolve the PSCs (delta
function kernel) with a PSP kernel, $\exp(-t/\tau_v) - \exp(-t/\tau_r)$, with $\tau_v = .01$ and $\tau_r = .001$ ms to
describe synaptic summation. We assume that the instantaneous postsynaptic spike probability is
simply a nonlinear function of the distance to a threshold voltage $\sigma\big(5(V(t) - V_{th})\big)$ where $\sigma(x) =$
$1/(1 + e^{-x})$ and $V_{th} = .5, .75,$ and 1 correspond to strong, moderate, and weak inputs
respectively. The spike transmission probability sums this instantaneous probability over a
window of 20ms after each presynaptic spike. Finally, we adjust the spike transmission probability
for the second PSP to account for potential post-spike history effects. Namely, we assume that the
adjusted spike transmission probability for the second spike is $p_2^* = (1 - p_1)p_2 + p_1 p_2 f_{ahp}$ where
p_1 is the transmission probability for the first spike, p_2 is the unadjusted probability for the second
spike, and f_{ahp} is the effect of the after-hyperpolarization. Here we use $f_{ahp}(\Delta t) = (\sigma\big(150(\Delta t -$
$0.02)\big) - c)/d$ where Δt is the presynaptic ISI, and c and d are constants ensuring that $f_{ahp}(0) =$
0 and $f_{ahp}(\infty) = 1$. Although this simulation is highly simplified, it demonstrates how the
observed spike transmission pattern depends, not just on the type and timescale of STP, but on the
interaction between STP, synaptic summation, after-hyperpolarization effects, and the spike
nonlinearity.

Simulation of non-connections

The TM-GLM relies on correctly identifying monosynaptic connections. To investigate how our
model performs when there is no actual synapse, we simulated a microcircuit with three neurons
where a presynaptic neuron provides excitatory input to two postsynaptic neurons with different
delays (1 and 3 ms). Here we test how different combinations of STP (depression and facilitation)
in connections between pre- and postsynaptic neurons would impact the overall estimation of spike
"transmission" probability in the spurious connection between the two postsynaptic neurons. Here
the spikes of the presynaptic neuron were simulated from an inhomogeneous Poisson process with
random, smooth rate fluctuations (5Hz average, 4.6Hz sd). The postsynaptic neurons were then
simulated using a leaky integrate-and-fire neuron with spike frequency adaptation (parameters are
from (Ghanbari et al., 2017)) that received a white noise current as input as well as a current-based
synapse from the presynaptic neuron (double exponential with rise time 1ms, decay time 10ms).
The PSCs of the input then vary according to the Tsodyks-Markram model (parameters for
depression/facilitation are as in (Ghanbari et al., 2017)).

Results

Short-term synaptic plasticity directly affects synaptic information processing by altering the
amplitude of presynaptic currents (Abbott and Regehr, 2004). However, in most neural systems it
remains unclear how these presynaptic effects translate to modified postsynaptic spike probability.

Postsynaptic spiking is affected by many factors including short-term plasticity, postsynaptic spike
history, summation of PSPs, and slow fluctuations in excitability. Here we develop a statistical
model that includes each of these factors and allows their effects to be estimated solely using pre-
and postsynaptic spiking activity. We examined the model's ability to capture the observed
patterns of spike transmission probability for three strong putative or identified synapses. We then
use one of these systems (the endbulb of Held synapse in the auditory brainstem), to explore how
the short-term dynamics of spike transmission depend on an external stimulus and compare the
results with a model without short-term synaptic plasticity. Finally, we apply our model to spiking
data from a large-scale, multi-electrode array recorded from multiple areas in an awake mouse.
Here we investigate the STP dynamics in putative synapses from excitatory neurons onto two
putative inhibitory neuron subtypes. We find that these two types of connections have distinct
patterns of spike transmission, consistent with previous experimental observations.

Spike transmission probability varies strongly as a function of presynaptic ISIs

Cross-correlograms of excitatory monosynaptic connections show a rapid, transient increase in the
postsynaptic spiking probability shortly after the presynaptic spike, with a latency of ~2-4ms
(Perkel et al., 1967; Fetz and Gustafsson, 1983; Fetz et al., 1991; Poliakov et al., 1996). The timing
and shape of the cross-correlogram depends on the presynaptic axonal conduction delay, the
synaptic delay, and the strength of the connection. However, in the overall cross-correlogram the
effects of all presynaptic spikes are averaged and any variations in spike transmission, such as
dependence on the history of presynaptic spiking, are hidden (Fig. 2A). To quantify how the history
of presynaptic spiking influences spike transmission probability, the probability of observing a
postsynaptic spike shortly after a presynaptic spike, previous studies have compared the cross-
correlograms for specific subsets of presynaptic spikes. For instance, comparing the cross-
correlograms calculated for presynaptic spikes within defined inter-spike intervals (ISI)
demonstrates how spike transmission probability varies depending on recent presynaptic spiking
(Swadlow and Gusev, 2001; English et al., 2017). Here, to illustrate the diversity of short-term
dynamics in spike transmission, we examine three strong synapses from three distinct neural
systems: (i) a pair of neurons in thalamus in a male mouse, (ii) a projection from ventrobasal
thalamus to somatosensory barrel cortex (VB-Barrel) in a female rabbit, and (iii) the auditory nerve
fiber to spherical bushy cell projection in a female gerbil (ANF-SBC), the endbulb of Held. The
short-term synaptic dynamics of thalamocortical projections, have been extensively characterized
in vivo (Swadlow and Gusev, 2001; Stoelzel et al., 2008, 2009). Similarly, ANF-SBC synapses
have been extensively studied in previous experiments and are well-characterized *in vitro*
(Thomson et al., 2002; Yang and Xu-Friedman, 2008, 2009). The presynaptic neurons in each of
these pairs have distinct ISI distributions (Fig. 2B), and, after splitting the spikes into ISI quantiles
and calculating the correlogram for each quantile, we find that postsynaptic responses differ
following short and long presynaptic ISIs (Fig. 2C). For the pair of thalamic neurons, spike
transmission probability is increased at short and long intervals and reduced for mid-range ISIs

(based on n=62661 presynaptic spikes). For the VB-Barrel connection, transmission probability is
higher for longer ISIs (based on n=68345 presynaptic spikes), while for ANF-SBC the highest
transmission probability occurs at intermediate intervals (based on n=20547 presynaptic spikes).
These three cases illustrate that the short-term dynamics of spike transmission can be highly
diverse between neurons and brain regions.

Fig. 2: Spike transmission probability depends on the presynaptic ISI and differs between synapses.
A) Cross-correlograms between pre- and postsynaptic spiking at three different synapses show an increase
in the postsynaptic spike count (or probability) after a short latency, indicative of a monosynaptic
connection. The efficacy (Eff.) for each synapse is calculated as the ratio between the number postsynaptic
spikes that are above baseline in the transmission interval (denoted by the horizontal bar) and the number
of presynaptic spikes. **B)** Inter-spike interval distributions (log-scale) for the presynaptic neurons. The
distributions are color-coded into 5 quantiles with equal numbers of presynaptic spikes. **C)** We calculate a
separate cross-correlogram using the subset of presynaptic spikes where the preceding spike fell within
each ISI range. Colors correspond to (B) going from shorter presynaptic ISIs (left) to longer ISIs (right).
Note that both the baseline firing rate and the synaptic peak for each connection change as a function of
presynaptic ISI.

The shape of spike transmission patterns depends on multiple factors

One potential explanation for the diverse dynamics of short-term spike transmission (Fig. 2) may
be that some synapses are depressing while others are facilitating. Short-term synaptic plasticity
directly alters postsynaptic currents such that the response after each presynaptic spike depends on
the recent history of presynaptic spiking (Markram et al., 1998; Ghanbari et al., 2017). However,
many factors can influence spike timing in addition to the dynamics of a single synapse. At short
presynaptic ISIs, membrane potential summation can lead to larger PSPs and increased spike
probability, even in absence of short-term synaptic plasticity (Carandini et al., 2007). Additionally,

the spiking nonlinearity and the history of postsynaptic spiking can alter how a given pattern of
presynaptic input is transformed into postsynaptic spiking (Pillow et al., 2008; Huang et al., 2016).
To illustrate how STP, synaptic summation, and postsynaptic history interact to create a particular
spike transmission pattern we performed simulations using a simplified spiking model with linear
voltage summation, short-term plasticity, a soft spiking nonlinearity, and an after-
hyperpolarization (Fig. 3).

Similar to experimental data (Markram et al., 1998; Ghanbari et al., 2017), the spike transmission
probability in this simplified model depends on the presynaptic ISI as well as the type of STP. For
depressing synapses, the spike transmission probability increases for longer presynaptic ISIs while
for facilitating synapses it increases for mid-range ISIs. Independent of STP type, PSPs sum at
short ISIs (Fig. 3A). However, in this model, the exact shape of transmission probabilities also
depends on the strength of the synapse and the history of postsynaptic spiking. An after-
hyperpolarization current following each postsynaptic spike, for instance, can briefly decrease the
probability of subsequent spikes. In our simulation, we find that "spike interference" from previous
postsynaptic activity can counteract membrane potential summation (Fig. 3B). This type of
postsynaptic spike interference generally decreases the spike probability for shorter presynaptic
ISIs, but the magnitude of this decrease depends on the synaptic strength and type of STP (Fig.
3C). Together, these simulations illustrate how patterns of spike transmission probability are the
result of, not just STP, but of the complex interaction between the membrane potential, the spike
nonlinearity, the post-spike history, and short-term synaptic plasticity.

Fig. 3: A simulation of a simplified spiking model shows how spike transmission probability depends
on multiple factors. A: For different types of short-term synaptic plasticity, postsynaptic summation
increases the amplitudes of the postsynaptic potentials (PSP) at shorter ISIs. Lines denote the membrane
potential of a postsynaptic neuron in a simplified model as it responds to short (dark traces) and long (light)
paired presynaptic pulses. Relative amplitudes of excitatory PSPs increase or decrease under the simplified
model depending on the type of STP. **B:** Spike generation changes with synaptic strength. In this paired-

pulse stimulation paradigm, stronger synapses are more likely to generate a spike following the first
presynaptic impulse which can then decrease the spiking probability following the second impulse if there
are post-spike history effects. As in (A) traces denote postsynaptic membrane potential responses to short
(dark) and long (light) presynaptic ISIs. Dashes denote example postsynaptic spiking, with "spike
interference" occurring for strong synapses and short ISIs. **C:** The pattern of spike transmission probability
under the simplified model changes depending on the type of STP, the coupling strength, and presence of
post-spike interference. Dashed lines show transmission probability without interference from previous
postsynaptic spikes, while solid lines show how post-spike history effects can decrease the spike
transmission probability.

Spike transmission patterns are diverse across regions and species

The combination of synaptic and non-synaptic factors could be one explanation for the diversity
of spike transmission patterns in experimental data. Here we aim to model these contributions and
extend a previously developed generalized linear model (GLM) framework for static functional
connections (Harris et al., 2003; Truccolo et al., 2005; Pillow et al., 2008). In the previous, static
GLM the probability of postsynaptic spiking is modeled as a linear combination of a baseline firing
rate parameter, a post-spike history filter to capture the postsynaptic spike dynamics, such as
refractoriness and burstiness, and a coupling filter describing the fixed influence of presynaptic
spikes. The sum of these effects is then passed through a spiking non-linearity. In our extended
model we added a linear term that allows changes in the excitability of the postsynaptic neuron as
a function of time (timescale >1 min) and allow the coupling term to change for each presynaptic
spike according to the Tsodyks and Markram (TM) model of STP (Markram et al., 1998). We fit
the parameters of this TM-GLM using only the pre- and postsynaptic spike observations and obtain
parameters for each effect using approximate maximum likelihood estimation (see Methods). This
provides estimates of the history and coupling filters, as in a static GLM, as well as additional
parameters for the dynamical synapse (TM model), including facilitation, depression, membrane
time-constants, and release probability. Given these parameters, this TM-GLM model provides
estimates of the postsynaptic spiking probability following each observed presynaptic spike and
can also predict spike transmission probabilities in response to arbitrary patterns of presynaptic
inputs.

After fitting the model to pre- and postsynaptic spike-trains, we compared its behavior to
experimentally observed patterns of spike transmission probability. In particular, we compare
peaks in the split cross-correlograms to the average model prediction for the same sets of
presynaptic spikes (see Methods). We find that our model is flexible enough to explain the changes
in spiking transmission probability observed in spiking statistics for all three synapses above
(Fig. 4A). Moreover, using the model-based approach, the contributions of the synaptic and non-
synaptic component can be disentangled. Our results suggest that the pattern of spike transmission
probability for the thalamus connection is dominated by a combination of membrane potential
summation and short-term depression. Although depression decreases spike transmission

probability at shorter ISIs, membrane summation acts to increase postsynaptic spiking. The ANF-
SBC synapse, in contrast, shows an increase in spike transmission probability for a medium range
of ISIs that is explained by a model dominated by short-term facilitation. Lastly, the VB-Barrel
connection shows a higher postsynaptic response for spikes following longer ISIs (isolated) that is
explained by the model as an effect of short-term synaptic depression.

In addition to estimating the contributions of synaptic and non-synaptic factors that affect spike
transmission, the model also improves the prediction of postsynaptic spiking. Although the cross-
correlogram provides an average efficacy for spike transmission, our models provide detailed
predictions of the postsynaptic spike probability following each presynaptic spike. Here we
measure the Receiver Operating Characteristics (ROC curves) of our models during this short
window of time following a presynaptic spike (see Methods). We compare the prediction of
postsynaptic spiking activity in the full, dynamic synapse model and a static synapse model
containing all components except STP. In all three datasets, a model with short-term synaptic
plasticity provides substantially better predictions of the postsynaptic spiking activity. For the
model with short-term synaptic plasticity accuracies were AUC=0.75$\pm$0.005, 0.69$\pm$0.002, and
0.79 $\pm$ 0.011 (mean $\pm$ SE) for the Thalamus pair, VB-Barrel, and ANF-SBC connections,
respectively; compared to a model without STP where the model accuracies were
AUC=0.54$\pm$0.003, 0.48$\pm$0.002, and 0.56$\pm$0.003 (mean$\pm$SE, bootstrapping over presynaptic
spikes). Note that, although static synapse models do account for the average increased probability
of spiking following a presynaptic spike, the fact that the AUC values are near chance (0.5)
indicates that they do not accurately predict which presynaptic spikes will lead to a postsynaptic
response and which will not.

In our model, the short-term dynamics of spike transmission are described by two coupled
differential equations with five parameters: $\theta_{stp} = \{\tau_d, \tau_f, U, f, \tau_s\}$ (see Methods). Here we
estimate values for depression, facilitation, and membrane time-constants along with release
probability, U, and magnitude of facilitation, f, (Fig. 4B). Since these values are estimated from
spikes and in observational settings rather than controlled experiments, the parameter estimates
are likely to be biased by omitted variables (Stevenson, 2018). However, the parameter estimates
do provide accurate predictions of postsynaptic spiking during natural, ongoing pre- and post-
synaptic spiking, and may provide an initial, approximate description of synaptic dynamics.
Comparing the estimates for the three model synapses – the thalamus pair has the highest release
probability (0.29$\pm$0.04 SE) and the largest membrane (14$\pm$2 ms) and depression time-constants
(410$\pm$107 ms). The VB-Barrel connection has a small membrane time-constant (0.3$\pm$0.003 ms)
and a larger depression (182$\pm$8 ms) time-constant than facilitation time-constant (105$\pm$9 ms). The
ANF-SBC synapse has the lowest release probability of the three connections (0.068$\pm$0.006) and
small depression (67$\pm$6 ms) and membrane time-constant (0.25$\pm$0.02 ms). Due to the potential
for omitted variable bias and differences in experimental preparations comparing these values

directly to measurements from intracellular recordings is difficult. However, the values estimated
from ongoing spiking and the results from intracellular recordings are generally in agreement. For
instance, previous *in vitro* studies of thalamocortical projections found that paired-pulse ratios
ranged from 0.3-0.9 consistent with depressing VB-Barrel synapses (Gil et al., 1997).
Additionally, *in vitro* observations of ANF-SBC connections report depression time-constants on
the order of 2-25 ms in response to a 100 Hz stimulus train (Wang and Manis, 2005, 2008). These
previous estimates are substantially faster than the time-constants estimated by the TM-GLM for
the ANF-SBC connection here. However, different patterns of presynaptic input (e.g. regular,
Poisson, natural) or differences in calcium concentration and temperature may make it difficult to
compare *in vitro* and *in vivo* STP parameters directly. One parameter that may be more readily
comparable across preparations is the membrane time-constant. We find that the estimated
membrane time-constant from the TM-GLM for the thalamus pair is consistent with thalamus relay
cells observed intracellularly (12.2 ± 1.1 ms, n=8) (Paz et al., 2007), and the estimated membrane
time-constant for ANF-SBC is close to *in vitro* measurements (1.05 ± 0.09 ms) as well (Wang and
Manis, 2005).

The TM-model used here is one of many possible parametric descriptions of short-term plasticity
(Hennig, 2013). Previous work modeling intracellular recordings suggests that the full TM model
may not be necessary to explain STP at some, purely depressing synapses (Costa et al., 2013).
Therefore, we explored how simplified TM models of STP, with fewer parameters, compare with
the full model using the Akaike information criterion (AIC; see Methods and Fig 4C). AIC
evaluates model accuracy (log-likelihood) penalized by the number of parameters, and lower AIC
may indicate that a simplified model with fewer parameters is preferred over a more complex
model. Generally, the synaptic dynamics in this class of models can be described by four
parameters: a time-constant for depression τ_d, a time-constant for facilitation τ_f, a baseline release
probability U, and facilitation parameter f. When modeling spike transmission we additionally
include a parameter for the membrane time-constant τ_s and consider the possibility that the
membrane potential "resets" following a post-synaptic spike (see Methods). For each of these
models, it is important to note there may be many possible parameter settings that are consistent
with the data, particularly when the recording time is limited (Costa et al., 2013). These
redundancies are present even in simple quantal analysis methods (Bykowska et al., 2019). Here,
altogether, we compare our full model to five reduced models: 1) a model with only membrane
integration, without dynamic release probability and resources, 2) a facilitation only model, 3) a
depression only model, 4) a 3-parameter TM model where the magnitude of facilitation is fixed,
and 5) the full TM model, but without post-spike reset of integration (Table 1). The full TM model
performs competitively in all cases, but, for some synapses, just as with previous results modeling
PSPs (Costa et al., 2013), the full model may be overly flexible and simpler models, with fewer
parameters, may be preferred. For the thalamus pair and VB-Barrel projection, the 3-parameter
TM-model with fixed magnitude of facilitation has the lowest AIC ($p<10^{-9}$ and p=0.07 compared

to model 6 with a paired t-test). For the ANF-SBC connection the full model gives the lowest AIC
($p<10^{-6}$ compared to model 4). For all three connections, models 4-6 perform statistically
significantly better than both the model without STP (e.g. $\Delta AIC<0$, Bonferroni-corrected paired t-
test $p<0.001$) and model 1 (Bonferroni-corrected paired t-test, $p<0.001$). These results provide
further evidence for STP-like changes in spike transmission at these connections.

Fig. 4 Including short-term dynamics substantially improves the model of spike transmission. A:
Spike transmission patterns are diverse across different connections. For three different connections
(between a pair of neurons in thalamus, a projection from ventrobasal thalamus to somatosensory cortex,
and an auditory nerve fiber projection onto a spherical bushy cell) transmission patterns are modeled by a
combination of different factors. For each synapse, top panels show the presynaptic ISI distributions (log-
spaced). In the second/third row, the observed spike transmission probability (red data points) and model
predictions (blue with 95% confidence bands) for training and test set (2-fold cross-validation). We then
used the estimated TM parameters for each synapse and simulated responses to paired presynaptic pulses.
Blue curves denote the PPRs of the full model, and gray lines denote PPRs by taking synaptic summation
out. (bottom row) TM-GLM (blue) are superior in predicting individual postsynaptic transmission events
compared to GLM (orange, without STP) for each synapse type. For each individual presynaptic spike, we
compare the model transmission probability with the observed binary outcome. ROC curves show the
prediction accuracy with positive deviations from the diagonal indicating better performance. **B:** Estimates
for the four STP parameters of the model for each synapse. Dots represent estimates from bootstrap sampled
data. **C:** Model comparison for 6 different models (Akaike information criteria, AIC, relative to a model

without plasticity). Models: 1) Integration only, 2) Facilitation only, 3) Depression only, 4) 3-parameter
TM, 5) 4-parameter TM without resetting integration, and 6) Full model. Boxplots denote the difference in
AIC values for bootstrap samples in (B).

Model	Description	τ_d	τ_f	f	U	Reset
1	Integration only	0	0	1	1	Yes
2	Facilitation only	0	No constraints			Yes
3	Depression only	No constraints	0	No constraints		Yes
4	3-parameter TM	No constraints		$f = U$		Yes
5	TM without reset	No constraints				No
6	Full model	No constraints				Yes

624 **Table 1: Parameters included each model**. Note that τ_s is not constrained in any of the 6 models.
625

Recent patterns of pre- and postsynaptic spiking shape the synaptic transmission probability

Although previous studies have focused largely on how spike transmission probability varies as a
function of the single ISI preceding the most recent presynaptic, synaptic dynamics depend on the
full sequence of presynaptic spiking. Unlike *in vitro* experiments where the state of the synapse
can, to some extent, be controlled before studying responses to a specific presynaptic pattern, *in*
vivo measurements of spike transmission can be heavily influenced by higher-order correlations
between successive ISIs (Stoelzel et al., 2008). Additionally, it is difficult to assess the effects of
multi-spike patterns empirically by splitting the correlograms, since the number of observations
for any given presynaptic spike pattern rapidly decreases with the number of spikes in the pattern.
Here we examine how spike transmission depends, not just on the preceding presynaptic ISI, but
on triplets of spikes. We compare the empirically observed spike transmission probability
following triplets to the estimated spike transmission probability from the TM-GLM. Using the
model fits for TM-GLM, we then simulate postsynaptic responses to isolated patterns of spikes
and determine to what extent the observed spike transmission patterns are influenced by higher-
order correlations between successive ISIs.

First, in addition to the timing of the two preceding presynaptic spikes (separated by the interval
ISI_1), we split correlograms based on the timing of the three preceding presynaptic spikes (Fig.
5A), separated by the most recent interval and the one before (ISI_2). Since the TM-GLM provides
estimates of the post-synaptic spike probability following every presynaptic spike, we can split

both the data and model fits the same way (Fig. 5C). We find that the spike transmission patterns clearly depend on the triplet patterns of presynaptic spikes in ongoing spiking activity. That is, the spike transmission probability is influenced by both ISI_1 and ISI_2, and the interaction between the two ISIs differs between synapses. However, as with spike transmission as a function of ISI_1 alone, the TM-GLM accurately captures the patterns of spike transmission for triplets of presynaptic spikes for the three synapses. In the thalamus pair, spike transmission probability is most influenced by ISI_1, and the effect of ISI_2 appears to be weak or, at least, does not appear to be monotonic. Spike transmission probability at the VB-Barrel connection depends on both ISI_1 and ISI_2, with higher spike transmission probability for longer ISI_2, consistent with recovery from depression. Lastly, for the ANF-SBC connection, transmission probabilities decrease for shorter ISI_2, but there also appears to be a strong interaction between ISI_1 and ISI_2, where transmission probability is high for multiple combinations of these two intervals (e.g. intervals of 10 ms then 100 ms and intervals of 100 ms then 10 ms both result in high probability transmission).

Although these empirical results suggest that spike transmission probability is influenced by triplet patterns of presynaptic spikes, these triplets are not isolated events but are embedded in longer sequences of spikes with higher-order correlations between successive ISIs. To examine to what extent the model predictions are affected by higher-order correlations between successive ISIs, we again use the estimated parameters in the TM-GLM to simulate postsynaptic responses to hypothetical, isolated triplets of presynaptic spikes (Fig. 5C, bottom). In these simulations we fix the post-spike history effect and the excitability in the model to their average values from model fits, and we fix the initial STP state (initial values of R and u in TM model) for the first spike in triplets to the average R and u values from the model fits. Although the initial states of the pre- and postsynaptic neurons in the experimental data are not matched for different values of ISI_1 and ISI_2, by simulating, we can assess the isolated influence of different triplets (ISI_1 and ISI_2) on the model. Here we find that for the thalamus pair, although the empirical data showed no clear effect for ISI_2, the simulated spike transmission probability increases with short ISI_2, consistent with strong synaptic summation. One reason that this effect may be masked in the empirical transmission probabilities is that post-spike history effects could act to decrease the probability of future postsynaptic spikes. For the VB-Barrel simulations, we find that short ISI_2 decreases transmission probability, consistent with the empirical transmission patterns, although less pronounced. Serial correlations in the sequence of presynaptic spikes (such as long bursts) could act to accentuate the depression in the empirical observations beyond what we see with the simulated responses to isolated triplets. Finally, for the ANF-SBC, although the empirical transmission probability showed decreased transmission for short ISI_2, the simulated responses to isolated patterns have increasing transmission at short ISI_2 (due to synaptic summation). This difference is likely due to the post-spike history filter, which has been fixed for the simulations, but can have a large effect in the experimental data. Since the overall efficacy of this synapse is

quite high (>0.7), is likely that a postsynaptic spike follows the first or second presynaptic spike
which then influences the response to the third spike.

To better understand the effects of post-spike history, we examined how the postsynaptic spiking
history changes the spike transmission patterns with a similar approach. In addition to splitting the
correlograms based on ISI_1, we also split based on the previous postsynaptic ISI, ISI_{post} (Fig. 5D).
Here, as with the triplets of presynaptic spikes, we find that the spike transmission patterns depend
on the triplet patterns of 2 pre- and 1 postsynaptic spike in data and that the TM-GLM accurately
captures the patterns of spike transmission at our three synapses (Fig. 5F). Here, for both thalamus
and VB-Barrel pairs, synaptic transmission probability decreases after a long postsynaptic ISI for
all values of ISI_1. In contrast, the ANF-SBC connection shows decreased transmission probability
at short postsynaptic ISIs.

As with the triplets of presynaptic spikes, we then simulate (Fig. 5F, bottom) how patterns of 2
pre- and 1 postsynaptic spike change spike transmission probability when the neurons start from
the same initial conditions (average values of excitability, post-spike history, R and u). For the
thalamus and VB-Barrel pairs, the simulations of isolated patterns match the general trends of
empirical spike transmission. However, for the VB-Barrel synapse, the effect of ISI_{post} in the
empirical transmission patterns is stronger than in the simulations, suggesting that serial
correlations in ISIs could again play a role and amplify the effects of isolated patterns.

Fig. 5: Pre- and postsynaptic spiking history determine transmission probability. A) Schematic of 4 different patterns of presynaptic spike triplets with a fixed interval between the two most recent presynaptic spikes (spikes denoted by black lines separated by ISI_1). B) We then split the presynaptic ISI distribution into 8 quantiles, denoted by the different colors. C) We then assess how ISI_2 influences the spike transmission previously described for ISI_1. Using the natural occurrence of different ISI_1 and ISI_2 in the data, each data point shows the observed spike transmission probability for each pattern (colors correspond to ISI_2 quantiles). Lines denote the average estimated transmission probability for each pattern under the model (based on the natural sequence of observed spikes). To examine the influence of serial correlations, we then simulate model responses to the isolated triplet pattern, assuming the synapse is initially in an average state (bottom panels). D) Synaptic transmission patterns change depending on the history of postsynaptic spiking, as well. E) Note that the postsynaptic ISI distributions need not match the presynaptic distributions. F) Here each data point in the scatter plots shows the spike transmission probability following different combinations of ISI_1 and ISI_{post}. Here, colors denote quantiles of the postsynaptic ISI distribution. Solid lines show the estimated transmission probability for each pattern under the model (based on the natural sequence of observed spikes). The bottom panels show model responses to isolated patterns using the estimated STP parameters and fixing the excitability from the model fits to their average values.

Spike transmission patterns change depending on stimulus type

The results above suggest that the presynaptic spike pattern has a complex effect on spike transmission probability. In sensory systems, one factor that affects the presynaptic spike pattern is the external stimulus. To examine how differences in stimulus statistics might alter spike

transmission, we fitted our model to a dataset recorded juxtacellularly from an ANF-SBC synapse,
presented with *natural sounds*, a range of randomized frequency-level pure-tones (*tuning stimuli*),
and s*pontaneous activity* in the absence of acoustic stimulation. Note that this dataset was partially
(tuning stimuli) used in the first section of the results. We merged these three datasets and fitted
the model to the merged dataset. As with the previous fits of the ANF-SBC connection (based on
a different set of *tuning stimuli*), the transmission probability under all three conditions exhibits a
bandpass-like pattern in mid-range ISIs suggesting facilitation and little to no synaptic summation.
However, spike transmission during natural stimuli was markedly different from that during pure
tone stimulation. During natural sounds, transmission probability is maximized at 100 ms rather
than 10 ms found in the *tuning stimuli* and during *spontaneous activity*. Further, *natural stimuli*
have much lower transmission probability at short ISIs. Interestingly, the TM-GLM captures the
overall facilitation, but also captures differences due to the different stimuli (Fig 6A). In contrast,
a static GLM captures almost none of the variations in spike transmission probability. Together,
these results suggest that the combination of STP, synaptic summation, history, and excitability is
sufficient to explain the observed differences spike transmission between stimuli, without
requiring any additional adaptation or plasticity.

Since these recordings were performed juxtacellularly, we also have access to the slope of
individual (extracellularly observed) PSPs, which are correlated with the intracellular PSP
amplitudes. We compared patterns of individual PSP slopes for each stimulus type and examine
how these slopes correlate with the estimated coupling amplitude following individual presynaptic
spikes in our model (Fig. 6B, 6C). Note that patterns of PSP slopes do not have the same pattern
as spike transmission probability, since there are other factors (e.g. postsynaptic spiking history)
contributing to postsynaptic spiking. However, as with spike transmission, we find that the PSP
amplitudes are stimulus-dependent and that a static GLM without STP cannot account for these
variations. Additionally, although the correlation is not perfect, the individual coupling effects in
the model do correlate with the measured PSP slope, even though the model is only fit to spikes.
By modeling dynamic functional connectivity, we can approximately reconstruct the amplitude of
individual synaptic events.

We then analyze how much the TM-GLM can generalize to other stimulus types when fit to one
stimulus type. We find that, although the model can describe the spike transmission patterns for
all three stimuli when fit to all stimuli, the model does not generalize to natural stimuli when fit
exclusively to one of the other stimulus types (and vice versa, Fig. 6D). The parameters from each
of these models are distinct – occupying different regions of the parameter space. Notably, the
model fit to all stimuli has a lower release probability and a higher facilitation time-constant
compared to the models fit to individual stimuli (Fig. 6E).

Fig. 6: The TM-GLM captures stimulus-dependent changes in spike transmission probability at the ANF-SBC synapse. A) The TM-GLM captures stimulus-dependent spike transmission probability patterns better than a static model without short-term synaptic plasticity. Dots show spike transmission probability for (log-spaced) presynaptic ISIs during two types of auditory stimuli

and during spontaneous activity: Natural Sounds (yellow), Spontaneous Activity (red), and Tuning Stimuli
(blue). Solid lines and 95% confidence bands show model predictions for each stimulus type.
Corresponding inter-spike interval distributions are shown on the right. **B)** The TM-GLM captures changes
in extracellularly recorded PSPs. Here the observed PSP slope (dots) approximately matches the coupling
term in the TM-GLM (solid lines) for each three stimuli. Although the spike transmission probability of the
static GLM can vary as a function of presynaptic ISI due to non-synaptic factors, the coupling term is fixed.
C) Estimates of individual PSP amplitudes predicted by the model and their PSP slopes in the juxtacellular
recording. Black lines denote linear fits and the bar plot shows the corresponding Spearman correlations.
D) After fitting each stimuli condition separately, in each column we plotted the estimated spike
transmission probability of each type using the estimated STP parameters of others. **E)** Distribution of
parameters from bootstrap samples with the TM-GLM fit for individual stimuli and all stimuli combined.

Postsynaptic cell-type specific changes in spike transmission patterns

We also applied our model to spiking data from a large-scale multi-electrode array recording to
investigate the spike transmission dynamics in synapses from putative excitatory neurons to two
different putative inhibitory subtypes. We detected putative synapses using the log-likelihood ratio
(LLR < -6, ~200 synapses) between a full model of the correlogram that includes the synaptic
effect and smooth model of the correlogram that only captures the slow structure (see Methods).
We then found excitatory-inhibitory microcircuits where putative excitatory neurons (based on the
cross-correlogram and spike waveform) give inputs to putative inhibitory neurons (41 excitatory
synapses onto 9 inhibitory neurons in total). To identify inhibitory neurons as inhibitory, we
required the neuron to have an outgoing connection to a third neuron with a fast, transient decrease
in the cross-correlogram. Each of the 9 putative inhibitory neurons here had at least one outgoing
connection where the spiking probability of a downstream neuron decreases >18% relative to
baseline following its spiking (Fig. 7A). We then categorized each neuron as a putative fast-spiking
(FS, n=5) or regular-spiking (RS, n=4) unit based on the spike waveform and firing rate (Fig. 7B).
Putative FS units had narrow-width spike waveforms (half-width of the trough = 0.08 ± 0.02 ms)
and higher firing rates (26.07 ± 9.6 Hz) compared to putative RS neurons (n=4) with broader
waveforms (half-width = 0.14 ± 0.02 ms) and lower firing rate (10.18 ± 10.01 Hz).

We identified these microcircuits in different regions with 4 putative excitatory-inhibitory
microcircuits recorded in hippocampus (depth differences: 77.2 ± 49.4 μm), 3 in thalamus
(49.4 ± 26.2 μm), and 2 in motor cortex (36.4 ± 23.5 μm). Putative excitatory neurons showed a
wide spike waveform (half-width = 0.18 ± 0.04 ms) similar to the putative regular-spiking
inhibitory neurons, but these two classes can be distinguished by their outgoing connection types
(e.g. inhibitory/excitatory) (Moore and Wehr, 2013) (Fig. 7B). Average efficacies from putative
excitatory-FS connections (0.22 ± 0.12, n=22) were larger, on average, compared to putative

excitatory-RS efficacies (0.13±0.13, n=19). We then fit the TM-GLM to data from these 41
putative synapses, similar to the three identified synapses analyzed above. Again, due to omitted
variable bias, the interpretation of the parameter values for the model fits is not necessarily straight-
forward. However, we find that there is substantial overlap between the estimated STP parameters
for excitatory connections onto these two inhibitory subtypes (Fig. 7C). The depression time-
constant for excitatory-RS connections is 215±219 ms (mean±SD, median 96 ms) and for
excitatory-FS is 411±459 ms (median 191 ms). The facilitation time-constant for excitatory-RS
connections is 820±745 ms (median 588 ms) and 406±552 ms (median 236 ms) for excitatory-
FS connections. And the membrane time-constant for excitatory-RS connection is 84±116 ms
compared to 72±196 ms for excitatory-FS. Interestingly, the estimates for membrane time-
constant (median 10 ms for FS, 45 ms for RS) are similar to the parameters measured using
intracellular recordings *in vitro* (Perrenoud et al., 2013).

Previous *in vitro* studies of postsynaptic cell-type specific STP concluded that putative excitatory-
RS connections show facilitation and putative excitatory-FS connections show depression
(Thomson and Lamy, 2007). Moreover, few *in vivo* studies characterized stimulated activities in
these connections (Pala and Petersen, 2015, 2018; Sedigh-Sarvestani and Vigeland, 2017). A cell-
type-specific study of somatosensory connections *in vivo* using 50Hz optogenetic stimulation
found little short-term plasticity in connections to Parvalbumin-expressing neurons (putative
excitatory-FS here), while excitatory to Somatostatin-expressing neurons (putative excitatory-RS
here) showed facilitation (Pala and Petersen, 2015). However, we are not aware of any *in vivo*
experiments that measured depression or facilitation time-constants for these systems during
ongoing spiking activity. Here we find that both connection types are somewhat facilitating but
excitatory-FS connections having a slightly shorter facilitation time-constant. However, unlike
what would be expected if excitatory-FS connections were depressing, the release probability of
excitatory-FS connections is lower than excitatory-RS connections (Fig. 7C, 0.34±0.19 for FS,
0.46±0.17 for RS). To better understand synaptic transmission *in vivo* it is important to consider
not just the parameters of the synapse but the full history of presynaptic spiking in the individual
presynaptic neurons. We use the estimated model parameters to simulate responses to a train of
regular presynaptic spikes with the frequency matched to the average firing rate of the
corresponding excitatory input. In simulating postsynaptic responses to the spike train, we fix the
excitability and postsynaptic history to their average values from model fits and set the initial STP
state of the first spike in the train to the average R and u values from model fits. With these input-
matched simulations, excitatory-RS connections show higher amplitude postsynaptic potentials
compared to excitatory-FS connections (Fig. 7D, the effect of membrane potential integration is
included). This is in accordance with the previously observed small degree of facilitation in
connections to Somatostatin-expressing neurons and small degree of short-term plasticity in
connections to Parvalbumin cells in (Pala and Petersen, 2015).

We also calculated spike transmission probabilities for all connections. On average, connections
to regular-spiking inhibitory neurons show a higher spike transmission probability across
interspike intervals (Fig. 7E). For all connections, we then evaluated the spike prediction accuracy
of a model without STP (e.g. static GLM) with our TM-GLM using the Area Under the ROC
Curve (Fig. 7F). The model with STP (TM-GLM) gives more accurate predictions for which
presynaptic spikes will lead to postsynaptic spiking for our population of 41 putative excitatory-
inhibitory connections (AUC=.69$\pm$.05) in comparison with the static GLM (AUC=.50$\pm$.03).
Altogether, these results illustrate how a dynamic model of functional connectivity, such as the
TM-GLM, can provide a detailed functional description of the short-term dynamics of spike
transmission in awake, behaving animals.

Fig. 7: Distinctive short-term dynamics for spike transmission in connections between excitatory
neurons to putative Regular-Spiking (RS) and Fast-Spiking (FS) inhibitory neurons. A) Here we
examine putative synapses between excitatory neurons and inhibitory neurons (identified by their cross-
correlations) and separate the putative inhibitory neurons into two classes: fast-spiking, which have narrow
spike waveforms and high rates (left), and regular-spiking (right), which have wide waveforms and lower
rates. Identifying these synapses requires both finding both a putative excitatory input and a putative
inhibitory output for the same neuron. B) Half-widths (of the trough) of the spike waveforms and firing
rates for the FS (orange) and RS (blue) inhibitory neurons, as well as, their excitatory inputs (grey).
Individual blue and orange waveforms (maximum amplitude across the MEA) are shown for all 9 putative
inhibitory neurons. C) Estimated depression, facilitation, and membrane time-constants for excitatory-RS
and excitatory-FS connections, along with the release probability (right). The purple error-bar next to the
membrane time-constant estimations show the median and standard deviations from *in vitro* experiments
(Perrenoud et al., 2013). D) Simulated postsynaptic potential amplitudes estimated from Tsodyks-Markram
model of short-term synaptic plasticity using estimated parameters. For each synapse, PSPs are estimated
in response to a pulse train with inter-pulse intervals set to their corresponding average presynaptic inter-
spike intervals. Dots and error bars denote the median and inter-quartile range for excitatory-RS (blue) and
excitatory-FS (red) connections. These responses include the effect of membrane potential integration. E)
Spike transmission probability patterns for individual synapses of excitatory-RS (blue) and excitatory-FS
(red) connections normalized by long interval probabilities as a function of the presynaptic ISI. F) Area
Under the Curve (AUC) of postsynaptic spiking prediction using the static GLM without short-term
synaptic plasticity (green) and the TM-GLM with short-term synaptic plasticity (blue). G-H) Spike-
transmission probabilities (left) and corresponding cross-correlograms (right) of 4 putative excitatory inputs
to putative FS (G) and RS (H) inhibitory neurons show cell-type specific similarities.

Synapse	n	τ_s (ms)	τ_d (ms)	τ_f (ms)	U
Thalamus	1	14 ± 2	410 ± 107	37 ± 12	0.29 ± 0.04
VB-Barrel	1	0.3 ± 0.003	182 ± 8	105 ± 9	0.10 ± 0.05
ANF-SBC	1	0.25 ± 0.02	67 ± 6	71 ± 3	0.068 ± 0.006
excitatory-RS	19	84 ± 116	215 ± 219	820 ± 745	0.46 ± 0.17
excitatory-FS	22	72 ± 196	411 ± 459	406 ± 552	0.34 ± 0.19

Table 2: Summary of parameter estimates from the full TM-GLM. Sample size (n), membrane
time-constant (τ_s), depression time-constant (τ_d), and facilitation time-constant (τ_f), and release
probabilities (U) for the identified and putative synapses from our three case studies and multi-
electrode recordings. For the cases studies, the mean±standard deviation is shown for the bootstrap
samples. For the MEA data, the mean±sd is shown across putative connections. In all cases, the
parameters are estimated from ongoing, in vivo spiking activity.
877

Spike "transmission" patterns between unconnected pairs of neurons

It is important to note that the dynamic functional connectivity model presented here assumes that, before fitting the model, we have accurately identified a monosynaptic connection. In some settings, it is possible to identify connections using optogenetic stimulation (English et al., 2017) or juxtacellular recording, however, in cases where we can only identify putative connections, it is important to consider the possibility that we are modeling a spurious correlation between neurons that are not actually monosynaptically connected. In general, the detection of monosynaptic connections from multielectrode spiking activity is far from perfect (Kobayashi et al., 2019).

To examine how the TM-GLM might be influenced by spurious correlations, we first simulated a small circuit with common drive that would likely lead to a falsely detected monosynaptic connection (Fig 8A). Here an unobserved presynaptic (inhomogeneous Poisson process) neuron provides strong excitatory input to two leaky integrate-and-fire postsynaptic neurons. Due to a difference in the latencies of these connections, there is a spurious peak in the correlogram between the two postsynaptic neurons where one postsynaptic neuron appears to excite the other. We find that when we measure the amplitude of this spurious peak, there are some variations as a function of the presumed presynaptic neuron's ISI, and the spike "transmission" pattern varies depending on whether the projections from the true presynaptic are both depressing, both facilitating, or a mixture of depressing and facilitating (Fig 8B). However, the TM-GLM is nearly constant (~0.1% variation) and does not accurately fit the observed variation. Despite a spurious correlation, the detailed pattern of spikes between the two postsynaptic neurons is unstructured and not well described by the TM model.

We also fit the TM-GLM to several (n=38) pairs of neurons from the MEA data all with average firing rates in range of 3-15 Hz and where there was no clear peak in the cross-correlogram (0-5ms following the spikes of one neurons). In these cases, although the coupling filter is likely fitting noise and does not describe a realistic synaptic effect (median latency 0.7 ms, median time-constant 0.02 ms), the TM-GLM does describe small variations in the ISI-dependent pattern of spike "transmission" probability (Fig 8C). These patterns are not as pronounced as the patterns observed in the identified and putative monosynaptic connections described above, but they also appear to have structure that the TM-GLM can account for. Altogether, these results illustrate how the TM-GLM simply aims to account for short-term dynamics in the spiking probability of one neuron in reference to the spikes of another neuron. Correctly identifying monosynaptic connections is a necessary first step before the short-term dynamics can be meaningfully interpreted.

Fig. 8: Short-term changes in spike probability for neurons that are not monosynaptically connected. A) In a simulated circuit, we generated a spurious connection between two neurons (Post$_1$ and Post$_2$) receiving common excitatory drive from a single presynaptic neuron (Pre) with different delays (orange cross-correlogram) B) Scatter plots show normalized spike "transmission" probabilities from different sets of simulations where the true connections to the postsynaptic neurons have different types of short-term synaptic plasticity (both Depressing, both Facilitating, and one Depressing; one Facilitating). Lines with same colors as scatter plots show the estimated spike probability from the TM-GLM. Here the data and model fits are averaged across 150 rounds of simulations (50 for each combination) and are normalized in order to have a spiking probability of 1 for the longest ISIs C) We then fit spike "transmission" probability for 38 pairs of neurons from the multi-electrode array (MEA) recording where there was no clear monosynaptic connection (putative non-connections). Observed (left) spike transmission probabilities show relatively little variation as a function of one neuron's ISIs, but the TM-GLM (right) does describe

927 what variation there is. Insets show example cross-correlograms from two of these putative non-
928 connections.

Discussion

Here we developed a dynamic model of functional connectivity, the TM-GLM, and applied this
model to disentangle synaptic and nonsynaptic contributions to excitatory spike transmission in
vivo. Short-term synaptic plasticity (STP) has been extensively studied with intracellular
recordings where the amplitudes of individual postsynaptic potential/currents (PSP/PSCs) can be
directly measured. However, the relationship between STP and *in vivo* spike transmission patterns
is complex. Patterns of postsynaptic spike transmission are highly diverse and multiple factors
beyond STP shape these patterns (Swadlow and Gusev, 2001; English et al., 2017). Here, using a
model-based approach, we characterized these diverse spike transmission patterns at identified and
putative excitatory synapses and attribute this diversity to different combinations of short-term
synaptic plasticity, synaptic summation, and post-spike history effects. We then showed how this
modeling framework has the potential to capture stimulus-specific and cell-type-specific changes
in spike transmission *in vivo*.

Estimating static functional connectivity using spike times has revealed network structure in the
retina (Pillow et al., 2008) and hippocampus (Harris et al., 2003), can reconstruct true
physiological circuitry (Gerhard et al., 2013), and improves encoding and decoding (Truccolo et
al., 2005; Pillow et al., 2008; Stevenson et al., 2012). However, synaptic weights can change
dramatically over time and can also depend on external stimuli and behavior (Fujisawa et al.,
2008). Although, standard GLMs can partially capture the first-order effects of recent presynaptic
spikes on postsynaptic spiking probability, they fail to capture the nonlinear dynamics of synaptic
transmission affected by longer sequences of presynaptic spikes. With a static coupling term the
GLM can account for the average change in the postsynaptic spiking probability following a
presynaptic spike, but it does not make detailed predictions about the variations in this probability.
Here we show that, by including a dynamical model of short-term plasticity, we can capture diverse
pattern of spike transmission probability and substantially improve prediction of postsynaptic
spiking. In a recording from the endbulb of Held (ANF-SBC) we further found that spike
transmission patterns differed between stimuli, and that these differences were well-described by
a single TM-GLM. Although the STP-parameters were the same for all stimuli, the different
presynaptic spike patterns yield different patterns of spike transmission. Since spike transmission
probability in the TM-GLM depends on the full history of presynaptic spiking, this model can
account for changes on behavioral timescales even in the absence of adaptation or other forms of
plasticity (e.g. STDP, LTP). Using the models for the short-term dynamics of spike transmission

estimated in one setting we may also be able to more accurately predict responses to novel
presynaptic patterns and, in sensory systems, novel stimuli.

Previous *in vitro* studies have shown that STP dynamics depend on both presynaptic and
postsynaptic cell-types (Thomson and Lamy, 2007). Using a large multi-electrode recording from
a freely behaving mouse, we investigated the dynamics of synaptic connections from putative
excitatory neurons to two different subtypes of putative inhibitory neurons: putative fast-spiking
(FS) and putative regular-spiking (RS). Using only spike times, we find that spike transmission
shows slightly higher facilitation for excitatory-RS compared to the excitatory-FS connections.
Although drawing strong conclusions about the parameters of the model is difficult due to potential
confounds, the STP dynamics reflect this same pattern and are in line with previous *in vitro*
findings (Thomson and Lamy, 2007). Including short-term dynamics into the model also
significantly improves the prediction of postsynaptic spiking. As large-scale extracellular
recording techniques advance, models such as the TM-GLM may allow us to characterize and
compare the short-term dynamics of spike transmission of many different cell types, brain regions,
and species.

Several details of the model may impact our results. Here we employed an extended GLM with a
logistic spike nonlinearity, since it appears to better describe strong connections, such as the ANF-
SBC, better than the traditional exponential nonlinearity. However, other nonlinearities may be
better for other neurons (McFarland et al., 2013). There are also alternatives to the Tsodyks-
Markram model for modeling synaptic dynamics (Hennig, 2013). Although the TM model is
biologically plausible, it only tracks average, deterministic dynamics of postsynaptic potentials,
while ignoring the stochasticity of synaptic release (Barri et al., 2016; Bird et al., 2016). Finally,
there are many covariates that could be added to improve model performance, including local field
potentials (Kelly et al., 2010), connections to other simultaneously observed presynaptic neurons
(Harris et al., 2003), higher-order history or coupling terms (Robinson et al., 2016; Song et al.,
2018), and covariates related to other types of plasticity (Stevenson et al., 2011; Linderman et al.,
2014; Robinson et al., 2016; Amidi et al., 2018; Bayat Mokhtari et al., 2018). Despite these
simplifying assumptions and the fact that we only observe a fraction of inputs to the neuron, the
TM-GLM captures a wide diversity of *in vivo*, excitatory spike transmission patterns.

Although our model provides a tool to characterize the dynamics of spike transmission, there may
be fundamental limitations to how well true synaptic dynamics can be estimated from spike
observations. Firstly, functional connections inferred from spikes do not necessarily guarantee
anatomical connections. A peak in the cross-correlogram does not conclusively indicate the
presence of a monosynaptic connection (Moore et al., 1970). In most cases, we assume that the
transient, short-latency increase in postsynaptic spiking activity following a presynaptic spike
indicates the presence of an excitatory monosynaptic connection (Perkel et al., 1967).
Nevertheless, verifying connections using optogenetics (English et al., 2017), juxtacellular

recordings (Pinault, 2011), or imaging (Weiler et al., 2008) may provide more confidence in determining true monosynaptic connections. Secondly, we employ a spiking model that does not explicitly account for the detailed membrane potential of the postsynaptic neuron. Although there are links between the GLM and voltage-based models (Latimer et al., 2014, 2018), other approaches to modeling synaptic transmission with realistic spike-generation mechanisms, currents, and even dendritic morphology may more accurately reflect subthreshold dynamics (Ladenbauer et al., 2018). Thirdly, long-term changes in the synaptic weight may alter the short-term dynamics. Experiments show that short-term depression may be reduced after long-term depression and increased after long-term potentiation (Markram and Tsodyks, 1996; Sjöström et al., 2007; Costa et al., 2015, 2017). Accounting for these long-term changes in synaptic strength may allow for more accurately estimation of STP. Finally, there are many other factors that are likely to affect short-term spike transmission dynamics including, dendritic spikes (Bono and Clopath, 2017), receptors nonlinearities (Magee, 2000), such as those in NMDA receptors, changes in spike threshold due to sodium inactivation (Naud et al., 2011) or coupled to the subthreshold activity (Mensi et al., 2016), feed-forward inhibition (Pouille and Scanziani, 2001), feedback inhibition (Suzuki and Bekkers, 2012), or disinhibition (Letzkus et al., 2015). With intracellular observations these effects can generally be separated from the synaptic dynamics based on the timing of the signals. However, since these effects directly alter spike timing, they may act as confounders for models based on spike observations. Although they could potentially be incorporated in future models, omitting these effects from the model presented here may result in biased parameter estimates for both the synaptic and non-synaptic effects that are included (Stevenson, 2018).

Intracellular observations in controlled settings have found that short-term synaptic dynamics vary depending on the pre- and postsynaptic cell type (Thomson and Lamy, 2007; Lee et al., 2019) as well as brain region (Dittman et al., 2000; Wang et al., 2006), age (Reyes et al., 1998), and species (Testa-Silva et al., 2014). Additionally, short-term synaptic dynamics appear to vary with stimulus type and the larger computational function of the neural circuit (Karmarkar and Buonomano, 2007). To link synaptic dynamics to circuit-level neural computations we will need to study these dynamics during natural ongoing activity (Klyachko and Stevens, 2006) and ultimately during natural behavior. Since short-term synaptic plasticity affects not only the postsynaptic membrane potential but also the probability of postsynaptic spiking (Markram et al., 1998; Swadlow and Gusev, 2001; London et al., 2002; English et al., 2017), it may be possible to indirectly observe the effects of synaptic dynamics on spike transmission. Here we examined this possibility by including the effects of short-term synaptic plasticity in models of functional connectivity. Using this approach, we characterized diverse, stimulus-dependent, and cell-type-specific patterns of excitatory spike transmission using spike observations alone.

Data and software availability

1036 All data and software central to the conclusion of this study are available at
1037 https://github.com/abedghanbari2/TM-GLM.

Acknowledgements

1039 Thanks to Nick Steinmetz for providing MEA dataset. AG, NR, and IHS were supported by NSF
1040 CAREER 1651396 to IHS. BE was partially funded via an NWO Open Grant (ALWOP.346) and
1041 an NWO VIDI grant (016.Vidi.189.052).

References

Abbott LF, Regehr WG (2004) Synaptic computation. Nature.

Amidi Y, Nazari B, Sadri S, … UE-2018 40th A, 2018 U (2018) Parameter Estimation in Synaptic
Coupling Model Using a Point Process Modeling Framework*. ieeexplore.ieee.org.

Barri A, Wang Y, Hansel D, Mongillo G (2016) Quantifying Repetitive Transmission at Chemical
Synapses: A Generative-Model Approach. eNeuro 3:Suppl:1-40.

Barthó P, Hirase H, Monconduit L, Zugaro M, Harris KD, Buzsáki G (2004) Characterization of
neocortical principal cells and interneurons by network interactions and extracellular features.
J Neurophysiol 92:600–608.

Bayat Mokhtari E, Lawrence JJ, Stone EF (2018) Data Driven Models of Short-Term Synaptic
Plasticity. Front Comput Neurosci.

Bird AD, Wall MJ, Richardson MJE (2016) Bayesian Inference of Synaptic Quantal Parameters
from Correlated Vesicle Release. Front Comput Neurosci 10:116.

Bono J, Clopath C (2017) Modeling somatic and dendritic spike mediated plasticity at the single
neuron and network level. Nat Commun 8:706.

Boyd S, Vandenberghe L (2004) Convex optimization. Cambridge university press.

Bykowska O, Gontier C, Sax A-L, Jia DW, Montero ML, Bird AD, Houghton C, Pfister J-P, Costa
RP (2019) Model-Based Inference of Synaptic Transmission. Front Synaptic Neurosci 11:21
Available at: https://www.frontiersin.org/article/10.3389/fnsyn.2019.00021/full [Accessed
February 17, 2020].

Carandini M, Horton JC, Sincich LC (2007) Thalamic filtering of retinal spike trains by
postsynaptic summation. J Vis 7:20.

Costa RP, Froemke RC, Sj??str??m PJ, van Rossum MCW (2015) Unified pre- and postsynaptic
long-term plasticity enables reliable and flexible learning. Elife 4:1–16.

Costa RP, Mizusaki BEP, Sjöström PJ, van Rossum MCW (2017) Functional consequences of
pre- and postsynaptic expression of synaptic plasticity. Philos Trans R Soc B Biol Sci 372.

Costa RP, Sjöström PJ, van Rossum MCW (2013) Probabilistic inference of short-term synaptic
plasticity in neocortical microcircuits. Front Comput Neurosci 7:75 Available at:
http://journal.frontiersin.org/article/10.3389/fncom.2013.00075/abstract [Accessed February

1071  17, 2020].

1072 Csicsvari J, Hirase H, Czurko A, Buzsáki G (1998) Reliability and state dependence of pyramidal
1073   cell-interneuron synapses in the hippocampus: An ensemble approach in the behaving rat.
1074   Neuron 21:179–189.

1075 Dittman JS, Kreitzer AC, Regehr WG (2000) Interplay between facilitation, depression, and
1076   residual calcium at three presynaptic terminals. J Neurosci.

1077 English DF, McKenzie S, Evans T, Kim K, Yoon E, Buzsáki G (2017) Pyramidal Cell-Interneuron
1078   Circuit Architecture and Dynamics in Hippocampal Networks. Neuron 96:505–520.

1079 Fetz E, Toyama K, Smith W (1991) Synaptic Interactions between Cortical Neurons. In, pp 1–47.

1080 Fetz EE, Gustafsson B (1983) Relation between shapes of post-synaptic potentials and changes in
1081   firing probability of cat motoneurones. J Physiol 341:387–410.

1082 Fujisawa S, Amarasingham A, Harrison MT, Buzsáki G (2008) Behavior-dependent short-term
1083   assembly dynamics in the medial prefrontal cortex. Nat Neurosci.

1084 Gerhard F, Kispersky T, Gutierrez GJ, Marder E, Kramer M, Eden U (2013) Successful
1085   Reconstruction of a Physiological Circuit with Known Connectivity from Spiking Activity
1086   Alone. PLoS Comput Biol 9.

1087 Ghanbari A, Malyshev A, Volgushev M, Stevenson IH (2017) Estimating short-term synaptic
1088   plasticity from pre- and postsynaptic spiking. PLOS Comput Biol 13:e1005738 Available at:
1089   https://doi.org/10.1371/journal.pcbi.1005738.

1090 Gil Z, Connors BW, Amitai Y (1997) Differential regulation of neocortical synapses by
1091   neuromodulators and activity. Neuron.

1092 Harris K, Csicsvari J, Hirase H, Dragoi G, Buzsáki G (2003) Organization of cell assemblies in
1093   the hippocampus. Nature 424:552–556.

1094 Hatsopoulos NG, Xu Q, Amit Y (2007) Encoding of movement fragments in the motor cortex. J
1095   Neurosci 27:5105–5114.

1096 Hennig MH (2013) Theoretical models of synaptic short term plasticity. Front Comput Neurosci
1097   7:45.

1098 Henze DA, Buzsáki G (2001) Action potential threshold of hippocampal pyramidal cells in vivo
1099   is increased by recent spiking activity. Neuroscience 105:121–130.

1100 Herz AVM, Gollisch T, Machens CK, Jaeger D (2006) Modeling single-neuron dynamics and
1101   computations: A balance of detail and abstraction. Science (80-).

1102 Huang C, Resnik A, Celikel T, Englitz B (2016) Adaptive Spike Threshold Enables Robust and
1103   Temporally Precise Neuronal Encoding. PLoS Comput Biol.

1104 Jun JJ et al. (2017) Fully integrated silicon probes for high-density recording of neural activity.
1105   Nature 551:232–236.

1106 Karmarkar UR, Buonomano D V. (2007) Timing in the Absence of Clocks: Encoding Time in
1107   Neural Network States. Neuron.

Kawaguchi Y (2001) Distinct firing patterns of neuronal subtypes in cortical synchronized
activities. J Neurosci.

Keine C, Rubsamen R, Englitz B (2016) Inhibition in the auditory brainstem enhances signal
representation and regulates gain in complex acoustic environments. Elife 5.

Keine C, Rübsamen R, Englitz B (2017) Signal integration at spherical bushy cells enhances
representation of temporal structure but limits its range. Elife 6.

Kelly RC, Smith MA, Kass RE, Lee TS (2010) Local field potentials indicate network state and
account for neuronal response variability. J Comput Neurosci 29:567–579.

Klyachko VA, Stevens CF (2006) Excitatory and feed-forward inhibitory hippocampal synapses
work synergistically as an adaptive filter of natural spike trains. PLoS Biol 4:1187–1200.

Kobayashi R, Kurita S, Kurth A, Kitano K, Mizuseki K, Diesmann M, Richmond BJ, Shinomoto
S (2019) Reconstructing neuronal circuitry from parallel spike trains. Nat Commun.

Ladenbauer J, McKenzie S, English DF, Hagens O, Ostojic S (2018) Inferring and validating
mechanistic models of neural microcircuits based on spike-train data. bioRxiv Prepr.

Latimer KW, Chichilnisky EJ, Rieke F, Pillow JW (2014) Inferring synaptic conductances from
spike trains with a biophysically inspired point process model. Neural Inf Process Syst
27:954–962.

Latimer KW, Rieke F, Pillow JW (2018) Inferring synaptic inputs from spikes with a conductance-
based neural encoding model. bioRxiv:281089.

Lee JH, Campagnola L, Seeman SC, Jarsky T, Mihalas S (2019) Functional synapse types via
characterization of short-term synaptic plasticity. bioRxiv:648725.

Letzkus JJ, Wolff SBE, Lüthi A (2015) Disinhibition, a Circuit Mechanism for Associative
Learning and Memory. Neuron 88:264–276.

Linderman S, Stock C, Adams R (2014) A framework for studying synaptic plasticity with neural
spike train data. Adv Neural Inf.

London M, Schreibman A, Haä;usser M, Larkum ME, Segev I (2002) The information efficacy of
a synapse. Nat Neurosci 5:332–340.

Magee JC (2000) Dendritic integration of excitatory synaptic input. Nat Rev Neurosci.

Markram H, Tsodyks M (1996) Redistribution of synaptic efficacy between neocortical pyramidal
neurons. Nature 382:807–810.

Markram H, Wang Y, Tsodyks M (1998) Differential signaling via the same axon of neocortical
pyramidal neurons. Proc Natl Acad Sci U S A 95:5323–5328.

McFarland JM, Cui Y, Butts DA (2013) Inferring Nonlinear Neuronal Computation Based on
Physiologically Plausible Inputs. PLoS Comput Biol 9.

Mensi S, Hagens O, Gerstner W, Pozzorini C (2016) Enhanced Sensitivity to Rapid Input
Fluctuations by Nonlinear Threshold Dynamics in Neocortical Pyramidal Neurons. PLoS
Comput Biol.

Moore AK, Wehr M (2013) Parvalbumin-Expressing Inhibitory Interneurons in Auditory Cortex
Are Well-Tuned for Frequency. J Neurosci 33:13713–13723.

Moore GP, Segundo JP, Perkel DH, Levitan H (1970) Statistical Signs of Synaptic Interaction in
Neurons. Biophys J 10:876–900.

Mora Lopez C, Putzeys J, Raducanu BC, Ballini M, Wang S, Andrei A, Rochus V, Vandebriel R,
Severi S, Van Hoof C, Musa S, Van Helleputte N, Yazicioglu RF, Mitra S (2017) A Neural
Probe with Up to 966 Electrodes and Up to 384 Configurable Channels in 0.13 μm SOI
CMOS. IEEE Trans Biomed Circuits Syst 11:510–522.

Naud R, Petersen CCH, Pozzorini C, Mensi S, Avermann M, Gerstner W (2011) Parameter
extraction and classification of three cortical neuron types reveals two distinct adaptation
mechanisms. J Neurophysiol.

Pachitariu M, Steinmetz N, Kadir S, Carandini M, Harris KD (2016) Kilosort: realtime spike-
sorting for extracellular electrophysiology with hundreds of channels.

Pala A, Petersen CC (2018) State-dependent cell-type-specific membrane potential dynamics and
unitary synaptic inputs in awake mice. Elife.

Pala A, Petersen CCH (2015) In Vivo Measurement of Cell-Type-Specific Synaptic Connectivity
and Synaptic Transmission in Layer 2/3 Mouse Barrel Cortex. Neuron 85:68–75.

Paz JT, Chavez M, Saillet S, Deniau J-M, Charpier S (2007) Activity of Ventral Medial Thalamic
Neurons during Absence Seizures and Modulation of Cortical Paroxysms by the
Nigrothalamic Pathway. J Neurosci.

Perkel DH, Gerstein GL, Moore GP (1967) Neuronal spike trains and stochastic point processes.
II. Simultaneous spike trains. Biophys J 7:419–440.

Perrenoud Q, Rossier J, Geoffroy H, Vitalis T, Gallopin T (2013) Diversity of gabaergic
interneurons in layer VIa and VIb of mouse barrel cortex. Cereb Cortex.

Pillow JW, Shlens J, Paninski L, Sher A, Litke AM, Chichilnisky EJ, Simoncelli EP (2008) Spatio-
temporal correlations and visual signalling in a complete neuronal population. Nature
454:995–999 Available at: http://www.nature.com/doifinder/10.1038/nature07140.

Pinault D (2011) The juxtacellular recording-labeling technique. Neuromethods.

Poliakov A V, Powers RK, Sawczuk A, Binder MD (1996) Effects of background noise on the
response of rat and cat motoneurones to excitatory current transients. J Physiol 495:143–157.

Pouille F, Scanziani M (2001) Enforcement of temporal fidelity in pyramidal cells by somatic
feed-forward inhibition. Science (80-).

Regehr WG (2012) Short-term presynaptic plasticity. Cold Spring Harb Perspect Biol.

Reyes A, Lujan R, Rozov A, Burnashev N, Somogyi P, Sakmann B (1998) Target-cell-specific
facilitation and depression in neocortical circuits. Nat Neurosci.

Robinson BS, Berger TW, Song D (2016) Identification of stable spike-timing-dependent
plasticity from spiking activity with generalized multilinear modeling. Neural Comput.

Sedigh-Sarvestani M, Vigeland L (2017) Intracellular, In Vivo, Dynamics of Thalamocortical

Synapses in Visual Cortex. J.

Sjöström PJ, Turrigiano GG, Nelson SB (2007) Multiple forms of long-term plasticity at unitary neocortical layer 5 synapses. Neuropharmacology 52:176–184.

Song D, Robinson BS, Berger TW (2018) Identification of Short-Term and Long-Term Functional Synaptic Plasticity From Spiking Activities. In: Adaptive Learning Methods for Nonlinear System Modeling, pp 289–312. Elsevier.

Stevenson IH (2018) Omitted Variable Bias in GLMs of Neural Spiking Activity. Neural Comput.

Stevenson IH, Kording K, Koerding K (2011) Inferring spike-timing-dependent plasticity from spike train data. In: Advances in Neural Information Processing Systems (Shawe-Taylor J, Zemel RS, Bartlett P, Pereira FCN, Weinberger KQ, eds), pp 1–9.

Stevenson IH, London BM, Oby ER, Sachs NA, Reimer J, Englitz B, David S V., Shamma SA, Blanche TJ, Mizuseki K, Zandvakili A, Hatsopoulos NG, Miller LE, Kording KP (2012) Functional Connectivity and Tuning Curves in Populations of Simultaneously Recorded Neurons. PLoS Comput Biol 8:e1002775.

Stevenson IH, Rebesco JM, Miller LE, Körding KP (2008) Inferring functional connections between neurons. Curr Opin Neurobiol 18:582–588.

Stoelzel CR, Bereshpolova Y, Gusev AG, Swadlow HA (2008) The Impact of an LGNd Impulse on the Awake Visual Cortex: Synaptic Dynamics and the Sustained/Transient Distinction. J Neurosci.

Stoelzel CR, Bereshpolova Y, Swadlow HA (2009) Stability of Thalamocortical Synaptic Transmission across Awake Brain States. J Neurosci.

Suzuki N, Bekkers JM (2012) Microcircuits Mediating Feedforward and Feedback Synaptic Inhibition in the Piriform Cortex. J Neurosci 32:919–931.

Swadlow HA (2002) Thalamocortical control of feed-forward inhibition in awake somatosensory "barrel" cortex. Philos Trans R Soc Lond B Biol Sci 357:1717–1727.

Swadlow HA, Gusev AG (2001) The impact of "bursting" thalamic impulses at a neocortical synapse. Nat Neurosci 4:402–408 Available at: http://www.ncbi.nlm.nih.gov/pubmed/11276231.

Swadlow HA, Gusev AG (2002) Receptive-field construction in cortical inhibitory interneurons. Nat Neurosci.

Testa-Silva G, Verhoog MB, Linaro D, de Kock CPJ, Baayen JC, Meredith RM, De Zeeuw CI, Giugliano M, Mansvelder HD (2014) High Bandwidth Synaptic Communication and Frequency Tracking in Human Neocortex. PLoS Biol 12.

Thomson AM, Lamy C (2007) Functional maps of neocortical local circuitry. Front Neurosci 1:19–42.

Thomson AM, West DC, Wang Y, Bannister AP (2002) Synaptic Connections and Small Circuits Involving Excitatory and Inhibitory Neurons in Layers 2-5 of Adult Rat and Cat Neocortex: Triple Intracellular Recordings and Biocytin Labelling In Vitro. Cereb Cortex 12:936–953.

Truccolo W, Eden UT, Fellows MR, Donoghue JP, Brown EN, John P (2005) A Point Process
Framework for Relating Neural Spiking Activity to Spiking History, Neural Ensemble, and
Extrinsic Covariate Effects. J Neurophysiol 93:1074–1089.

Tsodyks M, Pawelzik K, Markram H (1998) Neural networks with dynamic synapses. Neural
Comput 1998 May;10(4)821-835 835:821–835.

Tsodyks M V., Markram H (1997) The neural code between neocortical pyramidal neurons
depends on neurotransmitter release probability. Proc Natl Acad Sci 94:719–723.

Usrey WM, Alonso JM, Reid RC (2000) Synaptic interactions between thalamic inputs to simple
cells in cat visual cortex. J Neurosci 20:5461–5467.

Wang Y, Manis PB (2005) Synaptic Transmission at the Cochlear Nucleus Endbulb Synapse
During Age-Related Hearing Loss in Mice. J Neurophysiol.

Wang Y, Manis PB (2008) Short-term synaptic depression and recovery at the mature mammalian
endbulb of Held synapse in mice. J Neurophysiol.

Wang Y, Markram H, Goodman PH, Berger TK, Ma J, Goldman-Rakic PS (2006) Heterogeneity
in the pyramidal network of the medial prefrontal cortex. Nat Neurosci 9:534–542.

Weiler N, Wood L, Yu J, Solla SA, Shepherd GMG (2008) Top-down laminar organization of the
excitatory network in motor cortex. Nat Neurosci.

Yang H, Xu-Friedman MA (2008) Relative Roles of Different Mechanisms of Depression at the
Mouse Endbulb of Held. J Neurophysiol.

Yang H, Xu-Friedman MA (2009) Impact of Synaptic Depression on Spike Timing at the Endbulb
of Held. J Neurophysiol.

Zucker RS, Regehr WG (2002) Short-Term Synaptic Plasticity. Annu Rev Physiol 64:355–405.

1243