Title: Ecological networks: Pursuing the shortest path, however narrow and crooked

Andrea Costa[1,2]*, Ana M. Martín González[3]*, Katell Guizien[4], Andrea M. Doglioli[5], José María Gómez[6],

Anne A. Petrenko[5], Stefano Allesina[7,8]

* shared first authorship and corresponding authors

Correspondence to: ana.maria.martingonzalez@gmail.com and andrea.costa@pusan.ac.kr

1- Center for Climate Physics, Institute for Basic Science, Busan, South Korea, 46241

2- Department of Atmospheric Sciences, Pusan National University, Busan, South Korea, 46241

3- Center for Macroecology, Evolution and Climate, Natural History Museum of Denmark, University of

Copenhagen, Copenhagen, Denmark

4- CNRS, Sorbonne Universités (UPMC Univ. Paris 06), Laboratoire d'Ecogéochimie des

Environnements Benthiques (LECOB), 1 avenue Pierre Fabre Observatoire Océanologique,

Banyuls/Mer,

France

5- Aix Marseille Université, CNRS, Université de Toulon, IRD, OSU Pythèas, Mediterranean Institute of

Oceanography (MIO), UM 110, 13288, Marseille, France

6- Departamento de Ecología Funcional y Evolutiva, Estación Experimental de Zonas Áridas (EEZA-

CSIC), Almería, Spain

7- Department of Ecology and Evolution, University of Chicago, Chicago, IL, USA

8- Northwestern Institute on Complex Systems, Northwestern University, Evanston, IL, USA

Representing data as networks cuts across all sub-disciplines in ecology and evolutionary biology. Besides providing a compact representation of the interconnections between agents, network analysis allows the identification of especially important nodes, according to various metrics that often rely on the calculation of the shortest paths connecting any two nodes. While the interpretation of a shortest paths is straightforward in binary, unweighted networks, whenever weights are reported, the calculation could yield unexpected results. We analyzed 129 studies of ecological networks published in the last decade and making use of shortest paths, and discovered a methodological inaccuracy related to the edge weights used to calculate shortest paths (and related centrality measures), particularly in interaction networks. Specifically, 49% of the studies do not report sufficient information on the calculation to allow their replication, and 61% of the studies on weighted networks may contain errors in how shortest paths are calculated. Using toy models and empirical ecological data, we show how to transform the data prior to calculation and illustrate the pitfalls that need to be avoided. We conclude by proposing a five-point check-list to foster best-practices in the calculation and reporting of centrality measures in ecology and evolution studies.

INTRODUCTION

The last two decades have witnessed an exponential increase in the use of graph analysis in ecological and conservation studies (see refs 1&2 for recent introductions to network theory in ecology and evolution). Networks (graphs) represent agents as nodes linked by edges representing pairwise relationships. For instance, a food web can be represented as a network of species (nodes) and their feeding relationships (edges)[3]. Similarly, the spatial dynamics of a metapopulation can be analyzed by connecting the patches of suitable habitat (nodes) with edges measuring dispersal between patches[4]. Data might either simply report the presence/absence of an edge (binary, unweighted networks), or provide a strength for each edge (weighted networks). In turn, these weights can represent a variety of ecologically-relevant quantities, depending on the system being described. For instance, edge weights can quantify interaction frequency (e.g., visitation networks[5]), interaction strength (e.g., per-capita effect of one species on the growth rate of another[3]), carbon-flow between trophic levels[6], genetic similarity[7], niche overlap (e.g., number of shared resources between two species[8]), affinity[9], dispersal probabilities (e.g., the rate at which individuals of a population move between patches[10]), cost of dispersal between patches (e.g., resistance[11]), etc.

Despite such large variety of ecological network representations, a common task is the identification of nodes of high importance, such as keystone species in a food web, patches acting as stepping stones in a dispersal network, or genes with pleiotropic effects. The identification of important nodes is typically accomplished through centrality measures[5,12]. A large number of centrality measures has been proposed, each probing complementary aspects of node-to-node relationships[13]. For instance, Closeness centrality[14,15] highlights nodes that are "near" to all other nodes in the network in terms of average distance (calculated as number of edges) from all other nodes. Whenever the effects of a node on another weaken along the path[16], then central nodes are those having the largest capacity to influence the others. Consider however highly modular networks, in which tightly knit communities of nodes are loosely connected to one another; then, one may be interested in identifying nodes that act as bridges connecting the different communities, allowing for the spread of perturbations across the

entire network. Stress centrality[17], and Betweenness centrality[15] serve this purpose. The choice of a

centrality measure thus depends on the research question at hand, and on the characteristics of the

data being analyzed. Different centrality measures have been used to identify keystone species in

networks of biotic interactions[5,18], to explore the robustness of metapopulations[19], to describe

connectivity patterns across fragmented habitats[20], to explore social behavior and pathogen spread

within populations[21], and to provide a theoretical background to support decision-making in

conservation planning and urban management[11,22] (see Supporting Information for a complete list).

Given the wide array of available techniques and the span of ecological applications, confusion may

arise when performing and reporting centrality analysis. An understanding of how the calculations are

performed, as well as a clear and sufficient reporting of the details of the analysis, are necessary in

order to avoid misinterpretation of the results and ensure the reproducibility of published studies.

In particular, edge weights exert a substantial influence on all measures of centrality. Many centrality

metrics rely on calculating the shortest path connecting any two nodes, and for weighted edges this

translates into finding the path with the smallest sum of weights. Edge weights definition is crucial in all

measures of centrality. When edge weights represent cost, resistance, or in general scale inversely to

the strength of the relationship between two nodes, then the definition of "shortest" paths retains its

simple interpretation. However, whenever edge weights are proportional to the strength of the

relationship between two nodes (e.g., probability of dispersal, interaction frequency, contact rate, etc.),

then minimizing the sum of edges along the path makes no sense: the data need to be transformed

prior to the analysis, or one has to choose an appropriate method able to deal with this situation. This

issue has been raised before[23,24] and is widely acknowledged among studies describing community

structure, where measures of network structure have been specifically developed to account for

weighted edges[25-27]. However, our analysis of the literature suggests that this issue is not fully

resolved, and that incorrect interpretations of centrality measures linger on, particularly among

weighted networks.

Furthermore, most published studies do not report with sufficient detail the calculation of the node-to-

node distance definition used in conjunction with the calculated centrality measures. As a

consequence, it is often impossible to evaluate the correctness of the calculations. To quantify the

extent of this problematic, we performed a systematic analysis of the ecological network literature. We

selected all ecological studies from the Web of Science that use a network approach, by searching on

topics TS=(network AND ecolog*), and limiting our search to Articles written in English in the science-

related citation indexes (SCI-EXPANDED, CPCI-S, BKCI-S, ESCI) since 2006. We further added 22

records obtained from other sources. From the resulting list of articles, we refined our search to those

mentioning "centrality", and from this final list of 210 articles we selected studies that studied

ecological communities using centrality metrics requiring the calculation of shortest paths, discarding

purely methodological studies. Finally, armed with a list of 129 articles, we checked whether the

analysis was reported with sufficient detail to determine whether the calculations were appropriate. In

Figure 1 we summarize this information with a frequency chart and in the Supporting Information we

provide the full list of articles. 63 articles (49%) did not report sufficient information on the calculation

of centrality to allow their replication, six of which were unclear even whether edges were binary or

weighted. Moreover, 61% of the studies using weighted edges may contain errors in how shortest path

centralities are calculated, a figure that grows to 89% if we limit the analysis to the case of weighted

interaction networks. Noticeably, 88% of the studies that correctly accounted for weighted edges in the

calculation of shortest paths, considered networks in which weights are inversely proportional to the

strength of association – thus not requiring transformation. Furthermore, nine studies (eight of which

examine interaction networks) calculated centrality using the binary version of the weighted data,

without providing any justification for this methodological choice. Certain ecological sub-disciplines use

alternative metrics for the study of energy flow across networks, such as food-chain metrics to

describe energy flow along food webs. Such metrics operate differently than shortest-path centralities,

and are not reviewed in this article.

The final goal of this study is to offer a precise and detailed protocol for the calculation of shortest path

centralities in ecological networks. Given that the correct way to calculate shortest path-based centrality measures depends on the type of network considered (binary or weighted) and on the meaning of edge weights, we use simple examples as well as real data to show how different approaches can result in unreliable estimates of centrality. Finally, since most studies omit to report key aspects of the definition of the edge weights, we propose a simple checklist to foster best-practices in the calculation and reporting of shortest path-related centrality analysis.

THE SHORTEST PATH IS FULL OF PITFALLS

In network analysis, the interaction between nodes can be thought of as a flow of information between the nodes that are linked by edges. The sequence of edges that information must cross in order to reach a specific node is called a path. It is generally assumed that the bulk of information between any two given nodes (among all the possible paths between these two nodes) passes through the shortest path connecting them (i.e., the one with "lowest weight"). However, it should be emphasized that while the concept of information flow is general, its immanence can differ dramatically from case to case, depending on which network feature weights quantify. For reference, in Table 1 we list the principal types of networks and the proportionality of the edges to the information flow between nodes found in the ecological literature.

The interpretation of a shortest path as the path that funnels the bulk of information flow relies on it being the least weight path (i.e., the path of least resistance) between two nodes. Indeed, all the shortest path algorithms currently available[28,29] and generally implemented in graph theory software (Table 2) seek to minimize the value of the path between two nodes calculated as the sum of the edge weights. The reason being that a minimization problem converges, while maximization can fail (see ref 24 for a detailed explanation). Nevertheless, the identification of the shortest paths is far from trivial, as one must pay attention to what edge weights represent. That is, one has to ensure that the edge weight is inversely proportional to the flow of information between the nodes. This condition is automatically fulfilled if the natural weight suggested by the network at study is already inversely

proportional to the information flow (e.g., resistance distance, dispersal time). However, when the

weight is directly proportional to the information flow (e.g., interaction frequency, individual transfer,

dispersal probability, pathogen transmission, energy transfer across food webs), it is necessary to

transform the edge weight in order to calculate the shortest paths, and the centrality measures that

rely on them (Table 3). In particular, this is important when using user-friendly software packages that

automate the calculation of centrality measures (Table 2).

There is a wide range of functions that can accomplish this transformation. For example, if a_{ij}

measures the flow of information between nodes i and j, the functions $1-a_{ij}$, $\exp(-a_{ij})$, $1/a_{ij}$, $\log(1/a_{ij})$ and

$\log(a_{ij}/(1-a_{ij}))$ are all found in the graph-theoretical literature[29-32]. Note that one must pay attention to

the range of values that the edge weights can span, and to the values induced by the transformation –

for example, one must avoid the use of negative edges (e.g., log transformation of values between 0

and 1), as these can greatly hamper the interpretation of the results. Finally, if the edge weights

represent probabilities, one should account for the independence (or lack of independence) of the

different edges.

AVOIDING THE PITFALLS

When computing any shortest path-related measure, various decisions need to be made, and the

wrong decision could lead to unexpected results, as we will illustrate in the following examples. We

focus on the effects on Betweenness (BC) and Closeness centralities (CC) as these are the most

commonly used centrality measures. In Figure 2 we summarize this decision process.

Binary or weighted?

The first methodological choice when computing shortest paths is whether to consider edge weights.

Binary data can be highly informative: for example, it has been used to identify species fundamental

niches[33], and key species in pollination networks[18]. Nonetheless, studies must clearly state whether

the analysis is performed on weighted or unweighted networks, and provide an ecological justification

supporting either choice[34]. Seminal studies[35] as well as recent ones[10] analyzed unweighted versions

of their data arguing that paths with fewer edges would inherently be stronger than those composed of

multiple ones. In our analysis of the literature, we have found nine studies that, despite using weighted

data for some calculations, revert to binary versions of the network for the calculation of centrality

measures without providing a justification (Figure 1). Indeed, note that unipartite projections of

bipartite networks result in weighted networks even when the original network is binary. Seven articles

out of nine use binary unipartite projections without any justification. Although calculations in these are

technically correct, one must be aware that the calculation of shortest paths and related measures in

binary and weighted versions of the same network can lead to dramatically different conclusions.

To see how discarding the edge weights can significantly change the results of network analysis, let us

start with a highly idealized example. In Figure 3a we show a network such that there are two possible

paths between from Busan (South Korea) to Almería (Spain): one containing two edges (Busan-

Chicago-Almería), and the other one three (Busan-Copenhagen-Marseille-Almería). If one considers

only the binary data (implicitly assuming that the distance between any cities is the same), the shortest

path from Busan to Almería would be crossing Chicago (one stopover vs. the two stopovers of the

other possible path). However, if the distance between cities is considered, it can be easily seen that

the path Busan-Copenhagen-Marseille-Almería is much shorter (~11000 vs. ~18000 Km).

For an ecologically relevant example, we constructed a connectivity matrix for a hypothetical bird

species living between 500 and 2000 m above sea-level, and with a typical habitat size of about 15

km^2. For this purpose, the Global Relief Database ETOPO1[36] data in the region we considered were

coarse-grained to 15 km^2 horizontal resolution, resulting in 787 habitat patches (Figure 4a). Dispersal

probability between patches was calculated as $p_{ij} = \exp(-\alpha\, d_{ij})$ following ref 35, where d_{ij} is the

geographical distance between the patches boundaries, and α is a parameter chosen to be 0.03 in

order to have a (hypothetical) median dispersal distance of 100 km. This probability can be stored in a

connectivity matrix (see Supporting Information, Figure S1) and graph theory can be used to identify

the habitat patches that have high Betweenness and Closeness centrality scores. Calculating BC and

CC on binary and weighted versions of this dataset resulted in markedly different outcomes. None of

the 20 habitat patches with highest BC and CC in the binary network (Figure 4b-c) match the ones

obtained from the analysis of the weighted network (Figure 4d-e). Note that to calculate Betweenness

and Closeness centralities in weighted data we first inverted the edge weights using $\log(1/p_{ij})$ (see

next section for a detailed explanation).

Modifying edge weights

The next question one should answer when computing shortest paths is whether edge weights are

inversely proportional to the information flow between the nodes in the network (Figure 2). If this is the

case, the shortest path between nodes can be calculated directly using the edge weights. If, on the

other hand, edge weights are proportional to the flow of information, one must transform them before

using shortest path algorithms. If one does not modify the edge weights, the shortest path algorithms

will either fail (and identify the longest, rather than shortest path), or will be unable to identify a

shortest path at all. We use the transformations $1/a_{ij}$ or $\log(1/a_{ij})$ for the edges (although for this

operation there are several alternative options reviewed below).

For example, let us consider a small network of four primates sharing a certain number of parasites

(Figure 5a). In order to detect the primate mediating the transmission of infectious diseases in this

network, one could identify the primate displaying the largest number of parasites common to other

primates – indicated by stronger edges. In this simple network it is easy to verify that most of the paths

with the highest edge weight (largest number of shared parasites) pass through primate A. However, if

BC and CC were calculated directly on unmodified edge weights, we would conclude that primate B is

the key primate in this network (Figure 5a). If, on the other hand, we transform edge weights using the

function $1/a_{ij}$, we correctly identify primate A as the primate with the highest BC and CC (Figure 5b).

Using a real-world case, the impact of not reversing edge weights can be illustrated using the Global

Mammal Parasites Database (GMPD, https://parasites.nunn.lab.org) containing data on 542 primate

species and their 750 parasites (ref 37 and references therein). To identify the species mediating the

transmission of parasites (similarly to ref 38), a connectivity matrix linking primates that have been

found to host the same parasite was built (see Supporting Information, Figure S2). Edge weights were

defined as the number of shared parasites between any two species. Therefore, edge weight is

directly proportional to the relationship strength between two species and, as in the previous example,

the quantity should be transformed before calculating the shortest paths. In Figure 6a we show the

lack of correlation between species rankings based on the centrality scores calculated on modified

and unmodified edge weights. For this example, we use the inverse of the edge weights, e.g., $1/a_{ij}$.

Interestingly, the BC scores calculated using the modified edge weights highlight only few species,

one of which has by far the highest BC score. Instead, if we directly use the unmodified values,

several more species have comparable BC scores. This is not surprising if we consider that, when

using the raw weights, shortest paths pass through weak connections, which are likely to be

numerous. Differences in ranks are substantial. For instance, among the top ten high-Betweenness

species identified using modified weights, only one is also among the high-Betweenness species

identified using raw weights (see Supporting Information, Table S1).

Likewise, the results from the Closeness-based rankings (Figure 6b) show that species rankings

based on CC also differ significantly between modified and unmodified edge weights. The CC scores

calculated on modified edge weights also support the importance of a handful of species (see

Supporting Information, Table S1). Unlike with BC, nine of the top 10 high-Closeness species are the

same ones when using the unmodified weights (also in Supporting Information) but the exact ranking

differs between the two cases.

Other modifying functions

Adding constants

When edges are directly proportional to the information flow, it is frequent practice to make them

inversely proportional by subtracting their value from a theoretical maximum or some other meaningful

constant[29,32]. For example, in the case of transfer probabilities (migration, mass, energy, networks),

one could choose to subtract the edge weights a_{ij} from 1. However, the new edge weight $1-a_{ij}$ biases

the calculation of the shortest paths towards the path with the lowest number of edges (because

probabilities sum to one, nodes with many edges tend to have lower values).

Again, we will use the simple toy matrix presented in Figure 3, where we show a network connecting

different cities. We then consider three different quantities to weight the edges that represent the

movement of researchers between these cities. In the first case (Figure 3b), edge weights quantify the

number of researchers who moved from one site to another. In this case, the path sustaining the

largest "flow of researchers" between Busan and Almería is the three-steps path Busan-Copenhagen-

Marseille-Almería (30 vs 10). However, applying a shortest path algorithm directly to this network

would identify the two-step path as more important. One possible way to reverse the edge weight is to

subtract the edge weights from a large (and in most instances arbitrary) constant C – de facto adding

a constant to all edges. For example, if one chooses C = 100, now the largest edge weights (those

representing largest flows) are the smallest and would hence be identified by the shortest-path

centrality algorithm as more central. However, one can easily verify that such a transformation did not

change the fact that the two-step path is the shortest between Busan and Almería (190 vs 270). The

reason is that adding a constant to all the edge weights biases the shortest paths algorithm towards

paths with fewer edges.

In Figure 3c we see that subtracting the edge weights from a theoretical maximum C (in the case of

transfer probabilities of interaction frequencies, C = 1) makes the three-step path with higher

information flow the shortest path ($3 \times (1-0.9)$ vs $2 \times (1-0.1)$). However, this transformation does not work

in all cases. In fact, if the edge weights, as frequently occurs in ecological studies, span different

orders of magnitude (e.g., Figure 3d), the three-step path will not be the shortest path anymore ($3 \times (1-$

$10^{-3})$ vs $2 \times (1-10^{-5})$), as in Figure 3b. Furthermore, we note that this type of transformation, even when

it is likely to work, cannot be used for all the edge weights. For example, it cannot be used for

probabilities, as the values of $\log(1-a_{ij})$ are negative and, consequently, cannot be used to find shortest

paths (see next section).

A real-world example of the effect of adding constants to the edge weights is provided in the

Supporting Information (Figures S3, S4 and S5).

Negative weights and loops

Another way to reverse the edge weights is to reverse the sign of the weights (i.e., using $-a_{ij}$).

However, as shortest path algorithms seek to minimize the value of a path, they would keep looping

closed paths (cycles) ad infinitum, without ever converging. It must be noted that there are alternative

algorithms that can handle negative edges values (e.g., the Bellman-Ford-Moore algorithm[39]) cannot

handle cycles. As cycles are essentially ubiquitous in ecological applications, edge weight

transformations that result in negative values should therefore be avoided. As an example, consider a

simple toy network depicting the carbon flow between different layers of a food chain (Figure 7a). In

this case, given that the flow of carbon is directly proportional to the strength of the connection

between two layers of the food chain, we need to transform the weights. However, if one uses $-a_{ij}$

(Figure 7b), the shortest path algorithms would never converge, and would keep circling the loop. On

the other hand, using another weight reversing function, such as $1/a_{ij}$, would correctly identify the 2^{nd}

order consumers as key species pivoting the carbon flow in this network example (Figure 7c).

Independence of probabilities

We should note an important aspect to consider when calculating the lengths of paths in networks:

when edges represent probabilities, as for instance dispersal probabilities, we must question the

independence of the edges in order to calculate meaningful values for the overall probability of the

entire path. From a practical point of view, this means that when calculating the value of a path from

node A to node C passing through node B (path ABC), we need to postulate that the path BC does not

depend on the path used to reach B. When edges represent independent probabilities, the probability

along a path containing multiple nodes is the product of the probabilities of all the paths linking the

nodes. Interestingly, in the case of independent probability edges, converting edge weight a_{ij} into

distance using $\log(1/a_{ij})$ nicely transform probability product along multiple nodes path into distances

addition along this path, conserving weights relative contribution to the path and avoiding weights

distortion in shortest paths algorithm. Without that edge transformation, the path ABC will be the sum

of the probabilities of paths AB and BC, resulting in the identification of most improbable paths as

those more central (see ref 24 for a detailed explanation, and Figure S6 for an example using the

ETOPO1 dataset).

One or all shortest paths?

In most networks, whether binary or weighted, there may be more than one shortest path connecting

any two nodes. To account for this fact, an alternative definition of Betweenness centrality based on

random walks[40] and a generalization of node centrality that considers both edge weight and number

when calculating centrality measures[32] have been developed. Although the discussion of the pros and

cons of the different formulations falls beyond the scope of this study, we encourage the reader to be

aware that considering a single or all shortest paths may also introduce differences in the resulting

centrality values, and the decision should hence also be reported. This is of particular importance

when comparing centrality metrics with food chain length related metrics, where all paths may be

considered.

WIDENING THE PATH

Network analysis has been developing quite independently in different branches of ecology. However,

dissemination between ecological disciplines and reproduction of published studies are being

hampered, at least partially, by the lack of transparency when describing the methodologies used.

Establishing a protocol for the analysis and reporting of calculations would ease these obstacles, and

boost the use of centrality metrics for unconventional uses. For example, in a species-interaction

network (where species are typically considered closer if they interact with higher frequencies), one

could purposely choose to calculate shortest path-centrality measures without transforming the

weights in order to study the effect of weak interactions across the network. For this reason, we urge

all the researchers applying graph theory to ecological data to pay special attention when reporting

their calculations, and, in particular, to provide a description of the network and edge weight they

used.

Here, we provide a checklist of crucial methodological information that should always be reported

(Figure 2). Following this guide ensures the study reports sufficient information to allow reproducibility,

a quick understanding of the methods by readers from other fields, and that the decision process prior

calculations is done sequentially.

1) *A clear definition of what nodes and edges represent.* Nodes and edges depict different entities and

relationships in different ecological studies. Nodes may represent proteins, genes, individuals,

populations, species, sites, etc., and edges may depict interactions of different kind, or movement

measured in numerous ways. A clear definition of nodes and edges enables a faster and deeper

understanding of the rationale and methodology of the analysis by readers from different disciplines.

2) *Are edges binary or weighted?* If edges are weighted, one needs to report the proportionality of the

edge weight to the information flow between the nodes in order to evaluate whether edges need to be

modified. In particular, one should ensure that there is no contradiction between the weights of a

network and the interpretation of shortest paths.

3) *Report the eventual transformation applied to edge weight before the calculation of centrality*

measures. Furthermore, carefully justify any conceptual reason to not transform edge weights, or to

use the unweighted versions of weighted data. These decisions result in the identification of different

central nodes or edges, and hence should be justified from an ecological perspective.

4) *Report the formula used for the calculation of the centrality measure, and whether it considers all*

shortest paths or only one. To ensure reproducibility and a deeper understanding of what the results

represent.

5) *Report the full version of the software or package used for the calculation of centrality*. To ensure

reproducibility and to account for potential future updates in the algorithms used by different packages.

CONCLUSIONS

Graph theory enables to achieve precious insights on ecological networks. For this reason, it has

gained popularity in ecology and has developed quite independently in different disciplines, becoming

a routine analysis in ecological studies. Our analysis of the literature evidenced that this familiarity is

however associated to a lack of methodological rigor in the published studies. Indeed, by reading the

methodological sections of a large portion of the published studies, we were not able to clearly

ascertain what edges represented when centrality measures calculations were carried out. The

increasing popularity of packages for the analysis of ecological networks will only boost the use of

tools and methodologies researchers may be unfamiliar with. Using both theoretical and real-world

case studies we showed that oversights in the methods and calculations can lead to radically different

results. Hence it is fundamental to establish a code of good practices that guides researchers through

the calculations, while ensuring the correct calculation of metrics across fields, aiding understanding

from other fields and the reproducibility of results. For that reason, in this article we provide an

overview of different methods to meaningfully calculate shortest paths and related centrality measures

in ecological systems, and a checklist to ensure clear and sufficient reporting of such calculations. We

hope that following the protocol we suggest will further increase the popularity of centrality measures

in ecology, and, at the same time, guarantee the reproducibility of these studies.

REFERENCES

1. Dale, M. R. T. Applying Graph Theory in Ecological Research. 344pp. Cambridge University
Press, 2017). ISBN 9781316105450.
2. Delmas, E., et al. Analysing ecological networks of species interactions. Biological Reviews
(2017).
3. Brose, U., et al. Spatial aspects of food webs. In: Dynamic Food Webs: Multispecies
Assemblages, Ecosystem Development, and Environmental Change. Eds. De Ruiter, P.C.,

388 Wolters, V. & Moore, J.C. Academic Press (2005).

389 4. Altermatt, F., Seymour, M. & Martinez, N. River network properties shape α☐diversity and
390  community similarity patterns of aquatic insect communities across major drainage basins. J.
391  Biogeog. 40, 2249-2260 (2013).

392 5. Martín González, A. M., et al. The macroecology of phylogenetically structured hummingbird-
393  plant networks. Glob. Ecol. Biogeogr. 24, 1212-1224 (2015).

394 6. Jordán, F. Keystone species and food webs. Philos. Trans. R. Soc. B Biol. Sci. 364, 1733–
395  1741 (2009).

396 7. Rozenfeld, A. F., et al. Network analysis identifies weak and strong links in a metapopulation
397  system. Proc. Natl. Acad. Sci. U.S.A. 105, 18824–9 (2008).

398 8. Pires, M. M., Marquitti, F. M. & Guimarães Jr, P. R. The friendship paradox in species-rich
399  ecological networks: Implications for conservation and monitoring. Biol. Cons. 209, 245-252
400  (2017).

401 9. Luthe, T. & Wyss, R. Resilience to climate change in a cross-scale tourism governance
402  context: a combined quantitative-qualitative network analysis. Ecol. Soc. 21(1) (2016).

403 10. Zamborain-Mason, J., Russ, G.R., Abesamis, R.A., Bucol, A.A. & Connolly, S.R. Network
404  theory and metapopulation persistence: incorporating node self-connections. Ecol. Lett. 20,
405  815–831 (2017).

406 11. Girardet, X., Conruyt-Rogeon, G. & Foltête, J. C. Does regional landscape connectivity
407  influence the location of roe deer roadkill hotspots? Eur. J. Wildl. Res. 61, 731–742 (2015).

408 12. Treml, E. A., Halpin, P. N., Urban, D. L. & Pratson, L. F. Modeling population connectivity by
409  ocean currents, a graph-theoretic approach for marine conservation. Landscape Ecol. 23(1),
410  19-36 (2008).

411 13. Kool, J. T., Moilanen, A. & Treml, E. A. Population connectivity: recent advances and new
412  perspectives. Landscape Ecol. 28, 165 (2013).

413 14. Bavelas, A. Communication Patterns in Task-Oriented Groups. J. Acoust. Soc. Am. 22, 725
414  (1950).

415 15. Freeman, L. C. Centrality in Social Networks Conceptual Clarification. Soc. Networks, 1, 215-
416  239 (1979).

417 16. Burt, R. S. Decay functions. Soc. Networks, 22, 1-28. Available from http://www.statnet.org/
418  (2000).

419 17. Shimbel, A. (1953). Structural parameters of communication networks. The bulletin of
420  mathematical biophysics, 15, 501-507.

421 18. Martín González, A. M., Dalsgaard, B. & Olesen, J. M. Centrality measures and the importance
422  of generalist species in pollination networks. Ecol. Complex. 7, 36–43 (2010).

423 19. Thompson, P. L., Rayfield, B. & Gonzalez, A. Robustness of the spatial insurance effects of

biodiversity to habitat loss. Evol. Ecol. Res. 16(6), 445-460 (2015).

20. Carroll, C., McRae, B. H. & Brookes, A. Use of Linkage Mapping and Centrality Analysis
Across Habitat Gradients to Conserve Connectivity of Gray Wolf Populations in Western North
America. Conserv. Biol. 26, 78–87 (2012).

21. Aplin, L. M., et al. Individual personalities predict social behaviour in wild networks of great tits
(Parus major). Ecol. Lett. 16, 1365–1372 (2013).

22. Poodat, F., Arrowsmith, C., Fraser, D. & Gordon, A. Prioritizing Urban Habitats for Connectivity
Conservation: Integrating Centrality and Ecological Metrics. Environ. Manage. 53(3), 664-674
(2015).

23. Foltête, J. C., Clauzel, C., Vuidel, G. & Tournant, P. Integrating graph-based connectivity
metrics into species distribution models. Landsc. Ecol. 27(4), 557-569 (2012a).

24. Costa, A., Petrenko, A. A., Guizien K. & Doglioli, A. M. On the calculation of betweenness
centrality in marine connectivity studies using transfer probabilities. PLOS ONE 12(12):
e0189021 (2017).

25. Newman, M.E.J. Analysis of weighted networks. Phys. Rev. E, 70, 56131 (2004).

26. Almeida-Neto, M. & Ulrich, W. A straightforward computational approach for measuring
nestedness using quantitative matrices. Environ. Model. Softw. 26, 173–178 (2011).

27. Dormann, C. F. & Strauss, R. A method for detecting modules in quantitative bipartite networks.
Meth. Ecol. Evol. 5(1), 90-98 (2014).

28. Dijkstra, E. W. A note on two problems in connexion with graphs. Numerische Mathematik, 1:
269–271 (1959).

29. Brandes, U. On Variants of Shortest-Path Betweenness Centrality and their Generic
Computation. Soc. Networks, 30, 136-145 (2008).

30. Augustin, N. H., Cummins, R. P. & French, D. Exploring spatial vegetation dynamics using
logistic regression and a multinomial logit model. J. Appl. Ecol. 38, 991-1006 (2001).

31. Estrada, E. & Bodin, O. Using network centrality measures to manage landscape connectivity.
A short path for assessing habitat patch importance. Ecol. Appl. 18, 1810-1825 (2008).

32. Opsahl, T., Agneessens, F. & Skvoretz, J. Node centrality in weighted networks: Generalizing
degree and shortest paths. Soc. Networks, 32, 245–251 (2010).

33. Fründ, J., McCann, K. S. & Williams, N. M. Sampling bias is a challenge for quantifying
specialization and network structure: lessons from a quantitative niche model. Oikos, 125(4),
502-513 (2016).

34. Mello, M. A. R., et al. Keystone species in seed dispersal networks are mainly determined by
dietary specialization. Oikos, 124(8), 1031-1039 (2015).

35. Minor, E. S. & Urban, D. L. Graph theory as a proxy for spatially explicit population models in
conservation planning. Ecol. Appl. 17, 1771–1782 (2007).

36. Amante, C. & Eakins, B. W. ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis. NOAA Technical Memorandum NESDIS NGDC-24. National Geophysical Data Center, NOAA. doi:10.7289/V5C8276M (2009).

37. Stephens, P. R., et al. Global mammal parasite database version 2.0. Ecology, 98(5), 1476-1476 (2017).

38. Gómez, J. M., Nunn, C. L. & Verdú, M. Centrality in primate-parasite networks reveals the potential for the transmission of emerging infectious diseases to humans. Proc. Natl. Acad. Sci. U.S.A. 110, 7738–7741 (2013).

39. Bang-Jensen, J. & Gutin, G. "Section 2.3.4: The Bellman-Ford-Moore algorithm". Digraphs: Theory, Algorithms and Applications (First ed.). ISBN 978-1-84800-997-4 (2000).

40. Newman, M.E.J. A measure of betweenness centrality based on random walks. Soc. Networks, 27(1), 39-54 (2005).

41. Magris, R. A., Treml, E. A., Pressey, R. L. & Weeks, R. Integrating multiple species connectivity and habitat quality into conservation planning for coral reefs. Ecography, 39, 649–664 (2016).

42. Emer, C., et al. Seed-dispersal interactions in fragmented landscapes – a metanetwork approach. Ecol. Lett. 21, 484–493 (2018).

43. Herrera-Arroyo, M. L., et al. Seed-mediated connectivity among fragmented populations of Quercus castanea (Fagaceae) in a Mexican landscape. Am. J. Bot. 100, 1663–1671 (2013).

44. Naujokaitis-Lewis, I. R., Rico, Y., Lovell, J., Fortin, M.J. & Murphy, M. A. Implications of incomplete networks on estimation of landscape genetic connectivity. Conserv. Genet. 14, 287–298 (2013).

45. Ruggera, R. A., Blendinger, P. G., Gomez, M. D. & Marshak, C. Linking structure and functionality in mutualistic networks: Do core frugivores disperse more seeds than peripheral species? Oikos, 125, 541–555 (2016).

46. Lozano, S., Mateos, A. & Rodríguez, J. Exploring paleo food-webs in the European Early and Middle Pleistocene: A network analysis. Quat. Int. 413, 44–54 (2016).

47. Reino, L., et al. Networks of global bird invasion altered by regional trade ban. Sci. Adv. 3, 1–9 (2017).

48. Blaszczyk, M. B. Consistency in social network position over changing environments in a seasonally breeding primate. Behav. Ecol. Sociobiol. 72 (2018).

49. Livi, C. M., Jordán, F., Lecca, P. & Okey, T. A. Identifying key species in ecosystems with stochastic sensitivity analysis. Ecol. Model. 222, 2542–2551 (2011).

50. Lai, S. M., Liu, W. C. & Jordán, F. On the centrality and uniqueness of species from the network perspective. Biol. Lett. 8(4), 570-573 (2012).

51. Csardi, G. & Nepusz, T. The igraph software package for complex network research, InterJournal Complex Systems, 1695 (2006).

52. Butts, C. T. Tools for Social Network Analysis, R Package "sna" (2016).

53. Opsahl, T. Software for Analysis of Weighted, Two-Mode, and Longitudinal Networks, R
Package "tnet" (2015).

54. Mrvar, A. & Batagelj, V. Analysis and visualization of large networks with program package
Pajek. Complex. Adapt. Syst. Model. 4:6 (2016).

55. Borgatti, S. P., Everett, M. G. & Freeman, L. C. Ucinet for Windows: Software for Social
Network Analysis. Harvard, MA: Analytic Technologies (2002).

56. Saura, S. & Torné, J. Conefor Sensinode 2.2: a software package for quantifying the
importance of habitat patches for landscape connectivity. Environ. Modell. Softw. 24(1), 135-
139 (2009).

57. Foltête, J. C., Clauzel, C. & Vuidel, G. A software tool dedicated to the modelling of landscape
networks. Environ. Model. Softw. 38, 316–327 (2012b).

58. Pascual-Hortal, L. & Saura, S. Comparison and development of new graph-based landscape
connectivity indices: towards the priorization of habitat patches and corridors for conservation.
Landscape Ecol., 21(7), 959-967 (2006).

59. Saura, S. & Pascual-Hortal, L. A new habitat availability index to integrate connectivity in
landscape conservation planning: comparison with existing indices and application to a case
study. Landsc. Urban Plan. 83(2-3), 91-103 (2007).

ACKNOWLEDGMENTS

This work was supported by IBS-R028-D1. AMMG is supported through a Marie Skłodowska-Curie

Individual Fellowship (H2020-MSCA-IF-2015-704409), and thanks the Danish National Research

Foundation for its support of the Center for Macroecology, Evolution and Climate (Grant number

DNRF96). We thank Sonia Agüera-González (@immunosoni) for the illustrations in Figure 3

(https://www.behance.net/soniaguera6595).

AUTHOR'S CONTRIBUTIONS

AC, AMMG, SA, AAP, KG & AMD designed the study; AC conducted the data analysis, AMMG

performed the literature search and AC & AMMG performed the literature analysis and wrote the first

draft of the manuscript. All authors contributed substantially to the drafts and gave final approval for

publication.

ADDITIONAL INFORMATION

Competing interests

The authors declare no competing interests.

Data availability

This article uses no data.

 Tables

534 Table 1. Summary of different network types, describing the type of information flowing along edges

535 and the consequences for weight transformation. Landscape networks are spatially-

536 explicit, while interaction networks depict relationships among entities. Social networks are here

537 included as interaction networks, where shared traits include shared interacting organisms, foraging

538 time, etc. depicting social relationships or common characteristics.

Network type	Edge weight	Proportionality to information flow	Requires transformation	Example references
Landscape	Cost-distance	Inverse	No	11, 20
	Dispersal probability	Direct	Yes	10, 41
	Dispersal time	Inverse	No	12
	Exchange of individuals	Direct	Yes	42
	Genetic similarity	Direct	Yes	43, 44
Interaction	Frequencies	Direct	Yes	45, 46
	Shared traits, affinity	Direct	Yes	47, 48
	Trophic (or energy) flow	Direct	Yes	49, 50

540   Table 2. Analytical packages commonly used for the calculation of centrality in ecological studies and

541   references for each package. Only *sna* and *Ucinet* packages caution in their documentation about

542   edge transformations when calculating shortest paths.

Package	References
igraph	51, http://igraph.org/
sna	52, https://cran.r-project.org/web/packages/sna/index.html
tnet	53, https://cran.r-project.org/web/packages/tnet/index.html
Pajek	54, http://mrvar.fdv.uni-lj.si/pajek/
Ucinet	55, https://sites.google.com/site/ucinetsoftware/home
CONEFOR	56, http://www.conefor.org/
Graphab	57, https://sourcesup.renater.fr/graphab/en/home.html

543 Table 3. Measures of shortest path-related centrality measures commonly used in ecological network analysis. Notice that other common

544 centrality measures are based on eigenvectors or dissimilarity scores instead of on the identification of shortest paths and are hence not

545 considered in this work.

Centrality Measure	Definition	Formula	Intended Network type	Reference
Betweenness, BC	Quantifies the proportion of shortest paths g between any two nodes i, j, that pass through a focal node v.	$$BC(v) = \frac{\sum_{i \neq v \neq j} g_{ij}(v)}{g_{ij}}$$	All types	15
Stress, SC	Measures the number of shortest paths g between any two nodes i, j, that pass through a focal node v.	$$SC(v) = \sum_{i \neq v \neq j} g_{ij}(v)$$	All types	17
Closeness, CC	Measures the average length of the shortest paths from a node v to all the other nodes in the network.	$$CC(v) = \frac{N-1}{\sum_{j} g_{vj}}$$	All types	15
Integral Index of Connectivity, IIC	Measures the degree of connectivity of the entire landscape (of total area A_L) through the calculation of the number of edges in the shortest path nl_{ij} between patches with area a_i and a_j.	$$IIC(v) = \frac{\sum_{i=1}^{n} \sum_{j=1}^{n} a_i a_j /(1+nl_{ij})}{A_L^2}$$	Binary landscape networks	58
Probability of Connectivity Index, PC	Quantifies the probability that two species randomly placed across a patchy landscape (of total area A_L) fall into habitat patches a_i and a_j that are reachable from each other with a maximum connectivity probability p_{ij}, defined as the maximum product probability of all possible paths between patches i and j (including single-step paths).	$$PC(v) = \frac{\sum_{i=1}^{n} \sum_{j=1}^{n} a_i a_j \dot{p}_{ij}}{A_L^2}$$	Landscape networks	59

Figure 1. Summary of our literature analysis, detailing the number of studies by (a) edge weights, and

(b) whether the information provided was sufficient or insufficient, and network type (as "landscape",

"interaction", and "others", which include social, co-occurrence, etc. networks). We only categorize as

"correct" or as "wrong" studies on which we had enough information to support such a claim, and as

"seems correct" and "seems wrong" studies on which the insufficient information available suggest that

calculations are correct or wrong, respectively. Notice that (1) most of the correct or probably correct

studies use unweighted edges (n = 52, 66%); (1) for all network types half of the studies do not report

enough information to validate whether calculations were correct (n = 63, 49%); and that (3) numerous

studies report unclear calculations (n = 19). Differences in the number of oversights in centrality

calculations between weighted and landscape and interaction networks are probably due to the fact

that in interaction networks weighted edges typically require transformation (see Table 1), whereas in

landscape networks edge weights tend to be inversely proportional, requiring no transformation.

Figure 2. Scheme illustrating the step-by-step decision process for the calculation of shortest path-

centrality measures in ecological networks and the 5-point guide of information require for good-

practices.

Figure 3. Toy networks depicting two possible paths to arrive from the city of Busan to the city of

Almería. Edges connecting the different cities quantify distance in Km (a), or using different

measurements of the movement of researchers between these cities (b-d; see text for explanation).

Figure 4. Land topography and ocean bathymetry (m) of the Italian region (data from the ETOPO1

Global Relief database; doi:10.7289/V5C8276M). A hypothetical bird species lives between 500 and

2000 m a.s.l., with a typical habitat size of about 15 km^2 and a dispersal distance of 100km. a) Pixels

in lighter tones denote patches of suitable habitat (colored in black in the following panels). After

computing Betweenness (BC) and Closeness centrality (CC), we highlighted in red the 20 pixels with

highest BC in the binary (b) and weighted network (c), and the 20 pixels with highest CC in the binary

(d) and weighted network (e). Note the differences in the identification of the pixels between weighted

and binary versions of the data.

Figure 5. Toy network describing the social affinity between four primates. Edges quantify proportion

of shared parasites. Drawings are public domain (https://commons.wikimedia.org/wiki/File:Primates-

drawing.jpg).

Figure 6. Scatter plot showing species ranks based on Betweenness (a) and Closeness centrality (b)

values calculated on untransformed and inversed edge weights.

Figure 7. Toy network describing the carbon flow through a marine food web. Drawings by Siyavula

Education under a CC BY 2.0 license (https://www.flickr.com/photos/121935927@N06/13578843423).

a_{ij}

Describe network, report what nodes and edges represent

Be aware of the proportionality to the information flow

Are you calculating shortest-path based measures?

No

Yes

Report whether edges are binary or weighted and their proportionality to the strength of the relationship between nodes

Are a_{ij} weighted?

Use it as it is

No

Yes → Make network binary and justify

Use it as it is

Are a_{ij} directly proportional to the relationship strength between nodes?

Use it as it is

No

Yes

Are a_{ij} a probability?

No

Yes

Reverse edges using any suitable function

Report which transformation you used

Are a_{ij} independent from each other?

No

Yes

Report the formula used for the calculation of centrality

Reverse edges using any suitable function

Reverse edges using any suitable function that takes into account the dependency of the edges

Report full version of software or package used for the analysis

a)

Primate A

Primate B

Primate D

Primate C

b)

Primate A

1/2

1/2

Primate B

Primate D

Primate C