Metapopulation connectedness and the evolution of multiple life-history traits

Masier, Stefano[1,2*]; Bonte, Dries[1,3]

1: Terrestrial Ecology Unit, UGent, Ghent (Belgium)

2: Stefano.Masier@UGent.be; phone number: +32 (0)9 264 5258

3: Dries.Bonte@UGent.be

*: corresponding author

Abstract

Loss of habitat connectivity is a renowned threat to metapopulations and anticipated to impact

evolutionary dynamics. Dispersal-related traits are expected to be selected at metapopulation

level, but local demographic changes may also influence them. However, the exact

consequences have never been quantified in detail.

Using experimental metapopulations of two-spotted spider mites, we investigated how

connectedness loss affects trait evolution. We also implemented a randomization treatment to

separate local- and metapopulation-level selection.

We demonstrate selection on multiple traits related to population growth rate and dispersal,

but not dispersal rate per se. With longer dispersal distances, starvation resistance improved;

simultaneously, shorter distances translated into a quicker decision to disperse. Intrinsic

growth rates were lower in individuals from the most connected patches; as randomization did

not disrupt this pattern, fast epigenetic evolution must have been involved.

We demonstrate the importance of multivariate trait evolution at multiple spatial scales to

understand metapopulations eco-evolutionary dynamics.

Introduction

Besides habitat loss, habitat fragmentation is one of the most important human-induced drivers of declines in biodiversity (Collinge, 1998; Forman, 2000; Heilman *et al.*, 2002). Most species live in spatially heterogeneous landscapes of suitable habitat patches interspersed with areas of less suitable matrix. In such landscapes, dispersal enables the exchange of individuals among populations of different habitat patches, resulting in an overarching metapopulation. Loss of habitat connectivity influences metapopulation dynamics at every level of biological organization, from individuals to ecosystems (Gibbs and Stanton, 2001; Crooks, 2002; Cushman, 2006).

Connectivity is a species-specific metric, determined by the species' movement capacity and its dispersal between populations living in different patches (Tischendorf and Fahrig, 2001). In contrast, the connectedness of resource patches within spatial networks is only dependent on the extent to which a landscape facilitates or impedes the movements of organisms and their genes among these patches (Taylor *et al.*, 1993; Rudnick *et al.*, 2012). It is therefore affected by the structural properties –the number, shape, dimensions and interdistance– of the suitable patches within a landscape (Tischendorf and Fahrig, 2000; Wang, Blanchet and Koper, 2014). Connectedness thus impacts connectivity by mediating both the ecological and evolutionary determinants of the flows among patches of both individuals and genes (Bonte, Masier and Mortier, 2018). Loss of habitat connectivity decreases the chances of reestablishment of extinct populations, thereby putting the metapopulation at risk (Fuller, Doyle and Strayer, 2015; Thompson, Rayfield and Gonzalez, 2017). Habitat connectivity is thus central to metapopulation conservation.

While the ecological effects of reduced habitat connectivity on single species, and those integrated into food and mutualistic networks are well studied from a theoretical (Melián and

Bascompte, 2002; Kondoh, 2003; Liao *et al.*, 2017) and empirical perspective (Dobson *et al.*, 2006; Fenoglio *et al.*, 2010; Valladares, Cagnolo and Salvo, 2012), evolutionary impacts are anticipated as well, but rarely directly tested (Cheptou *et al.*, 2017). Studies on the evolutionary consequences of connectivity loss have mostly focused on the evolution of inbreeding, mating systems or dispersal rates (Stow *et al.*, 2001; Fahrig, 2003; Andersen, Fog and Damgaard, 2004; Aguilar *et al.*, 2006; Keyghobadi, 2007; Bonte, Masier and Mortier, 2018). Dispersal is anticipated to be a prime target of selection in metapopulations (Bonte and Dahirel, 2017), but should not be treated as a simple trait related to emigration rate, as it integrates multiple processes during the phases of departure, transience and settlement (Bowler and Benton, 2005; Clobert *et al.*, 2009). Costs and trade-offs may manifest themselves during each of these three phases, often in interaction with other life history traits (Bonte *et al.*, 2012), leading to life-history syndromes at both the individual, population and metapopulation-level (Clobert *et al.*, 2012). It can be assumed that connectedness loss mostly influences the costs of the transfer phase, thereby imposing selection by lowering departure rates (Bonte *et al.*, 2003). Evolutionary adaptations to reduce costs of the other dispersal phases may, however, be prominent as well and result in dispersal syndromes (Clobert *et al.*, 2012). Evolutionary changes in (pre-) departure behaviours (Derr, Aldent and Dinglet, 1981; Bell *et al.*, 2005; Cañete *et al.*, 2007), dispersal timing (Lakovic, Poethke and Hovestadt, 2015), movement (Korb and Linsenmair, 2002; Baguette and Van Dyck, 2007) and post-settlement strategies (Vessby and Wiktelius, 2003; Hansson, Bensch and Hasselquist, 2004) in response to altered habitat connectedness are becoming well-documented. Because interactions and/or trade-offs between two or more of these traits are also possible (Hazell *et al.*, 2005; van der Linde and Sevenster, 2006), complex multivariate selection is expected to be the rule in metapopulations.

Habitat patches are usually not organized in regular, symmetrical spatial networks. In reality,

spatial networks are heterogeneous and modular with respect to their topology (Fahrig, 2003;

Urban and Skelly, 2006; Ferrari, Lookingbill and Neel, 2007). A few patches are typically

more connected to others and thereby serve as hubs in the network (Van Langevelde, 2000;

Calabrese and Fagan, 2004). Such asymmetries in connectedness impact local

immigration/emigration balances and lead to source-sink dynamics and the emerging

metapopulation dynamics (Poethke, Dytham and Hovestadt, 2011; Dey, Goswami and Joshi,

2014; Wang, Haegeman and Loreau, 2015). When dispersal rates are equal among patches,

highly connected patches experience a net immigration rate, thereby imposing an increased

resource competition (Baguette and Schtickzelle, 2003; Amarasekare, 2008). Such local

competition can then impose selection on traits related to stress-resistance, or select for a

reduced population growth rate to counter resources overshooting in kin-structured

populations (McClain, 1995). Finally, if emigration-immigration balances change across

patches, different levels of relatedness may affect local reproductive (Macke *et al.*, 2011) and

dispersal strategies (Van Petegem *et al.*, 2018). As a result, even in metapopulations

consisting of a network of patches with identical habitat, local selective pressures can be

different (Rossetti *et al.*, 2014) and local selection on traits related to density dependence, kin

competition and stress resistance may act in concert with metapopulation-level selection on

dispersal.

Given the importance of trait evolution for metapopulation dynamics, persistence and rescue,

a deep understanding of the eco-evolutionary effects of connectivity loss is essential to guide

future conservation actions (Travis *et al.*, 2013; Urban *et al.*, 2016). We studied experimental

evolution in the spider mite *Tetranychus urticae* (Fronhofer *et al.*, 2014; De Roissart, Wang

and Bonte, 2015; Van Petegem *et al.*, 2018), to test whether and how different levels of

habitat connectedness affect trait evolution in a mite model system. We particularly focused

on traits related to dispersal and reproduction as theory predicts these to be under regional

and/or local selection in metapopulations. We expected individuals evolving in more

fragmented habitats to be characterised by a lower propensity of dispersal, as it is a more

costly event in such habitats, or to evolve lower dispersal costs by means of a higher

resistance to the environmental conditions during transfer (e.g., food deprivation). In order to

separate local and metapopulation-level selection, we disrupted local selection by reshuffling

mites among local patches in part of the metapopulations, to remove both *kin* (genetic

relatedness) and *kind* (phenotypic similarity) structure (Van Petegem *et al.*, 2018) and leave

the pure environmental effect of the habitat connectedness level.

Materials & Methods

Experimental evolution

Individuals from our focal species, the two-spotted spider mite *Tetranychus urticae* Koch were collected from our stock population and introduced in 24 artificial habitats composed by a 3-by-3 grid of food patches (bean leaves cuts) connected by Parafilm bridges (see fig.1).

Due to patch network design, each patch had a different level of connectedness inside the metapopulation (further referred to as *local connectedness*), with central patches having the highest number of links (8), corner patches the lowest (3) and side patches as intermediate (5) value.

The bridges varied in length between metapopulations between 4, 8 or 16 cm, thus determining the *metapopulation connectedness* of each setup: each connectedness level was replicated 8-fold, three of which were subjected to a randomisation procedure of the adult females. We implemented this randomisation procedure to test whether performance would evolve in response to local connectedness as well.

In the randomization treatments, all females from a single metapopulation were collected once per week using a thin pen brush, pooled together on a fresh bean leaf to reduce stress as much as possible and then randomly redistributed among the patches of the same metapopulation, maintaining the same local densities but reshuffling the individuals so that any one of them could randomly end up on any patch of its metapopulation. In order to ensure comparable levels of handling stress between treatments, the females from the non-randomization systems were also collected, not randomized and placed back on their original patch briefly afterwards. Contrasting trait values between the randomised and non-randomised experimental metapopulations indicate whether performance evolved at the metapopulation-level (no randomisation effects expected), and/or the local level (randomisation would then remove any signature of local evolution, hence rendering this effect statistically significant).

All experimental metapopulations (24 in total) were initialized in September 2016 with five

freshly inseminated females and 2 additional adult males on each patch (hence 1272

individuals overall) using a soft-haired brush and monitored for two weeks to ensure the

settlement of the population. This standardised initialisation ensures equal starting conditions

and sufficient genetic variation (as for instance demonstrated in Van Petegem *et al.*, 2018).

A more detailed outline of the focal species and the long term experimental setup can be

found in Supplementary Materials SI 1 and Masier & Bonte, *in prep*.

Trait evolution following common garden

25 weeks from the start of the experiment (approx.12 generations under the experimental

conditions; (Hance and Impe, 1999), we collected individuals from all the different patches to

quantify whether relevant ecological traits had evolved in response to metapopulation

connectedness. We quantified performance and dispersal in adult females that were raised for

one generation under common garden conditions (see further). We used five adults from each

patch in each experimental metapopulation, hence 45 F0 individuals per experimental

metapopulation (45×24 individuals in total), as the starting material for a generation of F1

females reared under identical conditions. We placed the five females coming from a same

patch together into the same common garden and we kept them separated from all the others.

Each F0 female was allowed to lay eggs for two days on a standardised bean leaf, and

offspring were subsequently raised under comparable densities (~ 2 individuals $\times$ cm^{-2}), at

30°C, a 16-8 L/D photoperiod and food *ad libitum*. This procedure enabled us to study traits

independent of plastic responses to the experienced conditions in the metapopulation. We

interpret, as others, any trait variation as genetic variation although intergenerational effects

cannot be ruled-out. Given the extent of the trait-quantification experiments, F1 offspring

were distributed across experiments to quantify (1) dispersal, (2) starvation resistance and (3)

reproductive performance. For reasons of feasibility, we chose to maximise trait

quantification across rather than within experimental metapopulations to ensure maximal

independent replication: in some cases, we then chose to pool together offspring originating

from different patches of the same metapopulation to insure a sufficiently high number of

individuals was used.

Dispersal Propensity

This test was performed at metapopulation level.

The propensity and the timing of dispersal were tested using a standardized setup of two-

patch systems, each composed of two 2.5 cm × 1.5 cm patches connected by an 8 cm long, 1

cm wide Parafilm bridge. On the starting patch, we placed females from F1 at densities

varying between 5 and 40 individuals, thereby generating a density gradient between 1.3 and

10.4 individuals/cm². The two patches were then connected and monitored for four days. Each

day, the number of individuals on the starting and the other leaf was recorded, along with the

number of individuals on the bridge at the time of the census. The target leaf to which

dispersal took place was refreshed after each count to prevent successfully dispersed females

from moving back to the starting patch. Freshly moulted F1 females (the dispersing sex)

reared for one generation under common garden conditions from the same experimental

metapopulation replicate were put together on the source patch for the dispersal trial, in order

to simulate a composition similar to the original metapopulation. Therefore, data could only

be analyzed in function of the metapopulation typology as this is the anticipated unit of

selection.

We ran in total 78 two-patches dispersal trials, for a grand total of 1330 monitored

individuals. We analyzed both the fraction of emigrants (individuals leaving the source patch)

and the fraction of immigrants (individuals successfully settling on the target patch). These

fractions were analyzed using generalized linear mixed model with a binomial error structure

and logit-link; metapopulation connectedness, presence/absence of randomization

(categorical) and density on the starting patch (continuous), as well as their two- and three-

ways interactions were considered as explanatory variables. The metapopulation replicate was

added as random effect to control for dependency in responses from individuals originating

from the same experimental metapopulation.

Finally we used the day at which each individual dispersed as a dependent variable in a linear

model, using again metapopulation connectedness, presence/absence of randomization

(categorical) and density on the starting patch (continuous covariate) as explanatory variables,

along with their two- and three-ways interactions. We assumed a Poisson distribution for

count data with low mean. The metapopulation replicate and dispersal trial replicate were here

added as random factors to control for interdependence of data because of shared evolutionary

history and experimental setup. Only individuals that performed a dispersal event were

included in this analysis so to render this analysis complementary to the emigration analysis

outlined higher. We complemented this analysis with a population-level test of the day at

which each replicate reached or passed the 50% of its final number of dispersers ("Time50"

from now on) and used this as a proxy for the dispersal timing of the population. As this

analysis yields complementary and similar insights as the individual-based test, we provide

these results in supplement (Supplementary Materials SI 2).

Starvation Resistance

This test was performed at metapopulation level.

One experimental arena was prepared for each metapopulation as well as one for the stock

population as control (total: 25 arenas). Each arena consisted of a 25 cm^2 square cut from a

black plastic sheet mounted on a wet cotton bed mounted into a plastic Petri plate and fixed in

position by wet paper strips: the strips partially overlapped with the plastic (~3mm) to form a

wet barrier and deter mites from falling into the cotton. For each experimental metapopulation

replicate, 5 1-2 day old F1 adult females from the common gardens were placed into an arena.

Individuals coming from different patches of the same metapopulation were pooled together

as to represent the composition of the original metapopulation. The five out of nine most

abundant local populations at the moment of the F0 individuals collection were selected to

collect the F1 females. The Petri plates were then stored at room temperature (~25°C).

Based on exploratory trials, we compared the survival percentage after 48h, as this time

appeared to be most discriminatory regarding the death/survival balance. Data were analyzed

by generalized linear mixed modelling using a binomial distribution, with the connectedness-

level and presence/absence of randomization as explanatory categorical variables. As we were

only able to have one experimental trial per experimental metapopulation replicate, no

random effects were added.

Reproductive performances

This test was performed at local population level.

Single fertilized females from the F1 generation reared under standardised conditions were

collected and placed individually on 2.5 cm × 1.5 cm bean leaf cuttings, mounted on a bed of

wet cotton and kept in place with paper strips. From each experimental metapopulation, 1-2

females were collected from each patch, for a grand total of 370 females across the 24

metapopulation replicates. The leaves were placed in a heated cabinet, kept at 30°C for ten

days with a 16-8 L/D photoperiod and moisturized daily to prevent desiccation of the leaves

and the escape of the mites through the cotton layer. Each female was monitored daily,

recording her status (dead or alive), as well as the number of eggs, larvae and juveniles

present on the patch. The number of individuals from these three age classes were

subsequently summed as a measure of reproductive performance. We ran a GLM assuming a

quasi-Poisson distribution for the count data, to correct for overdispersion (Cameron and

Trivedi, 1990). Reproductive performance at day 6 was used as the dependent variable to

avoid effects of overlapping generations. As reproductive performance was hypothesised to

evolve in relation to local conditions as well, we added patch location of the F0 females

(centre, corner or edge) as an independent categorical variable in addition to the earlier used

metapopulation-level connectedness and randomisation treatments. The experimental

metapopulation replicate was added as random effect to control for dependency by shared

evolutionary history.

 RESULTS

 Dispersal Propensity

Evolved emigration and immigration rate (success rate) were not dependent on the

connectedness or randomization treatment; only the starting density had a significant and

positive (see figure in Supplementary Materials SI 3) impact on the emigration rates (Table

1).

Individuals that evolved in the most disconnected metapopulation showed a delayed timing of

dispersal (day of departure) (Analysis of Deviance, Type III test: $\chi^2 =17.16$, P= 0.00018). The

same pattern was observed in the values of "Time50" at the population level (see

Supplementary Material SI 2).

Starvation resistance

The proportion of individuals surviving 48 hours of starvation only depended on the

connectedness experienced during experimental evolution, with highest survival rates for

individuals that evolved in the least connected metapopulation relative to the two other

treatments (Analysis of Deviance, Type III test: $\chi^2=11.89$, P=0.003; Fig. 3). Survival rates did

not differ between the randomization procedures (Analysis of Deviance, Type III test:

$\chi^2=0.12$, P=0.720) nor on the interaction between both (Analysis of Deviance, Type III test:

$\chi^2=0.85$, P=0.651).

Reproductive performance

The local connectedness of the patch the individuals originated from affected reproduction

significantly (Table 2, Fig. 4). In general, reproductive performance was overall the lowest in

offspring from individuals that were collected in the central patch (Posthoc Tukey test:

corner-center contrast: t-ratio = -2.895, P=0.0112; edge-center contrast: t-ratio = -3.073;

P=0.0065). Reproductive performance was not affected by the metapopulation connectedness

or the randomization procedure (see table 2).

DISCUSSION

Dispersal is a complex life history trait encompassing multiple, often tightly coupled stages.

To date, most evidence for dispersal evolution in metapopulations stems from studies on

emigration rate (Olivieri, Michalakis and Gouyon, 1995; Mathias, Kisdi and Olivieri, 2001;

Gyllenberg, Parvinen and Dieckmann, 2002; Ronce, 2007). However, simultaneous evolution

of traits related to the multiple dispersal stages, including transience and settlement, has been

theoretically demonstrated to lead to often complex dynamics which may or may not impact

the emigration rate (Travis *et al.*, 2013; Delgado *et al.*, 2014). Our work on experimental

evolution showed that changes in metapopulation connectedness altered the evolution of

dispersal timing and resistance against starvation, a possible dispersal cost, and not of

emigration rates *per se.* The evolution of dispersal timing has been theoretically addressed in

response to either dispersal at adulthood (breeding dispersal) or at birth (natal dispersal) (Johst

and Brandl, 1999; Hirota, 2004; Lakovic, Poethke and Hovestadt, 2015; Lakovic, Mitesser

and Hovestadt, 2017).

Here, we found breeding dispersal was delayed in the least connected metapopulations. This

can be mechanistically interpreted as a delayed density dependence as individuals delaying

dispersal are more likely to be engaged in local competitive interactions. In this sense,

delayed dispersal reflects density-dependent emigration strategies (Saether, Engen and Lande,

1999; Hovestadt and Poethke, 2006) with density thresholds evolving to higher levels in

disconnected metapopulations. Starvation resistance evolved in parallel with dispersal timing,

with individuals from the least connected metapopulation showing the largest starvation

resistance. Such an increased resistance on one hand reduced costs of local competition

following delayed dispersal. As the dispersal event in our experimental metapopulations was

costly, individuals may have adapted an "endurance-oriented" strategy in the least connected

metapopulations, as opposed to the "early leave" strategy of the more tightly connected

metapopulations. Alternatively, it may have evolved from higher transfer costs through the

hostile matrix: increased starvation resistance should lead to enhanced survival when

dispersing through a matrix void of food resources, and therefore compensates for the

prolonged exposure to nutritional stress (Matsumura and Miyatake, 2018). When integrated,

evolution of delayed dispersal and the costs of dispersal then release selection on the

emigration rate, equalizing connectivity at the metapopulation-level among the different

metapopulation despite variation in connectedness (Bélisle, 2005; Baguette and Van Dyck,

2007). We here thus illustrate the prevalence of expected but not yet documented eco-

evolutionary feedbacks in metapopulations (Bonte, Masier and Mortier, 2018).

The higher starvation resistance found in the least connected metapopulations can not be

explained by higher (local) population densities during the 6 months of the experiment, as no

significant difference was found in local dynamics or population size (S3, S4 and Masier, *in*

prep.).

While increased dispersal costs are tempting explanation of the increased starvation resistance

in the most disconnected metapopulations, it may also have evolved in response to the

metapopulation demography if local densities are on average higher in less connected

metapopulations, or if high densities and thus resource competition are more frequently

experienced. However, both at local and metapopulation level, no difference in the average

number of individuals per week was found between treatments over the 6 months of the

experiment, and neither were local fluctuations (alpha variability *sensu latu*) in densities

significantly different among the connectedness treatments (see supplementary material SI 4

and SI 5).

Building up energy reserves to survive long periods of food shortage is a costly process that

allocates resources, subtracting them from other processes. Starvation resistance is commonly

associated with bigger body mass, and usually has consequences on fertility, reproduction

timing and longevity (Tessier *et al.*, 1983; Hoffmann and Harshman, 1999; Rion and

Kawecki, 2007). However, we did not detect such trade-offs as reproductive performance

only diverged in response to the local patch type and not in relation to the connectedness of

the metapopulation in which they evolved (see also Supplementary Material SI 6).

The systematic lower reproductive performances in individuals from central patches in the

network is intriguing as (1) patches were identical in terms of quality and type throughout the

network, and (2) dispersal is maintained as an individual trait in all metapopulation types,

hence swamping putative local selection (Alzate *et al.*, 2017; Alzate, Etienne and Bonte,

2019).

The signature of local selection in the central patch is intriguing given the putative gene flow

within networks. Local selection can thus only be attributed to emerging feedbacks from the

metapopulation dynamics. Reduced reproductive performances are expected to evolve in

response to intense competition (Krüger, Liversidge and Lindström, 2002; Agrawal,

Underwood and Stinchcombe, 2004). Given the equal emigration rates across the different

levels of metapopulation connectedness, but differences in timing and starvation resistance at

the metapopulation level, we anticipate a similar rate of population decrease between patches

due to emigration and resources deprivation. Patches with higher local connectedness

however, will always receive more individuals than those with less connections, independent

of the metapopulation-level connectivity (see Box 1 in Supplementary Material). Although we

could not statistically detect systematically higher densities in the central patches, the

expected asymmetrical dispersal rates may nevertheless impose higher levels of competition

following local overshooting. At the same time, reduced individual growth rates could

compensate locally for an increased input of immigrants, thus explaining why the local

densities were found not significantly different.

Even more paradoxical, this signature was not destroyed by randomization. This implies that

the observed lower reproductive performance, as measured after one generation of common

garden, must originate from persisting intergenerational plasticity rather than genetic

evolution. Such an intergenerational plasticity enables populations to respond fast to

environmental change (Galloway, 2005; Drummond and Ancona, 2015), and has been

recorded in relation to dispersal-related traits before (Bitume *et al.*, 2011, 2014; Van Petegem

et al., 2015) as well as for diapause (Oku, Yano and Takafuji, 2003). The exact mechanisms

behind these epigenetic responses remain, however, unclear.

Habitat fragmentation is a central problem in conservation biology as it is expected to reduce

connectivity and eventually lead to species loss and disintegration of local food webs

(Thompson, Rayfield and Gonzalez, 2017). However, connectivity between patches can be

evolutionary rescued (Kendrick *et al.*, 2017). Given that evolution of dispersal-related traits

shifts cost-benefit balances towards a more beneficial ratio in metapopulations (Bonte *et al.*,

2012), these evolutionary dynamics adaptations will feedback on other traits (e.g. different

allocation of energy reserves to reproduction) as well and thereby impact the ecological

processes (Bonte, Masier and Mortier, 2018).

Such an integrated view on dispersal, life history evolution and costs shifts in response to

habitat fragmentation is lacking (Cheptou *et al.*, 2017) but needed to advance the predictive

ecology of species and ecosystems, especially in the fast-changing world of today (Urban *et*

al., 2016). While theory focuses, for reasons of tractability, on simple dynamics (Govaert *et*

al., 2018), we are able to demonstrate more realistic multivariate life history evolution caused

by altered habitat connectedness. We demonstrate that these evolutionary dynamics do not

follow simple theoretical predictions based on dispersal alone. One would expect lower

dispersal rates between local populations with reduced connectedness, possibly coupled with

a higher success rate as dispersing individuals are highly specialized in finding and reaching

new suitable patches in the matrix (Bonte, Hovestadt and Poethke, 2010; Brisson, 2010).

Traits other than dispersal are jointly evolving as well in response to both local- and

metapopulation-level selection pressures, thereby causing potentially unpredictable changes in

population dynamics.

Local and metapopulation-level dynamics interact with each other and influence the

adaptation of the individuals at both levels, with ecological population dynamics influencing

evolutionary dynamics and the other way around (Hanski, 2012). Stress resistance, for

instance, which we found to evolve in response to habitat fragmentation, can potentially turn

into pre-adaptation to other kind of stressors, i.e. suboptimal host plants or novel habitats and

enhance persistence under environmental change (Jenkins, Chaisson and Matin, 1990; De

Roissart *et al.*, 2016). Moreover, we show that evolutionary dynamics resulting from

fragmentation also include epigenetic inheritance or intergenerational plasticity. The reduced

intrinsic population growth of the most locally connected population was pertinent within the

treatments where any local adaptation and genetic structure had been removed. As these

responses are usually very fast (Herman *et al.*, 2014), their role should not be underestimated

and deserves more in-depth study. Lastly, despite the different levels of connectedness

between patches, we found no differences in net dispersal rates after four days, meaning that

the connectivity and the gene flow between local populations were basically maintained.

This decoupling of habitat connectedness and population connectivity could impact

predictions traditionally derived from theoretical ecological models, that assume a direct and

positive link between global dispersal and habitat connectedness (Legrand *et al.*, 2017). The

evolution of often-overlooked traits like dispersal timing or individual growth rates rather

than dispersal rates themselves could then act as a rescue mechanism for fragmented

populations through their impact on metapopulation variability, local stabilizing and spatial

synchronizing effects (Wang, Haegeman and Loreau, 2015).

Bibliography

Agrawal, A. A., Underwood, N. and Stinchcombe, J. R. (2004) 'Intrspecific variation in the strength of density dependence', *Ecological Entomology*, 29, pp. 521–526.

Aguilar, R. *et al.* (2006) 'Plant reproductive susceptibility to habitat fragmentation: Review and synthesis through a meta-analysis', *Ecology Letters*, 9(8), pp. 968–980. doi: 10.1111/j.1461-0248.2006.00927.x.

Alzate, A. *et al.* (2017) 'Interspecific competition counteracts negative effects of dispersal on adaptation of an arthropod herbivore to a new host', *Journal of Evolutionary Biology*, 30(11), pp. 1966–1977. doi: 10.1111/jeb.13123.

Alzate, A., Etienne, R. S. and Bonte, D. (2019) 'Experimental island biogeography demonstrates the importance of island size and dispersal for the adaptation to novel habitats', *Global Ecology and Biogeography*, 28(2), pp. 238–247. doi: 10.1111/geb.12846.

Amarasekare, P. (2008) 'Spatial Dynamics of Foodwebs', *Annual Review of Ecology, Evolution, and Systematics*, 39(1), pp. 479–500. doi: 10.1146/annurev.ecolsys.39.110707.173434.

Andersen, L. W., Fog, K. and Damgaard, C. (2004) 'Habitat fragmentation causes bottlenecks and inbreeding in the European tree frog (Hyla arborea)', *Proceedings of the Royal Society B: Biological Sciences*, 271(1545), pp. 1293–1302. doi: 10.1098/rspb.2004.2720.

Baguette, M. and Van Dyck, H. (2007) 'Landscape connectivity and animal behavior: Functional grain as a key determinant for dispersal', *Landscape Ecology*, 22(8), pp. 1117–1129. doi: 10.1007/s10980-007-9108-4.

Baguette, M. and Schtickzelle, N. (2003) 'Local population dynamics are important to the conservation of metapopulations in highly fragmented landscapes', *Journal of Applied Ecology*, 40(2), pp. 404–412. doi: 10.1046/j.1365-2664.2003.00791.x.

Bélisle, M. (2005) 'Measuring Landscape Connectivity: the Challenge of Behavioral Landscape Ecology', *Ecology*, 86(8), pp. 1988–1995. doi: 10.1890/04-0923.

Bell, J. R. *et al.* (2005) 'Ballooning dispersal using silk: world fauna, phylogenies, genetics and models', *Bulletin of Entomological Research*, 95(02), pp. 69–114. doi: 10.1079/BER2004350.

Bonte, D. *et al.* (2003) 'Low propensity for aerial dispersal in specialist spiders from fragmented landscapes', *Proceedings of the Royal Society B: Biological Sciences*, 270(1524), pp. 1601–1607. doi:

10.1098/rspb.2003.2432.

Bonte, D. *et al.* (2012) 'Costs of dispersal', *Biological Reviews*, 87(2), pp. 290–312. doi: 10.1111/j.1469-
185X.2011.00201.x.

Bonte, D. and Dahirel, M. (2017) 'Dispersal: a central and independent trait in life history', *Oikos*, 126(4), pp.
472–479. doi: 10.1111/oik.03801.

Bonte, D., Hovestadt, T. and Poethke, H. J. (2010) 'Evolution of dispersal polymorphism and local adaptation of
dispersal distance in spatially structured landscapes', *Oikos*, 119(3), pp. 560–566. doi: 10.1111/j.1600-
0706.2009.17943.x.

Bonte, D., Masier, S. and Mortier, F. (2018) 'Eco-evolutionary feedbacks following changes in spatial
connectedness', *Current Opinion in Insect Science*. Elsevier Inc, 29, pp. 64–70. doi: 10.1016/j.cois.2018.06.003.

Bowler, D. E. and Benton, T. G. (2005) 'Causes and consequences of animal dispersal strategies: Relating
individual behaviour to spatial dynamics', *Biological Reviews of the Cambridge Philosophical Society*, 80(2),
pp. 205–225. doi: 10.1017/S1464793104006645.

Brisson, J. A. (2010) 'Aphid wing dimorphisms: Linking environmental and genetic control of trait variation',
Philosophical Transactions of the Royal Society B: Biological Sciences, 365(1540), pp. 605–616. doi:
10.1098/rstb.2009.0255.

Calabrese, J. M. and Fagan, W. F. (2004) 'A comparison-shopper's guide to connectivity metrics', *Frontiers in
Ecology and the Environment*, 2(10), pp. 529–536. doi: 10.1890/1540-
9295(2004)002[0529:ACGTCM]2.0.CO;2.

Cameron, A. C. and Trivedi, P. K. (1990) 'Regression-based tests for overdispersion in the Poisson model',
Journal of Econometrics, 46(3), pp. 347–364. doi: 10.1016/0304-4076(90)90014-K.

Cañete, J. I. *et al.* (2007) 'Planktonic drifting dispersal of early juvenile Trochita calyptraeaformis Born 1778
[Gastropoda: Calyptraeidae]', *Journal of Experimental Marine Biology and Ecology*, 346(1–2), pp. 1–7. doi:
10.1016/j.jembe.2007.01.013.

Cheptou, P.-O. *et al.* (2017) 'Adaptation to fragmentation: evolutionary dynamics driven by human influences',
Philosophical Transactions of the Royal Society B: Biological Sciences, 372(1712), p. 20160037. doi:
10.1098/rstb.2016.0037.

Clobert, J. *et al.* (2009) 'Informed dispersal, heterogeneity in animal dispersal syndromes and the dynamics of

spatially structured populations', *Ecology Letters*, 12(3), pp. 197–209. doi: 10.1111/j.1461-0248.2008.01267.x.

Clobert, J. *et al.* (2012) *Dispersal ecology and evolution*.

Collinge, S. K. (1998) 'Spatial arrangement of habitat patches and corridors: Clues from ecological field

experiments', *Landscape and Urban Planning*, 42(2–4), pp. 157–168. doi: 10.1016/S0169-2046(98)00085-1.

Crooks, K. R. (2002) 'Relative Sensitivites of Mammalian Carnivores to Habitat Fragmentation', *Conservation*

Biology, 16(2), pp. 488–502. doi: 10.1046/j.1523-1739.2002.00386.x.

Cushman, S. A. (2006) 'Effects of habitat loss and fragmentation on amphibians: A review and prospectus',

Biological Conservation, 128(2), pp. 231–240. doi: 10.1016/j.biocon.2005.09.031.

Delgado, M. M. *et al.* (2014) 'Prospecting and dispersal: their eco-evolutionary dynamics and implications for

population patterns', *Proceedings of the Royal Society B: Biological Sciences*, 281(1778), pp. 20132851–

20132851. doi: 10.1098/rspb.2013.2851.

Derr, J. A., Aldent, B. and Dinglet, H. (1981) 'Insect Life Histories in Relation to Migration, Body Size, and

Host Plant Array: A Comparative Study of Dysdercus', *Journal of Animal Ecology*, 50(1), pp. 181–193.

Dey, S., Goswami, B. and Joshi, A. (2014) 'Effects of symmetric and asymmetric dispersal on the dynamics of

heterogeneous metapopulations: Two-patch systems revisited', *Journal of Theoretical Biology*. Elsevier, 345, pp.

52–60. doi: 10.1016/j.jtbi.2013.12.005.

Dobson, A. *et al.* (2006) 'Habitat loss, trophic collapse, and the decline of ecosystem services', *Ecology*, 87(8),

pp. 1915–1924. doi: 10.1890/0012-9658(2006)87[1915:HLTCAT]2.0.CO;2.

Drummond, H. and Ancona, S. (2015) 'Observational field studies reveal wild birds responding to early-life

stresses with resilience, plasticity, and intergenerational effects', *The Auk*, 132(3), pp. 563–576. doi:

10.1642/AUK-14-244.1.

Fahrig, L. (2003) 'Effects of Habitat Fragmentation on Biodiversity', *Annual Review of Ecology, Evolution, and*

Systematics, 34(1), pp. 487–515. doi: 10.1146/annurev.ecolsys.34.011802.132419.

Fenoglio, M. S. *et al.* (2010) 'Plant patch structure modifies parasitoid assemblage richness of a specialist

herbivore', *Ecological Entomology*, 35(5), pp. 594–601. doi: 10.1111/j.1365-2311.2010.01218.x.

Ferrari, J. R., Lookingbill, T. R. and Neel, M. C. (2007) 'Two measures of landscape-graph connectivity:

Assessment across gradients in area and configuration', *Landscape Ecology*, 22(9), pp. 1315–1323. doi:

10.1007/s10980-007-9121-7.

Forman, R. T. T. (2000) 'Estimate of the area affected ecologically by the road system in the United States',

Conservation Biology, 14(1), pp. 31–35. doi: 10.1046/j.1523-1739.2000.99299.x.

Fronhofer, E. A. *et al.* (2014) 'Spatially correlated extinctions select for less emigration but larger dispersal

distances in the spider mite tetranychus urticae', *Evolution*, 68(6), pp. 1838–1844. doi: 10.1111/evo.12339.

Fuller, M. R., Doyle, M. W. and Strayer, D. L. (2015) 'Causes and consequences of habitat fragmentation in

river networks', *Annals of the New York Academy of Sciences*, 1355(1), pp. 31–51. doi: 10.1111/nyas.12853.

Galloway, L. F. (2005) 'Maternal effects provide phenotypic adaptation to local environmental conditions', *New*

Phytologist, 166(1), pp. 93–100. doi: 10.1111/j.1469-8137.2004.01314.x.

Gibbs, J. P. and Stanton, E. J. (2001) 'Habitat fragmentation and arthropod community change: Carrion beetles,

phoretic   mites,   and   flies',   *Ecological   Applications*,   11(1),   pp.   79–85.   doi: 10.1890/1051-

0761(2001)011[0079:HFAACC]2.0.CO;2.

Govaert, L. *et al.* (no date) 'Eco-evolutionary feedbacks — theoretical models and perspectives'.

Gyllenberg, M., Parvinen, K. and Dieckmann, U. (2002) 'Evolutionary suicide and evolution of dispersal in

structured metapopulations', *Journal of Mathematical Biology*, 45(2), pp. 79–105. doi: 10.1007/s002850200151.

Hance, T. and Impe, G. Van (1999) 'The influence of initial age structure on predator – prey interaction',

Ecological Modelling, 114, pp. 195–211. doi: 10.1016/S0304-3800(98)00148-3.

Hanski, I. (2012) 'Eco-evolutionary dynamics in a changing world', *Annals of the New York Academy of*

Sciences, 1249(1), pp. 1–17. doi: 10.1111/j.1749-6632.2011.06419.x.

Hansson, B., Bensch, S. and Hasselquist, D. (2004) 'Lifetime Fitness of Short-and Long-Distance Dispersing

Great Reed Warblers', *Evolution*, 58(11), pp. 2546–2557. doi: 10.1111/j.0014-3820.2004.tb00883.x.

Hazell, S. P. *et al.* (2005) 'Competition and dispersal in the pea aphid: Clonal variation and correlations across

traits', *Ecological Entomology*, 30(3), pp. 293–298. doi: 10.1111/j.0307-6946.2005.00703.x.

Heilman, G. E. *et al.* (2002) 'Forest fragmentation of the conterminous United States: Assessing forest intactness

through road density and spatial characteristics.', *BioSience*, 52(5), pp. 411–422.

Herman, J. J. *et al.* (2014) 'How stable "should" epigenetic modifications be? Insights from adaptive plasticity

and bet hedging', *Evolution*, 68(3), pp. 632–643. doi: 10.1111/evo.12324.

Hirota, T. (2004) 'The evolution of sex-biased dispersal by pre-dispersal copulation and fluctuating

environment', *Journal of Animal Ecology*, 73(6), pp. 1115–1120. doi: 10.1111/j.0021-8790.2004.00878.x.

Hoffmann, A. A. and Harshman, L. G. (1999) 'Desiccation and starvation resistance in Drosophila: Patterns of

variation at the species, population and intrapopulation levels', *Heredity*, 83(6), pp. 637–643. doi:

10.1046/j.1365-2540.1999.00649.x.

Hovestadt, T. and Poethke, H. J. (2006) 'The control of emigration and its consequences for the survival of

populations', *Ecological Modelling*, 190(3–4), pp. 443–453. doi: 10.1016/j.ecolmodel.2005.03.023.

Jenkins, D. E., Chaisson, S. A. and Matin, A. (1990) 'Starvation-induced cross-protection against osmotic

challenge in *Escherichia coli*', *J. Bacteriol.*, 172(5), pp. 2779–2781. doi: 0021-9193/88/093910-05$02.00/0.

Johst, K. and Brandl, R. (1999) 'Natal versus breeding dispersal: Evolution in a model system', *Evolutionary*

Ecology Research, 1(8), pp. 911–921.

Kendrick, G. A. *et al.* (2017) 'Demographic and genetic connectivity: The role and consequences of

reproduction, dispersal and recruitment in seagrasses', *Biological Reviews*, 92(2), pp. 921–938. doi:

10.1111/brv.12261.

Keyghobadi, N. (2007) 'The genetic implications of habitat fragmentation for animals', *Canadian Journal of*

Zoology, 85(10), pp. 1049–1064. doi: 10.1139/Z07-095.

Kondoh, M. (2003) 'Habitat fragmentation resulting in overgrazing by herbivores', *Journal of Theoretical*

Biology, 225(4), pp. 453–460. doi: 10.1016/S0022-5193(03)00279-0.

Korb, J. and Linsenmair, K. E. (2002) 'Evaluation of predation risk in the collectively foraging termite

Macrotermes bellicosus', *Insectes Sociaux*, 49(3), pp. 264–269. doi: 10.1007/s00040-002-8312-0.

Krüger, O., Liversidge, R. and Lindström, J. (2002) 'Statistical modelling of the population dynamics of a raptor

community in a semi-desert enviroment', *Journal of Animal Ecology*, 71, pp. 603–613.

Lakovic, M., Mitesser, O. and Hovestadt, T. (2017) 'Mating timing, dispersal and local adaptation in patchy

environments', *Oikos*, 126(12), pp. 1804–1814. doi: 10.1111/oik.04369.

Lakovic, M., Poethke, H. J. and Hovestadt, T. (2015) 'Dispersal timing: Emigration of insects living in patchy

environments', *PLoS ONE*, 10(7), pp. 1–15. doi: 10.1371/journal.pone.0128672.

Van Langevelde, F. (2000) 'Scale of habitat connectivity and colonization in fragmented nuthatch populations',

Ecography, 23(5), pp. 614–622. doi: 10.1111/j.1600-0587.2000.tb00180.x.

Legrand, D. *et al.* (2017) 'Eco-evolutionary dynamics in fragmented landscapes', *Ecography*, 40(1), pp. 9–25.

doi: 10.1111/ecog.02537.

Liao, J. *et al.* (2017) 'Robustness of metacommunities with omnivory to habitat destruction: disentangling patch

fragmentation from patch loss', *Ecology*, 98(6), pp. 1631–1639. doi: 10.1002/ecy.1830.

van der Linde, K. and Sevenster, J. G. (2006) 'Local adaptation of development time and starvation resistance in

eight Drosophila species of the Phillipines', *Biological Journal of the Linnean Society*, 87(87), pp. 115–125. doi:

10.1111/j.1095-8312.2006.00560.x.

Macke, E. *et al.* (2011) 'Experimental Evolution of Reduced Sex Ratio Adjustment Under Local Mate

Competition', *Science*, 334(November), pp. 1127–1130. doi: https://doi.org/10.1126/science.1212177.

Mathias, A., Kisdi, É. and Olivieri, I. (2001) 'Divergent Evolution of Dispersal in a Heterogeneous Landscape',

Evolution, 55(2), pp. 246–259. doi: http://dx.doi.org/10.1554/0014-3820(2001)055[0246:DEODIA]2.0.CO;2.

Matsumura, K. and Miyatake, T. (2018) 'Costs of walking: differences in egg size and starvation resistance of

females between strains of the red flour beetle (Tribolium castaneum) artificially selected for walking ability',

Journal of Evolutionary Biology, pp. 0–3. doi: 10.1111/jeb.13356.

McClain, W. R. (1995) 'Effects of Population Density and Feeding Rate on Growth and Feed Consumption of

Red Swamp Crawfish Procam barns clarkii', *Journal Of The World Aquaculture Society*, 26(1), pp. 14–23. doi:

10.1111/j.1749-7345.1995.tb00204.x.

Melián, C. J. and Bascompte, J. (2002) 'Food web structure and habitat loss', *Ecology Letters*, 5(1), pp. 37–46.

doi: 10.1046/j.1461-0248.2002.00280.x.

Oku, K., Yano, S. and Takafuji, a (2003) 'Different maternal effects on diapause induction of tetranychid mites,

Tetranychus urticae and T. kanzawai (Acari: Tetranychidae)', *Applied Entomology and Zoology*, 38(2), pp. 267–

270. doi: 10.1303/aez.2003.267.

Olivieri, I., Michalakis, Y. and Gouyon, P. H. (1995) 'Metapopulation Genetics and the Evolution of Dispersal',

American Naturalist, 146(2), pp. 202–228. doi: 10.1086/285795.

Van Petegem, K. H. P. *et al.* (2015) 'Empirically simulated spatial sorting points at fast epigenetic changes in

dispersal behaviour', *Evolutionary Ecology*, 29(2), pp. 299–310. doi: 10.1007/s10682-015-9756-9.

Van Petegem, K. *et al.* (2018) 'Kin competition accelerates experimental range expansion in an arthropod

herbivore', *Ecology Letters*, 21(2), pp. 225–234. doi: 10.1111/ele.12887.

Poethke, H. J., Dytham, C. and Hovestadt, T. (2011) 'A Metapopulation Paradox: Partial Improvement of

Habitat May Reduce Metapopulation Persistence', *The American Naturalist*, 177(6), pp. 792–799. doi:

10.1086/659995.

Rion, S. and Kawecki, T. J. (2007) 'Evolutionary biology of starvation resistance: What we have learned from

Drosophila', *Journal of Evolutionary Biology*, 20(5), pp. 1655–1664. doi: 10.1111/j.1420-9101.2007.01405.x.

De Roissart, A. *et al.* (2016) 'Life-history evolution in response to changes in metapopulation structure in an

arthropod herbivore', *Functional Ecology*, 30(8), pp. 1408–1417. doi: 10.1111/1365-2435.12612.

De Roissart, A., Wang, S. and Bonte, D. (2015) 'Spatial and spatiotemporal variation in metapopulation

structure affects population dynamics in a passively dispersing arthropod', *Journal of Animal Ecology*, 84(6), pp.

1565–1574. doi: 10.1111/1365-2656.12400.

Ronce, O. (2007) 'How Does It Feel to Be Like a Rolling Stone? Ten Questions About Dispersal Evolution',

Annual Review of Ecology, Evolution, and Systematics, 38(1), pp. 231–253. doi:

10.1146/annurev.ecolsys.38.091206.095611.

Rossetti, M. R. *et al.* (2014) 'Not all in the same boat: trends and mechanisms in herbivory responses to forest

fragmentation differ among insect guilds', *Arthropod-Plant Interactions*, 8(6), pp. 593–603. doi:

10.1007/s11829-014-9342-z.

Rudnick, D. *et al.* (2012) 'The Role of Landscape Connectivity in Planning and Implementating Conservation

and Restoration Priorities', *Issue in Ecology*, 13(16), pp. 1–16. doi: 10.1371/JOURNAL.PONE.0110984.

Saether, B.-E., Engen, S. and Lande, R. (1999) 'Finite metapopulation models with density-dependent migration

and stochastic local dynamics', *Proceedings of the Royal Society B: Biological Sciences*, 266(1415), pp. 113–

118. doi: 10.1098/rspb.1999.0610.

Stow, A. J. *et al.* (2001) 'The impact of fragmentation on dispersalof Cunningham's sink (Egernia unninghami):

evidence from allelic and genotypic analyses of microsatellites', *Molecular Ecology*, 10, pp. 867–878.

Taylor, P. D. *et al.* (1993) 'Connectivity Is a Vital Element of Landscape Structure', *Oikos*, 68(3), pp. 571–573.

doi: 10.1111/j.2007.0906-7590.05378.x.

Tessier, A. J. *et al.* (1983) 'Starvation in Daphnia: Energy Reserves and Reproductive Allocation', 28(4), pp.

667–676.

Thompson, P. L., Rayfield, B. and Gonzalez, A. (2017) 'Loss of habitat and connectivity erodes species

diversity, ecosystem functioning, and stability in metacommunity networks', *Ecography*, 40(1), pp. 98–108. doi:

10.1111/ecog.02558.

Tischendorf, L. and Fahrig, L. (2000) 'On the usage and measurement of landscape connectivity', *Oikos*, 90(1),

pp. 7–19. doi: 10.1034/j.1600-0706.2000.900102.x.

Tischendorf, L. and Fahrig, L. (2001) 'On the use of connectivity measures in spatial ecology. A reply', *Oikos*,

95(1), pp. 152–155. doi: 10.1034/j.1600-0706.2001.950117.x.

Travis, J. M. J. *et al.* (2013) 'Dispersal and species' responses to climate change', *Oikos*, 122(11), pp. 1532–

1540. doi: 10.1111/j.1600-0706.2013.00399.x.

Urban, M. C. *et al.* (2016) 'Improving the forecast for biodiversity under climate change', *Science*, 353(6304).

doi: 10.1126/science.aad8466.

Urban, M. C. and Skelly, D. K. (2006) 'Evolving metacomunities: toward an evolutionary perspective on

metacommunities', *Ecology*, 87(7), pp. 1616–1626. doi: 10.1890/07-1861.1.

Valladares, G., Cagnolo, L. and Salvo, A. (2012) 'Forest fragmentation leads to food web contraction', *Oikos*,

121(2), pp. 299–305. doi: 10.1111/j.1600-0706.2011.19671.x.

Vessby, K. and Wiktelius, S. (2003) 'The influence of slope aspect and soil type on immigration and emergence

of some northern temperate dung beetles', *Pedobiologia*, 47(1), pp. 39–51. doi: 10.1078/0031-4056-00168.

Wang, S., Haegeman, B. and Loreau, M. (2015) 'Dispersal and metapopulation stability', *PeerJ*, 3, pp. 1–16.

doi: 10.7717/peerj.1295.

Wang, X., Blanchet, F. G. and Koper, N. (2014) 'Measuring habitat fragmentation: An evaluation of landscape

pattern metrics', *Methods in Ecology and Evolution*, 5(7), pp. 634–646. doi: 10.1111/2041-210X.12198.

Figures

Fig. 1: schematic representation of the long-term (6 months) experimental setups. Nine identical patches were arranged in a 3-by-3 grid and connected using Parafilm bridges. The length of the bridge (X cm in the picture) was dependent on the treatment; the diagonal bridges varied depending on the value of X.

Figure 2: the effect of connectedness on the individual day of dispersal in the different treatments (means ± sd).

Fig. 3: proportion of alive individuals (means ± sd) after 48h depending on the connectedness of the source metapopulation.

Fig. 4a (upper panel): Number of alive offspring after 6 days (mean ± sd), for each combination of fragmentation treatment (colors) and relative position of the patch of origin of the individuals (before common garden period). Fig. 4b (lower panel): pairwise comparisons between offspring originating from patches with a different position within the metapopulation (with confidence intervals at 95%).

626 **Tables**

Dependent variable	Explanatory variable	df	χ^2	Pr(>χ^2)
Emigration rate	Connectedness	2	1.5251	0.6765
	Randomization	1	0.0172	0.8957
	Starting Density	1	7.96	0.0048 (**)
	Connectedness * Randomization	2	0.1978	0.9058
	Connectedness * Starting Density	2	2.0713	0.3549
	Randomization * Starting Density	1	0.3847	0.5351
	Connectedness * Randomization * Starting Density	2	3.5738	0.1675
Immigration rate	Connectedness	2	0.6346	0.8885
	Randomization	1	0.5902	0.4423
	Starting Density	1	0.3421	0.5586
	Connectedness * Randomization	2	0.6865	0.7095
	Connectedness * Starting Density	2	1.3771	0.5023
	Randomization * Starting Density	1	0.0099	0.9208
	Connectedness * Randomization * Starting Density	2	1.1632	0.5590
Individual dispersal day	Connectedness	2	17.1631	0.0001875 (***)
	Randomization	1	1.7121	0.1907190
	Starting Density	1	0.0001	0.9916746
	Connectedness * Randomization	2	7.0688	0.0291764 (*)
	Connectedness * Starting Density	2	0.7331	0.6931065
	Randomization * Starting Density	1	0.5802	0.4462486
	Connectedness * Randomization * Starting Density	2	4.4365	0.1087995

Table 1: Full Model Analysis of Deviance table (Type III Wald χ^2-tests) for emigration and immigration rate and dispersal timing in relation to the connectedness, randomization of the metapopulation in which individuals evolved and density in the dispersal trial.

Explanatory variable	df	χ^2	Pr($>\chi^2$)
Position	9.6736	2	0.007933(**)
Connectedness	3.2587	2	0.196056
Randomization	0.4209	1	0.516487
Position * Connectedness	1.6079	4	0.806379
Position * Randomization	0.8291	2	0.660647
Connectedness * Randomization	3.7884	2	0.150437
Position * Connectedness * Randomization	1.1641	4	0.883968

Table 2: Full Model Analysis of Deviance table (Wald χ^2 test) for intrinsic reproductive performances as determined by the number of offspring produced by a single female after 6 days.