Mammalian SWI/SNF collaborates with a polycomb-associated protein to regulate male

germ line transcription in the mouse

Debashish U. Menon, Yoichiro Shibata, Weipeng Mu and Terry Magnuson

Department of Genetics, and Lineberger Comprehensive Cancer Center, The University of North

Carolina at Chapel Hill, Chapel Hill, North Carolina, NC 27599-7264, USA

*Correspondence: trm4@med.unc.edu, Tel 919-962-1319, Fax 919-843-4682

Abstract:

A deficiency in BRG1, the catalytic subunit of the SWI/SNF chromatin remodeling complex,

results in a meiotic arrest during spermatogenesis. Here, we explore the causative mechanisms.

BRG1 is preferentially enriched at active promoters of genes essential for spermatogonial

pluripotency and meiosis. In contrast, BRG1 is also associated with the repression of somatic

genes. Chromatin accessibility at these target promoters is dependent upon BRG1. These

results favor a model where BRG1 coordinates spermatogenic transcription to ensure meiotic

progression. In spermatocytes, BRG1 interacts with SCML2, a testes specific PRC1 factor that

is associated with the repression of somatic genes. We present evidence to suggest that BRG1

and SCML2 concordantly regulate genes during meiosis. Furthermore, BRG1 is required for the

proper localization of SCML2 and its associated deubiquitinase, USP7, to the sex chromosomes

during pachynema. SCML2 associated, mono ubiquitination of histone H2A lysine 119

(H2AK119ub1) and acetylation of histone lysine 27 (H3K27ac) are elevated in $Brg1^{cKO}$ testes.

Coincidentally, the PRC1 ubiquitin ligase, RNF2 is activated while a histone H2A/H2B

deubiquitinase, USP3 is repressed. Thus, BRG1 impacts the male epigenome by influencing the

localization and expression of epigenetic modifiers. This mechanism highlights a novel

paradigm of co-operativity between SWI/SNF and PRC1.

Keywords: SWI/SNF chromatin remodeling,BRG1, transcriptional regulation, SCML2

35   **Summary statement**

36   BRG1, a catalytic subunit of SWI/SNF chromatin remodeling complex, interacts with SCML2

37   (Sex comb on midleg-like 2), a member of the polycomb repressive 1 (PRC1) complex, to

38   regulate transcription during spermatogenesis. This represents a novel paradigm of co-

39   operativity between SWI/SNF and PRC1 that is critical to gametogenesis.

Introduction:

Spermatogenesis is a developmental cascade in which genetic information is passed on from mitotic precursors to meiotically derived haploid gametes. This process is particularly sensitive to the activity of several epigenetic regulators, known to influence meiotic recombination. Examples include the role of the meiosis specific H3K4 methyltransferase PRDM9 (PR domain zinc finger protein9) in double stranded break (DSB) formation and the roles of polycomb repressive complex2 (PRC2) and the H3K9 methyl transferases, EHMT2- Euchromatic histone lysine N-methyltransferase 2 (G9a); Suppressor of variegation 3-9 1 or 2 (SUV39H1/H2) in homolog pairing (synapsis) (Brick et al., 2012; Diagouraga et al., 2018; Hayashi et al., 2005; Mu et al., 2014; Tachibana et al., 2007; Takada et al., 2011). In addition to meiotic recombination, the temporal regulation of transcription is also essential for normal spermatogenesis. Defects in male fertility have been attributed to the mis-regulation of genes governed by polycomb repressive complexes, PRC1 and PRC2 (Hasegawa et al., 2015; Mu et al., 2014). Put together, these studies illustrate the importance of proteins that catalyze or respond to specific histone modifications in the germ line. Another class of proteins that modulate chromatin accessibility, have also been shown to be essential for spermatogenesis. These include the INO80, SWI/SNF and CHD5 family of ATP dependent nucleosome remodeling complexes. Their impact spans the gamut of spermatogenesis, beginning at meiosis and ending at spermiogenesis (Kim et al., 2012; Li et al., 2014; Serber et al., 2015).

Our lab previously reported the role of BRG1 (SMARCA4 – SWI/SNF catalytic subunit) during meiosis-I in males (Kim et al., 2012). Normally, meiosis-I is initiated by the formation of DSBs induced by the SPO11 endonuclease at leptonema, following which, an orderly process of chromosomal synapsis during zygonema, crossover formation at pachynema and asynapsis at diplonema ensues, culminating in DNA repair and recombination (reviewed in Handel and Schimenti, 2010). The germ line depletion of BRG1 results in a pachytene arrest. Briefly, mutant spermatocytes display unrepaired DNA DSBs evidenced by persistent γH2Ax, chromosomal asynapsis and reduced MLH1 foci, a marker of crossovers (Wang et al. 2012). Additionally, mutant pachytene spermatocytes also display enhanced levels of repressive chromatin, which could either be a physiological consequence of apoptotic initiation or a product of perturbed epigenetic processes. SWI/SNF influences chromatin accessibility by either sliding or evicting

nucleosomes (Clapier et al., 2017). It is therefore plausible that the meiotic defects seen in the

Brg1cKO testes might be a consequence of altered chromatin structure, which may either perturb

transcription or DNA repair. Both processes are known to be influenced by SWI/SNF.

SWI/SNF is associated with both gene activation and repression. In mouse embryonic fibroblasts,

BRG1 and the core SWI/SNF member, SNF5 are known to co-ordinate nucleosome occupancy

at promoters to achieve transcriptional regulation (Tolstorukov et al., 2013). The transcriptional

outcome is often dictated by the subunit composition of the complex, which in turn can influence

its genome-wide association and interactions with other gene regulators (Euskirchen et al., 2011;

Raab et al., 2015; Raab et al., 2017). SWI/SNF complexes are biochemically heterogeneous and

are broadly categorized into the *Brahma* Associated Factor (BAF) and *polybromo*-BAF (PBAF)

complex, based on the identity of mutually exclusive DNA binding subunits and associated

ATPase (BRG1/BRM) (Masliah-Planchon et al., 2015). This SWI/SNF heterogeneity is a theme

that features frequently in the regulation of tissue and cell type specific transcriptional programs

that impact critical processes such as embryonic stem (ES) cell pluripotency and differentiation,

neuronal and cardiac cell fate specification (Ho and Crabtree, 2010). SWI/SNF subunits also

constitute some of the most frequently mutated genes in various cancers (Masliah-Planchon et

al., 2015). Thus, SWI/SNF directed gene regulation plays a critical role in development and

disease.

Apart from gene regulation, SWI/SNF has also been implicated in several DNA repair

mechanisms. In Yeast and cell culture models, SWI/SNF is recruited to sites of DNA damage to

promote accessibility and stimulate γH2Ax, a key component of the DNA damage response

(DDR) signaling pathway (Kwon et al., 2015; Lee et al., 2010; Ogiwara et al., 2011). Other

activities include the recruitment of homologous recombination (HR) and non-homologous end

joining (NHEJ) repair factors (Ogiwara et al., 2011; Qi et al., 2014; Watanabe et al., 2014), the

transcriptional silencing of regions adjacent to DNA DSB (Kakarougkas et al., 2013). Unlike its

role in mitotic cells, BRG1 is dispensable for γH2Ax formation during meiosis but has been

reported to affect the distribution of DDR factors like RAD51 (DNA recombinase) and RPA

(Replication protein A1) (Kim et al., 2012).

In this study we present evidence to show that BRG1 influences spermatogenesis by

coordinating germ line transcription. BRG1 activates genes that are essential for maintaining

spermatogonial pluripotency and meiotic progression and represses somatic genes. Our data

suggest that gene repression is achieved through an interaction with SCML2 (Sex comb on

midleg-like 2), a known testis specific PRC1(Polycomb Repressive Complex 1) member known

to regulate somatic genes (Luo et al., 2015). We show that that loss of BRG1 results in the

activation of somatic genes that are normally repressed by SCML2. Additionally, mutant

pachytene spermatocytes display the mis-localization of SCML2 and its interacting partner,

USP7 (Ubiquitin Specific Peptidase 7) from the sex body (sub nuclear γH2Ax enriched

compartment containing the sex chromosomes), indicating a failure in recruitment. Furthermore,

histone modifications associated with SCML2, such as the repressive mono ubiquitination of

histone H2A lysine 119 (H2AK119ub1) and activating acetylation of histone lysine 27 (H3K27ac)

are perturbed in *Brg1^{cKO}* testes. Coincidentally, BRG1 also activates the expression of the H2A

ubiquitin ligase, RNF2 (Ring Finger Protein 2) and represses the histone H2A/H2B

deubiquitinase, USP3 (Ubiquitin Specific Peptidase 3). Thus SWI/SNF can potently shape the

epigenetic landscape, in an SCML2 dependent and independent fashion to regulate germ line

transcription.

Results:

BRG1 associates with transcriptionally active and poised chromatin

To understand the functions of SWI/SNF activity during meiosis, we determined the genome-wide

association of BRG1 by ChIP-seq. We reasoned that the examination of BRG1 binding dynamics

concurrent to critical meiotic processes such as DSB repair and homologous chromosome

synapsis would yield insight into the molecular mechanisms underlying the meiotic defects

observed in the *Brg1$^{fl/\Delta}$; Mvh-cre$^{Tg/0}$* (*Brg1^{cKO}*; see materials and methods) males. For this purpose

we isolated chromatin from spermatogenic cells obtained from P12 (mostly pre-pachytene germ

cells) and P18 testes (predominantly pachytene spermatocytes) (Bellve et al., 1977; Goetz et al.,

1984). At both P12 and P18, BRG1 appears to be enriched promoter proximally (Fig. 1A, panels

1 & 4). This is consistent with the fact that more than 50% of BRG1 peaks occupy promoters of

genes in P12 and P18 spermatogenic cells, with only a minority mapping to distal sites (Fig. S1A).

By performing K-means clustering we were able to categorize transcription start sites (TSS) into

three different classes (class1: Cl-1, class2: Cl-2, class: Cl-3) based upon their association with

BRG1. These TSS's are characterized by high (Cl-1), medium (Cl-2) and negligible (Cl-3) BRG1

enrichment.

To gain insight into the activity of genes associated with promoters occupied by BRG1 in

spermatogenic cells, we surveyed the chromatin environment surrounding these TSS. We

specifically monitored the enrichment of trimethylation of histone H3 lysine 4 (H3K4me3),

associated with gene activation and trimethylation of histone H3 lysine 27 (H3K27me3),

associated with gene repression. We compared BRG1 ChIP profiles from P12 and P18 testes to

stage-equivalent H3K4me3 and H3K27me3 ChIP profiles generated from P12 and P17 testes

(Mu et al., 2014) (Fig. 1A). Robust H3K4me3 enrichment appeared to be associated stably with

Cl-1 and Cl-2 promoters and was near-absent at Cl-3 promoters from P12 to P17 (Fig. 1A, panels

2 & 5). On the other hand, H3K27me3 levels at P12 and P17 appear depleted at Cl-1 (high BRG1)

promoters. Such antagonism is a well-established feature of SWI/SNF and PRC2 genomic

associations (Wilson et al., 2010). Cl-3 (low to no BRG1) promoters only displayed basal levels

of H3K27me3, which is particularly discernable at P17. In contrast to Cl-1 and Cl-3, Cl-2 promoters

that feature medium BRG1 enrichment display significantly higher levels of H3K27me3, which

appears progressively enhanced from P12 to P17 (Fig. 1A, panels 3 & 6). The co-occurrence of

H3K27me3 and H3K4me3 at Cl-2 promoters resemble features of bivalent promoters, which are

usually associated with transcriptionally quiescent genes poised to be reactivated at later stages

of development (Bernstein et al., 2006; Hammoud et al., 2014; Lesch et al., 2013). Thus genes

associated with Cl-2 promoters likely represent repressed target genes. In contrast potential

activated gene targets appear to be associated with Cl-1 BRG1 promoters.

To understand BRG1 directed gene regulation in the context of spermatogenesis, we categorized

the genes associated with each K-means cluster (Fig. 1A), by their temporal expression profiles,

previously determined from whole-testes RNA seq data (Margolin et al., 2014). The genes were

split between the five major categories, which include those maximally expressed in testes, at P6

(pre-meiotic: Pmei), from P8-P20 (meiotic: Mei), at P38 (Late, adult testis), from P6-P38

(Constantly; Const) and those expressed at low levels in testes from P6-P38 (Low, < 2 Reads per

Kilobase per Million of Reads; RPKM). The majority of Cl-1 associated genes were meiotic with

relatively fewer pre-meiotic genes. In contrast, Cl-2 and Cl-3 were mostly comprised of pre-meiotic

and low genes (Fig. S1B). This is particularly interesting given that Cl-1 gene promoters appear

to be exclusively marked by H3K4me3 indicative of transcriptional activation, while the Cl-2

promoters display bivalent chromatin modifications (H3K4me3/H3K27me3), which is indicative of

transcriptional silencing. We observed similar trends when monitoring H3K4me3 and H3K27me3

dynamics, specifically at BRG1 target promoters (P12 peaks) categorized by their temporal

expression profile during meiosis (P12 to P17) (Fig. 1B). Over this duration all target gene

categories experience a 4-fold decrease in BRG1 enrichment in the presence of relatively

abundant levels of H3K4me3 (Fig. 1B, top and middle row). Only pre-meiotic and low gene targets

display elevated levels of H3K27me3 from P12 to P17 (Fig. 1B, bottom row), distinguishing them

from, meiotic, constant and late gene targets, that appear exclusively marked by H3K4me3 (Fig.

1B, middle and bottom row). Candidate pre-meiotic targets such as *Zbtb16* and *Id4*, which are

markers of undifferentiated spermatogonial cells along with *Pdgfra*, a somatic signaling receptor,

that are normally repressed during meiosis, display promoters bivalently modified with H3K4me3

and H3K27me3 (Basciani et al., 2002; Green et al., 2018; Hammoud et al., 2014). In contrast, the

promoter of BRG1 target, *Sycp1*, which is expressed during meiosis and is essential for

synaptonemal complex assembly is marked exclusively by H3K4me3 (Fig. S1C) (De Vries et al.,

2005). Thus BRG1 might co-ordinate the expression of genes over the measured course of

spermatogenesis.

Given its robust association with active (H3K4me3) and poised (H3K4me3/H3K37me3)
chromatin, we were curious to examine BRG1 localization to the sex chromosomes, which are
transcriptionally silenced during pachynema (reviewed in Turner, 2007). We examined BRG1
occupancy at the TSS's of X – linked genes in P12 (Pre-pachytene stages) and P18 (Pachytene
stages) spermatogenic cells. BRG1 enrichment at X – linked TSS's appeared reduced at P18
relative to P12, but not absent (Fig. 1C). Thus BRG1 associates with regions that are subject to
transcriptional repression during meiosis.
194
Apart from TSS's, H3K4me3 is also enriched at DSB/recombination hotspots, known to be
associated with a meiosis specific histone methyl transferase, PRDM9 (Brick et al., 2012;
Diagouraga et al., 2018; Hayashi et al., 2005). We therefore examined BRG1 association at these
PRDM9 sites previously mapped by ChIP-seq in P12 testes (Baker et al., 2015). The lack of any
association with PRDM9 peaks makes it unlikely that BRG1 directly affects DSB formation (Fig.
S1D). Thus BRG1 might play a major role in gene regulation during meiosis.
201
202

BRG1 co-ordinates spermatogenic gene expression
The promoter centric association of BRG1 prompted us to examine its influence on the
transcription of target genes by RNA-seq. We compared transcript abundance between
spermatogenic cells isolated from P12 $Brg1^{fl/+}$ ($Brg1^{WT}$) and $Brg1^{fl/\Delta}; Mvh\text{-}cre^{Tg/0}$ ($Brg1^{cKO}$) testes.
The quantification of spermatocytes per tubule present at various stages of prophase I from
$Brg1^{WT}$ and $Brg1^{cKO}$ testes cryosections, staged by γH2Ax staining, revealed similar proportions
of pre-pachytene populations up to P13 (Fig. S2A, left panel). Beyond this stage we began to
notice differences in the cellular profiles between $Brg1^{WT}$ and $Brg1^{cKO}$ testes at P14 (Fig. S2A,
left panel). This is consistent with the incidence of pachytene arrest induced apoptosis in
previously reported in $Brg1^{cKO}$ testes (Kim et al., 2012). Thus we considered P12 to be an
appropriate stage at which RNA-seq could be performed between $Brg1^{WT}$ and $Brg1^{cKO}$ testes. In
agreement with the cell quantification, an examination of prophase I substage specific protein
coding transcripts (Ball et al., 2016) revealed no significant differences in the abundance of pre-
pachytene staging markers between P12 $Brg1^{WT}$ and $Brg1^{cKO}$ testes (Fig. S2A). Only, early and
late pachytene specific transcripts were slightly less abundant in the P12 $Brg1^{cKO}$ relative to
$Brg1^{WT}$ testes (Fig. S2A, right panel). This is suggestive of a minor difference in the onset of

pachynema. Overall we do not expect the analysis of gene expression to be impacted significantly

by secondary effects such as developmental delays.

To identify genes significantly mis-expressed (FDR ≤ 0.05) upon the loss of BRG1, we performed

an edgeR analysis on the RNA-seq data. An equivalent number of genes are either

transcriptionally down regulated (n=1100) or up regulated (n=983) in P12 *Brg1^cKO* relative to

Brg1^WT testes (Fig. 2A). More genes were down regulated (n=310) by a magnitude of 2-fold or

higher, relative to those up regulated (n=75), upon the loss of BRG1 (Fig. 2A) . Nearly half of all

the differentially regulated genes are occupied by BRG1 peaks (P12 peaks). The down regulated

genes were associated with transcripts normally expressed in gonadal tissue and the nervous

system and were enriched for Gene ontology (GO) terms relevant to meiotic processes (Fig. 2B).

In contrast ,the up regulated genes are normally expressed in limb and muscle and were enriched

for GO terms relevant to somatic developmental processes (Fig. 2B). Thus, BRG1 coordinates

germ line transcription by activating meiotic genes while concomitantly repressing somatic genes.

Since BRG1 is essential for meiotic progression (Kim et al., 2012), we monitored the expression

of genes associated with abnormal spermatogonia proliferation and meiotic arrest phenotypes,

curated from the mouse genome database (Blake et al., 2003). We integrated the expression

data into a heatmap that displays z-scores associated with transcript abundance of each

candidate gene, measured across P12 *Brg1^WT* and *Brg1^cKO* replicates   (Fig. 2C). With the

exception of *Androgen receptor* (*Ar*) and *Microtubule associated protein 7* (*Map7*), all other germ

cell factors appeared less abundant upon the loss of BRG1 (Fig. 2C). Next, we adopted a reverse

genetic approach to test whether the mis-expression of candidate genes were associated with

specific phenotypes in the *Brg1^cKO*. To do this we choose a pre-meiotic candidate, *Zbtb16* and

meiotic candidate, *Sycp2*.

Zbtb16 is essential for the maintenance of a pool of undifferentiated spermatogonia (typeA, SpgA)

(Buaas et al., 2004) . While BRG1 has been shown to be dispensable for the establishment of

spermatogonia, its role in the maintenance of SpgA is unknown (Kim et al., 2012). We quantified

the number of SpgA in P10 *Brg1^WT* and *Brg1^cKO* testes cryosections by immunofluorescence

staining for ZBTB16. The *Brg1^cKO* testes contain 46% fewer SpgA, when compared to *Brg1^WT*

testes (Fig. 2D). Since a reduction in the number of SpgA in the *Brg1^cKO* can skew the measured

abundance of *Zbtb16* mRNA, we performed qRT-PCR to measure transcript abundance in

purified, THY1+ fraction containing spgA cells obtained from P8 *Brg1^WT*, *Brg1^Het* (*Brg1^fl/+;Mvh-*

cre$^{Tg/0}$) and *Brg1^{cKO}* testes. In addition to *Zbtb16*, we also monitored the expression of other

established stem cell factors such as, *Inhibitor of DNA binding 4* (*Id4*), *POU domain, class 5,*

transcription factor 1 (*Pou5f1*, also known as *Oct4*)*, POU domain, class 3, transcription factor 1*

(*Pou3f1, also known as Oct6*) and *Forkhead box O1* (*Foxo1*) (Dann et al., 2008; Goertz et al.,

2011; Oatley et al., 2011; Wu et al., 2010). The loss of BRG1 resulted in a significant decrease in

the abundance of transcripts associated with *Zbtb16*, *Id4* and *Pou5f1* in the purified THY1^{+}

fractions (Fig. S2B). The fact that *Thy1* mRNA levels remain abundant argues against a loss of

spermatogonial cells early in development (Fig. S2B). Thus, BRG1 influences the maintenance

of a pool of undifferentiated spermatogonial cells by activating the expression of critical stem cell

factors.

The meiotic gene candidate, *Sycp2*, constitutes the structural component of the lateral element

of meiotic chromosomal axes and is essential for synapsis (Yang et al., 2006). Coincidentally,

Brg1^{cKO} spermatocytes display an increase in asynapsis (Kim et al., 2012). In the *Brg1^{cKO}*, SCYP2

levels are distinctly lower relative to the controls (Fig. 2E, Fig. S2C) and abnormally assemble

into short lateral filaments and aggregates in about 37% of the mutant zygotene spermatocytes

(Fig. 2F, Fig. S2D). While *Sycp2$^{-/-}$* spermatocytes fail to form SYCP3 elements , we did not

observe a similar defect in the *Brg1^{cKO}* spermatocytes (Fig. S2D) (Yang et al., 2006). We reason

that the reduced levels of SYCP2 might be sufficient to facilitate apparently normal SYCP3

assembly. Thus, a paucity in SYCP2 may limit meiotic progression by potentially compromising

the formation of a functional synaptonemal complex. In addition to the mis-regulation of essential

germ cell factors, we also identified a significant increase in the expression of several somatic

genes in the *Brg1^{cKO}* (Table S1). *Pdgfra* a signaling receptor, generally associated with somatic

(Basciani et al., 2018; Schmahl et al., 2008) and pre-meiotic spermatogonia (Hammoud et al.,

2014) , was up-regulated by more than 2-fold over a period spanning meiosis-I in the *Brg1^{cKO}* (Fig.

S2E).

BRG1 is required to maintain chromatin accessibility at promoters

Chromatin remodelers reposition nucleosomes, consequently regulating accessibility to

transcription factors. Thus BRG1 might influence transcription by modulating the structure of the

germ line epigenome. To investigate this, we performed Assay for Transposase-Accessible

Chromatin (ATAC)-seq to map open chromatin in pre-pachytene and pachytene spermatogenic

cells isolated from P12 and P18 testes respectively. Similar to the RNA-seq data, we surveyed

differences in chromatin accessibility between P12 $Brg1^{WT}$ and $Brg1^{cKO}$ testes (Fig. S3A, left

panel). Since P18 $Brg1^{cKO}$ testes are characterized by severe pachytene arrest, we compared

chromatin accessibility between P18 $Brg1^{WT}$ and $Brg1^{Het}$ ($Brg1^{fl/\Delta}$) testes (Fig. S3A, right panel).

In normal spermatogenic cells, strong ATAC read coverage was detected promoter proximally at

both P12 and P18 (Fig. S3A). This promoter accessibility is significantly diminished in P12

$Brg1^{cKO}$ testes and also under conditions of haploinsufficiency in P18 $Brg1^{Het}$ testes (FigS3A).

Since promoter proximal chromatin responds to the loss of BRG1, we first monitored the changes

in chromatin accessibility at promoters of target genes differentially regulated by BRG1 (Fig. 3A).

Promoters of BRG1 target genes that are normally activated (down regulated upon BRG1 loss),

display a significant decrease in promoter accessibility in the $Brg1^{cKO}$ relative to $Brg1^{WT}$ testes.

While the promoters of repressed gene targets (up regulated upon BRG1 loss) are less accessible

relative to their activated counterparts in $Brg1^{WT}$ testes, their accessibility appears further reduced

in the $Brg1^{cKO}$ testes, albeit at levels that fail to meet statistical significance (Fig. 3A). Thus only

activated gene promoters experience a significant change in accessibility.

To identify genome-wide changes in chromatin accessibility at P12, we performed edgeR on the

ATAC read counts obtained from wild type and mutant samples. The vast majority of regions that

displayed significant differences in chromatin accessibility appear to be less accessible (Closed;

n= 549), leaving only a few regions that acquire greater accessibility (Opened; n= 99) in the

$Brg1^{cKO}$ relative to the $Brg1^{WT}$ testes (Fig. 3B). Consistent with the general decrease in promoter

accessibility in the $Brg1^{cKO}$ (Fig. S3A), the closed regions are overwhelmingly associated with

promoters (Fig. 3C, Fig. S3B; panel 1). In contrast, the opened regions were prominently featured

within introns and intergenic regions (Fig. 3C, Fig. S3B, panel 2 & 3). This overall decrease in

chromatin accessibility genome-wide is consistent with the previously observed increase in

repressive epigenetic modifications in $Brg1^{cKO}$ spermatocytes (Kim et al., 2012). The genes

associated with closed promoters in $Brg1^{cKO}$ testes were mostly meiotic in function (Fig. 3D, Table

S2). Thus BRG1 probably activates meiotic genes by maintaining chromatin accessibility at

cognate promoters. In contrast a few genes associated with the distal sites that appear more

accessible in the $Brg1^{cKO}$ testes, represent somatic factors (Fig. 3D, Table S2).

BRG1 physically interacts with SCML2, a non-canonical PRC1 factor

To investigate further mechanisms governing SWI/SNF mediated epigenetic regulation, we monitored BRG1 interactions during meiosis. BRG1 Immuno-pulldowns (IP) were prepared from nuclear extracts obtained from 3-week-old testes (Fig. 4A). Proteins isolated from a BRG1 IP and control nonspecific (ns) IgG pulldown were identified by mass spectrometry (MS). Peptides belonging to known SWI/SNF sub-units were identified only in the BRG1 IP, which demonstrates the efficacy of this procedure (Fig. 4A, Fig. S4A). Furthermore the presence of both SWI/SNF sub complexes; *Brahma* Associated Factor (BAF) and *polybromo*-BAF (PBAF) are detected in the germ line (Fig. 4A,Fig. S4A). More interestingly we identified peptides associated with SCML2, a known testes specific PRC1 factor (Hasegawa et al., 2015; Luo et al., 2015) (Fig. 4A). Since candidate peptides were also detected in the non-specific IgG IP, we validated these interactions by performing a reverse IP with an antibody against SCML2 (Fig. 4B). The IP was conducted on nuclear lysates treated with universal nuclease (benzonase) to eliminate the possibility of non-specific, DNA mediated interactions. BRG1 was specifically detected in the SCML2 IP, compared to the nsIgG (Fig. 4B, lane 2 & 3). Furthermore we also detected a previously established interaction between SCML2 and RNF2 (Fig. 4B, lane3) (Hasegawa et al., 2015), which builds our confidence in the results. However, the smear like appearance of SCML2 in the nuclear extracts (Fig 4B) prompted us to confirm the specificity of the SCML2 antibody. We did this by performing western blots on nuclear extracts obtained from spermatogenic cells and ovaries. Consistent with its male specific expression pattern, we fail to detect a SCML2 signal in nuclear extracts obtained from ovaries (Fig.S4B, lane 2). Hence the smearing might be a product of protein instability or indicative of isoforms.

In pachytene spermatocytes, SCML2 is also known to interact with USP7, a known deubiquitinase and non-canonical member of the mammalian PRC1.4 complex (Lecona et al., 2015; Luo et al., 2015). Hence we were curious to determine its interaction with BRG1. We were unable to detect an interaction between USP7 and BRG1 (Fig. S4C, lane 2-4). Thus the BRG1 interaction appears to be limited to SCML2 during meiosis.

BRG1 and SCML2 concordantly regulate genes during meiosis

In pachytene spermatocytes, SCML2 is known to repress somatic and spermatogonial genes, while concurrently activating certain meiotic and late spermatogenic genes (Hasegawa et al.,

2015). This is similar to the epigenetic role of BRG1 in the germ line. Hence it might be possible
that BRG1 interacts with SCML2 to mutually regulate the germ line transcriptome. In support of
such a possibility, we identified robust enrichment of BRG1 at TSS's of genes differentially
regulated (FDR < 0.05) by SCML2, during pachynema (Fig. 4C). The relative enrichment was
greater at P12 (8 fold over input) compared to P18 (6 fold over input), indicating that BRG1
localizes prior to pachytene at this sites. Furthermore, chromatin accessibility at TSS's associated
with SCML2 regulated genes is reduced upon the loss of BRG1 at P12 (Fig. S5A).

Next, we looked for commonly mis-expressed genes (FDR<0.05) between the P12 BRG1 and
pachytene SCML2 RNA-seq data. About 65% (n=597) of genes commonly mis-expressed in the
absence of either BRG1 or SCML2 exhibited concordant expression changes that were highly
correlated (r^2=0.72). The remaining 35% (n=330) showed discordant changes (r^2=0.51) (Fig. 4D).
Of the concordantly regulated genes, 64% (n=388) were repressed, while the remainder (n=209)
were activated. The concordantly repressed genes accounted for 40% of the genes up-regulated
in the P12 $Brg1^{cKO}$ testes. In contrast, only 19% of genes down regulated in the P12 $Brg1^{cKO}$ testes
overlapped with concordantly activated genes. The commonly repressed genes were mostly
somatic in function (Fig. S5B) and included the somatic signaling receptor, PDGFRA, associated
with BRG1 (Fig. S1C, Fig. S2E). Thus during normal prophase-I, BRG1 might achieve gene
repression by recruiting SCML2. On the other hand the co-activated genes were associated with
GO terms relevant to the mitotic spindle checkpoint (Fig S5B). Previous studies have
demonstrated the existence of a meiotic spindle checkpoint in spermatocytes, and candidate
genes such as $Smc2$, a member of the condensin I and II complexes, is known to influence female
meiosis (Eaker et al., 2001; Lee et al., 2011). Thus SCML2 might activate these spindle
checkpoint factors, at later stages of meiosis I, in a BRG1 dependent manner.

374
375

376 **BRG1 influences SCML2 and USP7 localization to the sex body**
377 During pachynema, both SCML2 and its interacting partner, USP7 paint the sex body, a γH2Ax
378 enriched, sub-nuclear compartment containing the sex linked chromosomes (Hasegawa et al.,
379 2015; Luo et al., 2015). Therefore we determined whether SCML2 localization to the sex body
380 was dependent on BRG1.
381

We first compared SCML2 localization in *Brg1^WT* and *Brg1^cKO* testes cryo-sections from 2- and

3-week-old males by IF (Fig. 5A,Fig. S6A). Mutant pachytene spermatocytes were identified by

staining for γH2Ax, given that its association with the sex body remains unperturbed in the *Brg1^cKO*

(Kim et al., 2012). The loss of BRG1 appeared to impact the localization of SCML2 in *Brg1^cKO*

pachytene spermatocytes (Fig. 5A). The SCML2 signal in the mutants appeared abnormally

distributed genome-wide, without displaying the strong sex body association, typically observed

in *Brg1^WT* (Fig. 5A, panel 3 insets). We confirmed these defects, by co-staining for ATR (Ataxia

Telangiectasia and Rad3 Related), a DDR factor enriched on the sex body (Royo et al., 2013)

and also stained for BRG1, to demonstrate protein loss in pachytene spermatocytes displaying

SCML2 mis localization (Fig. S6A). Given that SCML2 is known to be physically associated with

γH2Ax (Hasegawa et al., 2015; Luo et al., 2015), it may be that in the absence of BRG1, it is de-

localized to autosomal sites harboring persistent γH2Ax (Fig 5A, panel insets). Surprisingly, subtle

defects in SCML2 localization were also seen in   *Brg1^Het* pachytene spermatocytes, where it

appeared more homogenously distributed genome-wide (Fig. S6B, panel insets).

We next examined SCML2 localization in *Brg1^WT* and *Brg1^cKO* meiotic spreads, co-stained with

a synaptonemal complex marker, SYCP3, to visualize chromosomes. Mutant meiotic spreads

were obtained from *Brg1^cKO* testes generated using two independent germ line specific CRE's,

namely the *Mvh-cre* and *Stra8-cre*. Similar to the cryosections, SCML2 association with the sex

chromosomes was perturbed in *Brg1^cKO*   pachytene spreads (Fig. S7). However, *Brg1^cKO*

pachytene spermatocyte generated with *Stra8-cre*, exhibited a less severe defect in SCML2

localization to the sex chromosomes, relative to their *Mvh-cre* derived counterparts (Fig. S7).

Such differences might be a consequence of the distinct temporal activity of each CRE (see

materials and methods) (Gallardo et al., 2007; Sadate-Ngatchou et al., 2008).

In addition to SCML2 we also monitored the localization of USP7 in pachytene spermatocytes

obtained from *Brg1^WT* and *Brg1^cKO* testes. Even though BRG1 does not directly interact with USP7

(Fig. S4C), we posited that the mis-localization of SCML2 in *Brg1^cKO* pachytene spermatocytes

might also affect USP7 enrichment on the sex body. In fact this seems to be the case in *Brg1^cKO*

pachytene spermatocytes immunofluorescently stained for USP7 and γH2Ax (Fig. 5B).

From previous studies it is clear that the mis localization of SCML2 does not affect processes,

critical to the initiation of meiotic sex chromosome inactivation (MSCI) (Hasegawa et al., 2015).

Since the loss of BRG1 has been previously reported to influence MSCI (Wang et al., 2012), we

examined its impact on the localization of known MSCI factors such as BRCA1, ATR and MDC1

to the sex body (Ichijima et al., 2011; Turner et al., 2004). By immunofluorescence, BRCA1, ATR

and MDC1 appeared stably associated with the sex body in the $Brg1^{cKO}$ pachytene spermatocytes

(Fig. 5C). Thus BRG1 like SCML2 does not affect the initiation of MSCI. Interestingly, it was

previously reported that the loss of MDC1 abrogates SCML2 recruitment to the sex chromosomes

(Hasegawa et al., 2015). Therefore, the stable association of MDC1 with the sex body in $Brg1^{cKO}$

spermatocytes (Fig. 5C), suggests that SWI/SNF may function downstream MDC1 in the

recruitment of SCML2 to the sex body.

BRG1 influences the abundance of SCML2 associated histone modifications

Since SCML2 has been shown to regulate both H2AK119ub1 and H3K27ac during meiosis

(Adams et al., 2018; Hasegawa et al., 2015), we investigated whether BRG1 influences these

histone post translational modifications (PTM). We first, monitored the abundance of

H2AK119ub1 and H3K27ac in P12 $Brg1^{WT}$, $Brg1^{Het}$ and $Brg1^{cKO}$ spermatogenic cells by

performing western blots on acid extracted histones (Fig. 6A, top panel). In addition to these

histone PTM's, we also examined the abundance of H3K4me3, associated with BRG1 target

promoters and PRC2 catalyzed H3K27me3 (Fig. 6A, bottom panel). While neither H3K4me3 nor

H3K27me3 levels were perturbed, both H2AK119ub1 and H3K27ac were dramatically elevated

in P12 $Brg1^{Het}$ and $Brg1^{cKO}$, relative to the $Brg1^{WT}$ spermatogenic cells (Fig. 6A, top panel).

Interestingly, these epigenetic perturbations were undetectable at P8 and P10, which coincide

with the initiation of meiosis and pairing (Leptonema to zygonema) (Fig. S8A). Thus BRG1

suppresses H2AK119ub1 and H3K27ac prior to the onset of pachynema. The dose dependency

of such epigenetic regulation, prompted us to verify whether it was a consequence of the $Brg1$

hypomorph or due to the mere presence of the Mvh-cre. The former scenario seems to be true,

as evidenced by lack of any difference in the levels of H2AK119ub1, between males with the CRE

and their littermate control (Fig. S8B). Another cause for concern was that, the H2AK119ub1

antibody (clone E6C5) used in this study was recently reported to recognize non-histone epitopes

(Hasegawa et al., 2015). Hence, we validated the specificity of clone E6C5 by performing western

blots on acid-extracted histones obtained from RNF2 (PRC1 E3-ubiquitiun ligase) knock out

($Rnf2^{KO}$) embryonic stem (ES) cells, engineered using CRISPR-CAS9. The near-absence of

H2AK119ub1 signal in the *Rnf2^KO* relative to *Rnf2^WT* ES cell histone extracts confirms the

specificity of clone E6C5 (Fig. S8C). Furthermore, H2AK119ub1 is clearly detected in chromatin

fractions obtained from P12 *Brg1^WT* , *Brg1^Het* and *Brg1^cKO* testes (Fig. S8D). Thus the changes in

H2AK119ub1 reported here are genuine.

BRG1 suppresses H2AK119ub1 and enhances H3K27ac at target promoters

Based on the genome-wide increase in H2AK119ub1 and H3K27ac in *Brg1^cKO* testes, we were

encouraged to survey the changes in these histone PTM's at BRG1 target promoters by ChIP-

seq. We compared the enrichment of H2AK119ub1 and H3K27ac between P12 *Brg1^WT* and

Brg1^cKO spermatogenic cells at promoters normally associated with BRG1 peaks. First, we

categorized the BRG1 targets by their chromosomal location, the rationale being that both

H2AK119ub1 and H3K27ac are known to be differentially regulated between the autosomes and

sex chromosomes at pachynema (Adams et al., 2018). The BRG1 target promoters displayed

contrasting changes in H2AK119ub1 and H3K27ac enrichment in the *Brg1^cKO* spermatogenic

cells. Irrespective of chromosomal location, target promoters displayed enhanced H2AK119ub1

combined with reduced H3K27ac in the *Brg1^cKO* relative to *Brg1^WT* spermatogenic cells (Fig. 6B,

top panel). This mirrors the changes in H2AK119ub1 and H3K27ac that occur on the sex

chromosomes in *Scml2^KO* pachytene spermatocytes (Adams et al., 2018; Hasegawa et al., 2015).

Thus, the regulation of H2AK119ub1 and H3K27ac at regions occupied by BRG1 may be

influenced by SCML2, with which it is physically associated.

Next, we analyzed the changes in H2AK119ub1 and H3K27ac at the promoters of BRG1 target

genes that are differentially expressed. We hypothesized that these epigenetic modifications

might influence the transcriptional status of genes regulated by BRG1. Consistent with changes

observed at all BRG1 occupied promoters (Fig. 6B, top panel), H2AK119ub1 was elevated while

H3K27ac was reduced at promoters of mis-expressed target genes, irrespective of their

transcriptional status in the *Brg1^cKO* relative to *Brg1^WT* spermatogenic cells (Fig. 6B, bottom panel).

These epigenetic changes are typically associated with gene repression and thus only account

for the silencing of mis-expressed genes in *Brg1^cKO* spermatogenic cells. Therefore, the activation

of mis-expressed target genes occurs by mechanisms independent of H2AK119ub1 or H3K27ac.

The loss of H3K27ac at promoters normally occupied by BRG1 in *Brg1^cKO* spermatogenic cells is

inconsistent with its genome-wide increase (Fig.6A; top panel , Fig. 6B). To determine whether

other genomic regions acquire greater H3K27ac enrichment upon the loss of BRG1, we

performed a differential peak calling analysis using the macs2 bdgdiff algorithm (Zhang et al.,

2008). This analysis revealed several intronic and intergenic regions that lost, gained or

maintained (common) H3K27ac peaks in *Brg1^cKO* relative to the *Brg1^WT* spermatogenic cells (Fig

6C; left panel, Fig. S9). Compared to the common peaks, regions that loose or gain H3K27ac are

normally depleted of BRG1 (Fig. 6C, right panel). Thus while total H3K27ac levels are elevated,

its local distribution appears more heterogeneous upon the loss of BRG1.

BRG1 can also influence the epigenome in a SCML2 independent manner

While our data suggest that BRG1 regulates H2AK119ub1 and H3K27ac, through its interaction

with SCML2, we cannot rule out the possibility that BRG1 might also influence these histone

PTM's by directly regulating the expression of cognate epigenetic modifiers. In fact, our ChIP-seq

data identifies BRG1 peaks at promoters of epigenetic modifiers known to influence H2AK119ub1,

H2BK120ub1 and H3K27ac (Fig. S10A). These include the H2A ubiquitin ligase, RNF2 (Wang et

al., 2004), USP3 a H2A/H2B deubiquitinase (Nicassio et al., 2007) and the histone deacetylases,

HDAC1 and HDAC2 (Gallinari et al., 2007) (Fig. S10A). The transcript abundances of all these

targets are significantly altered in response to the loss of BRG1 (Fig.S10B).

In the case of the histone ubiquitin modifiers, *Rnf2* and *Usp3*, the former displayed an increase

while the latter displayed a decrease in transcript abundance in *Brg1^cKO* relative to the *Brg1^WT*

spermatogenic cells at P12 (Fig. S10B, row 1). This is consistent with significant increases in

protein levels in *Brg1^cKO* (RNF2: 88% increase, USP3:59% decrease) relative to the *Brg1^WT*

chromatin fractions prepared from testes (Fig. 6D, lane 4-6). Thus BRG1 may influence

H2AK119ub1 by maintaining a balanced expression of RNF2 and USP3. Furthermore this

represents an SCML2 independent mechanism of epigenetic regulation. Since USP3 has also

been associated with mono ubiquitinylated H2B (Nicassio et al., 2007), we monitored the levels

of mono ubiquitination of H2B lysine 120 (H2BK120ub1), a mark associated with gene activation

and chromatin relaxation (Fierz et al., 2011; Minsky et al., 2008; Pavri et al., 2006). At P12,

H2BK120ub1 appeared elevated genome-wide in $Brg1^{cKO}$, relative to $Brg1^{WT}$ spermatogenic cells

(Fig. S10C). The changes in the $Brg1^{Het}$ spermatogenic cells appear to be a product of unequal

protein loading.

Similar to the histone ubiquitin modifiers, H3K27ac associated modifiers, *Hdac1* and *Hdac2*

displayed reduced transcript abundance in the $Brg1^{Het}$ and $Brg1^{cKO}$ (Fig. S10B, row 2). At least

in the case of *Hdac1*, we were unable to identify a corresponding depletion in nuclear protein

levels by western blot (Fig. 6D, lane 1-6). Since HDAC1 has been shown to compensate for

HDAC2 in various developmental scenarios (Ma et al., 2012; Montgomery et al., 2009; Yamaguchi

et al., 2010) it is unlikely that the perturbation in H3K27ac levels in the $Brg1^{cKO}$ spermatogenic

cells, is associated with mis-expression of histone deacetylases.

Discussion:

In this study we integrate genomic and proteomic approaches to show that SWI/SNF directed regulation of transcription, influences meiotic progression in males. In spermatogenic cells, BRG1 is overwhelmingly promoter associated which is distinct from what has been observed in other mammalian cell types and embryonic tissue (Alexander et al., 2015; Alver et al., 2017; Attanasio et al., 2014). This re-enforces the notion that cell or tissue specific associations influence SWI/SNF function during development (reviewed in Ho and Crabtree 2010). We propose a model in which the SWI/SNF ATPase, activates essential spermatogenic genes, while maintaining the repression of somatic genes. We show that BRG1 facilitates chromatin accessibility at the promoters of these differentially regulated genes, which is consistent with the generally accepted role of SWI/SNF chromatin remodeling (reviewed in Clapier et al., 2017). The activated genes play critical roles in the maintenance of undifferentiated spermatogonial cell populations and facilitate meiotic progression. Target stem cell factors include, *Zbtb16* and *Id4*. The latter specifically identifies a self-renewing population of spermatogonia, generally referred to as spermatogonial stem cells (SSC) (Green et al., 2018; Oatley et al., 2011). Thus BRG1 might influence SSC maintenance. The role of BRG1 in the maintenance of pluripotency while not novel, was previously unknown in the germ line, thus demonstrating a conserved role for SWI/SNF across various stem cell lineages (Ho et al., 2009; Lessard et al., 2007) . In the case of meiotic gene targets, several of these have been associated with meiotic arrest phenotypes. The role of BRG1 in the activation of *Sycp2*, which is associated with synaptonemal complex assembly and homolog synapsis (Yang et al., 2006), potentially explains the incomplete synapsis and subsequent meiotic arrest seen in *Brg1^cKO* spermatocytes (Kim et al., 2012). Thus the male sterility associated with *Brg1^cKO* adults is a consequence of a shortage of germ line progenitors and essential meiotic factors.

The repression of the somatic transcriptome during meiosis is a feature shared with two other epigenetic regulators, namely, PRC2 and SCML2, a testes specific PRC1 factor (Hasegawa et al., 2015; Mu et al., 2014). Here we propose that BRG1 achieves the repression of its target genes by recruiting SCML2 activity. This is supported by the following observations; BRG1 (i) physically interacts with SCML2, (ii) is enriched at the promoters of genes regulated by SCML2 (iii) concordantly represses most genes commonly regulated by SCML2 and (iv) influences SCML2 localization during pachynema, based on a pronounced effect on sex body association.

Interestingly, the transition to pachynema is accompanied by a reduction but not depletion of
BRG1 from the X chromosome. This reduced level of BRG1 might be sufficient to recruit SCML2
to the sex body. Alternatively, the enrichment of SCML2 on the sex body may be dependent on a
temporally defined interaction with BRG1 that precedes pachynema. In contrast to the repressed
genes, BRG1 concordantly activates fewer SCML2 regulated genes. Most of these are associated
with the regulation of mitotic spindle checkpoints, which in the context of meiosis occurs late into
meiosis-I (Eaker et al., 2001).

In addition to these observations, BRG1 also appears to influence SCML2 associated histone
PTM's, H2AK119ub1 and H3K27ac in the male germ line (Adams et al., 2018; Hasegawa et al.,
2015). Similar to SCML2, we find that BRG1 is required to suppress H2AK119ub1 and enhance
H3K27ac at target promoters in pre-pachytene spermatogenic cells. However unlike SCML2,
BRG1 did not appear to differentially regulate autosomal and sex linked chromatin (Hasegawa et
al., 2015). It is possible that such differential regulation does not manifest at pre pachytene stages.
Alternatively, this may be indicative of distinct epigenetic outcomes associated with SCML2
dependent and independent mechanisms. The latter is demonstrated by the effect of BRG1 on
the expression of potent H2AK119ub1 modifiers, such as RNF2 and USP3.

As counter-intuitive as it may seem, repressed BRG1 targets genes are associated with low
H2AK119ub1 and higher H3K27ac levels. Hence, the repression of target genes might occur by
alternative mechanisms. Several BRG1 target promoters are bivalently modified
(H3K4me3/H43K27me3) during the course of meiosis from P12 to P17. These bivalent promoters
appear to distinguish somatic and pre-meiotic targets from meiotic targets. Hence, the silencing
of these genes might occur through the formation of bivalent promoters, which are commonly
associated with somatic developmental genes in the germ line (Hammoud et al., 2014; Lesch et
al., 2013; Lesch et al., 2016). Interestingly, SCML2 has been recently shown to influence the
formation of bivalent domains during germ line development (Maezawa et al., 2018). Hence, such
SCML2 activity might potentially influence the repression of BRG1 target genes.

In addition to H2AK119ub1, BRG1 also suppresses H2BK120ub1, which is associated with
transcriptional activation and chromatin relaxation (Fierz et al., 2011; Minsky et al., 2008; Pavri et
al., 2006). Hence, an increase in H2BK120ub1 might potentially activate genes mis-expressed in
the *Brg1cKO* spermatogenic cells. This might occur due to the mis-regulation of USP3 or mis-

localization of USP7, two deubiquitinases, whose activities have been shown to influence mono

ubiquitination of H2B (Nicassio et al., 2007; Van Der Knaap et al., 2005). The evidence for USP7

is based on its function in *Drosophila* (Van Der Knaap et al., 2005). In contrast, mammalian USP7

does not seem to catalyze the deubiquitnation of H2B (Lecona et al., 2015; Maertens et al., 2010).

However there appears to be an effect on H2BK120ub1 that is independent of its enzymatic

activity (Huether et al., 2014).

In conclusion we reveal the transcriptional basis for the meiotic defects previously described in

Brg1^{cKO} males, and present a new paradigm for studying co-operation between SWI/SNF and

PRC1 factors in the regulation of the epigenome. While recent studies have illustrated the

propensity of BRG1 to evict canonical PRC1 members from chromatin in normal and oncogenic

cell culture models (Kadoch et al., 2016; Stanton et al., 2016), the relationship between SWI/SNF

and variant PRC1 factors remain unexplored. More recent reports have described the roles of

variant PRC1 factors in gene activation in ES cells and the central nervous system (CNS) (Creppe

et al., 2014; Gao et al., 2014). Whether SWI/SNF influences these activities in the germ line

remains a topic of future investigation. Interestingly, a recent report has revealed a role for PRC1

in gene activation in spermatogonial cells (Maezawa et al., 2017).

Materials and Methods:

Generation of *Brg1* conditional deletion and genotyping:

Brg1 floxed (Sumi-Ichinose et al., 1997) , *Mvh-Cre* (activated at ~ E15) (Gallardo et al., 2007) and *Stra8-Cre* (activated only in males at P3) (Sadate-Ngatchou et al., 2008) were maintained on a outbred background genetic background using CD-1 mice. $Brg1^{fl/fl}$ females were crossed to $Brg1^{fl+}$;Mvh-$Cre^{Tg/0}$ males to obtain $Brg1^{fl\Delta}$;Mvh-$Cre^{Tg/0}$ ($Brg1^{cKO}$,) , $Brg1^{fl+}$;Mvh-$Cre^{Tg/0}$ ($Brg1^{Het}$) and $Brg1^{fl+}$ ($Brg1^{WT}$) littermate controls. Similar crosses were made to generate the *Stra8-Cre* induced conditional knockouts and littermate controls. Genotyping primers used in this study include: $Brg1^{fl/+}$ alleles - (F)- 5$^{'}$ -CCTAGCCAAGGTAGCGTGTCCTCAT-3$^{'}$; (R) 5$^{'}$ - CCAGGACCACATACAAGGCCTTGTCT-3$^{'}$, the excised allele (Δ)- reverse primer used above in combination with (F) 5$^{'}$ -CTAACCGTGTATGTAGCCAGTTCTGCCT-3$^{'}$, *Mvh-Cre* - (F) 5$^{'}$ - CACGTGCAGCCGTTTAAGCCGCGT-3$^{'}$, (R) 5$^{'}$ -TTCCCATTCTAAACAACACCCTGAA-3$^{'}$ and *Stra8-Cre* - (F) 5$^{'}$ -GTGCAAGCTGAACAACAGGA-3$^{'}$, (R) 5$^{'}$ -AGGGACACAGCATTGGAGTC-3$^{'}$. All animal work was carried out in accordance with approved IACUC protocols at the University of North Carolina at Chapel Hill.

Disruption of Rnf2 by CRISPR-Cas9:

The sequences of sgRNAs for Rnf2 are 5'CACCGTGTTTACATCGGTTTTGCG3' and 5'AAACCGCAAAACCGATGTAAACAC3'. sgRNAs were cloned into pX330-U6-Chimeric_BB-CBh-hSpCas9 (42230, Addgene) using Golden Gate assembly cloning strategy (Bauer et al., 2015). Briefly, 5 x 10^4 E14 ES cells were cultured on 60 mm dishes for one day and then transfected with plasmids expressing Cas9 and sgRNAs, along with a plasmid expressing PGK-PuroR (Addgene, cat. no. 31937) using FuGENE HD reagent (Promega) according to the manufacturer's instructions. The cells were treated with 2 µg/ml puromycin for two days and recovered in normal culture medium until ES cell colonies grew up. *Ezh1* targeted colonies were verified by DNA sequencing.

Immunofluorescence staining:

Spermatocyte spreads were prepared as described (Peters et al., 1997) or by using a protocol adapted from (Wojtasz et al., 2009) , that was used to generate "3D-preserved" spermatocytes. The latter entails a detergent spreading technique in which single- cell preparations obtained as described (Biswas 2018) were treated with 0.25% NP-40 for not more than 2 minutes. These

spreads were used to view SYCP2 staining. All spermatocyte spreads were generated from 2- to

3-week-old mice. Prepared slides were either dried down and stored at -80^0 C or stored in

phosphate buffered saline (PBS) at 4^0 C in the case of "3D-preserved" spermatocytes.

Testis cryosections were prepared essentially as described before (Kim et al., 2012) with a few

modifications. Briefly, juvenile/adult testes were fixed in 10% neutral buffered formalin at 4^{0}C.

After 20 minutes of incubation in NBF the tissue was sliced into half and then fixed up to 1 hour.

Fixed tissue were washed 3 times in PBS at room temperature and then saturated through a

sucrose series - 10% (30 min), 20% (30 min) and 30% (1 hr). The tissue were then incubated

overnight in 30% sucrose/optimum cutting temperature (OCT) formulation at 4^0 C and

subsequently embedded in OCT. Frozen sections were cut at 9 μm thickness. Antigen retrieval

was performed for all antibodies used in this study. Briefly slides were incubated in boiling 10mM

citric acid (pH6.0) for 10 minutes. Over this period, citrate buffer was replaced every 2 minutes

with fresh boiling buffer and then allowed to cool down gradually for up to 20 minutes.

Tissue section and spreads were washed in PBS followed by permeabilization in 0.1% Triton-X

100 and then blocked in Antibody dilution buffer (10% bovine serum albumin; 10% goat/donkey

serum; 0.05% Triton-X 100) diluted 1:10 in PBS for 20 minutes before incubation with primary

antibody overnight at 4^{0}C. The following day, samples were again washed, permeabilized and

blocked in ADB/PBS after which they were incubated for 1 hour with Alexa fluor- conjugated

secondary antibodies. Immuno-stained slides were finally washed twice in PBS/0.32% photoflo

(Kodak), once in H$_2$O/ 0.32% photoflo and then counterstained with DAPI before mounting in

Prolong Gold antifade medium (P-36931; Life Technologies). A list of all the primary antibodies

used in this study are provided (Table S4). We used alexa fluor conjugated secondary antibodies

at a dilution of 1:500. All imaging in this study was done on a Zeiss AxioImager-M2.

Isolation of Spermatogonial stem cells and RNA extraction:

Testes cell suspension was generated from 8-day-old *Brg1^{WT}* (n=5), *Brg1^{Het}* (n=1)*, Brg1^{cKO}* (n=2)

mice as described previously (Kubota and Brinster, 2008). Conditional deletions were generated

using the *Mvh-Cre* transgene. Spermatogonial stem cells (SSC) were enriched using THY1$^+$

microbeads (Miltenyi Biotec; 131-049-101) followed by their isolation on magnetic activated cell

sorting (MACS) columns (Miltenyi Biotec; 131-090-312). RNA from SSC's were isolated and

purified using the PicoPure^{TM} RNA isolation kit (Life technologies; KIT0204).

RT-PCR and qPCR:

676 cDNA was synthesized using random primer mix (NEB) and ProtoScript® II reverse transcriptase

678 (NEB). Real time qPCR was performed using Sso Fast EvaGreen supermix (Bio-Rad) on a CFX96

679 thermocycler (Bio-Rad). A list of qRT-PCR primers used in this study are provided (Table S5).

Isolation of spermatogenic cells:

682 Spermatocyte enriched populations were isolated by methods described previously (Chang et al.,

683 2011; Mu et al., 2014). Spermatocyte populations were enriched using percoll and cell strainers.

684 Cell populations were then used for downstream applications such as RNA-seq, ATAC-seq, ChIP-

685 seq and nuclear lysate preparations for immunoprecipitations.

RNA-seq:

RNA was extracted from spermatogenic cells obtained from 4 biological replicates of P12 $Brg1^{WT}$

and $Brg1^{cKO}$ mice each. Conditional deletions were generated using the Mvh-Cre transgene. Cells

were treated with the TRIzol reagent (Invitrogen) and total RNA was isolated and cleaned up

using the Direct-zol RNA kit (Zymo). Sequencing libraries were prepared using a Kapa mRNA

library kit as per manufacturer's instruction and then sequenced on a Illumina Hiseq 4000 (50 bp

reads, single end). Scml2 RNA-seq was previously published and is publically available under

GEO accession number GSE55060 (Hasegawa et al., 2015).

RNA-seq data analysis:

Gene expression was quantified using kallisto (Bray et al., 2016). Transcript levels (counts) were

summarized per gene using tximport (Soneson et al., 2016) and then imported to perform a

differential analysis of gene expression using edgeR (Robinson et al., 2010). The mouse (mm9)

gene/transcript annotations were retrieved using the ensembldb R package

(https://github.com/jotsetung/ensembldb). Low abundance genes (counts per million <1 across

4 replicates) were filtered out in edgeR and significant differences in counts were called at a false

discovery rate (FDR)≤ 0.05. The lists of differentially expressed genes are provided (Table S1).

Anatomy ontology terms were curated from EMAPA (The Edinburgh Mouse Atlas Project)

(Hayamizu et al., 2013) and the analysis was done on MouseMine (www.mousemine.org)

(Motenko et al., 2015). Gene ontology analysis were performed using clusterProfiler (Yu et al.,

2012).

ChIP-seq:

BRG1 ChIP was performed exactly as described previously (Raab et al., 2015). We performed

the ChIP in duplicates on 4×10^7 wild type (CD-1) spermatogenic cells obtained from P12 and P18

mice each. H2AK119ub1 and H3K27ac ChIPs were also performed in duplicate on spermatogenic

cells obtained from P12 $Brg1^{WT}$ and $Brg1^{cKO}$ mice using a method described previously for low

chromatin inputs (Brind'Amour et al., 2015), with minor modifications (See supplemental materials

for details). The BRG1 ChIP samples were sequenced on a Hiseq 2500 using v4 chemistry (50

bp reads, single end), whereas the H3K27ac and H2AK119ub1 ChIP samples were sequenced

on an Illumina Hiseq 4000 (50 bp reads, single end). The antibodies used for ChIP are listed

(table S4). H3K4me3, H3K27me3 ChIP-seq data were previously published and are publically

available under GEO accession number GSE61902 (Mu et al., 2014). ChIP data analysis methods

can be found in the supplemental materials.

ATAC-seq:

ATAC-seq was performed on spermatogenic cells isolated from 2 biological replicates of P12

$Brg1^{WT}$ and $Brg1^{cKO}$ mice each. Only a single sample from 18 day old $Brg1^{WT}$ and $Brg1^{Het}$ mice

were processed for ATAC-seq. Conditional deletions were generated using the Mvh-Cre

transgene. ATAC-seq libraries were made as previously reported (Buenrostro et al., 2013) , with

the exception of a double-sided SPRI bead size selection step of each library, using 0.5x and 1x

ratio of SPRI beads to obtain a library size range of ~150bp to ~2kb. All libraries were combined

and sequenced on a single lane of Illumina Hiseq 2500 using v4 chemistry (50bp reads, single).

Preparation of nuclear lysates:

Nuclear extracts were prepared from spermatocyte enriched preparations as previously described

(Chandler et al., 2012; Li et al., 1991) with minor modifications. These nuclear lysates were used

for co-immunoprecipitations (co-IP) and for the mass spectrometric analysis of BRG1

immunopulldowns (supplemental materials).

Identification of BRG1 interacting proteins by mass spectrometry:

Proteins isolated from BRG1 IP and IP with a non-specific rabbit IgG were run approximately 2 cm below the bottom of the well of a precast SDS polyacrylamide gel (short gel). The short gel was stained with GelCode blue protein stain (ThermoFisher) and the lanes containing each sample were cut out and analyzed by mass spectrometry at the University of Massachusetts Medical School Mass spectrometry Core Facility (see supplemental materials).

Preparation of sub cellular protein fractions:

Cytosolic, nucleoplasmic (soluble) and chromatin (insoluble) fractions were prepared as described (Méndez and Stillman, 2000) from spermatogenic cells obtained from P12 and P21 $Brg1^{WT}$,$Brg1^{Het}$ and $Brg1^{cKO}$ mice. Conditional deletions were generated using the $Mvh\text{-}Cre$ transgene.

Preparation of acid extracted histones:

Histones were extracted from spermatogenic cells obtained from P12 $Brg1^{WT}$,$Brg1^{Het}$ and $Brg1^{cKO}$ mice using acid extraction protocol described previously (Shechter et al., 2007). Conditional deletions were generated using the $Mvh\text{-}Cre$ transgene.

Western blotting:

Protein samples were separated by polyacrylamide gel electrophoresis and then transferred to PVDF (Polyvinylidene Difluoride) membranes (Bio-Rad) using wet/ semi-dry transfer apparatus (Bio-Rad). Western blots were generated using the Li-COR Bioscience Odyssey fluorescent western blotting reagents. All the antibodies used in this study and their corresponding dilutions are listed (Table S4).

767 **Acknowledgements**

768 We thank all members of the Magnuson laboratory for helpful comments on manuscript

769 preparation. This work was supported by National Institutes of Health grants

770 R01GM101974 and U42OD010924.

772
773 **Competing interests statement**

774 The authors declare no competing financial interests.

776 **Data availability:**
777 Raw and processed data have been deposited under GEO accession number GSE119179 (For

778 access, please enter token number: wjyxqkgozrgrpir).

References:

References:

Adams, S. R., Maezawa, S., Alavattam, K. G., Abe, H., Sakashita, A., Shroder, M., Broering, T.

J., Sroga Rios, J., Thomas, M. A., Lin, X., et al. (2018). RNF8 and SCML2 cooperate to

regulate ubiquitination and H3K27 acetylation for escape gene activation on the sex

chromosomes. *PLoS Genet.* **14**,.

Alexander, J. M., Hota, S. K., He, D., Thomas, S., Ho, L., Pennacchio, L. A. and Bruneau, B. G.

(2015). Brg1 modulates enhancer activation in mesoderm lineage commitment.

Development **142**, 1418–1430.

Alver, B. H., Kim, K. H., Lu, P., Wang, X., Manchester, H. E., Wang, W., Haswell, J. R., Park, P. J.

and Roberts, C. W. M. (2017). The SWI/SNF chromatin remodelling complex is required for

maintenance of lineage specific enhancers. *Nat. Commun.* **8**, 1–10.

Attanasio, C., Nord, A. S., Zhu, Y., Blow, M. J., Biddie, S. C., Mendenhall, E. M., Dixon, J.,

Wright, C., Hosseini, R., Akiyama, J. A., et al. (2014). Tissue-specific SMARCA4 binding at

active and repressed regulatory elements during embryogenesis. *Genome Res.* **24**, 920–

929.

Baker, C. L., Kajita, S., Walker, M., Saxl, R. L., Raghupathy, N., Choi, K., Petkov, P. M. and

Paigen, K. (2015). PRDM9 Drives Evolutionary Erosion of Hotspots in Mus musculus

through Haplotype-Specific Initiation of Meiotic Recombination. **11**,.

Ball, R. L., Fujiwara, Y., Sun, F., Hu, J., Hibbs, M. A., Handel, M. A. and Carter, G. W. (2016).

Regulatory complexity revealed by integrated cytological and RNA-seq analyses of meiotic

substages in mouse spermatocytes. *BMC Genomics* **17**, 1–17.

Basciani, S., Mariani, S., Arizzi, M., Ulisse, S., Rucci, N., Jannini, E. A., Rocca, C. Della,

Manicone, A., Carani, C., Spera, G., et al. (2002). Expression of Platelet-Derived Growth

Factor-A (PDGF-A), PDGF-B, and PDGF Receptor-α and -β during Human Testicular

Development and Disease. *J. Clin. Endocrinol. Metab.* **87**, 2310–2319.

Basciani, S., Mariani, S., Arizzi, M., Ulisse, S., Rucci, N., Jannini, E. A., Rocca, C. Della,

Manicone, A., Carani, C., Spera, G., et al. (2018). Expression of Platelet-Derived Growth

Factor-A. **87**, 2310–2319.

Bellve, a R., Cavicchia, J. C., Millette, C. F., O'Brien, D. a, Bhatnagar, Y. M. and Dym, M.

(1977). Spermatogenic cells of the prepubertal mouse. Isolation and morphological

characterization. *J Cell Biol* **74**, 68–85.

Bernstein, B. E., Mikkelsen, T. S., Xie, X., Kamal, M., Huebert, D. J., Cuff, J., Fry, B., Meissner,

A., Wernig, M., Plath, K., et al. (2006). A Bivalent Chromatin Structure Marks Key

Developmental Genes in Embryonic Stem Cells. *Cell* **125**, 315–326.

Blake, J. A., Richardson, J. E., Bult, C. J., Kadin, J. A., Eppig, J. T., Baldarelli, R. M., Beal, J. S.,

Bradt, D. W., Burkart, D. L., Butler, N. E., et al. (2003). MGD: The mouse genome

database. *Nucleic Acids Res.* **31**, 193–195.

Bray, N. L., Pimentel, H., Melsted, P. and Pachter, L. (2016). Near-optimal probabilistic RNA-

seq quantification. *Nat. Biotechnol.* **34**, 525–527.

Brick, K., Smagulova, F., Khil, P., Camerini-Otero, R. D. and Petukhova, G. V. (2012). Genetic

recombination is directed away from functional genomic elements in mice. *Nature* **485**,

642–645.

Brind'Amour, J., Liu, S., Hudson, M., Chen, C., Karimi, M. M. and Lorincz, M. C. (2015). An

ultra-low-input native ChIP-seq protocol for genome-wide profiling of rare cell

populations. *Nat. Commun.* **6**,.

Buaas, F. W., Kirsh, A. L., Sharma, M., McLean, D. J., Morris, J. L., Griswold, M. D., De Rooij, D.

G. and Braun, R. E. (2004). Plzf is required in adult male germ cells for stem cell self-

renewal. *Nat. Genet.* **36**, 647–652.

Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. and Greenleaf, W. J. (2013).

Transposition of native chromatin for fast and sensitive epigenomic profiling of open

chromatin, DNA-binding proteins and nucleosome position. *Nat. Methods* **10**, 1213–8.

Chandler, R. L., Brennan, J., Schisler, J. C., Serber, D., Patterson, C. and Magnuson, T. (2012).

ARID1a-DNA Interactions Are Required for Promoter Occupancy by SWI/SNF. *Mol. Cell.*

Biol. **33**, 265–280.

Chang, Y. F., Lee-Chang, J. S., Panneerdoss, S., MacLean, J. a. and Rao, M. K. (2011). Isolation

of Sertoli, Leydig, and spermatogenic cells from the mouse testis. *Biotechniques* **51**, 341–

344.

Clapier, C. R., Iwasa, J., Cairns, B. R. and Peterson, C. L. (2017). Mechanisms of action and

regulation of ATP-dependent chromatin-remodelling complexes. *Nat. Rev. Mol. Cell Biol.*

Creppe, C., Palau, A., Malinverni, R., Valero, V. and Buschbeck, M. (2014). A Cbx8-Containing

Polycomb Complex Facilitates the Transition to Gene Activation during ES Cell

Differentiation. *PLoS Genet.* **10**,.

Dann, C. T., Alvarado, A. L., Molyneux, L. A., Denard, B. S., Garbers, D. L. and Porteus, M. H.

(2008). Spermatogonial Stem Cell Self-Renewal Requires OCT4, a Factor Downregulated

During Retinoic Acid-Induced Differentiation. *Stem Cells* **26**, 2928–2937.

De Vries, F. A. T., De Boer, E., Van Den Bosch, M., Baarends, W. M., Ooms, M., Yuan, L., Liu, J.

G., Van Zeeland, A. A., Heyting, C. and Pastink, A. (2005). Mouse Sycp1 functions in

synaptonemal complex assembly, meiotic recombination, and XY body formation. *Genes*

Dev. **19**, 1376–1389.

Diagouraga, B., Clément, J. A. J., Duret, L., Kadlec, J., de Massy, B. and Baudat, F. (2018).

PRDM9 Methyltransferase Activity Is Essential for Meiotic DNA Double-Strand Break

Formation at Its Binding Sites. *Mol. Cell* **69**, 853–865.e6.

Eaker, S., Pyle, A., Cobb, J. and Handel, M. A. (2001). Evidence for meiotic spindle checkpoint

from analysis of spermatocytes from Robertsonian-chromosome heterozygous mice. *J. Cell*

Sci. **114**, 2953–2965.

Euskirchen, G. M., Auerbach, R. K., Davidov, E., Gianoulis, T. a., Zhong, G., Rozowsky, J.,

Bhardwaj, N., Gerstein, M. B. and Snyder, M. (2011). Diverse roles and interactions of the

SWI/SNF chromatin remodeling complex revealed using global approaches. *PLoS Genet.* **7**,.

Fierz, B., Chatterjee, C., McGinty, R. K., Bar-Dagan, M., Raleigh, D. P. and Muir, T. W. (2011).

Histone H2B ubiquitylation disrupts local and higher-order chromatin compaction. *Nat.*

Chem. Biol. **7**, 113–119.

Gallardo, T., Shirley, L., John, G. B. and Castrillon, D. H. (2007). Generation of a germ cell-

specific mouse transgenic Cre line, Vasa-Cre. *Genesis* **45**, 413–7.

Gallinari, P., Di Marco, S., Jones, P., Pallaoro, M. and Steinkühler, C. (2007). HDACs, histone

deacetylation and gene transcription: From molecular biology to cancer therapeutics. *Cell*

Res. **17**, 195–211.

Gao, Z., Lee, P., Stafford, J. M., Schimmelmann, M. Von, Schaefer, A. and Reinberg, D. (2014).

An AUTS2 – Polycomb complex activates gene expression in the CNS. *Nature* **516**, 349–

354.

Goertz, M. J., Wu, Z., Gallardo, T. D., Hamra, F. K. and Castrillon, D. H. (2011). Foxo1 is

required in mouse spermatogonial stem cells for their maintenance and the initiation of

spermatogenesis. *J. Clin. Invest.* **121**, 3456–3466.

Goetz, P., Chandley, a C. and Speed, R. M. (1984). Morphological and temporal sequence of

meiotic prophase development at puberty in the male mouse. *J. Cell Sci.* **65**, 249–263.

Green, C. D., Ma, Q., Manske, G. L., Shami, A. N., Zheng, X., Marini, S., Moritz, L., Sultan, C.,

Gurczynski, S. J., Moore, B. B., et al. (2018). A Comprehensive Roadmap of Murine

Spermatogenesis Defined by Single-Cell RNA-Seq. *Dev. Cell* **46**, 651–667.e10.

Hammoud, S. S., Low, D. H. P., Yi, C., Carrell, D. T., Guccione, E. and Cairns, B. R. (2014).

Chromatin and Transcription Transitions of Mammalian Adult Germline Stem Cells and

Spermatogenesis. *Cell Stem Cell* **15**, 239–253.

Handel, M. A. and Schimenti, J. C. (2010). Genetics of mammalian meiosis: regulation,

dynamics and impact on fertility. *Nat. Rev. Genet.* **11**, 124–136.

Hasegawa, K., Sin, H., Hasegawa, K., Sin, H., Maezawa, S., Broering, T. J., Kartashov, A. V and

Alavattam, K. G. (2015). SCML2 Establishes the Male Germline Epigenome through

Regulation of Histone H2A Ubiquitination Article SCML2 Establishes the Male Germline

Epigenome through Regulation of Histone H2A Ubiquitination. *Dev. Cell* **32**, 574–588.

Hayamizu, T. F., Wicks, M. N., Davidson, D. R., Burger, A., Ringwald, M. and Baldock, R. A.

(2013). Open Access EMAP / EMAPA ontology of mouse developmental anatomy : 2013

update. *J. Biomed. Semantics* 1–5.

Hayashi, K., Yoshida, K. and Matsui, Y. (2005). A histone H3 methyltransferase controls

epigenetic events required for meiotic prophase. *Nature* **438**, 374–378.

Ho, L. and Crabtree, G. R. (2010). Chromatin remodelling during development. *Nature* **463**,

474–484.

Ho, L., Ronan, J. L., Wu, J., Staahl, B. T., Chen, L., Kuo, A., Lessard, J., Nesvizhskii, A. I., Ranish,

J. and Crabtree, G. R. (2009). An embryonic stem cell chromatin remodeling complex,

esBAF, is essential for embryonic stem cell self-renewal and pluripotency. *Proc. Natl. Acad.*

Sci. U. S. A. **106**, 5181–5186.

Huether, R., Dong, L., Chen, X., Wu, G., Parker, M., Wei, L., Ma, J., Edmonson, M. N., Hedlund,

E. K., Rusch, M. C., et al. (2014). The landscape of somatic mutations in epigenetic

regulators across 1,000 paediatric cancer genomes. *Nat. Commun.* **5**, 1–7.

Ichijima, Y., Ichijima, M., Lou, Z., Nussenzweig, A., Daniel Camerini-Otero, R., Chen, J.,

Andreassen, P. R. and Namekawa, S. H. (2011). MDC1 directs chromosome-wide silencing

of the sex chromosomes in male germ cells. *Genes Dev.* **25**, 959–971.

Kadoch, C., Williams, R. T., Calarco, J. P., Miller, E. L., Weber, C. M., Braun, S. M. G., Pulice, J.

L., Chory, E. J. and Crabtree, G. R. (2016). Dynamics of BAF–Polycomb complex opposition

on heterochromatin in normal and oncogenic states. *Nat. Genet.* **49**, 213–222.

Kakarougkas, A., Ismail, A., Chambers, A., Riballo, E., Herbert, A., Künzel, J., Löbrich, M.,

Jeggo, P. and Downs, J. (2013). Requirement for PBAF in Transcriptional Repression and

Repair at DNA Breaks in Actively Transcribed Regions of Chromatin. *Mol. Cell* 1–10.

Kim, Y., Fedoriw, A. M. and Magnuson, T. (2012). An essential role for a mammalian SWI/SNF

chromatin-remodeling complex during male meiosis. *Development* **139**, 1133–1140.

Kubota, H. and Brinster, R. L. (2008). Culture of Rodent Spermatogonial Stem Cells, Male

Germline Stem Cells of the Postnatal Animal. *Methods Cell Biol.* **86**, 59–84.

Kwon, S.-J., Park, J.-H., Park, E.-J., Lee, S.-A., Lee, H.-S., Kang, S. W. and Kwon, J. (2015). ATM-

mediated phosphorylation of the chromatin remodeling enzyme BRG1 modulates DNA

double-strand break repair. *Oncogene* **34**, 303–313.

Lecona, E., Narendra, V. and Reinberg, D. (2015). USP7 Cooperates with SCML2 To Regulate

the Activity of PRC1. *Mol. Cell. Biol.* **35**, 1157–1168.

Lee, H.-S., Park, J.-H., Kim, S.-J., Kwon, S.-J. and Kwon, J. (2010). A cooperative activation loop

among SWI/SNF, gamma-H2AX and H3 acetylation for DNA double-strand break repair.

EMBO J. **29**, 1434–1445.

Lee, J., Ogushi, S., Saitou, M. and Hirano, T. (2011). Condensins I and II are essential for

construction of bivalent chromosomes in mouse oocytes. *Mol. Biol. Cell* **22**, 3465–3477.

Lesch, B. J., Dokshin, G. a, Young, R. a, McCarrey, J. R. and Page, D. C. (2013). A set of genes

critical to development is epigenetically poised in mouse germ cells from fetal stages

through completion of meiosis. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 16061–6.

Lesch, B. J., Silber, S. J., McCarrey, J. R. and Page, D. C. (2016). Parallel evolution of male

germline epigenetic poising and somatic development in animals. *Nat. Genet.* **48**, 1–10.

Lessard, J., Wu, J. I., Ranish, J. a., Wan, M., Winslow, M. M., Staahl, B. T., Wu, H., Aebersold,

R., Graef, I. a. and Crabtree, G. R. (2007). An Essential Switch in Subunit Composition of a

Chromatin Remodeling Complex during Neural Development. *Neuron* **55**, 201–215.

Li, Y. C., Ross, J., Scheppler, J. A. and Franza Jr., B. R. (1991). An in vitro transcription analysis

of early responses of the human immunodeficiency virus type 1 long terminal repeat to

different transcriptional activators. *Mol Cell Biol* **11**, 1883–1893.

Li, W., Wu, J., Kim, S.-Y., Zhao, M., Hearn, S. a, Zhang, M. Q., Meistrich, M. L. and Mills, A. a

(2014). Chd5 orchestrates chromatin remodelling during sperm development. *Nat.*

Commun. **5**, 3812.

Luo, M., Zhou, J., Leu, N. A., Abreu, C. M., Wang, J., Anguera, M. C., de Rooij, D. G., Jasin, M.

and Wang, P. J. (2015). Polycomb Protein SCML2 Associates with USP7 and Counteracts

Histone H2A Ubiquitination in the XY Chromatin during Male Meiosis. *PLoS Genet.* **11**, 1–

17.

Ma, P., Pan, H., Montgomery, R. L., Olson, E. N. and Schultz, R. M. (2012). Compensatory

functions of histone deacetylase 1 (HDAC1) and HDAC2 regulate transcription and

apoptosis during mouse oocyte development. *Proc. Natl. Acad. Sci.* **109**, E481–E489.

Maertens, G. N., El Messaoudi-Aubert, S., Elderkin, S., Hiom, K. and Peters, G. (2010).

Ubiquitin-specific proteases 7 and 11 modulate Polycomb regulation of the INK4a tumour

suppressor. *EMBO J.* **29**, 2553–2565.

Maezawa, S., Hasegawa, K., Yukawa, M., Sakashita, A., Alavattam, K. G., Andreassen, P. R.,

Vidal, M., Koseki, H., Barski, A. and Namekawa, S. H. (2017). Polycomb directs timely

activation of germline genes in spermatogenesis. *Genes Dev.* **31**, 1693–1703.

Maezawa, S., Hasegawa, K., Yukawa, M., Kubo, N. and Sakashita, A. (2018). Polycomb protein

SCML2 facilitates H3K27me3 to establish bivalent domains in the male germline [Genetics].

Pnas 3–8.

Margolin, G., Khil, P. P., Kim, J., Bellani, M. a and Camerini-Otero, R. D. (2014). Integrated

transcriptome analysis of mouse spermatogenesis. *BMC Genomics* **15**, 39.

Masliah-Planchon, J., Bièche, I., Guinebretière, J.-M., Bourdeaut, F. and Delattre, O. (2015).

SWI/SNF Chromatin Remodeling and Human Malignancies. *Annu. Rev. Pathol. Mech. Dis.*

10, 145–171.

Méndez, J. and Stillman, B. (2000). Chromatin association of human origin recognition

complex, cdc6, and minichromosome maintenance proteins during the cell cycle: assembly

of prereplication complexes in late mitosis. *Mol. Cell. Biol.* **20**, 8602–12.

Minsky, N., Shema, E., Field, Y., Schuster, M., Segal, E. and Oren, M. (2008).

Monoubiquitinated H2B is associated with the transcribed region of highly expressed

genes in human cells. *Nat. Cell Biol.* **10**, 483–488.

Montgomery, R. L., Hsieh, J., Barbosa, A. C., Richardson, J. A. and Olson, E. N. (2009). Histone

deacetylases 1 and 2 control the progression of neural precursors to neurons during brain

development. *Proc. Natl. Acad. Sci.* **106**, 7876–7881.

Motenko, H., Neuhauser, S. B., O'Keefe, M. and Richardson, J. E. (2015). MouseMine: a new

data warehouse for MGI. *Mamm. Genome* **26**, 325–330.

Mu, W., Starmer, J., Fedoriw, A. M., Yee, D. and Magnuson, T. (2014). Repression of the soma-

specific transcriptome by Polycomb-repressive complex 2 promotes male germ cell

development. *Genes Dev.* **28**, 2056–2069.

Nicassio, F., Corrado, N., Vissers, J. H. A., Areces, L. B., Bergink, S., Marteijn, J. A., Geverts, B.,

Houtsmuller, A. B., Vermeulen, W., Di Fiore, P. P., et al. (2007). Human USP3 Is a

Chromatin Modifier Required for S Phase Progression and Genome Stability. *Curr. Biol.* **17**,

1972–1977.

Oatley, M. J., Kaucher, A. V, Racicot, K. E. and Oatley, J. M. (2011). Inhibitor of DNA binding 4

is expressed selectively by single spermatogonia in the male germline and regulates the

self-renewal of spermatogonial stem cells in mice. *Biol. Reprod.* **85**, 347–356.

Ogiwara, H., Ui, A., Otsuka, A., Satoh, H., Yokomi, I., Nakajima, S., Yasui, A., Yokota, J. and

Kohno, T. (2011). Histone acetylation by CBP and p300 at double-strand break sites

facilitates SWI/SNF chromatin remodeling and the recruitment of non-homologous end

joining factors. *Oncogene* **30**, 2135–2146.

Pavri, R., Zhu, B., Li, G., Trojer, P., Mandal, S., Shilatifard, A. and Reinberg, D. (2006). Histone
H2B Monoubiquitination Functions Cooperatively with FACT to Regulate Elongation by
RNA Polymerase II. *Cell* **125**, 703–717.

Peters, A. H. F. M., Plug, A. W., Van Vugt, M. J. and De Boer, P. (1997). A drying-down
technique for the spreading of mammalian melocytes from the male and female germline.
Chromosom. Res. **5**, 66–68.

Qi, W., Wang, R., Chen, H., Wang, X., Xiao, T., Boldogh, I., Ba, X., Han, L. and Zeng, X. (2014).
BRG1 promotes the repair of DNA double-strand breaks by facilitating the replacement of
RPA with RAD51. *J. Cell Sci.* **128**, 317–330.

Raab, J. R., Resnick, S. and Magnuson, T. (2015). Genome-Wide Transcriptional Regulation
Mediated by Biochemically Distinct SWI/SNF Complexes. *PLoS Genet.* **11**, 1–26.

Raab, J. R., Runge, J. S., Spear, C. C. and Magnuson, T. (2017). Co-regulation of transcription by
BRG1 and BRM, two mutually exclusive SWI/SNF ATPase subunits. *Epigenetics and*
Chromatin **10**, 1–15.

Robinson, M. D., McCarthy, D. J. and Smyth, G. K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–40.

Royo, H., Prosser, H., Ruzankina, Y., Mahadevaiah, S. K., Cloutier, J. M., Baumann, M.,
Fukuda, T., Höög, C., Tóth, A., de Rooij, D. G., et al. (2013). ATR acts stage specifically to
regulate multiple aspects of mammalian meiotic silencing. *Genes Dev.* **27**, 1484–1494.

Sadate-Ngatchou, P. I., Payne, C. J., Dearth, A. T. and Braun, R. E. (2008). Cre recombinase
activity specific to postnatal, premeiotic male germ cells in transgenic mice. *Genesis* **46**,
738–742.

Schmahl, J., Rizzolo, K. and Soriano, P. (2008). The PDGF signaling pathway controls multiple
steroid-producing lineages. *Genes Dev.* **22**, 3255–3267.

Serber, D. W., Runge, J. S., Menon, D. U. and Magnuson, T. (2015). The Mouse INO80
Chromatin Remodeling Complex Is an Essential Meiotic Factor for Spermatogenesis. *Biol.*
Reprod.

Shechter, D., Dormann, H. L., Allis, C. D. and Hake, S. B. (2007). Extraction, purification and
analysis of histones. *Nat. Protoc.* **2**, 1445–1457.

Soneson, C., Love, M. I. and Robinson, M. D. (2016). Differential analyses for RNA-seq:

transcript-level estimates improve gene-level inferences. *F1000Research* **4**, 1521.

Stanton, B. Z., Hodges, C., Calarco, J. P., Braun, S. M. G., Ku, W. L., Kadoch, C., Zhao, K. and

Crabtree, G. R. (2016). Smarca4 ATPase mutations disrupt direct eviction of PRC1 from

chromatin. *Nat. Genet.* **49**, 282–288.

Sumi-Ichinose, C., Ichinose, H., Metzger, D. and Chambon, P. (1997). SNF2beta-BRG1 is

essential for the viability of F9 murine embryonal carcinoma cells. *Mol. Cell. Biol.* **17**, 5976–

5986.

Tachibana, M., Nozaki, M., Takeda, N. and Shinkai, Y. (2007). Functional dynamics of H3K9

methylation during meiotic prophase progression. *EMBO J.* **26**, 3346–3359.

Takada, Y., Naruse, C., Costa, Y., Shirakawa, T., Tachibana, M., Sharif, J., Kezuka-Shiotani, F.,

Kakiuchi, D., Masumoto, H., Shinkai, Y. -i., et al. (2011). HP1 links histone methylation

marks to meiotic synapsis in mice. *Development* **138**, 4207–4217.

Tolstorukov, M. Y., Sansam, C. G., Lu, P., Koellhoffer, E. C., Helming, K. C., Alver, B. H., Tillman,

E. J., Evans, J. a, Wilson, B. G., Park, P. J., et al. (2013). Swi/Snf chromatin

remodeling/tumor suppressor complex establishes nucleosome occupancy at target

promoters. *Proc. Natl. Acad. Sci.* **110**, 10165–10170.

Turner, J. M. a. (2007). Meiotic sex chromosome inactivation. *Development* **134**, 1823–1831.

Turner, J. M. A., Aprelikova, O., Xu, X., Wang, R., Kim, S., Chandramouli, G. V. R., Barrett, J. C.,

Burgoyne, P. S. and Deng, C. X. (2004). BRCA1, histone H2AX phosphorylation, and male

meiotic sex chromosome inactivation. *Curr. Biol.* **14**, 2135–2142.

Van Der Knaap, J. A., Kumar, B. R. P., Moshkin, Y. M., Langenberg, K., Krijgsveld, J., Heck, A. J.

R., Karch, F. and Verrijzer, C. P. (2005). GMP synthetase stimulates histone H2B

deubiquitylation by the epigenetic silencer USP7. *Mol. Cell* **17**, 695–707.

Wang, H., Wang, L., Erdjument-Bromage, H., Vidal, M., Tempst, P., Jones, R. S. and Zhang, Y.

(2004). Role of histone H2A ubiquitination in Polycomb silencing. *Nature* **431**, 873–878.

Wang, J., Gu, H., Lin, H. and Chi, T. (2012). Essential Roles of the Chromatin Remodeling Factor

Brg1 in Spermatogenesis in Mice. *Biol. Reprod.* **86**, 186–186.

Watanabe, R., Ui, A., Kanno, S. I., Ogiwara, H., Nagase, T., Kohno, T. and Yasui, A. (2014).

SWI/SNF factors required for cellular resistance to dna damage include arid1a and arid1b

and show interdependent protein stability. *Cancer Res.* **74**, 2465–2475.

Wilson, B. G., Wang, X., Shen, X., McKenna, E. S., Lemieux, M. E., Cho, Y. J., Koellhoffer, E. C.,

Pomeroy, S. L., Orkin, S. H. and Roberts, C. W. M. (2010). Epigenetic antagonism between

polycomb and SWI/SNF complexes during oncogenic transformation. *Cancer Cell* **18**, 316–

328.

Wojtasz, L., Daniel, K., Roig, I., Bolcun-Filas, E., Xu, H., Boonsanay, V., Eckmann, C. R., Cooke,

H. J., Jasin, M., Keeney, S., et al. (2009). Mouse HORMAD1 and HORMAD2, two conserved

meiotic chromosomal proteins, are depleted from synapsed chromosome axes with the

help of TRIP13 AAA-ATPase. *PLoS Genet.* **5**,.

Wu, X., Oatley, J. M., Oatley, M. J., Kaucher, A. V., Avarbock, M. R. and Brinster, R. L. (2010).

The POU Domain Transcription Factor POU3F1 Is an Important Intrinsic Regulator of GDNF-

Induced Survival and Self-Renewal of Mouse Spermatogonial Stem Cells1. *Biol. Reprod.* **82**,

1103–1111.

Yamaguchi, T., Cubizolles, F., Zhang, Y., Reichert, N., Kohler, H., Seiser, C. and Matthias, P.

(2010). Histone deacetylases 1 and 2 act in concert to promote the G1-to-S progression.

Genes Dev. **24**, 455–469.

Yang, F., De La Fuente, R., Leu, N. A., Baumann, C., McLaughlin, K. J. and Wang, P. J. (2006).

Mouse SYCP2 is required for synaptonemal complex assembly and chromosomal synapsis

during male meiosis. *J. Cell Biol.* **173**, 497–507.

Yu, G., Wang, L.-G., Han, Y. and He, Q.-Y. (2012). clusterProfiler: an R Package for Comparing

Biological Themes Among Gene Clusters. *Omi. A J. Integr. Biol.* **16**, 284–287.

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nussbaum, C.,

Myers, R. M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS).

Genome Biol. **9**,.

1067

1068

 Figure Legends:

 Figure 1. BRG1 is enriched at transcriptionally active and poised regions. (A) Heatmaps with K-
 means clustering depicting the relative enrichment of BRG1, H3K4me3 and H3K27me3 across
 RefSeq gene TSS ± 4 Kb in P12, P17 and P18 testes. TSS: Transcription start site, Cl-1:Class 1,
 Cl-2: Class 2, Cl-3: Class 3. Number (n) of TSS's sites within each cluster are indicated (B)
 Metaplots depicting the relative enrichment of BRG1 at P12 and P18 (top row), H3K4me3
 (middle row) and H3K27me3 (bottom row), at P12 and P17, at TSS ± 4 Kb, associated with
 P12 BRG1 ChIP-seq peaks categorized by the testes temporal expression profile of cognate
 genes. (C) P12 and P18 BRG1 enrichment across an 8 Kb window centered at TSS's of X -
 linked genes.

 Figure 2. BRG1 influences transcription during spermatogenesis. (A) Smear plot showing the
 Log2 fold change in transcript abundance (y-axis) of genes in the P12 $Brg1^{cKO}$ relative to $Brg1^{WT}$
 and Log2 transcript abundance in counts per million (CPM; x-axis). Red dots denote
 significantly mis-expressed genes (FDR≤0.05). The horizontal blue lines demarcate the 2-fold
 change in transcript abundance. (B) Table listing enriched anatomy and gene ontology terms
 associated with genes differentially regulated by BRG1. Benjamini – Hochberg, adjusted p
 values are indicated within parenthesis next to the corresponding ontology term. (C) Heatmap
 depicting the change in transcript abundance of genes associated with arrest of male meiosis
 (MP:0008261) and abnormal spermatogonia proliferation (MP:0002685) phenotype ontologies
 (labeled along vertical axis), across individual P12 Brg^{WT} (WT1-WT4) and Brg^{cKO} (KO1-KO4)
 replicates (horizontal axis). Transcript abundances are represented as z- scores. (D) P10 Brg^{WT}
 and $Brg1^{cKO}$ testes cryosections, immunofluorescently labeled for ZBTB16 (green) and counter
 stained with DAPI (blue). The average number of ZBTB16$^+$ cells (type A spermatogonia; SpgA)
 ± standard error of mean (SEM) per tubule are indicated below each representative image.
 Images were captured using a 25x objective Scale bar: 20 μm. (E) Western blot showing the
 abundance of SYCP2 in sub-cellular fractions obtained from P21 $Brg1^{WT}$, Brg^{Het} and $Brg1^{cKO}$
 spermatogenic cells. Histone- H3 was used as a nuclear loading control. (F) Brg^{WT} and $Brg1^{cKO}$
 zygotene spermatocytes immunofluorescently labeled for SYCP2 (red) and γH2Ax (green). The
 percentage of $Brg1^{cKO}$ zygotene spermatocytes (n=212) displaying abnormal SYCP2 filaments
 is indicated. Images were captured using a 100x objective Scale bar: 10 μm.

1101

Figure 3. BRG1 directly regulates chromatin accessibility at promoters. (A) Bar plot depicting

the Log2 values of normalized ATAC-seq read coverage with added pseudocount (y-axis) at the

promoters (TSS± 0.5Kb) of BRG1 activated and repressed genes (x-axis) in P12 *Brg1^wt^* (red)

and *Brg1^cKO^* (blue) spermatogenic cells. *** : Significant p-value (p<0.001); NS: not significant,

as calculated by Wilcoxon rank sum test (B) Smear plot depicting the Log2 fold change in

chromatin accessibility (y-axis) and Log2 ATAC read abundance in counts per million (CPM; x-

axis) in the P12 *Brg1^cKO^* relative to *Brg1^WT^*. Red dots denote regions displaying significant

differences in chromatin accessibility (FDR≤0.05). Horizontal blue lines denote a 2- fold change

in read abundance. Closed: sites less accessible; Opened: sites more accessible, in the P12

*Brg1^cKO^* relative to *Brg1^WT^* (C) Genomic associations of closed and opened sites. Total number

either class of sites are indicated within parenthesis (D) Gene ontology analysis of closed and

opened sites. Benjamini – Hochberg, adjusted p values are indicated.

1114

1115

Figure 4. BRG1 physically interacts with SCML2 to regulate gene expression. (A) Silver stained

gel (left) and table (right) summarizing proteins identified by performing mass spectrometry(MS)

on BRG1 Immuno-pulldowns (IP) obtained from nuclear lysates which were prepared from 3-

week-old wild type testes. Peptide counts for proteins detected in IgG pulldown are indicated

within parenthesis (B) SCML2 Co-IP . Nonspecific (ns) IgG was used as negative control.

Lysates were treated with 25 U of universal nuclease (benzonase) prior to IP. Red arrowheads

label proteins detected in IP. Lane numbers are indicated at bottom (C) Metaplots depicting

BRG1 enrichment at P12 (green) and P18 (orange) over at TSS ± 2 Kb, associated with SCML2

activated (pach *Scml2* Act) and repressed (pach *Scml2* Rep) genes in pachytene

spermatocytes (D) Quadrant plot depicting the Log2 fold change in expression (KO/WT),

associated with genes commonly regulated by BRG1 (y-axis) and SCML2 (x-axis). Genes

displaying concordant and discordant expression changes are denoted by green and orange

dots respectively. The number of concordantly expressed genes (n) are indicated. *r²* values

were derived from pearson's correlation test.

1130

1131

Figure 5. BRG1 influences the localization of SCML2 to the sex body. P21 *Brg1^WT^* and *Brg1^cKO^*

cryosections, immunofluorescently labeled for (A) SCML2 (green) and γH2Ax (red), (B) USP7

(green) and γH2Ax (red). DNA (blue) was stained using DAPI. Arrowheads denote the sex

body labeled with γH2Ax. Magnified image of pachytene spermatocytes are shown in panel

insets (Pa: Pachynema). All images were captured using a 63x objective. Scale bar: 20 µm. (C)

Pachytene spermatocytes spreads from *Brg1^{WT}* and *Brg1^{cKO}* testes, immunofluorescently

labeled for SYCP3 (red) and markers of meiotic sex chromosome inactivation (green) -

BRCA1(left-most), ATR (middle) and MDC1 (right-most) . Magnified image of the sex

chromosomes are shown in panel insets. Images were captured using a 100x objective. Scale

bar: 20 µm.

1142

1143

Figure 6. BRG1 regulates H2AK119ub1 and H3K27ac during spermatogenesis. (A) Western

blots depicting H2AK119ub1, H3K27ac, H3K4me3 and H3K27me3, abundance in acid

extracted histones obtained from P12 *Brg^{WT}*, *Brg^{Het}* and *Brg1^{cKO}* testes. H3 was used as a

loading control. (B) Log2 fold change in H2AK119ub1 (green box) and H3K27ac (Orange box)

at BRG1 occupied promoters (TSS± 0.5Kb) categorized by chromosomal location (top panel)

and transcriptional status (bottom panel) in P12 *Brg1^{cKO}* relative to *Brg1^{WT}*. *** :Significant p-

value (p<0.001); NS: not significant, as calculated by Wilcoxon rank sum test. (C) The

enrichment of H3K27ac (left) and BRG1 (right) at lost, gained and common H3K27ac peaks

identified by a differential analysis of ChIP-seq peaks between P12 *Brg1^{WT}* and *Brg1^{cKO}*

spermatogenic cells. (D) Western blot depicting the abundance (top panel) and quantification

(bottom) of BRG1, RNF2 , USP3 and HDAC1 in nuclear fractions obtained from P12 *Brg^{WT}* ,

Brg^{Het} and *Brg1^{cKO}* spermatogenic cells. Bar plot represents the mean of protein band intensities

in the insoluble nuclear fractions determined from at least two independent trials. Protein levels

are relative to the wild type control and normalized to H3. Error bars represent the standard

error of mean (SEM). * denotes a significant students t-test p value.

1159

Figure 7. Model describing the role of SWI/SNF in spermatogenic gene regulation. (A) In the

developing male germ line BRG1 activates genes essential for the maintenance of pluripotent

spermatogonial precursors and several meiotic factors during meiosis, while repressing pre-

meiotic and somatic genes in spermatocytes. (B) During meiosis BRG1 activated gene

promoters (meiotic genes) display robust H3K4me3 enrichment while repressed gene

promoters display hallmarks of bivalent chromatin (H3K4me3 and H3K27me3). BRG1 is

required to maintain promoter accessibility and suppresses H2AK119ub1 while maintaining a

relatively higher level of H3K27ac at target genes. We propose that the repression of somatic

and pre-meiotic genes is achieved through the activity of SCML2, a PRC1 factor that is testes

specific. BRG1 exclusively interacts with SCML2 and influences its localization during meiosis.

As a consequence of this interaction, BRG1 also influences the localization of USP7, a factor

known to interact with SCML2.

1172

1173

A

C

B

A

B

	Anatomy Ontology (adj. p value)	Gene Ontology (adj. p value)
Down regulated	Brain (0.0049)	Meiotic cell cycle (2.292e-12)
	Central nervous system (0.0051)	meiotic nuclear division (4.727e-11)
	Gonad (0.0052)	chromosome segregation (1.747e-10)
	Molar (0.0057)	chromosome organization (2.756e-08)
Up regulated	Limb (4.077e-0.9)	Regulation of cell morphogenesis (0.00013)
	Mesenchyme (5.17e-0.9)	Axonogenesis (0.00013)
	Musculoskeletal system (5.27e-0.9)	Developmental cell growth (0.00019)
	Oral region (9.21e-0.9)	Regulation of actib filament-based process (0.00019)

C

D

E

F

A

Promoter accesibility

Log$_2$(Normalized coverage + 1)

*** NS

P12 *Brg1^{WT}*
P12 *Brg1^{cKO}*

Activated Repressed

P12 BRG1 targets

C

Genomic Associations

% distribution

Closed (n= 549)   Opened (n= 99)

UTR
Intergenic
Exon
Intron
Promoter (TSS±1Kb)

B

Differential analysis of accessibility (P12)

Log$_2$FC (*Brg1^{cKO}*/*Brg1^{WT}*)

● FDR≤0.05
● FDR≥0.05

Opened   Closed

log$_2$CPM (Read Count)

D

	Gene Ontology (Biological Process)	Adj.p value	Candidate Genes
Closed	Meiotic cell cycle	0.0003	*Top6bl, Tex19.1, Brca2*
	DNA repair	0.0009	*Yy1, Ube2n, Fanca, Mus81*
	Chromosome organization	0.0009	*Stag3, Phf13, Ercc3, Ccne2*
Opened	Cardiac muscle tissue development	0.06	*Angpt1,Zfpm2, Pln*
	Fat cell differentiation	0.06	*Jag1, Fndc3b, Igf1*
	Cytoplasm organization	0.06	*Zmiz1,Etv6*

A

	Protein	# Unique Peptide	% Coverage
Core	BRG1	63	45
	SNF5	24	70
	BAF155	55	61
	BAF170	49	52
PBAF	ARID2	46	38
	PBRM1	43	35
	BRD7	17	44
BAF	ARID1A	68	42
	ARID1B	45	26
	BRD9	12	33
	BCL7A	5	41
	BCL7B	5	41
	BCL7C	5	46
PRC1	SCML2	11(14)	16(17)

BRG1-IP   IgG

250 kDa
150 kDa
100 kDa
75 kDa
50 kDa
37 kDa
25 kDa
15 kDa

3 week old testes

B

+ Univ.Nuclease (25U)

20% Inp   SCML2 IP   IgG

BRG1
SCML2 (1D12)
RNF2

1   2   3

C

BRG1 Enrichment

pach *Scml2* Act   pach *Scml2* Rep

$\log_2$ (BRG1/Input)

3.0
2.5
2.0
1.5
1.0

-2kb   TSS   +2kb   -2kb   TSS   +2kb

━━ P12 BRG1   ━━ P18 BRG1

D

Transcript Abundance

Log_2 FC ($Brg1^{KO}/Brg1^{WT}$)

n=388

n=209

-1

-2

-3

-2   0   2   4

Log_2 FC ($Scml2^{KO}/Scml2^{WT}$)

● Concordant (r2=0.72)
● Discordant (r^2=0.51)

A

BRG1 co-ordinates spermatogenic gene expression

B

BRG1 collaborates with SCML2 to regulate meiotic transcription

Supplemental Information

Mammalian SWI/SNF collaborates with a polycomb-associated protein to regulate male germ line transcription in the mouse

Debashish U. Menon, Yoichiro Shibata, Weipeng Mu and Terry Magnuson

Department of Genetics, and Lineberger Comprehensive Cancer Center, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, NC 27599-7264, USA
*Correspondence: trm4@med.unc.edu, Tel 919-962-1319, Fax 919-843-4682

List of supplemental information:

Supplemental figures S1-S10

Figure S1. Features of BRG1 genomic associations.

Figure S2. Transcriptional response to the loss of BRG1 in the male germ line.

Figure S3. BRG1 directed changes in chromatin accessibility

Figure S4. Summary and validation of BRG1 interactions.

Figure S5. Features of BRG1 association with genes differentially regulated by SCML2.

Figure S6. Validation of the role of BRG1 in the recruitment of SCML2 to the sex body.

Figure S7. Analysis of SCML2 localization in meiotic spreads

Figure S8. Temporal regulation of H2AK119ub1 in testes and validation of anti- H2AK119ub1 (clone E6C5).

Figure S9. Genomic associations of lost, gained and common H3K27ac peaks.

Figure S10. BRG1 regulates the expression of epigenetic modifiers of H2AK119ub1, H2BK120ub1 and H3K27ac.

Supplemental tables

Table S1. List of genes differentially regulated by BRG1 at P12 (Excel sheet).

Table S2. List of regions displaying significant differences in chromatin accessibility in the $Brg1^{cKO}$ relative to $Brg1^{WT}$ (Excel sheet).

Table S3. List of BRG1 ChIP-seq peak calls and the genomic location of sites differentially bound by BRG1 between P12 and P18 testes (Excel sheet).

Table S4. List of antibodies used in this study

Table S5. Sequences of qRT-PCR primers used in this study

Supplemental figure legends

Figure S1. Features of BRG1 genomic associations. (A) Annotation of regions associated with P12 and P18 BRG1 peaks. Number (n) of P12 and P18 peaks are indicated within parenthesis. (B) Distribution of temporal expression profile of genes associated with class 1-3 TSS's. PMei: pre-meiotic, Mei: Meiotic, Const: constant. (C) UCSC genome browser views depicting H3K4me3 and H3K27me3 peak associations with promoters of candidate BRG1 target genes. Thick black bars denote BRG1 Macs2 peak calls. (D) P12 (green line) and P18 (orange line) BRG1 enrichment across an 8 Kb window centered at PRDM9 peaks (purple line).

Figure S2. Transcriptional response to the loss of BRG1 in the male germ line. (A) Quantification of spermatocyte populations per tubule (left panel) staged by γH2Ax immunostaining of Brg^{WT} and $Brg1^{cKO}$ cryosections obtained from P10 , P13 and P14 testes. The total number of tubules analyzed (n) at each stage is indicated. Abundance (right panel) of spermatogonial and meiotic sub stage specific protein-coding genes (x-axis) between P12 Brg^{WT} (red box) and $Brg1^{cKO}$ (blue boxes). Transcript abundance is expressed as the Log2 value of counts per million (CPM) added with a pseudo count (y–axis). (B) Quantitative RT-PCR analysis to determine the transcript abundance (y-axis) of candidate stem cell factors (x-axis) in Brg^{Het} and $Brg1^{cKO}$ relative to Brg^{WT} spermatogonia (THY1$^+$). The transcript abundance of candidate factors was normalized to genes whose transcript levels are constantly expressed (*Sdha* and *Ywhaz*). * denotes a p-value < 0.05, calculated using an unpaired students t-test. (C) Western blot showing the abundance of SYCP2 in sub-cellular fractions obtained from P12 Brg^{Het} and $Brg1^{cKO}$ spermatogenic cells. Histone- H3 was used as a nuclear loading control. (D) Brg^{WT} and $Brg1^{cKO}$ zygotene spermatocytes immunofluorescently labeled for SYCP2 (red) and SYCP3 (green). Images were captured using a 100x objective Scale bar: 10 μm. (E) Western blot showing the abundance of PDGFRA in cytosolic fractions obtained from P12, P14, P21 Brg^{WT}, Brg^{Het} and $Brg1^{cKO}$ spermatogenic cells. α TUBULIN was used as a loading control.

Figure S3. BRG1 directed changes in chromatin accessibility (A) Metaplots (top) and corresponding heatmaps (bottom) depicting the pairwise comparisons of normalized ATAC-seq signal at RefSeq genes ± 2Kb, between P12 $Brg1^{WT}$ and $Brg1^{cKO}$ spermatogenic cells, P18 $Brg1^{WT}$ and $Brg1^{Het}$ ($Brg1^{fl/Δ}$) spermatogenic cells. TSS: Transcription start site, TES: Transcription end site. (B) UCSC browser view of candidate closed and opened regions.

Figure S4. Summary and validation of BRG1 interactions. (A) List of known SWI/SNF subunits identified by IP-MS that are common to BAF and PBAF subcomplexes. (B) Validation of SCML2 antibody (1D12) specificity on nuclear extracts prepared from testes, ovaries and purified spermatogenic cells. Nucleolin (NCL) was used as a nuclear marker. (C) Co-immunoprecipitation of BRG1 and USP7. Nonspecific IgG was used as negative control. All lysates were treated with 25 U of universal nuclease (benzonase) prior to IP.

Figure S5. Features of BRG1 association with genes differentially regulated by SCML2. (A) ATAC seq coverage from P12 *Brg1^WT^* and *Brg1^cKO^* at TSS ± 4 Kb, associated with genes regulated by SCML2 in pachytene spermatocytes. (B) Enriched gene ontology terms associated with genes concordantly activated or repressed by BRG1 (P12) and SCML2 (pachynema).

Figure S6. Validation of the role of BRG1 in the recruitment of SCML2 to the sex body. Cryosections prepared from (A) P14 and P21 *Brg1^WT^* and *Brg1^cKO^* testes, immunofluorescently labeled for SCML2 (cyan), BRG1(red) and ATR (green), (B) P14 *Brg1^WT^* and *Brg1^Het^* testes, immunofluorescently labeled for SCML2 (green) and γH2Ax (red), DNA was stained with DAPI. Arrowheads label pachytene spermatocytes with completely formed sex body. Images were captured using a 63x objective. Scale bar: 20 μm.

Figure S7. Analysis of SCML2 localization in meiotic spreads. Pachytene spermatocyte from *Mvh-Cre* (left panel) and *Stra8-Cre* (right panel) induced *Brg1^cKO^* and *Brg1^WT^* testes, immunofluorescently labeled for SCML2 (green) and SYCP3 (red). Arrowheads denote the sex chromosomes and dotted circle outlines the SCML2 signal around the sex body. Images were captured using a 100x objective. Scale bar: 20 μm.

Figure S8. Temporal regulation of H2AK119ub1 in testes and validation of anti- H2AK119ub1 (clone E6C5) . (A) Western blot depicting H2AK119ub1, H3K27ac abundance in acid extracted histones obtained from P8,P10 *Brg^WT^*, *Brg^Het^* and *Brg1^cKO^* testes. H3 was used as a loading control (B) Western blot showing H2AK119ub1 and H3 abundance in acid extracted histones obtained from testes of P12 wild type mice carrying a *Mvh-Cre* transgene and their littermate controls (+/+). (C) Validation of anti-H2AK119ub1 (clone E6C5) specificity by immunoblotting for total H2AK119ub1 in acid extracted histones obtained from *Rnf2^WT^* and two different *Rnf2^KO^* ES cell lines. Blots were stained with Ponceau S to show total histone levels. (D) Western blot using anti-H2AK119ub1 (clone E6C5), to show H2AK119ub1 (green) levels in sub-cellular fractions of spermatogenic cells obtained from P12 *Brg^WT^*, *Brg^Het^* and *Brg1^cKO^* testes. H3 (red) was used as nuclear loading control. Dotted rectangle labels the nuclear levels of H2AK119ub1 levels. L: ladder.

Figure S9. Genomic associations of lost, gained and common H3K27ac peaks.

Figure S10. BRG1 regulates the expression of epigenetic modifiers of H2AK119ub1, H2BK120ub1 and H3K27ac. (A) UCSC browser view depicting BRG1 occupancy at *Rnf2*, *Usp3*, *Hdac1* and *Hdac2* promoters. Thick black bars label BRG1 peaks. (B) Transcript abundance (y-axis) of *Rnf2*, *Usp3*, *Hdac1* and *Hdac2* in P12 *Brg1^WT^* (red box) and *Brg1^cKO^* (blue box). Transcript abundance is expressed as the Log2 value of normalized abundance in counts per million (CPM) added with a pseudo count. (C) Western blot depicting H2BK120ub1 levels in acid extracted histones obtained from P12 *Brg^WT^* , *Brg^Het^* and *Brg1^cKO^* testes. H3 was used as a loading control.

Antibody	Vendor (catalog #)	Application
Rabbit anti-BRG1	Abcam (ab110641)	ChIP-seq (10 µg) IF (1:500)
Rabbit anti- YH2Ax	Cell signaling (9718)	IF (1:1000)
Mouse anti- YH2Ax	Millipore (05-636)	IF (1:1000)
Mouse anti-SYCP3	Abcam (ab97672)	IF (1:500)
Rabbit anti-SYCP3	Abcam (ab154255)	IF (1:500) WB (1:1000)
Rabbit anti-SYCP2	Millipore (ABE2622)	IF (1:500) WB (1:1000)
Goat anti- ATR (N-19)	Santa Cruz Biotech (sc-1887)	IF (1:50)
Mouse anti-SCML2 (1D12)	Developmental Studies Hybridoma Bank (PCRP-SCML2-1D12)	IF (1:10) IP (10 µg)
Rat anti- RNAPII subunit B1 (phosphor CTD Ser-2)	Millipore (04-1571)	IF (1:200)
Mouse anti- H2AK119ub1 (E6C5)	Millipore (05-678)	ChIP-seq (10 µg) WB (1:1000)
Rabbit anti- H3K27ac	Active Motif (39133)	ChIP-seq (5 µg) WB (1:1000)
Rabbit anti- H2BK120ub1	Cell signaling (5546)	WB (1:1000)
Mouse anti- H3K27me3	Abcam (ab6002)	WB (1:1000)
Rabbit anti– H3K4me3	Abcam (ab1012)	WB (1:1000)
Rabbit anti- H3	Abcam (ab1791)	WB (1:5000)
Mouse anti- RNF2	Santa Cruz Biotech (sc-1887)	WB (1:1000)
Mouse anti- USP3	GeneTex (GTX128238)	WB (1:1000)
Mouse anti- USP7	Santa Cruz Biotech (sc-1887)	IF (1:100) IP (2 µg) WB (1:1000)
Mouse anti- HDAC1	Abcam (ab7028)	WB (1:1000)
Rabbit anti- PDGFRA	Cell signaling (3164)	WB (1:1000)
Mouse anti α TUBULIN	Developmental Studies Hybridoma Bank (12G10)	WB (1:1000)
Rabbit anti NCL (Nucleolin)	Bethyl (A300-711A)	WB (1:5000)

Table. S4. List of antibodies used in this study

Gene Symbol	Forward Primer (5' - 3')	Reverse Primer (5' - 3')
Zbtb16	TATCTCGAAGCATTCCAGCGAGGA	ACTCATGGCTGAGAGACCGAAAGA
Id4	CCCGCGCCACCTCTCCAC	CAGAGAATGCTGTCA CCCTG
Pou3f1	TGGGGGCTGTCACTTTATTC	GGAGTTAGAAGGACCCCAGG
Pou5f1	AGCTGCTGAAGCAGAAGAGG	GTGGTCTGGCTGAACACCTT
Foxo1	AAGAGCGTGCCCTACTTCAA	TGCTGTGAAGGGACAGATTG
Sdha	TGGACCTTGTAGTCTTTGGCA	AACCGATTCTTCTCCAGCATT
Ywhaz	GAGAAAAAGCAGCAGATGGC	CTTTCTGGTTGCGAAGCATT
Brg1	GAAGACCATCCAGACCATCG	TTCATACGCCCAGTTTGACA

Table. S5. Sequences of qRT-PCR primer used in this study

Supplemental Materials

ChIP from low input chromatin:

Spermatogenic cells obtained from 12-day-old *Brg1^WT* and *Brg1^cKO* mice were fixed as described (Raab et al., 2015). Frozen pellets (10^6 cells each) were thawed on ice and then resuspended in 50 µl of nuclear isolation buffer (Sigma NUC-101). For the H3K27ac ChIPs, 5mM sodium butyrate was added to the nuclear suspensions. Samples were mixed by pipetting the cell suspension 15-20 times. 50 µl of nuclear preparations were mixed with 10 µl of MNase digestion buffer [6x MNase buffer (NEB), 8.8 mM DTT, 12 gel units MNase (NEB)] and incubated at 37 C for 15 minutes. The digestion was stopped by adding 1/10^th volume of 100 µM EDTA (Ethylenediaminetetraacetic acid). 6.6 µl of a 1% Triton X-100/1% sodium deoxycholate solution was added to the digested chromatin and vortexed gently for 30 seconds after which the samples were held on ice until the next step. In the meantime the DNA content of each chromatin preparation was quantified from 5 ul of chromatin. Briefly, chromatin was mixed with 95 µl H_2O and 100 µl 10% chelex-100 (Bio-Rad) and incubated at 95 C for 10 min. Following RNaseA digestion for 15 min at 37 C, ProteinaseK digestion for 1 hr at 56 C, the DNA was purified using ChIP-DNA clean and concentrator kit (Zymo) and quantified on a qubit. This way we ensured that for each pairwise comparison (wild type versus mutant ChIP), we began with an equal amount of input. We started with 145 ng and 285 ng of input DNA for each H3K27ac and H2AK119ub1 ChIP respectively. Digested chromatin was mixed with complete immnuoprecipitation (IP) buffer (20mM Tris-HCl pH 8.0, 2mM EDTA, 150 mM NaCl, 0.1% Triton X-100, 1x Protease inhibitor cocktail -PIC, 1mM Phenylmethanesulfonyl fluoride -PMSF) such that it comprised less than 25 % of the total volume. Chromatin was pre-cleared with 10 ul of magnetic Protein A (Bio-Rad) or G (Invitrogen/Dynalbeads) beads for 1 hour on a rotator at 4 C. After pre-clearing, 10% chromatin was set aside as input. Pre-cleared lysates were then mixed with antibodies to perform the ChIP. 5µg of rabbit anti-H3K27ac (Active Motif – 39133) conjugated to Protein A beads (Bio-Rad) and 10 µg of mouse anti-H2AK119ub1 IgM antibodies (Millipore E6C5, 05-678) were added to the lysates and left to bind chromatin by rotating the tubes overnight at 4 C. The following day anti-mouse IgM (Millipore, 12-488) conjugated to proteinA/G beads were added to the H2AK119ub1 ChIP samples and rotated at 4 C for 3 hours to capture anti-H2AK119ub1 bound chromatin. Chromatin bound to bead-antibody conjugates from all ChIP samples were isolated using a magnetic separator. Beads were then washed twice in low salt buffer followed by two washes in high salt buffer and eventually re-suspended in 100 µl of elution buffer (1%SDS/100mM $NaHCO_3$) in a 1.5 ml eppendorf tube. DNA was

eluted at 65 C on a shaking incubator (Eppendorf) at 800 rpm for 30 min. Eluate was separated from beads on a magnetic separator. 5 µl of 5M NaCl was added to eluate and incubated in 65 C water bath overnight to reverse crosslinks. ChIP DNA was digested with RNAseA for 30 min at 37 C followed by ProteinaseK digestion for 1 hr at 56 C and then purified using a ChIP-DNA clean and concentrator kit (Zymo) and quantified with a qubit. Libraries were prepared from ChIP and Input samples using the Kapa Hyperprep kit.

ChIP-seq data analysis:

Reads were aligned to mm9 using bowtie/bowtie2 (Langmead and Salzberg, 2012; Langmead et al., 2009). The resulting sam output files were converted to the BAM format using Samtools, version 1.6.0 (Li et al., 2009). The bam files were filtered to remove PCR duplicates using Picard tools, MarkDuplicates (**http://broadinstitute.github.io/picard**). The BAM files were converted to bigwig files for visualization on the UCSC browser (Kent 2002). The bigwig files were generated using DeepTools (Ramírez et al., 2016), bamCompare with a bin size of 10 bp ,extending fragments to 150bp (nucleosome size), filtered for mm9 blacklisted regions and normalized to 1X depth of coverage. Each bigwig file represents the $\log_2$ ratio of ChIP to the corresponding Input sample. For comparison across samples the bigwig files were normalized to effective library size. Replicates were merged into a single bigwig using UCSC tools, bigWigMerge (http://hgdownload.soe.ucsc.edu/admin/exe/). Read coverage over regions of interest were generated from matrix files generated using DeepTools, computeMatrix following which metagene plots were made in R using ggplo2 (Wilkinson, 2011). Peaks from each sample were called using Macs2, version2.1.0, (Zhang et al., 2008) in broadpeak mode with --broad-cutoff 0.05 (FDR ≤ 0.05). Overlapping peaks between replicates were identified using bedtools, intersectBed and were used for subsequent analysis. A list of all BRG1 peak calls are provided (Table S3) Peaks were annotated using HOMER, peakannotate.pl (Heinz et al., 2010). Differential peak analysis was performed on bedgraph files using Macs2, bdgdiff run with default parameters.

ATAC-seq data analysis:

Reads were aligned to the mm9 genome using bowtie, with following paramters: -S –q –m 1 –p 2 –best –strata –chunkmbs 256 (Langmead et al., 2009). The outputed sam files were converted to BAM files using Samtools version 1.3.1. Next Bedtools version 2.25.0 (Quinlan and Hall, 2010), bamtobed was used to convert BAM output files to bed file format to do the subsequent steps . BAM files were converted to bigWig files for visualization on UCSC browser and to generate metagene plots as described above (see supplementary materials on ChIP-seq data analysis).

Differential analysis of chromatin accessibility:

For the differential analysis of chromatin accessibility, we adopted a method described for analyzing significant differences in counts between DNA hypersensitive sites identified by Dnase seq (Shibata et al., 2012). We compared differences in chromatin accessibility between 12 day old (P12) $Brg1^{WT}$ and $Brg1^{cKO}$ spermatogenic cells . Here the pairwise comparison was performed using edgeR, after obtaining the read counts from each replicate across defined 300 bp windows generated from a union set of the top 100,000 peaks (ranked by F-Seq). We called windows with significantly different counts from the pairwise comparison at a FDR ≤ 0.05. The significantly altered regions were annotated using HOMER, peakannotate.pl (Heinz et al., 2010). Regions with significant differences in open chromatin are provided (Table S2).

Preparation of nuclear lysates for immunoprecipitation:

Spermatogenic cells isolated from 2- to 3-week-old males were washed once in PBS and centrifuged at 600g for 5 min at 4 C. The resulting cell pellet was re-suspended in 20 PCV (packed cell volumes) of buffer A (10mM HEPES-KOH pH7.9, 1.5mM MgCl2, 10mM KCl, 0.1% NP-40, 0.5mM DTT, 0.5mM PMSF, 1x PIC, 0.5) and left on ice to swell for 10 min. Cells were centrifuged at 600g for 5 min at 4 C, resuspended in 2 PCV of buffer A and then homogenized using a dounce (type B). The homogenate containing nuclei were centrifuged at 700g for 10 min at 4 C. Nuclear preparations were washed once again in 10 PCV of buffer A and pelleted at 5000 rpm for 10 min at 4 C. Lysates were extracted from pelleted nuclei at least twice for 1 hour each with an equal volume of high salt buffer C (20mM HEPES-KOH pH7.9, 1.5mM MgCl2,

420mM NaCl, 10mM KCl, 25 % glycerol, 0.2mM EDTA, 0.5mM DTT, 0.5mM PMSF, 1x PIC) on a nutator at 4 C. The resulting nuclear lysates were mixed with 2.8x volume of buffer D (20mM HEPES-KOH pH7.9, 20 % glycerol, 0.2mM EDTA ,0.5mM DTT, 0.5mM PMSF, 1x PIC) following which they were clarified by spinning at 14,000 rpm for 10 min at 4 C, pooled together and snap-frozen in liquid nitrogen and stored at -80 C until further use.

Co-Immunoprecipitation (Co-IP):

100 µl of magnetic protein A beads were initially washed twice in PBS for 5 min each at 4 C followed by one wash in PBS + 0.5% BSA for 10 min at 4 C. The antibodies (tableS3) were then mixed with beads resuspended in 1x volume of PBS + 0.5% BSA and allowed to conjugate for at least 2 hr at 4 C. After this, antibody-bead conjugates were washed once with 500 µl of PBS + 0.5% BSA , followed by two washes in IP buffer (20mM HEPES-KOH pH7.9, 0.15mM KCl, 10 % glycerol, 0.2mM EDTA, 0.5mM DTT, 0.5mM PMSF, 1x PIC) for 5 min each at 4 C. The antibody-coupled beads were stored in IP buffer until the nuclear lysates were processed for IP. Briefly, the lysates were thawed on ice and centrifuged at 14,000 rpm at 4 C to remove any precipitates. 500 µg of nuclear lysate was diluted in IP buffer to make up the volume to 1.3 ml and then incubated with unconjugated protein A beads to pre-clear the lysate. A fraction of lysate was set aside as input and the remaining was transferred to a tube containing the antibody coupled beads and left on a rotator overnight at 4 C. The following day the protein bound antibody-bead conjugates were separated using a magnetic separator and washed on a rotator with a series of buffers in the following order at 4 C for 5 min each: twice in IP buffer, twice in high salt wash buffer (20mM HEPES-KOH pH7.9, <u>300mM KCl</u>, 10 % glycerol, 0.2mM EDTA, 0.1 % Tween-20, 0.5mM DTT, 0.5mM PMSF, 1x PIC), twice in a low salt wash buffer (20mM HEPES-KOH pH7.9, <u>100mM KCl</u>, 10 % glycerol, 0.2mM EDTA, 0.1 % Tween-20, 0.5mM DTT, 0.5mM PMSF, 1x PIC), once in final wash buffer (20mM HEPES-KOH pH7.9, 60mM KCl, 10 % glycerol, 0.5mM DTT, 0.5mM PMSF, 1x PIC). The proteins were eluted by resuspending the beads in 2X Laemmli buffer and incubating them at 70 C for 10 min followed by boiling the samples at 95 C for 5 min.

BRG1-IP and mass spectrometry:

Prior to mass spectrometry the IPs were performed with anti-BRG1 antibody and non-specific IgG (table S4) as described above with a few modifications. After conjugation each antibody

was crosslinked to protein A beads with BS^3 (bis[sulfosuccinimidyl] suberate) crosslinking agent, prepared as per the manufacturer's instructions (Thermo scientific, 21585). Crosslinking with BS^3 significantly reduces IgG elution and improves signal to noise ratio in samples obtained by boiling beads in Laemlli buffer (Sousa et al., 2011). Briefly, antibody coupled beads were washed once in PBS for 5 min at 4 C, followed by two washes in conjugation buffer pH 7.9 (20 mM Sodium Phosphate, 0.15M NaCl). The antibody-coupled beads were resuspended in 250 ul of conjugation buffer containing 5mM BS^3 in a tube and incubated on a rotator for 40 min at room temperature. Antibody crosslinked beads were separated and the crosslinker was quenched with an equal volume of 1M glycine (1x volume of beads). The crosslinked beads were then washed once in PBS at 4 C for 5 min, followed by incubation in 100 ul of 0.11M glycine (pH2.5) for 10 min at 4 C to remove any uncrosslinked antibody. The crosslinked beads were then washed thrice in PBS tween-20 (0.1%) + 0.5 % BSA, followed by another three washes in IP buffer before transferring them to a 15 ml falcon tube containing 4 mg of nuclear lysate made up in IP buffer up to a final volume of 10ml and left on rotator overnight at 4 C. The following day the beads were washed and proteins were eluted as described above. Samples were run on a short gel from which the regions containing the proteins were cut out and processed for mass spectrometry and peptide identification.

In-gel Digestion

Gel slices were cut into 1x1 mm pieces and placed in 1.5 mL eppendorf tubes with 1 mL of water for 30 min. The water was then removed and 50 µL of 250 mM ammonium bicarbonate followed by10 µL of 45 mM 1, 4 dithiothreitol (DTT) were added prior to incubation at 50 °C for 30 min. The samples were cooled to room temperature and then alkylated with 10 µL of 100 mM iodoacetamide for 30 min. The gel pieces were washed 2x with 1 mL of water, removed and added 1 mL of 50 mM ammonium bicarbonate:acetonitrile (1:1) and allowed to incubate at room temperature for 1 hr. The solution was then removed and 200 µL of acetonitrile was added, removed, and the gel pieces dried by SpeedVac. Gel pieces were rehydrated in 70 µL of 2 ng/µL trypsin (Sigma) and 0.01% ProteaseMAX surfactant (Promega) in 50mM ammonium bicarbonate and incubated at 37 °C for 21hrs. Supernatant was removed, gel pieces added 100 µL of 80:20 (1% (v/v) formic acid in acetonitrile), combined with the former supernatant, and dried on a SpeedVac. Samples were reconsituted in 25 µL of 5% acetonitrile (0.1% (v/v) trifluroacetic acid) for LC-MS/MS analysis.

LC-MS/MS

Tryptic peptides were dissolved in 0.1% trifluoroacetic acid and directly loaded at 4 µL/min for 7 minutes onto a custom-made trap column (100 µm I.D. fused silica with Kasil frit) containing 2 cm of 200Å, 5 µm Magic C18AQ particles (Michrom Bioresources). Peptides were then eluted onto a custom-made analytical column (75 µm I.D. fused silica) with gravity-pulled tip and packed with 25 cm 100Å, 5 µm Magic C18AQ particles (Michrom). Peptides were eluted with a linear gradient from 100% solvent A (0.1% (v/v) formic acid in water:0.1% formic acid in acetonitrile (95:05)) to 35% solvent B (0.1% (v/v) formic acid in acetonitrile) in 90 minutes at 300 nanoliters per minute using a Waters NanoAcquity UPLC system directly coupled to a Thermo Scientific Q Exactive hybrid mass spectrometer. Data were acquired using a data-dependent acquisition routine of acquiring one mass spectrum (*m/z* 300 -1750) in the Orbitrap (resolution 70,000, 1e6 charges, 30 ms maximum fill time) followed by 10 tandem mass spectrometry scans (resolution 17,500, 1e5 charges, 110 ms maximum fill time, HCD collision energy 27 eV NCE). Dynamic exclusion was employed to maximize the number of peptide identifications and minimize data redundancy.

Data Analysis

Raw data files were processed into peak lists using Proteome Discoverer (version 1.4; Thermo Scientific) and then searched against the Uniprot mouse database with Mascot (version 2.5; Matrix Science) using precursor mass tolerances of 10 ppm and fragment mass tolerances of 0.5 Da. Full tryptic specificity was specified considering up to 2 missed cleavages; variable modifications of acetylation (protein N-term), pyro-glutamination (N-term glutamine), and oxidation (methionine) were considered and fixed modifications of carbamidomethylation (cysteine) were considered. Mascot search results were loaded into Scaffold (Proteome Software) with threshold values of 80% for peptides (1.0% false-discovery rate) and 90% for proteins (2 peptide minimum) for final annotation.

Supplemental references:

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H. and Glass, C. K. (2010). Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Mol. Cell* **38**, 576–589.

Langmead, B. and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359.

Langmead, B., Trapnell, C., Pop, M. and Salzberg, S. (2009). Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G. and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079.

Quinlan, A. R. and Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842.

Raab, J. R., Resnick, S. and Magnuson, T. (2015). Genome-Wide Transcriptional Regulation Mediated by Biochemically Distinct SWI/SNF Complexes. *PLoS Genet.* **11**, 1–26.

Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., Heyne, S., Dündar, F. and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res.* **44**, W160–W165.

Shibata, Y., Sheffield, N. C., Fedrigo, O., Babbitt, C. C., Wortham, M., Tewari, A. K., London, D., Song, L., Lee, B. K., Iyer, V. R., et al. (2012). Extensive evolutionary changes in regulatory element activity during human origins are associated with altered gene expression and positive selection. *PLoS Genet.* **8**,.

Sousa, M. M. L., Steen, K. W., Hagen, L. and Slupphaug, G. (2011). Antibody cross-linking and target elution protocols used for immunoprecipitation significantly modulate signal-to noise ratio in downstream 2D-PAGE analysis. *Proteome Sci.* **9**, 45.

Wilkinson, L. (2011). ggplot2: Elegant Graphics for Data Analysis by WICKHAM, H. *Biometrics* **67**, 678–679.

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nussbaum, C., Myers, R. M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**,.

A

Staging spermatocytes

Expression of staging markers

B

THY1+ spermatogonia

D

C

Cytosolic fractions

E

A

B

A

Protein	# Unique Peptide	% Coverage
BAF60A	37	82
BAF60B	26	60
BAF60C	19	45
BAF45A	14	36
BAF45B	10	38
BAF45C	8	33
BAF45D	15	49
BAF57	20	51
BAF53A	22	55

Shared SWI/SNF sub units

B

C

A

B

A

B

A

Acid extracted histones (P12 Testes)

$Brg1^{WT}$ $Brg1^{Het}$ $Brg1^{cKO}$ $Brg1^{WT}$ $Brg1^{Het}$ $Brg1^{cKO}$

H2AK119ub1 (E6C5)

H3K27ac

H3

P8 Testes   P10 Testes

C

ES cells

$Rnf2^{WT}$ $Rnf2^{KO1}$ $Rnf2^{KO2}$

H2AK119ub1 (E6C5)

Ponceau S

B

P12 Testes

$+/+$ $+/+;Mvhcre^{Tg/0}$

H2AK119ub1 (E6C5)

H3

D

$Brg1^{WT}$ $Brg1^{Het}$ $Brg1^{cKO}$ $Brg1^{WT}$ $Brg1^{Het}$ $Brg1^{cKO}$ $Brg1^{WT}$ $Brg1^{Het}$ $Brg1^{cKO}$

L

250 KDa
150 KDa
75 KDa
50 KDa
25 KDa
15 KDa

H2AK119ub1

H3

Cytosol (10 ug)   Nucleoplasm (3 ug)   Chromatin (3 ug)

A

B

C