Genotyping-by-sequencing and microarrays are complementary for detecting

quantitative trait loci by tagging different haplotypes in association studies

Sandra Silvia Negro[1,3], Emilie Millet[2,4], Delphine Madur[1], Cyril Bauland[1], Valérie Combes[1],

Claude Welcker[2], François Tardieu[2], Alain Charcosset[1], Stéphane Dimitri Nicolas[1]

Affiliations :

[1] GQE – Le Moulon, INRA, Univ. Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay,

91190 Gif-sur-Yvette, France.

[2] Laboratoire d'Ecophysiologie des Plantes sous Stress Environnementaux (LEPSE),

UMR759, INRA-SupAgro, 34060 Montpellier, France.

[3] Present address: GIGA-R Medical Genomics, University of Liège, Belgium.

[4] Present address: Biometris, Department of Plant Science, Wageningen University &

Research, 6700 AA Wageningen, The Netherlands.

E-mail addresses:

snegro@uliege.be; emilie.millet@wur.nl; delphine.madur@inra.fr; cyril.bauland@inra.fr;

valerie.combes@inra.fr; claude.welcker@inra.fr; francois.tardieu@inra.fr;

alain.charcosset@inra.fr; stephane.nicolas@inra.fr

Corresponding author:

stephane.nicolas@inra.fr

Abstract

Background: Single Nucleotide Polymorphism (SNP) array and re-sequencing technologies

have different properties (*e.g.* calling rate, minor allele frequency profile) and drawback (*e.g.*

ascertainment bias), which lead us to study the complementarity and consequences of using

them separately or combined in diversity analyses and Genome-Wide Association Studies

(GWAS). We performed GWAS on three traits (grain yield, plant height and male flowering

time) measured in 22 environments on a panel of 247 diverse dent maize inbred lines using

three genotyping technologies (Genotyping-By-Sequencing, Illumina Infinium 50K and

Affymetrix Axiom 600K arrays).

Results: The effects of ascertainment bias of both arrays were negligible for deciphering

global genetic trends of diversity in this panel and for estimating relatedness. We developed

an original approach based on linkage disequilibrium (LD) extent in order to determine

whether SNPs significantly associated with a trait and that are physically linked should be

considered as a single QTL or several independent QTLs. Using this approach, we showed

that the combination of the three technologies, which have different SNP distribution and

density, allowed us to detect more Quantitative Trait Loci (QTLs, gain in power) and

potentially refine the localization of the causal polymorphisms (gain in position).

Conclusions: Conceptually different technologies are complementary for detecting QTLs by

tagging different haplotypes in association studies. Considering LD, marker density and the

combination of different technologies (arrays and re-sequencing), the genotypic data presently

available were most likely enough to well represent polymorphisms in the centromeric

regions, whereas using more markers would be beneficial for telomeric regions.

Keywords: GWAS, linkage disequilibrium, genome coverage, maize, high-throughput

genotyping technologies.

Background

Understanding the genetic bases of complex traits involved in the adaptation to biotic and abiotic stress in plants is a pressing concern, with world-wide drought due to climate change as a major source of human food and agriculture threats. Recent progress in next generation sequencing and genotyping array technologies contribute to a better understanding of the genetic basis of quantitative trait variation by performing Genome-Wide Association Studies (GWAS) on large diversity panels [1]. Single Nucleotide Polymorphism (SNP)-based techniques became the most commonly used genotyping methods for GWAS because SNPs are cheap, numerous, codominant and can be automatically analysed with SNP-arrays or produced by genotyping-by-sequencing (GBS), or sequencing [2-4]. The decreasing cost of genotyping technologies have led to an exponential increase in the number of markers used for the GWAS in association panels, thereby raising the question of computation time to perform the association tests. Computational issues were addressed by using either approximate methods by avoiding re-estimating variance component for each SNP [5] or exact methods using mathematical tools for sparing time in matrix inversion [6, 7]. It is noteworthy that using approximate computation in GWAS can produce inaccurate p-values when the SNP effect size is large or/and when the sample structure is strong [8].

Several causes may impact the power of Quantitative Trait Locus (QTL, locus involved in quantitative trait variation) detection in GWAS. Highly diverse panels have in general undergone multiple historical recombinations, leading to a low extent of linkage disequilibrium (LD). However, these panels can present different average and local patterns of LD [9-11]. A high marker density and a proper distribution of SNPs are therefore essential to capture causal polymorphisms. Furthermore, minor allele frequencies (MAF), population stratification and cryptic relatedness are three other important parameters affecting power and

false positive detection [12, 14]. These last two factors are substantial in several cultivated

species such as maize [15] and grapevine [16], and their impact on LD can be statistically

evaluated [17]. Population structure and kinship can be estimated using molecular markers

[18-21] and can be modelled to efficiently detect marker-trait associations due to linkage only

[12, 22, 23]. These advances have largely increased the power and effectiveness of linear

mixed models that can now efficiently account for population structure and relatedness in

GWAS [12, 8].

In maize, an Illumina Infinium HD 50,000 SNP-array (50K array) was developed by Ganal *et*

al. [3] and has been used extensively for diversity and association studies [24, 25]. For

example, GWAS were conducted to unravel the genetic architecture of phenology, yield

component traits and to identify several flowering time QTLs linked to adaptation of tropical

maize to temperate climate [26, 27]. With the same array, Rincent *et al.* [11] showed that LD

occurs over a longer distance in a dent than in a flint panel, with appreciable effects on the

power of QTL detection. Comparison of LD extent between association panels suggests that

genotyping with 50K markers causes a limited power of GWAS in many panels due to the

low LD extent and the correlation between allelic values at some SNPs due to the kinship and

population structure [14, 27]. Therefore, higher marker densities are desirable because the

maize genome size is large (2.4 Gb), the level of diversity is high, and LD extent is low (more

than one substitution per hundred nucleotides) [28]. As a consequence, an Affymetrix Axiom

600,000 SNP-array (600K array) was developed and used in association genetics [29, 30] and

detection of selective sweep [4]. Another possibility is whole genome sequencing, but this is

currently unpractical for large genomes such as maize because of the associated cost. Hence, a

Genotyping-By-Sequencing (GBS) procedure has been developed [2] that targets low-copy

genomic regions by using cheap restriction enzymes. Genotyping-by-sequencing has been

successfully used in maize for genomic prediction [31]. Romay *et al.* [32] and Gouesnard *et*

al. [33] highlighted the interest of the GBS for (i) deciphering and comparing the genetic

diversity of the inbred lines in seedbanks and (ii) identifying QTLs by GWAS for kernel

colour, sweet corn and flowering time. To our knowledge, the respective interests of DNA

arrays and GBS for diversity analyses and GWAS have never been compared in plants.

The main drawback of the DNA arrays is that they do not allow to discover new SNPs. This

possibly leads to some ascertainment bias in diversity analysis when the SNPs selected for

building arrays come from (i) the sequencing of a set of individuals who did not well

represent the diversity explored in the studied panel, (ii) a subset of SNPs that skews the

allelic frequency profile towards the intermediate frequencies [27, 34]. Ascertainment bias

can compromise the ability of the SNP arrays to reveal an exact view of the genetic diversity

[34]. Genotyping-by-sequencing can overcome ascertainment bias since it is based on

sequencing and therefore allows the discovery of alleles in the diversity panel analysed. It can

be generalized to any species at a low cost providing that numerous individuals have been

sequenced in order to build a representative library of short haplotypes to call SNPs [35].

Non-repetitive regions of genomes can be targeted with two- to three-fold higher efficiency,

thereby considerably reducing the computationally challenging problems associated with

alignment in species with high repeat content. However, GBS may have a low-coverage

leading to a high missing data rate (65% in both studies; [32, 33]) and heterozygote under-

calling, depending on genome size and structure, and on the number of samples combined in

the flow-cell. Furthermore, GBS requires the establishment of demanding bioinformatic

pipelines and imputation algorithms [36]. Pipelines have been developed to call SNP

genotypes from raw GBS sequence data and to impute the missing data from a haplotype

library [35, 36].

Here, we investigated the impact of using GBS and DNA arrays on the quality of the

genotyping data, together with the biological properties of data generated by these

technologies, and the potential complementarity of these approaches. In particular, we

analyzed the impact of increasing the marker density and using different genotyping

technologies (sequencing *vs* array) on (i) the estimates of relatedness and population structure,

(ii) the detection of QTLs (power. To address these issues, we performed a GWAS based on

genotypic datasets obtained using either GBS or DNA arrays with low (50k) or high (600k)

densities on a diversity panel of maize hybrids obtained from a cross of dent lines with a

common flint tester. Three traits were considered, namely grain yield, plant height and male

flowering time (day to anthesis), measured in 22 different environments (sites $\times$ years $\times$

treatments) over Europe. We developed an original approach based on LD extent in order to

determine whether SNPs significantly associated with a trait should be considered as a single

QTL or several independent QTLs.

Results

Combining Tassel and Beagle imputations improved the genotyping quality for GBS

We estimated the genotyping and imputation concordance of the GBS based on common

markers with the 50K or 600K arrays (Additional file 1: Figure S1 and Table S1). After SNP

calling from reads using AllZeaGBSv2.7 database (direct reads, GBS_1, Additional file 1:

Figure S1), the call rate was 33.81% on the common SNPs with the 50K array, *vs* 37% for the

whole GBS dataset. The genotyping concordance rate was 98.88% (Additional file 1: Table

S1). After imputation using *TASSEL* by Cornell Institute (GBS$_2$), the concordance rate was

96.04% on the common markers with the 50K array and 11.91% of missing data remained for

the whole GBS dataset. In GBS$_3$, all missing data were imputed by *Beagle* but yielded a lower

concordance rate (92.14% and 91.58% for the 50K and the 600K arrays). In an attempt to

increase the concordance rate of the genotyping while removing missing data, we tested two

additional methods, namely GBS$_4$ where the missing data and heterozygotes of Cornell

imputed data (GBS$_2$) were replaced by Beagle imputation, and GBS$_5$ where Cornell

homozygous genotypes (GBS$_2$) were completed by imputations from GBS$_3$ (Additional file 1:

Figure S1 and Table S1). GBS$_5$ displayed a slightly better concordance rate than GBS$_2$

(96.25% vs 96.04%) and predicted heterozygotes with a higher quality than GBS$_4$. GBS$_5$ was

therefore used for all genetic analyses and named GBS hereafter.

GBS displayed more rare alleles and lower call rate than microarrays

The SNP call rate was higher for the arrays (average values of 96% and >99% for the 50K

and 600K arrays, respectively), than for the GBS (37% for the direct reads). The MAF

distribution differed between the technologies (Figure 1): while the use of arrays resulted in a

near-uniform distribution, GBS resulted in an excess of rare alleles with a L-shaped

distribution (22% of SNPs with MAF < 0.05 for the GBS *versus* 6% and 9% for the 50K and

600K, respectively). This is not surprising since the 50K array was based on 27 sequenced

lines for SNPs discovery [3], the 600K array was based on 30 lines for [4], whereas GBS was

based on 31,978 lines, thereby leading to higher discovery of rare alleles. Consistent with

MAF distribution, the average gene diversity (*He*) was lower for GBS (0.27) than for arrays

(0.35 and 0.34 for the 50K and 600K arrays, respectively). The distribution of SNP

heterozygosity was similar for the three technologies, with a mean of 0.80%, 0.89% and 0.22

% for the 50K and 600K arrays and GBS, respectively. The heterozygosity of inbred lines was

highly correlated between technologies with large coefficients of Spearman correlation: $r_{50K\text{-}}$

$_{600K}$ = 0.90, $r_{50K\text{-}GBS}$ = 0.76, $r_{600K\text{-}GBS}$ = 0.83. The distribution of the SNPs along the genome

was denser in the telomeres for the GBS and in the peri-centromeric regions for the 600K

array, whereas the 50K array exhibited a more uniform distribution (top graph in Figure 2 and

in Additional file 2: Figure S2).

Population structure and relatedness were consistent between the three technologies

We used the ADMIXTURE software to analyse the genetic structure within the studied panel

based on SNPs from the three technologies, by using two to ten groups. Based on a K-fold

cross-validation, the clustering in four genetic groups (N_Q = 4) was identified as the best one

in the datasets resulting from the three technologies. Considering a threshold of 0.5 (ancestral

fraction), the assignation to the four groups was identical except for a few admixed inbred

lines (Additional file 3: Figure S3). Based on the 50K, the four groups were constituted by (i)

39 lines in the Non Stiff Stalk (Iodent) family traced by PH207, (ii) 46 lines in the Lancaster

family traced by Mo17 and Oh43, and (iii) 55 lines in the stiff stalk families traced by B73

and (iv) 107 lines that did not fit into the three primary heterotic groups, such as W117 and

F7057. This organization appeared consistent with the organization of breeding programs into

heterotic groups, generally related to few key founder lines.

We compared two estimators of relatedness between inbred lines, IBS (Identity-By-State) and

K_Freq (Identity-By-Descend), calculated per technology. For IBS, pairs of individuals were

on average more related using the GBS data than those from arrays (Additional file 3: Table

S2). Relatedness estimated with the two arrays were highly correlated: r = 0.95 and 0.98 for

IBS and *K_Freq*, respectively (Figure 3 and Additional file 3: Figure S4b). The differences

between the kinships estimated from the three technologies were reduced if the excess of rare

alleles in the GBS was removed (Additional file 3: Figure S4c).

We further carried out diversity analyses by performing Principal Coordinate Analyses

(PCoA) on IBD (*K_Freq*, weights by allelic frequency) estimated from the three technologies.

The two first PCoA axes explained 12.9%, 15.6% and 16.3% of the variability for the GBS,

50K and 600K arrays, respectively. The same pattern was observed regardless of the

technology with the first axis separating the Stiff Stalk from the Iodent lines and the second

axis separating the Lancaster from the Stiff Stalk and Iodent lines (see illustration with the

50K kinship, Additionnal File 3: Figure S5). Key founders lines of the three heterotic groups

(Iodent: PH207, Stiff Stalk: B73, Lancaster: Mo17) were found at extreme positions along the

axes, which was consistent with the admixture groups previously described.

Long distance linkage disequilibrium was removed by taking into account population structure or relatedness

In order to evaluate the effect of kinship and the genetic structure on linkage disequilibrium

(LD), we studied genome-wide LD between 29,257 PANZEA markers from the 50K array

within and between chromosomes before and after taking into account the kinship (*K_Freq*

estimated from the 50K array), structure (Number of groups = 4) or both (Additional file 4:

Figure S6). Whereas inter-chromosomal LD was only partially removed when the genetic

structure was taken into account, it was mostly removed when either the kinship or both

kinship and structure were considered (Additional file 4: Figure S6b and c). Accordingly, long

distance intra-chromosomal LD was almost totally removed for all chromosomes by

accounting for the kinship, structure or both. Interestingly, some pairs of loci located on

different chromosomes or very distant on a same chromosome remained in high LD despite

correction for genetic structure and kinship (Additional file 4: Figure S6). This can be

explained either by genome assembly errors, by chromosomal rearrangements such as

translocations or by strong epistatic interactions. Linkage disequilibrium decreased with

genetic or physical distance, Additional file 4: Figure S7). The majority of pairs of loci with

high LD (r^2K>0.4) in spite of long physical distance (>30Mbp), were close genetically

(<3cM), notably on chromosome 3, 5, 7 and to a lesser extent 9 and 10. These loci were

located in centromeric and peri-centromeric regions that displayed low recombination rate,

suggesting that this pattern was due to variation of recombination rate along the chromosome.

Only very few pairs of loci in high LD were genetically distant (>5cM) but physically close

(<2Mbp). Linkage disequilibrium (r^2K and r^2KS) was negligible beyond 1 cM since 99% of

LD values were less than 0.12 in this case. Note that some unplaced SNPs remained in LD

after taking into account the kinship and structure with some SNPs with known positions on

chromosome 1, 3 and 4 (Additionnal File 4: Figure S6). Therefore, LD measurement

corrected by the kinship can help to map unplaced SNPs.

Linkage disequilibrium strongly differed between and within chromosomes

We combined the three technologies together to calculate the r^2K for all pairs of SNPs, which

were genetically distant by less than 1 cM. For any chromosome region, LD extent in terms of

genetic and physical distance showed a limited variation over the 100 sets of 500,000 loci

pairs (cf. Material). This suggests that the estimation of LD extent did not strongly depend on

our set of loci. LD extent varied significantly between chromosomes for both high

recombinogenic (>0.5 cM/Mbp) and low recombinogenic regions (<0.5 cM/Mbp, Table 1).

Chromosome 1 had the highest LD extent in high recombination regions (0.062 ±0.007 cM)

and chromosome 9 the highest LD extent in low recombinogenic regions (898.6±21.7 kbp)

(Table 1). Linkage disequilibrium extent relative to genetic and physical distances was highly

and positively correlated in high recombinogenic regions (r = 0.86), whereas it was not in low

recombinogenic regions (r = -0.64).

Table 1: Variation of LD extent, and percentage of genome covered.

		Chromosome										Whole Genome
		1	2	3	4	5	6	7	8	9	10	
Physical Size (kbp)		301,354	237,069	232,140	241,474	217,873	169,174	176,765	175,794	156,751	150,189	2,058,583
Genetic Size (cM)		268	211	188	150	205	129	148	182	145	139	1766
Physical LD extent (kbp) in Low Recombination regions (<0.5 cM / Mbp)		306	491	846	808	658	418	547	497	899	815	629
Genetic LD Extent (cM) in High Recombination regions (>0.5 cM / Mbp)		0.062	0.027	0.033	0.022	0.031	0.019	0.012	0.038	0.023	0.019	0.029
Percent of physical genome covered	50K	81%	72%	76%	77%	74%	67%	71%	73%	71%	71%	74%
	600K	98%	88%	91%	89%	90%	84%	81%	90%	87%	84%	89%
	GBS	92%	81%	84%	83%	83%	77%	77%	81%	79%	76%	82%
	Combined	98%	90%	92%	90%	91%	87%	83%	92%	88%	85%	90%
Percent of genetic map covered	50K	72%	41%	44%	38%	41%	32%	24%	46%	32%	27%	42%
	600K	96%	71%	76%	68%	72%	62%	47%	78%	63%	53%	71%
	GBS	86%	58%	61%	53%	61%	48%	37%	63%	48%	40%	58%
	Combined	97%	74%	78%	72%	74%	65%	51%	81%	66%	57%	74%

Genetic and Physical LD extent were obtained by adjusting Hill and Weir model's on

100 different sets of 500,000 loci randomly sampled in high and low recombination

regions, respectively. The value represented the average across these 100 sets. The

percentage of genome coverage was estimated using markers with MAF > 5% and

E(r^2k) = 0.1, for each technology and for the three technologies combined

(GBS+600K+50K).

The effective population size (*Ne*) estimated from the Hill and Weir's model [37] using

genetic distance varied from 7.9±0.04 (Chromosome 1) to 41.2±0.12 (Chromosome 7) in high

recombinogenic regions. Noteworthy, the same approach lead to unrealistic values in low

recombinogenic regions (from 961 on Chromosome 6 to more than 1 million for chromosome

2 and 10), thereby confirming that the use of genetic distance is not well suited to model local

LD in low recombinogenic regions.

Finally, we studied the variation of local LD extent along each chromosome by adjusting the

Hill and Weir's model against genetic distance within a sliding windows of 1cM (Additional

file 4: Figure S8). After removing intervals that did not reach our criteria (Absence of model

convergence, effective population size > 247, low number of loci), the 3,205 remaining

intervals (90%) showed a high variation for genetic LD extent along each chromosome, with

LD extent varying from 0.019 (Ne = 246) to 0.997 cM (Ne = 0.06) (Additional file 4: Figure

S8).

Large differences in genome coverage between technologies

We estimated the percentage of the genome that was covered by LD windows around SNPs,

calculated by using either physical or genetic distances (Table 1). We observed a strong

difference in coverage between the three technologies at both genome-wide and chromosome

scale, as illustrated in Figure 2 on chromosome 3 (Table 1, and Additional file 2: Figure S2).

For a LD extent of r^2K = 0.1, 74%, 82% and 89% of the physical map, and 42%, 58% and

71% of the genetic map were covered by the 50K array, the GBS and the 600K array,

respectively (Table 1). For the combined data (50K + 600K + GBS), the coverage strongly

varied between chromosomes, ranging from 83% (chromosome 7) to 98% (chromosome 1) of

the physical map, and from 51% (chromosome 7) to 97% (chromosome 1) of the genetic map

(Table 1). For the physical map, increasing the LD extent threshold to r^2K=0.4 reduced the

genome coverage from 89% to 49% for 600K, 82% to 28% for GBS, 74% to 20% for 50K

and 90% to 52% for the combined data. Increasing the MAF threshold reduced slightly the

genome coverage, with smaller reduction for the physical map than genetic map. Surprisingly,

increasing the SNP number by combining the markers from the arrays and GBS did not

strongly increase the genome coverage as compared to the 600K, regardless of the threshold

for LD extent (Figure 2 and Additional file 2: Figure S2).

We observed a strong variation of genome coverage along each chromosome with contrasted

patterns in low and high recombinogenic regions (Figure 2 and Additional file 2: Figure S2).

While low recombinogenic regions were totally covered with all the technologies (except for

few intervals using the 50K array), the genome coverage in high recombinogenic regions

varied depending on both technology and SNP distribution. 47% of the 2Mbp intervals in

high recombination regions were better covered by the 600K array than the GBS against only

1%, which were better covered by GBS than 600K.

Number of QTLs detected using genome-wide association studies increases with markers density

We observed a strong variation in the number of SNP significantly associated with the three

traits across the 22 environments (Table 2). The mean number of significant SNPs per

environment and trait was 3.7, 44.7, 17.9 and 62.3 for the 50K, 600K, GBS and the three

technologies combined, respectively (Table 3). Considering the p-value threshold used, 28,

303 and 204 false positives were expected among the 243, 2,953 and 1,182 associations

detected for 50K, 600K and GBS, respectively. False discovery rate appeared therefore higher

for GBS (17.2%) than for DNA arrays (11.5% and 10.2% for 50K and 600K, respectively). It

could be explained by the higher genotyping error rate of GBS due to imputation and/or by its

higher number of makers with a lower MAF. Both reduce the power of GBS compared to

DNA arrays and therefore lead to a higher false discovery rate. Proportionally to the SNP number, 50K and 600K arrays resulted in 1.5- and 1.7-fold more associated SNPs per situation (environment × trait) than GBS (p-value<2x10^{-6}, Table 3). This difference between arrays and GBS was higher for grain yield (GY) and plant height (plantHT) than for male flowering time (DTA, Table3).

Table 2: Number of significant SNPs per environment, per technology and for the combined technologies.

Environment	Flowering Time					Plant Height					Grain Yield				
	Heritability	50K	600K	GBS	Combined	Heritability	50K	600K	GBS	Combined	Heritability	50K	600K	GBS	Combined
Gai12R	0.62	8	80	24	104	0.15	1	47	41	89	0.31	0	23	8	31
Gai12W	0.66	5	42	15	59	0.40	0	1	3	4	0.56	3	71	14	85
Gai13R	0.58	0	24	8	31	0.56	0	6	6	11	0.66	0	4	5	9
Gai13W	0.78	1	45	9	54	0.40	0	1	3	4	0.81	2	7	1	9
Ner12R	0.70	7	23	18	43	0.22	0	7	3	10	0.53	1	10	6	16
Ner12W	0.80	1	80	30	108	0.28	0	2	2	4	0.60	1	8	6	15
Ner13R	0.77	3	60	26	88	0.22	1	25	13	38	0.35	0	13	7	20
Ner13W	0.81	2	23	17	42	0.27	0	8	5	13	0.76	2	28	4	32
Cra12R	0.64	1	40	18	59	0.26	0	6	3	9	0.23	3	57	45	100
Cra12W	0.69	3	25	19	44	0.18	10	69	16	83	0.54	12	98	53	149
Kar12R	0.71	4	30	21	51	0.26	0	4	3	7	0.73	0	5	6	11
Kar12W	0.77	8	60	10	73	0.22	1	10	4	14	0.54	2	19	11	29
Kar13R	0.66	3	65	11	77	0.27	0	4	2	6	0.92	4	37	24	62
Kar13W	0.81	0	17	12	29	0.26	0	2	7	9	0.67	4	12	6	19
Mur13R	0.85	3	48	19	68	0.26	7	61	7	68	0.84	14	90	28	116
Mur13W	0.8	0	11	8	19	0.33	3	4	2	9	0.74	10	80	25	103
Deb12R	0.58	1	14	16	30	0.25	0	1	0	1	0.56	2	14	0	16
Deb12W	0.73	0	25	38	61	0.41	0	6	7	12	0.47	0	6	2	8
Deb13R	0.60	1	17	5	23	0.08	0	33	9	42	0.35	1	22	15	37
Cam12R	0.36	0	3	1	4	0.52	23	209	88	289	0.35	21	286	102	379
Cam12W	0.60	1	18	13	31	0.43	22	270	72	338	0.52	41	525	167	684
Cam13R	0.45	0	9	7	16	0.15	0	2	3	5	0.18	0	1	3	4
Average		2.4	34.5	15.7	50.6		3.1	35.4	13.6	48.4		5.6	64.4	24.5	87.9
Median		1	25	15.5	47.5		0	6	4.5	10.5		2	20.5	7.5	30

The average, median and standard deviation (SD) per environment are calculated for each trait (male Flowering Time, Plant Height, Grain Yield).

Table 3: Comparison of associated SNPs and QTLs detected between traits and three technologies.

		Significant SNPs				QTLs			
Technology		50K	600K	GBS	Combined	50K	600K	GBS	Combined
Marker Nb		42049	459191	308946	810580	42049	459191	308946	810580
Total Nb	DTA	52	759	345	1114	20	130	133	226
	plantHT	68	778	299	1065	16	96	90	160
	GY	123	1416	538	1934	33	166	120	238
	Per trait	81	984	394	1371	23	131	114	208
Average per envir.	DTA	2.4	34.5	15.7	50.6	0.9	5.9	6.0	10.3
	plantHT	3.1	35.4	13.6	48.4	0.7	4.4	4.1	7.3
	GY	5.6	64.4	24.5	87.9	1.5	7.5	5.5	10.8
	Per trait	3.7	44.7	17.9	62.3	1.0	5.9	5.2	9.5
Average per SNP tested	DTA	5.70E-5	7.50E-5	5.10E-5	2.10E-5	2.20E-5	1.30E-5	2.00E-5	1.30E-5
	plantHT	7.40E-5	7.70E-5	4.40E-5	2.00E-5	1.70E-5	9.50E-6	1.30E-5	9.00E-6
	GY	1.30E-4	1.40E-4	7.90E-5	3.60E-5	3.60E-5	1.60E-5	1.80E-5	1.30E-5
	Per trait	8.80E-5	9.70E-5	5.80E-5	7.70E-5	2.50E-5	1.30E-5	1.70E-5	3.50E-5

QTLs were obtained by grouping associated SNPs with overlapping LD windows

(*LD_win*) for the three traits (DTA: male flowering time; PlantHT: plant height; GY:

grain yield). *Marker Nb*: is the number of markers tested in GWAS. *Total number*: is

the sum of associated SNPs or QTLs across environments. *Average per envir.*: is the

average number of QTLs obtained in 22 environments for three traits (66 trait-

environments combinations). *Average per SNP tested*: is the number of associated

SNPs or QTLs detected divided by the number of SNP tested.

We used two approaches based on LD for grouping significant SNPs: (i) considering that all

SNPs with overlapping LD windows for $r^2K=0.1$ belong to the same QTL (*LD_win*) and (ii)

grouping significant SNPs that are adjacent on the physical map and are in LD ($r^2K > 0.5$,

LD_adj). The QTLs defined by using the two approaches were globally consistent since

significant SNPs within QTLs were in high LD whereas SNPs from different adjacent QTLs

were not (Additional file 6: Figure S9-LD-Adjacent and Additional file 7: Figure S9-LD-

Windows). *LD_adj* detected more QTLs than *LD_win* for flowering time (242 vs 226), plant

height (240 vs 160) and grain yield (433 vs 237). The number of QTLs detected with the

LD_adj approach increased strongly when the LD threshold was set above 0.5. Differences in

QTL groupings between the two methods were observed for specific LD and recombination

patterns. This occurred for instance on chromosome 6 for grain yield (Additional file 6:

Figure S9-LD-Adjacent and Additional file 7: Figure S9-LD-Windows). Within this region,

the recombination rate was low and the LD pattern between associated SNPs was complex.

While *LD_adj* splitted several SNPs in high LD into different QTLs (for instance QTL 232,

235, 237, 249), *LD_win* grouped together associated SNPs that are genetically close but

displayed a low LD (Additional file 6: Figure S9-LD-Adjacent and Additional file 7: Figure

S9-LD-Windows). Reciprocally, for flowering time, we observed different cases where

LD_win separated distant SNPs in high LD into different QTLs whereas *LD_adj* grouped

them (QTL 25 and 26, 51 to 53, 95 to 97, 208 and 209, 218 and 219). As these differences

were specific to complex LD and recombination patterns, we used the *LD_win* approach for

the rest of the analyses.

Although a large difference in number of associated SNPs was observed between 600K and

GBS, little difference was observed between QTL number after grouping SNPs (Table 2,

Table 3). The mean number of QTLs was indeed 1.0, 5.9 and 5.2 and 9.5 for the 50K, 600K,

GBS, and the three technologies combined, respectively (Table 3). Note that the number of

QTLs continued to increase with marker density when SNPs from GBS, 50K and 600K were

combined (Figure 4). The number of SNPs associated with each QTL varied according to the

technology (on average 3.7, 7.6, 3.4 and 6.6 significant SNPs for the 50K, 600K, GBS, and

the combined technologies, respectively). The total number of QTLs detected over all

environments by using the 600K array and GBS was close for flowering time (130 *vs* 133)

and plant height (96 *vs* 90). It was 1.4-fold higher for the 600K than GBS for grain yield (166

vs 120).

The 600K and GBS were highly complementary for association mapping

The 600K and GBS technologies were highly complementary to detect QTLs for the three

traits: 78%, 76% and 71% of the QTLs of flowering time, plant height and grain yield,

respectively, were specifically detected by 600K or GBS (Figure 5). On the contrary, 50K

displayed very few specific QTLs. While only 9 out 69 QTLs from the 50K array were not

detected when the 600K array was used, 39 QTLs detected using the 50K array were not

detected when using GBS. When we combined the GBS and 600K markers, 7% of their

common QTLs had $-log_{10}(Pval)$ increased by 2 and 21% by 1 potentially indicating a gain in

accuracy of the position of the causal polymorphism (Additional file 8: Table S3).

This complementarity between GBS and 600K is well exemplified with two strong

association peaks for flowering time on chromosome 1 (QTL32) and 3 (QTL95) detected in

several environments (Additional file 8: Table S3 and Figure 6a). In order to better understand

the origin of the complementarity between GBS and 600K technologies for GWAS, we

scrutinized the LD between SNPs and the haplotypes within these two QTLs (Figure 6b and c,

and Additional file 9: Figure S10 for other examples). For example, QTL95 showed a gain in

power. It was only identified by the 600K array although the region included numerous SNPs

from GBS close to the associated peak. None of these SNPs was in high LD with the most

associated marker of the QTL95 (Figure 6b). Another example is QTL32, which was detected

by 1 to 10 GBS markers in 9 environments with $–log(p\text{-}value)$ ranging from 5 to 7.6, whereas

it was detected by only two 600K markers in one environment (Ner12W) with $–log(p\text{-}value)$

slightly above the significance threshold (Additional file 8: Table S3 and Figure 6b).

Haplotype analyses showed that the SNPs from the GBS within QTL95 were not able to

discriminate all haplotypes (Figure 6c). In QTL95, using the 600K markers allowed one to

discriminate the three main haplotypes (H1, H2, H3), whereas using the GBS markers did not

allow discrimination of H3 against H1 + H2. As H1 contributed to an earlier flowering time

than H2 or H3, associations appeared more significant for the 600K than for GBS (Figure 6c).

In QTL32, the use of GBS markers allowed identifying late individuals that mostly displayed

H1, H2 and H3 haplotypes, against early individuals that mostly displayed H4 and H5

haplotypes (Figure 6c). The gain of power for GBS markers as compared to 600K markers for

QTL32 originated from the ability to discriminate late individuals (black alleles) from early

individuals (red alleles) within H4 haplotypes (Figure 6c).

Stability, pleiotropy and distribution of QTL detected across environments

After combining the three technologies, we identified 226, 160, 238 QTLs for the flowering

time, plant height and grain yield, respectively (Table 3 and Additional file 8: Table S3). We

highlighted 23 QTLs with the strongest effects on flowering time, plant height and grain yield

($-log_{10}(Pval) \geq 8$, Table 4). The strongest association corresponded to the QTL95 for

flowering time ($-log_{10}(p\text{-}value) = 10.03$) on chromosome 3 (158,943,646 – 159,005,990 bp),

the QTL135 for GY ($-log_{10}(p\text{-}value) = 18.7$) on chromosome 6 (12,258,527 – 29,438,316 bp)

and QTL78 on chromosome 6 (12,258,527 – 20,758,095 bp) for plant height ($-log_{10}(p\text{-}value)$

$= 17.31$). The QTL95 for flowering time trait was the most stable QTLs across environments

since it was detected in 19 environments (Additional file 8: Table S3). Moreover, this QTL

showed a colocalization with QTL74 for grain yield in 5 environments and QTL30 for plant

height in 1 environment suggesting a pleiotropic effect. More globally, 472 QTLs appeared

trait-specific whereas 70 QTLs overlapped between at least two traits (6,3%, 5.2% and 3.0%

for GY and plantHT, GY and DTA, and DTA and plantHT, respectively) suggesting that

some QTLs are pleiotropic (Additional file 10: Figure S11). This is not surprising since

average corresponding correlations within environments for these traits were moderate (0.47,

0.54 and 0.45, respectively). Only 0.7% overlapped between the three traits (Additional file

10: Figure S11). Twenty percent of QTLs were detected in at least two environments and 9%

in at least three environments (Additional file 10: Figure S12 and Table S4). We observed no

significant differences of stability between the three traits (*p-value* = 0.2). However, 6 out 7

most stable QTLs (Number of environments >5) were found for flowering time and a higher

proportion of QTLs were specific for plant height than grain yield and flowering time (85% vs

77% for both flowering time and grain yield, *p-value* = 0.09, *p-value* = 0.2), respectively. We

observed that QTLs that displayed a significant effect in more than one environment had

larger effects and *-log(p-value)* values than those significant in a single environment. This

difference in *-log(p-value)* values was stronger for grain yield and plant height than flowering

time.

Table 4: Summary of the main QTLs ($-log_{10}(Pval) \geq 8$) identified for the three traits.

Trait	QTL	Chr	LowerLimit	UpperLimit	Pos	R2	Effect	Log	MinorAll	MajorAll	MAF	EnvMax	NbDiffEnv
DTA	95	3	158943646	159005990	158974594	0.15	1.27	10.03	G	C	0.41	Ner13R	19
plantHT	21	2	129971437	130912039	130441738	0.15	-6.74	9.21	A	G	0.14	Cam12W	3
plantHT	71	6	6593785	6636807	6614012	0.13	-4.76	8.39	G	A	0.18	Cam12R	2
plantHT	72	6	6793841	6837747	6807230	0.14	-4.87	8.77	T	G	0.18	Cam12R	2
plantHT	78	6	12258527	20758095	20330595	0.27	-8.99	17.31	C	T	0.26	Cam12W	4
plantHT	79	6	21037721	23951687	22905376	0.19	-5.45	11.42	T	G	0.31	Cam12R	3
plantHT	80	6	24184017	26606537	25317825	0.13	-4.41	8.17	T	C	0.2	Cam12R	2
plantHT	81	6	26695327	28659766	28130108	0.16	-5.14	9.84	C	G	0.44	Cam12R	2
plantHT	94	6	101463249	101501936	101482646	0.14	-5.98	8.22	T	A	0.17	Cam12W	2
plantHT	110	8	12767198	12798330	12782777	0.19	-7.67	12.44	C	T	0.22	Cam12W	3
GY	65	3	140505559	144210207	141621777	0.12	0.42	8.13	A	C	0.27	Gai12W	5
GY	85	3	186994852	187057772	187028970	0.12	-0.48	8.49	A	C	0.28	Kar12W	1
GY	120	6	5131927	5177694	5155708	0.12	-0.58	8.24	C	T	0.42	Cam12W	2
GY	122	6	5623945	5659803	5638516	0.12	-0.57	8.11	T	G	0.23	Cam12W	2
GY	124	6	5855407	5887383	5871000	0.12	-0.56	8.4	T	G	0.42	Cam12W	2
GY	127	6	6593785	6636807	6612654	0.16	-0.65	10.44	C	A	0.32	Cam12W	2
GY	128	6	6793841	6837747	6807462	0.12	-0.54	8.41	A	C	0.26	Cam12W	2
GY	129	6	6878877	6930838	6890199	0.12	0.63	8.06	T	A	0.48	Cam12W	3
GY	130	6	7027497	7088575	7046773	0.13	0.62	8.71	A	G	0.4	Cam12W	3
GY	131	6	7113662	7200479	7159714	0.12	0.59	8.24	T	C	0.39	Cam12W	3
GY	135	6	12258527	29438316	18528943	0.28	-0.78	18.7	G	C	0.31	Cam12W	6
GY	147	6	101463249	101501936	101482646	0.22	-0.65	15.04	T	A	0.17	Cam12W	5
GY	173	8	12767198	12798330	12782777	0.17	-0.61	11.8	C	T	0.22	Cam12W	4

"LowerLimit" and "UpperLimit" are the lower and upper physical limits for each QTL,

followed by the physical position (*Pos*, bp), proportion of the variance explained (R^2),

the effect of the major allele (*Effect*) as outputted by FastLMM, $-log_{10}($***Pval***$)$ (*Log*), the

minor and major alleles and the minor allele frequency (*MAF*) of the most significant

SNP within the QTL. The environment for which the most associated QTL was

observed (EnvMax) and the number of different environments (NbDiffEnv) that

detected the QTL are shown. Note that QTLs 71-72 for the plant height and QTLs

129-130 for the grain yield are genetically close (<1cM) and display high mean LD

(r^2K>0.5). Hence, QTLs 71-72 and 129-130 can potentially be merged.

The distribution of QTLs was not homogeneous along the genome since 82%, 77% and 79%

of flowering time, plant height and grain yield QTLs, respectively, were located in the high

recombinogenic regions, whereas they represented 46% of the physical genome (Additional

file 10: Table S5 and Figure S13). The QTLs were more stable (≥ 2 environments) in low than

in high recombinogenic regions (12.8% vs 5.8%, *p-value* = 0.03).

Discussion

GBS required massive imputation but displayed similar global trends than DNA arrays for genetic diversity organization

In order to reduce genotyping cost, GBS is most often performed at low depth leading to a

high proportion of missing data, thereby requiring imputation in order to perform GWAS.

Imputation can produce genotyping errors that can cause false associations and introduce bias

in diversity analysis [33]. We evaluated the quality of genotyping and imputation obtained by

different approaches, taking the 50K or 600K arrays as references. The best imputation

method that yielded a fully genotyped matrix with a low error rate for the prediction of both

heterozygotes and homozygotes was the approach merging the homozygous genotypes from

Tassel and the imputation of Beagle for the other data (GBS$_5$ in Additional file 1: Table S1).

The quality of imputation was high with 96% of allelic values consistent with those of the

50K and 600K arrays. This level of concordance is identical than in a study of USA national

maize inbred seed bank by Romay *et al.* [32]. It is higher than in a diversity study of

European flint maize collection (93%) by Gouesnard *et al.* [33], which was more distant from

the reference AllZeaGBSv2.7 database than for the panel presented here.

The ascertainment bias of arrays due to the limited number of lines used for SNP discovery

was reinforced by counter-selection of rare alleles during the design process of DNA arrays

[3, 4]. For GBS, the polymorphism database to call polymorphisms included thousands of

diverse lines [35]. In our study, we used AllZeaGBSv2.7 database. After a first step of GBS

imputation (GBS$_2$), missing data dropped to 11.9% *i.e.* only slightly more than in Romay *et*

al. (10%) [34]. This confirms that the polymorphism database (AllZeaGBSv2.7) covered

adequately the genetic diversity of our genetic material.

Although, we observed differences of allelic frequency spectrum between GBS and DNA

arrays, these technologies revealed similar trends in the organization of population structure

and relatedness (Figure 1, Additional file 3: Figure S3 and S4 and Table S2) suggesting no

strong ascertainment bias for deciphering global genetic structure trends in the panel.

However, although highly correlated, level of relatedness differed between GBS and DNA

arrays, especially when the lines were less related as showed by the deviation (to the left) of

the linear regression from the bisector (Figure 3).

The extent of linkage disequilibrium strongly varied along and between chromosomes

Linkage disequilibrium extent in high recombinogenic regions varied to a large extent among

chromosomes, ranging from 0.012 to 0.062 cM. Similar variation of genetic LD extent

between maize chromosomes has been previously observed by Rincent et al. [14], but their

classification of chromosomes was different from ours. This difference could be explained by

the fact that we analyzed specifically high and low recombination regions. According to Hill

and Weir Model [37], the physical LD extent in a genomic region increased when the local

recombination rate decreased. As a consequence, chromosome 1 and 9 had the lowest and

highest physical LD extent and displayed the highest and one of the lowest recombination rate

in pericentromeric regions, respectively (0.26 vs 0.11 cM / Mbp, Table 1 and Additional file

10: Table S5). Unexpectedly, the genetic LD extent also correlated negatively with the

recombination rate. It suggested that chromosomes with a low recombination rate also display

a low effective population size. Background selection for deleterious alleles could explain this

pattern since it reduced the genetic diversity in low recombinogenic regions [38, 39]. Finally,

we observed a strong variation of the LD extent along each chromosome (Additional file 4:

Figure S8). As we used a consensus genetic map [40] that represents well the recombination

within our population, it suggested, according to Hill and Weir's model, that the number of

ancestors contributing to genetic diversity varied strongly along the chromosomes. This likely

reflects the selection of genomic regions for adaptation to environment or agronomic traits

[38], that leads to a differential contribution of ancestors according to their allelic effects.

Ancestors with strong favorable allele(s) in a genomic region may lead ultimately to large

identical by descent genomic segments [41].

SNPs were clustered into QTL highlighting interesting genomic regions

In previous GWAS, the closest associated SNPs were grouped into QTLs according to

either a fixed physical distance [1] or a fixed genetic distance [30, 42]. These approaches

suffer of two drawbacks. First, the physical LD extent can vary strongly along chromosomes

according to the variation of recombination rate (Additional file 2: Figure S2). Second, the

genetic LD extent depends both on panel composition and the position along the genome

(Table 1). These approaches may therefore strongly overestimate or underestimate the number

of QTLs. To address both issues Cormier *et al.* [43] proposed to group associated SNPs by

using a genetic window based on the genetic LD extent estimated by Hill and Weir model in

the genomic regions around the associated peaks [37]. In our study, we improved this last

approach (*LD_win):*

- First, we used *r^2K* that corrected r^2 for kinship rather than the classical r^2 since *r^2K*

reflected the LD addressed in our GWAS mixed models to map QTL [17].

- Second, we took advantage of the availability of both physical and genetic maps of

maize to project the genetic LD extent on the physical map. This physical window was useful

to retrieve the annotation from B73 reference genome, decipher local haplotype diversity

(Figure 6) and estimate physical genome coverage (Table 1, Figure 2).

- Third, we considered an average LD extent estimated separately in the high and low

recombinogenic genomic regions. This average was estimated by using several large random

sets of pairs of loci in these regions rather than the local LD extent in the genomic regions

around each associated peaks.

We preferred this approach rather than using local LD extent in order to limit the effect of (i)

the strong variation of marker density along the chromosome (Additional file 2: Figure S2),

(ii) the local ascertainment bias due to the markers sampling (iii) the poor estimation of the

local recombination rate using a genetic map, notably for low recombination regions [3, 41]

(iv) errors in locus order due to assembly errors or chromosomal rearrangements.

We compared *LD_win* with *LD_adj,* another approach based on LD to group the SNPs

associated to trait variation into QTL. The discrepancies between the two approaches can be

explained by the local recombination rate and LD pattern. Since *LD_adj* approach was based

on the grouping of contiguous SNPs according to their LD, this approach was highly sensitive

to (i) error in marker order or position due to genome assembly errors or structural variations,

which are important in maize [44] (ii) genotyping or imputation errors, which we estimated at

ca. 1% and *ca.* 4%, respectively, for GBS (Additional file 1: Table S1), (iii) presence of

allelic series with contrasted effects in different experiments which are currently observed in

maize [40], (iv) LD threshold used. On the other hand, *LD_win* lead either to inflate the

number of QTLs in high recombinogenic regions in which SNPs were too distant genetically

to be grouped, or deflated their number by grouping associated SNPs in low recombinogenic

regions. Since *LD_win* considered the average LD extent, this method could conduct either to

separate or group abusively SNPs when local LD extent were different than the global LD

extent.

Note that LD windows should not be considered as confidence intervals since the

relationship between LD and recombination is complex due to demography, drift and

selection in association panels, contrary to linkage based QTL mapping [17]. The magnitude

of the effect of causal polymorphism in the estimation of these intervals which is well

established for linkage mapping should be explored further [45]. Other approaches have been

proposed to cluster SNPs according to LD [46, 47]. These approaches aim at segmenting the

genome in different haplotype blocks separating by high recombination regions. These

methods are difficult to use for estimating putative windows inside which the causal

polymorphisms are because such approaches are not centered on the associated SNP.

Several QTLs identified by *LD_win* in our study correspond to regions previously

identified: in particular six regions associated with female flowering time [27] and 30 regions

associated with different traits in the Cornfed dent panel [11]. Conversely, we did not identify

in our study any QTL associated to the florigen *ZCN8*, which showed significant effect in

these two previous studies. This relates most likely to the fact that we narrowed the flowering

time range in our study, in particular by eliminating early lines. This reduced the

representation of the early allele in the Zcn8, leading to a MAF of 0.27 in our study vs. 0.35 in

Rincent *et al.* [11], which can diminishes the power of the tests [14].

Complementarity of 600K and GBS for QTL detection resulted mostly from the tagging of different haplotypes rather than the coverage of different genomic regions.

Number of significant SNPs and QTLs increased with the increase in marker number (Table

3, Figure 4). This could be explained partly by a better coverage of some genomic regions by

SNPs, notably in high recombinogenic regions which showed a very short LD extent and were

enriched in QTLs (Additional file 10: Figure S13). Numerous new QTLs identified by the

600K array and GBS as compared with those identified by the 50K array were detected in

high recombinogenic regions that were considerably less covered by the 50K array than the

600K array or GBS (Additional file 2: Figure S2).

The high complementarity for QTL detection between GBS and 600K array was only

explained to a limited extent by the difference of the SNP distribution and density along the

genome, since these two technologies targeted similar regions as showed by coverage analysis

(Figure 2 and Additional file 2: Figure S2). However, at a finer scale, SNPs from the 600K

array and GBS could tag close but different genomic regions around genes. SNPs from the

600K array were mostly selected within coding regions of genes [4], whereas SNP from GBS

targeted more largely low copy regions, which included coding but also regulatory regions of

genes [32, 35]. To further analyse the complementarity of the technologies, we analysed local

haplotypes. We showed that both technologies captured different haplotypes when similar

genomic regions were targeted (Figure 6). Hence, we pinpointed that GBS and DNA arrays

are highly complementary for QTL detection because they tagged different haplotypes rather

than tagging different regions (Figure 6). Based on the L-shaped MAF distribution, which

suggest no ascertainment bias, and the high number of sequenced lines used for the GBS, we

expect a closer representation of the variation present in our panel by this technology

compared to the 600K array, but this comes to the cost of an enrichment in rare alleles. Both

factors tend to counterbalance each other in terms of GWAS power.

Our results suggest that we did not reach saturation with our *c.* 800,000 SNPs because (i)

some haplotypes certainly remain not tagged (ii) the genome coverage was not complete, and

(iii) the number of significant SNPs and QTLs continued to increase with marker density

(Figure 4). Considering LD and marker density, the genotypic data presently available were

most likely enough to well represent polymorphisms in the centromeric regions, whereas

using more markers would be beneficial for telomeric regions. New approaches based on

resequencing of representative lines and imputation are currently developed to achieve this

goal.

Methods

Plant Material and Phenotypic Data

The panel of 247 genotypes (Additional file 11: Table S6) includes 164 lines from a wider

panel of the CornFed project, composed of dent lines from Europe and America [11] and 83

additional lines derived from public breeding programs in Hungary, Italy and Spain and

recent lines free of patent from the USA. Lines were selected within a restricted window of

flowering time (10 days). Candidate lines with poor sample quality, i.e. high level of

heterozygosity, or high relatedness with other lines were discarded. The lines selection was

also guided by pedigree to avoid as far as possible over-representation of some parental

materials. These inbred lines were crossed with a common flint tester (UH007) and the

hybrids were evaluated for male flowering time (Day To Anthesis, DTA), plant height

(plantHT), and grain yield (GY) at seven sites in Europe, during two years (2012 and 2013),

and for two water treatments (watered and rainfed) [30]. The adjusted mean (Best Linear

Unbiased Estimation, BLUEs, https://doi.org/10.15454/IASSTN) of the three traits were

estimated per environment (site × year × treatment) using a mixed model with correction for

blocks, repetitions and rows and columns in order to take into account spatial variation of

micro-environment in each field trial [30]. Variance components and heritability of each traits

in each environment were also estimated [30] (Additional file 12: Table S7). Adjusted means

of hybrids were combined with genotyping data of the lines to perform GWAS.

Genotyping and Genotyping-By-Sequencing Data

The inbred lines were genotyped using three technologies: a maize Illumina Infinium HD 50K

array [3], a maize Affymetrix Axiom 600K array [4], and Genotyping-By-Sequencing [2, 35].

In the arrays, DNA fragments are hybridized with probes attached to the array that flanked

SNPs that have been previously identified between inbred lines (Additional file 5:

Supplementary Text 1 for the description of the data from the two arrays). Genotyping-by-

sequencing technology is based on multiplex resequencing of tagged DNA from different

individuals for which some genomic regions were targeted using restriction enzyme (Keygene

N.V. owns patents and patent applications protecting its Sequence Based Genotyping

technologies) [2]. Cornell Institute (NY, USA) processed raw sequence data using a multi-

step Discovery and a one-top Production pipeline (*TASSEL-GBS*) in order to obtain genotypes

(Additional file 5: Supplementary Text 2). An imputation step of missing genotypes was

carried out by Cornell Institute [36], which utilized an algorithm that searches for the closest

neighbour in small SNP windows across the haplotype library [35], allowing for a 5%

mismatch. If the requirements were not met, the SNP was left ungenotyped for individuals.

We applied different filters (heterozygosity rate, missing data rate, minor allele frequency) for

a quality control of the genetic data before performing the diversity and association genetic

analyses. For GBS data, the filters were applied after imputation using the method

"Compilation of Cornell homozygous genotypes and Beagle genotypes" (GBS$_5$ in Additional

file 1: Figure S1; See section "Evaluating Genotyping and Imputation Quality"). We

eliminated markers that had an average heterozygosity and missing data rate higher than 0.15

and 0.20, respectively, and a Minor Allele Frequency (MAF) lower than 0.01 for the diversity

analyses and 0.05 for the GWAS. Individuals which had heterozygosity and/or missing data

rate higher than 0.06 and 0.10, respectively, were eliminated.

Evaluating Genotyping and Imputation Quality

Estimating the genotyping and imputation quality were performed using 245 lines since two

inbred lines have different seedlots between technologies. The 50K and the 600K arrays were

taken as reference to compare the concordance of genotyping (genotype matches) with the

imputation of GBS based on their position. While SNP position and orientation from GBS

were called on the reference maize genome B73 AGP_v2 (release 5a) [48], flanking

sequences of SNPs in the 50K array were primary aligned on the first maize genome reference

assembly B73 AGP_v1 (release 4a.53) [49]. Both position and orientation scaffold carrying

SNPs from the 50K array can be different in the AGP_v2, which could impair correct

comparison of genotype between the 50K array and GBS. Hence, we aligned flanking

sequences of SNPs from the 50K array on maize B73 AGP_v2 using the Basic Local

Alignment Search Tool (BLAST) to retrieve both positions and genotype in the same and

correct strand orientation (forward) to compare genotyping. The number of common markers

between the 50K/600K, 50K/GBS, GBS/600K and 50K/600K/GBS was 36,395, 7,018,

25,572 and 5,947 SNPs, respectively. The comparison of the genotyping and imputation

quality between the 50K/GBS, 50K/600K and 600K/GBS was done on 5,336 and 24,286

PANZEA markers [50] in common, and 26,154 markers in common, respectively. The

genotyping concordance of the 600K with the 50K array was extremely high (99.50%) but

slightly lower for heterozygotes (92.88%). In order to achieve these comparisons, we

considered the direct reads from GBS (**GBS$_1$**) and four approaches for imputation (GBS$_2$ to

GBS$_5$, Additional file 1: Figure S1). **GBS$_2$** approach consisted in one imputation step from the

direct read by Cornell University, using *TASSEL* software, but missing data was still present.

GBS$_3$ approach consisted in a genotype imputation of the whole missing data of the direct

read by *Beagle v3* [13]. In **GBS$_4$,** genotype imputation by Beagle was performed on Cornell

imputed data after replacing the heterozygous genotypes into missing data. **GBS$_5$**, consisted in

homozygous genotypes of GBS$_2$ completed by values imputed in GBS$_3$ (Additional file 1:

Figure S1).

Diversity Analyses

After excluding the unplaced SNPs and applying the filtering criteria for the diversity

analyses (MAF $\geq$ 0.01), we obtained the final genotyping data of the 247 lines with 44,732

SNPs from the 50K array, 506,662 SNPs from the 600K array, and 395,053 SNPs from the

GBS (Figure 1). All markers of the 600K array and GBS$_5$ that passed the quality control were

used to perform the diversity analyses (estimation of Q genetic groups and K kinships). For

the 50K, we used only the PANZEA markers (29,257 SNPs) [50] in order to reduce the

ascertainment bias noted by Ganal *et al.* [3] when estimating Nei's index of diversity [51] and

relationship coefficients. Genotypic data generated by the three technologies were organized

as *G* matrices with *N* rows and *L* columns, *N* and *L* being the panel size and number of

markers, respectively. Genotype of individual *i* at marker *l* ($G_{i,l}$) was coded as 0 (the

homozygote for an arbitrarily chosen allele), 0.5 (heterozygote), or 1 (the other homozygote).

Identity-By-Descend (IBD) was estimated according to Astle and Balding [19]:

$$K\_Freq_{i,j} = \frac{1}{L}\sum_{l=1}^{L}\frac{(G_{i,l}-p_l)(G_{j,l}-p_l)}{p_l(1-p_l)},$$

where p_l is the frequency of the allele coded 1 of marker *l* in the panel of interest, *i*

and *j* indicate the inbred lines for which the kinship was estimated. We also estimated the

Identity-By-State (IBS) by estimating the proportion of shared alleles. For GWAS, we used

K_Chr [14] that are computed using similar formula as *K_Freq*, but with the genotyping data

of all the chromosomes except the chromosome of the SNP tested. This formula provides an

unbiased estimate of the kinship coefficient and weights by allelic frequency assuming Hardy-

Weinberg equilibrium. Hence, relatedness is higher if two individuals share rare alleles than

common alleles.

Genetic structure was analysed using the sofware *ADMIXTURE v1.22* [18] with a

number of groups varying from 2 to 10 for the three technologies. We compared assignation

by *ADMIXTURE* of inbred lines between the three technologies by estimating the proportion

of inbred lines consistently assigned between technologies two by two (50K *vs* GBS$_5$, 50K *vs*

600K, 600K *vs* GBS$_5$) using a threshold of 0.5 for admixture.

Expected heterozygosity (He) [51] was estimated at each marker as $2p_l(1 - p_l)$ and

was averaged on all the markers for a global characterization of the panel for the three

technologies. Principal Coordinate Analyses (PCoA) were performed on the genetic distance

matrices [52], estimated as $1_{N,N} - K\_Freq$, where $1_{N,N}$ is a matrix of ones of the same size as

K_Freq.

Linkage Disequilibrium Analyses

We first analyzed the effect of the genetic structure and kinship on linkage disequilibrium

(LD) extent within and between chromosomes by estimating genome-wide linkage

disequilibrium using the 29,257 PANZEA SNPs from the 50K array. Four estimates of LD

were used: the squared correlation (r^2) between allelic dose at two markers [53], the squared

correlation taking into account global kinship with *K_Freq* estimator (*r²K*), the squared

correlation taking into account population structure (*r²S*), and the squared correlation taking

into account both (*r²KS*) [17].

To explore the variation of LD decay and the stability of LD extent along the chromosomes,

we estimated LD between a non-redundant set of 810,580 loci from the GBS, the 50K and

600K arrays. To save computation time, we calculated LD between loci within a sliding

window of 1 cM. Genetic position was obtained by projecting the physical position of each

locus using a *smooth.spline* function R calibrated on the genetic consensus map of the

Cornfed Dent Nested Association Mapping (NAM) design [40]. We used the estimator r^2 and

r²K using 10 different kinships *K_Chr*. This last estimator was calculated because it

corresponds exactly to LD used to map QTL in our GWAS model. It determines the power of

GWAS to detect QTL considering that causal polymorphisms were in LD with some

polymorphisms genotyped in our panel [17]. To study LD extent variation, we estimated LD

extent by adjusting Hill and Weir's model [37] using non-linear regression (*nls* function in R-

package *nlme*) against both physical and genetic position within each chromosome. Since

recombination rate (cM / Mbp) varied strongly along the genome (Figure 2 and Additional

file 2: Figure S2), we defined high (>0.5 cM / Mbp) and low (<0.5 cM / Mbp)

recombinogenic genomic regions within each chromosome. We adjusted Hill and Weir's

model [37] separately in low and high recombinogenic regions (Additional file 10: Table S5)

by randomly sampling 100 sets of 500,000 pairs of loci distant from less than 1 cM. This

random sampling avoided over-representation of pairs of loci from low recombinogenic

regions due to the sliding-window approach (Additional file 12: Figure S14). 500,000 pairs of

loci represented 0.36% (Chromosome 3 / High rec) to 1.20% of all pairs of loci (Chromosome

8 / High rec).

For all analyses, we estimated LD extent by calculating the genetic and physical distance for

the fitted curve of Hill and Weir's Model that reached r^2K=0.1, r^2K=0.2 and r^2K=0.4.

Genome coverage estimation

In order to estimate the genomic regions in which the effect of an underlying causal

polymorphisms could be captured by GWAS using LD with SNP from three technologies, we

developed an approach to define LD windows around each SNP with MAF ≥ 5% based on

LD extent (Additional file 12: Figure S14). To set the LD window around each SNP, we used

LD extent with r^2K=0.1 (negligible LD), r^2K=0.2 (intermediate LD) and r^2K=0.4 (high LD)

estimated in low and high recombinogenic regions for each chromosome. We used the global

LD decay estimated for these large chromosomal regions rather than local LD extent (i) to

avoid bias due to SNP sampling within small genomic regions, (ii) to reduce computational

time, and (iii) to limit the impact of possible local error in genome assembly. In low

recombinogenic regions, we used the physical LD extent, hypothesizing that recombination

rate is constant along physical distance in these regions. In high recombinogenic regions, we

used the genetic LD extent since there is a strong variation of recombination rate by base pair

along the physical position (Additional file 2: Figure S2). We then converted genetic LD

windows into physical windows by projecting the genetic positions on the physical map using

the *smooth.spline* function implemented in R calibrated on the NAM dent consensus map

[40]. Reciprocally, we obtained the genetic positions of LD windows in low recombinogenic

regions by projecting the physical boundaries of LD windows on the genetic map.

To estimate coverage of the three technologies to detect QTLs based on their SNP distribution

and density, we calculated cumulative genetic and physical length that are covered by LD

windows around the markers, considering different LD extents for each chromosome

(r^2K=0.1, r^2K=0.2, r^2K=0.4). In order to explore variation of genome coverage along the

chromosome, we estimated the proportion of genome covered using a sliding-windows

approach based on physical distance (2Mbp).

Statistical Models for Association Mapping

We used four models to determine the statistical models that control best the confounding

factors (*i.e.* population structure and relatedness) in GWAS (Additional file 5: Supplementary

Texts 3 and 4). We tested different software implementing either approximate (EMMAX) [8]

or exact computation of standard test statistics (ASReml and FaST-LMM) [6, 54] for

computational time and GWAS results differences (Additional file 5: Supplementary Text 5).

Single-trait, single-environment GWAS was performed for each marker for each environment

and all traits using FaST-LMM. We selected the mixed model using *K_Chr*, estimated from

PANZEA markers of the 50K array to perform GWAS on 66 situations (environment $\times$ trait)

(Additional file 5: Supplementary Text 4, Additional file 12: Figure S15 and Additional file

12: Figure S16). We developed a GWAS pipeline in *R v3.2.1* [55] calling FaST-LMM

software and implementing [14] approaches to conduct single trait and single environment

association tests.

Multiple testing is a major challenge in GWAS using large numbers of markers. The

experiment-wise error rate (α_e) increases with the number of tests (number of markers) carried

out, even when the point-wise error rate (α_p) is maintained low. Popular methods [56, 57] are

overly conservative and can result in overlooking true positive associations. In addition, these

corrections assume that the hypothesis tests are independent. To take into account the

dependence of the tests in GWAS, α_p has to be adjusted in order to keep α_e at a nominal level.

Moskvina and Schmidt [58] and Gao *et al.* [59, 60] corrections can correctly infer the number

of independent tests and use the Bonferroni formula to rapidly adjust for multiple testing.

Using Gao approaches, we estimated the number of independent tests for GWAS at 15,780

for the 50K, 92,752 for the 600K, 109,117 for the GBS_5 and 191,026 for the combined genetic

data, leading to different *-log₁₀(p-value)* thresholds: 5.49, 6.27, 6.34 and 6.58, respectively.

Because of these differences, we used two thresholds of *-log₁₀(p-value) = 5* (less stringent)

and 8 (hightly conservative and slightly above Bonferroni) for comparing GWAS to avoid the

differences of identification of significant SNPs between the technologies due to the choice of

the threshold.

Methods for grouping associated SNPs into QTLs

We used two approaches based on LD for grouping significant SNPs. The first approach

(*LD_win*) used LD windows, previously described, to group significant SNPs into QTLs

considering that all SNPs with overlapping LD windows of $r^2K=0.1$ belong to the same QTL.

We hypothesized that significant SNPs with overlapping LD windows at $r^2K=0.1$ captured the

same causal polymorphism and were therefore a single and unique QTL. The second

approach (*LD_adj*) grouped into single QTL significant SNPs that are adjacent on the

physical map providing that their LD were above a LD threshold ($r^2K > 0.5$). We used LD

heatmaps for comparing the SNP grouping produced by the two approaches on the three

different traits across all environments (Additional file 6: Figure S9-LD-Adjacent and

Additional file 7: Figure S9-LD-Windows). All scripts are implemented in R software [55].

List of abreviations

DTA = Day to Anthesis

GY = Grain Yield adjusted at 15% moisture

plantHT = Plant Height

GBS = Genotyping By Sequencing

LD = Linkage disequilibrium

GWAS = Genome-Wide Association Studies

MAF = Minimum Allelic Frequency

SNP = Single Nucleotide Polymorphism

HRR = High Recombinogenic Regions

LRR = Low Recombinogenic Regions

QTL = Quantitative Trait Locus

Declarations

Ethics approval and consent to participate

773   Not applicable.

Consent for publication

774   Not applicable.

Availability of data and material

775   The following links toward the data will be available upon publication of this paper.

776   All the genotyping data used in this study can be found at https://doi.org/10.15454/AEC4BN.

777   The GWAS results can be found at https://doi.org/10.15454/6TL2N4.

778   The phenotypic dataset can be found at https://doi.org/10.15454/IASSTN.

Competing interests

779   The authors declare that they have no competing interests.

Funding

780   This project (Project ID: 244374) was funded under the European FP7- KBBE (CP – IP –

781   Large-scale integrating project, DROPS) and the *Agence Nationale de la Recherche* project

782   ANR-10-BTBR-01 (ANR-PIA AMAIZING).

Authors' contributions

S.S.N., S.D.N. and A.C., designed the studied and wrote the article. S.S.N. performed

genotyping data quality control, imputation and genetic analyses. S.D.N. developed and

performed LD analyses. A.C. designed the association panel with the help of S.D.N. and C.W.

C.B. participated in assembling the dent inbred lines panel, organizing the germplasms and

field work for seeds production. E.J.M., C.W. and F.T. collected and analysed the phenotypic

data. V.C. and D.M. performed DNA extraction and prepared the samples. All authors

critically reviewed and approved the final manuscript.

Acknowledgements

We are grateful to key partners from the field: Pierre Dubreuil, Cécile Richard, Jérémy Lopez

(Biogemma), Tamás Spitkó (MTA ATK), Therese Welz (KWS), Franco Tanzi, Ferenc Racz,

Vincent Schlegel (Syngenta) and Maria Angela Canè (UNIBO). We also acknowledge Björn

Usadel and Axel Nagel (MPI) for data management. We thank Willem Kruijer, Fred Van

Eeuwijk (WUR), Tristan Mary-Huard and Laurence Moreau (INRA) for helpful discussions

and statistical advice. We are grateful to Chris-Carolin Schön (TUM) for providing an early

access to the Affymetrix Axiom 600K array and Edward Buckler (USDA) for providing

genotyping using GBS. We are also grateful to partners of the CornFed project, Univ.

Hohenheim (Germany), CSIC (Spain), CRAG (Spain), MTA ATK (Hungary), NCRPIS

(USA), CRB Maize (France) and CRA-MAC (Italy) who contributed to the genetic material.

Authors' information (optional)

800   Not applicable.

801   **References**

804   1.   Tian F, Bradbury PJ, Brown PJ, Hung H, Sun Q, Flint-Garcia S, Rocheford TR,

805   McMullen MD, Holland JB, Buckler ES: **Genome-wide association study of leaf**

806   **architecture in the maize nested association mapping population.** *Nature Genetics*

807   2011, **43:**159.

808   2.   Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE: **A**

809   **Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity**

810   **Species.** *PLoS ONE* 2011, **6:** e19379.

811   3.   Ganal MW, Durstewitz G, Polley A, Bérard A, Buckler ES, Charcosset A, Clarke JD,

812   Graner E-M, Hansen M, Joets J, et al: **A Large Maize (Zea mays L.) SNP**

813   **Genotyping Array: Development and Germplasm Genotyping, and Genetic**

814   **Mapping to Compare with the B73 Reference Genome.** *PLoS ONE* 2011, **6:**

815   e28334.

816   4.   Unterseer S, Bauer E, Haberer G, Seidel M, Knaak C, Ouzunova M, Meitinger T,

817   Strom TM, Fries R, Pausch H, et al: **A powerful tool for genome analysis in maize:**

818   **development and evaluation of the high density 600 k SNP genotyping array.**

819   *BMC Genomics* 2014, **15:**823.

820   5.   Zhou X, Stephens M: **Genome-wide efficient mixed-model analysis for association**

821   **studies.** *Nature Genetics* 2012, **44:**821.

822   6.   Lippert C, Listgarten J, Liu Y, Kadie CM, Davidson RI, Heckerman D: **FaST linear**

823   **mixed models for genome-wide association studies.** *Nature Methods* 2011, **8:**833.

7.   Listgarten J, Lippert C, Kadie CM, Davidson RI, Eskin E, Heckerman D: **Improved**

linear mixed models for genome-wide association studies. *Nature Methods* 2012,

9:525.

8.   Kang HM, Sul JH, Service SK, Zaitlen NA, Kong S-y, Freimer NB, Sabatti C, Eskin

E: **Variance component model to account for sample structure in genome-wide**

association studies. *Nature Genetics* 2010, **42:**348.

9.   Flint-Garcia SA, Thornsberry JM, Buckler ES: **Structure of Linkage Disequilibrium**

in Plants. *Annual Review of Plant Biology* 2003, **54:**357-374.

10.   Remington DL, Thornsberry JM, Matsuoka Y, Wilson LM, Whitt SR, Doebley J,

Kresovich S, Goodman MM, Buckler ES: **Structure of linkage disequilibrium and**

phenotypic associations in the maize genome. *Proceedings of the National Academy*

of Sciences 2001, **98:**11479.

11.   Rincent R, Nicolas S, Bouchet S, Altmann T, Brunel D, Revilla P, Malvar RA,

Moreno-Gonzalez J, Campo L, Melchinger AE, et al: **Dent and Flint maize diversity**

panels reveal important genetic potential for increasing biomass production.

Theoretical and Applied Genetics 2014, **127:**2313-2331.

12.   Yu J, Pressoir G, Briggs WH, Vroh Bi I, Yamasaki M, Doebley JF, McMullen MD,

Gaut BS, Nielsen DM, Holland JB, et al: **A unified mixed-model method for**

association mapping that accounts for multiple levels of relatedness. *Nature*

Genetics 2006, **38:**203.

13.   Browning SR, Browning BL. **Rapid and Accurate Haplotype Phasing and Missing-**

Data Inference for Whole-Genome Association Studies By Use of Localized

Haplotype Clustering. The American Journal of Human Genetics 2007, **81** (5): 1084-

1097

14.   Rincent R, Moreau L, Monod H, Kuhn E, Melchinger AE, Malvar RA, Moreno-

Gonzalez J, Nicolas S, Madur D, Combes V, et al: **Recovering Power in Association**

Mapping Panels with Variable Levels of Linkage Disequilibrium. *Genetics* 2014,

197:375.

15. Van Inghelandt D, Melchinger AE, Lebreton C, Stich B: **Population structure and**

genetic diversity in a commercial maize breeding program assessed with SSR and

SNP markers. *Theoretical and Applied Genetics* 2010, **120:**1289-1299.

16. Nicolas SD, Péros J-P, Lacombe T, Launay A, Le Paslier M-C, Bérard A, Mangin B,

Valière S, Martins F, Le Cunff L, et al: **Genetic diversity, linkage disequilibrium**

and power of a large grapevine (Vitis vinifera L) diversity panel newly designed

for association studies. *BMC Plant Biology* 2016, **16:**74.

17. Mangin B, Siberchicot A, Nicolas S, Doligez A, This P, Cierco-Ayrolles C: **Novel**

measures of linkage disequilibrium that correct the bias due to population

structure and relatedness. *Heredity* 2012, **108:**285.

18. Alexander DH, Novembre J, Lange K: **Fast model-based estimation of ancestry in**

unrelated individuals. *Genome Research* 2009, **19:**1655-1664.

19. Astle W, Balding DJ: **Population Structure and Cryptic Relatedness in Genetic**

Association Studies. *Statist Sci* 2009, **24:**451-471.

20. Pritchard JK, Stephens M, Rosenberg NA, Donnelly P: **Association Mapping in**

Structured Populations. *The American Journal of Human Genetics* 2000, **67:**170-

181.

21. VanRaden PM: **Efficient Methods to Compute Genomic Predictions.** *Journal of*

Dairy Science 2008, **91:**4414-4423.

22. Bernardo R: **Genomewide Markers for Controlling Background Variation in**

Association Mapping. *The Plant Genome* 2013, **6**.

23.   Thornsberry JM, Goodman MM, Doebley J, Kresovich S, Nielsen D, Buckler ES:

874   **Dwarf8 polymorphisms associate with variation in flowering time.** *Nature*

875   *Genetics* 2001, **28:**286.

876   24.   Hufford MB, Lubinksy P, Pyhäjärvi T, Devengenzo MT, Ellstrand NC, Ross-Ibarra J:

877   **The Genomic Signature of Crop-Wild Introgression in Maize.** *PLOS Genetics*

878   2013, **9:**e1003477.

879   25.   Rincent R, Laloë D, Nicolas S, Altmann T, Brunel D, Revilla P, Rodríguez VM,

880   Moreno-Gonzalez J, Melchinger A, Bauer E, et al: **Maximizing the Reliability of**

881   **Genomic Selection by Optimizing the Calibration Set of Reference Individuals:**

882   **Comparison of Methods in Two Diverse Groups of Maize Inbreds (*Zea Mays L.*).**

883   *Genetics* 2012, **192:**715.

884   26.   Bouchet S, Bertin P, Presterl T, Jamin P, Coubriche D, Gouesnard B, Laborde J,

885   Charcosset A: **Association mapping for phenology and plant architecture in maize**

886   **shows higher power for developmental traits compared with growth influenced**

887   **traits.** *Heredity* 2017, **118:**249-259.

888   27.   Bouchet S, Servin B, Bertin P, Madur D, Combes V, Dumas F, Brunel D, Laborde J,

889   Charcosset A, Nicolas S: **Adaptation of Maize to Temperate Climates: Mid-**

890   **Density Genome-Wide Association Genetics and Diversity Patterns Reveal Key**

891   **Genomic Regions, with a Major Contribution of the Vgt2 (ZCN8) Locus.** *PLoS*

892   *ONE* 2013, **8:**e71377.

893   28.   Messing J, Dooner HK: **Organization and variability of the maize genome.** *Current*

894   *Opinion in Plant Biology* 2006, **9:**157-163.

895   29.   Hu H, Schrag TA, Peis R, Unterseer S, Schipprack W, Chen S, Lai J, Yan J, Prasanna

896   BM, Nair SK, et al: **The Genetic Basis of Haploid Induction in Maize Identified**

with a Novel Genome-Wide Association Method. *Genetics* 2016, **202:**1267.

30.   Millet EJ, Welcker C, Kruijer W, Negro S, Coupel-Ledru A, Nicolas SD, Laborde J,

Bauland C, Praud S, Ranc N, et al: **Genome-Wide Analysis of Yield in Europe:**

Allelic Effects Vary with Drought and Heat Scenarios. *Plant Physiology* 2016,

172:749.

31.   Crossa J, Beyene Y, Kassa S, Pérez P, Hickey JM, Chen C, de los Campos G,

Burgueño J, Windhausen VS, Buckler E, et al: **Genomic Prediction in Maize**

Breeding Populations with Genotyping-by-Sequencing. *G3:*

Genes|Genomes|Genetics 2013, **3:**1903.

32.   Romay MC, Millard MJ, Glaubitz JC, Peiffer JA, Swarts KL, Casstevens TM, Elshire

RJ, Acharya CB, Mitchell SE, Flint-Garcia SA, et al: **Comprehensive genotyping of**

the USA national maize inbred seed bank. *Genome Biology* 2013, **14:**R55.

33.   Gouesnard B, Negro S, Laffray A, Glaubitz J, Melchinger A, Revilla P, Moreno-

Gonzalez J, Madur D, Combes V, Tollon-Cordet C, et al: **Genotyping-by-sequencing**

highlights original diversity patterns within a European collection of 1191 maize

flint lines, as compared to the maize USDA genebank. *Theoretical and Applied*

Genetics 2017, **130:**2165-2189.

34.   Frascaroli E, Schrag TA, Melchinger AE: **Genetic diversity analysis of elite**

European maize (Zea mays L.) inbred lines using AFLP, SSR, and SNP markers

reveals ascertainment bias for a subset of SNPs. *Theoretical and Applied Genetics*

2013, **126:**133-141.

35.   Glaubitz JC, Casstevens TM, Lu F, Harriman J, Elshire RJ, Sun Q, Buckler ES:

TASSEL-GBS: A High Capacity Genotyping by Sequencing Analysis Pipeline.

PLoS ONE 2014, **9:**e90346.

36. Swarts K, Li H, Romero Navarro JA, An D, Romay MC, Hearne S, Acharya C,

Glaubitz JC, Mitchell S, Elshire RJ, et al: **Novel Methods to Optimize Genotypic**

Imputation for Low-Coverage, Next-Generation Sequence Data in Crop Plants.

The Plant Genome 2014, **7**.

37. Hill WG, Weir BS: **Variances and covariances of squared linkage disequilibria in**

finite populations. *Theoretical Population Biology* 1988, **33:**54-78.

38. Charlesworth D, Willis JH: **The genetics of inbreeding depression.** *Nature Reviews*

Genetics 2009, **10:**783.

39. Hudson RR, Kaplan NL: **Deleterious background selection with recombination.**

Genetics 1995, **141:**1605.

40. Giraud H, Bauland C, Falque M, Madur D, Combes V, Jamin P, Monteil C, Laborde J,

Palaffre C, Gaillard A, et al: **Reciprocal Genetics: Identifying QTL for General**

and Specific Combining Abilities in Hybrids Between Multiparental Populations

from Two Maize (*Zea mays L.*) Heterotic Groups. *Genetics* 2017, **207:**1167.

41. Bauer E, Falque M, Walter H, Bauland C, Camisan C, Campo L, Meyer N, Ranc N,

Rincent R, Schipprack W, et al: **Intraspecific variation of recombination rate in**

maize. *Genome Biology* 2013, **14:**R103.

942 41. Tian F, Bradbury PJ, Brown PJ, Hung H, Sun Q, Flint-Garcia S, Rocheford TR,

943 McMullen MD, Holland JB, Buckler ES: **Genome-wide association study of leaf**

944 **architecture in the maize nested association mapping population.** *Nature Genetics*

945   2011, **43:**159.

946   42.   Le Gouis J, Bordes J, Ravel C, Heumez E, Faure S, Praud S, Galic N, Remoué C,

947   Balfourier F, Allard V, Rousset M: **Genome-wide association analysis to identify**

948   **chromosomal regions determining components of earliness in wheat.** *Theoretical*

949   *and Applied Genetics* 2012, **124:**597-611.

950   43.   Cormier F, Le Gouis J, Dubreuil P, Lafarge S, Praud S: **A genome-wide**

951   **identification of chromosomal regions determining nitrogen use efficiency**

952   **components in wheat (Triticum aestivum L.).** *Theoretical and Applied Genetics*

953   2014, **127:**2679-2693.

954   44.   Springer NM, Ying K, Fu Y, Ji T, Yeh C-T, Jia Y, Wu W, Richmond T, Kitzman J,

955   Rosenbaum H, et al: **Maize Inbreds Exhibit High Levels of Copy Number**

956   **Variation (CNV) and Presence/Absence Variation (PAV) in Genome Content.**

957   *PLOS Genetics* 2009, **5:**e1000734.

958   45.   Darvasi A, Weinreb A, Minke V, Weller JI, Soller M: **Detecting marker-QTL**

959   **linkage and estimating QTL gene effect and map location using a saturated**

960   **genetic map.** *Genetics* 1993, **134:**943.

961   46.   Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J,

962   DeFelice M, Lochner A, Faggart M, et al: **The Structure of Haplotype Blocks in the**

963   **Human Genome.** *Science* 2002, **296:**2225.

964   47.   Wang H, Chu WS, Hemphill C, Elbein SC: **Human Resistin Gene: Molecular**

965   **Scanning and Evaluation of Association with Insulin Sensitivity and Type 2**

966   **Diabetes in Caucasians.** *The Journal of Clinical Endocrinology & Metabolism* 2002,

967   **87:**2520-2524.

48.   Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, Liang C, Zhang J,

Fulton L, Graves TA, et al: **The B73 Maize Genome: Complexity, Diversity, and**

Dynamics. *Science* 2009, **326:**1112.

49.   Ganal MW, Altmann T, Röder MS: **SNP identification in crop plants.** *Current*

Opinion in Plant Biology 2009, **12:**211-217.

50.   Gore MA, Chia J-M, Elshire RJ, Sun Q, Ersoz ES, Hurwitz BL, Peiffer JA, McMullen

MD, Grills GS, Ross-Ibarra J, et al: **A First-Generation Haplotype Map of Maize.**

Science 2009, **326:**1115.

51.   Nei M: **Estimation of Average Heterozygosity and Genetic Distance from a Small**

Number of Individuals. *Genetics* 1978, **89:**583.

52.   Gower JC: **Some distance properties of latent root and vector methods used in**

multivariate analysis. *Biometrika* 1966, **53:**325-338.

53.   Hill WG, Robertson A: **Linkage disequilibrium in finite populations.** *Theoretical*

and Applied Genetics 1968, **38:**226-231.

54.   Butler DG, Cullis BR, Gilmour AR, Gogel BJ: **ASReml-R reference manual.** *The*

State of Queensland, Department of Primary Industries and Fisheries, Brisbane,

Australia 2009.

55.   R Core Team: **R: A language and environment for statistical computing.** *R*

Foundation for Statistical Computing, Vienna, Austria 2015**:**URL https://www.R-

project.org/.

56.   Bonferroni CE: **Teoria statistica delle classi e calcolo delle probabilità.**

Pubblicazioni dell'Istituto Superiore di Scienze Economiche e Commerciali di Firenze

1936, **8:**3-62.

57.   Šidák Z: **Rectangular Confidence Regions for the Means of Multivariate Normal**

Distributions. *Journal of the American Statistical Association* 1967, **62:**626-633.

58.   Moskvina V, Schmidt KM: **On multiple-testing correction in genome-wide**

association studies. *Genetic Epidemiology* 2008, **32:**567-573.

59.   Gao X, Becker LC, Becker DM, Starmer JD, Province MA: **Avoiding the high**

Bonferroni penalty in genome-wide association studies. *Genetic Epidemiology*

2010, **34:**100-105.

60.   Gao X, Starmer J, Martin ER: **A multiple testing correction method for genetic**

association studies using correlated single nucleotide polymorphisms. *Genetic*

Epidemiology 2008, **32:**361-369.

1001

1002

Figure legends

Figure 1: Comparison of genotyping data between 50K and 600K arrays, and GBS. (a)

Distribution of minor allele frequency per SNP before filtering (monomorphic SNPs

removed). (b) Distribution of SNP missing data proportion for the 50K array, 600K array,

GBS direct reads (GBS$_1$) and GBS after imputation by Cornell Institute (GBS$_2$, note that the

scale of the x-axes is different). (**c**) Relatedness distribution (Identity-By-State, IBS) after QC

filtering with MAF≥1% (IBS using GBS$_1$ was not estimated because of the low calling rate).

1010

Figure 2: Variation of the markers density (top), the recombination rate (middle) and the

genome coverage (bottom). Non-overlapping 2Mbp windows along the chromosome 3 were

used. The percentage of genome coverage used the cumulative length of LD windows

calculated around each SNP. Markers along chromosome 3 have MAF ≥ 5%. Green, blue, red

and black lines represent variation of GBS, 600K, 50K and combined technologies,

respectively.

1017

1018 **Figure 3:** Correlation (r) of the Identity-By-State (IBS) between the three technologies (after

1019 imputation). (**a**) IBS_{600K} vs IBS_{50K}, (**b**) IBS_{GBS} vs IBS_{50K}, (**c**) IBS_{GBS} vs IBS_{600K}. The red line

1020 indicates the bisector.

1021

Figure 4: Number of significant SNPs (blue line) and QTLs (red line) identified as a function

of SNP density (x-axis) for the three traits (DTA, male flowering time; plantHT, plant height;

GY, grain yield).

Figure 5: Complementarity of the three technologies to detect QTLs. The numbers of specific

QTLs detected by each technologies for the three traits (flowering time, plant height, grain

yield) are shown.

Figure 6: Complementarity of QTLs detection between the 600K array and the GBS for two

regions (QTL 32/QTL95). (**a**) Manhattan plot of the $-log_{10}(p$-value) along the genome. Dotted

red lines correspond to QTL32 and QTL95 located on chromosome 1 and 3, respectively, for

the flowering time in one environment (Ner13R). (**b**) Local manhattan plot of the $-log_{10}(p$-

value) (top) and linkage disequilibrium corrected by the kinship (r^2K) (bottom) of all SNPs

with the strongest associated marker within QTL 32 (left) and QTL 95 (right). (**c**) Local

haplotypes displayed by SNPs within the QTLs 32 (left) and 95 (right). Inbred lines are in

rows and SNPs are in columns. Inbred lines were ordered by hierarchical clustering based on

local dissimilarity estimated by all SNPs within each QTLs. Genotyping matrix is colored

according to their allelic dose at each SNP. Red and black represent homozygotes and gray

represent heterozygotes. The associated peaks (red vertical lines) and other associated SNPs

with $-log_{10}(p$-*value*$) > 5$ (orange vertical lines) are indicated above the genotyping matrix. H1,

H2, H3, H4, H5 represent the 5 and 3 haplotypes obtained by cutting the dendograms with the

most 5 and 3 dissimilar clusters within QTL32 and QTL95, respectively.

1042

1043   **Additional file legends**

Additional file 1 (.docx):

Figure S1: Different approaches used to compare the quality of genotyping and imputation of

the GBS. We considered the direct reads from GBS (**GBS$_1$**) and four approaches for

imputation (GBS$_2$ to GBS$_5$). **GBS$_2$** approach consisted in one imputation step from the direct

read by Cornell University, using *TASSEL* software, but missing data was still present. **GBS$_3$**

approach consisted in a genotype imputation of the whole missing data of the direct read by

Beagle v3. In **GBS$_4$,** genotype imputation by Beagle was performed on Cornell imputed data

after replacing the heterozygous genotypes into missing data. **GBS$_5$,** consisted in homozygous

genotypes of GBS$_2$ completed by values imputed in GBS$_3$.

Table S1: Percentage of GBS concordance based on the 50K and 600K arrays (Reference).

Call rate of SNPs from GBS are in brackets. * After Beagle inference of missing data, the call

rate is 100%. Here the call rate is <100% because the comparison was made against the 50K

and the 600K arrays that include few missing data.

1057

Additional file 2 (.pdf):

Figure S2: Variation of the markers density, the recombination rate and the genome coverage

in non-overlapping 2 Mbp windows along each chromosome. The percentage SNP coverage

(bottom) used the cumulated length of physical LD windows around each SNP. Markers have

MAF $\geq$ 5%. Green, blue, red and black lines represent variation of GBS, 600K, 50K and

combined technologies, respectively.

1064

Additional file 3 (.docx):

Figure S3: Contribution of four ancestral populations to 247 inbred lines after ADMIXTURE

analysis. Markers from the 50K array (top), 600K array (middle) and GBS (bottom) were

used. One vertical bar corresponds to one individual. Lines were ordered according to

contributions observed for the 50K array. From left to right, we have Stiff Stalk lines type

B73 and B14a (red), Iodent lines type PH207 (green), Lancaster lines type Mo17 and Oh43

(turquoise), a group of lines assembling W117, F7057 type lines (blue).

Table S2: Means and ranges of the two relatedness estimators (IBS and IBD *i.e.* K_Freq)

from the 50K (29,257 PANZEA SNPs only) and 600K arrays, and GBS.

Figure S4: Correlation (r) between the IBS and IBD (K_Freq) for each technology (A) and

correlation of IBD between the three technologies (B). (C) Correlation of IBD between the

three technologies after removing the excess of rare alleles in the GBS to have the same

distribution of MAF as in the 50K and the 600K arrays. The red line is the bisector.

Figure S5: Principal coordinate analyses (PCoA) of the DROPS panel. The PCoA were based

on the covariance matrix K_Freq estimated from the 50K Illumina array. The genetic groups

identified by ADMIXTURE ($N_Q = 4$) are colored (differently than in Fig. S6). Three key

founders are indicated (Iodent: PH207 in red, Stiff Stalk: B73 in blueviolet, Lancaster: Mo17

in turquoise).

1083

Additional file 4 (.docx):

Figure S6: Heatmap of genome-wide linkage disequilibrium (LD) between all markers within

and between chromosomes using PANZEA SNPs from the 50K array. All SNPs were ordered

according to their position on the genome. Dots represented LD between two loci and were

colored according to their strength. Classical LD measurement r^2 between loci were

represented within triangle below the diagonal. Linkage disequilibrium corrected for structure

(r^2S, A), relatedness (r^2K, B) or both (r^2KS, C) were represented within triangle above the

diagonal.

Figure S7: Linkage disequilibrium (r^2, top) and LD corrected for relatedness (r^2k, bottom) as

a function of physical distance (left) and genetic distance (right): example of chromosome 1.

Figure S8: Variation of genetic LD extent (Dm, cM), effective population size (N), along the

physical map. A sliding window of 1 cM moving by 0.5 cM at each step was used. Local

genetic LD extent (cM) and local effective size (N) were estimated by adjusting the Hill and

Weir model's using r^2K between all loci that are located in sliding windows of 1 cM. Each

values were plotted on the physical map of each chromosome by projecting the genetic

position of the windows on the physical map.

1100

Additional file 5 (.docx):

Supplementary Text 1: Differences between microarrays.

Supplementary Text 2: GBS pipelines.

Supplementary Text 3: Statistical models for GWAS.

Supplementary Text 4: Effects of confounding factors on GWAS.

Supplementary Text 5: Performance of different software.

1107

Additional file 6 (.pdf):

Figure S9-LD_Windows: QTL limits obtained by the *LD_win* approach projected on

heatmaps representing the level of LD between associated SNPs for each trait (DTA: male

flowering time, plantHT: plant height and GY: grain yield) and each chromosome. Upper and

lower triangles on the heatmaps represented the r^2 and r^2K values between associated SNPs,

respectively. Linkage disequilibrium between loci was colored according to values from weak

LD (yellow) to high LD (red). The significant markers were ordered according to their

physical positions on the chromosome and were represented by ticks on the four sides of the

heatmaps. Limits of QTLs were displayed by gray dotted lines. QTL numbers were indicated

in gray on the top and the right of each heatmap.

1118

Additional file 7 (.pdf):

Figure S9-LD_Adjacent: QTL limits obtained by the *LD_Adj* approach projected on

heatmaps representing the level of LD between associated SNPs for each trait (DTA: male

flowering time, plantHT: plant height and GY: grain yield) for each chromosome. Upper and

lower triangles on the heatmaps represented the r^2 and r^2K values between associated SNPs,

respectively. Linkage disequilibrium between loci was colored according to values from weak

LD (yellow) to high LD (red). The significant markers were ordered according to their

physical positions on the chromosome and were represented by ticks on the four sides of the

heatmaps. Limits of QTLs were displayed by gray dotted lines. QTL numbers were indicated

in gray on the top and the right of each heatmap.

1129

Additional file 8 (.pdf):

Table S3: Summary of all the QTLs identified for the male flowering time (DTA), plant

height (plantHT) and grain yield (GY). "LowerLimit" and "UpperLimit" columns are the

lower and upper physical limits for each QTL. The "Rec" column indicates if the QTL is

located in a high or low region of recombination. "NbSNP50", "LogPvaMax50",

"NbSNP600", "LogPvaMax600", "NbSNPGBS", "LogPvaMaxGBS" are the number of

significant SNPs and the most significant $-log_{10}(Pval)$ within the QTL for each technology

across all environments. The physical position ("PosMax"), the proportion of the variance

explained ("R2_LDMax") and the effect ("EffectMax") of the most significant SNP within

the QTL is shown. "NbDiffEnv" gives the number of different situations that detected the

QTL.

1141

Additional file 9 (.docx):

Figure S10: Examples of comparison of QTLs detection on Chromosome 1, 6 and 8 for the

different traits. Local distribution of the $-log_{10}(p$-$value)$ and linkage disequilibrium (bottom)

corrected by the kinship (r^2k) of all SNPs with the strongest associated marker within the

chosen QTL for the three technologies. Ticks on different x-axes show the marker density of

the three technologies (red for the 50K, blue for the 600K and green for the GBS). The

vertical red line spots the position of the SNP with the maximum $-log_{10}(p$-$value)$ within the

QTL.

1150

Additional file 10 (.docx):

Figure S11: Pleiotropy of QTLs between the traits. Number of QTLs specific and shared by

the three traits across all environments. Note that several QTLs from one trait were sometimes

included in a single QTL of another trait.

Figure S12: Percentage of stable QTLs across environments for the three traits (DTA: male

flowering time, plantHT: Plant Height, GY: Grain Yield).

Table S4: Stability of QTL across environments. DTA: male flowering time, plantHT: plant

height, GY: grain yield traits.

Table S5: Recombination rate and proportion of low and high recombination regions.

Average recombination rate ("RecRate") and proportion of the physical ("Phys") and genetic

("Genetic) map in low ("LowRec", <0.5 cM / Mbp) and high ("HighRec", >0.5 cM / Mbp)

recombination regions for each chromosomes. "Chr" indicates the chromosome. Physical and

genetic size columns indicated the size of each chromosome in bp and cM, respectively.

Figure S13: Percentage of QTLs located in high (darkgrey) and low (lightgrey)

recombinogenic regions. (a) male flowering time, (b) plant height and (c) grain yield.

1166

1167 **Additional file 11 (.pdf):**

1168 **Table S6:** Description of inbred lines. Variety and accession along with the breeders, seeds

1169 providers and genetic groups obtained using ADMIXTURE for K=4 (Stiff Stalk, Iodent,

1170 Lancaster, Other).

1171

Additional file 12 (.docx):

Table S7: Narrow sense heritability (h²) and variance components (V_g, genetic variance; V_e,

residual variance). The heritability and variance components were estimated for all traits

(grain yield, male flowering time and plant height) using the R package Heritability [1].

Figure S14: Linkage disequilibrium based approach to delineate a physical window around

each SNP, examplified with chromosome 3. Linkage disequilibrium (LD) windows were

defined per chromosome for each SNP based on physical LD extent in low recombinogenic

regions (left part) and based on genetic LD extent in high recombinogenic regions (right part).

These LD windows were used (i) to group significant SNPs into QTLs when they overlapped,

(ii) to estimate genome coverage to detect QTLs by GWAS considering region not covered by

LD windows, (iii) identify putative underlying genes involved in trait variations.

Figure S15: QQ-plots representing observed $-log_{10}(p$-$value)$ against expected $-log_{10}(p$-$value)$

under null hypothesis (No association, black line). We tested association between 44,729

SNPs from the 50K array and the male flowering time trait in one environment (Gai12R)

using different GWAS models, kinship estimators and programs. (A) Comparisons between

statistical models: M1 is the model without correction (green dots), M2 takes into account the

group structure (blue dots), M3 takes into account kinship (IBD: K_freq) between individuals

(purple dots) and M4 takes into account both group structure and kinship (red dots). (B)

Comparison between mixed models using different estimates (IBS and IBD, K_freq) of

kinship. (C) Comparison of using or not Rincent *et al.* 2014 approach (using K_Chr vs

K_freq). (D) Comparison between different informatics tools (EMMAX, ASReml, FasST-

LMM) that perform GWAS.

Figure S16: Correlations between the GWAS results from the GBS genetic data using a

kinship estimated from the PANZEA 50K array (x-axis) and a kinship estimated from the

GBS (y-axis). The horizontal and vertical lines are the threshold $-log_{10}(p$-$value) = 5$. The

correlations were done using the flowering time (DTA) and plant height (plantHT) traits and

the two sites, two years and two treatments (Gai12R, Gai12W, Gai13R, Gai13W, Ner12R,

Ner12W, Ner13R, Ner13W).

Chromosome 3