The 'zoo' accompanying *Botryococcus braunii*:

Unraveling the fundamentals of algae-bacteria biocoenosis

Running title: Fundamentals of algae-bacteria biocoenosis

Olga Blifernez-Klassen[1,2], Viktor Klassen[1,2], Daniel Wibberg[2], Swapnil Chaudhari[1,2],

Christian Henke[2,3], Anika Winkler[2], Arwa Al-Dilaimi[2], Oliver Rupp[4], Jochen Blom[4],

Alexander Goesmann[4], Alexander Sczyrba[2,3], Andrea Bräutigam[2,5], Jörn Kalinowski[2],

Olaf Kruse[1,2*]

[1]Algae Biotechnology and Bioenergy, Faculty of Biology, Bielefeld University,
Universitätsstrasse 27, 33615 Bielefeld, Germany
[2]Center for Biotechnology (CeBiTec), Bielefeld University, Universitätsstrasse 27, 33615
Bielefeld, Germany
[3]Computational Metagenomics, Faculty of Technology, Bielefeld University,
Universitätsstrasse 25, 33615 Bielefeld, Germany
[4]Bioinformatics and Systems Biology, Justus-Liebig-University, Heinrich-Buff-Ring 58,
35392 Gießen, Germany
[5]Computational Biology, Faculty of Biology, Bielefeld University, Universitätsstrasse 27,
33615 Bielefeld, Germany

*Address correspondence to: Olaf Kruse, Algae Biotechnology and Bioenergy, Faculty of
Biology, Center for Biotechnology (CeBiTec), Bielefeld University, Universitätsstrasse 27,
33615 Bielefeld, Germany; phone: +49 521 106-12258; email: olaf.kruse@uni-bielefeld.de

Competing interests
The authors declare that they have no competing interests.
This work was supported by the European Union Seventh Framework Programme (FP7/2007-
2013) under grant agreement 311956 and by the BMBF-funded project "Bielefeld-Gießen
Center for Microbial Bioinformatics" – BiGi (grant 031A533).

Abstract

The interactions between microalgae and bacteria represent a fundamental ecological

relationship within aquatic environments, by controlling nutrient cycling and biomass

production at the food web base. However, the extent of these interactions and their metabolic

significance are poorly understood. To elucidate the complexity of microalga-bacteria

interactions, *Botryococcus braunii* consortia were chosen as a suitable model system with an

organic carbon-rich phycosphere. Here we present a comparative metagenome analysis in

conjunction with physiological data that allow a deep insight into the community relationship.

The reconstruction of ten high-quality draft and two full genomes of the abundant associates

clarify consortia's basic functional characteristics based on the bacterial genetic repertoire.

Implying a strong mutualistic algae-bacteria relationship, most abundant members of the

Botryococcus phycosphere belong to a distinct niche of B-vitamin auxotrophs, essentially

depending on the microalga to provide these important cofactors. This high dependency

allows the microalga to effectively control its own phycosphere and impacts the bacteria-

bacteria interactions within the consortium. Overall, our results provide highly valuable

insights into the fundamentals of microbial ecology where metabolic networks cross species

boundaries and define complex interactions. The knowledge of these interactions could enable

the design of functional synthetic ecosystems consisting of photosynthetic and heterotrophic

microorganisms.

Introduction

Microalgae are the dominant primary producers and the base of the food web within

aquatic ecosystems and sustain in the natural environment in close relationship with multiple

associated microorganisms [1, 2]. Algae-bacteria interactions, still poorly understood, are

multifaceted and complex, and often take place in the microenvironment of individual algal

cells, the phycosphere [2, 3]. The spectrum of ecological interrelations within the specific

phycosphere ranges from cooperative to competitive [1], where algae and bacteria 'live

together' within a framework of symbiosis [4]. The nature of the exchange of micro- and

macronutrients, diverse metabolites including complex algal products such as

polysaccharides, and infochemicals defines the relationship of the symbiotic partners [5, 6],

which span mutualism, commensalism and parasitism [2]. Within a parasitic association, the

bacteria are known to adversely affect the algae (*e.g.* cell wall degradation [7]), while in

commensalism the partners can survive independently, with the commensal feeding on algae-

provided substrates, but without harming the host [2]. Mutualistic interactions include *i.a.* the

obligate relationships between vitamin-synthesizing bacteria and vitamin-auxotrophic

phytoplankton species [8]. Many microalgae cannot synthesize several growth-essential

vitamins (like B_{12} and B_1) [8–10] and obtain them through a symbiotic relationship with

bacteria in exchange for organic carbon [11]. Not only the microalgae require these vitamins,

but also many bacteria have to compete with other organisms for the B-vitamins [12].

However, in microbial environments, where the distinction between host and symbionts is

less clear, the identification of the associated partners and the nature of their relationship is

challenging. Looking for a suitable model system to elucidate the complex interactions

between algae and bacteria, we focussed on the lifestyle of the colony-forming and

hydrocarbon producing microalga *Botryococcus braunii*. This green microalga is known to

co-exist with heterotrophic microorganisms [13]. The alga-bacteria consortium may represent

an example for an association already existing during the early stages of life development on

earth, since it is generally considered that a progenitor of *Botryococcus* impacted the

development of today's oil reserves and coal shale deposits [14]. Nowadays, this cosmopolitan

microalga is widely considered of enormous biotechnological importance because of its

unique ability to produce and secrete large quantities of aliphatic (C_{20}-C_{40}) hydrocarbons and

exo-polysaccharides [15]. Due to the excretion of large quantities of organic carbon into its

phycosphere, naturally occurring *B. braunii* cultures are generally not axenic but are

associated with a variety of microorganisms [13]. The interactions between *B. braunii* and the

associated bacteria are not fully understood, but are known to influence the microalgal growth

performance and product formation capacity [16]. The associated bacteria may stimulate algal

growth and product formation [17] or, conversely restrict algal growth by competing for

nutrients, degrading algal exopolysaccharides and hydrocarbons, releasing anti-algal

substances, or affecting algal morphology [13, 15].

To understand the complex nature of the unique habitat of a microalgal phycosphere and to

unravel the food web between alga and bacteria, we investigated the consortium

accompanying the hydrocarbon/carbohydrate-producing green microalga *Botryococcus*

braunii. Two *B. braunii* races (A and B) were sequenced at different time points (linear and

stationary growth) for the analysis and identification of the bacterial key players within

communities and their ecological impact on the microalga.

Materials and Methods

Strains, cultivation conditions and growth determination parameters. Liquid, xenic

cultures of *Botryococcus braunii* race A (CCALA778, isolated in Serra da Estrela (Baragem

da Erva da Fome) Portugal) and race B (AC761, isolated from Paquemar, Martinique, France)

as well as the axenic *B. braunii* race A strain (SAG30.81, Peru, Dpto.Cuzco, Laguna Huaypo)

used for co-cultivation studies. The cultures were phototrophically cultivated under 16:8 light-

dark illumination of 350 ± 400 μmol photons m^{-2} s^{-1} white light in modified Chu-13 media

(without the addition of vitamin and organic carbon source) as described before [18]. Carbon

supply and agitation were achieved by bubbling the cultures with moisture pre-saturated

carbon dioxide-enriched air (5%, v/v) with a gas flow rate of 0.05 vvm. The axenic strain was

regularly checked for contaminations.

The bacterial isolates were obtained by plating highly diluted xenic *B. braunii* cultures

(CCALA778 and AC761) on LB agar plates. After two weeks single cell colonies were

picked, separately plated and analyzed *via* 16S rDNA Sanger-sequencing by using the primer

pair 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1385R (5'-

CGGTGTGTRCAAGGCCC-3'). Two isolates, deriving from algal races A and B, were

classified based on full length 16S rDNA similarity as *Mycobacterium* sp. and *Pimelobacter*

sp., respectively and were used for further analysis.

Cultures growth was monitored by measurements of biomass organic dry weight (DW) by

drying 10 mL culture per sample (in three biological and technical replicates) following the

protocol according to Astals et al [19]. For the co-cultivation experiments, 500 mL axenic

cultures (~0.25 gL^{-1}) were inoculated with the isolates (*Mycobacterium* sp. BbA or

Pimelobacter sp. Bb-B (~$2x10^6$ cell/mL)) and cultured under photoautotrophic conditions for

21 days. The bacterial cell density was determined by manual cell counting using

hemocytometer. Additionally, microalgae and bacteria cells were regularly checked by optical

microscopy.

Hydrocarbon extraction and analysis. Hydrocarbon extraction was performed according to

the protocol published previously [18]. In brief, 3.0 mL of culture was mixed with equal

volume of acetone and *n*-hexane and vortexed for 2-3 min. After a heating (5 min at 85 °C)

and centrifugation (5 min at 3020 $\times g$) step the upper *n*-hexane phase was collected and dried

in a N_2-atmosphere. The dry extracted hydrocarbons were resuspended in 500 µL of *n*-

hexane, containing an internal standard *n*-hexatriacontane ($C_{36}H_{74}$) and analysed *via* GC-MS

and GC-FID. GC-MS analysis was performed with a TraceGC gas chromatograph and a ITQ

ion trap mass spectrometer equipped with an AS2000 auto sampler (ThermoFinnigan,

Dreieich, Germany) using a 30-m×0.25-mm VF-5ms column with a 0.25 μm 5% diphenyl and

95% dimethylsiloxane coating (Varian Deutschland GmbH, Darmstadt, Germany). The

method consisted of a temperature gradual ramp from 80 to 300 °C at 6 °C min^{-1}; at the final

temperature of 300 °C, 1 μL sample was splitlessly injected. Mass spectra were recorded at 20

scans s^{-1} with a scanning range of 50–750 m/z. The evaluation of the chromatograms was

accomplished with the Xcalibur software (ThermoFinnigan). Additionally, all samples were

analyzed in a Shimadzu GC-2010 Plus (Shimadzu, Kyoto, Japan) unit with a FID detector as

described previously [20]. Briefly, the separation was performed with a Macherey-Nagel

OPTIMA FFAPplus column (30 m×0.25 mm ID, 0.40 mm OD and 0.50 μm film) and a 10 m

pre-column (carbowax-deactivated, 0.25 mm ID, 0.40 mm OD, Macherey-Nagel). The sample

injection volume was 1 μL working in split mode (1:5) and injected with an auto-injector

(Shimadzu AOC-20i) at a constant pressure-mode of with 160 kPa with helium as carrier gas

and nitrogen as makeup gas (Linde, Munich, Germany). The method consisted of a

temperature ramp from 55 to 232 °C at 6 °C min^{-1} and to 250 °C at 3 °C min^{-1}, with keeping

the final temperature of 250 °C constant for 29.5 min. For identification and quantification of

all hydrocarbons external standards (alkane standard solution C_{21}-C_{40}, Sigma-Aldrich/Fluka)

were used.

DNA sample collection and preparation. The samples from xenic *B. braunii* race A and B

cultures (three biological replicates) were taken in the linear and stationary growth phases

(after 9 and 24 days). Total community genomic DNA was extracted as previously described

by Zhou et al [21]. The quality of the DNA was assessed by gelelectrophoresis and the

quantity was estimated using the Quant-iT PicoGreen dsDNA Assay Kit (Invitrogen) and a

Tecan Infinite 200 Microplate Reader (Tecan Deutschland GmbH).

High-throughput 16S rDNA amplicon sequencing. To get insight into the community

composition of *B. braunii* Race A and B, high-throughput 16S rDNA amplicon sequencing

was performed as described recently by Maus et al [22]. The primer pair Pro341F (5'-

CCTACGGGGNBGCASCAG-3') and Pro805R (5'-GACTACNVGGGTATCTAATCC-3')

was used to amplify the hypervariable regions V3 and V4 of diverse bacterial and archaeal

16S rRNAs [23]. In addition, the primers also cover the 16S rDNA gene of algal chloroplasts

and eukaryotic mitochondrial genomes. Furthermore, in a two PCR steps based approach,

multiplex identifier (MID) tags and Illumina-specific sequencing adaptors were added to each

amplicon. Only amplicons featuring a size of ~460 bp were purified using AMPureXP®

magnetic beads (Beckman Coulter GmbH, Brea, California, USA). Resulting amplicons were

qualitatively and quantitatively analyzed using the Agilent 2100 Bioanalyzer system (Agilent

Inc., Santa Clara, California, USA) and pooled in equimolar amounts for paired-end

sequencing on the Illumina MiSeq system (Illumina, San Diego, California USA). This

sequencing approach provided ~150,000 reads per sample. An in-house pipeline as described

previously [24] was used for adapter and primer trimming of all samples. For amplicon

processing, a further in house amplicon processing pipeline including FLASH [25],

USEARCH [26] (v8.1), UPARSE [27] and the RDP [28] classifier (v2.9) was applied as

described recently [29, 30]. In summary, unmerged sequences resulting by FLASH [25]

(default settings +-M 300) were directly filtered out. Furthermore, sequences with >1Ns

(ambiguous bases) and expected errors >0.5 were also discarded. Resulting data was further

processed and operational taxonomic units (OTUs) were clustered by applying USEARCH

[26] (v8.1). The resulting OTUs were taxonomically classified using the RDP [28] classifier

(v2.9) in 16S modus (Threshold >0.8) and compared to the nt database by means of BLASTN

[31]. In the last step, raw sequence reads were mapped back onto the OTU sequences in order

to get quantitative assignments (Supplementary data 1).

(Meta)Genome sequencing, assembly, binning and annotation. For each of the four

samples (*B. braunii* race A and B from linear (T1) and stationary (T2) growth phase), the

genomic DNA from three replicates was pooled. To obtain the metagenome sequence, four

whole-genome-shotgun PCRfree sequencing libraries (Nextera DNA Sample Prep Kit;

Illumina, Munich, Germany) were generated based on the manufacturer's protocol

representing different time points of the cultivation and different algae communities. The four

libraries were sequenced in paired-end mode in a MiSeq run (2 x 300 bp) resulting in

11,963,558 and 11,567,552 reads for race A T1 and T2; 6,939,996 and 12,031,204 reads for

race B T1 and T2, respectively (total 12.75 Gb). The *de novo* metagenome assembly was

performed using the Ray Meta assembler [32] (v2.3.2) using a k-mer size of 41 and default

settings. For contigs with length larger than 1 kb (n=127,914), the total assembly size was

338.5 Mb, the N50=2,884 and the largest contig 1.59 Mb.

All processed raw sequence reads of the four samples were aligned to the assembled

metagenome contigs by means of Bowtie 2 [33] (v2.2.4). By using SAMtools [34] (v1.6), the

SAM files were converted to BAM files. These files were sorted and read mapping statistics

were calculated. The following portions of reads could be assembled to the draft

metagenomes: 70.1% and 77.5% for race A (T1 and T2); 51.0% and 62.9% for race B (T1 and

T2). For binning of the metagenome assembled genomes (MAGs), the tool MetaBAT [35]

(v0.21.3) was used with default settings. Completeness and contamination of the resulting

MAGs were tested with BUSCO [36, 37] (v3.0.), using the Bacteria and Metazoa data set

(Table S1).

For sequencing of the isolated species, 4 µg of purified chromosomal DNA was used to

construct PCR-free sequencing library and sequenced applying the paired protocol on an

Illumina MiSeq system, yielding ~1.71 Mio Reads (517 Mbp in 3 Scaffolds, 58 Contigs) for

Mycobacterium sp. Bb-A and ~1.70 Mio Reads (503 Mbp in 6 Scaffolds, 65 Contigs) for

Pimelobacter sp. Bb-B. Obtained sequences were *de novo* assembled using the GS *de novo*

Assembler software (v2.8, Roche). An *in silico* gap closure approach was performed as

described previously [38]. Annotation of the draft chromosomes was performed within the

PROKKA [39] (v1.11) pipeline and visualized using the GenDB 2.0 system [40].

Gene prediction, taxonomic assignment and functional characterization. Gene prediction

on the assembled contigs of the MAGs was performed with Prodigal [41] (v2.6.1) using the

metagenome mode ("-p meta"). All genes were annotated and analyzed with the in-house

EMGB annotation system using the databases NCBI NR [31], Pfam [42] and KEGG [43].

Based on DIAMOND [44] (v0.8.36) hits against the NCBI-NR [31] database, all genes were

subject to taxonomic assignment using MEGAN [45]. Relationships to metabolic pathways

were assigned based on DIAMOND [44] hits against KEGG [43]. For Pfam [42] annotations,

pfamscan was used with default parameters. The genomes were manually inspected using a E-

value cutoff of $1x10^{-10}$ for several specific sequences listed in Table 1 and Supplementary

data 4.

The profiling of the carbohydrate-active enzymes (CAZy) encoded by the *B. braunii*

community members was accomplished using dbCAN2 [46] metaserver (Supplementary data

5). The identification, annotation and analysis of the secondary metabolite biosynthesis gene

clusters within the MAGs and complete genomes were accomplished *via* the antiSMASH [47]

(v4.0) web server with default settings (Supplementary data 7).

Phylogenetic analyses. To assign and phylogenetically classify the MAGs, a core genome

tree was created by applying EDGAR [48] (v2.0). For calculation of a core-genome-based

phylogenetic tree, the core genes of all MAGs and the two isolates *Mycobacterium* sp. Bb-A

and *Pimelobacter* sp. Bb-B, corresponding to selected reference sequences were considered.

The implemented version of FASTtree [49] (using default settings) in EDGAR [48] (v2.0)

was used to create a phylogenetic Maximum-Likelihood tree based on 53 core genes (see

Supplementary data 2) of all genomes. Additionally, the MAGs were taxonomically classified

on species level by calculation of average nucleic and amino acid identities (ANI, AAI) by

applying EDGAR [48] (v2.0) (Figure S3). For phylogenetic analysis of the 16S rDNA

amplicon sequences, MEGA [50] (v7.0.26) was used. Phylogenetic tree was constructed by

means of Maximum-Likelihood Tree algorithm using Tamura 3-parameter model. The

robustness of the inferred trees was evaluated by bootstrap (1000 replications).

Availability of data and materials

The high-throughput 16S rDNA amplicon sequencing datasets obtained during the present

work are available at the EMBL-EBI database under BioProjectID PRJEB21978. The

metagenome datasets of *Botryococcus braunii* as well as the resulting MAGs have been

deposited under the BioProjectIDs PRJEB26344 and PRJEB26345, respectively. The

communities most abundant ten high-quality draft genomes are accessible via EMGB

platform (https://emgb.cebitec.uni-bielefeld.de/Bbraunii-bacterial-consortium/). The genomes

of the isolates *Mycobacterium* sp. BbA or *Pimelobacter* sp. Bb-B are accessible under

BioProjectIDs PRJEB28031 und PRJEB28032, respectively.

Results

Effect of the bacterial community on the *B. braunii* host

We started our investigations with the analysis of an axenic *Botryococcus braunii* strain to

assess algal capacities for growth and hydrocarbon production under strict phototrophic

conditions (*e.g.* without the supplementation of vitamins or organic carbon). Axenic *B.*

braunii showed a continuous increase in cell biomass and hydrocarbon content, yielding

2.1 ± 0.2 gL^{-1} and 0.7 ± 0.1 gL^{-1}, respectively (Figure 1a).

In a next step, we co-cultivated the axenic strain with two isolated actinobacterial

associates from *B. braunii* (*Mycobacterium* sp. Bb-A and *Pimelobacter* sp. Bb-B, see

Methods). *B. braunii* in bacteria-free conditions (axenic culture) accumulated hydrocarbons

up to 38% of dry weight (DW) during the cultivation, while the cultures containing bacteria

(xenic cultures) accumulated only very limited hydrocarbons during the bacterial exponential

growth phase (up to 5% of DW for co-cultures with bacteria, Figure 1). In the stationary

phase of bacterial growth, the alkadiene $C_{31}H_{60}$ accumulated again to a low level. Of

particular note was that the algal cultures changed their morphology during bacteria co-

cultivation by reducing colony formation when compared to axenic growth.

This dramatic decrease in hydrocarbon content and disturbed colony formation was not

observed in the xenic *Botryococcus braunii* cultures, from which both bacterial isolates are

derived (Figure S1). To test the metabolic capabilities and possible interactions in the *B.*

braunii consortia of races A and B at linear and stationary algal growth, we assessed the

bacteria associated with each culture using metagenomics.

Metagenomic survey of the *Botryococcus braunii* consortia

We profiled four different communities using high-throughput 16S rDNA gene amplicon

data and added 12.75 Gb of paired-end metagenome data. The analysis of the 16S rDNA

amplicons and a read-based metagenome approach showed that each of the *B. braunii*

communities represents a distinct large consortium, consisting of a single algal and various

bacterial species as well as traces of Archaea and viruses (Figure S2). Each community is

comprised of four bacterial phyla, consisting of at least 25 species with the members of the

phylum Proteobacteria being the most prominent in both *B. braunii* races. Of 33 different

bacterial genera, four were specific to race A, seven were specific to race B and 22 were

shared (Figure 2). The relative abundance of plastid 16S rDNA sequences accounted up to 49

and 59% of all detected amplicon reads in race A and B samples in the linear growth phase

and declined to approximately 26% and 44% at stationary phase, respectively (Figure S2a). At

the same time, the 16S rDNA contribution of the accompanying microbial community

increased. The community members varied quantitatively based on both race and time of

sampling, but not qualitatively.

Both *B. braunii* races were dominated by the genera *Devosia* and *Dyadobacter* (up to 26%

and 22% of the bacterial community) in the logarithmic growth phase, followed by

Dokdonella in race A (16%) and *Acidobacteria* in race B (22%) (Figure 2). During the

cultivation, the relative abundancy of the dominant species in the race A consortium such as

Devosia and *Dokdonella* increased at the expense of *Dyadobacter* in comparison to the linear

growth phase. In the race B consortium, less abundant taxa such as *Mesorhizobium,*

Brevundimonas and *Hydrogenophaga* increased at the expense of *Dyadobacter* and *Devosia*.

The detected relative levels of both actinobacterial isolates, which had been investigated in

co-cultivation experiments with the axenic algae (Figure 1) were comparatively low, although

both *B. braunii* races are capable to provide a wealth of organic substrate (in form of

hydrocarbons and/or carbohydrates) to ensure their proliferation (Figure S1). *Mycobacterium*

sp. Bb-A isolate, obtained from the *B. braunii* race A community, was almost exclusively

propagated therein and increased during stationary growth to 7% of all detected bacterial

amplicons (Figure 2). *Pimelobacter* sp. Bb-B was isolated from the race B consortium, and is

present in both communities in equally (≤1%) low abundances which increased slightly

during stationary phase.

To test the relationships between *B. braunii*, *Mycobacterium*, *Pimelobacter* and the

remaining consortium, we explored the genetic potential of the abundant representatives of

algal-bacterial consortia. We combined metagenome *de novo* assembly and differential

coverage and tetranucleotide signature based binning of the *B. braunii* race A and B samples.

This approach resulted in the reconstruction of ten high-quality (completeness >80%,

contamination <10%) and ten fragmentary metagenome assembled genomes (MAGs) as well

as seven eukaryotic genome fragments (for details see Methods, Table S1). To identify taxa

representing the ten high-quality draft MAGs, we constructed a phylogenetic tree built out of

a core of 53 highly conserved marker genes per genome (Figure 3). All abundant taxa shared

by both races were recovered, including the most abundant alphaproteobacterial MAG of the

order *Rhizobiales*, designated Dev-G (similar to *Devosia* sp. 66-14, Table S2 and Figure S3).

Additionally, MAGs of *Dyadobacter* and *Hydrogenophaga* were constructed (Dya-G and

Hyd-G, similar to *Dyadobacter* sp. 50-39 and *Hydrogenophaga* sp. PML113, respectively).

Another two alphaproteobacterial MAGs Bre1-G and Bre2-G of the order *Caulobacteriales*

(similar to *Brevundimonas* sp. DS20), clustered together within the 16S rDNA amplicon

approach (Figure 2) but were separated through the binning procedure (Tables S1 and S2).

We reconstructed sphingobacterial MAGs of the order *Chitinophagales*, Ter1-G and Ter2-G

similar to different extent to *Terrimonas ferruginea* DSM30193. Further two MAGs, basically

occurring in race A strain, included a gammaproteobacterial bacterium similar to *Dokdonella*

koreensis DS-123 (Dok-G) and a member of *Corynebacteriales* similar to *Mycobacterium* sp.

Root135 (Myc-G). An exclusively in *B. braunii* race B occurring member of the subdivision

IV of the phyla Acidobacteria (Gp4-G) is similar to *Pyrinomonas methylaliphatogenes* K22.

We also sequenced the two actinobacterial isolates, *Mycobacterium* sp. Bb-A and

Pimelobacter sp. Bb-B, used for the co-cultivation experiments (Figure 1). After assembly

and *in silico* finishing, both genomes were completed and represented by a single

chromosome, each (Figure S4). The *Mycobacterium* sp. Bb-A genome sequence was identical

to the Myc-G, indicating that both represent the same species (Figure S3h), thus underlining

the quality of our metagenome assembly and binning approach.

Elucidation of the genetic portfolio of the bacterial community

To identify genes that enabled the bacterial community members to contribute to and to

benefit from the food web of the *B. braunii* algae based consortium, we performed a KEGG-

based quantitative functional assignment of the annotated high-quality MAGs and sequenced

genomes by applying the elastic metagenome browser (EMGB, https://emgb.cebitec.uni-

bielefeld.de/Bbraunii-bacterial-consortium/).

Six of twelve genomes carried genes for bacterial chemotaxis and five of twelve genes for

flagella synthesis, hinting at active motility and ability to move towards substrates in the

medium (Figure S5). *Botryococcus braunii* synthesizes aliphatic long-chain hydrocarbons,

which accumulate in large quantities in the extracellular matrix of this species [15]. Seven of

twelve genomes carried gene homologs of already characterized hydrocarbon hydroxylases

(*alkB*, *ladA*, *almA* and/or cytochrome P450 (CYP153), Table 1) for the initial

oxyfunctionalization of long-chain hydrocarbons [51]. While the most abundant consortia

members Dev-G and Dya-G exhibited a low enzyme number, the more rare *Mycobacterium*

and *Pimelobacter* genomes contained a large portfolio of hydrocarbon monooxygenase/

hydroxylase genes, and were the only genomes to carry non-heme membrane-associated

monooxygenase *alkB* genes (Table 1).

Mucilaginous (exo-)polysaccharides as part of the internal fibrillar layer of the *B. braunii*

cell wall represent another prospective carbon source for the bacterial community, consisting

mainly of galactose but also of fucose and rhamnose [15, 18]. The profiling of the

carbohydrate-active enzymes (CAZymes [52]), encoded by the members of *B. braunii*

communities using dbCAN2 [46] metaserver, revealed the presence of several putative

enzyme families involved in the complex carbohydrate metabolism (Table 1).

Eight out of twelve genomes coded for several carbohydrases for the degradation of poly-

and oligosaccharides, generally present in the cell walls of microalgae, and contained genes

for chitin disintegration and assimilation (Supplementary data 5). However, six species

showed a clear prevalence in the number of enzymes involved in the biosynthesis of

carbohydrates (glycosyltransferases) compared to degradation (glycoside hydrolases). Some

species like the members of Bacteroidetes contained a large number of CAZyme-encoding

genes in their genomes (up to 5.9% of total gene content) with a clear prevalence to hydrolytic

activity of complex carbohydrates (Table 1). For instance, the Dya-G genome contains more

genes encoding for the glycoside hydrolase (GH) families involved in the degradation of

complex polymers (cellulose, xylan, chitin), while others, such as the genome of Dev-G,

harbor enzymes acting on oligosaccharides [52] (Supplementary data 5).

Eight of twelve genomes encode genes for the synthesis of cyanophycin granule

polypeptides (CGPs) [53], however only four contained genes for the degradation of this C-

and N-storage compound (Table 1). Related to this, all genomes carry genes for the synthesis

but not for the degradation of polyphosphate granules (Poly-P), a polymeric reserve with a

significant role in the regulation of enzymatic activities, gene expression and stress adaptation

processes [54]. Suggesting an adaptation to phosphorous and iron limitation, all genomes

contained genes relevant for phosphate starvation (*pst* system) or siderophore biosynthesis

(*fhu* system).

Looking at the primary metabolism, all genomes encode genes for processes essential for

aerobic respiration (Figure S5, Supplementary data 3). Genes that enable pyruvate

fermentation to lactate, acetate and ethanol are also present, showing a versatility of the

community members to switch between aerobic and anaerobic metabolisms depending on the

oxygen availability. All except *Mycobacterium* genomes contained genes for both the aerobic

cobalamin (vitamin B_{12})-independent form and the oxygen-independent but B_{12}-dependent

type of ribonucleotide reductases [55] (Table 1), demonstrating adaptations to a facultative

aerobic life style of the *B. braunii* associates.

We were unable to reconstruct the complete cobalamin (vitamin B_{12}) synthesis pathway for

eight out of twelve assembled genomes. The absence of the *metE* gene for the B_{12}-

independent methionine synthesis pathway in these genomes (except Dya-G, Bre1-G and

Hyd-G) implies the need of exogenous supply of vitamin B_{12} for survival (Table 1). Seven of

twelve genomes lack the complete gene portfolio for the biotin (vitamin B_7) synthesis, but

contained the complete biosynthesis pathway of fatty acids for which biotin is essential [12].

Four of twelve genomes were incomplete in the synthesis of thiamin (vitamin B_1), which is

essential for citrate cycle and pentose phosphate pathway [12]. In addition to vitamin B_1 and

B_7-auxotrophy, the most abundant Dev-G species was found to comprise incomplete *de novo*

biosynthesis pathways of riboflavin (vitamin B_2, essential as the FMN/FAD precursor) and

pantothenate (vitamin B_5, essential for CoA synthesis) [56] (Table 1).

Additionally, eleven out of twelve genomes encoded for bacteriocins, antimicrobial

peptides with significant potency against other bacteria, which can exhibit narrow spectra of

activity (targeting members of the same species) or display broader activity spectra (targeting

other species and genera) [57]. Especially, Hyd-G, *Mycobacterium* sp. Bb-A and

Pimelobacter sp. Bb-B exhibited a large portfolio of metabolic gene cluster highly similar to

antibiotic compounds with a broad spectrum of activity against Gram-positive bacteria which

are exceedingly rare in the communities of *B. braunii*.

Collectively, within our survey of the microbial community accompanying the

Botryococcus braunii races A and B, we made complex observations, which hint at

interconnected food webs and are summarized in Figure 4. The genetic portfolio of

Mycobacterium sp. Bb-A and *Pimelobacter* sp. Bb-B reflect the strong metabolization of

hydrocarbons (Figure 1, Table 1). However, despite the apparent vitamin-autotrophy and the

pronounced proliferation in co-cultivation experiments with the microalga, both

actinobacterial strains were poorly represented in the consortia of both *B. braunii* races. On

the other hand, Gram-negative bacterial species such as *Devosia* (Dev-G) and *Dyadobacter*

(Dya-G) that displayed several vitamin auxotrophies (especially B_7) and would not be viable

on their own, dominated the algal communities. These results clearly indicate that both

Actinobacteria are actively restrained by the abundant consortia members, possibly realized

by the synthesis and secretion of secondary metabolites (*e.g.* bacteriocines).

Discussion

Algae exist in the natural environment as part of a complex microbial consortium, where

the different species may exert considerable influence on each other for the exploitation of

unique biological functions and/or for the metabolite exchange [1].

B. braunii bacterial communities appear to be highly conserved and illustrate relatively

low species biodiversity, with most of the detected genera occurring in both communities,

albeit differentially abundant (Figure 2). The re-processing of the *B. braunii* Guadeloupe

strain metagenome data [58] also shows data qualitatively but not quantitatively similar to the

two *Botryococcus* consortia analyzed in this study. The direct comparison of the different

metagenome datasets revealed the coinciding presence of the individual genera (especially

members of Bacteroidetes, Alpha- and Betaproteobacteria), although massively varying in the

abundance pattern (Figure S6). The communities did differ regarding the presence of the

Gram-positive bacterial representatives: while the consortia obtained from our studies

harbored the members of *Actinobacteria*, the members of the phyla *Firmicutes* were strongly

represented in the Guadeloupe strain [58]. This evaluation of three *B. braunii* consortia under

changing conditions has emphasized that depending on the physiological condition (*e.g.*

supplementation of vitamins or antibiotics [58]), the relative abundance of certain species may

widely vary, suggesting the existence of many bacterial taxa with functional redundancy [59].

Many genera identified within the *Botryococcus* communities (such as *Devosia*,

Dyadobacter, *Mesorhizobium* etc.) also occur in other algae-bacterial consortia like biofilms

[60], perhaps indicating preferential coexistence between certain bacteria species and

microalgae. The interdependence of certain bacterial phyla such as Alphaproteobacteria and

Bacteroidetes observed in this study, which had been frequently found to be associated with

phytoplankton, underlines the proposed existence of the 'archetypal microalga/phytoplankton-

associated bacterial taxa' [61, 62].

At first glance, the *Botryococcus* alga does not appear obligatory dependent on a bacterial

community, since the axenic strains can be fully photoautotrophically cultured without

vitamin supplements (Figure 1). However, *Botryococcus* readily releases huge amounts of

organic carbon (up to 60% of DW in form of hydrocarbons and carbohydrates) into the

extracellular milieu (Figure S1), which creates a phycosphere that naturally attracts many

microorganisms, including those with potential damaging effects. Evolutionary fitness will

depend on efficient management of the "bacteria zoo" by *B. braunii*.

From the perspective of a microalga, the associated bacteria can be providers of micro- and

macronutrients but also competitors for limiting nutrients [6]. Based on their observed generic

potential, individual species appear to be involved in the acquisition and remineralization of

macronutrients and in the storage of nitrogen and phosphorous (in form of cyanophycin and

Poly-P, Table 1), which benefit the microalga.

We post that the microalga supplies vitamins B_1, B_2, B_5 and especially B_7 to its resident

community [12]. The abundant community members of the *B. braunii* consortia are

auxotrophs for various B-vitamins making them dependent on an exogenous supply of

vitamins for survival (Table 1). The consortium's most abundant bacterial species *Devosia*

(Dev-G, Figure 2) is indeed auxotroph for vitamins B_1, B_2, B_5 and B_7, but can synthesize

vitamin B_{12}. In view of the fact that the bacterial abundance determines their functional

relevance for the community [63, 64], this species is also likely to provide both the bacterial

community and the microalga with this vitamin [8], thus confirming the recent proposition

that the B_{12} supply is realized only by a few certain bacteria within the alga-bacteria consortia

[60]. In this context, since the B_7-auxotrophs are prevalent within the detected consortia

(except for Dok-G, appearing only in race A), it is doubtful that the large required supply of

vitamin B_7 is of bacterial origin. Because of its predominance within the community (Figure

S2), the microalga is more likely to secrete and provide this vitamin to the bacteria, at least

the secretion of the B-vitamins (*e.g.* B_5 and B_7) has already been observed in species such as

Chlamydomonas, *Chlorella* and *Coccomyxa* [65, 66]. The high requirement [64] of Dev-G for

other B-vitamins will most likely also be supplemented by the microalga. In fact, the bacterial

dependence on algal vitamins has been suggested earlier in connection with marine

phytoplankton communities [12]. However, it can not be ruled out that the missing cofactors

may be complemented by other bacteria in exchange for *e.g.* B_{12}, suggesting also strong

bacteria-bacteria relations in addition to the algae-bacteria interactions. Considering this view,

specific bacteria that are requiring particular B-vitamins are deliberately promoted by

Botryococcus braunii in exchange for *e.g.* vitamin B_{12}, micro- and macronutrients as well as

essential metabolites. The presence of particular antibacterial and probiotic compounds (Table

1) explains, why *B. braunii* communities consist mainly of Gram-negative bacteria and only a

few low abundant Gram-positive species (*e.g. Mycobacterium* and *Pimelobacter*, Figure 2).

Thus, the *B. braunii* associative consortia appears restrained by the microalgae, yet displaying

strong bacteria-bacteria interactions and capable to synthesize and provide a great deal of

substances to "fight" the potential competitors.

Based on the genetically encoded features, the bacterial associates of *B. braunii* can be

apportioned into two roughly framed categories of *i*) B-vitamin-auxotrophic (controllable),

and *ii*) B-vitamin-independent (uncontrollable) species (Figure 4). The first category includes

the members of the class of Alphaproteobacteria (like Dev-G), which display several B-

vitamin-auxotrophies, but are delivering B_{12} to the algae (and bacteria). Bacterial associates,

which are incapable to synthesize essential cofactors (including B_{12}), rely heavily on both, the

microalga and the associated bacteria. Although non-viable on their own, these phylotypes

belonging to the Proteobacteria, Bacteroidetes and Acidobacteria are dominating the

consortiae, indicating a very strong mutualistic relationship with the microalga. Completely

autonomous species like our isolates *Mycobacterium* sp. Bb-A and *Pimelobacter sp.* Bb-B are

assigned to the second category. These phylotypes are known to be hydrocarbonoclastic

parasitic microorganisms targeting mainly the organic carbon source and therefore pose a

potential risk to the microalga since they are uncontrollable, strongly implying a parasitic

relationship. And although both isolates can in principle also supply the microalgae with

vitamin B_{12}, nutrients and lots of antibacterial metabolites, they are harmful to the alga due to

their destructive effect through uncontrolled proliferation and substrate consumption, as

demonstrated in the co-cultivation experiments (Figure 1). Their activity causes a dramatic

change/damage to the algal colony morphology, resulting in a less complex colony structure,

thus having a negative impact on survival in natural environments, *i.e.* a higher susceptibility

to grazers [67].

Conclusions

While *B. braunii* can be grown axenically, naturally occurring populations exist together

with their bacterial consortia. The interaction between the microalgae and bacteria within the

Botrycoccus braunii consortiae display a complex interplay between all members, however

based on our observtions the microalgae seems to be the "conductor" in this play, attracting

specifically vitamin-auxotrophic bacterial species and keeping them in close proximity by

controlling through the subsequent vitamin supply. Metagenome analysis indicated a "zoo" of

largely supportive bacteria which supply vitamin B_{12}, and assimilated N and P in exchange for

other vitamins and carbon from the alga. In addition to this benign members, autark bacteria

without vitamin auxotrophies are present in small proportions, likely decimated by

antibacterial agents from the supportive community, and attack the hydrocarbon/

polysaccharide coat of the alga. War and peace play out amongst this microscopic community

that keeps the alga well enough to sustain collective growth.

Based on the nature of the presented observations, we unraveled only the obvious

interrelations, which are encoded in the bacterial genomes of the conserved consortiae of *B.*

braunii. The interactions between the bacteria and microalgae (phytoplankton in general) are

far more complex and multifarious, and may involve many physiological exchanges and

communications, which are not solely educible from the genetic predisposition and should be

subject for future research. But, with the present work we provide a rough framework for the

basic interdependence between a microalga (phytoplankton) and its associated bacteria, which

greatly expands the current knowledge in this area.

Acknowledgements

We are thankful to Joao Gouveia and Douwe van der Veen (Wageningen University, The

Netherlands) for providing *Botryococcus braunii* race A and B (CCALA778 and AC761)

stock cultures. This work was supported by the European Union Seventh Framework

Programme (FP7/2007-2013) under grant agreement 311956 (relating to project "SPLASH –

Sustainable PoLymers from Algae Sugars and Hydrocarbons"). The bioinformatics support by

the BMBF-funded project "Bielefeld-Gießen Center for Microbial Bioinformatics" – BiGi

(grant 031A533) within the German Network for Bioinformatics Infrastructure (de. NBI) is

gratefully acknowledged.

Competing interests

The authors declare that they have no competing interests

Authors' contributions

O.B.K., V.K, and O.K. conceived this study. O.B.K., A.S., J.K. and O.K. supervised

experiments and analyses. O.B.K., V.K. and S.C. accomplished all physiological studies.

O.B.K., D.W., A.S., C.H., A.B., V.K., A.A.D., A.W., J.K., O.R., J.B. and A.G. performed the

sequencing and the bioinformatic analyses. O.B.K., V.K. and O.K. wrote the original draft

manuscript with contributions from all authors. O.B.K., V.K., A.B and O.K. reviewed and

edited the manuscript. All authors read and approved the final manuscript.

Supplementary information

Supplementary information is available.

References

1. Seymour JR, Amin SA, Raina JB, Stocker R. Zooming in on the phycosphere: The
ecological interface for phytoplankton-bacteria relationships. *Nat Microbiol* . 2017. , **2**

2. Ramanan R, Kim B-H, Cho D-H, Oh H-M, Kim H-S. Algae–bacteria interactions:
Evolution, ecology and emerging applications. *Biotechnol Adv* 2016; **34**: 14–29.

3. Bell W, Mitchell R. Chemotactic and Growth Responses of Marine Bacteria to Algal
Extracellular Products. *Biol Bull* 1972; **143**: 265–277.

4. Cooper MB, Smith AG. Exploring mutualistic interactions between microalgae and
bacteria in the omics age. *Curr Opin Plant Biol* . 2015. , **26**: 147–153

5. Amin SA, Hmelo LR, Van Tol HM, Durham BP, Carlson LT, Heal KR, et al.
Interaction and signalling between a cosmopolitan phytoplankton and associated
bacteria. *Nature* 2015; **522**: 98–101.

6. Cole JJ. Interactions Between Bacteria and Algae in Aquatic Ecosystems. *Annu Rev*
Ecol Syst 1982; **13**: 291–314.

7. Wang X, Li Z, Su J, Tian Y, Ning X, Hong H, et al. Lysis of a red-tide causing alga,
Alexandrium tamarense, caused by bacteria from its phycosphere. *Biol Control* 2010;
52: 123–130.

8. Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. Algae acquire
vitamin B12 through a symbiotic relationship with bacteria. *Nature* 2005; **438**: 90–93.

9. Croft MT, Warren MJ, Smith AG. Algae need their vitamins. *Eukaryot Cell* . 2006. , **5**:
1175–1183

10. Tang YZ, Koch F, Gobler CJ. Most harmful algal bloom species are vitamin B1 and
B12 auxotrophs. *Proc Natl Acad Sci* 2010; **107**: 20756–20761.

11. Grant MAA, Kazamia E, Cicuta P, Smith AG. Direct exchange of vitamin B12 is
demonstrated by modelling the growth dynamics of algal–bacterial cocultures.
ISME J 2014; **8**: 1418–1427.

12. Sañudo-Wilhelmy SA, Gómez-Consarnau L, Suffridge C, Webb EA. The Role of B
Vitamins in Marine Biogeochemistry. *Ann Rev Mar Sci* 2014; **6**: 339–367.

13. Chirac C, Casadevall E, Largeau C, Metzger P. Bacterial Influence Upon Growth and
Hydrocarbon Production of the Green Alga Botryococcus Braunii. *J Phycol* . 1985. ,
21: 380–387

14. McKirdy DM, Cox RE, Volkman JK, Howell VJ. Botryococcane in a new class of
Australian non-marine crude oils. *Nature* 1986; **320**: 57–59.

15. Banerjee A, Sharma R, Chisti Y, Banerjee UC. Botryococcus braunii: A renewable
source of hydrocarbons and other chemicals. *Crit Rev Biotechnol* 2002; **22**: 245–279.

16. Fuentes JL, Garbayo I, Cuaresma M, Montero Z, González-Del-Valle M, Vílchez C.
Impact of microalgae-bacteria interactions on the production of algal biomass and

associated compounds. *Mar Drugs* . 2016. , **14**

17.   Tanabe Y, Okazaki Y, Yoshida M, Matsuura H, Kai A, Shiratori T, et al. A novel
alphaproteobacterial ectosymbiont promotes the growth of the hydrocarbon-rich green
alga Botryococcus braunii. *Sci Rep* 2015; **5**.

18.   Blifernez-Klassen O, Chaudhari S, Klassen V, Wördenweber R, Steffens T, Cholewa
D, et al. Metabolic survey of Botryococcus braunii: Impact of the physiological state on
product formation. *PLoS One* 2018; **13**: e0198976.

19.   Astals S, Nolla-Ardèvol V, Mata-Alvarez J. Anaerobic co-digestion of pig manure and
crude glycerol at mesophilic conditions: Biogas and digestate. *Bioresour Technol* 2012;
110: 63–70.

20.   Schwarzhans JP, Cholewa D, Grimm P, Beshay U, Risse JM, Friehs K, et al.
Dependency of the fatty acid composition of Euglena gracilis on growth phase and
culture conditions. *J Appl Phycol* 2014; **27**: 1389–1399.

21.   Zhou J, Bruns MA, Tiedje JM. DNA recovery from soils of diverse composition. *Appl*
Environ Microbiol 1996; **62**: 316–322.

22.   Maus I, Kim YS, Wibberg D, Stolze Y, Off S, Antonczyk S, et al. Biphasic Study to
Characterize Agricultural Biogas Plants by High-Throughput 16S rRNA Gene
Amplicon Sequencing and Microscopic Analysis. *J Microbiol Biotechnol* 2017; **27**.

23.   Takahashi S, Tomita J, Nishioka K, Hisada T, Nishijima M. Development of a
prokaryotic universal primer for simultaneous analysis of bacteria and archaea using
next-generation sequencing. *PLoS One* 2014; **9**: e105592.

24.   Wibberg D, Andersson L, Rupp O, Goesmann A, Pühler A, Varrelmann M, et al.
Genome analysis of the sugar beet pathogen Rhizoctonia solani AG2-2IIIB revealed
high numbers in secreted proteins and cell wall degrading enzymes. *BMC Genomics*
2016; 245.

25.   Magoč T, Salzberg SL. FLASH: Fast length adjustment of short reads to improve
genome assemblies. *Bioinformatics* 2011; **27**: 2957–2963.

26.   Edgar RC. Search and clustering orders of magnitude faster than BLAST.
Bioinformatics 2010; **26**: 2460–2461.

27.   Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads.
Nat Methods 2013; **10**: 996–8.

28.   Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ*
Microbiol 2007; **73**: 5261–5267.

29.   Liebe S, Wibberg D, Winkler A, Puehler A, Schlueter A, Varrelmann M. Taxonomic
analysis of the microbial community in stored sugar beets using high-throughput
sequencing of different marker genes. *FEMS Microbiol Ecol* 2016; **92**: 1–12.

30.   Klassen V, Blifernez-Klassen O, Wibberg D, Winkler A, Kalinowski J, Posten C, et al.
Highly efficient methane generation from untreated microalgae biomass. *Biotechnol*

Biofuels 2017; **10**: 186.

31. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference
sequence (RefSeq) database at NCBI: Current status, taxonomic expansion, and
functional annotation. *Nucleic Acids Res* 2016; **44**: D733–D745.

32. Boisvert S, Raymond F, Godzaridis E, Laviolette F, Corbeil J. Ray Meta: scalable de
novo metagenome assembly and profiling. *Genome Biol* 2012; **13**: R122.

33. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
2012; **9**: 357–359.

34. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. *Bioinformatics* 2009; **25**: 2078–2079.

35. Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately
reconstructing single genomes from complex microbial communities. *PeerJ* 2015; **3**:
e1165.

36. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO:
Assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics 2015; **31**: 3210–3212.

37. Waterhouse RM, Seppey M, Simao FA, Manni M, Ioannidis P, Klioutchnikov G, et al.
BUSCO applications from quality assessments to gene prediction and phylogenomics.
Mol Biol Evol 2018; **35**: 543–548.

38. Wibberg D, Blom J, Jaenicke S, Kollin F, Rupp O, Scharf B, et al. Complete genome
sequencing of Agrobacterium sp. H13-3, the former Rhizobium lupini H13-3, reveals a
tripartite genome consisting of a circular and a linear chromosome and an accessory
plasmid but lacking a tumor-inducing Ti-plasmid. *J Biotechnol* 2011; **155**: 50–62.

39. Seemann T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* 2014; **30**:
2068–2069.

40. Meyer F, Goesmann A, McHardy AC, Bartels D, Bekel T, Clausen J, et al. GenDB -
An open source genome annotation system for prokaryote genomes. *Nucleic Acids Res*
2003; **31**: 2187–2195.

41. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:
Prokaryotic gene recognition and translation initiation site identification. *BMC*
Bioinformatics 2010; **11**.

42. Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, et al. Pfam: The
protein families database. *Nucleic Acids Res* . 2014. , **42**

43. Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: New
perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res* 2017; **45**:
D353–D361.

44. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND.
Nat Methods 2015; **12**: 59–60.

45.   Huson DH, Beier S, Flade I, Górska A, El-Hadidi M, Mitra S, et al. MEGAN
662   Community Edition - Interactive Exploration and Analysis of Large-Scale Microbiome
663   Sequencing Data. *PLoS Comput Biol* 2016; **12**.

664   46.   Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, et al. dbCAN2: a meta server
665   for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 2018; 1–7.

666   47.   Blin K, Wolf T, Chevrette MG, Lu X, Schwalen CJ, Kautsar SA, et al. antiSMASH
667   4.0—improvements in chemistry prediction and gene cluster boundary identification.
668   *Nucleic Acids Res* 2017; **45**: W36–W41.

669   48.   Blom J, Kreis J, Spänig S, Juhre T, Bertelli C, Ernst C, et al. EDGAR 2.0: an enhanced
670   software platform for comparative gene content analyses. *Nucleic Acids Res* 2016; **44**:
671   W22–W28.

672   49.   Price MN, Dehal PS, Arkin AP. FastTree 2 - Approximately maximum-likelihood trees
673   for large alignments. *PLoS One* 2010; **5**.

674   50.   Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis
675   Version 7.0 for Bigger Datasets. *Mol Biol Evol* 2016; **33**: 1870–1874.

676   51.   Rojo F. Enzymes for Aerobic Degradation of Alkanes. In: Timmis KN (ed). *Handbook
677   of Hydrocarbon and Lipid Microbiology*. 2010. Springer Berlin Heidelberg, Berlin,
678   Heidelberg, pp 781–797.

679   52.   Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. The
680   carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res* 2014; **42**.

681   53.   Rehm BHA. Bacterial polymers: Biosynthesis, modifications and applications. *Nat Rev
682   Microbiol* . 2010. , **8**: 578–592

683   54.   Kulakovskaya T V., Vagabov VM, Kulaev IS. Inorganic polyphosphate in industry,
684   agriculture and medicine: Modern state and outlook. *Process Biochem* . 2012. , **47**: 1–
685   10

686   55.   Torrents E. Ribonucleotide reductases: essential enzymes for bacterial life. *Front Cell
687   Infect Microbiol* 2014; **4**.

688   56.   Webb ME, Marquet A, Mendel RR, Rébeillé F, Smith AG. Elucidating biosynthetic
689   pathways for vitamins and cofactors. *Nat Prod Rep* . 2007. , **24**: 988–1008

690   57.   Cotter PD, Ross RP, Hill C. Bacteriocins-a viable alternative to antibiotics? *Nat Rev
691   Microbiol* . 2013. , **11**: 95–105

692   58.   Sambles C, Moore K, Lux TM, Jones K, Littlejohn GR, Gouveia JD, et al.
693   Metagenomic analysis of the complex microbial consortium associated with cultures of
694   the oil-rich alga Botryococcus braunii. *Microbiologyopen* 2017; **6**.

695   59.   Allison SD, Martiny JBH. Resistance, resilience, and redundancy in microbial
696   communities. *Proc Natl Acad Sci* 2008; **105**: 11512–11519.

697   60.   Krohn-Molt I, Alawi M, Förstner KU, Wiegandt A, Burkhardt L, Indenbirken D, et al.
698   Insights into Microalga and bacteria interactions of selected phycosphere biofilms

using metagenomic, transcriptomic, and proteomic approaches. *Front Microbiol* 2017;
8.

61.   Buchan A, LeCleir GR, Gulvik CA, González JM. Master recyclers: features and
functions of bacteria associated with phytoplankton blooms. *Nat Rev Microbiol* . 2014.
, **12**: 686–698

62.   Ramanan R, Kang Z, Kim B-H, Cho D-H, Jin L, Oh H-M, et al. Phycosphere bacterial
diversity in green algae reveals an apparent similarity across habitats. *Algal Res* 2015;
8: 140–144.

63.   Reed H, Martiny J. Testing the functional significance of microbial composition in
natural communities. *FEMS Microbiol Ecol* 2007; **62**: 161–170.

64.   Rivett DW, Bell T. Abundance determines the functional role of bacterial phylotypes in
complex communities. *Nat Microbiol* 2018; **3**: 767–772.

65.   Bednar TW, Holm-Hansen O. Biotin liberation by the lichen aga Coccomyxa sp. and
by Chlorella pyrenoidos. *Plant Cell Physiol* 1964; **5**: 297–303.

66.   Aaronson S, Dhawale SW, Patni NJ, Deangelis B, Frank O, Baker H. The cell content
and secretion of water-soluble vitamins by several freshwater algae. *Arch Microbiol*
1977; **112**: 57–59.

67.   Pančić M, Kiørboe T. Phytoplankton defence mechanisms: traits and trade-offs. *Biol
Rev* 2018; **93**: 1269–1303.

Figure legends

Figure 1: Co-cultivation of axenic *B. braunii* with bacterial associates *Mycobacterium* sp.

Bb-A and *Pimelobacter* sp. Bb-B. Determination of algal growth and product formation

performance of **a.** axenic cultures and **b.** xenic cultures supplemented with bacterial isolates.

The progress of the bacterial growth is represented by the number of cells on a logarithmic

($\log_{10}$) scale.The error bars represent standard error of mean values of three biological and

three technical replicates (SE; n=9).

Figure 2: Bacterial representatives accompanying the *B. braunii* consortiae. Illustration

of the relative abundance and phylogenetic distribution of bacterial genera detected *via* high-

throughput 16S rDNA amplicon sequencing approach (see Supplementary data 1). Bacterial

genera labeled green or red are represented only in the consortia of race A or B, respectively.

The abundant members among the communities are marked bold. Black, grey and white

circles represent the nodes with calculated bootstrap values of 100, 75-100 and <75% (1,000

replicates).

Figure 3: Phylogenetic characterization of the high-quality draft and complete genomes

of *B. braunii* accompanying bacterial community. Maximum-likelihood-based

phylogenetic tree built out of a core of 53 conserved marker genes (Supplementary data 2) per

analyzed genome (bold). Black, grey and white circles represent the nodes with calculated

bootstrap values of 100, 99 and <75%, respectively (1,000 replicates).

Figure 4: Alga-bacteria interactions within the *Botryocccus braunii* consortia.

Summarizing view on the observed generic potential encoded in the high-quality draft and

complete genomes of the associates (Table 1, Figure S5), and the resultant interactions

between alga and bacteria. The size of the illustrated bacteria indicates the relative abundance

(averaged) of the respective species within the consortia. The species marked with stripes,

such as Dok-G and Gp4-G, indicate that these occur only within *B. braunii* race A or B,

respectively. The incomplete B-vitamin *de novo* biosynthesis pathways are shown by the red

capital letter B with the number of the subsequent vitamin. The arrows indicate the possible

external supply of B-vitamins within the consortia. The apparent ability of the particular

associates to store nitrogen and phosphorous as polymeric reserves is indicated by N in the

green square (cyanophycin granule polypeptides) and P in the yellow circle (PolyP,

Polyphosphate granules), respectively. The estimated potential for degradation of

hydrocarbons and carbohydrates of the bacterial associates is illustrated based on the

identified enzyme number (Table 1) in conjunction with the relative abundance of the

respective species via yellow and striped pacmans, respectively. The genetic predisposition to

the synthesis of bacteriocins and antibiotics encoded by the secondary metabolite biosynthesis

gene clusters, is indicated by the blue filled or empty triangles, respectively. *M*.Bb-A,

Mycobacterium sp. Bb-A; *P*.Bb-B, *Pimelobacter* sp. Bb-B; HC, Hydrocarbons; CH,

carbohydrates; B1, Thiamin; B2, Riboflavin; B5, Pantothenate; B7, Biotin; B12, Cobalamin.

Table 1: Selected genetic features of the *B. braunii* bacterial associates. The the high-

quality draft and complete bacterial genomes were manually inspected using a E-value cutoff

of 1×10^{-10} for specific gene groups (see Supplementary data 4). Carbohydrate-active enzymes

(CAZymes) were profiled using dbCAN2 [46] metaserver (Supplementary data 5). The

values in parentheses represent the results acquired with two tools. The key enzymes involved

in   B-vitamin   *de   novo*   biosynthesis   were   searched   in   the   EMGB   database

(https://emgb.cebitec.uni-bielefeld.de/Bbraunii-bacterial-consortium/) and manually verified

via BLASTp (Supplementary data 6). The analysis of the secondary metabolite biosynthesis

gene clusters within the high-quality draft and complete genomes was accomplished via

antiSMASH [47] 4.0 web server (Supplementary data 7). *M*. Bb-A, *Mycobacterium* sp. Bb-A;

P.Bb-B, *Pimelobacter* sp. Bb-B; RNR, Ribonucleotide reductase; GH, Glycoside hydrolases;

GT, Glycosyltransferases.

a

B. braunii
axenic control

b

B. braunii xenic cultures with
Mycobacterium sp.   *Pimelobacter* sp.

Table 1: Selected genetic features of *B. braunii* bacterial associates.

	Dev-G	Dok-G	Gp4-G	Dya-G	Ter1-G	Ter2-G	Bre1-G	Bre2-G	Hyd-G	Myc-G	*M*.Bb-A	*P*.Bb-B
Enzymes oxidizing long-chain hydrocarbons												
AlkB	-	-	-	-	-	-	-	-	-	3	3	2
AlmA	2	-	-	1	-	-	-	-	1	17	17	11
LadA	3	-	-	-	-	-	-	-	-	9	11	8
CYP153	1	-	-	2	-	-	1	-	1	40	40	12
CAZymes (Enzymes involved in carbohydrate metabolism)												
total number	164 (98)	109 (59)	111 (50)	395 (264)	240 (170)	195 (125)	86 (61)	68 (45)	96 (47)	183 (127)	182 (127)	131 (79)
GHs class	79 (56)	19 (10)	40 (26)	183 (146)	138 (117)	91 (73)	32 (27)	26 (21)	24 (14)	51 (43)	50 (42)	36 (28)
GTs class	47 (23)	55 (44)	39 (14)	103 (77)	46 (30)	52 (31)	27 (22)	20 (17)	47 (24)	60 (47)	60 (47)	53 (38)
% of total gene content	3.1 (1.9)	3.0 (1.7)	2.7 (1.2)	5.9 (3.9)	5.2 (3.7)	4.7 (3.0)	2.7 (1.9)	2.2 (1.4)	2.0 (1.0)	3.2 (2.2)	3.2 (2.2)	2.3 (1.4)
Methionine synthases												
MetH [a]	√	√	√	√	√	√	√	√	√	√	√	√
MetE	√		√				√		√	√	√	
Ribonucleoside diphosphate reductase												
RNR class I [b]	√	√	√	√	√	√	√	√	√	√	√	√
RNR class II [a]	√	√	√	√	√	√	√	√	√			√
Nutrient storage												
cyanophycin synthase		√	√		√	√	√		√	√	√	
cyanophycinase		√	√		√	√						
polyphosphate kinase	√	√	√	√	√	√	√	√	√	√	√	√
exopolyphosphotase	√	√		√		√	√	√	√	√	√	√
Vitamin *de novo* synthesis pathways												
Cobalamin (vitamin B_{12})	√									√	√	√
Vitamin B_6	√	√	√	√	√	√	√	√	√	√	√	√
Riboflavin (vitamin B_2)		√	√	√	√	√	√	√	√	√	√	√
Biotin (vitamin B_7)		√							√	√	√	√
Thiamin (vitamin B_1)		√	√				√	√	√	√	√	√
Pantothenate (vitamin B_5)		√	√	√	√	√	√	√	√	√	√	√
Nicotinate (vitamin B_3)	√		√	√	√	√	√	√	√	√	√	√
Secondary metabolite gene cluster with putative												
herbicidal activity	√									√	√	
antiviral activity												√
antifungal activity									√	√	√	√
insecticidal activity									√			
antibacterial activity	√	√	√	√			√		√	√	√	√
against Gram$^+$-bacteria									√	√	√	√
bacteriocins	√	√	√	√	√	√	√	√	√	√	√	
total cluster number	37	34	31	60	22	23	18	15	42	91	93	72

[a] Vitamin B_{12} (Cobalamin) dependent enzyme
[b] *Mycobacterium* sp. Bb-A (Myc-G and *M*.Bb-A) encodes genes for RNR class Ib and all other genomes for RNR class Ia

The high-quality draft and complete bacterial genomes were manually inspected using a E-value cutoff of 1×10^{-10} for specific gene groups (see Supplementary data 4). Carbohydrate-active enzymes (CAZymes) were profiled using dbCAN2 [46] metaserver (Supplementary data 5). The values in parentheses represent the results acquired with two tools. The key enzymes involved in

B-vitamin *de novo* biosynthesis were searched in the EMGB database (https://emgb.cebitec.uni-bielefeld.de/Bbraunii-bacterial-consortium/) and manually verified via BLASTp (Supplementary data 6). The analysis of the secondary metabolite biosynthesis gene clusters within the high-quality draft and complete genomes was accomplished via antiSMASH [47] 4.0 web server (Supplementary data 7). *M*. Bb-A, *Mycobacterium* sp. Bb-A; *P*.Bb-B, *Pimelobacter* sp. Bb-B; RNR, Ribonucleotide reductase; GH, Glycoside hydrolases; GT, Glycosyltransferases.

References

46.   Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, et al. dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 2018; 1–7.

47.   Blin K, Wolf T, Chevrette MG, Lu X, Schwalen CJ, Kautsar SA, et al. antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids Res* 2017; **45**: W36–W41.