Phytoplankton consortia as a blueprint for mutually beneficial eukaryote-bacteria ecosystems: Biocoenosis of *Botryococcus* consortia

Olga Blifernez-Klassen[1,2*], Viktor Klassen[1,2], Daniel Wibberg[2], Enis Cebeci[1], Christian Henke[2,3], Christian Rückert[2], Swapnil Chaudhari[1,2], Oliver Rupp[4], Jochen Blom[4], Anika Winkler[2], Arwa Al-Dilaimi[2], Alexander Goesmann[4], Alexander Sczyrba[2,3], Jörn Kalinowski[2], Andrea Bräutigam[2,5], Olaf Kruse[1,2*]

[1]Algae Biotechnology and Bioenergy, Faculty of Biology, Bielefeld University, Universitätsstrasse 27, 33615 Bielefeld, Germany

[2]Center for Biotechnology (CeBiTec), Bielefeld University, Universitätsstrasse 27, 33615 Bielefeld, Germany

[3]Computational Metagenomics, Faculty of Technology, Bielefeld University, Universitätsstrasse 25, 33615 Bielefeld, Germany

[4]Bioinformatics and Systems Biology, Justus-Liebig-University, Heinrich-Buff-Ring 58, 35392 Gießen, Germany

[5]Computational Biology, Faculty of Biology, Bielefeld University, Universitätsstrasse 27, 33615 Bielefeld, Germany

*Address correspondence to: Olaf Kruse, olaf.kruse@uni-bielefeld.de or

Olga Blifernez-Klassen olga.blifernez@uni-bielefeld.de

Running title: Biocoenosis of *Botryococcus* consortia

Abstract

Bacteria occupy all major ecosystems and maintain an intensive relationship to the
eukaryotes, developing together into complex biomes (i.e., phycosphere and rhizosphere).
Interactions between eukaryotes and bacteria range from cooperative to competitive, with the
associated microorganisms affecting their host`s development, growth, health and disease.
Since the advent of non-culture dependent analytical techniques such as metagenome
sequencing, consortia have been described but owing to the complex interactions rarely
functionally dissected. Multifaceted analysis of the microbial consortium of the ancient
phytoplankton *Botryococcus* as an attractive model food web revealed that its all abundant
bacterial members belong to a distinct niche of biotin auxotrophs, essentially depending on
the microalga. In addition, hydrocarbonoclastic bacteria without vitamin auxotrophies, which
adversely affect the algal cell morphology, appear evidently decimated. Synthetic
rearrangement of a minimal community consisting of alga, mutualistic and parasitic bacteria
underpins the model of a eukaryote that domesticates its own mutualistic bacterial "zoo" to
manipulate and control its surrounding biosphere. This model of domestication of mutualistic
bacteria for the defense against destruents by a eukaryotic host could represent ecologically
relevant interactions that cross species boundaries. Metabolic and system reconstruction
disentangles the relationships and provide a blueprint for the construction of mutually
beneficial synthetic ecosystems.

Keywords: eukaryote-bacteria interactions, phycosphere, mutualism, parasitism, microbiome,
biotin (B-vitamin) auxotrophy, *Botryococcus braunii*, synthetic ecosystem, (meta)genome
reconstruction

Background

Bacteria are omnipresent and affect likely every eukaryotic system through their presence

(Grosberg and Strathmann, 2007). Microalgae are the dominant primary producers and the

base of the food web within aquatic ecosystems and sustain in the natural environment in

close relationship with multiple associated microorganisms (Ramanan et al., 2016; Seymour

et al., 2017). Thus, the interactions between phytoplankton and bacteria represent a

fundamental ecological relationship within aquatic environments (Seymour et al., 2017), by

controlling nutrient cycling and biomass production at the food web base. The nature of the

exchange of micro- and macronutrients, diverse metabolites including complex products such

as polysaccharides, and infochemicals defines the relationship of the symbiotic partners (Cole,

1982; Amin et al., 2015; Zengler and Zaramela, 2018), which span mutualism, commensalism

and parasitism (Ramanan et al., 2016). Within a parasitic association, the bacteria are known

to adversely affect the algae (Wang et al., 2010), while in commensalism the partners can

survive independently, with the commensal feeding on algae-provided substrates, but without

harming the host (Ramanan et al., 2016). Mutualistic interactions include *i.a.* the obligate

relationships between vitamin-synthesizing bacteria and vitamin-auxotrophic phytoplankton

species (Croft et al., 2005). Many microalgae cannot synthesize several growth-essential

vitamins (like B_{12} and B_1) (Croft et al., 2005; Croft et al., 2006; Tang et al., 2010) and obtain

them through a symbiotic relationship with bacteria in exchange for organic carbon (Grant et

al., 2014). Recent observations suggest that not only certain microalgae require vitamins, but

many bacteria also have to compete with other organisms for the B-vitamins (Sañudo-

Wilhelmy et al., 2014; Gómez-Consarnau et al., 2018).

However, in microbial environments, where the distinction between host and symbionts is

less clear, the identification of the associated partners and the nature of their relationship is

challenging. Looking for a suitable model food web system to elucidate the complexity of

eukaryote-prokaryote interactions, we focused on the microalga-bacteria consortia with an

organic carbon-rich phycosphere. The planktonic chlorophyta *Botryococcus braunii* excretes

chemically inert long-chain hydrocarbons and a variety of exo-polysaccharides (Banerjee et

al., 2002) which allows this single celled alga to form large agglomerates (Grosberg and

Strathmann, 2007). Both organic carbon-based products are exposed to the surrounding

ecosystem and enable the alga to sustain in the natural environment (phycosphere) in close

relationship with multiple associated microorganisms (Ramanan et al., 2016; Seymour et al.,

2017). *Botryococcus* exudes up to 46% of their photosynthetically fixed carbon into the

phycosphere (Blifernez-Klassen et al., 2018), similarly to processes, however with lesser

extent, observed for seedlings as well as adult plants within the rhizosphere in the form of

poorly characterized rhizodeposits (Nguyen, 2003; Bisseling et al., 2009; Marschner, 2011).

B. braunii consortia, naturally accompanied with a variety of microorganisms (Chirac et al.,

1985), may represent an example for an association already existing during the early stages of

life development on earth, since it is generally considered that a progenitor of *Botryococcus*

impacted the development of today's oil reserves and coal shale deposits (McKirdy et al.,

1986). Thus, the microbiome associated with *Botryococcus braunii* presents a unique

opportunity to dissect the roles of the eukaryote and prokaryotes within the consortium and

test if and how the eukaryote manipulates its biome.

The interactions between *B. braunii* and the associated bacteria are not fully understood,

but are known to influence the microalgal growth performance and product formation

capacity (Chirac et al., 1985; Banerjee et al., 2002; Tanabe et al., 2015; Fuentes et al., 2016).

To understand the complex nature of the unique habitat of a microalgal phycosphere and to

unravel the food web between alga and bacteria, we investigated the consortia accompanying

the hydrocarbon/carbohydrate-producing green microalga *Botryococcus braunii*.

Results and discussion

Metagenomic survey of the *Botryococcus braunii* consortia

To determine the inhabitants of the *B. braunii* ecosystem, we profiled four different

Botryococcus communities (races A and B) using high-throughput 16S rDNA gene amplicon

and read-based metagenome sequencing approaches (Figures S1-3; Figure 1a). Assembly and

functional annotation of the metagenomes together with single strain isolation and sequencing

reconstructed the capabilities of the most abundant consortial microorganisms and were used

to design and test synthetic alga-bacteria ecosystems. The analysis of the 16S rDNA

amplicons and a read-based metagenome approach showed that each of the *B. braunii*

communities represents a comparable conserved consortium of a single algal and various

bacterial species as well as traces of Archaea and viruses (Figure S2). 16S rDNA amplicon

sequencing identified 33 bacterial taxa (Figure S3). The relative abundance of plastid 16S

rDNA sequences accounted up to 49 and 59% of all detected amplicon reads in race A and B

samples, respectively. Metagenomic assembly resulted in ten high quality draft genomes with

>80% assembled. Three additional genomes were closed by single strain sequencing (Figure

1; Tables S1 and S2; Supplementary Discussion, Chapter 1). The bacterial community varied

quantitatively, but not qualitatively between growth phases and strains (Figure S3). *B. braunii*

consortia are inhabited by the same bacterial phyla (*Proteobactreria*, *Bacteroidetes*,

Acidobacteria and *Actinobacteria*), albeit in different relative abundances, compared to those

phyla which occupy the plant rhizosphere (Xu et al., 2018) and the mammals gut (Ley et al.,

2008).

Figure 1: **Taxonomy of the *B. braunii* bacterial community and selected functional categories**
encoded in reconstructed MAGs and complete genomes. (**a**) Normalized read-based comparison
and taxonomic assignment of different *B. braunii* metagenome datasets. The circle size represents the
amount of classified reads for the respective taxonomic group (minimum 50 reads per taxonomic
group). (**b**) Abundance of selected functional categories encoded in the high-quality draft and
complete bacterial genomes. Numbers in parentheses represent the maximal total number of genes
within each pathway. Percentage: per cent of total number of genes per pathway identified within each
genome. All annotated genes within the respective pathways are presented on the EMGB platform
(https://emgb.cebitec.uni-bielefeld.de/Bbraunii-bacterial-consortium/). The pathways statistics are
summarized in the Table S5.

During the growth phase, the *Botryococcus* phycosphere is dominated by the genera of
Devosia, Dyadobacter, Dokdonella, Acidobacteria, Brevundimonas, Mesorhizobium and
Hydrogenophaga which account for over 80% of the bacteria (Table S1). The re-processing
of the *B. braunii* Guadeloupe strain metagenome data (Sambles et al., 2017) and direct
comparison of the different datasets revealed the coinciding presence of the individual
bacterial genera, although massively varying in the abundance pattern in dependence of
cultivation conditions (Figure S5).The same taxa are also present *i.a.* in the active microbiome
associated roots and rhizosphere of oilseed rape (Gkarmiri et al., 2017), as well as in the
lettuce (Schreiter et al., 2014) and global citrus rhizosphere (Xu et al., 2018), strongly
suggesting an overall mutual dependence or preferred coexistence.

To identify genes that enable the bacterial community members to contribute to and to
benefit from the food web of the consortium, we performed a KEGG-based quantitative
functional assignment of the annotated high-quality MAGs (metagenome assembled
genomes). Based on these data sets, we sequenced genomes of isolated associates and
imported them to the EMGB platform (elastic metagenome browser, https://emgb.cebitec.uni-
bielefeld.de/Bbraunii-bacterial-consortium/; for details see Methods; Supplementary
Discussion, Chapter 2). We originally expected a uniform distribution of abilities and a high
degree of interconnectedness between all members of the consortium. Surprisingly, the
consortium divides into two functionally distinct groups of bacteria: The dominant consortia
members such as *Devosia* (DevG), *Dyadobacter* (DyaG) and *Brevundimonas* (Bre1G, *B*.Bb-
A) do not oxyfunctionalize hydrocarbons (Rojo, 2010), but their genomes are rich in genes
involved in the degradation/assimilation of carbohydrates (Figure 1b). On the contrary, the
bacterial strains *Mycobacterium* and *Pimelobacter* (*M*.Bb-A and *P*.Bb-B), which are of low
abundance within the consortia (Figure S3), contain a large portfolio of genes encoding
monooxygenases/hydroxylases for hydrocarbon degradation (Rojo, 2010) and hold the
potential for a pronounced catabolic capacity towards lipids, fatty acids and xenobiotics
(Figure 1b). Besides, *Mycobacterium* and *Pimelobacter* genomes were the only genomes
among the abundant community members to carry non-heme membrane-associated
monooxygenase *alkB* genes (Table S6). To test these observed genetic predisposition of the
consortia, we assessed algal capacities for growth and hydrocarbon production in co-
cultivation of the axenic *B. braunii* with one representative of each group.

Effect of the bacterial community on the *B. braunii* host

One representative of the core as well as less abundant community members were isolated

in axenic culture (*Brevundimonas* sp. Bb-A and *Mycobacterium* sp. Bb-A, respectively).

Synthetic algae-bacteria consortia were formed with an axenic hydrocarbon- and carbohydrate

producing *B. braunii* race A strain under strict phototrophic conditions (*e.g.* without the

supplementation of vitamins or organic carbon). Axenic *Botryococcus braunii* showed a

continuous increase in cell biomass and hydrocarbon content, yielding 2.1±0.2 gL^{-1} and

0.7±0.1 gL^{-1}, respectively (Figure 2a, b; Figure S6). Co-cultivation of the *Brevundimonas* sp.

Bb-A promotes growth of the alga and increases biomass by 60.5±16.9% compared to axenic

controls (Figure 2a) while co-cultivation with the *Mycobacterium* sp. Bb-A reduces growth by

30.6±4.8% (Figure 2b).

Figure 2: **Physiological effect of the individual bacterial isolates on the microalgae during co-**
cultivation. (**a-d**) Determination of algal growth and product formation performance of axenic and
xenic *B. braunii* cultures, supplemented with bacterial isolates. Shown are growth analysis of *B.*
braunii in the presence of (**a**) *Mycobacterium* sp. Bb-A and (**b**) *Brevundimonas* sp. Bb-A; (**c**)
comparison of the hydrocarbon levels observed in samples cultivated with bacterial isolates in relation
to the axenic *B. braunii* culture in the course of the cultivation; (**d**) morphological characteristic of
algal cells during the cultivation (9 days) with the bacterial isolates (red arrows indicate the
hydrocarbons produced by the microalga). Graphs imbedded in the pictures show the progress of the
bacterial growth (cell number on log$_{10}$ scale).The error bars represent standard error of mean values of
three biological and three technical replicates (SE; n=9).

B. braunii accumulates hydrocarbons up to 38% of its dry weight (DW) in a bacteria-free

environment. In contrast, co-cultivation with *Mycobacterium* limits hydrocarbon

accumulation down to $4.4\pm0.2\%$ of DW (Figure 2c) and alters colony morphology (Figure

2d). Loss of the hydrocarbon matrix reduces colony size (Figure 2d) and likely increases

susceptibility to grazers (Pančić and Kiørboe, 2018). The malfunction of colony formation

can thus be attributed to imbalanced and dysbiotic host-microbiome interaction similar to

imbalanced interactions previously observed (Gilbert et al., 2016; Durán et al., 2018) and

Mycobacterium essentially acts as a pathogen. The dramatic decrease in hydrocarbon content

and disturbed colony formation cannot be observed in cultures supplemented with

Brevundimonas or in the xenic *Botryococcus braunii* algae-bacteria communities (Figure 2c,

d; Figures S1 and S6), pointing to a balanced equilibrium between the alga and its

microbiome in natural environments.

Since *Botryococcus* readily releases huge amounts of organic carbon into the extracellular

milieu (Blifernez-Klassen et al., 2018), creating a phycosphere that naturally attracts many

microorganisms, including those with potential damaging effects, its evolutionary fitness will

depend on efficient management of the "bacterial zoo". We hypothesized that there is a strong

control mechanism that allows the algae host to attract and control symbiotic / beneficial

bacteria. Therefore, we examined the genetic portfolio of the bacterial community for

essential cofactors and interdependencies.

The core community of *Botryococcus* phycosphere belongs to a distinct niche of biotin

auxotrophs

At first glance, the *Botryococcus* alga does not appear to be obligatory dependent on a

bacterial community, since axenic strains can be fully photoautotrophically cultured without

vitamin supplements (Figure 2 a, b) and contains the vitamin B_{12}-independent form of

methionine synthase (*metE*) (Molnár et al., 2012). (Meta)Genome reconstruction revealed that

the *B. braunii* bacterial core community is auxotrophic for various B-vitamins (Figure 3a),

making them dependent on an exogenous supply of vitamins for survival. All abundant genera

within the *B. braunii* consortia (including the growth-promoting *Brevundimonas*) lack the

complete gene portfolio for biotin (vitamin B_7) synthesis (Figure 3a), but contained the

complete biosynthesis pathway of fatty acids for which biotin is essential (Sañudo-Wilhelmy

et al., 2014). The complete thiamin synthesis pathway (vitamin B_1, essential for citrate cycle

and pentose phosphate pathway (Sañudo-Wilhelmy et al., 2014)) could not be reconstructed

for four assembled genomes. Five genera are auxotrophic for cobalamin (vitamin B_{12}).

Although, auxotrophic for various B-vitamins including biotin, the consortia's most abundant

species *Devosia* (DevG) contains the complete pathway for vitamin B_{12} synthesis. In view of

the fact that the bacterial abundance determines their functional relevance for the community

(Reed and Martiny, 2007; Rivett and Bell, 2018), this species is also likely to provide both the

bacterial community and the microalga with this vitamin (Croft et al., 2005), thus confirming

the recent proposition that the B_{12} supply is realized only by a few certain bacteria within the

alga-bacteria consortia (Krohn-Molt et al., 2017). In this context, since the B_7-auxotrophs are

prevalent within the detected consortia, it is doubtful that the large required supply of biotin is

of bacterial origin. Because of its predominance within the community (Figure S2), the

microalga is more likely to secrete and provide this vitamin to the bacteria.

Figure 3: Essential cofactors (inter-)dependencies of the bacterial core community and isolates of *B. braunii*. (**a**) Heatmap of the reconstructed *de novo* biosynthesis of B-vitamins encoded in the high-quality draft and complete bacterial genomes. Numbers in parentheses represent the maximal total number of genes within each pathway. Percentage: per cent of total number of genes per pathway identified within each genome. All annotated genes within the respective pathways are presented on the EMGB platform (https://emgb.cebitec.uni-bielefeld.de/Bbraunii-bacterial-consortium/). The pathways statistics are summarized in the Table S5 and S8. (**b**) Growth assay to confirm B-vitamin auxotrophy and prototrophy in axenic *Brevundimonas* sp. Bb-A (blue) and *Mycobacterium* sp. Bb-A (orange) strains, respectively. (**c**) Biotin concentration levels detected in cell-free supernatants from the cultures of axenic *B. braunii* as well as in the presence of *Mycobacterium* sp. Bb-A (orange) and *Brevundimonas* sp. Bb-A (blue). The error bars represent standard error of mean values of three biological and three technical replicates (SE; n=9).

It is well known, that many microalgae rely on the external supply of growth-essential

vitamins like cobalamin and thiamin (Croft et al., 2005; Croft et al., 2006; Tang et al., 2010),

however, biotin auxotrophy is rare among phytoplankton species (Croft et al., 2006). Here,

our genome reconstruction results strongly suggests that also bacteria have to compete with

other organisms for the B-vitamins and live in a close mutualistic relationship with the

microalga strongly depending on the biotin supply. The growth promoting effect of

Brevundimonas (Figure 2b), which depends on an exogenous supply of biotin for survival, is

thus based on the close interaction with the microalga. In contrast, low abundant consortial

members such as *Mycobacterium* are B-vitamin prototrophs (Figure 3a).

Extended experimental analysis confirmed the biotin (vitamin B_7) auxotrophy in

Brevundimonas sp. Bb-A and vitamin B prototrophy in *Mycobacterium* sp. Bb-A as deduced

from the assembled genomes (Figure 3b). *Botryococcus braunii* secrets biotin into the

surrounding environment (Figure 3c) and secretion is increasingly induced by the presence of

bacteria, particularly in the cultures with the B_7-auxotrophs (compare to Figure 3c at

timepoint 3 days). The secretion of some B-vitamins has already been observed in microalgae

(Aaronson et al., 1977), and also plants support their rhizosphere by active riboflavin (vitamin

B_2) excretion from their roots (Higa et al., 2008). However, the eukaryotic B-vitamin

secretion (especially of biotin), which was increasingly induced by the presence of bacteria, as

well as their simultaneous active uptake, remained so far unrecognized.

Collectively, in a healthy (natural) community, non-hydrocarbon degrading bacterial

genera which are auxotrophic for one or more vitamins dominate the phycosphere, while fully

prototrophic hydrocarbonoclastic bacteria are present but not abundant. We hypothesized that

B. braunii cultivates its own auxotrophic "bacterial zoo" to combat hydrocarbonoclastic

bacteria. To test the hypothesis, synthetic consortia of alga–"good bacteria"–"bad bacteria"

were formed and analyzed.

Bacterial biotin-auxotrophic symbionts fend their eukaryotic host

292   We hypothesized that the alga maintains its balanced, 'healthy' microbiome by recruiting

293   beneficial, vitamin B-auxotrophic microorganisms with the provision of photosynthetically

294   fixed carbon and biotin and thus counteracts pathogen assault. *Brevundimonas* and

295   *Mycobacterium* were co-cultivated with the axenic *Botryococcus* alga under photoautrophic

296   conditions with three different inoculum ratios (Figure 4a).

Figure 4: **Bacterial symbionts fend their eukaryotic host from hydrocarbonoclastic invaders.**
Physiology of a synthetic alga-bacteria-bacteria community, consisting of *Brevundimonas* sp. Bb-A
(blue) and *Mycobacterium* sp. Bb-A (orange) in co-culture with *B. braunii* (green) under
photoautrophic conditions with three different inoculum ratios. Shown are (**a**) the progression of
bacterial proliferation (cfu, colony-forming units) and (**b**) detected relative hydrocarbon levels
(normalized to the axenic control) during the co-cultivation. The error bars represent standard error of
mean values of three biological and three technical replicates (SE; n=9).

Independently from the initial amount of inoculum, biotin-auxotrophic *Brevundimonas*

proliferated strongly within this synthetic community. *Mycobacterium* did not proliferate in

the presence of *Brevundimonas* (Figure 4a), but was clearly capable of growth in the presence

of the alga alone (Figure 2b). *Brevundimonas* apparently produces bioactive compounds with

bacteriostatic properties, which likely decimate the proliferation of *Mycobacterium*. Indeed, all

reconstructed genomes encode metabolic gene cluster for antibiotic compounds (especially

against Gram-positive bacteria) and for bacteriocins (Figure 1b, Table S9) with significant

antimicrobial potency (Cotter et al., 2013), the potential of which should be subject of further

research. The protective effect of *Brevundimonas* is evident from the hydrocarbon content of

cultures which increases with decreasing proportions of *Mycobacterium* (Figure 4b).

Experimental analyses (Figures 2-4) thus confirmed key conjectures derived from

metagenomic assemblies (Figure 1).

In conclusion, the interactions between the microalga and bacteria within the *Botryococcus*

braunii consortia depends on a complex interplay between all members, with the microalga as

the "conductor" in this play (Figure 5). Our observations indicate that the microalga is

associated with a "zoo" of largely supportive bacteria and suggest a framework in which the

microalga specifically attracts vitamin-auxotrophic bacterial species and domesticates them

by controlling vitamin supply. Since autark bacteria without vitamin auxotrophies are present

in small proportions only (Figure S3; Figure 3a), they are likely decimated by antibacterial

agents from the supportive community.

Figure 5: **Alga-bacteria interactions within the *Botryococcus braunii* phycosphere.** The depiction
summarizes the observed generic potential encoded in the (meta)genomes of bacterial associates
(Figures 1and 3), along with experimental validation within the minimal synthesis community
(Figures 2 and 4). The color code indicates the functional grouping of the community members:
supportive B-vitamin-auxotrophic (blue) and hydrocarbon-clastic prototrophic (red). The potential for
degradation of hydrocarbons and carbohydrates of the bacterial associates is illustrated via yellow and
striped pacmans, respectively. The genetic predisposition to the synthesis of bacteriocins and
antibiotics encoded by the secondary metabolite biosynthesis gene clusters, is indicated by the halos
and triangles, respectively. Abbreviation: HC, Hydrocarbons; CH, carbohydrates; B_7, Biotin; N,
nitrogen; P, phosphorus.

The analyses mechanistically resolved biotin as a critical metabolite and indicated the

presence of at least two additional communication signals: a bacterial signal increasing biotin

production (Figure 3c) and a bacteriostatic compound directed from *Brevundimonas* at

Mycobacterium (Figure 4a). The metagenomes hint at additional beneficial interactions

between the alga and its cultivated "zoo": vitamin B_{12}, and assimilated N and P from the

bacteria in exchange for biotin (and other vitamins) as well as carbon from the alga (Figure 5).

The successful design of minimal synthetic alga-bacteria cultures consisting only of two

bacteria species and the alga precisely demonstrates a functional triangle relationship for

preventing uncontrolled overgrowing of hydrocarbonoclastic destruents. However, the

elucidation of the communication signals exchanged between the participating parties will be

the subject of further research.

The metagenomic analysis of the *Botryococcus* consortium presents a path towards

mechanistic resolution of ecological interactions at the microscopic scale. Metabolic and

system reconstruction unravels the relations in the eukaryote-bacteria consortia and provides a

blueprint for the construction of mutually beneficial synthetic ecosystems.

Methods

Strains, cultivation conditions and growth determination parameters. Liquid, xenic

cultures of *Botryococcus braunii* race A (CCALA778, isolated in Serra da Estrela, Baragem

da Erva da Fome, Portugal) and race B (AC761, isolated from Paquemar, Martinique, France)

as well as the axenic *B. braunii* race A strain (SAG30.81, Peru, Dpto.Cuzco, Laguna Huaypo)

were used for co-cultivation studies. The cultures were phototrophically cultivated under 16:8

light-dark illumination of 350±400 μmol photons m^{-2} s^{-1} white light in modified Chu-13

media (without the addition of vitamin and organic carbon source) as described before

(Blifernez-Klassen et al., 2018). Carbon supply and agitation were achieved by bubbling the

cultures with moisture pre-saturated carbon dioxide-enriched air (5%, v/v) with a gas flow

rate of 0.05 vvm. Cultures growth was monitored by measurements of biomass organic dry

weight (DW) by drying 10 mL culture per sample (in three biological and technical replicates)

following the protocol described before (Astals et al., 2012). Additionally, microalgae and

bacteria cells were regularly checked by optical microscopy.

The bacterial isolates were obtained by plating highly diluted xenic *B. braunii* cultures

(CCALA778 and AC761) on LB agar plates supplied with vitamins (0.1 nM cobalamin, 3.26

nM thiamine and 0.1 nM biotin). After 1-2 weeks single cell colonies were picked, separately

plated and analyzed *via* 16S rDNA Sanger-sequencing by using the universal primer pair 27F

(5'-AGAGTTTGATCCTGGCTCAG-3') and 1385R (5'-CGGTGTGTRCAAGGCCC-3').

Two isolates, deriving from algal race A and one from race B, were classified based on full

length 16S rDNA similarity as *Mycobacterium* sp., *Brevundimonas* sp. and *Pimelobacter* sp.,

respectively. *Mycobacterium* sp. and *Brevundimonas* sp. were used for further analysis. The

purity of the axenic alga and bacterial strains was regularly verified by polymerase chain

reaction (PCR) amplification and Sanger-sequencing of the 16S rRNA gene using a universal

primer pair 27F and 1385R.

For experiments testing the requirement of B-vitamins, mono-cultures of *Mycobacterium* sp.

Bb-A and *Brevundimonas* sp. Bb-A were pre-washed three times in minimal medium

(Schlegel et al., 1961) and then starved of vitamins for two days prior to the start of the

experiment. Then, bacterial cells were diluted to 0.2 (oD_{600}) with minimal medium

supplemented with $1 gL^{-1}$ glucose and B-vitamins (at the following concentrations: 0.1 nM

(each) of cobalamin, riboflavin, biotin and pantothenate as well as 3.26 nM thiamine) and

grown for seven days.

For the co-cultivation experiments, 500 mL axenic algal cultures (~0.25g L^{-1}) were inoculated

with the isolates (*Mycobacterium* sp. Bb-A or *Brevundimonas* sp. Bb-A, with ~$2x10^6$ cell mL^{-1}

for single isolate cultivation and with ~$2x10^6 – 2x10^7$ cell mL^{-1} for co-cultivation with both

bacterial strains) and cultured under photoautotrophic conditions. The bacterial cell density

was determined by manual cell counting using hemocytometer or by determination of the

colony-forming units (cfu) as a proxy for viable population density using the replica plating

method (Jett et al., 1997).

The quantitative determination of the vitamin B_7 (biotin) concentration in cell-free

supernatants of the co-cultures and the axenic control was accomplished by using

microbiological microtiter plate test (VitaFast, r-biopharm, Darmstadt, Germany) according to

manufacturer instructions.

Hydrocarbon extraction and analysis. Hydrocarbon extraction was performed according to

the protocol published previously (Blifernez-Klassen et al., 2018). The dry extracted

hydrocarbons were resuspended in 500 µL of *n*-hexane, containing an internal standard *n*-

hexatriacontane ($C_{36}H_{74}$) and analysed *via* GC-MS and GC-FID as described before

(Schwarzhans et al., 2014; Blifernez-Klassen et al., 2018).

DNA sample collection and preparation. The samples from xenic *B. braunii* race A and B

cultures (three biological replicates) were taken in the linear and stationary growth phases

(after 9 and 24 days). Total community genomic DNA was extracted as previously described

(Zhou et al., 1996). The quality of the DNA was assessed by gel electrophoresis and the

quantity was estimated using the Quant-iT PicoGreen dsDNA Assay Kit (Invitrogen) and a

Tecan Infinite 200 Microplate Reader (Tecan Deutschland GmbH).

High-throughput 16S rDNA amplicon sequencing. To get insight into the community
composition of *B. braunii* Race A and B, high-throughput 16S rDNA amplicon sequencing
was performed as described recently by Maus et al (Maus et al., 2017). The primer pair
Pro341F   (5′-CCTACGGGGNBGCASCAG-3′)   and   Pro805R   (5′-
GACTACNVGGGTATCTAATCC-3′) was used to amplify the hypervariable regions V3 and
V4 of diverse bacterial and archaeal 16S rRNAs (Takahashi et al., 2014). In addition, the
primers also cover the 16S rDNA gene of algal chloroplasts and eukaryotic mitochondrial
genomes. Furthermore, in a two PCR steps based approach, multiplex identifier (MID) tags
and Illumina-specific sequencing adaptors were added to each amplicon. Only amplicons
featuring a size of ~460 bp were purified using AMPureXP® magnetic beads (Beckman
Coulter GmbH, Brea, California, USA). Resulting amplicons were qualitatively and
quantitatively analyzed using the Agilent 2100 Bioanalyzer system (Agilent Inc., Santa Clara,
California, USA) and pooled in equimolar amounts for paired-end sequencing on the Illumina
MiSeq system (Illumina, San Diego, California USA). This sequencing approach provided
~150,000 reads per sample. An in-house pipeline as described previously (Wibberg et al.,
2016) was used for adapter and primer trimming of all samples. For amplicon processing, a
further in house amplicon processing pipeline including FLASH (Magoč and Salzberg, 2011),
USEARCH (Edgar, 2010) (v8.1), UPARSE (Edgar, 2013) and the RDP (Wang et al., 2007)
classifier (v2.9) was applied as described recently (Liebe et al., 2016; Klassen et al., 2017). In
summary, unmerged sequences resulting by FLASH (Magoč and Salzberg, 2011) (default
settings +-M 300) were directly filtered out. Furthermore, sequences with >1Ns (ambiguous
bases) and expected errors >0.5 were also discarded. Resulting data was further processed and
operational taxonomic units (OTUs) were clustered by applying USEARCH (Edgar, 2010)
(v8.1). The resulting OTUs were taxonomically classified using the RDP (Wang et al., 2007)
classifier (v2.9) in 16S modus (Threshold >0.8) and compared to the nt database by means of
BLASTN (O'Leary et al., 2016). In the last step, raw sequence reads were mapped back onto
the OTU sequences in order to get quantitative assignments.

(Meta)Genome sequencing, assembly, binning and annotation. For each of the four
samples (*B. braunii* race A and B from linear (T1) and stationary (T2) growth phase), the
genomic DNA from three replicates was pooled. To obtain the metagenome sequence, four
whole-genome-shotgun PCR-free sequencing libraries (Nextera DNA Sample Prep Kit;
Illumina, Munich, Germany) were generated based on the manufacturer's protocol
representing different time points of the cultivation and different algae communities. The four
libraries were sequenced in paired-end mode in a MiSeq run (2 x 300 bp) resulting in

11,963,558 and 11,567,552 reads for race A T1 and T2; 6,939,996 and 12,031,204 reads for

race B T1 and T2, respectively (total 12.75 Gb). The *de novo* metagenome assembly was

performed using the Ray Meta assembler (Boisvert et al., 2012) (v2.3.2) using a k-mer size of

41 and default settings. For contigs with length larger than 1 kb (n=127,914), the total

assembly size was 338.5 Mb, the N50=2,884 and the largest contig 1.59 Mb. All processed

raw sequence reads of the four samples were aligned to the assembled metagenome contigs by

means of Bowtie 2 (Langmead and Salzberg, 2012) (v2.2.4). By using SAMtools (Li et al.,

2009) (v1.6), the SAM files were converted to BAM files. These files were sorted and read

mapping statistics were calculated. The following portions of reads could be assembled to the

draft metagenomes: 70.1% and 77.5% for race A (T1 and T2); 51.0% and 62.9% for race B

(T1 and T2). For binning of the metagenome assembled genomes (MAGs), the tool MetaBAT

(v0.21.3) (Kang et al., 2015) was used with default settings. Completeness and contamination

of the resulting MAGs were tested with BUSCO (Simão et al., 2015; Waterhouse et al., 2018)

(v3.0.), using the Bacteria and Metazoa data set (Table S1).

For sequencing of the isolated species, 4 µg of purified chromosomal DNA was used to

construct PCR-free sequencing library and sequenced applying the paired protocol on an

Illumina MiSeq system, yielding ~1.71 Mio Reads (517 Mb in 3 Scaffolds, 58 Contigs) for

Mycobacterium sp. Bb-A and ~1.70 Mio Reads (503 Mb in 6 Scaffolds, 65 Contigs) for

Pimelobacter sp. Bb-B. Obtained sequences were *de novo* assembled using the GS *de novo*

Assembler software (v2.8, Roche). An *in silico* gap closure approach was performed as

described previously (Wibberg et al., 2011). MinION sequencing library with genomic DNA

from *Brevundimonas* sp. Bb-A was prepared using the Nanopore Rapid DNA Sequencing kit

(SQK-RAD04) according to the manufacturer's instructions with the following changes: The

entry DNA amount was increased to 800 ng and an AMPure XP bead cleanup was carried out

after transposon fragmentation. Sequencing was performed on an Oxford Nanopore MinION

Mk1b sequencer using a R9.5 flow cell, which was prepared according to the manufacturer's

instructions. MinKNOW (v1.13.1) was used to control the run using the 48h sequencing run

protocol; base calling was performed offline using albacore (v2.3.1). The assembly was

performed using canu v1.7 (Koren et al., 2017), resulting in a single, circular contig (170 Mb,

1kb to 17.5kb). This contig was then polished with Illumina short read data from the

metagenome data set using pilon (Walker et al., 2014), run for sixteen iterative cycles. bwa-

mem (Li and Wren, 2014) was used for read mapping in the first eight iterations and bowtie2

v2.3.2 (Langmead and Salzberg, 2012) in the second set of eight iterations. Annotation of the

draft chromosomes was performed within the PROKKA (Seemann, 2014) (v1.11) pipeline

and visualized using the GenDB 2.0 system (Meyer et al., 2003).

Gene prediction, taxonomic assignment and functional characterization. Gene prediction

on the assembled contigs of the MAGs was performed with Prodigal (Hyatt et al., 2010)

(v2.6.1) using the metagenome mode ("-p meta"). All genes were annotated and analyzed

with the in-house EMGB annotation system using the databases NCBI NR (O'Leary et al.,

2016), Pfam (Finn et al., 2014) and KEGG (Kanehisa et al., 2017). Based on DIAMOND

(Buchfink et al., 2015) (v0.8.36) hits against the NCBI-NR (O'Leary et al., 2016) database, all

genes were subject to taxonomic assignment using MEGAN (Huson et al., 2016).

Relationships to metabolic pathways were assigned based on DIAMOND (Buchfink et al.,

2015) hits against KEGG (Kanehisa et al., 2017). For Pfam (Finn et al., 2014) annotations,

pfamscan was used with default parameters. The genomes were manually inspected using an

E-value cutoff of 1×10^{-10} for several specific sequences listed Additional file 2, Table S6. The

profiling of the carbohydrate-active enzymes (CAZy) encoded by the *B. braunii* community

members was accomplished using dbCAN2 (Zhang et al., 2018) metaserver (Table S7). The

identification, annotation and analysis of the secondary metabolite biosynthesis gene clusters

within the MAGs and complete genomes were accomplished *via* the antiSMASH (Blin et al.,

2017) (v4.0) web server with default settings (Table S9).

Phylogenetic analyses. To assign and phylogenetically classify the MAGs, a core genome

tree was created by applying EDGAR (Blom et al., 2016) (v2.0). For calculation of a core-

genome-based phylogenetic tree, the core genes of all MAGs and the three isolates

Mycobacterium sp. Bb-A, *Brevundimonas* sp. Bb-A and *Pimelobacter* sp. Bb-B,

corresponding to selected reference sequences were considered. The implemented version of

FASTtree (Price et al., 2010) (using default settings) in EDGAR (Blom et al., 2016) (v2.0)

was used to create a phylogenetic Maximum-Likelihood tree based on 53 core genes (see

Table S4) of all genomes. Additionally, the MAGs were taxonomically classified on species

level by calculation of average nucleic and amino acid identities (ANI, AAI) by applying

EDGAR (Blom et al., 2016) (v2.0). For phylogenetic analysis of the 16S rDNA amplicon

sequences, MEGA (Kumar et al., 2016) (v7.0.26) was used. Phylogenetic tree was

constructed by means of Maximum-Likelihood Tree algorithm using Tamura 3-parameter

model. The robustness of the inferred trees was evaluated by bootstrap (1000 replications).

Availability of data and materials

The high-throughput 16S rDNA amplicon sequencing datasets obtained during the present

work are available at the EMBL-EBI database under BioProjectID PRJEB21978. The

metagenome datasets of *Botryococcus braunii* consortia as well as the resulting MAGs have been deposited under the BioProjectIDs PRJEB26344 and PRJEB26345, respectively. The genomes of the isolates *Brevundimonas* sp. Bb-A, *Mycobacterium* sp. BbA or *Pimelobacter* sp. Bb-B are accessible under BioProjectIDs PRJNA528993, PRJEB28031 und PRJEB28032, respectively. The communities most abundant ten high-quality draft as well as isolate genomes are also accessible via EMGB platform ([https://emgb.cebitec.uni-bielefeld.de/Bbraunii-bacterial-consortium/).](https://emgb.cebitec.uni-bielefeld.de/Bbraunii-bacterial-consortium/)

Authors' contributions

O.B.K., V.K, and O.K. conceived this study. O.B.K., A.S., A.B., J.K. and O.K. supervised experiments and analyses. O.B.K., V.K., E.C. and S.C. accomplished all physiological studies. O.B.K., D.W., A.S., C.H., C.R., A.B., V.K., A.A.D., A.W., J.K., O.R., J.B. and A.G. performed the sequencing and the bioinformatic analyses. O.B.K., V.K., A.B. and O.K. wrote the original draft manuscript with contributions from all authors. O.B.K., V.K., A.B and O.K. reviewed and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We are thankful to Joao Gouveia and Douwe van der Veen (Wageningen University, The Netherlands) for providing *Botryococcus braunii* race A and B (CCALA778 and AC761) stock cultures. Furthermore, we are grateful to the Center for Biotechnology (CeBiTec) at Bielefeld University for access to the Technology Platforms.

Funding

This work was supported by the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement 311956 (relating to project "SPLASH – Sustainable PoLymers from Algae Sugars and Hydrocarbons"). The bioinformatics support by the BMBF-funded project "Bielefeld-Gießen Center for Microbial Bioinformatics" – BiGi (grant 031A533) within the German Network for Bioinformatics Infrastructure (de. NBI) is gratefully acknowledged.

Ethics approval and consent to participate

Not applicable.

Competing interests

The authors declare that they have no competing interests

References

Aaronson, S., Dhawale, S. W., Patni, N. J., Deangelis, B., Frank, O., and Baker, H.
(1977). The cell content and secretion of water-soluble vitamins by several freshwater
algae. *Arch. Microbiol.* **112**:57–59.

Amin, S. A., Hmelo, L. R., Van Tol, H. M., Durham, B. P., Carlson, L. T., Heal, K. R.,
Morales, R. L., Berthiaume, C. T., Parker, M. S., Djunaedi, B., et al. (2015).
Interaction and signalling between a cosmopolitan phytoplankton and associated
bacteria. *Nature* **522**:98–101.

Astals, S., Nolla-Ardèvol, V., and Mata-Alvarez, J. (2012). Anaerobic co-digestion of pig
manure and crude glycerol at mesophilic conditions: Biogas and digestate. *Bioresour.*
Technol. **110**:63–70.

Banerjee, A., Sharma, R., Chisti, Y., and Banerjee, U. C. (2002). Botryococcus braunii: A
renewable source of hydrocarbons and other chemicals. *Crit. Rev. Biotechnol.* **22**:245–
279.

Bisseling, T., Dangl, J. L., and Schulze-Lefert, P. (2009). Next-Generation Communication.
Science (80-.). **324**:691 LP – 691.

Blifernez-Klassen, O., Chaudhari, S., Klassen, V., Wördenweber, R., Steffens, T.,
Cholewa, D., Niehaus, K., Kalinowski, J., and Kruse, O. (2018). Metabolic survey of
Botryococcus braunii: Impact of the physiological state on product formation. *PLoS One*
13:e0198976.

Blin, K., Wolf, T., Chevrette, M. G., Lu, X., Schwalen, C. J., Kautsar, S. A.,
Suarez Duran, H. G., de los Santos, E. L. C., Kim, H. U., Nave, M., et al. (2017).
antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary
identification. *Nucleic Acids Res.* **45**:W36–W41.

Blom, J., Kreis, J., Spänig, S., Juhre, T., Bertelli, C., Ernst, C., and Goesmann, A.
(2016). EDGAR 2.0: an enhanced software platform for comparative gene content
analyses. *Nucleic Acids Res.* **44**:W22–W28.

Boisvert, S., Raymond, F., Godzaridis, E., Laviolette, F., and Corbeil, J. (2012). Ray
Meta: scalable de novo metagenome assembly and profiling. *Genome Biol.* **13**:R122.

Buchfink, B., Xie, C., and Huson, D. H. (2015). Fast and sensitive protein alignment using
DIAMOND. *Nat. Methods* **12**:59–60.

Chirac, C., Casadevall, E., Largeau, C., and Metzger, P. (1985). Bacterial Influence Upon

Growth and Hydrocarbon Production of the Green Alga Botryococcus Braunii. *J. Phycol.* **21**:380–387.

Cole, J. J. (1982). Interactions Between Bacteria and Algae in Aquatic Ecosystems. *Annu. Rev. Ecol. Syst.* **13**:291–314.

Cotter, P. D., Ross, R. P., and Hill, C. (2013). Bacteriocins-a viable alternative to antibiotics? *Nat. Rev. Microbiol.* **11**:95–105.

Croft, M. T., Lawrence, A. D., Raux-Deery, E., Warren, M. J., and Smith, A. G. (2005). Algae acquire vitamin B12 through a symbiotic relationship with bacteria. *Nature* **438**:90–93.

Croft, M. T., Warren, M. J., and Smith, A. G. (2006). Algae need their vitamins. *Eukaryot. Cell* **5**:1175–1183.

Durán, P., Thiergart, T., Garrido-Oter, R., Agler, M., Kemen, E., Schulze-Lefert, P., and Hacquard, S. (2018). Microbial Interkingdom Interactions in Roots Promote Arabidopsis Survival. *Cell* Advance Access published 2018, doi:10.1016/j.cell.2018.10.020.

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**:2460–2461.

Edgar, R. C. (2013). UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat. Methods* **10**:996–8.

Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., Heger, A., Hetherington, K., Holm, L., Mistry, J., et al. (2014). Pfam: The protein families database. *Nucleic Acids Res.* **42**.

Fuentes, J. L., Garbayo, I., Cuaresma, M., Montero, Z., González-Del-Valle, M., and Vílchez, C. (2016). Impact of microalgae-bacteria interactions on the production of algal biomass and associated compounds. *Mar. Drugs* **14**.

Gilbert, J. A., Quinn, R. A., Debelius, J., Xu, Z. Z., Morton, J., Garg, N., Jansson, J. K., Dorrestein, P. C., and Knight, R. (2016). Microbiome-wide association studies link dynamic microbial consortia to disease. *Nature* **535**:94.

Gkarmiri, K., Mahmood, S., Ekblad, A., Alström, S., Högberg, N., and Finlay, R. (2017). Identifying the Active Microbiome Associated with Roots and Rhizosphere Soil of Oilseed Rape. *Appl. Environ. Microbiol.* **83**:e01938-17.

Gómez-Consarnau, L., Sachdeva, R., Gifford, S. M., Cutter, L. S., Fuhrman, J. A., Sañudo-Wilhelmy, S. A., and Moran, M. A. (2018). Mosaic patterns of B-vitamin synthesis and utilization in a natural marine microbial community. *Environ. Microbiol.*

20:2809–2823.

Grant, M. A. A., Kazamia, E., Cicuta, P., and Smith, A. G. (2014). Direct exchange of

vitamin B12 is demonstrated by modelling the growth dynamics of

algal–bacterial cocultures. *ISME J.* **8**:1418–1427.

Grosberg, R. K., and Strathmann, R. R. (2007). The Evolution of Multicellularity: A

Minor Major Transition? *Annu. Rev. Ecol. Evol. Syst.* Advance Access published 2007,

doi:10.1146/annurev.ecolsys.36.102403.114735.

Higa, A., Miyamoto, E., Rahman, L. ur, and Kitamura, Y. (2008). Root tip-dependent,

active riboflavin secretion by Hyoscyamus albus hairy roots under iron deficiency. *Plant*

Physiol. Biochem. Advance Access published 2008, doi:10.1016/j.plaphy.2008.01.004.

Huson, D. H., Beier, S., Flade, I., Górska, A., El-Hadidi, M., Mitra, S., Ruscheweyh, H.

J., and Tappu, R. (2016). MEGAN Community Edition - Interactive Exploration and

Analysis of Large-Scale Microbiome Sequencing Data. *PLoS Comput. Biol.* **12**.

Hyatt, D., Chen, G. L., LoCascio, P. F., Land, M. L., Larimer, F. W., and Hauser, L. J.

(2010). Prodigal: Prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**.

Jett, B. D., Hatter, K. L., Huycke, M. M., and Gilmore, M. S. (1997). Simplified Agar

Plate Method for Quantifying Viable Bacteria. *Biotechniques* **23**:648–650.

Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y., and Morishima, K. (2017). KEGG:

New perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res.*

45:D353–D361.

Kang, D. D., Froula, J., Egan, R., and Wang, Z. (2015). MetaBAT, an efficient tool for

accurately reconstructing single genomes from complex microbial communities. *PeerJ*

3:e1165.

Klassen, V., Blifernez-Klassen, O., Wibberg, D., Winkler, A., Kalinowski, J., Posten, C.,

and Kruse, O. (2017). Highly efficient methane generation from untreated microalgae

biomass. *Biotechnol. Biofuels* **10**:186.

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., and Phillippy, A. M.

(2017). Canu: Scalable and accurate long-read assembly via adaptive κ-mer weighting

and repeat separation. *Genome Res.* Advance Access published 2017,

doi:10.1101/gr.215087.116.

Krohn-Molt, I., Alawi, M., Förstner, K. U., Wiegandt, A., Burkhardt, L., Indenbirken,

D., Thieß, M., Grundhoff, A., Kehr, J., Tholey, A., et al. (2017). Insights into

Microalga and bacteria interactions of selected phycosphere biofilms using

metagenomic, transcriptomic, and proteomic approaches. *Front. Microbiol.* **8**.

Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics
Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**:1870–1874.

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods **9**:357–359.

Ley, R. E., Hamady, M., Lozupone, C., Turnbaugh, P. J., Ramey, R. R., Bircher, J. S.,
Schlegel, M. L., Tucker, T. A., Schrenzel, M. D., Knight, R., et al. (2008). Evolution
of Mammals and Their Gut Microbes. *Science (80-.).* **320**:1647 LP – 1651.

Li, H., and Wren, J. (2014). Toward better understanding of artifacts in variant calling from
high-coverage samples. *Bioinformatics* Advance Access published 2014,
doi:10.1093/bioinformatics/btu356.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and
SAMtools. *Bioinformatics* **25**:2078–2079.

Liebe, S., Wibberg, D., Winkler, A., Puehler, A., Schlueter, A., and Varrelmann, M.
(2016). Taxonomic analysis of the microbial community in stored sugar beets using
high-throughput sequencing of different marker genes. *FEMS Microbiol. Ecol.* **92**:1–12.

Magoč, T., and Salzberg, S. L. (2011). FLASH: Fast length adjustment of short reads to
improve genome assemblies. *Bioinformatics* **27**:2957–2963.

Marschner, P. (2011). *Marschner's Mineral Nutrition of Higher Plants: Third Edition*.

Maus, I., Kim, Y. S., Wibberg, D., Stolze, Y., Off, S., Antonczyk, S., Pühler, A., Scherer,
P., and Schlüter, A. (2017). Biphasic Study to Characterize Agricultural Biogas Plants
by High-Throughput 16S rRNA Gene Amplicon Sequencing and Microscopic Analysis.
J. Microbiol. Biotechnol. **27**.

McKirdy, D. M., Cox, R. E., Volkman, J. K., and Howell, V. J. (1986). Botryococcane in a
new class of Australian non-marine crude oils. *Nature* **320**:57–59.

Meyer, F., Goesmann, A., McHardy, A. C., Bartels, D., Bekel, T., Clausen, J.,
Kalinowski, J., Linke, B., Rupp, O., Giegerich, R., et al. (2003). GenDB - An open
source genome annotation system for prokaryote genomes. *Nucleic Acids Res.* **31**:2187–
2195.

Molnár, I., Lopez, D., Wisecaver, J. H., Devarenne, T. P., Weiss, T. L., Pellegrini, M.,
and Hackett, J. D. (2012). Bio-crude transcriptomics: gene discovery and metabolic
network reconstruction for the biosynthesis of the terpenome of the hydrocarbon oil-
producing green alga, Botryococcus braunii race B (Showa). *BMC Genomics* **13**:576.

Nguyen, C. (2003). Rhizodeposition of organic C by plants: mechanisms and controls
Advance Access published 2003.

O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R.,
Rajput, B., Robbertse, B., Smith-White, B., Ako-Adjei, D., et al. (2016). Reference
sequence (RefSeq) database at NCBI: Current status, taxonomic expansion, and
functional annotation. *Nucleic Acids Res.* **44**:D733–D745.

Pančić, M., and Kiørboe, T. (2018). Phytoplankton defence mechanisms: traits and trade-
offs. *Biol. Rev.* **93**:1269–1303.

Price, M. N., Dehal, P. S., and Arkin, A. P. (2010). FastTree 2 - Approximately maximum-
likelihood trees for large alignments. *PLoS One* **5**.

Ramanan, R., Kim, B.-H., Cho, D.-H., Oh, H.-M., and Kim, H.-S. (2016). Algae–bacteria
interactions: Evolution, ecology and emerging applications. *Biotechnol. Adv.* **34**:14–29.

Reed, H., and Martiny, J. (2007). Testing the functional significance of microbial
composition in natural communities. *FEMS Microbiol. Ecol.* **62**:161–170.

Rivett, D. W., and Bell, T. (2018). Abundance determines the functional role of bacterial
phylotypes in complex communities. *Nat. Microbiol.* **3**:767–772.

Rojo, F. (2010). Enzymes for Aerobic Degradation of Alkanes. In *Handbook of Hydrocarbon*
and Lipid Microbiology (ed. Timmis, K. N.), pp. 781–797. Berlin, Heidelberg: Springer
Berlin Heidelberg.

Sambles, C., Moore, K., Lux, T. M., Jones, K., Littlejohn, G. R., Gouveia, J. D., Aves, S.
J., Studholme, D. J., Lee, R., and Love, J. (2017). Metagenomic analysis of the
complex microbial consortium associated with cultures of the oil-rich alga Botryococcus
braunii. *Microbiologyopen* **6**.

Sañudo-Wilhelmy, S. A., Gómez-Consarnau, L., Suffridge, C., and Webb, E. A. (2014).
The Role of B Vitamins in Marine Biogeochemistry. *Ann. Rev. Mar. Sci.* **6**:339–367.

Schlegel, H. G., Kaltwasser, H., and Gottschalk, G. (1961). Ein Submersverfahren zur
Kultur wasserstoffoxydierender Bakterien: Wachstumsphysiologische Untersuchungen.
Arch. Mikrobiol. **38**:209–222.

Schreiter, S., Ding, G.-C., Heuer, H., Neumann, G., Sandmann, M., Grosch, R., Kropf,
S., and Smalla, K. (2014). Effect of the soil type on the microbiome in the rhizosphere
of field-grown lettuce. *Front. Microbiol.* **5**:144.

Schwarzhans, J. P., Cholewa, D., Grimm, P., Beshay, U., Risse, J. M., Friehs, K., and
Flaschel, E. (2014). Dependency of the fatty acid composition of Euglena gracilis on
growth phase and culture conditions. *J. Appl. Phycol.* **27**:1389–1399.

Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**:2068–
2069.

Seymour, J. R., Amin, S. A., Raina, J. B., and Stocker, R. (2017). Zooming in on the
phycosphere: The ecological interface for phytoplankton-bacteria relationships. *Nat.*
Microbiol. **2**.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., and Zdobnov, E. M.
(2015). BUSCO: Assessing genome assembly and annotation completeness with single-
copy orthologs. *Bioinformatics* **31**:3210–3212.

Takahashi, S., Tomita, J., Nishioka, K., Hisada, T., and Nishijima, M. (2014).
Development of a prokaryotic universal primer for simultaneous analysis of bacteria and
archaea using next-generation sequencing. *PLoS One* **9**:e105592.

Tanabe, Y., Okazaki, Y., Yoshida, M., Matsuura, H., Kai, A., Shiratori, T., Ishida, K. I.,
Nakano, S. I., and Watanabe, M. M. (2015). A novel alphaproteobacterial
ectosymbiont promotes the growth of the hydrocarbon-rich green alga Botryococcus
braunii. *Sci. Rep.* **5**.

Tang, Y. Z., Koch, F., and Gobler, C. J. (2010). Most harmful algal bloom species are
vitamin B1 and B12 auxotrophs. *Proc. Natl. Acad. Sci.* **107**:20756–20761.

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.
A., Zeng, Q., Wortman, J., Young, S. K., et al. (2014). Pilon: An integrated tool for
comprehensive microbial variant detection and genome assembly improvement. *PLoS*
One Advance Access published 2014, doi:10.1371/journal.pone.0112963.

Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R. (2007). Naive Bayesian classifier
for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ.*
Microbiol. **73**:5261–5267.

Wang, X., Li, Z., Su, J., Tian, Y., Ning, X., Hong, H., and Zheng, T. (2010). Lysis of a
red-tide causing alga, Alexandrium tamarense, caused by bacteria from its phycosphere.
Biol. Control **52**:123–130.

Waterhouse, R. M., Seppey, M., Simao, F. A., Manni, M., Ioannidis, P., Klioutchnikov,
G., Kriventseva, E. V., and Zdobnov, E. M. (2018). BUSCO applications from quality
assessments to gene prediction and phylogenomics. *Mol. Biol. Evol.* **35**:543–548.

Wibberg, D., Blom, J., Jaenicke, S., Kollin, F., Rupp, O., Scharf, B., Schneiker-Bekel, S.,
Sczcepanowski, R., Goesmann, A., Setubal, J. C., et al. (2011). Complete genome
sequencing of Agrobacterium sp. H13-3, the former Rhizobium lupini H13-3, reveals a
tripartite genome consisting of a circular and a linear chromosome and an accessory

plasmid but lacking a tumor-inducing Ti-plasmid. *J. Biotechnol.* **155**:50–62.

Wibberg, D., Andersson, L., Rupp, O., Goesmann, A., Pühler, A., Varrelmann, M.,
Dixelius, C., and Schlüter, A. (2016). Genome analysis of the sugar beet pathogen
Rhizoctonia solani AG2-2IIIB revealed high numbers in secreted proteins and cell wall
degrading enzymes. *BMC Genomics* Advance Access published 2016,
doi:10.1186/s12864-016-2561-1.

Xu, J., Zhang, Y., Zhang, P., Trivedi, P., Riera, N., Wang, Y., Liu, X., Fan, G., Tang, J.,
Coletta-Filho, H. D., et al. (2018). The structure and function of the global citrus
rhizosphere microbiome. *Nat. Commun.* **9**:4894.

Zengler, K., and Zaramela, L. S. (2018). The social network of microorganisms - How
auxotrophies shape complex communities. *Nat. Rev. Microbiol.* Advance Access
published 2018, doi:10.1038/s41579-018-0004-5.

Zhang, H., Yohe, T., Huang, L., Entwistle, S., Wu, P., Yang, Z., Busk, P. K., Xu, Y., and
Yin, Y. (2018). dbCAN2: a meta server for automated carbohydrate-active enzyme
annotation. *Nucleic Acids Res.* Advance Access published 2018,
doi:10.1093/nar/gky418.

Zhou, J., Bruns, M. A., and Tiedje, J. M. (1996). DNA recovery from soils of diverse
composition. *Appl. Environ. Microbiol.* **62**:316–322.

Additional files

Supplementary Materials.

Figure S1: Characteristics and taxonomic profiling of xenic *Botryococcus braunii* race A and
B cultures. Illustrated are the typical colony-forming cells of *B. braunii* races A and B with a
complex structure size of approximately >100µm in diameter. The table shows the mean
values (three biological and technical replicates, SE; n=9) of algal content of biomass,
hydrocarbons and carbohydrates at the sampling time points (linear and stationary growth
phases). DW, culture dry weight.

Figure S2: Taxonomic profiling of *Botryococcus braunii* associating community. Relative
abundance of detected taxa (phylum level) based on (**a**) high-throughput 16S rRNA gene
amplicons and (**b**) metagenomic reads. The evaluation of 16S rDNA amplicons are detailed in
Table S3. Metagenome datasets were taxonomically assigned via MEGAN (200,000
randomly subsampled reads, minimum 50 reads per taxonomic group). The relatively high

proportion of unassigned reads likely results from the fact that the algal and some bacterial

genomes are not disposed in the used NCBI-NR database. The analysis revealed the

occurrence of the phyla Euryarchaeota and Viruses belonging to the family of *Microviridae*

(bacteriophages with a single-stranded DNA genome).

Figure S3: Microbiome accompanying the *Botryococcus braunii* consortia. Illustration of the

relative abundance and phylogenetic distribution of bacterial genera detected via high-

throughput 16S rDNA amplicon sequencing approach (for details see Table S3).

Figure S4: Phylogenetic classification of selected *Botryococcus braunii* consortia members.

Illustrated is the phylogenetic characterization of the high-quality draft and complete genomes

of *B. braunii* accompanying bacterial community. Maximum-likelihood-based phylogenetic

tree built out of a core of 53 conserved marker genes (Table S4) per analyzed genome (bold).

Black, grey and white circles represent the nodes with calculated bootstrap values of 100, 99

and <75%, respectively (1,000 replicates).

Figure S5: Comparison and taxonomic assignment of different *Botryococcus braunii*

metagenomes. Datasets obtained during the present study (races A and B, T1 and T2 (linear

and stationary growth phases, respectively) were compared to the datasets of Guadeloupe race

B strain (Condition A-C supplemented with citric acid as organic carbon source and vitamins

(B_1, B_7 and B_{12}), with A: initial consortium, B: washed culture, C: antibiotics-treated

(ciprofloxacin)) (Sambles et al. 2017). For each sample, 200,000 metagenomic reads were

randomly subsampled and subjected to the taxonomic assignment using DIAMOND and

MEGAN (minimum 50 reads per taxonomic group). The circle size represents the amount of

classified reads for the respective taxonomic group.

Figure S6: Hydrocarbon accumulation profiles of axenic and xenic *B. braunii* cultures. Shown

is the hydrocarbon content observed in the samples of (**a**) axenic *B. braunii* (green) as well as

in presence of (**b**) *Mycobacterium* sp. Bb-A (orange) and (**c**) *Brevundimonas* sp. Bb-A (blue)

during the course of cultivation. The data serves as additional information for Figure 2.

Table S1: Features of the metagenome assembled genomes (MAGs). Assembly and

annotation characteristics (*e.g.* Total size, N50, etc.) are listed according to the respective

MAG. The completeness as well as the contamination degree of the MAGs was assessed via

BUSCO (v3.0.). P- and E-Modi refer to prokaryotic MAGs and eukaryotic genome fragments,

respectively. The high-quality MAGs were classified via single-copy marker genes, *e.g. rpoB*,

rpoD etc., detected by BUSCO. The metagenomic datasets of *B. braunii* race A and B were

mapped against the MAGs, and represented as percentage of mapped reads.

Table S2: Characteristics of high-quality draft and complete genomes. Summary of the

taxonomy distribution and annotation features of ten high-quality MAGs and two complete

genomes (completeness >80%, contamination <10%, Table S2) obtained from the *B. braunii*

race A and B consortia. Shown are similarity mean values based on AAI (genome to genome

comparison). Genomes with similarity values >95% are regarded as the same species as the

reference. MAG, metagenome assembled genome; AAI, average amino acid identities; CDS,

Coding DNA sequences.

Supplementary Discussion

Chapter 1 | Metagenomic survey of the *Botryococcus braunii* consortia

Chapter 2 | Elucidation of the genetic portfolio of the bacterial community

Supplementary Tables. (XLSX**,** 330 KB)

Table S3: Statistics and taxonomic classification of 16S rRNA gene amplicon sequencing of

the *B. braunii* consortia. Datasets include the analysis of the high-throughput 16S rDNA

amplicon sequencing results and are supplemental information for Figures S2a and S3.

Abbreviations: OTU, operational taxonomic unit; T, time point.

Table S4: List of conserved core genes. Dataset serves as supplemental information for Figure

S4 and includes conserved marker (core) genes shared among the 31 genomes, used for the

construction of the Maximum-likelihood-based (ML) phylogenetic tree.

Table S5: List of all detected functional pathways encoded in the high-quality draft and

complete bacterial genomes. Datasets include KEGG-based functionally assigned pathways of

the annotated high-quality MAGs and complete genomes imported to the EMGB platform.

(elastic metagenome browser, https://emgb.cebitec.uni-bielefeld.de/Bbraunii-bacterial-

consortium/). Displayed are the total number of the enzymes and the corresponding EC

numbers encoded within the particular genome for the respective pathway. Bold marked

pathway maps have been selected for Figure 1b.

Table S6: List of selected manually inspected genes within the high-quality draft and
complete bacterial genomes. Datasets include manual BLASTp screening results (E-value
threshold 1×10^{-10}) of selected genes.

Table S7: List of Carbohydrate-active enzymes (CAZy) encoded by the *B. braunii* bacterial
community members. Datasets include Carbohydrate-active enzymes (CAZy) encoded within
the high-quality draft and complete bacterial genomes and serve as additional information for
Figure 1b. The gene search within the high-quality draft and complete genomes was
accomplished via dbCAN2 metaserver (Zhang et al. 2018).

Table S8: List of all detected B-vitamin biosynthesis genes encoded by the *B. braunii*
bacterial community members. Datasets include the function description (EC number) as well
as the corresponding genes of the key enzymes involved in the de novo biosynthesis of B-
vitamins and serve as supplemental information for Figure 3a. The genes involved in the
particular B-Vitamin pathway were searched (based on EMGB annotation system and manual
BLASTp search) within the high-quality draft and complete bacterial genomes. √*, alternative
gene identified via manual BLASTp search. Abbreviations: *M*. Bb-A, *Mycobacterium* sp. Bb-
A; *B*. Bb-A, *Brevundimonas* sp. Bb-A; *P*.Bb-B, *Pimelobacter* sp. Bb-B.

Table S9: List of all detected secondary metabolite biosynthetic gene clusters. The datasets
include the identified secondary metabolite biosynthetic gene clusters encoded by the high-
quality draft and complete genomes of the *B. braunii* associates and serve as additional
information for Figure 1b. The analysis was accomplished using the antibiotics & Secondary
Metabolite Analysis Shell (antiSMASH, Version 4.1.0, Blin et al. 2017).