1 **Genetic dissection of a *Leishmania* flagellar proteome**

2 **demonstrates requirement for directional motility in sand fly**

3 **infections**

5 Running Head: *L. mexicana* flagellar mutant screen

6 Authors

7 Tom Beneke[1], François Demay[2], Edward Hookway[3], Nicole Ashman[1], Heather Jeffery[1],

8 James Smith[1], Jessica Valli[1], Tomas Becvar[4], Jitka Myskova[4], Tereza Lestinova[4], Shahaan

9 Shafiq[1,5], Jovana Sadlova[4], Petr Volf[4], Richard Wheeler[6,1] and Eva Gluenz[1*]

10 Affiliations

[1]Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford, OX1

3RE, UK

[2] University of Lille 1, Cité Scientifique, 59655 Villeneuve d'Ascq, France

[3] Research Department of Pathology, University College London, 72 Huntley Street, London,

WC1E 6JD

[4] Department of Parasitology, Faculty of Science, Charles University, Vinicna 7, Prague 2,

128 44, Czech Republic

[5] School of Life Sciences, Oxford Brookes University, Gipsy Lane, Oxford, OX3 0BP, UK

[6] Peter Medawar Building for Pathogen Research, Nuffield Department of Medicine,

University of Oxford, South Parks Road, Oxford OX1 3SY, UK

*Corresponding author: eva.gluenz@path.ox.ac.uk

23 Abstract

The protozoan parasite *Leishmania* possesses a single flagellum, which is remodelled during

the parasite's life cycle from a long motile flagellum in promastigote forms in the sand fly to a

short immotile flagellum in amastigotes residing in mammalian phagocytes. This study

examined the protein composition and *in vivo* function of the promastigote flagellum. Protein

mass spectrometry and label free protein enrichment testing of isolated flagella and

deflagellated cell bodies defined a flagellar proteome for *L. mexicana* promastigote forms

(available via ProteomeXchange with identifier PXD011057). This information was used to generate a CRISPR-Cas9 knockout library of 100 mutants to screen for flagellar defects. This first large-scale knockout screen in a *Leishmania* sp. identified 56 mutants with altered swimming speed (52 reduced and 4 increased) and defined distinct mutant categories (faster swimmers, slower swimmers, slow uncoordinated swimmers and paralysed cells, including aflagellate promastigotes and cells with curled flagella and disruptions of the paraflagellar rod). Each mutant was tagged with a unique 17-nt barcode, providing a simple barcode sequencing (bar-seq) method for measuring the relative fitness of *L. mexicana* mutants *in vivo*. In mixed infections of the permissive sand fly vector *Lutzomyia longipalpis*, paralysed promastigotes and uncoordinated swimmers were severely diminished in the fly after defecation of the bloodmeal. Subsequent examination of flies infected with a single mutant lacking the central pair protein PF16 showed that these paralysed promastigotes did not reach anterior regions of the fly alimentary tract. These data show that *L. mexicana* need directional motility for successful colonisation of sand flies.

Author Summary

Leishmania are protozoan parasites, transmitted between mammals by the bite of phlebotomine sand flies. Promastigote forms in the sand fly have a long flagellum, which is motile and used for anchoring the parasites to prevent clearance with the digested blood meal remnants. To dissect flagellar functions and their importance in life cycle progression, we generated here a comprehensive list of >300 flagellar proteins and produced a CRISPR-Cas9 gene knockout library of 100 mutant *Leishmania*. We studied their behaviour *in vitro* before examining their fate in the sand fly *Lutzomyia longipalpis*. Measuring mutant swimming speeds showed that about half behaved differently compared to the wild type*:* a few swam faster, many slower and some were completely paralysed. We also found a group of uncoordinated swimmers. To test whether flagellar motility is required for parasite migration from the fly midgut to the foregut from where they reach the next host, we infected sand flies with a mixed mutant population. Each mutant carried a unique tag and tracking these tags up to nine days after infection showed that paralysed and uncoordinated *Leishmania* were rapidly lost from flies. These data indicate that directional swimming is important for successful colonisation of sand flies.

Introduction

Eukaryotic flagella / cilia are complex multifunctional organelles conserved from protists to humans [1]. Protists use flagella for swimming, feeding, cell-to-cell communication, adherence to substrates and morphogenesis [2]. Single-celled organisms, most commonly among them the green algae *Chlamydomonas reinhardtii,* have served as important model organisms to study molecular mechanisms of ciliogenesis and ciliary function [3], spurred on by the recognition that ciliary defects cause human genetic disorders collectively termed "ciliopathies" [4]. The eukaryotic flagellum is a complex, highly structured organelle and dissection of the molecular mechanisms underpinning its diverse functions requires detailed knowledge of its component parts. Proteomic studies of isolated flagella or axonemes from diverse species typically identified at least 300 distinct proteins [5-9] and phylogenetic profiling identified a set of 274 evolutionarily conserved ciliary genes [10]. All of these datasets comprise many "hypothetical" proteins still awaiting functional characterisation in addition to well-characterised core components of the microtubule axoneme, associated motor proteins and regulatory complexes.

Insights into conserved ciliary biology have helped elucidation of flagellar function in eukaryotic microbes, with a particular focus on human pathogens [11,12]. Among these, flagella have been most extensively studied in the causative agent of African trypanosomiasis, *Trypanosoma brucei* [13], which uses flagellar motility for locomotion and immune evasion [14] and exhibits close spatio-temporal coordination between flagellum assembly and cell morphogenesis during division [15]. The *T. brucei* bloodstream form is particularly sensitive to the loss of flagellar function [6,16], highlighting a potential Achilles' heel that might be exploitable for new anti-parasitic treatments.

The *Leishmania* flagellum is also a multi-functional organelle, which undergoes striking structural changes during the parasite's life cycle [17-19]. Amastigote forms proliferating in mammalian macrophages possess a short sensory-type 9+0 microtubule axoneme, which is remodelled to a canonical long motile 9+2 axoneme during differentiation to promastigote forms, which live in blood-feeding phlebotomine sand flies (Diptera: Psychodidae). In the fly, nectomonad promastigote forms attach via their flagella to the microvilli of the posterior midgut [20] to protect the parasites from being cleared during defecation of remnants of the blood meal. In the oesophageal valve, broad haptomonad forms attach to the cuticular lining via their flagellar tips, forming hemidesmosomes [20]. These life cycle descriptions [21] imply that periods of attachment must be followed by migration to more anterior regions of the alimentary tract and the propulsive function of the *Leishmania* flagellum is presumed to drive this forward migration but this has not been directly tested.

To enable a detailed genetic dissection of flagellar functions and mechanisms in *Leishmania*,
we defined here a flagellar proteome for motile *L. mexicana* promastigotes. We used new
CRISPR-Cas9 genome editing methods [22] to generate a *Leishmania* knockout library of
100 mutants, over half of which showed altered swimming speed. We also developed a
barcode sequencing (bar-seq) protocol to test the fitness of mutants in the permissive sand
fly vector *Lutzomyia longipalpis*. This study identified new genes required for flagellar motility
and shows that whilst culture-form promastigotes tolerated loss of the flagellum, paralysed
mutants and uncoordinated swimmers failed to colonise sand flies indicating that directional
flagellar motility is required for completion of the parasite's life cycle.

Results

Defining the promastigote flagellar proteome

To enable a systematic genetic dissection of flagellar functions we sought to isolate *L.*
mexicana promastigote flagella comprising the axoneme, extra-axonemal structures and the
surrounding membrane for subsequent analysis by protein mass spectrometry (MS).
Mechanical shearing in the presence of 75 mM Ca^{2+} successfully separated cells into flagella
(F) and deflagellated cell bodies (CB) (Figure 1A, B). Subsequent centrifugation on sucrose
gradients allowed isolation of F and CB fractions with little cross-contamination: the CB
fraction contained only 2.03% (±0.69%) isolated flagella and the F fractions contained 0.56%
(±0.15%) deflagellated cell bodies (S1 Figure). Isolated flagella still retained their membrane:
First, examination of F fractions by transmission electron microscopy (TEM) confirmed that
most axonemes were bounded by a membrane (S2 Figure) and second, tracking an
abundant promastigote flagellar membrane protein, the small myristoylated protein 1 (SMP-1,
[23]) tagged with enhanced green fluorescent protein (eGFP) showed that it remained
associated with isolated flagella (Figure 1). Analysis of the SMP-1::eGFP signal also
facilitated flagellar length measurements in whole cells, F and CB fractions, which showed
that flagella were separated from the cell body near the exit point from the flagellar pocket.
The average break point was 2.7 μm distal to the base of the flagellum. The length of the
isolated flagella was similar to those on intact cells, indicating that isolated flagella remained
in one piece, with little fragmentation (S3 Figure). Two independently prepared sets of F and
CB fractions were separated into detergent soluble (s) and insoluble fractions (i), yielding four
fractions, F_S, F_I, CB_S and CB_I (Figure 1C). All four fractions for both replicates were analysed
by liquid chromatography tandem mass spectrometry (MS), which detected a total of 2711
distinct proteins (Figure 1D). Enrichment of detected proteins between biological replicates

correlated well (Pearson's r > 0.72, Spearman's r_s > 0.83, S4 Figure). To discover proteins

enriched in each of the four fractions, we used a label-free normalized spectral index

quantitation method (SINQ, [24]; S1, S2 and S3 Table) to generate a SINQ enrichment plot

(Figure 2A). The promastigote flagellar proteome, defined as proteins enriched in F vs. CB

fractions consisted of 701 unique proteins detected in at least one MS run; 352 of these were

enriched in F vs. CB fractions in both MS runs.

Comparison with existing datasets validates flagellar proteome

To validate the data, we mapped well-characterised flagellar proteins onto the enrichment

data plot (Figure 2A). Axonemal, paraflagellar rod (PFR), flagellar tip and flagellar membrane

proteins mapped to the F_I and F_S quadrants. Basal body, FAZ and tripartite attachment

complex (TAC) proteins mapped to the CB_I and CB_S quadrants because F fractions

contained exclusively the cell-external portion of the flagellum. Intraflagellar transport (IFT)

proteins clustered around the midpoint of the plot, indicating their abundance was similar in

the F and CB fractions, which is consistent with their known dynamic association with the

flagellar basal body and axoneme.

We also found substantial overlaps between *L. mexicana* proteins in the F_I quadrant and

proteins detected in previously published flagellar proteomes of *L. donovani* and *T. brucei*

(S5 Figure A-D). However, *L. mexicana* proteins in the F_S quadrant showed only a moderate

overlap with reported soluble *T. brucei* flagellar proteins (S5 Figure C). We designed a

website (www.leishgedit.net/leishgedit_db) for interactive browsing of proteins in the

enrichment plots shown in Figures 2 and S5.

Proteins with predicted trans-membrane domains (TMD) were predominantly detected in the

detergent soluble fractions (S5 Figure F). Overall, TMD proteins were underrepresented in

the proteome compared to their frequency in the genome (Chi-squared test, p < 0.0001), as

were proteins smaller than 10 kDa (S6 Figure), most likely due to well-known technical

limitations of protein MS [25]. Although ribosomal proteins were detected in individual F

fractions, the enrichment plot clustered them around the midpoint, with many enriched in the

cell body fractions (S5 Figure E). Our simple strategy of testing for enrichment thus

successfully filtered out likely contaminating proteins from the promastigote flagellar

proteome, as recently observed for enrichment of other cytoskeleton structures in *T. brucei*

[26,27].

Interestingly, a comparison of these proteomics data with *L. mexicana* RNA-seq data from

promastigotes and amastigotes [28] showed that proteins enriched in the flagellar fractions

were significantly more likely to have higher RNA abundance in promastigotes vs.

amastigotes, compared to proteins detected in the cell body fraction (Figure 2B; Chi-squared

test, p < 0.0001). This is consistent with the disassembly of the motile axoneme during
differentiation from promastigotes to amastigotes [17]. Whilst on a global scale transcript
levels correlate poorly with protein abundance in *Leishmania* spp. [29] these data indicate
that modulation of mRNA levels is a key regulatory step in *Leishmania* flagellar biogenesis
and differentiation from a 9+2 to a 9+0 flagellum.

Selection of candidates for a motility mutant screen

Many of the proteins detected in the F fractions had orthologs in previously defined flagellar
and ciliary proteomes yet lacked any functional characterisation. Arguably, endowing cells
with motility is the primary function of the promastigote flagellum and we took advantage of
our high-throughput CRISPR-Cas9 toolkit [22] to identify proteins required for motility and
subsequently study the phenotypes of the mutant *Leishmania*. In our knockout (KO) library
(S4 Table) we included 19 highly conserved axonemal proteins involved in the regulation of
flagellar beating, three intraflagellar transport (IFT) proteins, 60 flagellar proteins with
transcript enrichment in promastigotes [28] and eight additional soluble and four insoluble
flagellar proteins. Twenty of the selected proteins were detected in the promastigote flagellar
proteome but have to our knowledge not been linked to flagella before. We also made
deletion mutants for two genes implicated in membrane protein trafficking, *BBS2* and
Kharon1. Flagellar localisation of a subset of proteins was independently examined by
generating cell lines expressing proteins tagged with a fluorescent protein at the N- and/or C-
terminus (S7 Figure).
Orthofinder [30] was used to generate genome-wide orthologous protein sequence families
using genome sequences of 33 ciliated and 15 non-ciliated species from across eukaryotic
life, including *L. mexicana* and *T. brucei* (S5 Table). Twenty-two proteins were kinetoplastid-
specific (*L. mexicana* and *T. brucei*), 30 were conserved specifically in ciliated organisms and
23 widely conserved across eukaryotes whilst the remainder showed no clear pattern. In the
following, we refer to genes of unknown function by their GeneID from TriTrypDB.org [31]
and where we identified named orthologs we used these gene names.

Screening CRISPR-Cas9 knockout mutants for motility defects

The target genes were then deleted as described previously [22]. To facilitate high-
throughput generation of knockout (KO) cell lines, PCR reactions and transfections were
performed in 96-well plates. Analysis of drug-resistant transfectants by PCR confirmed loss
of the target ORF and integration of the drug-resistance gene in 94 of 98 cell lines (S8
Figure). This 96% success rate highlights the power of our gene deletion strategy. The
reason for the presence of the target ORF in the remaining four cell lines was not further

investigated, but was confirmed by diagnostic PCR of two independently isolated samples of

genomic DNA from the relevant mutants.

The flagellar mutants generated in this study, the previously generated paralysed cell line

ΔPF16 [22], the parental line *L. mex* Cas9 T7, and wild type promastigotes were subjected

to motility assays using dark field microscopy to track the swimming behaviour of cells and

measure swimming speed and directionality as previously described [32]. Parental cells

immobilised though formaldehyde fixation were also measured. More than half of all mutant

lines showed a significant deviation from the normal average swimming speed measured for

the parental cell line and wild type controls (Figure 3A, B): 52 (53.6%) mutants showed a

significant reduction in speed and 4 (4.1%) swam faster (Student's t-test, $p < 0.005$; Figure 3A,

S4 Table). Plotting mean swimming speed against mean directionality (velocity/speed)

shows broad groups of mutants (Figure 3B): Those which are paralysed, slower swimmers,

slow uncoordinated swimmers, faster swimmers and a single mutant that had faster and

more directional swimming (Δ*LmxM.36.3620*). The mechanistic contribution to swimming

behaviour remains to be clarified for many proteins in this set. Loss of flagellar waveform

modulators would cause altered motility patterns, and this is exemplified by two mutants in

this set: the Δ*dDC2* mutant, which lacks the outer dynein arm docking complex protein dDC2

and can perform a ciliary beat but no flagellar beat [33] clusters with the uncoordinated

group. By contrast, Δ*LC4-like*, which lacks a distal regulator of outer dynein arms and spends

more time doing a flagellar beat at a higher beat frequency [33], was among the faster

swimmers.

The most severe loss of motility was observed in three cell lines that had no visible external

flagellum (Figure 4); all of these were deletions of conserved intraflagellar transport (IFT)

proteins (Δ*IFT122B*, Δ*IFT139* and Δ*IFT88*). Ablation of the central pair (CP) protein hydin

also resulted in almost complete paralysis, comparable to the deletion of the CP protein

PF16 [22].

In a subset of paralysed or slow-swimming uncoordinated mutants (Figure 3C) we noted that

the flagella tended to be in a curled rather than straight conformation. Δ*hydin* mutants had

the highest proportion of curled-up flagella (62.6%, Figure 4 and S9 Figure) while fewer than

1% of flagella were curled-up in the parental cell line and many other slow swimming mutants

(S9 Figure). A high proportion (>10%) of curled-up flagella was also found in four paralysed

KO lines (inner dynein arm intermediate chain protein mutant Δ*IC140*, 57%; Δ*PF16*, 14%;

tether and tether head complex protein mutants Δ*CFAP44*, 15% and Δ*CFAP43*, 19%) and

three uncoordinated KO lines (Δ*MBO2*, 26%; nexin-dynein regulatory complex protein mutant

Δ*DRC4*, 13%; Δ*LmxM.33.0560*, 12%). The curls were observed in aldehyde fixed cells as

well as in live cells in culture, indicating they were not an artefact of microscopy sample preparation. This novel phenotype might be caused by disrupted dynein regulation and warrants further investigation.

Null mutants for the major PFR protein PFR2, lacking the paracrystalline PFR lattice structure, are known to have impaired motility [34]. To compare motility of a Δ*PFR2* mutant with other mutants generated in this study, we used CRISPR-Cas9 to delete both allelic copies of the *PFR2* array (*PFR2A*, *PFR2B* and *PFR2C*) and confirmed loss of PFR2 expression by Western blot (S10 Figure). This Δ*PFR2* line had slower and less directional swimming compared to the parental cells, clustering with other slow swimming mutants defined in Figure 3B. To test whether gene deletion in other slow swimming mutants had a major disruptive effect on the PFR, which might explain their motility defect, we expressed PFR2::mNG in KO lines and looked for changes to PFR length or loss of PFR integrity (defined as gaps in the PFR2::mNG signal) (Figures 4 and 5). Three mutants had shorter flagella compared to the parental cell line, but the PFR remained proportional to the overall flagellar length and was uninterrupted (Δ*ARL-3A*, Δ*CFAP44*, and Δ*FLAM2*). Six mutants had PFR-specific defects (Figure 5B): a shorter flagellum with a disproportionately shorter PFR (Δ*LmxM.27.0860;* Δ*TTC29;* Δ*LmxM.14.1220*), a normal-length flagellum with a shorter PFR (Δ*FM458*) or a shorter PFR with gaps (Δ*LmxM.21.1110,* 25.3% of all flagella; Δ*MBO2* only 4.1% of all flagella). Interestingly, these comparatively subtle alterations to PFR length and integrity reduced swimming speed to similar levels as PFR2 deletion (Figure 3C).

We generated 13 add-back cell lines to rescue mutant phenotypes by transfecting episomes containing the deleted ORF. Four complemented mutants fully recovered parental swimming speed (Figure 3) and complemented Δ*CFAP44* and Δ*MBO2* lines showed fewer curled flagella (S9 Figure). Complementation of the other slow swimming mutants resulted in a significant increase in swimming speed close to parental levels (Figure 3) and reduction of curling compared to the KO lines (S9 Figure).

Thus, our screen readily identified promastigote mutants with impaired motility and even the most severe phenotype, ablation of flagellar assembly caused by loss of IFT components, was compatible with promastigote survival *in vitro*, in line with earlier reports [35], [36], [37].

Paralysed promastigotes are cleared from sand flies

Whilst flagellar motility is generally believed to be required for development in sand flies, enabling *Leishmania* migration from the midgut to the mouthparts [38-40], this has not been directly tested. To interrogate the phenotypes of larger cohorts of *Leishmania* mutants in parallel, we developed a multiplexed bar-seq strategy inspired by pioneering phenotyping

screens in yeast [41] and the malaria parasite *Plasmodium berghei* [42,43]. We pooled mutant *L. mexicana* lines that were each tagged with a unique 17 bp barcode. This enabled us to measure the relative abundance of each line at different time points after sand fly infection (S11 Figure). Seventeen were flagellar mutants described above and five were parental control cell lines tagged with unique barcodes in their small subunit (SSU) ribosomal RNA locus. We also generated a barcoded Δ*LPG1* KO mutant, which is only defective in LPG synthesis [22,44] and three barcoded mutants defective in the pathway leading to mannose activation for synthesis of LPG and other glycoconjugates: KOs of phosphomannose isomerase [45] (Δ*PMI*), phosphomannomutase [46] (ΔPMM) and GDP-mannose pyrophosphorylase [47] (ΔGDP-MP). These mutants were included as control lines expected to be outcompeted by the parental cell lines. The barcoded cell lines were pooled in equal proportions and used to infect *L. longipalpis*. The relative abundance of each line was determined by sequencing DNA isolated from the mixed promastigote pool and from flies at two, six and nine days after infection. The results show progressively diminishing proportions for the control mutants defective in LPG synthesis (Δ*LPG1*) or a broader range of glycoconjugates including LPG (Δ*PMI*, Δ*PMM* and Δ*GDP-MP*) (Figure 6, S9 Table) indicating that parasites lacking these molecules were at a competitive disadvantage in these infections. This effect was apparent as early as two days after infection, consistent with a protective role for PG-containing glycoconjugates in the digesting bloodmeal [48] and a role for LPG in *L. mexicana* attachment to *L. longipalpis* [49].

Paralysed and uncoordinated mutants also became noticeably scarcer as the infection progressed (Figure 6, S9 Table). The aflagellate Δ*IFT88* mutant showed the most severe phenotype and a significant decrease over time was also measured for Δ*PF16, ΔCFAP43, ΔCFAP44, ΔIC140, ΔdDC2* and Δ*RSP4/6.* By contrast, mutants with a mild swimming defect (slower swimmers Δ*LmxM.21.1110,* Δ*FM458* and Δ*LmxM.18.1090* and faster Δ*LC4-like*) (Figure 3D) remained as abundant as the normal swimmers throughout the infection (Figure 6, S9 Table). The exceptions were the slower swimmers Δ*Kharon1,* which is also defective in the transport of a flagellar glucose transporter [50], and Δ*ARL-3A,* which has a short flagellum (Figure 5). Both of these were rarer in the fly compared to the starting pool. To gain anatomical resolution and determine whether an immotile mutant fails to migrate to anterior portions of the fly gut, we infected separate batches of *L. longipalpis* with motile parasite lines and complemented KO lines as controls, with the motile Δ*BBS2* mutant, which lacks a component of the BBSome complex [51] which is expected to play a role in flagellar membrane trafficking, and with the paralysed Δ*PF16* mutant (Figure 7). The Δ*PF16* mutants are among the least motile cells that retain a long flagellum (Figure 5), while having only

moderate levels of flagellar curling (S9 Figure). The axonemal defect resulting in paralysis is

a well-characterised disruption of the central pair in kinetoplastids (Figure 3B and [22,52,53])

and is similar to the defect of the *pf16 Chlamydomonas reinhardtii* mutant [54] indicating it is

a well-conserved core axoneme component. Two days post blood-meal (PBM), the *L.*

mexicana wild type and *L. mex* Cas9 T7 [22] control cell lines and the Δ*BBS2* mutant

developed well, with infection rates above 70%; the Δ*PF16* mutant produced the lowest

infection rate (below 50%). The introduction of an add-back copy of *PF16* into the Δ*PF16* line

restored infection levels (Figure 7A). In all lines, promastigotes were localized in the

abdominal midgut, within the bloodmeal enclosed in the peritrophic matrix (Figure 7B). After

defecation (day 6 PBM), all control lines and the Δ*BBS2* mutant replicated well and

developed late-stage infections with colonisation of the whole mesenteron including the

stomodeal valve (Figure 7B) which is a prerequisite for successful transmission. Their

infection rates ranged from 56% to 83%. By contrast, Δ*PF16 Leishmania* failed to develop;

the infection rate was less than 2% (a single positive fly out of 62 dissected (Figure 7A), with

parasites restricted to the abdominal midgut (Figure 7B)), indicating that Δ*PF16* parasites

were lost during defecation and were unable to develop late stage infections in *L. longipalpis*.

Our data provide strong evidence that flagellar motility is an essential requirement for

successful development in sand flies and, by implication, transmission.

Discussion

This study demonstrates the power of high-throughput CRISPR-Cas9 knockout screens to

discover mutant phenotypes in *Leishmania*. We defined a high-confidence flagellar proteome

and used these data in conjunction with transcriptomics data and prior knowledge of

conserved axonemal proteins to demonstrate a role in motility for >50 genes from a set of

one hundred. We also show the importance of flagellar motility in the colonisation of sand

flies. The data from the pooled mutant population show a progressive loss of paralysed or

uncoordinated swimmers over nine days from infection. Because these data report total

abundance of each genotype in the whole fly without discriminating between regions of the

gut, we probed this question further in infections with the Δ*PF16* mutant, which is essentially

paralysed and incapable of sustained directional motility due to a defined defect in the central

pair complex of the axoneme [22]. The results show that Δ*PF16 Leishmania* were rapidly lost

from most of the dissected flies, consistent with the depletion of this mutant from the mixed

pool, and additionally shows that none of the few remaining parasites reached anterior parts

of the alimentary tract. These findings show that parasite motility is required for completion of

the *Leishmania* life cycle, in line with the essential role of motility in other vector-transmitted protists. For example, Rotureau *et al.,* [55] showed that loss of forward motility, caused by ablation of outer dynein arms though KO of *DNAI1*, rendered *T. brucei* unable to reach the tsetse fly foregut. It seems likely that loss of motility also contributed to the inability of *L. amazonensis* to progress beyond the abdominal midgut of *L. longipalpis* when the parasites overexpressed GTP-locked ADP-ribosylation factor-like protein 3A (Arl-3A) and as a result grew only short flagella [56].

Observations of attached *Leishmania* in dissected sand flies show adhesion specifically via the flagellum but the precise molecular interactions between flagellum and the microvillar gut lining remain to be clarified. The dominant cell surface glycoconjugate LPG which covers the entire parasite surface including the flagellum is known to be important in *Leishmania* attachment to sand fly guts [57] and our results support the view that LPG plays an important role in *L. mexicana* infection of *L. longipalpis* [49]. The proportion of ∆*LPG1* mutants had decreased by two days after infection and reduced further as infection progressed. The observed loss of fitness of the ∆*PMM*, ∆*GDP-MP* and ∆*PMI* mutants is likely the cumulative effect of the loss of LPG and a broader range of mannose-containing glycoconjugates which were shown to protect *Leishmania* in the digesting bloodmeal [48]. The consistency of the pooled mutant data with the reported phenotypes of individual glycoconjugate-deficient mutants demonstrates the power of this new rapid method for mutant phenotyping in *Leishmania*. However, whilst a role for LPG in *L. mexicana* attachment to the fly is well established, the possible contribution of flagellum-specific surface molecules [58] has not yet been conclusively resolved. Zauli et al., [36] reported isolation of *L. braziliensis* from a patient's skin lesion which differentiated to promastigotes with an "atypical" morphology. These cells had a short flagellum barely protruding from the flagellar pocket, with an amorphous tip suggestive of a defect in flagellum elongation. In experimental infections of *L. longipalpis*, these parasites persisted in the fly following defecation of the blood meal, suggesting that they remained sufficiently anchored without a long flagellum. It would be interesting to follow up the subsequent development of this mutant in the fly.

Several lines of evidence suggest a role for the trypanosomatid flagellum in environmental sensing [40,59-61]. Evidence for specific signal transduction pathways aiding promastigote navigation through the sand fly is however limited. Cyclic nucleotide signal transduction pathways may have important roles in coupling environmental sensing with regulation of flagellar beat patterns [62,63]. In our flagellar proteome we identified several adenylate cyclases (ACs), cAMP-specific phosphodiesterases (PDEs) and PKA subunits and mapped their localisations to distinct flagellar subdomains by protein tagging (S7 Figure). The motility

assays showed that deletion of PKA subunits (Δ*LmxM.34.4010* (partial KO only) and
Δ*FM458*) reduced swimming speed, whereas deletion of two different PDEs
(Δ*LmxM.18.1090* and Δ*LmxM.08_29.2440*) increased it, pointing to an activating role for
cAMP in *Leishmania* motility. Knockout of receptor-type adenylate cyclase a-like protein
LmxM.36.3180 had no effect on swimming speed in our motility assay but given the possible
redundancy with other flagellar ACs, this preliminary finding should be followed up by
examination of other AC mutants individually and in combinations. In our pooled KO screen
in sand flies, KOs of PDE LmxM.18.1090 and PKA RSU (FM458) remained as abundant as
the controls, indicating that the mild motility phenotypes measured *in vitro* did not significantly
impair colonisation of flies.
Perturbation of the flagellar membrane might be expected to interfere with sensory functions
mediated through the flagellum. Ablation of membrane proteins LmxM.17.0870 and
LmxM.23.1020 (S7 Figure) did not significantly enhance or reduce the relative abundance of
the respective mutants in sand flies over the nine-day observation period. BBS2 is an integral
part of the core BBSome complex which is highly conserved across ciliated eukaryotes [64]
and functions as a cargo adaptor for ciliary membrane protein trafficking in *Chlamydomonas*
flagella and metazoan cilia [65]. Our pooled mutant data and infections with the *BBS2*
deletion mutant alone found that loss of this gene had no discernible detrimental effect on
survival in sand flies and the parasites' ability to reach the anterior gut. By contrast, KO of
Kharon1, a protein shown to be required for trafficking of the glucose transporter LmGT1,
and perhaps other proteins, to the promastigote flagellum [50] led to slightly reduced fitness
in the flies from the earliest time point. The Δ*Arl-3A* mutants were also less abundant
compared to the controls. This is reminiscent of the previously published abortive phenotype
of *L. amazonensis* overexpressing the constitutively GTP bound *Ld*ARL-3A-Q70L [56]. This
mutant formed only a short flagellum, similar to the Δ*Arl-3A* mutant generated in the present
study. Failed attachment as a result of the shortened flagellum was thought to be a likely
cause for the rapid clearance of *Ld*ARL-3A-Q70L-expressing parasites but it was noted that
an inability to migrate at later stages of development would also lead to the disappearance of
the mutants [56]. In our study the phenotype of the Δ*Arl-3A* mutants was however mild
compared to the aflagellate (Δ*IFT88*) or paralysed mutants. Arl-3A acts as guanine
nucleotide exchange factor in the transport of lipidated proteins to the flagellar membrane
[66] and protein mis-targeting could contribute to the phenotype in addition to flagellar
shortening. Further insights into the contribution of flagellar membrane proteins to attachment
or directional swimming behaviour may be uncovered by further biochemical studies into

flagellar membrane composition and subjecting different mutants (with or without overt motility phenotypes in culture) to chemotaxis assays and fly infections.

In contrast to the absolute requirement of motility for movement through the sand fly vector, flagellar motility is dispensable for promastigote proliferation in culture. Promastigotes are viable and able to divide even if they fail to assemble a flagellum at all, as demonstrated originally by the deletion of cytoplasmic dynein-2 heavy chain gene *LmxDHC2.2* [35] and IFT140 [37] and the phenotypes of knockouts of anterograde and retrograde IFT components in the present study. The ensuing prediction that most gene deletions affecting flagellar function are expected to yield viable promastigotes in the laboratory is borne out by our high success rate of obtaining 96% of attempted knockouts. Thus, in *Leishmania*, flagellar mutant phenotypes can be observed in replication-competent cells over many cell cycles and our mutant library enables detailed systematic studies of KO phenotypes to probe protein functions in flagellum assembly, motility and signal transduction.

A fruitful area for further studies will be dissection of PFR function and assembly mechanisms. This extra-axonemal structure is required for motility as demonstrated through deletion of the major structural PFR components, PFR1 and PFR2 in *Leishmania* [34,67] and ablation of PFR2 by RNAi in *T. brucei* procyclic forms [68] but its precise role remains unclear. The PFR comprises more than 40 proteins, some with structural roles, others with roles in adenine nucleotide homeostasis, cAMP signalling, calcium signalling and many uncharacterised components [69,70] and it may anchor metabolic and regulatory proteins as well as influencing the mechanical properties of the flagellum. Our results showed that fragmentation of the PFR caused by loss of LmxM.21.1110 reduced swimming speed to levels similar to the structurally more severe PFR2 KO. Whether LmxM.21.1110 is required for correct PFR assembly or stabilisation is currently unknown.

Motility mutants analysed in our screen also included deletions of genes with human orthologs linked to ciliopathies (such as *hydin*) or male infertility (*CFAP43* and *CFAP44*) [71]. *Leishmania* offers a genetically tractable system to gain further mechanistic insight into their functions. The *hydin* mutant has been extensively characterised in other species: in mammals, mutations in the *hydin* gene cause early-onset hydrocephalus [72] and subsequent studies on *C. reinhardtii*, *T. brucei* and mice showed that hydin localises to the C2 projection of the central pair complex [73], and that loss of hydin function causes mispositioning and loss of the CP [74] and motility defects [73-75]. The motility phenotype in the *L. mexicana* Δ*hydin* mutant was consistent with these existing data and we made the new observation that the mutant flagella show extensive curling (Figure 4, S9 Figure). Interestingly, *hydin* knockdown in *C. reinhardtii* caused flagella to arrest at the switch point

between effective and recovery stroke, leaving cells with one flagellum pointing up and the other down, prompting speculation that this may indicate a role for hydin in signal transmission to dynein arms [73]. Consistent with this hypothesis, cilia of *hy3/hy3* mouse mutants frequently stalled at the transition point between the effective and recovery stroke [75]. Curling may represent the failure of flagellum bending to reverse during progression of the normal flagellum waveform down the flagellum, leaving the flagellum locked at one extreme of bending, analogous to the ciliary beat *hydin* phenotype. In *L. mexicana*, the Δ*hydin* mutant presented the most severe manifestation of the curling phenotype, which was also observed in a lower proportion of other mutants (S9 Figure). This phenotype may be a consequence of mis-regulated dyneins and the set of mutants exhibiting curling will facilitate further experiments to establish the mechanistic basis for flagellar curling.

Genetic, biochemical and structural studies have provided elegant and detailed models for the mechanisms of flagellar motility [76,77]. Phylogenetic profiling and comparative proteomics studies have yielded insights into the evolutionary history, core conserved structures and lineage-specific adaptations of eukaryotic flagella. Our CRISPR-Cas9 KO method enables rapid targeted gene deletion and characterisation of loss-of-function phenotypes for large cohorts of *Leishmania* genes *in vitro* and *in vivo* and hence new opportunities to interrogate the functions of hitherto poorly characterised flagellar proteins in motility regulation, environmental sensing and axoneme remodelling from 9+2 to 9+0. The bar-seq strategy for phenotyping of mutants can also be used to probe parasite-host interactions in mammals.

Materials and Methods

Cell culture

Promastigote-form *L. mexicana* (WHO strain MNYC/BZ/62/M379) were grown at 28°C in M199 medium (Life Technologies) supplemented with 2.2 g/L NaHCO$_3$, 0.005% haemin, 40 mM 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid (HEPES) pH 7.4 and 10% FCS. The modified cell line *L. mexicana SMP1:TYGFPTY* [17] was cultured in supplemented M199 with the addition of 40 µg/ml G-418 Disulfate. L*. mex* Cas9 T7 [22] was cultured in supplemented M199 with the addition of 50 µg/ml Nourseothricin Sulphate and 32 µg/ml Hygromycin B.

Deflagellation protocol

To avoid proteolytic degradation, all procedures were performed on ice. 2·10^9 *L. mexicana SMP1:TYGFPTY* cells were collected at 800*g* for 15 min at 4°C, washed once in 20 ml

phosphate buffered saline (PBS) and resuspended in 5 ml 10 mM PIPES [10 mM NaCl, 10

mM piperazine-N,N′-bis(2-ethanesulfonic acid, 1 mM $CaCl_2$, 1 mM $MgCl_2$, 0.32 M sucrose,

adjusted to pH 7.2]. 0.375 ml of 1 M Ca^{2+} solution (final conc. 0.075 M) and a protease

inhibitor cocktail [final concentration, 5 μM E-64, 50 μM Leupeptin hydrochloride, 7.5 μM

Pepstatin A and 500 μM Phenylmethylsulfonyl fluoride (PMSF)] were added to the cell

suspension. Cells were deflagellated by passing them 100 times through a 200 μl gel loading

pipette tip (Starlab) attached to a 10 ml syringe. Flagella and cell bodies were separated

through density gradient centrifugation, using a modified version of the protocol in [78]. The

sample was loaded on top of the first sucrose-bed containing three layers of 10 mM PIPES

with 33% (upper), 53% (middle) and 63% (bottom) w/v sucrose [10 mM NaCl, 10 mM

piperazine-N,N′-bis(2-ethanesulfonic acid, 1 mM $CaCl_2$, 1 mM $MgCl_2$, adjusted to pH 7.2 with

either 0.96M, 1.55M or 1.84M sucrose] and centrifuged at 800g for 15 min at 4°C. The pellet

in the 63% sucrose layer was diluted with 10 ml 10 mM PIPES and centrifuged at 800g for 15

min at 4°C. The supernatant was discarded and the pellet resuspended in 40 μl 10 mM

PIPES. This was the cell body fraction. The top layer of the first sucrose-bed, containing

flagella, was collected and sucrose sedimentation was repeated with a second sucrose-bed

containing only one layer of 10 mM PIPES with 33% w/v sucrose. The resulting top layer of

the second sucrose bed was transferred to an ultra-centrifugation tube (Beckmann tubes)

and collected by ultra-centrifugation at 100,000g for 1 h at 4°C (Beckman Coulter). The

supernatant was discarded and the pellet resuspended in 40 μl 10 mM PIPES. This was the

flagellar fraction. All other sucrose layers contained a mixture of flagella and cell bodies and

were discarded. 1 μl of flagellar and cell body fractions was used for counting and imaging

and 36 μl of each fraction were used for proteomic analysis.

Protein Mass spectrometry

Cell body and flagellar fractions were supplemented with 4 μl protease inhibitor cocktail (see

above) and 10 μl octylglycoside (1% (w/v) final conc.), incubated for 20 min on ice and

centrifuged at 18,500g for 1 h at 4°C to separate soluble (supernatant) from insoluble (pellet)

proteins. 50 μl ice cold reducing 2x Laemmli buffer was added to the resulting supernatant.

Pellets were dissolved in 100 μl 1x Laemmli buffer. To avoid aggregation of hydrophobic

proteins, fractions were not boiled prior to SDS-PAGE [79]. 20 μl of flagella fractions and 10

μl of cell body fractions (~5 – 20 μg protein) were pre-fractionated on a 10% polyacrylamide

gel, stained overnight with SYPRO Ruby Protein Gel Stain (Molecular Probes) and destained

for 30 min in 10% (v/v) Methanol / 7% (v/v) acetic acid. Sample preparation in the following

was carried out as described in [80]. Briefly, gel pieces were destained with 50% acetonitrile,

reduced with 10mM TCEP (Tris(2-carboxyethyl)phosphine hydrochloride) for 30 minutes at

RT, followed by alkylation with 55 mM Iodoacetamide for 60 minutes in the dark at RT.

Samples were deglycosylated with PNGase F over two days at RT and digested overnight at

37°C with 100 ng trypsin. Samples were acidified to pH 3.0 using 0.1% trifluoroacetic acid

and desalted by reversed phase liquid chromatography. Samples were analysed on an

Ultimate 3000 RSLCnano HPLC (Dionex, Camberley, UK) system run in direct injection

mode coupled to a QExactive Orbitrap mass spectrometer (Thermo Electron, Hemel

Hempstead, UK).

MS data analysis and SINQ enrichment plot

MS-data were converted from .RAW to .MGF file using ProteoWizard (S6 Table) and

uploaded to the Central Proteomics Facilities Pipeline (CPFP [81]). Protein lists were

generated by using CPFP meta-searches (S6 Table) against the predicted *L. mexicana*

proteome (gene models based on [28], followed by label-free SINQ quantification (S1 and S6

Table). For SINQ enrichment plots detected geneIDs were filtered (p ≥ 0.95, ≥ 2 peptides)

and plotted using normalized spectral indices. For missing indices pseudo spectral indices of

10^{-10} were inserted. The mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE [82] partner repository with the dataset

identifier PXD011057.

Gene tagging

Tagging was achieved by insertion of a drug-selectable marker cassette and fluorescent

protein gene into the endogenous gene to produce an in-frame gene fusion. The fusion PCR

method described in Dean *et al.*, [83] was used for tagging with eYFP, using pJ1170 (pLENT-

YB) as the template plasmid for PCR and selection with 5 µg/ml Blasticidin-S deaminase.

The CRISPR-Cas9 method described in Beneke et al., [22] was used for tagging with

mNeonGreen. The online primer design tool www.LeishGEdit.net was used to design primers

for amplification of the 5' or 3' sgRNA template and primers for amplification of donor DNA

from pPLOTv1 blast-mNeonGreen-blast or pPLOTv1 puro-mNeonGreen-puro. Transfectants

were selected with either 5 µg/ml Blasticidin-S deaminase or 20 µg/ml Puromycin

Dihydrochloride.

CRISPR-Cas9 gene knockouts

Gene deletions were essentially done as described in Beneke et al., [22]. The online primer

design tool www.LeishGEdit.net was used to design primers for amplification of the 5' and 3'

sgRNA templates and for amplification of donor DNA from pTBlast and either pTPuro or

pTNeo. Primers for deletion of *PFR2A-C* were designed with the EuPaGDT primer design

tool [84] using the *PFR2* array sequence from *L. mex* Cas9 T7. For amplification of the

sgRNA template DNA, primers:

Nsg: 5'-gaaattaatacgactcactataggTGCGTGCGGAGGTTTGCACGgttttagagctagaaatagc-3' /

Csg: 5'- gaaattaatacgactcactataggAAGGGTGGACGCCATCTCCGgttttagagctagaaatagc-3'.

For amplification of a pTNeo donor DNA fragment with 80 bp homology arms, primers:

F: 5'GCCCACCCCTTTCACTCTTTCGCTGCTCTCTCACCTCACCGACCCCCGCCTCTTT

CCATCTCTCACTGTGTGCTCCACCTgtataatgcagacctgctgc-3' /

R: 5'-AGCAGCCTTGAGACGACACCTGTAACAAAACCATCACCACAAGCTCCAAGGCGA

CAACATCGCGGGAAGACTTCGCCCCAccaatttgagagacctgtgc-3'.

For transfections on 96-well plates the protocol was modified as follows (similar to

descriptions in [85]): 52 x 10^7 late log phase *L. mex* Cas9 T7 cells (1 x 10^7 cells per reaction)

were collected by centrifugation at 800g for 15 min. A transfection buffer was prepared by

mixing 2 ml 1.5 mM $CaCl_2$, 6.5 ml modified 3x Tb-BSF buffer (22.3 mM Na_2HPO_4, 7.67 mM

NaH_2PO_4, 45 mM KCl, 450 mM sucrose, 75 mM HEPES pH 7.4) and 1.5 ml ddH_2O. The cells

were re-suspended in 3 ml of this transfection buffer and centrifuged again as above. The

heat-sterilised sgRNA and donor DNA PCR products were placed into 48 wells of a 96-well

disposable electroporation plate (4 mm gap, 250 µl, BTX) such that a given well combined all

of the donor DNAs and the sgRNA templates for a given target gene. After centrifugation,

cells were re-suspended in 5.2 ml transfection buffer and 100 µl of the cell suspension

dispensed into each of the 48 wells containing the PCR products. Plates were sealed with foil

and placed into the HT-200 plate handler of a BTX ECM 830 Electroporation System.

Transfection used the following settings: 1500 V, 24 pulses, 2 counted pulses, 500 ms

interval, unipolar, 100 µs. After transfection cells were recovered in 1 ml supplemented M199

in 24-well plates and incubated for 8-16 h at 28°C / 5% CO_2 before addition of the relevant

selection drugs by adding 1 ml of supplemented M199 with double the concentration of the

desired drugs. Mutants were selected with 5 µg/ml Blasticidin-S deaminase in combination

with either 20 µg/ml Puromycin Dihydrochloride or 40 µg/ml G-418 Disulfate and further

incubated. Drug resistant populations typically emerged after one week.

Diagnostic PCR for knockout validation

Drug-selected populations were passaged at least twice (one with at least a 1:100 dilution)

before extraction of genomic DNA using the protocol described in [86]. A diagnostic PCR was

done to test for the presence of the target gene ORF in putative KO lines and the parental

cell line (S8 Figure). Primer3 [87] was used to design, for the entire *L. mexicana* genome

(gene models based on [28]), primers to amplify a short 100 – 300 bp unique fragment of the

target gene ORF (S7 Table). In a second reaction, primers 518F: 5'-

CACCCTCATTGAAAGAGCAAC-3' and 518R: 5'-CACTATCGCTTTGATCCCAGGA-3' were

used to amplify the blasticidin resistance gene from the same genomic DNA samples. For

Δ*PFR2* additional primers were used to confirm deletions (S10 Figure;

1F: 5'-GCAGAAGGAGAAGAGCGAGC-3'; 1R: 5'-CCAGGAACTGCTGGTACTCC-3';

2F: 5'-CGCAGAAGGAGAAGAGCGAG-3'; 2R: 5'-GTTGTACACGGACAGCTCCA-3';

3F: 5'-ACCCCTTTCACTCTTTCGCTG-3'; 4R: 5'-ACCAACGACGTACACAGCAG-3').

Light and electron microscopy

Leishmania cells expressing fluorescent fusion proteins were imaged live. Samples were

prepared as described in [17]. Cells were immediately imaged with a Zeiss Axioimager.Z2

microscope with a 63× numerical aperture (NA) 1.40 oil immersion objective and a

Hamamatsu ORCA-Flash4.0 camera or a 63× NA 1.4 objective lens on a DM5500 B

microscope (Leica Microsystems) with a Neo sCMOS camera (Andor Technology) at the

ambient temperature of 25–28°C.

For transmission electron microscopy, deflagellated cell bodies and isolated flagella were

prepared with a modified version of the chemical fixation protocol described in Höög et al.,

[88]. Pellets of cell fractions were fixed with 2.5% glutaraldehyde in 10 mM PIPES (buffer as

described above) overnight at 4°C. Pellets were washed four times for 15 min in 10 mM

PIPES and overlaid with 10 mM PIPES, containing 1% osmium tetraoxide and incubated at

4°C for 1 h in darkness, then washed five times with ddH$_2$O for 5 min each time and stained

with 500 µl of 0.5% uranyl acetate in darkness at 4°C overnight. Samples were dehydrated,

embedded in resin, sectioned and on-section stained as described previously [88]. Electron

micrographs were captured on a Tecnai 12 TEM (FEI) with an Ultrascan 1000 CCD camera

(Gatan).

Image processing and analysis

598 Micrographs were processed using Fiji [89]. To enable comparisons between the parental

599 and tagged cell lines, the same display settings for the green fluorescence channel were

600 used. PFR length (defined by reporter signal) and flagellar length (distance between stained

601 kinetoplast DNA and flagellar tip) was measured with the ROI manager in Fiji [89].

Motility assays

603 Mutant and parental cell lines were tracked using the previously described method in [32]

604 with three modifications. Firstly, the scripts were modified for batch analysis of multiple files.

605 Secondly, prior to calculation of swimming statistics any 'drift' due to bulk fluid flow in the

606 sample was subtracted. As swimming direction of each cell in the population is in a random

607 direction any drift is visible as a mean population movement between successive frames. We

treated drift as an apparent translation and scaling of cell positions between successive video frames, which was then subtracted. Finally, the primary statistics we considered were mean speed (using the path at 200 ms resolution) and directionality (mean velocity as a fraction of mean speed). Swimming behaviour was measured in triplicates at approximately $6\cdot10^6$ cells/ml and analysed from 5 µl of cell culture on a glass slide in a 250-µm deep chamber covered with a 1.5 mm cover slip using darkfield illumination with a 10× NA 0.3 objective and a Hamamatsu ORCA-Flash4.0 camera on a Zeiss Axioimager.Z2 microscope at the ambient temperature of 25–28°C. The sample was stored inverted prior to use, then turned upright immediately prior to imaging to ensure consistent motion of immotile cells through sedimentation between samples. A sample of the parental cell line killed with a final concentration of 1% formaldehyde was used as a reference for motion of completely paralysed cells through sedimentation and Brownian motion alone.

Lutzomyia longipalpis infections

Sand flies were either infected with pooled mutant populations of *L. mexicana* or individually with *L. mexicana* MNYC/BZ/62/M379 wild type (WT), parental line *L. mex* Cas9 T7, knockout cell line Δ*PF16*, its add-back (*PF16*AB) [22], knockout cell line Δ*BBS2* and its add-back (*BBS2*AB). All parasites were cultivated at 23°C in M199 medium supplemented with 20% foetal calf serum (Gibco), 1% BME vitamins (Sigma), 2% sterile urine and 250 µg/ml amikin (Amikin, Bristol-Myers Squibb). Before experimental infections, logarithmic growing parasites were washed three times in PBS and resuspended in defibrinated heat-inactivated rabbit blood at a concentration of 10^6 promastigotes/ml. *Lutzomyia longipalpis* (from Jacobina, Brazil) were maintained at 26°C and high humidity on 50% sucrose solution and a 12h light/12h dark photoperiod. Sand fly females, 3-5 days old, were fed through a chick skin membrane as described previously [90]. Fully-engorged females were separated and maintained at 26° C with free access to 50% sucrose solution. They were dissected on days 2 or 6 post blood-meal (PBM) and the guts were checked for localisation and intensity of infection by light microscopy. Parasite load was graded as described previously by [91] into four categories: zero, weak (<100 parasites/gut), moderate (100–1000 parasites/gut) and heavy (>1000 parasites/gut).

DNA extraction for bar-seq screen

Mutant *Leishmania* lines were grown separately to late log phase and mixed in equal proportions ($1\cdot10^7$ cells per cell line). This pool was divided equally into three aliquots. DNA was extracted using the Roche High Pure Nucleic Acid Kit according to the manufacturers instructions. Each aliquot was then used to infect three separate batches of 150 sand flies.

The three batches were kept separate and DNA was extracted from 50 whole sand flies two,

six and nine days post blood meal, using the same kit as follows: Sand flies were placed in

two 1.5 ml microcentrifuge tubes (25 flies per tube), overlaid with 100 µl tissue lysis buffer

and frozen at -20°C. The dead flies were defrosted and after addition of 100 µl tissue lysis

buffer and 40 µl proteinase K, flies were disrupted with a disposable plastic pestle (Bel-Art)

and DNA purified according to the kit manufacturer's instructions. DNA was eluted with 50 µl

elution buffer and eluates from the same timepoint were combined, yielding 100 µl in total.

Bar-seq library preparation and sequencing

For each sample, 600 ng isolated DNA was incubated with exonuclease VII (NEB) overnight

at 37°C, purified using SPRI magnetic beads and amplified using PAGE purified custom

designed p5 and p7 primers (Life Technologies), containing indexes for multiplexing and

adapters for Illumina sequencing. Amplicons were again bead purified and multiplexed by

pooling them in equal proportions. The final sequencing pool was once again bead purified

and quantified by qPCR using NEB Library Quant Kit. Library size was determined using

Agilent High Sensitivity DNA Kit on a 2100 Bioanalyzer instrument. The pool was diluted to 4

nM and spiked with 30% single indexed *Leishmania* genomic DNA, prepared using Illumina

TruSeq Nano DNA Library kit according to the manufacturers instructions. The final library

was spiked with 1% PhiX DNA and the Illumina sequencer was loaded with 8 pM to allow low

cluster density (~800 K/mm²). Sequencing was performed using MiSeq v3 150 cycles kit

following the manufactures instructions with paired-end sequencing (2x75 cycles, 6 and 8

cycles index read).

MiSeq raw files were de-multiplexed using bcl2fastq (Illumina). De-multiplexed samples were

subjected to barcode counting using a bash script. Each gene in the *Leishmania* genome

was assigned a unique barcode – the number of times each of these barcodes appeared in

the sequencing output was recorded. The criteria for barcode counting was a 100% match to

the 17 nt total length. Counts for each barcode were normalized for each sample by

calculating their abundance relative to all 25 barcodes. One of the mutants selected for the

pooled screen was excluded from the analysis because sequencing of the cell line showed

eight nucleotide mismatches in the p5 sequence (likely introduced during oligonucleotide

synthesis) which precluded amplification of the barcode region. To calculate "fitness"

normalized barcode counts in the pooled population before feeding were divided by

normalized counts at the relevant time point post blood meal.

Orthofinder

Orthofinder 1.1.2 was used to generate orthogroups for predicted protein coding genes from
48 eukaryotic genomes (32 ciliated species and 16 non-ciliated species): The 45 previously
used by Hodges *et al*. [64] (with *Leishmania major* replaced with *Leishmania mexicana*) and
supplemented with *Aspergillus nidulans*, *Plasmodium berghei* and *Schistosoma mansoni*.

Acknowledgements

We like to thank Svenja Hester, Philip Charles and Benjamin Thomas Central Proteomics
Facility at the Sir William Dunn School of Pathology for help with protein mass spectrometry,
Errin Johnson (Dunn School Bioimaging Facility) for assistance with electron microscopy,
Amanda Williams (University of Oxford) for help with Illumina sequencing, Maxime Cesca
(Université de Paris Sud) and Laura Makin (University of Oxford) for assistance with mutant
characterisation, Oliver Billker (Wellcome Sanger Institute) and members of O.B.'s lab for
helpful discussions about the bar-seq method, Sébastien Pomel (Université de Paris Sud) for
drawing our attention to the PMI, PMM and GDP-MP mutants, Diane McMahon-Pratt (Yale
University) for antibody 2E10 (F-4) and Keith Gull (University of Oxford) for advice and
helpful comments on the manuscript, antibodies L8C4 and L13D6 and access to equipment,
and members of K.G.'s lab for helpful discussions, particularly Samuel Dean and Jack Sunter
for help with development of high-throughput transfection protocols.

Funding statement

This research was jointly funded by the UK Medical Research Council (MRC) and the UK
Department for International Development (DFID) under the MRC/DFID Concordat
agreement; grant no. MR/R000859/1.
EG was supported by a Royal Society University Research Fellowship (UF100435 and
UF160661).
RW was supported by the Wellcome Trust, grant nos. [211075/Z/18/Z, 103261/Z/13/Z,
104627/Z/14/Z]
PV, JSad and TL were supported by European Regional Development (ERD) Funds, project
CePaViP (CZ.02.1.01/0.0/0.0/ 16_019/0000759).
This work was supported by MRC PhD studentships to TB (15/16_MSD_836338) and to JV
(13/14_MSD_OSS_363238), by BBSRC Interdisciplinary Biosciences DTP studentships to
HJ and SS, by an Oxford Radcliffe Scholarship to HJ and by Erasmus grants to TB and FD.

NA was funded by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Moran J, McKean PG, Ginger ML (2014) Eukaryotic Flagella: Variations in Form, Function, and Composition during Evolution. BioScience 64: 1103-1114.

2. Ginger ML, Portman N, McKean PG (2008) Swimming with protists: perception, motility and flagellum assembly. Nat Rev Microbiol 6: 838-850.

3. Vincensini L, Blisnick T, Bastin P (2011) 1001 model organisms to study cilia and flagella. Biol Cell 103: 109-130.

4. Badano JL, Mitsuma N, Beales PL, Katsanis N (2006) The ciliopathies: an emerging class of human genetic disorders. Annu Rev Genomics Hum Genet 7: 125-148.

5. van Dam TJ, Wheway G, Slaats GG, Huynen MA, Giles RH (2013) The SYSCILIA gold standard (SCGSv1) of known ciliary components and its applications within a systems biology consortium. Cilia 2: 7.

6. Broadhead R, Dawe HR, Farr H, Griffiths S, Hart SR, et al. (2006) Flagellar motility is required for the viability of the bloodstream trypanosome. Nature 440: 224-227.

7. Pazour GJ, Agrin N, Leszyk J, Witman GB (2005) Proteomic analysis of a eukaryotic cilium. J Cell Biol 170: 103-113.

8. Mayer U, Kuller A, Daiber PC, Neudorf I, Warnken U, et al. (2009) The proteome of rat olfactory sensory cilia. Proteomics 9: 322-334.

9. Nakachi M, Nakajima A, Nomura M, Yonezawa K, Ueno K, et al. (2011) Proteomic profiling reveals compartment-specific, novel functions of ascidian sperm proteins. Mol Reprod Dev 78: 529-549.

10. Nevers Y, Prasad MK, Poidevin L, Chennen K, Allot A, et al. (2017) Insights into Ciliary Genes and Evolution from Multi-Level Phylogenetic Profiling. Mol Biol Evol 34: 2016-2034.

11. Krüger T, Engstler M (2015) Flagellar motility in eukaryotic human parasites. Semin Cell Dev Biol 46: 113-127.

12. Hochstetter A, Pfohl T (2016) Motility, Force Generation, and Energy Consumption of Unicellular Parasites. Trends Parasitol 32: 531-541.

13. Langousis G, Hill KL (2014) Motility and more: the flagellum of *Trypanosoma brucei*. Nat Rev Microbiol 12: 505-518.

14. Engstler M, Pfohl T, Herminghaus S, Boshart M, Wiegertjes G, et al. (2007) Hydrodynamic flow-mediated protein sorting on the cell surface of trypanosomes. Cell 131: 505-515.

15. Vaughan S (2010) Assembly of the flagellum and its role in cell morphogenesis in *Trypanosoma brucei*. Curr Opin Microbiol 13: 453-458.

16. Alsford S, Turner DJ, Obado SO, Sanchez-Flores A, Glover L, et al. (2011) High-throughput phenotyping using parallel sequencing of RNA interference targets in the African trypanosome. Genome Res 21: 915-924.

17. Wheeler RJ, Gluenz E, Gull K (2015) Routes to a 9+0 flagellum: Basal body multipotency and axonemal plasticity. Nature Communications 6: 8964.

18. Gluenz E, Höög JL, Smith AE, Dawe HR, Shaw MK, et al. (2010) Beyond 9+0: noncanonical axoneme structures characterize sensory cilia from protists to humans. FASEB J 24: 3117-3121.

19. Gadelha AP, Cunha-e-Silva NL, de Souza W (2013) Assembly of the *Leishmania*
amazonensis flagellum during cell differentiation. J Struct Biol 184: 280-292.
20. Killick-Kendrick R, Molyneux DH, Ashford RW (1974) *Leishmania* in phlebotomid
sandflies. I. Modifications of the flagellum associated with attachment to the mid-gut
and oesophageal valve of the sandfly. Proc R Soc Lond B Biol Sci 187: 409-419.
21. Lawyer PG, Ngumbi PM, Anjili CO, Odongo SO, Mebrahtu YB, et al. (1990) Development
of *Leishmania major* in *Phlebotomus duboscqi* and *Sergentomyia schwetzi* (Diptera:
Psychodidae). Am J Trop Med Hyg 43: 31-43.
22. Beneke T, Madden R, Makin L, Valli J, Sunter J, et al. (2017) A CRISPR Cas9 high-
throughput genome editing toolkit for kinetoplastids. R Soc Open Sci 4: 170095.
23. Tull D, Vince JE, Callaghan JM, Naderer T, Spurck T, et al. (2004) SMP-1, a member of
a new family of small myristoylated proteins in kinetoplastid parasites, is targeted to
the flagellum membrane in *Leishmania*. Mol Biol Cell 15: 4775-4786.
24. Trudgian DC, Ridlova G, Fischer R, Mackeen MM, Ternette N, et al. (2011) Comparative
evaluation of label-free SINQ normalized spectral index quantitation in the central
proteomics facilities pipeline. Proteomics 11: 2790-2797.
25. Lubec G, Afjehi-Sadat L (2007) Limitations and pitfalls in protein identification by mass
spectrometry. Chem Rev 107: 3568-3584.
26. Varga V, Moreira-Leite F, Portman N, Gull K (2017) Protein diversity in discrete
structures at the distal tip of the trypanosome flagellum. Proc Natl Acad Sci U S A
114: E6546-E6555.
27. Dean S, Moreira-Leite F, Varga V, Gull K (2016) Cilium transition zone proteome reveals
compartmentalization and differential dynamics of ciliopathy complexes. Proc Natl
Acad Sci U S A 113: E5135-5143.
28. Fiebig M, Kelly S, Gluenz E (2015) Comparative lifecycle transcriptomics revises
Leishmania mexicana genome annotation and links a chromosome duplication with
parasitism of vertebrates. PLoS Pathog 11: e1005186.
29. Lahav T, Sivam D, Volpin H, Ronen M, Tsigankov P, et al. (2011) Multiple levels of gene
regulation mediate differentiation of the intracellular pathogen *Leishmania*. FASEB J
25: 515-525.
30. Emms D, Kelly S (2015) OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthologous gene group inference accuracy. .
Genome Biol. 16:157.
31. Aslett M, Aurrecoechea C, Berriman M, Brestelli J, Brunk BP, et al. (2010) TriTrypDB: a
functional genomic resource for the Trypanosomatidae. Nucleic Acids Res 38: D457-
462.
32. Wheeler RJ (2017) Use of chiral cell shape to ensure highly directional swimming in
trypanosomes. PLoS Comput Biol 13: e1005353.
33. Edwards BFL, Wheeler RJ, Barker AR, Moreira-Leite FF, Gull K, et al. (2018) Direction of
flagellum beat propagation is controlled by proximal/distal outer dynein arm
asymmetry. Proc Natl Acad Sci U S A 115: E7341-E7350.
34. Maga JA, Sherwin T, Francis S, Gull K, LeBowitz JH (1999) Genetic dissection of the
Leishmania paraflagellar rod, a unique flagellar cytoskeleton structure. J Cell Sci 112
(Pt 16): 2753-2763.
35. Adhiambo C, Forney JD, Asai DJ, LeBowitz JH (2005) The two cytoplasmic dynein-2
isoforms in *Leishmania mexicana* perform separate functions. Molecular and
Biochemical Parasitology 143: 216-225.
36. Zauli RC, Yokoyama-Yasunaka JK, Miguel DC, Moura AS, Pereira L, et al. (2012) A
dysflagellar mutant of *Leishmania* (*Viannia*) *braziliensis* isolated from a cutaneous
leishmaniasis patient. Parasit Vectors 5: 11.
37. Fowlkes-Comninellis T, Beverley SM (2015) *Leishmania* IFT140 mutants show normal
viability but lack external flagella: a tool for the study of flagellar function through the
infectious cycle. Cilia 4 (Suppl 1): P49.

38. Sunter J, Gull K (2017) Shape, form, function and *Leishmania* pathogenicity: from
textbook descriptions to biological understanding. Open Biol 7.
39. Bates PA (2008) *Leishmania* sand fly interaction: progress and challenges. Curr Opin
Microbiol 11: 340-344.
40. Leslie G, Barrett M, Burchmore R (2002) *Leishmania mexicana*: promastigotes migrate
through osmotic gradients. Exp Parasitol 102: 117-120.
41. Smith AM, Heisler LE, Mellor J, Kaper F, Thompson MJ, et al. (2009) Quantitative
phenotyping via deep barcode sequencing. Genome Res 19: 1836-1842.
42. Gomes AR, Bushell E, Schwach F, Girling G, Anar B, et al. (2015) A genome-scale
vector resource enables high-throughput reverse genetic screening in a malaria
parasite. Cell Host Microbe 17: 404-413.
43. Bushell E, Gomes AR, Sanderson T, Anar B, Girling G, et al. (2017) Functional Profiling
of a *Plasmodium* Genome Reveals an Abundance of Essential Genes. Cell 170: 260-
272 e268.
44. Ryan KA, Garraway LA, Descoteaux A, Turco SJ, Beverley SM (1993) Isolation of
virulence genes directing surface glycosyl-phosphatidylinositol synthesis by functional
complementation of *Leishmania*. Proc Natl Acad Sci U S A 90: 8609-8613.
45. Garami A, Ilg T (2001) The role of phosphomannose isomerase in *Leishmania mexicana*
glycoconjugate synthesis and virulence. J Biol Chem 276: 6566-6575.
46. Garami A, Mehlert A, Ilg T (2001) Glycosylation defects and virulence phenotypes of
Leishmania mexicana phosphomannomutase and dolicholphosphate-mannose
synthase gene deletion mutants. Mol Cell Biol 21: 8168-8183.
47. Garami A, Ilg T (2001) Disruption of mannose activation in *Leishmania mexicana*: GDP-
mannose pyrophosphorylase is required for virulence, but not for viability. EMBO J
20: 3657-3666.
48. Sacks DL, Modi G, Rowton E, Späth G, Epstein L, et al. (2000) The role of
phosphoglycans in Leishmania-sand fly interactions. Proc Natl Acad Sci U S A 97:
406-411.
49. Jecna L, Dostalova A, Wilson R, Seblova V, Chang KP, et al. (2013) The role of surface
glycoconjugates in *Leishmania* midgut attachment examined by competitive binding
assays and experimental development in sand flies. Parasitology 140: 1026-1032.
50. Tran KD, Rodriguez-Contreras D, Vieira DP, Yates PA, David L, et al. (2013) KHARON1
mediates flagellar targeting of a glucose transporter in *Leishmania mexicana* and is
critical for viability of infectious intracellular amastigotes. J Biol Chem 288: 22721-
22733.
51. Nachury MV, Loktev AV, Zhang Q, Westlake CJ, Peranen J, et al. (2007) A core complex
of BBS proteins cooperates with the GTPase Rab8 to promote ciliary membrane
biogenesis. Cell 129: 1201-1213.
52. Branche C, Kohl L, Toutirais G, Buisson J, Cosson J, et al. (2006) Conserved and
specific functions of axoneme components in trypanosome motility. J Cell Sci 119:
3443-3455.
53. Ralston KS, Lerner AG, Diener DR, Hill KL (2006) Flagellar motility contributes to
cytokinesis in *Trypanosoma brucei* and is modulated by an evolutionarily conserved
dynein regulatory system. Eukaryot Cell 5: 696-711.
54. Dutcher SK, Huang B, Luck DJ (1984) Genetic dissection of the central pair microtubules
of the flagella of *Chlamydomonas reinhardtii*. J Cell Biol 98: 229-236.
55. Rotureau B, Ooi CP, Huet D, Perrot S, Bastin P (2014) Forward motility is essential for
trypanosome infection in the tsetse fly. Cell Microbiol 16: 425-433.
56. Cuvillier A, Miranda JC, Ambit A, Barral A, Merlin G (2003) Abortive infection of
Lutzomyia longipalpis insect vectors by aflagellated LdARL-3A-Q70L overexpressing
Leishmania amazonensis parasites. Cell Microbiol 5: 717-728.
57. Pimenta PF, Saraiva EM, Rowton E, Modi GB, Garraway LA, et al. (1994) Evidence that
the vectorial competence of phlebotomine sand flies for different species of

Leishmania is controlled by structural polymorphisms in the surface
lipophosphoglycan. Proc Natl Acad Sci U S A 91: 9155-9159.

58. Warburg A, Tesh RB, McMahon-Pratt D (1989) Studies on the attachment of *Leishmania*
flagella to sand fly midgut epithelium. J Protozool 36: 613-617.

59. Sharma AI, Olson CL, Engman DM (2017) The Lipid Raft Proteome of African
Trypanosomes Contains Many Flagellar Proteins. Pathogens 6: 39.

60. Rodriguez-Contreras D, Aslan H, Feng X, Tran K, Yates PA, et al. (2015) Regulation and
biological function of a flagellar glucose transporter in *Leishmania mexicana*: a
potential glucose sensor. FASEB J 29: 11-24.

61. Pozzo LY, Fontes A, de Thomaz AA, Santos BS, Farias PM, et al. (2009) Studying taxis
in real time using optical tweezers: applications for *Leishmania amazonensis*
parasites. Micron 40: 617-620.

62. Porter ME, Sale WS (2000) The 9 + 2 axoneme anchors multiple inner arm dyneins and
a network of kinases and phosphatases that control motility. J Cell Biol 151: F37-42.

63. Mukhopadhyay AG, Dey CS (2016) Reactivation of flagellar motility in demembranated
Leishmania reveals role of cAMP in flagellar wave reversal to ciliary waveform. Sci
Rep 6: 37308.

64. Hodges ME, Scheumann N, Wickstead B, Langdale JA, Gull K (2010) Reconstructing the
evolutionary history of the centriole from protein components. J Cell Sci 123: 1407-
1413.

65. Jin H, White SR, Shida T, Schulz S, Aguiar M, et al. (2010) The conserved Bardet-Biedl
syndrome proteins assemble a coat that traffics membrane proteins to cilia. Cell 141:
1208-1219.

66. Ismail SA, Chen YX, Miertzschke M, Vetter IR, Koerner C, et al. (2012) Structural basis
for Arl3-specific release of myristoylated ciliary cargo from UNC119. EMBO J 31:
4085-4094.

67. Santrich C, Moore L, Sherwin T, Bastin P, Brokaw C, et al. (1997) A motility function for
the paraflagellar rod of *Leishmania* parasites revealed by *PFR-2* gene knockouts. Mol
Biochem Parasitol 90: 95-109.

68. Bastin P, Sherwin T, Gull K (1998) Paraflagellar rod is vital for trypanosome motility.
Nature 391: 548.

69. Portman N, Lacomble S, Thomas B, McKean PG, Gull K (2009) Combining RNA
interference mutants and comparative proteomics to identify protein components and
dependences in a eukaryotic flagellum. J Biol Chem 284: 5610-5619.

70. Portman N, Gull K (2010) The paraflagellar rod of kinetoplastid parasites: from structure
to components and function. Int J Parasitol 40: 135-148.

71. Coutton C, Vargas AS, Amiri-Yekta A, Kherraf ZE, Ben Mustapha SF, et al. (2018)
Mutations in CFAP43 and CFAP44 cause male infertility and flagellum defects in
Trypanosoma and human. Nat Commun 9: 686.

72. Davy BE, Robinson ML (2003) Congenital hydrocephalus in hy3 mice is caused by a
frameshift mutation in Hydin, a large novel gene. Hum Mol Genet 12: 1163-1170.

73. Lechtreck KF, Witman GB (2007) *Chlamydomonas reinhardtii* hydin is a central pair
protein required for flagellar motility. J Cell Biol 176: 473-482.

74. Dawe HR, Shaw MK, Farr H, Gull K (2007) The hydrocephalus inducing gene product,
Hydin, positions axonemal central pair microtubules. BMC Biol 5: 33.

75. Lechtreck KF, Delmotte P, Robinson ML, Sanderson MJ, Witman GB (2008) Mutations in
Hydin impair ciliary motility in mice. J Cell Biol 180: 633-643.

76. Lindemann CB, Lesich KA (2010) Flagellar and ciliary beating: the proven and the
possible. J Cell Sci 123: 519-528.

77. Lin J, Nicastro D (2018) Asymmetric distribution and spatial switching of dynein activity
generates ciliary motility. Science 360.

78. Oberholzer M, Langousis G, Nguyen HT, Saada EA, Shimogawa MM, et al. (2011)
Independent analysis of the flagellum surface and matrix proteomes provides insight

911   into flagellum signaling in mammalian-infectious *Trypanosoma brucei*. Mol Cell
912   Proteomics 10: M111 010538.
913 79. Sagne C, Isambert MF, Henry JP, Gasnier B (1996) SDS-resistant aggregation of
914   membrane proteins: application to the purification of the vesicular monoamine
915   transporter. Biochem J 316 (Pt 3): 825-831.
916 80. Gundry RL, White MY, Murray CI, Kane LA, Fu Q, et al. (2009) Preparation of proteins
917   and peptides for mass spectrometry analysis in a bottom-up proteomics workflow.
918   Curr Protoc Mol Biol Chapter 10: Unit10 25.
919 81. Trudgian DC, Thomas B, McGowan SJ, Kessler BM, Salek M, et al. (2010) CPFP: a
920   central proteomics facilities pipeline. Bioinformatics 26: 1131-1132.
921 82. Vizcaino JA, Csordas A, Del-Toro N, Dianes JA, Griss J, et al. (2016) 2016 update of the
922   PRIDE database and its related tools. Nucleic Acids Res 44: 11033.
923 83. Dean S, Sunter J, Wheeler RJ, Hodkinson I, Gluenz E, et al. (2015) A toolkit enabling
924   efficient, scalable and reproducible gene tagging in trypanosomatids. Open Biol 5:
925   140197.
926 84. Peng D, Tarleton R (2015) EuPaGDT: a web tool tailored to design CRISPR guide RNAs
927   for eukaryotic pathogens. Microb Genom 1: e000033.
928 85. Dyer P, Dean S, Sunter J (2016) High-throughput Gene Tagging in *Trypanosoma brucei*.
929   J Vis Exp.
930 86. Rotureau B, Gego A, Carme B (2005) Trypanosomatid protozoa: a simplified DNA
931   isolation procedure. Exp Parasitol 111: 207-209.
932 87. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, et al. (2012) Primer3--new
933   capabilities and interfaces. Nucleic Acids Res 40: e115.
934 88. Höög JL, Gluenz E, Vaughan S, Gull K (2010) Ultrastructural investigation methods for
935   *Trypanosoma brucei*. Methods Cell Biol 96: 175-196.
936 89. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, et al. (2012) Fiji: an
937   open-source platform for biological-image analysis. Nat Methods 9: 676-682.
938 90. Volf P, Volfova V (2011) Establishment and maintenance of sand fly colonies. J Vector
939   Ecol 36 Suppl 1: S1-9.
940 91. Myskova J, Votypka J, Volf P (2008) *Leishmania* in sand flies: comparison of quantitative
941   polymerase chain reaction with other techniques to determine the intensity of
942   infection. J Med Entomol 45: 133-138.
943 92. Dupe A, Dumas C, Papadopoulou B (2015) Differential Subcellular Localization of
944   *Leishmania* Alba-Domain Proteins throughout the Parasite Development. PLoS One
945   10: e0137243.
946 93. Subota I, Julkowska D, Vincensini L, Reeg N, Buisson J, et al. (2014) Proteomic analysis
947   of intact flagella of procyclic *Trypanosoma brucei* cells identifies novel flagellar
948   proteins with unique sub-localization and dynamics. Mol Cell Proteomics 13: 1769-
949   1786.
950 94. Kozlowski LP (2016) IPC - Isoelectric Point Calculator. Biol Direct 11: 55.
951 95. Kohl L, Sherwin T, Gull K (1999) Assembly of the paraflagellar rod and the flagellum
952   attachment zone complex during the *Trypanosoma brucei* cell cycle. J Eukaryot
953   Microbiol 46: 105-109.
954 96. Ismach R, Cianci CM, Caulfield JP, Langer PJ, Hein A, et al. (1989) Flagellar membrane
955   and paraxial rod proteins of *Leishmania*: characterization employing monoclonal
956   antibodies. J Protozool 36: 617-624.
957

Legends

Figure 1. Isolation of flagella and deflagellated cell bodies

(A) Overview of the deflagellation and differential centrifugation protocol. Percentage sucrose concentration (w/v) is indicated. **(B)** Micrographs show merged phase and fluorescence channel (SMP-1::GFP, green; Hoechst DNA stain, red) for each isolation stage (i-iv) depicted in (A). **(i)** *L. mexicana* SMP1::GFP cells before deflagellation, **(ii)** cells after deflagellation, **(iii)** isolated flagella (F) and **(iv)** deflagellated cell bodies (CB). Scale bars represent 20 μm. **(C)** Protein gel stained with SYPRO RUBY. Numbers on the left indicate molecular weight in kDa, numbers below indicate cell equivalent of protein loaded on each lane. Each sample lane of F_S, F_I, C_S and C_I was cut into eight pieces and analysed by mass spectrometry (two biological replicates). **(D)** Venn diagram shows total number of all detected proteins (≥ 2 peptides detected, p-value > 0.95) of both replicates.

Figure 2. Enrichment plot of flagellar and cell body proteins

Proteins detected by MS were quantified with SINQ. Plotted on the X-axis is the log2 fold-change of the spectral index (C_S+C_I / F_S+F_I) and on the Y-axis the log2 fold change of the spectral index (F_S+C_S / F_I+C_I). To allow plotting of proteins that were only detected in one fraction, a value of 10^{-10} was inserted for missing spectral indices. The resulting diagonal lines in each quadrant represent proteins uniquely detected in the respective fraction. **(A)** Each data point represents one of 2414 proteins detected in MS run 1 and bubble size reflects protein abundance. Coloured circles indicate representative proteins for different flagellar sub-structures (GeneIDs in S2 Table). The plot can be interactively explored on http://www.leishgedit.net/leishgedit_db/. **(B)** Correlation with RNA-seq data. All 2414 proteins detected in MS run 1 were plotted as in (A) and colour-coded according to the log2 fold change of differentially expressed transcripts between promastigotes (PRO) and amastigotes (AMA) [28].

Figure 3. Identification of motility phenotypes

All deletion cell lines generated in this study and the Δ*PF16* mutant [22] were analysed for defects in swimming speed or directionality (the ratio of velocity to speed). **(A)** Mean swimming speeds. Speeds were measured three times and the mean of all three replicates (●) and the individual replicates (○) are shown. Error bars represent the standard deviation. Red dotted lines indicate two standard deviations above and below the parental cell line (Cas9 T7) mean swimming speed. Cas9 T7 cells killed with 1% formaldehyde (Cas9 T7 fixed) were used as a completely immotile control. *** p<0.0005, ** p<0.005, * p<0.05

(Student's t-test compared to the parental cell line). For a sub-set of mutants, an addback
copy of the deleted gene was introduced and swimming speeds restored toward the wild-
type. **(B)** The swimming speeds of all knockout mutants (○), as in (A), plotted against mean
directionality. Error bars represent the standard deviation of the three replicates. Four main
mutant phenotype clusters are apparent: Paralysed (including mutants lacking a long
flagellum), uncoordinated (which move slowly, but with greatly reduced directionality), slow
(which move slowly, but with reduced directionality and speed) and fast (which move faster,
with a tendency for higher directionality). **(C)** Speed and directionality of all knockout mutants
in (B) with deletion of IFT components, mutants with a tendency for curling of the flagellum
(S9 Figure) or disrupted PFR structure (Figure 5) highlighted. **(D)** Speed and directionality of
knockout mutants in (B) with those passaged though sand flies (Figure 6) highlighted.

Figure 4. Phenotype categories
Categories of mutant phenotypes: among cells with normal morphology there were normal
swimmers as well as slower or faster swimmers. Four categories of distinct morphological
defects were observed, which all lead to impaired motility: no flagellum, short flagellum,
curled flagellum and disrupted PFR. One representative mutant is shown for each category,
the deleted gene is indicated. Green, PFR2::mNG signal; red, Hoechst-stained DNA. Scale
bar, 2 μm.

Figure 5. Measurement of flagellar and PFR length in motility mutants
(**A**) Measurements of flagellar length (measured from kinetoplast DNA to flagellar tip; grey
bar) and PFR2::mNG signal (green bar; data in S8 Table). Error bars show standard
deviation. At least 70 measurements were recorded per cell line. The GeneIDs / gene names
indicate the deleted gene. Numbers above bars show PFR : flagellar length ratio.
Measurements were compared to the parental cell line expressing PFR2::mNG and p-values
(Students t-test) for flagellar length (grey) and PFR length (green) are indicated: *** p≤0.001.
(**B**) Fluorescence micrographs showing tagged mutant cell lines used for measurements in
(A). Green, PFR2::mNG signal; red, Hoechst-stained DNA. Scale bar, 5 μm. White arrows
indicate PFR defects.

Figure 6. Relative fitness of *Leishmania* mutants in *L. longipalpis* infections
The plots display abundance of barcodes at time points post bloodmeal (PBM) relative to the
abundance of this barcode in the mixed parasite population used to infect *L. longipalpis*.
Mutants are grouped according to the function of the deleted gene and severity of motility
phenotype (**A**) flagellar mutants with severe motility defects (paralyzed, uncoordinated

swimmers and aflagellate cells), (**B**) flagellar mutants with mild motility defects, (**C**) mutants

lacking flagellar membrane proteins or proteins involved in protein trafficking to the flagellar

membrane, (**D**) mutants lacking key enzymes for the synthesis of LPG and other

glycoconjugates, (**E**) control mutants with wild type motility. Data points represent the

average of three replicates. Error bars show the standard deviation of the mean of the three

replicates. Dotted red lines indicate two standard deviations above and below the parental

cell line (SBL1-5). Measurements were compared (two-sided t-test) to the average of all five

parental controls and p-values are indicated: *≤0.05, **≤0.005, ***≤0.0005.

Figure 7. Development of *L. mexicana* ΔPF16 in *L. longipalpis*

(A) Infection rates (% of infected females) and intensities of infections (four categories) of *L.*

mexicana in *L. longipalpis*. Numbers above the bars indicate numbers of dissected females

in the group. **(B)** Localization of *L. mexicana* infections in *L. longipalpis*. ESP: endoperitrophic

space, AMG: abdominal midgut, TMG: thoracic midgut, CAR: cardia, SV: stomodeal valve.

Numbers above the bars indicate numbers of infected females in the group.

Figure 1

A

Soluble

log2 fold change

Insoluble

Flagellum ⟷ Cell body
log2 fold change

○ 2414 detected proteins
● Tripartite attachment complex
● Flagellar attachment zone
● Flagellar tip
● Central pair
● Outer or Inner dynein arm

● Conserved ancestral centriolar protein
● Paraflagellar rod
● Intraflagellar transport
● Flagellar membrane
● Dynein regulatory complex
● Radial spoke protein

B

Soluble

log2 fold change

Insoluble

Flagellum ⟷ Cell body
log2 fold change

PRO 5 ●●●●●●●●●●●● AMA -5

Log2 fold change DEseq
Fiebig et al., 2015

Figure 2

Figure 4

Figure 5

Figure 6

Figure 7