Pervasive positive linkage between bacterial social traits suggests the

predominance of competitive interactions in natural *Pseudomonas* communities

Jos Kramer[1]*, Miguel Ángel López Carrasco[1,2] and Rolf Kümmerli[1]

[1] Department of Plant and Microbial Biology, University of Zürich, Switzerland

[2] Depto Biología Cellular, Genética y Fisiología, University of Málaga, Spain

* Corresponding author: Jos Kramer; phone: +41 (0)44 63 52902; e-mail: JosKramer@gmx.de

Words: 6885 | Figures: 3 | Tables: 3

ABSTRACT

Bacteria frequently cooperate by sharing secreted metabolites such as enzymes and siderophores. Many of these 'public good' traits are linked at the regulatory level, and studies on laboratory systems have shown that trait linkage is a key determinant of strain dynamics and community composition. Here, we test to what extent regulatory linkage among social traits occurs in natural *Pseudomonas* communities by investigating investment levels and correlations between five public goods: biosurfactants, biofilm matrix components, proteases, pyoverdines, and toxic compounds. Our analyses involving 315 isolates from soil and freshwater communities revealed that their investment into social traits varied dramatically, and that correlations between traits were frequent, exclusively positive, and sometimes habitat-specific. Our results indicate that *Pseudomonas* communities are dominated by strains lying on a continuum between a 'social' type producing multiple public goods, and an 'asocial' type showing low investment into social traits. This pattern suggest that competitive interactions, where asocial types act as 'supercheaters' and exploit social types, are more common than the mutualistic exchange of different public goods among strains with specialized trait repertoires. Overall, our work sheds light on the complex interdependencies among social traits, and indicates that expression constraints shape the evolution of microbial sociality in nature.

Keywords: public goods; bacterial communities; competition; mutualism; regulatory linkage

INTRODUCTION

Bacteria are social organisms. They form multicellular structures such as biofilms, communicate with each other via chemical signals, and engage in group-coordinated motility, resource acquisition and 'chemical warfare' against predators or competitors (West et al. 2007; Foster 2010). Many of these social behaviours involve the cooperative release of costly secondary metabolites. For instance, biofilm formation requires the production of structural polysaccharides, group-coordinated motility often involves the production of biosurfactants, and resource acquisition is based on the production of exo-enzymes and metal-chelating siderophores (Griffin et al. 2004; Nadell et al. 2009; Kearns 2010; Jousset et al. 2013). Once released, these compounds are – in principle – publicly accessible to all neighbouring cells. Such 'public goods' do therefore not only benefit (clonemates of) the producer, but can also affect other community members. The resulting interactions cover all aspects of bacterial life, ranging from 'cheating' (where strains deficient for the production of a public good free-ride on the common pool of this resource) (West et al. 2006; Özkaya et al. 2017) over inter-strain competition (Cordero et al. 2012; Bruce et al. 2017; Leinweber et al. 2018) to division of labour between bacterial strains exchanging different types of public good (van Gestel et al. 2015; Dragoš et al. 2018). There is increasing evidence that such social interactions play a crucial role during hosts infection (Rumbaugh et al. 2009; Pollitt et al. 2014; Andersen et al. 2015; Granato et al. 2018), in various biotechnological and environmental processes (Bachmann et al. 2016; Hesse et al. 2018) and within the microbiome (Rakoff-Nahoum et al. 2016).

While many aspects of bacterial social interactions are well understood for individual traits, it is important to consider that different traits are often linked at the regulatory level.

Regulatory linkage leads to an additional level of complexity, as selection on one trait can have

pleiotropic consequences for other traits (Jousset et al. 2009; Wilder et al. 2011; Dandekar et al.

2012; Wang et al. 2015). For example, for *Pseudomonas* bacteria it was shown that a reduction

of exo-protease production can increase sensitivity to predation (Jousset et al. 2009), and reduce

metabolic abilities and toxin resistance (Dandekar et al. 2012; Wang et al. 2015), thereby

compromising the spread of cheaters. Especially when regulatory linkage occurs between two or

more public good traits, the resulting pleiotropic effects can have far-reaching consequences.

Positive linkage – where mutations reducing the production of one public good also reduce the

production of a second public good – can accelerate the loss of cooperation, and could hence

favour the emergence of 'supercheaters' that simultaneously free-ride on multiple public goods

(Granato et al. 2018). Conversely, negative linkage – where a reduced production of one public

good leads to an increase in the production of another public good – can promote the emergence

of 'mosaic producers' with a limited public good repertoire, which can potentially complement

each other (Driscoll et al. 2011; Ross-Gillespie et al. 2015). In addition to these two scenarios,

both positive and negative linkage can stabilize cooperation, namely if changes in the expression

of linked traits have negative net fitness effects on the individual (Harrison and Buckling 2009).

While these studies on laboratory systems show that regulatory linkage and the resulting

constraints are key determinants of bacterial social interactions and strain co-existence, we know

little on whether trait linkage occurs in natural bacterial communities and whether it has

disruptive or stabilising effects on communities. Here, we tackle this question and investigate

whether and how the production of five different public goods is correlated across 315

Pseudomonas strains isolated from eight soil and eight freshwater communities (18-20 strains

per community). In a first step, we screened each strain for the production of: (i) pyoverdines, a

highly variable class of siderophores used to scavenge iron (Griffin et al. 2004); (ii) proteases used

to digest extracellular proteins (Diggle et al. 2007); (iii) biosurfactants used during group-motility

(Xavier et al. 2011); (iv) structural components used to form biofilms (Nadell et al. 2009); and (v)

toxic compounds used to fight predators and competitors (Jousset 2012). We then used our

phenotypic data set to ask whether: (a) strains vary in the extent to which they invest in the public

goods; (b) there is a phylogenetic signal for trait expression, where more closely related strains

produce similar amounts of a particular public good; (c) production levels of different public

goods correlate with one another, suggesting positive or negative regulatory linkage; and (d)

there are differences in linkage patterns between habitats.

MATERIALS AND METHODS

Sampling, isolation and identification of Pseudomonads. We sampled 16 *Pseudomonas*

communities from soil and pond habitats (n = 8 communities per habitat) located on the campus

of the University of Zürich (47.40° N, 8.54° E), Switzerland. We isolated 320 strains (n = 20 strains

per community) on a medium selective for fluorescent pseudomonads, and then verified their

affiliation to this group by amplifying and sequencing part of the housekeeping gene *rpoD* (a

commonly used phylogenetic marker for pseudomonads) (Mulet et al. 2009). The 315 isolates

that could be unequivocally identified as pseudomonads were preserved as freezer stocks until

their use in our experiments (see (Butaitė et al. 2017) for details on the sampling, isolation and

identification procedure).

Phenotypic screening of natural isolates. We used standard phenotypic assays to screen all

isolates for the production of the five public goods (see supplementary methods for details).

Before each assay, we grew our isolates from freezer stocks for 24h in rich medium, either in 50

ml Falcon tubes under shaking conditions (190 rpm) (for the protease assay), or in 96-well plates

under static conditions (for all other assays). We quantified biosurfactant production as the

extent of the biosurfactant-induced collapse of droplets of culture supernatant on a hydrophobic

surface (Bodour and Miller-Maier 1998). Specifically, we grew isolates for 24h in a glucose-

supplemented minimal medium, harvested the supernatant, spotted 5 µl supernatant-droplets

on an oil-coated lid of a 96-well plate, and then measured the droplet surface area after one

minute using the software ImageJ (Schneider et al. 2012). We used the crystal violet assay to

quantify the production of biofilm matrix components (Savoia and Zucca 2007). In particular, we

first grew isolates in Lysogeny Broth (LB) for 24h in 96-well plates, and then measured the amount

of crystal violet retained by the newly formed biofilm matrix at 570nm using an Infinite M200 Pro

microplate reader (Tecan Trading AG, Switzerland). To quantify protease production, we spotted

1 µL aliquots of bacterial culture onto skim milk agar in 24-well plates, and then used ImageJ to

measure the size of the proteolytic halo around colonies after 24h of growth (Jousset et al. 2013).

To quantify pyoverdine production, we grew isolates in iron-limited casamino acids medium for

24h, and measured the natural fluorescence of pyoverdines in the culture supernatants

(emission|excitation at 400|460 nm, respectively) (Kümmerli et al. 2009). Finally, for the

production of toxic compounds we developed a supernatant feeding assay to assess to which

extent isolates affect the growth of six reference strains (*P. aureofaciens* ATCC13985, *P.*

entomophila, *P. putida* IsoF, *P.syringae* B728a, *P.protegens* CHA0, and *P. aeruginosa* PA01). We

first generated supernatant from each isolate after 24h growth in LB, then exposed the reference

strains to each supernatant, and finally measured their growth in LB during 24h. For each

screening assay, we grew isolates under static conditions at 28°C in fourfold replication, and

measured bacterial growth of the public-goods producing cultures (as optical density at 600 nm,

or – in the case of protease production – as colony size).

Phylogenetic signal in public goods production. To explore the relationship between the

phylogeny of our *Pseudomonas* isolates and their public goods production, we constructed

maximum-likelihood phylogenetic trees based on partial rpoD gene sequences. We constructed

separate trees for soil and pond habitats, based on, respectively, 149 and 148 (of the 158 and

157) strains for which we obtained sequence lengths ≥ 600 bp. We used *P. aeruginosa* PAO1 as

an outgroup, and included 20 characterised fluorescent pseudomonads as references to cover

the broad phylogenetic diversity of our natural strains (see (Butaitė et al. 2017) for details on tree

construction). Next, we mapped the production profiles of strains onto the trees (using the web

tool iTOL (Letunic and Bork 2016)), and then calculated Blomberg's K for each trait and habitat,

to quantify the extent to which phylogeny predicts public good production profiles. K-values close

to one reflect strong phylogenetic signals, suggesting that phylogenetic history is the main factor

driving the trait covariance among strains). Conversely, K-values close to zero reflect weak

signals, suggesting that factors apart from phylogeny affect trait evolution (Blomberg et al. 2003).

Note that we ignored branch lengths for the calculation of the phylogenetic signal (cf. (Butaitė et

al. 2017, 2018)), because the high variability of branch lengths observed among our isolates led

to biased and underestimated phylogenetic signals.

Statistical analysis. All statistical analyses were conducted using R version 3.5.0 (www.r-project.org). We used a five-step procedure to investigate the production of – and correlation among – different public goods in our *Pseudomonas* communities. To preclude that differences in growth lead to spurious correlations among social traits, we first checked for associations between growth and public goods production. As growth was positively correlated with the production levels of biofilm matrix components (Spearman's rank correlation: $\rho = 0.341$, $p < 0.001$), proteases ($\rho = 0.238$, $p < 0.001$) and pyoverdines ($\rho = 0.750$, $p < 0.001$), we scaled the production levels of these traits relative to growth. Growth was independent of the production levels of biosurfactants ($\rho = -0.058$, $p = 0.301$) and toxic compounds ($\rho = 0.094$, $p = 0.097$). Next, we scaled the measurements of public goods production either relative to the corresponding measurements of the laboratory reference strain that had been grown alongside the natural isolates, or – in the case of toxin production – relative to the growth of the reference strains in the absence of our isolates' supernatants. Note that we transformed the supernatant effect (by calculating [supernatant effect - 1]*[-1]) so that values above and below zero indicate, respectively, an inhibition (i.e. toxicity) and stimulation of growth. In the third step of our analyses, we transformed the measurements of all five social traits using a rank-based inverse normal (RIN) transformation (Bishara and Hittner 2012). This allowed us to investigate the production of and correlations among public good traits using parametric analyses. The transformation was necessary, as preliminary analyses based on the untransformed measurements unequivocally yielded heavily-skewed residual distributions. In the fourth step, we used a nested-design permutational multivariate analysis of variance (PERMANOVA) to test whether the social profiles, including all five social traits, differed between soil and pond isolates.

Finally, we used linear mixed models to test for (habitat-specific) links between the social traits.

We used the 'false discovery rate' (Benjamini and Hochberg 1995) to correct the p-values of

correlations between pairs of social traits for multiple testing. Where appropriate, we included

'community' as a random effect to account for our nested sampling design. Further details on

the statistical analyses are provided in the Supplementary Material.

RESULTS

Natural Pseudomonas isolates greatly vary in their social profiles

The social profile of our 315 *Pseudomonas* isolates differed between habitats, and varied

substantially both among and within communities (figure 1). The difference between habitats

(PERMANOVA: n_{perm} = 1000; habitat: F_1 = 2.545, p = 0.045, figure 1a) was mainly driven by a

higher production of pyoverdines and marginally higher biofilm matrix component production in

freshwater as compared to soil isolates (table 1). By contrast, variation among communities (F_{14}

= 3.907, p = 0.001, figure 1b-f) was high for all five social traits (table 1). Within-community

variation followed trait-specific patterns (table S1). In particular, communities were dominated

by strains producing no or low amounts of biosurfactants, while harbouring only few members

producing high amounts of this public good (figure 1b). A similar albeit less pronounced pattern

governed variation in biofilm production, with most strains producing low to intermediate

amounts of matrix components, and only few strains investing more substantially in this social

trait (figure 1c). Within-community variation in the production of proteases was characterized by

a bimodal pattern, whereby 35.9 % of all isolates produced no proteases at all, while the

remaining isolates produced considerable amounts of this public good (figure 1d). By contrast,

variation in pyoverdine production spanned the entire continuum from zero to extremely high

production levels (figure 1e). Similarly, the extent to which isolates had toxic effects on our

references strains covered the entire continuum from being atoxic to strongly inhibitory (figure

1f). In summary, the high variation in the production of all five types of public good leaves ample

scope for the occurrence of both negative and positive trait linkage.

Phylogenetic signal for public goods production is higher in soil compared to pond

The production levels of all five public goods showed a phylogenetic signal (table 2, figure 2a-b).

Overall, phylogenetic signals were higher among soil than among freshwater isolates (paired t-

test; t_4 = 3.790, p = 0.0193). In both habitats, the signals were significantly higher than expected

from the null model for all but one trait. Specifically, biofilm production yielded a significant signal

only among soil but not among freshwater isolates (table 2). Signals were generally of

intermediate strength, indicating that both common ancestry and frequent changes in trait

expression profiles within clades of related strains shape public good production profiles.

Table 1 | Among-community variation in the expression of public good traits. Given are the grand median as

well as the minimal and maximal median of the production of biosurfactants, biofilm matrix components,

proteases, pyoverdines and toxic compounds in soil and freshwater communities, as well as the results of

tests for habitat- and community-specific differences in the production levels of these traits.

	grand [min|max] median		Δ habitat		Δ community	
	soil	pond	F_1	p	F_{14}	p
biosurfactants	0.017 [-0.111|0.388]	-0.005 [-0.194|0.132]	1.387	0.282	6.439	**0.001**
biofilm	0.200 [0.080|0.285]	0.301 [0.178|0.492]	3.795	0.065	1.971	**0.017**
proteases	0.402 [0.000|0.605]	0.485 [0.000|0.546]	0.152	0.705	3.278	**0.001**
pyoverdines	0.380 [0.090|0.803]	0.698 [0.551|1.127]	9.344	**0.009**	2.379	**0.004**
toxins	0.031 [-0.004|0.045]	0.039 [0.008|0.059]	1.505	0.245	5.855	**0.001**

Significant p-values (here and below) in bold.

Figure 1 | Social profile of *Pseudomonas* isolates. Depicted are (A) an overall representation of the social

profile of 315 natural *Pseudomonas* strains from eight soil and eight pond communities (in yellow and blue,

respectively), as well as the variation in the production of (B) biosurfactants, (C) biofilm components, (D)

proteases, (E) pyoverdines and (F) toxic compounds within and across communities. The depiction of the

social profile in (A) is based on the first two components extracted from a principal component analysis

based on the RIN-transformed values. Shaded areas are 95% confidence ellipses. The values of (B)-(E) are

relative to the respective value of a laboratory control (*P. aeruginosa* PAO1, indicated by the dashed line).

Values in (F) are the average toxic effect of the strains' supernatant on the growth of six reference strains.

Positive correlations between public good traits predominate

The production levels of the five examined public goods were positively correlated in seven out

of ten possible pairwise comparisons (table 3, figure 3). While four of these positive correlations

showed congruent patterns in soil and pond communities, the remaining three were habitat-

specific (table 3). In particular, the production of proteases and biosurfactants correlated only

among freshwater (slope ± se 0.183 ± 0.073, $t_{124.7}$ = 2.488, p = 0.021) but not soil isolates (-0.034

± 0.075, $t_{155.9}$ = -0.459, p = 0.700), whereas the production of pyoverdine and biofilm matrix

components was positively correlated among soil (0.348 ± 0.067, $t_{155.7}$ = 5.220, p < 0.001) but not

freshwater isolates (0.112 ± 0.085, $t_{154.3}$ = 1.319, p = 0.271). Moreover, pyoverdine production

increased with toxicity in both habitats, but this increase was more pronounced among soil

(0.456 ± 0.083, $t_{153.3}$ = 5.498, p < 0.001) as compared to freshwater isolates (0.167 ± 0.067, $t_{153.2}$

= 2.493, p = 0.021). When focusing on specific trait combinations, we found that biosurfactant

production was poorly linked to any of the other four traits, with only one positive correlation

occurring with protease production in ponds (table 3, figure 3). In stark contrast, we observed

pervasive positive correlations between the production of biofilm matrix components, proteases,

pyoverdines, and toxic compounds (table 3, figure 3).

Table 2 | Phylogenetic signal in the production of public goods. Given are the phylogenetic signal (Blomberg's

K) and the results of a randomization test (under the null hypothesis of the absence of a phylogenetic signal)

for the production of biosurfactants, biofilm matrix components, proteases, pyoverdines, and toxic

compounds among soil and freshwater isolates, respectively.

	soil		freshwater	
	Blomberg's K	p	Blomberg's K	p
biosurfactants	0.499	**0.001**	0.206	**0.023**
biofilm	0.349	**0.001**	0.181	0.063
proteases	0.516	**0.001**	0.357	**0.001**
pyoverdines	0.325	**0.001**	0.251	**0.001**
toxins	0.283	**0.001**	0.211	**0.004**

Figure 2 | Maximum-likelihood cladograms for soil (A) and pond (B) isolates based on partial rpoD sequences.

For both habitats, published *rpoD* sequences of 20 well-characterized fluorescent pseudomonads and *P.*

aeruginosa PAO1 were integrated into the cladograms to demonstrate taxonomic affiliation and diversity

of our environmental isolates. Yellow triangles and blue circles indicate bootstrap values (50-100%) for

branches in the soil and the pond cladogram, respectively. From the inside outwards, bars depict the

relative production levels of biosurfactants (red), biofilm components (grey), proteases (orange),

pyoverdine (green), and toxic compounds (blue), respectively. Colour strips around cladograms represent

the communities (in shades of grey) from which isolates originated.

Table 3 | Pairwise correlations among public good traits. Shown are the (habitat-specific) pairwise correlations of the production of biofilm matrix components, proteases, pyoverdine and toxic compounds with the respective other social traits.

	biofilm		proteases		pyoverdines		toxins	
	χ_1^2	p	χ_1^2	p	χ_1^2	p	χ_1^2	p
biosurfactants (bis)	0.013	0.908	7.531	**0.006**	0.212	0.700	1.226	0.349
habitat	3.840	0.050	0.291	0.590	9.096	**0.003**	1.629	0.202
bis : habitat	0.551[§]	0.458[§]	4.374	**0.037**	0.898[§]	0.343[§]	3.284[§]	0.070[§]
biofilm (bif)			7.222	**0.016**	2.190	0.139	26.245	**< 0.001**
habitat			0.032	0.859	7.108	**0.008**	0.829	0.363
bif : habitat			1.479[§]	0.224[§]	4.666	**0.031**	3.559[§]	0.059[§]
proteases (pro)					27.686	**< 0.001**	40.363	**< 0.001**
habitat					6.709	**0.010**	1.194	0.275
pro : habitat					1.225[§]	0.268[§]	0.126[§]	0.722[§]
pyoverdines (pvd)							6.040	**0.014**
habitat							0.375	0.541
pvd : habitat							7.484	**0.006**

[§] values prior to the removal of the interaction from the model. Note that we obtain the same results when the respective 'response' and the 'explanatory' social traits are swapped (Table S2).

Discussion

Although regulatory linkage between bacterial social traits occurs frequently and has important consequences for strain interactions, pathogenicity and community functioning in laboratory systems (Harrison and Buckling 2009; Driscoll et al. 2011; Ross-Gillespie et al. 2015; Granato et al. 2018), we know surprisingly little about the prevalence and direction of social trait linkage in natural bacterial communities. Here we addressed this knowledge gap by examining and comparing the production levels of five social traits across 315 *Pseudomonas* isolates originating

from eight soil and eight freshwater communities. The social traits we focused on included the

production of biosurfactants used during group motility; matrix components required for biofilm

formation; proteases used for extra-cellular protein digestion; pyoverdines required for iron

scavenging; and toxic compounds used to fight competitors and predators. We found that (i)

natural isolates varied dramatically in the extent to which they invest in social traits, and showed

that (ii) correlations between social traits occurred frequently, (iii) were exclusively positive, and

(iv) in some cases habitat-specific. A comparison across traits suggest that natural *Pseudomonas*

communities are dominated by strains that lie on a continuum between two main strain types –

a 'social' type that is proficient in expressing multiple social traits, and an 'asocial' type that

generally shows low investment levels into any of the social traits. Conversely, mosaic producers

with a specialized trait repertoire characterized by high investment in only some (but low

investment in other) social traits were rare. These patterns suggest that communities of natural

pseudomonads are dominated by competitive interactions with 'asocial' types potentially acting

as supercheaters that simultaneously exploit multiple social traits. In contrast, mutualistic

interactions based on the reciprocal exchange of different public goods seem less common.

At the mechanistic level, correlated trait expression could arise because public goods are

linked at the regulatory level. Well-known and wide-spread co-regulatory systems in

pseudomonads include quorum sensing systems, the GacS/GacA two-component system, and

systems where public goods themselves serve as signals to co-regulate other social traits (Lamont

et al. 2002; Jimenez et al. 2012; Dumas et al. 2013). Alternatively, trait expression could be

environmentally induced, which would mean that strains classified as non-producers for a

particular social trait might actually express this trait under different conditions. For instance,

some strains might grow poorly in our artificial culture media and thereby remain below the

quorum-sensing threshold required for social trait expression. Such environmental effects might

explain the generally low levels of biosurfactant production (and thus the scarcity of correlations

between the production of biosurfactants and the other public goods). But overall, we believe

that environmental trait induction plays a relatively minor role in our data set. This is because

the production of public goods (except for pyoverdines) was assessed in nutrient-rich media,

where strains grew well. Moreover, we corrected for differences in bacterial densities whenever

the production level of a public good correlated with growth. In the scenarios discussed below,

we therefore assume that the observed correlations at least partly reflect regulatory linkage.

At the evolutionary level, correlated trait expression could arise because high and low

public good producers occupy – and are locally adapted to – different niches. In this scenario, the

'asocial' type would occur in microhabitats where social traits are not required or where

coordinated actions among cells are difficult due to low bacterial densities (Kümmerli et al. 2014;

Butaitė et al. 2018). Conversely, the 'social' type should mainly occur in microhabitats supporting

higher bacterial densities and/or featuring conditions under which cooperation is beneficial.

However, we believe that such a clear-cut segregation of social profiles according to niches is

unlikely because pseudomonads are typically motile (Raaijmakers et al. 2010), and because

bacteria can occur at high density in both terrestrial (Whitman et al. 1998) and aquatic habitats

(where they aggregate locally on particles (Cordero et al. 2012)). Alternatively, correlated trait

expression could be the result of evolution driven by social interactions involving the exploitation

and/or cross-use of public goods among strains (Diggle et al. 2007; Nadell et al. 2009; Van Gestel

et al. 2014; Butaitė et al. 2017).

Figure 3 | Correlation patterns between social traits. Shown are the pairwise correlations between social

traits of 315 natural *Pseudomonas* strains from eight soil (in yellow) and eight pond (in blue) communities.

Solid and dashed lines depict significant and non-significant relationships, respectively. Bands in habitat-

specific coloration as well as grey bands are 95 % confidence intervals derived from linear models and

represent, respectively, habitat-specific correlations and correlations that are consistent across habitats.

The evolution of exploitation and cross-use of public goods at the community level can be

conceptualized within the framework of the so-called Black Queen Hypothesis (BQH) (Morris et

al. 2012). This hypothesis has been coined to explain the evolution of metabolic dependencies

among free-living microbes. It predicts that the loss of a metabolic function is selectively favoured

at the individual level as long as the function is costly, still accessible via other community

members, and available in sufficient amounts to support growth. Such conditions should

promote the evolution of two distinct phenotypes: so-called 'beneficiaries' that are deficient for

the production of one or multiple public goods, and so-called 'helpers' that produce these goods

for everyone in the community. The BQH predicts that the likelihood to lose a function

accelerates for strains that have already lost a function before, thereby passively locking

producing strains into their role as helpers (Morris et al. 2012). The observed co-existence of the

'social' and 'asocial' types in our *Pseudomonas* communities is well in line with this scenario.

Moreover, differences in linkage patterns among community members could explain why certain

strains become 'helpers', whereas others become 'beneficiaries'. As discussed above, positive

linkage of traits could lead to the simultaneous loss of multiple traits, and thus increase the

chances of becoming a 'beneficiary'. Conversely, no or negative linkage, but also positive linkage

involving fewer traits could increase the likelihood of strains being locked into the 'helper'-

function, as the simultaneous loss of multiple functions is impeded.

If social interactions drive the evolution of the 'social' and 'asocial' strain types, their co-

existence can be explained in multiple ways. In the case of competitive interactions, where

'asocial' types exploit the 'social' types, public goods cooperation and strain co-occurrence could

be maintained by negative frequency-dependent selection (Ross-Gillespie et al. 2007). Similar to

the BQH, the concept of frequency-dependent selection applies if the social traits are essential

and a certain amount of public goods is required for community survival. In this case cheaters

might only experience high relative fitness when rare, but not when common, leading to an

equilibrium frequency at which the two strain types can co-exist. Alternatively, antagonistic co-

evolution between the two types could explain the co-existence of asocial and social types

(Kümmerli et al. 2015; O'Brien et al. 2017). For instance, increased biofilm formation in strains

that highly invest in proteases and pyoverdine could be a direct response to cheating. This is

because biofilm formation induces spatial segregation between cell types (Van Gestel et al. 2014;

Irie et al. 2017) and can thus prevent cheaters from efficiently exploiting other public goods

(Drescher et al. 2014). Similarly, the increased toxicity of strains producing high amounts of

proteases and pyoverdine could represent an adaptation to selectively target asocial types, which

show lower toxin production and might thus be more susceptible to toxins (Jousset et al. 2009;

Wang et al. 2015). Finally, cheating 'asocial' bacteria might be constrained by pleiotropic effects

resulting from trait linkage. For example, it was shown that cheating on pyoverdine or proteases

can compromise the ability of *Pseudomonas* bacteria to, respectively, form biofilms or defend

themselves against predation (Harrison and Buckling 2009; Jousset et al. 2009).

In the case of commensalistic or mutualistic interactions, strain co-existence could be

favoured if a certain amount of public goods is required to support the community (Morris et al.

2012), or if strain-specific public-good specialization promotes division of labour between strains

via the exchange of public goods (Dragoš et al. 2018). Such interactions seem, however, relatively

rare in our communities. Commensalistic interactions require that the utilization of public goods

by the 'asocial' type does not affect the 'social' type's fitness (Morris et al. 2012) – an assumption

that is unlikely to be met because our isolates are generally closely related and therefore likely

have overlapping resource requirements. Mutualistic interactions involving the exchange of

different public goods require strains with specialised trait repertoires. However, the absence of

negative correlations among public good traits suggests that such specialized mosaic producers

are uncommon in our communities, such that cross-feeding scenarios, similar to those described

in laboratory systems (Driscoll et al. 2011; Pande et al. 2014; Kim et al. 2016; Dragoš et al. 2018),

might be rare among our natural isolates. Our findings are in line with other studies showing that

competitive interactions prevail in natural bacterial communities (Foster and Bell 2012), and that

mutualistic interactions only evolve under limited conditions (Oliveira et al. 2014).

Our observation that there are habitat-specific correlations suggest that trait linkage can

change in response to different environmental selection pressures. Changes in the local

environment, for instance when a soil strain is washed into a pond, could thus rapidly promote

the breaking of existing but superfluous linkage patterns or select for new ones that are beneficial

in the new habitat. The moderate phlyogenetic signals we detected are in line with this notion,

as they suggest that the expression levels of social traits can evolve relatively flexibly. In this

context, the overall higher phylogenetic signal in soil as compared to freshwater communities

could indicate faster and more flexible evolution of public good repertoires in ponds.

In conclusion, our findings indicate that social trait linkage (i) is pervasive in natural

Pseudomonas communities, (ii) commonly promotes competitive rather than commensalistic or

mutualistic interactions, and (iii) underlies the predominance of strains lying on a continuum

between two main strain types – a 'social' type that is proficient in expressing multiple social

traits, and an 'asocial' type that generally shows low investment levels into most of the social

traits. While our study solely focuses on pseudomonads, it would be interesting to see whether

the same patterns of trait linkage occur beyond the genus boundary, especially for traits such as

proteases and toxins, which have more broad-spectrum effects than pyoverdine, which is a

characteristic trait of the fluorescent pseudomonads. Moreover, while our use of standardized

laboratory assays allowed us to assess the strains' genetic potential to produce specific public

goods, we do not know to which extent they would do so under natural conditions, and whether

strains can plastically adjust public goods production in response to changes in environmental or

social conditions. Our work therefore only provide a first glimpse of the complexity of microbial

social life in natural communities. This glimpse, however, suggests that social trait linkage could

play an important role in explaining community composition and functioning.

Data Accessability. The dataset supporting this article has been uploaded as part of the

supplementary material, and will be made available in the Dryad Repository upon acceptance.

Competing interests. We have no competing interests.

Authors' contributions. JK and RK designed the experiments; JK and MALC conducted the

experiments; JK performed the statistical analyses; JK, MALC and RK wrote the paper. All authors

gave final approval for publication.

Acknowledgements. We thank Elena Butaitė for collecting the natural strains, and Stefan Wyder

for help with the phylogenetic analysis.

Funding. This work was funded by the Swiss National Science Foundation (grant no. 165835 to

RK), the European Research Council (grant no. 681295 to RK), the Forschungskredit of the

University of Zürich (FK-17-111 to JK) and the German Science Foundation (KR 5017/2-1 to JK).

REFERENCES

Andersen, S. B., R. L. Marvig, S. Molin, H. Krogh Johansen, and A. S. Griffin. 2015. Long-term social dynamics drive loss of function in pathogenic bacteria. Proc. Natl. Acad. Sci. U. S. A. 112:201508324.

Bachmann, H., F. J. Bruggeman, D. Molenaar, F. Branco dos Santos, and B. Teusink. 2016. Public goods and metabolic strategies. Curr. Opin. Microbiol. 31:109–115.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B 57:289–300.

Bishara, A. J., and J. B. Hittner. 2012. Testing the significance of a correlation with nonnormal data: Comparison of Pearson, Spearman, transformation, and resampling approaches. Psychol. Methods 17:399–417.

Blomberg, S. P., T. G. Jr, and A. R. Ives. 2003. Testing for phylogenetic signal in comparative data: Behavioral traits are more labile. Evolution (N. Y). 57:717–745.

Bodour, A. A., and R. M. Miller-Maier. 1998. Application of a modified drop-collapse technique for surfactant quantitation and screening of biosurfactant-producing microorganisms. J. Microbiol. Methods 32:273–280.

Bruce, J. B., G. A. Cooper, H. Chabas, S. A. West, and A. S. Griffin. 2017. Cheating and resistance to cheating in natural populations of the bacterium *Pseudomonas fluorescens*. Evolution (N. Y). 71:2484–2495.

Butaitė, E., M. Baumgartner, S. Wyder, and R. Kümmerli. 2017. Siderophore cheating and cheating resistance shape competition for iron in soil and freshwater *Pseudomonas* communities. Nat. Commun. 8:414.

Butaitė, E., J. Kramer, S. Wyder, and R. Kümmerli. 2018. Environmental determinants of

pyoverdine production, exploitation and competition in natural *Pseudomonas*

communities. Environ. Microbiol. 263004.

Cordero, O. X., L.-A. Ventouras, E. F. DeLong, and M. F. Polz. 2012. Public good dynamics drive

evolution of iron acquisition strategies in natural bacterioplankton populations. Proc. Natl.

Acad. Sci. 109:20059–20064.

Dandekar, A. A., S. Chugani, and E. P. Greenberg. 2012. Bacterial Quorum Sensing and

Metabolic Incentives to Cooperate. Science 338:264–266.

Diggle, S. P., A. S. Griffin, G. S. Campbell, and S. A. West. 2007. Cooperation and conflict in

quorum-sensing bacterial populations. Nature 450:411–414.

Dragoš, A., H. Kiesewalter, M. Martin, C. Y. Hsu, R. Hartmann, T. Wechsler, C. Eriksen, S. Brix, K.

Drescher, N. Stanley-Wall, R. Kümmerli, and Á. T. Kovács. 2018. Division of labor during

biofilm matrix production. Curr. Biol. 1–11.

Drescher, K., C. D. Nadell, H. A. Stone, N. S. Wingreen, and B. L. Bassler. 2014. Solutions to the

Public Goods Dilemma in Bacterial Biofilms. Curr. Biol. 24:50–55.

Driscoll, W. W., J. W. Pepper, L. S. Pierson, and E. A. Pierson. 2011. Spontaneous Gac Mutants of

Pseudomonas Biological Control Strains: Cheaters or Mutualists? Appl. Environ. Microbiol.

77:7227–7235.

Dumas, Z., A. Ross-Gillespie, and R. Kümmerli. 2013. Switching between apparently redundant

iron-uptake mechanisms benefits bacteria in changeable environments. Proc. R. Soc. B

Biol. Sci. 280:20131055.

Foster, K. R. 2010. Social behaviour in microorganisms. Pp. 331–357 *in* T. Székely, A. J. Moore,

421 and J. Komdeur, eds. Social behaviour: Genes, ecology and evolution. Cambridge

422 University Press, Cambridge.

423 Foster, K. R., and T. Bell. 2012. Competition, not cooperation, dominates interactions among

424 culturable microbial species. Curr. Biol. 22:1845–1850.

425 Granato, E. T., C. Ziegenhain, R. L. Marvig, and R. Kümmerli. 2018. Low spatial structure and

426 selection against secreted virulence factors attenuates pathogenicity in *Pseudomonas*

427 *aeruginosa*. ISME J., doi: 10.1038/s41396-018-0231-9.

428 Griffin, A. S., S. A. West, and A. Buckling. 2004. Cooperation and competition in pathogenic

429 bacteria. Nature 430:1024–1027.

430 Harrison, F., and A. Buckling. 2009. Siderophore production and biofilm formation as linked

431 social traits. ISME J. 3:632–634.

432 Hesse, E., S. O'Brien, N. Tromas, F. Bayer, A. M. Luján, E. M. van Veen, D. J. Hodgson, and A.

433 Buckling. 2018. Ecological selection of siderophore-producing microbial taxa in response to

434 heavy metal contamination. Ecol. Lett. 21:117–127.

435 Irie, Y., A. E. L. Roberts, K. N. Kragh, V. D. Gordon, J. Hutchison, R. J. Allen, G. Melaugh, T.

436 Bjarnsholt, S. A. West, and S. P. Diggle. 2017. The *Pseudomonas aeruginosa* PSL

437 polysaccharide is a social but noncheatable trait in biofilms. MBio 8:1–13.

438 Jimenez, P. N., G. Koch, J. A. Thompson, K. B. Xavier, R. H. Cool, and W. J. Quax. 2012. The

439 multiple signaling systems regulating virulence in *Pseudomonas aeruginosa*. Microbiol.

440 Mol. Biol. Rev. 76:46–65.

441 Jousset, A. 2012. Ecological and evolutive implications of bacterial defences against predators.

442 Environ. Microbiol. 14:1830–1843.

Jousset, A., N. Eisenhauer, E. Materne, and S. Scheu. 2013. Evolutionary history predicts the

stability of cooperation in microbial communities. Nat. Commun. 4:2573.

Jousset, A., L. Rochat, M. Péchy-Tarr, C. Keel, S. Scheu, and M. Bonkowski. 2009. Predators

promote defence of rhizosphere bacterial populations by selective feeding on non-toxic

cheaters. ISME J. 3:666–674.

Kearns, D. B. 2010. A field guide to bacterial swarming motility. Nat. Rev. Microbiol. 8:634–44.

Kim, W., S. B. Levy, and K. R. Foster. 2016. Rapid radiation in bacteria leads to a division of

labour. Nat. Commun. 7:1–10.

Kümmerli, R., N. Jiricny, L. S. Clarke, S. A. West, and A. S. Griffin. 2009. Phenotypic plasticity of a

cooperative behaviour in bacteria. J. Evol. Biol. 22:589–598.

Kümmerli, R., L. A. Santorelli, E. T. Granato, Z. Dumas, A. Dobay, A. S. Griffin, and S. A. West.

2015. Co-evolutionary dynamics between public good producers and cheats in the

bacterium *Pseudomonas aeruginosa*. J. Evol. Biol. 28:2264–2274.

Kümmerli, R., K. T. Schiessl, T. Waldvogel, K. Mcneill, and M. Ackermann. 2014. Habitat

structure and the evolution of diffusible siderophores in bacteria. Ecol. Lett. 17:1536–

1544.

Lamont, I. L., P. A. Beare, U. Ochsner, A. I. Vasil, and M. L. Vasil. 2002. Siderophore-mediated

signaling regulates virulence factor production in *Pseudomonas aeruginosa*. Proc. Natl.

Acad. Sci. U. S. A. 99:7072–7077.

Leinweber, A., M. Weigert, and R. Kümmerli. 2018. The bacterium *Pseudomonas aeruginosa*

senses and gradually responds to inter-specific competition for iron. Evolution (N. Y). 1–27.

Letunic, I., and P. Bork. 2016. Interactive tree of life (iTOL) v3: an online tool for the display and

annotation of phylogenetic and other trees. Nucleic Acids Res. 44:W242–W245.

Morris, J. J., R. E. Lenski, and E. R. Zinser. 2012. The Black Queen Hypothesis : Evolution of

Dependencies through Adaptative Gene Loss. MBio 3:1–7.

Mulet, M., A. Bennasar, J. Lalucat, and E. García-Valdés. 2009. An rpoD-based PCR procedure

for the identification of *Pseudomonas* species and for their detection in environmental

samples. Mol. Cell. Probes 23:140–147.

Nadell, C. D., J. B. Xavier, and K. R. Foster. 2009. The sociobiology of biofilms. FEMS Microbiol.

Rev. 33:206–224.

O'Brien, S., A. M. Luján, S. Paterson, M. A. Cant, and A. Buckling. 2017. Adaptation to public

goods cheats in *Pseudomonas aeruginosa*. Proc. R. Soc. B Biol. Sci. 284.

Oliveira, N. M., R. Niehus, and K. R. Foster. 2014. Evolutionary limits to cooperation in microbial

communities. Proc. Natl. Acad. Sci. U. S. A. 111:201412673.

Özkaya, Ö., K. B. Xavier, F. Dionisio, and R. Balbontín. 2017. Maintenance of microbial

cooperation mediated by public goods in single- and multiple-trait scenarios. J. Bacteriol.

199:1–14.

Pande, S., H. Merker, K. Bohl, M. Reichelt, S. Schuster, L. F. De Figueiredo, C. Kaleta, and C. Kost.

2014. Fitness and stability of obligate cross-feeding interactions that emerge upon gene

loss in bacteria. ISME J. 8:953–962.

Pollitt, E. J. G., S. A. West, S. A. Crusz, M. N. Burton-Chellew, and S. P. Diggle. 2014.

Cooperation, quorum sensing, and evolution of virulence in *Staphylococcus aureus*. Infect.

Immun. 82:1045–1051.

Raaijmakers, J. M., I. de Bruijn, O. Nybroe, and M. Ongena. 2010. Natural functions of

lipopeptides from Bacillus and Pseudomonas: More than surfactants and antibiotics. FEMS

Microbiol. Rev. 34:1037–1062.

Rakoff-Nahoum, S., K. R. Foster, and L. E. Comstock. 2016. The evolution of cooperation within

the gut microbiota. Nature 533:255–259.

Ross-Gillespie, A., Z. Dumas, and R. Kümmerli. 2015. Evolutionary dynamics of interlinked public

goods traits: An experimental study of siderophore production in *Pseudomonas*

aeruginosa. J. Evol. Biol. 28:29–39.

Ross-Gillespie, A., A. Gardner, S. A. West, and A. S. Griffin. 2007. Frequency dependence and

cooperation: theory and a test with bacteria. Am. Nat. 170:331–342.

Rumbaugh, K. P., S. P. Diggle, C. M. Watters, A. Ross-Gillespie, A. S. Griffin, and S. A. West. 2009.

Quorum Sensing and the Social Evolution of Bacterial Virulence. Curr. Biol. 19:341–345.

Savoia, D., and M. Zucca. 2007. Clinical and Environmental Burkholderia Strains: Biofilm

Production and Intracellular Survival. Curr. Microbiol. 54:440–444.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of image

analysis. Nat. Methods 9:671–675.

van Gestel, J., H. Vlamakis, and R. Kolter. 2015. From Cell Differentiation to Cell Collectives:

Bacillus subtilis Uses Division of Labor to Migrate. PLoS Biol. 13:1–29.

Van Gestel, J., F. J. Weissing, O. P. Kuipers, and Á. T. Kovács. 2014. Density of founder cells

affects spatial pattern formation and cooperation in *Bacillus subtilis* biofilms. ISME J.

8:2069–2079.

Wang, M., A. L. Schaefer, A. A. Dandekar, and E. P. Greenberg. 2015. Quorum sensing and

policing of *Pseudomonas aeruginosa* social cheaters. Proc. Natl. Acad. Sci. 112:2187–2191.

West, S. A., S. P. Diggle, A. Buckling, A. Gardner, and A. S. Griffin. 2007. The Social Lives of

Microbes. Annu. Rev. Ecol. Evol. Syst. 38:53–77.

West, S. A., A. S. Griffin, A. Gardner, and S. P. Diggle. 2006. Social evolution theory for

microorganisms. Nat. Rev. Microbiol. 4:597–607.

Whitman, W. B., D. C. Coleman, and W. J. Wiebe. 1998. Prokaryotes: The unseen majority. Proc.

Natl. Acad. Sci. U. S. A. 95:6578–6583.

Wilder, C. N., S. P. Diggle, and M. Schuster. 2011. Cooperation and cheating in Pseudomonas

aeruginosa: the roles of the las, rhl and pqs quorum-sensing systems. ISME J. 5:1332–1343.

Xavier, J. B., W. Kim, and K. R. Foster. 2011. A molecular mechanism that stabilizes cooperative

secretions in Pseudomonas aeruginosa. Mol. Microbiol. 79:166–179.