1

The (p)ppGpp-binding GTPase Era promotes rRNA processing and cold shock survival in *Staphylococcus aureus*

Short title: Era promotes rRNA processing and bacterial survival

Alison Wood, Sophie E. Irving, Daniel J. Bennison and Rebecca M. Corrigan*

The Florey Institute, Department of Molecular Biology and Biotechnology, University of Sheffield, Sheffield, United Kingdom

* Correspondence: Email: r.corrigan@sheffield.ac.uk

Abstract

Ribosome assembly cofactors are widely conserved across all domains of life. One such group, the ribosome-associated GTPases (RA-GTPase), act as molecular switches to coordinate ribosome assembly. We previously identified the *Staphylococcus aureus* RA-GTPase Era as a target for the stringent response alarmone (p)ppGpp, although the function of Era in ribosome assembly is unclear. Era is highly conserved throughout the bacterial kingdom and is essential in many species. Here we show that Era is not essential in *S. aureus* but is important for 30S ribosomal subunit assembly. Protein interaction studies reveal that Era interacts with the endonuclease YbeY and the DEAD-box RNA helicase CshA. We determine that both Era and CshA are required for growth at suboptimal temperatures, virulence and 16S rRNA processing. Era and CshA also form direct interactions with the (p)ppGpp synthetase Rel_{SA}, with Rel_{SA} positively impacting the GTPase activity of Era. We propose that Era acts to direct enzymes involved in rRNA maturation and ribosome subunit assembly to their site of action, an activity that is regulated by components of the stringent response.

Author summary

The bacterial ribosome is an essential cellular component and as such is the target for a number of currently used antimicrobials. Correct assembly of this complex macromolecule requires a number of accessory enzymes, the functions of which are poorly characterised. Here we examine the function of Era, a GTPase enzyme involved in 30S ribosomal subunit biogenesis in the important human pathogen *S. aureus*. We uncover that Era is not an essential enzyme in *S. aureus*, as it is in many other species, but is important for correct ribosome assembly. In a bid to uncover the cellular function of this enzyme, we identify a number of protein interaction partners with roles in ribosomal RNA maturation and degradation, supporting the idea that Era acts as an intermediary protein facilitating ribosomal biogenesis. We also uncover a link between Era and the (p)ppGpp synthetase Rel_{SA}, revealing an additional level of control of rRNA levels by the stringent response. With this study we elaborate on the functions of GTPases in ribosomal assembly, processes that are controlled at multiple points by the stringent response.

Introduction

Ribosomes are macromolecular machines responsible for the synthesis of proteins in all living cells. In bacteria these complexes consist of two subunits, with the large 50S subunit containing 33 large proteins (L1-36) and two rRNAs, and the 30S small subunit containing 21 small proteins (S1-21) and the 16S rRNA. As such, assembly of the ribosome is a tightly regulated process, with correct maturation requiring the help of assembly cofactors, one class of which are the P-loop ribosome-associated GTPases (RA-GTPase). These enzymes are widely conserved across all domains of life and act as molecular switches, cycling between inactive GDP-bound, and active, effector-binding GTP-bound states. Within the P-loop GTPase class lies the Era family (*Escherichia coli* Ras-like protein), which is characterised by the presence of a distinct derivative of a KH domain [1]. Era, the protein after which this family is named, is highly conserved throughout the bacterial kingdom, although is missing in *Chlamydia* and mycobacterial species [1]. This GTPase is essential in *E. coli* [2-5], *Salmonella* Typhimurium [6] and in strains of *Bacillus subtilis* [7, 8]. The first indication that Era is involved in ribosomal biogenesis came from the ability of the 16S rRNA methyltransferase KsgA to suppress a cold-sensitive phenotype of an Era E200K mutation [9]. While important for ribosomal maturation, additional phenotypic defects associated with depleted cellular levels of Era include cell cycle control and chromosome segregation, as well as carbon and nitrogen metabolism [10-12].

Era is composed of two domains, an N-terminal GTPase domain and a C-terminal RNA-binding K-homology domain [13]. A cryo-electron micrograph structure of Era in complex with the 30S subunit of the ribosome reveals that Era binds into the same pocket as small subunit protein S1 [14]. In the absence of S1, Era interacts with proteins S2, S7, S11 and S18, as well as with a number of helices of the

16S rRNA. In addition, Era interacts with h45 and nearby residues 1530-1539

(GAUCACCUCC) in the 3' minor domain of the 16S rRNA via its CTD region [14,

15]. These residues include the anti-Shine Dalgarno sequence, critical for the

formation of the 30S preinitiation complex. Including Era in *in vitro* reconstitutions of

the 30S ribosome promotes the incorporation of several late-stage ribosomal proteins

for the RNA [16, 17]. Consequently, it has been proposed that Era functions as a

checkpoint protein, and that by binding to the 16S rRNA the formation of the initiation

complex is prevented until the appropriate time [14]. In addition to interacting with

ribosomal proteins, Era has also been reported to interact with a number of proteins

involved in 16S rRNA maturation. One of these, YbeY, is an endonuclease required in

E. coli for the maturation of the 3' end of the 16S rRNA [18, 19]. It is proposed that

the binding of YbeY to Era and S11 guide the endonuclease to its site of action [18].

The stringent response is a bacterial signalling system used by bacteria to cope

with a variety of environmental stresses, the best characterised of which is nutrient

deprivation. The opportunistic pathogen *Staphylococcus aureus* contains three

enzymes, Rel_{SA}, RelP and RelQ, which upon sensing a stress, synthesise the

nucleotides guanosine tetra- and pentaphosphate ((p)ppGpp) [20, 21]. Once produced,

this alarmone controls cellular responses to aid survival. Our previous work identified

Era, as well as three other GTPase enzymes from *S. aureus*, as target proteins for

(p)ppGpp and demonstrated that the production of (p)ppGpp has a negative impact on

mature 70S assembly [22]. Here, we examine the role of Era as an enzyme required for

ribosome biogenesis. Unlike in *E. coli*, this GTPase is not essential for the growth of *S.*

aureus, however mutant cells are defective in 30S subunit maturation. We identify the

S. aureus endonuclease YbeY and the DEAD-box RNA helicase CshA as interaction

partners for Era, and show that both Era and CshA are crucial for rRNA homeostasis

and growth at low temperatures. We additionally demonstrate that both Era and CshA

interact with the (p)ppGpp synthetase Rel$_{SA}$, and that cellular rRNA processing is

controlled in a stringent response-dependent manner. With this, we propose that Era is

a protein that facilitates the interactions between a number of rRNA processing and

degrading enzymes and the 30S subunit/16S rRNA, and show that under stress

conditions the stringent response is important for this process.

Results

Era is not essential in *S. aureus* but is required for normal growth and ribosome

assembly

Previous reports show that *era* is an essential gene in multiple bacterial species [3, 6].

In agreement with this, transposon insertions in this gene have not been identified in a

number of published *S. aureus* transposon libraries [23-25]. Closer inspection of the

transposon insertion hits within the *era* operon obtained in the Nebraska *S. aureus*

mutant library [23] reveals insertions in *ybeZ*, a gene of unknown function, and at the

3' end of the diacylglycerol kinase *dgkA* but not within *ybeY*, *cdd*, *era* or *recO*,

suggestive of an additional promoter upstream of *era* and *cdd* (Fig 1a). To rule out

polar effects and examine whether *era* in isolation is essential, we replaced the entire

coding sequence with the Tet-encoding *tetAM* open reading frame in the presence of

the anhydrotetracycline (Atet)-inducible covering plasmid pCN55iTET-*era*. Following

deletion of the chromosomal copy of *era*, we attempted to phage transduce this

mutation into *S. aureus* strains containing the empty vector pCN55iTET or the

complementing plasmid pCN55iTET-*era*. Transduction efficiencies were similar with

both recipients, albeit smaller colony sizes were observed in the absence of *era*,

indicating that this gene is not essential in *S. aureus*. We next transduced the *era*

deletion into the community-associated methicillin resistant *S. aureus* strain LAC* to

rule out secondary mutations and used this strain for further studies. Analysis of

growth rates revealed that an *era* mutant strain, while viable, does have a significant

growth defect, which can be complemented by the expression of Era from

pCN55iTET-*era* (Fig 1b).

A role for Era as a ribosomal subunit assembly cofactor has been reported in

both *E. coli* and *B. subtilis*, where depletion leads to a decrease in 70S ribosomes and

an accumulation of individual 50S and 30S subunits [26, 27]. We reasoned that the

growth defect observed in *S. aureus* may be due to defects in ribosome assembly. To

investigate this in the context of a complete *era* deletion, we analysed the cellular

ribosomal content of wildtype, mutant and complemented strains by sucrose density

centrifugation. This revealed that the *era* mutant strain contained fewer polysomes

with a concurrent build-up of 50S subunits, a defect that was reversed in the presence

of the complementing plasmid (Fig 1c). As cryo-electron microscopy has shown Era

interacting with the 30S subunit [14], an excess of free 50S subunits may be, in and of

itself, an indication that there is a defect in small subunit biogenesis, leading to a

build-up in free 50S. While the peak for polysomes disappeared, the levels of 70S

ribosomes remained similar between the wildtype and mutant, however it is not known

if these 70S ribosomes are fully functional. We reasoned that this defect in ribosome

assembly might make the *era* mutant strain more susceptible to ribosome-targeting

antibiotics. In agreement with this, we observed that the minimum inhibitory

concentration (MIC) for the Δ*era* strain decreased 2-4-fold when exposed to the 30S-

targeting antibiotic spectinomycin (Fig S1a). However, the MIC was unaffected by the

50S targeting antibiotic chloramphenicol (Fig S1b). Together this indicates that while

Era is not an essential protein in *S. aureus*, it is important for optimal growth and

ribosome maturation.

Uncovering Era protein interaction dynamics in a native background

In *S. aureus,* Era is encoded in an operon with five other genes, one of which encodes

YbeY (Fig 1a). YbeY is an endoribonuclease implicated in the maturation of the 3'

terminus of 16S rRNA in *E. coli*, as well as a quality control checkpoint protein that

together with RNase R is involved in eliminating defective 70S ribosomes [19, 28].

Era interacts with YbeY in *E. coli* [18], although these two genes are not encoded in

the same operon in that organism (Fig 1a). To determine whether Era from *S. aureus*

interacts with YbeY, and/or any other proteins encoded in the *S. aureus era* operon,

we first used a bacterial two-hybrid approach, heterologously expressing Era in

combination with YbeZ, YbeY, DgkA, Cdd or RecO in *E. coli*. Using this approach,

we observed that Era potentially interacts with YbeY and YbeZ (Fig 2a).

In order to look at these interactions natively in an *S. aureus* background, we

adapted a split-luciferase system recently developed to analyse protein-protein

interactions in *Clostridium difficile* for use in *S. aureus* (Fig 2b) [29]. The *era* gene, in

combination with itself and each of the operon genes, were cloned into the split-

luciferase plasmid pAP118 and introduced into *S. aureus* strain LAC*. Induction of

protein expression with Atet revealed a strong positive interaction for Era with itself

(Fig 2c), indicating that Era can form dimers. Dimerisation of Era has previously been

observed while solving the crystal structure of the *E. coli* protein [30], and so this

interaction confirms the functionality of the split-luciferase system in *S. aureus*. In

addition, a significant interaction was also observed for Era and YbeY (2 x above the

control), although not for YbeZ (1.40 x above the control) (Fig 2c). In agreement with

the bacterial two-hybrid results, no interactions occurred between Era and DgkA, Cdd

or RecO. A decrease in luminescence below the Era control pAP256-era_{SB}-LB occurred

when co-expressing Era with DgkA and RecO. The reason for this is unclear but could

be due to toxicity upon over-expression of a second copy of these genes from a multi-

copy plasmid. To confirm the validity of the Era-YbeY interaction *in vitro*, we

performed pulldown assays using glutathione beads coupled to either GST or GST-

tagged Era and incubated them with His-tagged YbeY (Fig 2d). Here we observed

low-level cross reactivity of YbeY with the gluthatione beads that could not be

reduced despite increased washes. However, His-YbeY was pulled down significantly

more in the presence of Era (Fig 2d). Altogether the bacterial two-hybrid, split

luciferase and affinity pulldown assays confirm an interaction between Era and the

endonuclease YbeY from *S. aureus*. With this data we have adapted a luciferase

system for confirmation of protein-protein interactions in a native *S. aureus*

background.

Era interacts with the DEAD-box RNA helicase CshA

We wished to shed further light on the role played by Era in the cell and sought to

identify unknown interaction partners using a genome-wide bacterial two-hybrid

screen. A library of *S. aureus* genomic DNA fragments cloned into pUT18C was

screened against pKNT25-*era*. Of the 17 hits obtained, 14 contained fragments

mapping to *cshA*, a gene encoding a DEAD-box RNA helicase. CshA functions in

mRNA protection and RNA decay in *S. aureus* [31, 32], while the homologue CsdA

from *E. coli* is involved in the cold shock degradosome [33]. To confirm the

interaction between Era and CshA, full-length *cshA* was cloned into both pUT18 and

pUT18C. Upon co-transformation, interactions occurred between CshA-T18 and both

T25-Era and Era-T25 (Fig 3a).

To assess this interaction *in vitro*, we performed pulldown assays using

glutathione beads coupled to either GST or GST-tagged CshA and incubated them

with His-tagged Era, revealing that while His-Era did not interact with the control

GST protein, it was pulled down in the presence of CshA (Fig 3b). Finally, to analyse

and confirm this interaction in the context of the native host, we used the split

luciferase assay (Fig 2b). Assaying for luminescence revealed a 4-fold increase in

luciferase activity upon co-expression of Era and CshA (Fig 3c), confirming the

identification of a novel Era interaction partner.

The N-terminal region of CshA is crucial for interacting with Era

CshA is a DEAD-box RNA helicase with an N-terminal helicase core containing two

RecA-like domains and a disordered C-terminal (CT) region involved in RNA binding

(Fig 3d) [32]. This protein is capable of unwinding both double stranded RNA and

RNA-DNA hybrids and is required for both the stabilisation and the degradation of

mRNA, the latter of which occurs via interactions of the CT region with components

of the RNA degradosome [31, 32].

To determine which regions of both Era and CshA are important for

interacting, shorter domain constructs comprising only RecA1 (aa 1-221), RecA2 (aa

222-382), the entire core helicase domain (aa 1-382) or just containing the disordered

CT-region (aa 383-506) were cloned, together with Era, into the split luciferase vector

pAP118. Interactions with full-length Era were only apparent in the presence of the

full core helicase domain (aa 1-382: Fig 3e), indicating that the RNA-binding CT

region of CshA is dispensable for this interaction. To determine whether the GTPase

domain of Era is required for binding, a shorter construct comprising the N-terminal G

domain (aa 1-180) was cloned into pAP118 alongside full-length CshA. Luciferase

assays reveal that the G domain is sufficient for this interaction (Fig 3f).

Interactions between Era and CshA do not affect enzymatic activity

CshA is an RNA helicase, while Era has GTPase activity. To investigate whether the

interaction between these two proteins affects the enzymatic activity of either one, we

first performed helicase assays. A double stranded RNA oligo was incubated with each

protein singly or in combination. While CshA was able to unwind the dsRNA, the

addition of Era had no effect on its activity (Fig S2a). In addition, the GTPase activity

of Era was unaltered in the presence of either CshA or YbeY (Fig S2b and S2c),

indicating that while these proteins do interact, this binding has little effect on their

enzymatic functions. As previously proposed, we suggest that these interactions fit

with a role for Era as a guide for rRNA/ribosome maturation enzymes to their

substrates [14, 18].

CshA and Era are both required for growth at suboptimal temperatures,

virulence and rRNA processing

In *E. coli,* CshA has been linked to survival at low temperatures [34], while strains

with depleted levels of Era are sensitive to both cold and heat shock [9, 10]. To

examine the importance of these enzymes in *S. aureus*, we utilised deletion strains in

the *S. aureus* MRSA strain LAC*. While both the *S. aureus era* and *cshA* null mutants

showed significant growth defects at 37°C (Fig 1b and S3a), the growth of both was

severely compromised at 25°C (Fig 4a and S3b). This defect was enhanced in a double

Δ*era cshA* mutant, which failed to grow at 25°C even after 48 h (Fig 4a and S3b),

highlighting that these two proteins are essential for bacteria to cope with growth at

suboptimal temperatures.

Era is an RA-GTPase that, along with a second RA-GTPase RsgA, binds to the

30S ribosomal subunit [14, 35]. In *E. coli,* overexpression of Era can complement a

deletion of *rsgA* [36]. After confirming that expression of Era can complement a

deletion of *rsgA* in *S. aureus* (Fig S4a), we used cross-complementation of the Δ*era*

and Δ*cshA* mutations to investigate whether Era and CshA act in the same pathway

and can functionally complement each other. However, cross-complementation did not

improve growth at 25°C (Fig S4b and S4c), indicating that while these proteins

interact and bacteria lacking both cannot survive cold temperatures, they have separate

cellular functions. We next assessed the contribution of each gene to the virulence of

S. aureus using *Galleria mellonella* as an invertebrate model of systemic infection.

While the wildtype bacteria were fully virulent, both the *era* and *cshA* mutant strains

displayed reduced killing, with the combined double mutant being completely

avirulent (Fig 4b).

To understand why Era and CshA are so important for growth at cold

temperatures and for virulence, we sought to assess the impact of deleting these genes

on the cellular rRNA. We observed that RNA extracted from both the single Δ*era* and

double Δ*era cshA* strains grown at 37°C had a processing or degradation intermediate

migrating slightly below the 16S band (Fig 4c). At 25°C we also observed a defect for

the Δ*cshA* mutant strain and multiple additional bands were evident in the Δ*era cshA*

double mutant (Fig 4c), which could represent either processing intermediates from

the 16S, or degradation intermediates from both the 23S and the 16S. A build-up of

degradation products are expected in the *cshA* mutant due to its role in the RNA

degradosome. However, to determine whether processing defects could be occurring

in these mutant strains when grown at low temperatures, we performed qRT-PCR with

probes that either amplified internally to the 16S rRNA sequence, or either flanked the

5' or 3' processing junctions required to convert 17S into mature 16S rRNA (Fig 4d).

While the levels of 16S rRNA were not significantly different between wildtype and

mutants, large increases in unprocessed 5' and 3' rRNA were evident in the Δ*era* strain

(Fig 4d). As Era is known to interact with the 30S ribosomal subunit, this could

potentially be due to mis-localisation of endonuclease YbeY in the absence of Era. It is

therefore, possible that Era acts as an intermediary protein on the 30S ribosomal

subunit, enabling proteins such as the endonuclease YbeY, and the RNA-degradosome

helicase CshA to access the 17S rRNA, allowing for processing (maturation or

degradation as appropriate) of rRNA under normal and stress conditions, such as

growth at cold temperatures.

Rel$_{SA}$ interacts with both Era and CshA

Our previous work demonstrated that the GTPase activity of Era is inhibited by the

stringent response alarmone ppGpp [22]. To examine whether ppGpp also interacts

with CshA we performed DRaCALA assays, revealing that this nucleotide does not

interact with CshA (Fig S5a), nor does the affinity of ppGpp for Era alter in the

presence of CshA (K_d of 3.1 ± 0.4 μM in the absence versus 3.2 ± 0.3 μM in the

presence of CshA) (Fig S5b).

To investigate whether the stringent response has any additional points of

interaction with either Era and CshA, we examined whether one of the (p)ppGpp

synthetases might directly interact with either protein via bacterial two-hybrid. This

revealed that Rel$_{SA}$, but not RelP or RelQ, interacts with CshA and that Rel$_{SA}$ interacts

with Era (Fig 5a). To confirm this interaction, we first used affinity pulldown assays,

showing that although no significant interaction occurred with Era, CshA and Rel$_{SA}$

interacted quite strongly (Fig 5b). Following this, we again used the split luciferase

assay, which confirmed an interaction between Rel$_{SA}$ and CshA (Fig 5c) or Era (Fig

5d). It was interesting to note that the CshA-Rel$_{SA}$ interaction only occurred in the

CshA$_{SB}$-Rel$_{LB}$ and not the Rel$_{SB}$-CshA$_{LB}$ orientation, highlighting the importance of

creating both types of luciferase fusions.

Rel$_{SA}$ positively affects the GTPase activity of Era and is important for rRNA

processing

We wished to ascertain whether the interactions between Rel$_{SA}$ and Era impact the

activity of either enzyme. Recombinantly purified full-length Rel$_{SA}$ is known to be in

the synthetase-off/hydrolase-on conformation *in vitro* [37]. Using purified Rel$_{SA}$, we

determined that Era and CshA have no effect on the pppGpp hydrolase activity of this

enzyme (Fig S5c). However, upon examination of the GTPase activity of Era, we

observed significantly increased activity in the presence of Rel$_{SA}$ (Fig 6a).

Through Rel$_{SA}$, the stringent response appears to interact with components of the

ribosome and 16S rRNA processing machinery. To analyse the importance of the

stringent response for rRNA processing, we grew the wildtype *S. aureus* LAC* at

25°C, a condition under which we have shown that both Era and CshA are important

for viability, and induced the stringent response with mupirocin, an antibiotic that

inhibits the isoleucyl-tRNA synthetase mimicking amino acid deprivation [38]. We

again performed qRT-PCR with the probes as in Fig 4d that either amplified internally

to the 16S rRNA sequence, or either flanked the 5' or 3' processing junctions of the

17S rRNA. For the probe internal to the 16S rRNA we observed a significant decrease

in 16S rRNA in the presence of mupirocin (Fig 6b). This is expected, as it is known

that rRNA levels decrease upon activation of the stringent response. However, when

we compare the amount of unprocessed 5' or 3' sequences relative to the level of 16S

in each strain, we observed a significant increase in the amount of unprocessed leader

17S rRNA upon induction of (p)ppGpp (Fig 6c: p = 0.0031). We have previously

shown that ppGpp interacts with Era and that mature ribosome formation is inhibited

upon induction of the stringent response [22]. Here we propose that the stringent

response can also impact RA-GTPase function through direct interactions of Rel$_{SA}$

with both Era and CshA, influencing the GTPase activity of Era and 17S rRNA

processing.

Discussion

The synthesis of ribosomes and proteins consumes approximately 40% of the energy

within a growing bacterial cell [39]. Ribosomal assembly cofactors are, therefore, an

essential group of enzymes for coordinating biogenesis as efficiently as possible.

Members of this group include RNA helicases, rRNA and protein modification

enzymes, chaperones and RA-GTPases. *S. aureus* contains over 11 RA-GTPases, each

with potentially varying roles in the biogenesis of the 50S and 30S subunits, although

the precise functions of each in this process is unclear.

While the RA-GTPase Era has been extensively studied, its precise function in

the cell is unknown. Cryo-electron microscopy shows Era interacting with the 30S

ribosomal subunit, however this protein has also been implicated in numerous other

cellular processes [14]. Era-depleted *B. subtilis* or *E. coli* cells are elongated, with

defects in septum formation. These cells also contain diffuse nucleoid material,

implicating Era in cell division and chromosome segregation [5, 12]. In *B. subtilis,*

Era-depleted cells have defects in spore formation [7], while growth defects in strains

containing Era variants with reduced nucleotide binding abilities can be rescued by

truncating either *rpoN* or *ptsN* [11]. RpoN is required for nitrogen assimilation and

fixation, while PtsN is involved in sugar transport, suggesting that Era is also involved

in regulating carbon and nitrogen metabolism. Localisation studies have indicated that

this protein is present at both the membrane and in the cytoplasm [40], and so it has

been suggested that this enzyme cycles between the membrane and the ribosome in

response to cellular triggers. In addition to ribosomal biogenesis factors, Era has been

reported to interact with MazG, Ndk, Pk and YggG [41-43]. Ndk, a nucleoside

diphosphate kinase and Pk, a pyruvate kinase, from *Pseudomonas aeruginosa* were

both shown to form a complex with Era, further implicating Era in energy metabolism

[42]. MazG is a nucleoside triphosphate pyrophosphohydrolase, while YggG is a

membrane-associated heat shock protein. The significance of these interactions with

Era is unknown.

Era has also been implicated in helping cells cope with cold stress. Conditional

cold-sensitive mutants of Era have been constructed in *E. coli* [10] and these mutations

could be suppressed by the overexpression of the 16S rRNA methyltransferase KsgA,

providing one of the first links between Era and ribosome biogenesis [9]. In addition,

Era interacts with YbeY in *E. coli* [18]. YbeY is a universally conserved endonuclease

that, while dispensable in *E. coli*, is essential in *B. subtilis* and potentially in *S. aureus*,

given the lack of transposon mutants available [19, 25, 28]. This endonuclease is

required for the maturation of the 3' end of the 16S rRNA, and in strains depleted of

YbeY the 70S ribosomes are targeted for degradation with the help of RNase R [19,

28]. Era and YbeY are fused in a polypeptide in clostridial species, further

highlighting a link between these two proteins. Although not encoded in the same

operon as *era* in *E. coli*, this prompted us to examine the interactions of all other

proteins encoded in the *era* operon in *S. aureus* with Era (Fig 2). As in *E. coli*, YbeY

from *S. aureus* interacts with Era. YbeZ also interacted by bacterial two-hybrid but not

significantly using the quantitative luciferase assay (Fig 2). YbeZ has been shown to

interact with YbeY in *E. coli* [18]. This protein has an ATPase domain and an RNA

binding motif, and so is likely to also bind to the 17S rRNA and may form a complex

with Era and YbeY to aid in processing the 17S rRNA to the mature 16S form.

Here we use interaction studies to show that Era interacts with a protein

involved in survival at suboptimal temperatures, the DEAD-box RNA helicase CshA

(Fig 3). In *B. subtilis,* CshA is one of the most abundant RNA helicases produced at

low temperatures [44] and in *B. subtilis* and *E. coli* it has been implicated in multiple

processes, including ribosome biogenesis and interacting with components of the RNA

degradosome [32-34, 44]. In *S. aureus,* CshA has been linked to controlling the

turnover of mRNA in the cell [31]. Deletion of *cshA* results in the stabilisation of some

mRNA transcripts, such as the *spa* mRNA, but CshA also protects a number of other

mRNA and sRNA transcripts under stress conditions [31]. In addition, the CT region

of CshA is required for interactions with the degradosome [32]. Our analysis shows

that Era and CshA interact *in vitro* and when expressed heterologously in *E. coli* (Fig

3a and 3b). We additionally used an adapted split luciferase assay to show this

interaction natively in *S. aureus* (Fig 3c). Using this assay has also allowed us to

determine that the CT disordered region of CshA, which is important for binding to

mRNA and the degradosome [32], is dispensable for interacting with Era (Fig 3).

Using *cshA* and *era* deletion strains we show that both single mutants are

important for growth at cold temperatures, with a double mutant unable to survive at

25°C (Fig 4). By examining the rRNA of these strains we observe that the *era* mutant

strain has increased 17S unprocessed rRNA intermediates (Fig 4d). Additionally, a

double Δ*era cshA* mutant has multiple processing and degradation defects (Fig 4c). It

has been proposed that Era may act to guide processing enzymes to their site of action

[18]. In keeping with this, we have observed that these proteins have little effect on the

enzymatic activity of each other. Therefore, we propose a model whereby Era acts as

an intermediary protein, allowing proteins such as the endonuclease YbeY and the

RNA helicase CshA, and potentially others, access to their substrate rRNA. It has been

suggested that CshA helps RNase components of the degradosome access cleavage

sites that may be otherwise inaccessible [45], and so it is tempting to speculate that Era

is responsible for recruiting CshA to target misfolded ribosomal subunits for

degradation under stress conditions.

Cellular levels of rRNA are controlled by the stringent response [46]. Upon

activation of the stringent response, transcription from rRNA promoters is reduced

(Fig 6b), either by direct binding of ppGpp to the RNA polymerase in Gram negatives,

or by tight control of cellular GTP levels in Gram positives [47, 48]. Here we see an

additional level of regulation by the stringent response on rRNA. We show a direct

interaction between the (p)ppGpp synthetase Rel_{SA} and both CshA and Era (Fig 5),

and show that Rel_{SA} has a positive effect on the GTPase activity of Era (Fig 6a).

Additionally, we observed that the activation of the stringent response results in

increased processing defects in a wildtype strain grown at 25°C (Fig 6c). RA-GTPases

are reported to have increased association to the ribosome in the GTP-bound form.

Hydrolysis of GTP then acts as a signal to promote dissociation, presumably following

a maturation event [49, 50]. The increased GTPase activity of Era in the presence of

Rel_{SA} could promote premature dissociation of Era from the ribosome, leading to

processing defects (Fig 6c) and contributing to the stalled growth phenotype

synonymous with the stringent response. The precise dynamics of processing-enzyme

association under stress conditions requires further in-depth research but is beyond the

scope of this study. For instance, under what growth conditions is Rel_{SA} binding to Era

and CshA promoted and how does this affect the localisation of processing proteins to

the ribosome?

Taken together, we demonstrate a cellular function for Era as a protein

important for coordinating 30S ribosomal biogenesis in a stringent response-dependent

manner. It is possible that defects in ribosomal assembly lead to altered protein

translation, which may be the reason for the plethora of other phenotypes associated

with depletion of Era. As Era can complement a defect in RsgA (Fig S4a, [36]), as

well as RbfA [26], we propose a broader role for RA-GTPases as intermediary

proteins. It is now of interest to determine what processing events are coordinated by

the other RA-GTPases in *S. aureus*, as well as more broadly in other prokaryotes.

Materials and methods

Bacterial strains and culture conditions. *E. coli* strains were grown in Luria Bertani

broth (LB) and *S. aureus* strains in tryptic soy broth (TSB) at 37°C with aeration.

Strains and primers used are listed in Tables S1 and S2. Information on strain

construction is provided in SI Methods.

Luciferase assay. Overnight cultures of *S. aureus* strains were diluted to an OD_{600} of

0.05 and grown for 90 min in the presence of 100 ng/ml anhydrotetracycline (Atet).

Strains were normalised to an OD_{600} of 0.1 and luciferase activity measured according

to the Nano-Glo Luciferase Assay System Protocol (Promega).

Pull-down experiments and western blotting. Glutathione beads were washed 5 times in 1 x wash solution (25 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA, 0.5% Triton X-100). 1 μM of glutathione-S-transferase (GST), GST-CshA or GST-Era were coupled to the beads by incubating in 1 x wash buffer for 4 h at 4^{o}C. Unbound protein was removed by washing in 1 x wash solution between 5-10 times. Protein-bound beads were incubated with 1 μM His-tagged Era, His-YbeY or His-Rel$_{SA}$ in the presence of 1 x binding buffer (25 mM Tris pH 7.5, 150 mM NaCl, 5 mM MgCl$_2$, 0.5% Triton X-100) at 4^{o}C overnight. After washing, attached proteins were eluted from the beads with 50 μl elution buffer (25 mM Tris pH 8, 150 mM NaCl, 1 mM EDTA, 0.5% Triton X-100, 10 mM reduced glutathione). Samples were mixed 1:1 with 2 x SDS protein sample buffer. Aliquots were separated on 12% SDS-polyacrylamide gels and proteins subsequently transferred to PVDF membranes. Bound His-tagged proteins were detected using HRP-conjugated anti-His antibodies (Sigma) at a 1:500 dilution. Blots were developed by enhanced chemiluminescence and imaged using a ChemiDoc MP imager (Bio-Rad).

Construction of bacterial two-hybrid library. Genomic DNA of *S. aureus* was extracted and partially digested by incubation with Sau3AI at 37^{o}C for 20 min. Digested DNA was run on a 0.8% agarose gel and fragments of 500 to 1000 bp and 1000 to 3000 bp gel extracted and purified. This was repeated five times from separate genomic preps and DNA fragments pooled. The vector pUT18C was digested overnight at 37^{o}C with BamHI and dephosphorylated with antarctic phosphatase for 90 min. Genomic DNA fragments were ligated into the pUT18C linearized vector using T4 ligase, transformed into *E. coli* DH5α competent cells (New England Biolabs) and

plated onto LB agar with carbenicillin. Plates were scraped and transformant plasmids

isolated using the GeneJET plasmid purification kit (Thermo scientific).

Bacterial two-hybrid. Bacterial two hybrid plasmids containing genes of interest were

co-transformed into *E. coli* BTH101 cells, plated on LB agar containing 150 μg/ml

carbenicillin and 30 μg/ml kanamycin and incubated at 30°C overnight. Colonies were

isolated, grown overnight at 30°C in LB broth containing 0.5 mM IPTG and 5 μl

spotted onto LB agar containing 0.5 mM IPTG and 40 μg/ml X-gal. Plates were

incubated at 30°C for up to 48 h. In some cases, BTH101 transformants were

incubated overnight with 0.5 mM IPTG and spotted directly onto LB agar containing

IPTG, X-gal and appropriate antibiotics.

***Galleria mellonella* survival assays.** *S. aureus* strains were grown overnight in TSB,

pelleted at 17,000 x g, washed twice in sterile PBS and diluted to an OD_{600} of 0.1 (2-4

x 10^7 cfu/ml). Colony numbers were confirmed by dilution and plating on TSA. Final

larval stage *G. mellonella* weighing 250-350 mg (Live foods direct, Sheffield, UK)

were stored in the dark at 4°C and used within 7 days. 5 μl of *S. aureus* inoculum was

injected into the hemocoel through the last pro-leg using a Hamilton syringe 701 RN.

Larvae were incubated in sealed plastic petri dishes at 37°C and the number of

surviving individuals recorded daily.

RNA extraction. Strains of *S. aureus* were grown overnight at 37°C and diluted to an

OD_{600} of 0.05. Cultures were grown to an OD_{600} of 0.4 at either 37 or 25°C and

harvested. RNA was extracted using the RiboPure RNA Purification Kit (Invitrogen)

as per guidelines. RNA was visualised using a modified agarose gel containing 0.7%

agarose and 0.9% Synergel (Diversified Biotech) in 0.5 X TBE (44.5 mM Tris, 1 mM

EDTA pH 8, 44.5 mM boric acid) as per Wachi *et al.* 1999 [51]. RNA was visualised

using EvaGreen fluorescent nucleic acid dye (Biotium) on a ChemiDoc MP imager

(Bio-Rad).

RT-PCR. RNA was extracted as described above. Complementary DNA was

synthesised from 1.5 μg RNA with transcriptor reverse transcriptase (Sigma) and

random primers. RT-PCR was performed on 100 ng of cDNA in triplicate using

SYBR Green Jumpstart Taq readymix (Sigma). Primers RMC379/380 (16S internal),

RMC570/571 (leader) and RMC572/548 (trailer) were designed to amplify 200-220 bp

target regions in the 17S rRNA. The housekeeping gene *rho* was amplified with

primers RMC573/574. The *rho* transcript was chosen as it has been shown to be highly

stable under nutrient-deficient conditions under which the stringent response would be

induced [52]. Cycle threshold values were determined for 3 biological repeats in

duplicate. For each reaction, the ratio to *rho* transcript number was calculated as

follows: $2^{-(Ct\ target\ -\ Ct\ rho)}$.

Growth curves. *S. aureus* strains were grown overnight in TSB medium containing

the appropriate antibiotics. Overnight cultures were diluted to a starting OD_{600} of 0.05

in the presence of 100 ng/ml Atet where appropriate, incubated at 25°C or 37°C with

aeration and OD_{600} values determined at 2 h intervals. Growth curves were performed

in triplicate, and averages and standard deviations are plotted.

Minimum Inhibitory Concentrations. Overnight cultures of wildtype LAC* and

Δ*era* strains were adjusted to an OD_{600} of 0.05 in Mueller-Hinton broth and 100 μl

incubated in 96 well plates with 2-fold dilutions of various antimicrobials at the

following starting concentrations: spectinomycin 250 μg/ml and chloramphenicol 64

μg/ml. Plates were incubated at 37°C overnight with shaking. Assays were performed

in triplicate, and averages and standard deviations are plotted.

Protein purifications. Proteins were purified from 1-2 L *E. coli* cultures. Cultures

were grown to an OD_{600} 0.5-0.7, expression induced with 1 mM IPTG and incubated

for 3 h at $37^{o}C$. Protein purifications were performed by either nickel or glutathione

affinity chromatography. For nickel purifications, cell pellets were resuspended in 5

ml Buffer A (50 mM Tris pH 7.5, 150 mM NaCl, 5% glycerol, 10 mM imidazole),

lysed with 100 μg lysozyme and sonication and the filtered cell lysate loaded onto a 1

ml HisTrap HP Ni^{2+} column (GE Healthcare) before elution using a gradient of Buffer

B (50 mM Tris pH 7.5, 150 mM NaCl, 5% glycerol, 500 mM imidazole). GST-tagged

proteins were resuspended in 5 ml PBS, lysed and the cell lysate was loaded onto a 1

ml GSTrap HP column (GE Healthcare) before elution using 50 mM Tris pH 8.0, 10

mM reduced glutathione. Protein containing fractions were dialysed in 50 mM Tris pH

7.5, 200 mM NaCl, 5% glycerol before storage at -80°C. Protein concentrations were

determined by A_{280} readings.

Synthesis of (p)ppGpp and differential radial capillary action of ligand assay

(DRaCALA). The synthesis of (p)ppGpp and DRaCALA binding assays were

performed as described previously [22].

Ribosomal profiles from *S. aureus* cell extracts. Crude isolations of ribosomes from

S. aureus cell extracts were achieved as described by Chin Loh *et al.* with some

modifications [27]. Briefly, 100 ml cultures of the different *S. aureus* strains were

grown to an OD_{600} of 0.4 in TSB 100 ng/ml Atet. 100 μg/ml chloramphenicol was

added to each culture, before being cooled to 4°C. Pelleted cells were suspended in

association buffer (20 mM Tris-HCl pH 7.6, 8 mM $MgCl_2$, 30 mM NH_4Cl and 2 mM

β-mercaptoethanol), normalized to an OD_{600} of 15, lysed by the addition of 0.2 μg/ml

lysostaphin and 75 ng/ml DNase and incubated for 30 min at 37°C. The extracts were

centrifuged at 17,000 x g for 5 min and subsequently 250 μl were layered onto 10-50%

sucrose density gradients made in association buffer. Gradients were centrifuged for

4.5 h at 192,100 x g. Gradients were fractionated by upwards displacement of 250 μl

aliquots, which were analysed for RNA content at an absorbance of 260 nm.

Helicase assay. Reactions were set up as described [53]. Briefly, 250 nM of Cy3-

labelled   duplex   RNA   (Cy3-GCUUUACGGUGCUA,

AACAAAACAAAAUAGCACCGUAAAGC) purchased from IDT, was incubated

with 0.5 μM recombinant protein in reaction buffer (20 mM Tris pH 7.5, 50 mM

potassium acetate, 5 mM magnesium acetate, 10 mM DTT, 0.1 mg/ml BSA, 10 U

RNasin). The 0.5 μM recombinant protein used is at a concentration similar to that

used in the affinity pulldown assays (1 μM) that show an Era-CshA interaction. The

reaction was initiated with the addition of 1 mM ATP and incubated at 25°C for 10

min. Reactions were terminated by mixing with an equal volume of stop buffer (1%

SDS, 0.5 mM EDTA, 20% glycerol) and loaded onto 20% native polyacrylamide gels

before being electrophoresed in 0.5 x TBE. Bands were imaged using a ChemiDoc MP

imager (Bio-Rad).

Nucleotide hydrolysis assays. The ability of proteins to hydrolyse GTP was

determined by incubating 100 nM recombinant Era with equimolar YbeY/CshA or

Rel_{SA}, 100 nM *S. aureus* ribosomes, 1 μM GTP and 2.78 nM α-^{32}P-GTP in 40 mM

Tris pH 7.5, 100 mM NaCl, 10 mM $MgCl_2$ at 37°C for the indicated times. All

reactions were set up in the absence of Era as controls. pppGpp hydrolysis assays were

performed by incubating 100 nM recombinant Rel_{SA} in the presence and absence of

equimolar GST, GST-Era or GST-CshA, 1 μM pppGpp and 2.78 nM α-^{32}P-pppGpp in

50 mM Tris pH 8.5, 0.1 mM $MnCl_2$, 20 mM KCl at 37°C for the indicated times. All

reactions were set up in the absence of Rel_{SA} as controls. Hydrolysis reactions were

inactivated with the addition of formic acid to a final concentration of 1.2 M.

Precipitated proteins were pelleted by centrifugation at 17,000 x g for 10 min.

Reaction products were then visualized by spotting 1 μl on PEI-cellulose thin layer

chromatography (TLC) plates (Merck Millipore) and separated using 1 M KH_2PO_4,

pH 3.6 buffer. The radioactive spots were visualised using an LA 7000 Typhoon

PhosphorImager and images quantified using ImageJ.

References

1. Leipe DD, Wolf YI, Koonin EV, Aravind L. Classification and evolution of P-
loop GTPases and related ATPases. J Mol Biol. 2002;317(1):41-72.

2. March PE, Lerner CG, Ahnn J, Cui X, Inouye M. The *Escherichia coli* Ras-
like protein (Era) has GTPase activity and is essential for cell growth. Oncogene.
1988;2(6):539-44.

3. Inada T, Kawakami K, Chen SM, Takiff HE, Court DL, Nakamura Y.
Temperature-sensitive lethal mutant of *era*, a G protein in *Escherichia coli*. J
Bacteriol. 1989;171(9):5017-24.

4. Takiff HE, Chen SM, Court DL. Genetic analysis of the rnc operon of
Escherichia coli. J Bacteriol. 1989;171(5):2581-90.

5. Gollop N, March PE. A GTP-binding protein (Era) has an essential role in
growth rate and cell cycle control in *Escherichia coli*. J Bacteriol. 1991;173(7):2265-
70.

6. Anderson PE, Matsunaga J, Simons EL, Simons RW. Structure and regulation
of the *Salmonella* Typhimurium rnc-*era-recO* operon. Biochimie. 1996;78(11-
12):1025-34.

7. Minkovsky N, Zarimani A, Chary VK, Johnstone BH, Powell BS, Torrance
PD, et al. Bex, the *Bacillus subtilis* homolog of the essential *Escherichia coli* GTPase
Era, is required for normal cell division and spore formation. J Bacteriol.
2002;184(22):6389-94.

8. Morimoto T, Loh PC, Hirai T, Asai K, Kobayashi K, Moriya S, et al. Six
GTP-binding proteins of the Era/Obg family are essential for cell growth in *Bacillus*
subtilis. Microbiology. 2002;148(Pt 11):3539-52.

9.   Lu Q, Inouye M. The gene for 16S rRNA methyltransferase (*ksgA*) functions

as a multicopy suppressor for a cold-sensitive mutant of *era*, an essential RAS-like

GTP-binding protein in *Escherichia coli*. J Bacteriol. 1998;180(19):5243-6.

10.   Lerner CG, Inouye M. Pleiotropic changes resulting from depletion of Era, an

essential GTP-binding protein in *Escherichia coli*. Mol Microbiol. 1991;5(4):951-7.

11.   Powell BS, Court DL, Inada T, Nakamura Y, Michotey V, Cui X, et al. Novel

proteins of the phosphotransferase system encoded within the *rpoN* operon of

Escherichia coli. Enzyme IIANtr affects growth on organic nitrogen and the

conditional lethality of an erats mutant. J Biol Chem. 1995;270(9):4822-39.

12.   Britton RA, Powell BS, Dasgupta S, Sun Q, Margolin W, Lupski JR, et al.

Cell cycle arrest in Era GTPase mutants: a potential growth rate-regulated checkpoint

in *Escherichia coli*. Mol Microbiol. 1998;27(4):739-50.

13.   Johnstone BH, Handler AA, Chao DK, Nguyen V, Smith M, Ryu SY, et al.

The widely conserved Era G-protein contains an RNA-binding domain required for

Era function *in vivo*. Mol Microbiol. 1999;33(6):1118-31.

14.   Sharma MR, Barat C, Wilson DN, Booth TM, Kawazoe M, Hori-Takemoto C,

et al. Interaction of Era with the 30S ribosomal subunit implications for 30S subunit

assembly. Mol Cell. 2005;18(3):319-29.

15.   Tu C, Zhou X, Tarasov SG, Tropea JE, Austin BP, Waugh DS, et al. The Era

GTPase recognizes the GAUCACCUCC sequence and binds helix 45 near the 3' end

of 16S rRNA. Proc Natl Acad Sci U S A. 2011;108(25):10156-61.

16.   Tamaru D, Amikura K, Shimizu Y, Nierhaus KH, Ueda T. Reconstitution of

30S ribosomal subunits *in vitro* using ribosome biogenesis factors. RNA.

2018;24(11):1512-9.

17.   Bunner AE, Nord S, Wikstrom PM, Williamson JR. The effect of ribosome

assembly cofactors on in vitro 30S subunit reconstitution. J Mol Biol. 2010;398(1):1-

7.

18.   Vercruysse M, Kohrer C, Shen Y, Proulx S, Ghosal A, Davies BW, et al.

Identification of YbeY-Protein Interactions Involved in 16S rRNA Maturation and

Stress Regulation in *Escherichia coli*. MBio. 2016;7(6).

19.   Baumgardt K, Gilet L, Figaro S, Condon C. The essential nature of YqfG, a

YbeY homologue required for 3' maturation of *Bacillus subtilis* 16S ribosomal RNA

is suppressed by deletion of RNase R. Nucleic Acids Res. 2018;46(16):8605-15.

20.   Lemos JA, Lin VK, Nascimento MM, Abranches J, Burne RA. Three gene

products govern (p)ppGpp production by *Streptococcus mutans*. Mol Microbiol.

2007;65(6):1568-81.

21.   Cassels R, Oliva B, Knowles D. Occurrence of the regulatory nucleotides

ppGpp and pppGpp following induction of the stringent response in staphylococci. J

Bacteriol. 1995;177(17):5161-5.

22.   Corrigan RM, Bellows LE, Wood A, Grundling A. ppGpp negatively impacts

ribosome assembly affecting growth and antimicrobial tolerance in Gram-positive

bacteria. Proc Natl Acad Sci U S A. 2016;113(12):E1710-9.

23.   Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, et al. A

genetic resource for rapid and comprehensive phenotype screening of nonessential

Staphylococcus aureus genes. MBio. 2013;4(1):e00537-12.

24.   Chaudhuri RR, Allen AG, Owen PJ, Shalom G, Stone K, Harrison M, et al.

Comprehensive identification of essential *Staphylococcus aureus* genes using

Transposon-Mediated Differential Hybridisation (TMDH). BMC Genomics.

2009;10:291.

25.   Santiago M, Matano LM, Moussa SH, Gilmore MS, Walker S, Meredith TC.

A new platform for ultra-high density *Staphylococcus aureus* transposon libraries.

BMC Genomics. 2015;16:252.

26.   Inoue K, Alsina J, Chen J, Inouye M. Suppression of defective ribosome

assembly in a *rbfA* deletion mutant by overexpression of Era, an essential GTPase in

Escherichia coli. Mol Microbiol. 2003;48(4):1005-16.

27.   Loh PC, Morimoto T, Matsuo Y, Oshima T, Ogasawara N. The GTP-binding

protein YqeH participates in biogenesis of the 30S ribosome subunit in *Bacillus*

subtilis. Genes Genet Syst. 2007;82(4):281-9.

28.   Jacob AI, Kohrer C, Davies BW, RajBhandary UL, Walker GC. Conserved

bacterial RNase YbeY plays key roles in 70S ribosome quality control and 16S rRNA

maturation. Mol Cell. 2013;49(3):427-38.

29.   Oliveira Paiva AM, Friggen AH, Qin L, Douwes R, Dame RT, Smits WK.

The Bacterial Chromatin Protein HupA Can Remodel DNA and Associates with the

Nucleoid in *Clostridium difficile*. J Mol Biol. 2019.

30.   Chen X, Court DL, Ji X. Crystal structure of ERA: a GTPase-dependent cell

cycle regulator containing an RNA binding motif. Proc Natl Acad Sci U S A.

1999;96(15):8396-401.

31.   Kim S, Corvaglia AR, Leo S, Cheung A, Francois P. Characterization of RNA

Helicase CshA and Its Role in Protecting mRNAs and Small RNAs of *Staphylococcus*

aureus Strain Newman. Infect Immun. 2016;84(3):833-44.

32.   Giraud C, Hausmann S, Lemeille S, Prados J, Redder P, Linder P. The C-

terminal region of the RNA helicase CshA is required for the interaction with the

degradosome and turnover of bulk RNA in the opportunistic pathogen *Staphylococcus*

aureus. RNA Biol. 2015;12(6):658-74.

33.   Prud'homme-Genereux A, Beran RK, Iost I, Ramey CS, Mackie GA, Simons

RW. Physical and functional interactions among RNase E, polynucleotide

phosphorylase and the cold-shock protein, CsdA: evidence for a 'cold shock

degradosome'. Mol Microbiol. 2004;54(5):1409-21.

34.   Charollais J, Dreyfus M, Iost I. CsdA, a cold-shock RNA helicase from

Escherichia coli, is involved in the biogenesis of 50S ribosomal subunit. Nucleic

Acids Res. 2004;32(9):2751-9.

35.   Daigle DM, Brown ED. Studies of the interaction of *Escherichia coli* YjeQ

with the ribosome *in vitro*. J Bacteriol. 2004;186(5):1381-7.

36.   Campbell TL, Brown ED. Genetic interaction screens with ordered

overexpression and deletion clone sets implicate the *Escherichia coli* GTPase YjeQ in

late ribosome biogenesis. J Bacteriol. 2008;190(7):2537-45.

37.   Gratani FL, Horvatek P, Geiger T, Borisova M, Mayer C, Grin I, et al.

Regulation of the opposing (p)ppGpp synthetase and hydrolase activities in a

bifunctional RelA/SpoT homologue from *Staphylococcus aureus*. PLoS Genet.

2018;14(7):e1007514.

38.   Reiss S, Pane-Farre J, Fuchs S, Francois P, Liebeke M, Schrenzel J, et al.

Global analysis of the *Staphylococcus aureus* response to mupirocin. Antimicrob

Agents Chemother. 2012;56(2):787-804.

39.   Wilson DN, Nierhaus KH. The weird and wonderful world of bacterial

ribosome regulation. Crit Rev Biochem Mol Biol. 2007;42(3):187-219.

40.   Gollop N, March PE. Localization of the membrane binding sites of Era in

Escherichia coli. Res Microbiol. 1991;142(2-3):301-7.

41.   Zhang J, Inouye M. MazG, a nucleoside triphosphate pyrophosphohydrolase,

interacts with Era, an essential GTPase in *Escherichia coli*. J Bacteriol.

2002;184(19):5323-9.

42.   Chopade BA, Shankar S, Sundin GW, Mukhopadhyay S, Chakrabarty AM.

Characterization of membrane-associated *Pseudomonas aeruginosa* Ras-like protein

Pra, a GTP-binding protein that forms complexes with truncated nucleoside

diphosphate kinase and pyruvate kinase to modulate GTP synthesis. J Bacteriol.

1997;179(7):2181-8.

43.   Huang Y, Zhang B, Dong K, Zhang X, Hou L, Wang T, et al. Up-regulation of

yggG promotes the survival of *Escherichia coli* cells containing Era-1 mutant protein.

FEMS Microbiol Lett. 2007;275(1):8-15.

44.   Lehnik-Habrink M, Rempeters L, Kovacs AT, Wrede C, Baierlein C, Krebber

H, et al. DEAD-Box RNA helicases in *Bacillus subtilis* have multiple functions and

act independently from each other. J Bacteriol. 2013;195(3):534-44.

45.   Lehnik-Habrink M, Pfortner H, Rempeters L, Pietack N, Herzberg C, Stulke J.

The RNA degradosome in *Bacillus subtilis*: identification of CshA as the major RNA

helicase in the multiprotein complex. Mol Microbiol. 2010;77(4):958-71.

46.   Cashel M, Gentry, D.R., Hernandez, V.J., and Vinella, D. The stringent

response. In *E. coli* and *Salmonella*: Washington DC: ASM press; 1996.

47.   Kriel A, Bittner AN, Kim SH, Liu K, Tehranchi AK, Zou WY, et al. Direct

regulation of GTP homeostasis by (p)ppGpp: a critical component of viability and

stress resistance. Mol Cell. 2012;48(2):231-41.

48.   Ross W, Sanchez-Vazquez P, Chen AY, Lee JH, Burgos HL, Gourse RL.

ppGpp Binding to a Site at the RNAP-DksA Interface Accounts for Its Dramatic

Effects on Transcription Initiation during the Stringent Response. Mol Cell.

2016;62(6):811-23.

49.   Himeno H, Hanawa-Suetsugu K, Kimura T, Takagi K, Sugiyama W, Shirata

S, et al. A novel GTPase activated by the small subunit of ribosome. Nucleic Acids

Res. 2004;32(17):5303-9.

50.   Achila D, Gulati M, Jain N, Britton RA. Biochemical characterization of

ribosome assembly GTPase RbgA in *Bacillus subtilis*. J Biol Chem.

2012;287(11):8417-23.

51.   Wachi M, Umitsuki G, Shimizu M, Takada A, Nagai K. *Escherichia coli cafA*

gene encodes a novel RNase, designated as RNase G, involved in processing of the 5'

end of 16S rRNA. Biochem Biophys Res Commun. 1999;259(2):483-8.

52.   Sihto HM, Tasara T, Stephan R, Johler S. Validation of reference genes for

normalization of qPCR mRNA expression levels in *Staphylococcus aureus* exposed to

osmotic and lactic acid stress conditions encountered during food production and

preservation. FEMS Microbiol Lett. 2014;356(1):134-40.

53.   Xu L, Wang L, Peng J, Li F, Wu L, Zhang B, et al. Insights into the Structure

of Dimeric RNA Helicase CsdA and Indispensable Role of Its C-Terminal Regions.

Structure. 2017;25(12):1795-808 e5.

Fig 1. Role of Era in bacterial growth and ribosome assembly. A) Schematic

representation of the *era*-containing operons from *S. aureus*, *B. subtilis* and *E. coli*.

The *era* operon in *S. aureus* consists of: *ybeZ* – encoding for a protein of unknown

function; *ybeY* – a 16S rRNA endoribonuclease; *dgkA* - a diacylglycerol kinase

involved in lipid metabolism; *cdd* - a cytidine deaminase involved pyrimidine

metabolism; *era*, and *recO* encoding for a DNA repair protein. Arrows indicate the site

of transposon insertions in the *S. aureus* Nebraska Transposon library. B) Growth of *S.*

aureus strains LAC* iTET, LAC* Δ*era* iTET and LAC* Δ*era* iTET-*era*. Overnight

cultures were diluted to an OD_{600} of 0.05 and grown in the presence of 100 ng/ml Atet

for 8 h. Growth curves were performed in triplicate (N=3), with averages and standard

deviations shown. C) Ribosome profiles from LAC* iTET, LAC* Δ*era* iTET and

783 LAC* Δ*era* iTET-*era.* Normalised extracts from each strain were layered onto 10-50%

784 sucrose gradients. Gradients were fractionated and analysed for RNA content at an

785 absorbance of 260 nm. 30S, 50S, 70S and polysomes-containing fractions are

786 indicated. Experiments were performed in triplicate (N=3) with one representative

787 graph shown.

Fig 2. Era interacts with other proteins encoded in the same operon. A) Bacterial

two-hybrid showing a positive interaction between Era-T25 and both T18-YbeZ and

YbeY. T25 and T18-encoding empty vectors are used as negative controls. N=3, with

one representative image shown. B) Schematic representation of the luciferase vectors

used for native protein-protein interaction studies. pAP118 contains an Atet-inducible

promoter upstream of two genes – SB encodes for the small bit of the luciferase

protein, while LB encodes for the large bit. When genes encoding for interacting

proteins (A & B) are translationally fused to SB and LB they associate to produce

luciferase. pAP256 and pAP257 are control vectors. C) Split luciferase assay

demonstrating a significant interaction between Era-Era and Era-YbeY in *S. aureus*.

The negative is pAP256-*era*$_{SB}$-LB, which has Era fused to the SB but nothing fused to

LB. The negative is normalised to a relative luminescence unit (RLU) of 1. The

average values and standard deviations of triplicate experiments are plotted (N=3).

Statistical analysis was performed using a one-way ANOVA, followed by Dunnett's

multiple comparisons test (** $P < 0.005$, **** $P < 0.0001$). D) Top: affinity pulldown

assay using GST or GST-tagged Era coupled to glutathione beads. Beads alone, GST

or GST-Era coupled beads were incubated with His-tagged YbeY. After washing,

bound YbeY-His was detected using HRP-conjugated anti-His antibodies. N=3 with

one representative image shown. Below: the mean signal intensities and standard

deviations from the 3 repeats are plotted. Statistical analysis was performed using a

one-way ANOVA, followed by Dunnett's multiple comparisons test (* $P < 0.05$).

Fig 3. Era interacts with the DEAD-box helicase CshA. A) Bacterial two-hybrid

showing an interaction between CshA-T18 and both Era-T25 and T25-Era. Era is

known to form a dimer and an Era-Era interaction is used as a positive control. N=3

with one representative image shown. B) Affinity pulldown assay using GST or GST-

tagged CshA coupled to glutathione beads. Beads were incubated with His-tagged Era

and after washing bound protein was detected using HRP-conjugated anti-His

antibodies. N=3 with one representative image shown. C) Split luciferase assay

demonstrating an interaction between Era and CshA in *S. aureus*. Dimers resulting

from Era fused to both the small (SB) and large bits (LB) of the luciferase gene

function as a positive control. The negative, which is normalised to a relative

luminescence unit (RLU) of 1, has Era fused to the SB but nothing fused to LB. Cells

were grown in the presence of 100 ng/ml Atet at 37°C before being normalised. The

average values and standard deviations of triplicate experiments (N=3) are plotted. D)

Schematic representation of CshA and Era. The sizes of each domain construct are

indicated by the black lines and numbering. E) Split luciferase assay with Era fused to

SB and various truncated domain constructs of CshA fused to LB. F) Split luciferase

assay with the G domain of Era (1-180) fused to SB and full-length CshA fused to LB.

The average values and standard deviations of triplicate experiments (N=3) for all

luciferase assays are plotted. All statistical analysis was performed using a one-way

ANOVA, followed by Dunnett's multiple comparisons test (** $P < 0.005$, **** $P <$

0.0001).

Fig 4. Era and CshA are required for cold shock survival, virulence and rRNA

processing. A) Serial dilutions of the wildtype LAC*, Δ*era*, Δ*cshA* and Δ*era cshA*

were spotted onto TSA agar plates and incubated at either 37 or 25°C for the times

indicated. N=3 with one representative image shown. B) Survival of *G. mellonella*

larvae inoculated into the hind leg with 1 x 10^5 cfu of wildtype LAC* and mutant

Δ*era*, Δ*cshA* and Δ*era cshA* strains. Larvae were incubated at 37°C for 4 days.

Experiments were repeats in triplicate (N=3), with one graph shown. Survival curves

were calculated by Kaplan-Meier analysis, with the log-rank (Mantel Cox) test

performed to compare different strains (* $P < 0.05$, *** $P < 0.0005$, **** $P < 0.0001$).

C) rRNA profiles from *era* and *cshA* mutant strains. 500 ng of RNA extracted from

LAC*, Δ*era*, Δ*cshA* and Δ*era cshA* grown to an OD_{600} of 0.4 at either 37 or 25°C,

were run on 0.7% agarose/0.9% synergel gels and stained with Evagreen dye. *

highlights the presence of either processing or degradation intermediates. N=3 with

one representative image shown. D) top: schematic of the 17S rRNA, with arrows

indicating processing sites that lead to the formation of the mature 16S rRNA. The

blue bars indicate the probe regions amplified by RT-PCR. Bottom: RT-PCR of total

RNA from the wildtype LAC*, Δ*era*, and Δ*cshA* strains grown at 25°C using probes

as indicated above. rRNA values are expressed as mean relative expression +/- SD

normalised to the wildtype and the expression of *rho* RNA as an internal control.

Statistical analysis was performed using a two-way ANOVA, followed by Tukey's

multiple comparisons test (**** *P* < 0.0001). Cycle threshold values were determined

for 3 biological repeats in duplicate (N=3).

Fig 5. The stringent response synthetase Rel$_{SA}$ interacts with Era and CshA. A)

Top: Bacterial two-hybrid showing an interaction between CshA-T18/T18-CshA and

both Rel$_{SA}$-T25 and T25-Rel$_{SA}$. CshA did not interact with the small (p)ppGpp

synthetase enzymes RelP or RelQ. Bottom: Bacterial two-hybrid showing an

interaction between Rel$_{SA}$-T25 and both Era-T18 and T18-Era. N=3 with one

representative image shown. B) Affinity pulldown assay using GST, GST-tagged Era

and GST-tagged CshA coupled to glutathione beads. Beads alone, GST, GST-Era and

GST-CshA coupled beads were incubated with His-tagged Rel$_{SA}$. After washing,

bound Rel$_{SA}$-His was detected using HRP-conjugated anti-His antibodies. N=3 with

one representative image shown above. Below, the mean signal intensities and

standard deviations from the 3 repeats are plotted. Statistical analysis was performed

using a one-way ANOVA, followed by Dunnett's multiple comparisons test (*** $P <$

0.0005). C & D) Split luciferase assay demonstrating an interaction between CshA (C)

or Era (D) and Rel_{SA} in *S. aureus*. The negative controls have genes singly fused to

either the SB or the LB. The average values and standard deviations of triplicate

experiments (N=3) are plotted. All statistical analysis was performed using a one-way

ANOVA, followed by Dunnett's multiple comparisons test (**** $P < 0.0001$).

Fig 6. The stringent response influences Era GTPase activity and rRNA

processing. A) The GTPase activity of 100 nM Era was measured in the presence of

an equal amount of ribosomes, 100 nM Rel_{SA} and 1 μM GTP. All reactions contained

ribosomes. The Rel_{SA} only control reaction contained Rel_{SA} and ribosomes but no Era.

Hydrolysis of ^{32}P-GTP was monitored by TLC and the percentage GDP formed

quantified using ImageJ. Experiments were repeated three times with means and

standard deviations shown. Statistical analysis was performed using a two-way

ANOVA, followed by Dunnett's multiple comparisons test (** $p < 0.005$, * $p < 0.05$).

B & C) rRNA extracted from LAC*. Strains were grown at 25°C until an OD_{600} of 0.4.

Cells were split and half exposed to 60 μg/ml mupirocin for 1 h (plus), which at 25°C

corresponds to one doubling. RT-PCR of total RNA using the probes as indicated in

Fig. 4d that bind internally to the 16S (B) or to the unprocessed leader and trailer

sequences (C) of the 17S rRNA. rRNA values are expressed as mean relative

expression +/- SD normalised to the expression of *rho* RNA as an internal control. For

comparisons between two groups (B - LAC* minus v plus) a two-tailed, unpaired

Student's t-test was performed (**** $p < 0.0001$). For comparisons between more than

two groups (C) a two-way ANOVA, followed by Sidak's multiple comparisons test

was performed (** $p < 0.005$). Cycle threshold values were determined for 3

biological repeats in duplicate (N=3).

Supporting information

S1 Fig. Antibiotic susceptibility testing

S2 Fig. Enzyme activity of CshA and Era

S3 Fig. Growth of Δ*era* and Δ*cshA* strains at 37 and 25°C

S4 Fig. Cross-complementation of Δ*rsgA,* Δ*era* and Δ*cshA* strains

S5 Fig. ppGpp does not interact with CshA

Supplemental methods – plasmid and strain construction

S1 Table. Bacterial strains used in this study

S2 Table. Primers used in this study