Climate gradients explain population-level divergence in drought-induced plasticity of functional traits and gene expression in a South African *Protea*

Melis Akman[1,2], Jane E. Carlson[3,4], Andrew M. Latimer[1]

[1] Department of Plant Sciences, UC Davis

[2] Department of Plant and Microbial Biology, UC Berkeley

[3] Department of Biology, Nicholls State

[4] Gulf Coast Network Inventory and Monitoring Program, National Park Services

Abstract

Long term environmental variation often drives local adaptation and leads to trait differentiation across populations. Additionally, when traits can change in an environment-dependent way through phenotypic plasticity, the underlying genetic variation will also be under selection, but only in the inducing environment. Both of these processes will create a landscape of differentiation across populations in trait means as well as their plasticity. However, studies uncovering environmental drivers of this variation are scarce. With this work, we studied drought responses in seedlings of a shrub species from the Cape Floristic Region, the common sugarbush (*Protea repens*). We measured morphological and physiological traits as well as whole transcriptomes in 8 populations that represent both the climatic and the geographic distribution of this species. We found that there is substantial variation in how populations respond to drought, but we also observe common patterns such as reduced leaf size and thickness and upregulation of stress- and down-regulation of growth-related gene groups. Both environmental heterogeneity and milder source site climates were associated with higher plasticity in various traits and co-expression gene networks. By uncovering associations between traits, trait plasticity, co-expression gene networks with source site climate, we showed that temperature plays a bigger role in shaping these patterns when compared to precipitation, in line with recent changes in the region due to climate change. We also found that traits respond to climatic variation in a context dependent manner: some associations between traits and climate were apparent only under certain growing conditions.

Keywords: phenotypic plasticity, climatic variation, trait variation, transcriptome, *Protea repens*, drought, climate change

INTRODUCTION

Populations can persist in changing environmental conditions through three mechanisms: by tracking their climatic niche in space through migration or dispersal, by shifting their niche through adaptation, and/or by matching traits to the new environment through functional phenotypic plasticity (Chevin *et al.* 2010). For plants, plasticity can be the fastest, since it involves rapid adjustments of morphology, physiology and gene expression within a single generation. Like mean trait values, the form and magnitude of trait plasticity can be under selection and thus plasticity can also vary among populations along environmental gradients (Geber & Griffen 2003). Plasticity in key growth-related plant traits, including stomatal density and conductance, growth rate, and leaf morphology has been observed to vary among populations in long-lived woody plants (e.g. Aubin *et al.* 2016) as well as herbaceous annuals (Jacobs & Latimer 2012). Recent studies have demonstrated that population differences in plasticity might have climatic drivers (Pratt & Mooney 2013; Lázaro-Nogal *et al.* 2015). However, examples of underlying environmental processes that maintain population differentiation in plasticity are still scarce.

Functional plasticity can provide at least short-term benefits to individual plants upon changing environmental conditions (Wennersten & Forsman 2012). The fitness consequences of plasticity will often be context dependent, however, as is demonstrated for many functional traits (e.g. Etterson 2004). For example, plants generally adjust many features of leaf morphology to optimize photosynthesis at different light levels, and tune stomatal behavior to adjust water use efficiency depending on soil moisture levels. While levels and patterns of plasticity can have strong, context-dependent fitness consequences, it remains less clear how often differential plasticity confers environment-specific fitness benefits (Forsman 2015). In a few cases, differential plasticity has been directly shown to be adaptive: for example, Dudley & Schmitt (1996) showed that plastic stem elongation was beneficial at high but not low plant densities. Other studies have characterized patterns of plasticity that suggest local adaptation: for example, Sultan (2003) showed that plasticity in allocation to leaf light capture was more pronounced in a generalist species than a high-light specialist. Nicotra *et al.* (2015) showed that plasticity in seedlings of an alpine plant was higher and also favored in low elevation but not in high elevation populations. Resurrection studies using historical and modern *Daphnia magna* populations showed evolution of plasticity for heat tolerance and fish predation (Stoks *et al.* 2016; Cuenca

Cambronero *et al.* 2018). In a field study using *Boechera stricta*, Wadgymar *et al.* (2018) showed differential germination plasticity along elevational gradients. Apart from these studies, our understanding of the role of adaptive evolution in differential plasticity remains largely theoretical (West-Eberhard 2003; Ghalambor *et al.* 2015).

In light of these studies and theoretical predictions (Bradshaw & Hardwick 1989; Grime 1994; Schlichting & Pigliucci 1995; Alpert & Simms 2002), we expect that local adaptation should promote differential plasticity under two main circumstances. First, plants from environments with strong and relatively predictable forms of spatial or temporal heterogeneity should be more plastic than those from homogeneous environments (Ackerly *et al.* 2000; Van Kleunen & Fischer 2005; Baythavong & Stanton 2010; Scheiner 2013; but see Palacio-López *et al.* 2015). Second, plants from harsh climates are expected to be less plastic than those from milder climates (Vitasse *et al.* 2013; Gugger *et al.* 2015), because the fitness costs of environmental mismatch should be greater in resource-poor conditions (Alpert & Simms 2002). Elevation contrasts provide one example of this latter pattern, as plants from colder, higher-elevation environments tend to have both lower and less plastic growth rates than plants from warmer sites (Nicotra *et al.* 2015).

The form and magnitude of functional plasticity which organisms express can be expected to influence the vulnerability of populations to environmental change. Thus, understanding the factors that drive differentiation among populations in functional plasticity, and the consequences of these plasticity differences for population vulnerability, therefore, is an important challenge for global change research. The importance of plasticity for population persistence is well-grounded in theory (Nicotra *et al.* 2010; Chevin *et al.* 2010) and borne out in some experimental work (e.g. Anderson & Gezon 2015; Lázaro-Nogal *et al.* 2015; Sgrò *et al.* 2016). Studies that have incorporated plasticity variation in predictive models of species distribution have yielded a range of results; in some cases, plasticity is projected to cause slow loss at the warmer range edge (Valladares *et al.* 2014) or open new habitats for colonization (Muñoz *et al.* 2014), while in others plasticity is projected to accelerate range loss (Duputié *et al.* 2015). The gaps in empirical knowledge about both the functional basis and environmental landscape of plasticity variation creates a challenge for these models to accurately predict how and under what scenarios plasticity will buffer populations in the face of climate change.

Our study species originates from the Cape Floristic Region (CFR), a biodiversity
hotspot with a climatically stable past (Cowling 2009), that is predicted to be strongly
affected by climate change. Climate models project warming up to 3 °C by 2070 (Wilson
et al. 2015), and precipitation projections being more variable, most models project
drying in the super-diverse western mountains (Hewitson & Crane 2006; Wilson *et al.*
2015). Our study species, *Protea repens* (common sugar bush), is part of a lineage that
has radiated over the past 5 to 18 million years to produce 112 species, ~85% of which
are endemic (Rebelo 2001; Sauquet *et al.* 2009). Along the environmental gradients
Protea species inhabit, we have evidence for local adaptation and context-dependent
benefits in several traits (Latimer *et al.* 2009; Carlson *et al.* 2011, 2016; Prunier *et al.*
2012; Akman *et al.* 2016; Mitchell *et al.* 2018). There is also evidence for differential
developmental plasticity and drought plasticity in leaf traits of several other *Protea*
species (*Protea sect. Exsertae*: Carlson & Holsinger 2012; Heschel *et al.* 2014),
however, the extent to which this reflects local adaptation remains unknown. Upon
rapidly shifting environmental conditions, plasticity will play an important role for survival
in this species considering its long generation time. Thus, uncovering patterns of
plasticity can inform vulnerability predictions for this species under climate change.

A particularly important form of environmental variation in the CFR is drought.
Some Mediterranean-climate regions including the CFR, have recently experienced
unprecedented drought. Such droughts are projected to be more likely in the near
future. Recent droughts have caused large mortality events in trees (Allen *et al.* 2010;
Young *et al.* 2017). Understanding the patterns and effects of functional plasticity to
drought is thus a vital issue for plant vulnerability in many semi-arid regions.
Measurements of plasticity in gene expression, in which tens of thousands of
"phenotypes" can be measured simultaneously, are particularly valuable for
investigating the causes and consequences of plasticity. Gene expression patterns can
change within seconds of a stimulus, long before larger biochemical shifts and
morphological changes become apparent. Gene expression patterns can also be linked
to morphological and physiological adjustments, providing insights into the genetic basis
of adaptation and/or plasticity (Whitehead 2012; Akman *et al.* 2016). Previous studies
have revealed patterns of adaptive plasticity in gene expression in fish (Morris *et al.*
2014; Schneider *et al.* 2014; Dayan *et al.* 2015), plants (Lasky *et al.* 2014; Colicchio *et*
al. 2015) and corals (Granados-Cifuentes *et al.* 2013; Barshis *et al.* 2013). These
studies demonstrate that gene expression profiling can usefully complement standard
functional trait measurements to help understand the genetic basis and physiological

significance of plasticity and how it varies across populations. With this study, we aim to answer two questions: 1) How do populations of *P. repens* respond to drought and how do these responses vary across populations? and 2) Do the climatic conditions of the source sites covary with traits, gene expression and their plasticity? To do so, we first uncover the consequences of a severe, experimentally imposed drought by measuring traits and gene expression patterns in seedlings in a dry-down experiment. Then we test for population differentiation in trait means and drought-response plasticity in functional traits, performance, and gene expression patterns, and evaluate associations between patterns of population differentiation and their potential climatic drivers.

MATERIALS AND METHODS

Species description

Protea repens L. is a common evergreen shrub endemic to South Africa, with particularly high abundance in semi-arid fynbos of the CFR (Rebelo 2001). This species has narrow, sclerophyllous leaves and reaches 2-4 m in height. Growth is sympodial, with new shoots and leaves generally emerging in the months before flowering. Plants flower between Feb-Aug for 1-3 months, with eastern populations blooming later than western populations. Inflorescences are dense heads of 70-150 protandrous florets, subtended by pink or white bracts. Each floret can produce a single-seeded fruit, although seed set per head averages under 30%. All viable and nonviable fruits remain within a woody, serotinous infructescence until the adult plant is killed by fire, which occurs at ~10-13 year intervals in the fynbos (with variation among geographic regions; (Van Wilgen *et al.* 2010). Fruits disperse by tumbling, and seeds germinate during the next rainy season (May-Aug for much of the CFR). Due to the Mediterranean rainfall pattern in the western CFR, some western population sites receive far more wintertime precipitation than do eastern population sites.

Field sampling and climate data

We collected seeds from 8 wild populations of *P. repens* spanning much of its geographic distribution in the CFR (**Figure 1A**). Between Feb. and May 2011, we visited these 8 sites as part of a larger sampling effort in *P. repens* (see Carlson *et al.* 2016). At each site, we collected infructescences from 20-40 plants along a transect through each population's center, and we air-dried infructescences until they opened and single-seeded fruits could be removed. We stored seeds in envelopes at room

temperature until the greenhouse study. At each seed collection site, we also recorded
GPS coordinates near the population center, and we used these points to extract
elevation and six climate metrics for rainfall and temperature from GIS layers. A
principal component analysis performed on these data confirmed that the sample sites
cover most of the range of variation in major environmental factors (**Figure 1B**). The 6
climate data layers were from the South African Atlas of Agrohydrology and Climatology
(Schulze 1997 with higher resolution for 4 and 5 below) (Schulze 2007 for the rest) and
were based on 30 year climate averages, extremes, or variances. The simple or
composite climate layers used were (1) mean annual rainfall, (2) average daily
maximum temperature for January (summer), (3) average daily minimum temperature
for July (winter), (4) rainfall seasonality, measured as the percent of annual precipitation
to fall within a single month (5) inter-annual rainfall variability, measured as the
coefficient of variation in mean annual rainfall across years and (6) extreme temperature
range within a year, measured as the difference between January maximum and July
minimum temperatures (2 minus 3), which is strongly correlated with the difference
between the maximum minus the minimum monthly temperature using 50 year
extremes (R=0.95). Our seventh variable was altitude, which was extracted separately
from ASTER GDEM (Daac 2011). Most of these layers have been used in our earlier
work on *P. repens*, except for (5) and (6), which were added to join rainfall seasonality
(4) as measures of environmental heterogeneity. Locations of each population along
these axes can be estimated from **Figure 1B** or viewed in **Table S1**. Only two pairs of
climate variables were strongly correlated with each other: July minimum temperature
with altitude (Pearson's R= -0.91) and January maximum temperature with annual
temperature range (annual max. minus min. temp., R=0.70; see **Table S2** for all
correlation coefficients). Rainfall seasonality in our dataset is positively associated with
wintertime rainfall (total for Dec-Feb, Pearson's R=0.80) and negatively associated with
summertime rainfall (total for June-August, Pearson's R=-0.90).

Experimental setup and dry-down treatments

In late July 2014, we germinated 938 seeds from 114 maternal lines (per population
means: 117 seeds from 14 lines) for a dry-down experiment. Seeds were sown on low
nutrient mix (50% bark, 50% sand) in trays of 100 individual plugs 2x2 cm in size. Low-
nutrient mix is standard for growing *Protea* species because they naturally occur on low-
nutrient soils. Trays were placed in a long-day growth chamber set for 8 hours of 8℃
(nighttime) and 16 hours of 20℃ (daytime). After 1.5 months, 419 seeds had

germinated, and these were subsequently transplanted into small, deep pots (5 cm diameter x 17 cm deep) in trays of 50 each, in the same low nutrient mix. Transplants were given an additional two weeks indoors under growth lights before being transferred to a single greenhouse bench at Nicholls State University Farm, Thibodaux, LA.

Two weeks post-greenhouse transfer and two weeks prior to the dry-down treatment, 316 seedlings from 8 populations were reorganized into thirteen trays of 25 in a completely randomized design. The drought experiment began in late October 2014, when greenhouse conditions were warm but relatively dry, and thus most similar to summer conditions in their native range. The 2.5 months-old seedlings were assigned to one of two treatments: regular watering (~every 3 days) or 12 days of water deprivation. Efforts were made to include a replicate from each maternal line within each treatment, which we succeed at for 80% of lines. Total sample size ranges from 19 to 20 plants per population per treatment.

Trait measurements and transcriptome sampling

At 6 days and 12 days into the drought experiment, plants were measured for functional traits and leaves were sampled for transcriptomics. Our sampling design is depicted in **Figure S1**. Functional traits that were measured at both time-points included plant height (cm), stomatal conductance, leaf area, leaf length:width ratio, specific leaf area (fresh leaf area/dry leaf mass), stomatal density (number per mm^2), stomatal pore length (mm) and stem pigmentation. Stem pigmentation, measured as stem hue, was derived using Endler's segment classification (Maia *et al.* 2013), based on the average of two reflectance spectra readings (400-700 nm) taken at mid-stem (JAZ, Ocean Optics). On the segment classification scale for hue, higher values represent redder stems that contain higher levels of anthocyanin. Stomatal conductance was measured using a steady-state leaf porometer (SC-1, Decagon) on one leaf per plant, with an additional adjacent leaf measured simultaneously if the chamber was not filled by a single leaf due to small leaf size. All leaf measurements were taken from the most recently fully expanded leaf from the stem apex. Immediately following conductance measurement, the leaf was removed to measure leaf area, width, length and stomatal traits. Fresh leaves were digitally scanned shortly after harvest at 400 dpi on a flatbed scanner. After scanning, acrylic peels were taken from the adaxial leaf surface and placed on cellophane, which were later viewed under a light microscope at 40x for stomatal measurements (see Carlson *et al.* 2016). Although height and pigmentation were measured on all plants on both day 6 and day 12 of the treatments (mean sample size

per population=39), all other traits were measured on only a subset of plants on each of these days, typically in association with the transcriptomic data (10-20 individuals per population for each day). Two additional variables were calculated by subtracting day 6 from day 12 values on the same plant: pigment accumulation (day 12 hue - day 6 hue) and growth (day 12 height - day 6 height). A principle component analysis was performed on the traits measured separately for day 6 and day 12, and shown in **Figure 2C-D**.

A smaller subset of plants had one leaf per plant harvested for whole transcriptome sequencing and gene expression analysis (n=120 plants total, or 4 per population per treatment per day, except no ALC on day 12 due to insufficient number of individuals because of lower germination in this population). Plants with leaves removed on day 6 were not sampled for transcriptomics on day 12, to avoid possible shifts in gene expression caused by leaf removal. As a consequence, gene expression and most traits were measured in one set of plants on day 6 and in a separate set on day 12. Transcriptome sampling took place prior to any other measurements on a given day and consisted of collecting one leaf for RNA extraction per plant (the 2nd or 3rd youngest leaf from the tip) into a 1 mL tube and snap-freezing immediately in liquid nitrogen. These samples were stored at -80℃ until library preparation. All sampling for transcriptome sequencing was done between 10 AM and 11 AM, and functional trait measurements were done between 10AM and 2PM each day. Both transcriptomics and other trait measurements were done in a random order. On day 12, we harvested a subset of plants for root length and carbohydrate storage measurements (n=9 per population including the 4 transcriptome plants and 5 additional plants).

After completing all trait and transcriptome sampling, we cut plants at the root-shoot junction and placed aboveground tissues in a 10 mL tube and snap-froze the samples in liquid nitrogen for future carbohydrate analyses. Belowground tissues were gently removed from the soil, rinsed clean and stored in a separate tube and also snap-frozen. For these 9 plants per population, we also measured the length of the longest root. Tubes containing plant tissues were stored in a -80℃ freezer for ~1 month, then freeze-dried for biomass and carbohydrate analyses. Dried tissues for above- and belowground material was measured and ground. Ten to hundred and fifty mg of ground material was used for carbohydrate analyses as described in (Akman *et al.* 2012). Briefly, carbohydrates and starch were extracted with 70% methanol solution. The extract was used for soluble carbohydrate content measurements and the pellet

including starch was treated enzymatically with α-amylase and amyloglucosidase

resulting in hexose sugars. Soluble carbohydrates and hexose sugar concentration was

measured with a modified version of anthrone method using fixed glucose standards.

We also measured biomass but biomass and carbohydrate content were highly

correlated, and thus only carbohydrate content results are shown (dry mass and soluble

sugars Pearson's R = 0.90 and 0.94; starch Pearson's R = 0.64 and 0.81 for

belowground and aboveground, respectively).

Gene expression analyses and bioinformatics

Please see Supplementary Materials and Methods for transcriptome library preparation

and sequencing details. Raw reads were trimmed and demultiplexed with trimmomatic

(Bolger *et al.* 2014) and fastx toolkit (Hannon, 2010). By using bowtie (Langmead &

Salzberg 2012), trimmed reads were mapped to a previously published de-novo

transcriptome of *P. repens* (Akman *et al.* 2016) assembled by using sequences of three

individuals from 19 populations that also include all of our focal populations. Before

reads were mapped to the transcriptome, a further clustering of the contigs were done

using cd-hit (Fu *et al.* 2012). The final number of contigs in this final assembly was

168515. Read counts were extracted using eXpress (Pachter 2011), and ranged

between 2.46 to 21.80 million reads per sample with an average of 6.02 million reads.

Differential gene expression analyses were performed using edgeR. Contigs with <1

counts per million (cpm) were excluded from the analyses. For each time point,

differentially expressed genes per population were revealed using pairwise comparisons

using negative binomial models. The common set of differentially expressed genes was

uncovered by extracting the overlap of differentially expressed genes for each

population at day 12 only. We then used linear models in edgeR, by including

sequencing lane as a fixed factor, to detect contigs that showed variation between

treatments and among populations (population x treatment). Multi-dimensional scaling

(MDS) analysis was performed for each time point to visualize general patterns of gene

expression for all the samples using the top 500 genes that explain the variation the

most (**Figure 2A-B and Figure S6**).

Co-expression gene networks were computed using WGCNA (Langfelder &

Horvath 2008) with consensus clustering option for the 2 time-points. Normalized and

cpm-filtered read counts were transformed with "voom" in limma package for using in

WGCNA. A soft-threshold of 6 was used for clustering and 14 co-expressed gene

networks were constructed. The number of genes within each gene network varied

between 49 (GN13) to 7682 (GN1). The gene network "grey" (called GN0 in this study) as constructed by WGCNA for genes that do not fit any gene network has 3124 genes. A representative "eigen gene" value, which is essentially the first principal component axis for all genes within the network, was used for further analyses of climate and trait associations. By using gene orthology of our genes with Arabidopsis (Akman *et al.* 2016), we also tested for Gene Ontology (GO) enrichment for the 13 networks. This analysis was performed in Gorilla (Eden *et al.* 2009), using the network genes as target set and the whole gene set as the background. Contigs within each module, GO enrichments and the genes with the highest module membership for a single network are provided in **Table S3**.

Statistical analyses

How do populations respond to drought and how do these responses vary?

We tested whether drought significantly affected functional and performance traits in *Protea repens* using linear mixed models. We used a total of 23 plant metrics in our analyses, including 8 traits on day 6, and 14 traits or performance indices on day 12. We analyzed each trait on each day separately, because preliminary analyses on both days combined revealed significant time-point effects for almost every response variable. Each model included the fixed effects of treatment, population, and the population by treatment interaction (G x E) and random effects of maternal line. Models for stomatal conductance included the added random effect of air temperature during measurement. In these models, the treatment-by-population interaction term indicates whether there was differential plasticity among populations, or in other words, that plants from different populations responded differently to the treatment (G x E effect). These analyses were performed using the lme4 package (Bates *et al.* 2012) in R 3.1 (Team 2014). To obtain p-values for each fixed effect, we successively dropped one predictor from the full model and used a chi-square test to compare AIC scores between models. Given the large number of statistical tests that this entailed, we also adjusted p-values according to estimated false discovery rates (R package "fdrtools").

Gene expression analyses followed the same general approach as for traits. As response variables, these analyses focused on levels of expression in the 14 co-expressed gene networks (GN) that were produced through the gene clustering analysis (see *Gene expression analyses and bioinformatics* section above). Models for gene networks included fixed effects of treatment, population, and their interaction, random

effects for maternal line, and an additional random effect for sequencing lane (1 or 2).
We also tested for genes that are differentially expressed among populations for the two
treatments and time-points separately. A Venn diagram (Oliveros 2016) showing the
genes in these four groups are given in **Figure 3B** . For example, the group of genes
that are differentially expressed among populations under drought at day 6 is shown in
purple color on the diagram. Gene ontology analysis was performed for detecting
enrichments in these four groups using GOrilla as explained above.

Do the climatic conditions of the source sites covary with traits, gene expression and
their plasticity?

To test whether characteristics of the home climate predicted cross-population variation
in traits and gene expression, we used mixed effect models. In these models, traits and
gene networks (GN) were response variables, the climate variables were fixed effect
terms and maternal line nested in population was used as random effect variables. For
testing the significance of the fixed factors, we calculated p-values based on chi-square
tests as explained above. FDR corrections were applied to p-values in three groups:
functional traits, performance traits and gene networks. Day 6 and Day 12 FDR
corrections were also applied separately. For variables with significant climate and
treatment interaction, each treatment was also analyzed separately to estimate slopes
and test their significance within treatments. These are depicted as divided squares in
Figure 4.

To characterize plasticity for individual traits and to quantify the direction and
magnitude of this plasticity, we used maternal line plasticity indices. For each maternal
family, we calculated plasticity as the family mean trait value in the wet treatment minus
the family mean trait value in the drought treatment, divided by the overall family mean
trait value. We used mixed effect models to determine whether characteristics of the
home climate predicted cross-population variation in plasticity of traits and gene
expression. In our analyses, plasticity in traits and gene networks (GN) were response
variables, the 7 climate variables were fixed effect variables and population was
included as a random effect. Each climate variable was tested against each response
variable in a separate model, with day 6 and day 12 measurements analyzed separately
(lmer, R program). FDR corrections were applied to p-values in three groups: functional
traits, performance traits and gene networks. Day 6 and Day 12 FDR corrections were
also applied separately.

RESULTS

How do populations respond to drought and how do these responses vary?

Drought responses are stronger in gene expression patterns compared to traits

Six days into the experiment, only three of the 8 traits differed significantly between treatments: unwatered plants had shorter stems, lower stomatal conductance and more variable SLA. At the later stage, treatment effects became pronounced for more traits measured (12 of 15): unwatered plants had lower stomatal conductance, decreased height, slower growth (from day 6 to day 12), larger SLA, smaller leaf area, denser stomata, smaller stomatal pore size, increasing pigment accumulation, less carbohydrates aboveground (but not belowground). In the earlier stages of the stress imposition, morphological and physiological traits did not show responses as pronounced as those observed in gene expression patterns (**Figure 2**). In contrast to most of the traits, the treatment effect was significant for over half of the gene networks already at day 6 into the treatment. This trend of more pronounced gene expression response compared to morpho-/physiological traits is visible both in co-expression gene networks (**Table 2**, **Figure S2**) and Multi-dimensional Scaling (MDS) analyses (**Figure 2A-B**), which indicates a lag between the adjustments in gene expression and trait onset. However, we should also note that the MDS for gene expression analyses is based on the top 500 genes that explain the most variation, and we might not have measured morpho-physiological traits that might also respond as quickly. For both gene expression patterns as seen in MDS and morpho-physiological traits in PC analyses, axis 1 and 2 explain the variation between treatments and among populations, respectively. Notably, population V (Baviaanskloof, a high-elevation eastern site) does not show a clear treatment effect in traits at the later stage, although the effect is quite pronounced in the gene expression data (**Figure 2B&D** and **3A**).

At day 6 into the treatment, unwatered plants had higher values for gene networks 2, 4, 5, 6, and 7 and lower values for gene networks 1, 9 and 11, relative to watered plants (**Table 2**, **Figure S4**). These modules show gene enrichment in GO categories for ATPase activity (GN2), oxidoreductase activity (GN4), cytoskeleton organization (GN5), photorespiration (GN6), response to high light intensity and temperature (GN7; note that p-value<0.0005, fdr-corrected p-value is not significant), cell wall organization (GN1) and pyridine-containing compound metabolic process (GN9). As the stress advanced after 12 days into the drought treatment, watered and

unwatered plants differed significantly for most traits and gene networks. Significant

treatment responses were detected for 10 of 14 gene networks, 8 of 10 functional traits,

and 4 of 5 performance measures (**Tables 1-3**, **Figures S3-5**). In addition to the gene

networks with treatments effects at day 6, GN3 (tropism and carbohydrate metabolism

related, and shows associations with leaf area, height and aboveground soluble

carbohydrates) and GN13 (homeostasis related) also had evident treatments effects at

day 12, with GN3 having lower values and GN13 higher values under drought.

Drought up-regulates stress related genes and down-regulates growth related genes

The number of genes differentially regulated between earlier and later stages of the

stress and among populations differed significantly (**Figure 3B**). For example,

population G (Garcia's Pass) does not have any genes differentially expressed at day 6,

indicating either a lower level of stress or lack of accurate perception and timely signal

transduction of the environmental cues to gene expression phenotypes. In contrast,

population B (Bredasdorp) had several hundred differentially expressed genes by day 6.

All populations had at least several hundred differentially expressed genes by day 12.

Two populations (C-Cederberg and S-Swartberg) showed few differentially expressed

genes initially, but over 1500 differentially expressed genes by day 12 (**Figure 3A**).

Although the number of genes regulated upon drought stress was highly variable, we

found 110 and 63 genes to be consistently up- or down- regulated, respectively, among

populations (**Table S4**). Homology of these genes with Arabidopsis show that up-

regulated genes include heat shock proteins, auxin signalling and ABA catabolism

genes, genes related to drought stress and photosynthesis, and a leaf amylase;

whereas down-regulated genes included ARF genes (involved in cell elongation and

division), and salt stress-, fatty acid and amino acid biosynthesis-related genes.

Population differentiation in stress response becomes more evident as the stress
advances

A population effect is evident in most of the traits, 5 out of 8 functional traits and 6 out of

7 performance traits were significantly different between the populations at day 6, and

all traits except for root length showed a population effect at day 12 (**Table 1 and 3**).

We also observed G x E effects (variation in population responses to drought) to a

limited extent on day 6, and to a greater extent on day 12 (**Table 1**). At day 6, only SLA

and gene network 13 (enriched for homeostasis GO term) showed variation in

responses to drought across populations (i.e., population x treatment interaction

significant; **Table 1**). Specific leaf area was lower under drought for four, mostly western populations (R, G, C and F), but it was higher under drought in the remaining 4 populations (V, A, B, S;**Table 1**, **Figure S3**). At day 12, populations showed variation in their response to drought for SLA, stomatal size, pigment accumulation, stomatal conductance, stem growth rate, starch biomass aboveground, sugar biomass above and belowground. In addition to GN13 at day 6, GN7 expression (linked to high light intensity and high temperature GO term) also showed a G x E effect at day 12 (**Table 2**).

In addition to gene network responses, we also tested for gene expression responses in single genes. Differential expression patterns of some genes varied across populations (significant G x E interaction), including 13 contigs on day 6, rising to 23 on day 12. These included genes associated with ATP synthase, cell redox homeostasis, translation, response to cadmium ion, cellulose biosynthesis and heat shock response. For example, contig 9970 showing homology with a heat shock gene in Arabidopsis (AT5G12020) is up-regulated in all the populations 12 days after the treatment started, but this increase is variable among populations (**Figure S7**).

For each treatment, we also tested for genes expressed differentially among populations (testing for population effect) and time-points separately. This analysis reveals which genes are differentially expressed among populations, indicating population differentiation, within a treatment and a timepoint. With these analyses, we find 851 and 668 genes to be differentially regulated among populations for the control treatment, and 717 and 786 for drought treatment at day 6 and 12, respectively (FDR-corrected p-value<0.01). We also observed some overlap between these groups (**Figure 3B**); 45 of the genes being differentially expressed among populations in both of the treatments for both days. Interestingly, the majority of genes differentially expressed in a treatment did not overlap between the two time-points and this holds true for both of the treatments (e.g. purple and pink in **Figure 3B**). Likewise, the genes differentially expressed in the different treatments did not show overlap within a single time-point (e.g. purple and yellow in **Figure 3B**). These results show that together with a strong treatment effect, there is a pronounced time-point effect in our experiment for gene expression (note that the plants were all harvested at the same time of the day but 6 days apart). This is also evident from our data for both traits and gene expression as there is a significant time-point effect or interaction involving the time-point for most traits and gene expression networks. We also found that the differentially expressed

genes among populations in watered conditions show enrichments for processes related to translation and peptide biosynthesis, whereas in dry conditions we find oxidative and heat stress related categories.

Do the climatic conditions of the source sites covary with traits, gene expression and their plasticity?

Gene networks and functional traits correlate with temperature-related source-site climate variables

In order to assess climatically driven trait and gene expression variation among populations, we tested for correlations between source-site climate variables and co-expression gene networks and functional traits. We found several gene networks that covary with climatic variables (**Figure 4**; boxed combinations). These include GN 3, 4, 5, 9 and 13 at day 6, and GN3, 4, 5, 6, 7, 10, 12 and 13 at day 12. Most of these gene networks showed strong correlations with elevation and the closely related variables July (winter) minimum temperature and January (summer) maximum temperature. We also observed correlations between mean annual precipitation and GN6, 10 and 13, and stomatal conductance. Of the performance traits, stem growth showed correlations with elevation, summer maximum and winter minimum temperature. Aboveground starch content co-varied with elevation, mean annual precipitation and intra-annual precipitation variation. Similarly, aboveground soluble carbohydrates showed associations with elevation.

We tested for climate by treatment interaction which revealed if associations between traits and the climate are significant for only one of the treatments. We found that gene expression patterns for GN7 at day 12 show a positive correlation with elevation (and minimum winter temperature) only for plants under drought conditions. Contrastingly, the correlations between GN12 and elevation, and GN13 and mean annual precipitation are significant only for watered plants. Similarly for stem growth and aboveground starch, the correlations with elevation are significant only for watered plants, and stomatal conductance shows a correlation with mean annual precipitation only under watered conditions.

Plasticity differentiation is associated with climate

We tested if plasticity in traits and gene expression reflect signatures of climatic effects. In order to do this, we used a subset of climate and trait pairs that already show

trait/gene expression variation among populations, as well as trait/ gene expression patterns that show a climate x treatment effect (associations between the climate holds true only at a single treatment). At day 6, plasticity in GN5 (related to cytoskeleton organization) is the only gene network that show a correlation with a climatic variable: low elevation populations showed higher plasticity in gene expression patterns of this gene network (**Table 4**). At day 12, we found that plasticity in GN10 (related to translation) and GN13 (linked to homeostasis) covaries with mean annual precipitation. For GN10, populations from both the driest and the wettest sites show higher plasticity (note that plasticity has both positive and negative values dependent on the directionality of the response), and populations from wetter sites show higher plasticity for GN13 (**Table 4**, **Figure 5D**). Additionally, plasticity in stem growth correlates with both elevation and summer maximum temperature. High plasticity in growth is observed in populations from low elevation and warmer sites (**Figure 5A**). Similar to GN13, plasticity in leaf area shows correlations with mean annual precipitation with higher plasticity in populations from wetter sites (**Figure 5B**). Leaf conductance and pigment accumulation shows correlations with temperature heterogeneity (temperature range within years). For both pigmentation and conductance, higher temperature heterogeneity, which is correlated with maximum summer temperature, leads to higher plasticity (**Figure 5C**). Finally, pigment accumulation plasticity also shows correlations with summer maximum temperature, with populations from coolest and warmest sites showing the highest plasticity. However, pigment-related associations are driven by a single population in Cederberg, the most northern population we studied. We should also note that this trend is driven by the direction of plasticity in this population for pigment content, which is opposite to what we observe in all the other populations:, plants from the Cederberg population have higher pigment under drought and lower pigments when watered.

DISCUSSION

Local adaptation can lead to divergence among populations along environmental gradients. This divergence in genetic variation is often reflected in the phenotypic variation in morphology, physiology as well as the underlying gene expression variation. Plasticity in these, as any other phenotypic character, can also be under selection, creating matching patterns between plasticity differentiation and the source site climatic variation among populations. In this study, we combined a greenhouse drought experiment with extensive measurements of physio/morphological traits and RNA-seq

data for 8 common sugarbush populations collected along an environmental gradient to uncover patterns of climatic selection on these traits and their plasticity. We show that functional traits and gene expression patterns are associated with the climate, and these associations are highly dependent on the environmental conditions the traits are measured in. More importantly, we show that plasticity in the functional traits and gene expression patterns also are associated with climatic conditions of the source population sites. Together, our results suggest that local adaptation affects evolution of both trait means and their plasticity along a climate gradient in *Protea repens*.

Drought imposes a complex stress regulating many gene expression networks

Drought has been a strong driver of plant evolution in the Cape Floristic Region and other Mediterranean climate zones (Allsopp *et al.* 2014). When water availability is limited, as it typically is for several months during the summer in most of the region, plants experience a suite of complex stresses such as accumulation of reactive oxygen species, starvation, lower photosynthesis rates and turgor loss (Ramachandra Reddy *et al.* 2004; West *et al.* 2012). Therefore, selection is expected to act on several genes involved in these stresses, enabling plants to cope with the severe effects of drought. Indeed, several studies have shown that various pathways including ABA-dependent pathways, carbohydrate metabolism, ROS scavenging and photosynthesis are regulated under drought (Miller *et al.* 2010; Heschel *et al.* 2014). In concordance with drought being a complex stress acting on several pathways, we observe regulation of 10 out of 13 gene networks under drought conditions. For example, the genes with the highest gene network membership of GN5 (related to cytoskeleton organization) are downregulated under drought in all the populations, and this gene network also shows correlations with both elevation and traits such as height, leaf area and aboveground sugar content. Cytoskeleton organization related pathways were also shown to be highly hampered during drought stress in Arabidopsis (Bhaskara *et al.* 2017), linking this process to reduced growth. These results make this gene network a prime candidate for controlling elevation-dependent growth patterns. A general pattern similar to that of GN5 genes is seen in other gene networks important for growth (such as GN1 related to cell wall organization and other carbohydrate metabolism related gene networks, and shows significant correlations with conductance). This "shut-off" of growth also explains the shorter stems and smaller plant carbohydrate biomass aboveground under 12 days of drought in all of the populations. Additionally, the genes that were consistently up-regulated among all the populations were genes involved in stress response. Together

these results indicate a trade-off in which resources are channeled from growth to stress acclimation. A similar trade-off has been observed across many plant species for herbivory (Kempel *et al.* 2011; Züst *et al.* 2015), cold stress (Loehle 1998) and drought (Haugen *et al.* 2008). We also detected reduced stomatal conductance in all populations in the drought treatment, and thus a potential drop in photosynthesis might also further contribute to the energy deficit, which might hamper growth and efficient acclimation to the stress. The only gene network, GN6 (enriched for photorespiration related genes) that shows a correlation with conductance at the later stage of the stress is also correlated with leaf area, aboveground starch and above and below ground soluble carbohydrates. Interestingly, we also find a correlation between this gene network and mean annual precipitation suggesting local adaptation to water availability via regulation of these genes.

Drought induces smaller and thinner leaves

Upon 12 days of drought stress, we see smaller leaves (except for population V) and higher SLA (thinner leaves). This reduction in leaf size overall is potentially due to a reduced investment in photosynthetic architecture. Previously, we showed that leaf area measured in a common garden increases with source site precipitation, potentially maximizing photosynthesis with water availability (Carlson *et al.* 2016). These patterns are observed in other species within micro and macro scales. For example, Peppe et al. (2011) showed that leaf size decreases with precipitation across many species which also holds true here within the single species *Protea repens*. Additionally, we also see a drop in stomatal pore size coupled with a higher stomatal density under drought conditions. Previous studies show both positive and negative trends between aridity and stomatal density or no relationship at all (Stenström *et al.* 2002; Pearce *et al.* 2006; Yu *et al.* 2008; Franks & Beerling 2009; Skelton *et al.* 2012; Carins Murphy *et al.* 2014). Our finding is consistent with a common garden study of 19 *P. repens* populations sourced from a range of climates, in that plants from hotter sites with drier summers had smaller and denser stomates (Carlson *et al.* 2016).

Associations with the climate are context dependent

When there are fitness consequences of a trait state that is induced conditionally in response to environmental stressors, with no fitness costs/benefits when those stressors are absent, genetic variation that controls this modified trait state (plastic response) will be under selection only under the inducing stressful environment (Van

Dyken & Wade 2010; Ledón-Rettig *et al.* 2014). Strikingly, we observed climatic correlations in gene expression and traits that only exist in either one of the treatments (**Figure 4**). Gene network 7 (GN7), for example, shows a correlation with elevation and winter minimum temperature only when the plants are under drought stress. Gene ontology enrichments in this module point to regulation by high light and temperature stress. Overall patterns show that best representative gene of this network (with the highest similarity to eigen gene of PC1) show a higher expression in the high elevation populations. A similar gene network in our previous study in a common garden was also enriched for GO terms related to high light intensity and temperature, and this module showed correlations with stem diameter and stomatal density, but surprisingly not with any climate variables. Gene network 7 in this study also shows significant correlations with height at both timepoints and SLA in the earlier timepoint (**Table S5**). Populations from warmer sites show lower GN7 values (note that within a gene network there could be genes that are regulated in opposite directions among populations, and thus it is not possible to infer if a gene network is up- or down-regulated) which translates to taller plants and thicker leaves. The top members of this gene network with Arabidopsis homology are down-regulated in populations coming from higher elevation; a nucleotide-sugar transporter gene (AT4G32390), a zinc-finger family protein (AT2G38970), MTP8 (AT3G58060), 4CL3 involved in last step of phenylpropanoid pathway - the upstream pathway anthocyanins are derived from (AT1G65060) and PRA1.B1 (AT3G56110). We also observe a weak correlation between stem pigmentation and temperature. Plants from warmer sites (higher maximum summer temperature) have higher pigmentation accumulation (p-value =0.029, fdr-corrected value = 0.106). Previous studies showed a negative effect of heat stress on plant growth (Olszyk *et al.* 1998; Hasanuzzaman *et al.* 2013; Bita & Gerats 2013), and anthocyanin accumulation as a protective mechanism against heat, cold, oxidative stress and UV radiation (Hatier & Gould 2009; Steyn *et al.* 2009; Zeng *et al.* 2010; Wang *et al.* 2013). The higher amount of anthocyanins accumulated in populations originating from warmer sites in this study and their taller stature may accordingly be due to stronger selection in warmer environments for production of anthocyanins to provide protection against reactive oxygen species. In a parallel finding, under drought conditions, differentially expressed genes among populations were enriched for only oxidative stress related genes, suggesting there has also been selection on expression of these genes arising from different levels of heat/drought stress along the climate gradient.

Contrastingly, GN13 shows a correlation with mean annual precipitation only in
watered conditions. This gene network also shows correlations with leaf area, SLA,
stomatal pore size and aboveground carbohydrate content. Gene network 13 is a
module related to homeostasis and the best representative of this module shows
homology to an Arabidopsis heat shock protein (AT5G59720). Heat shock proteins are
shown to be involved in responses to stresses such as heat, cold, UV radiation, salt and
drought (Chen *et al.* 2018). Under control conditions, plants from wetter source sites
express genes in this network less compared to populations from drier sites, and show
a bigger up-regulation, eventually reaching similar levels of expression for all
populations under drought. In drier environments, higher constitutive expression of
these genes might be crucial for a prompt response to drought via regulation of stomatal
pore size, leaf size and thickness. However, populations that do not experience drought
as often might not benefit from constitutive expression of these genes, creating a
mismatch between costs and benefits. These patterns indicate local adaptation in
plasticity of these genes through a relaxed selection under wet environments, and
directional selection under drought.

Similar to GN13, we also observe a correlation between GN12 and elevation
under wet conditions: populations from high elevation show higher expression of genes
in this network. Although this gene network is not enriched for gene ontologies, the
gene network is constituted of photosynthesis and chloroplast related genes, and shows
correlations with leaf area. Unlike GN13, we observe a variable population response
under drought: some populations have lower eigen gene values and some have higher
under drought conditions compared to controls. This implies that some genes are up
and some are down-regulated in a population specific manner. Under drought it is likely
that most populations reduced their photosynthetic rate to very low values and thus
patterns are not present, as evidenced from the generally low conductance values in the
drought treatment.

Temperature exerts a stronger selection pressure than precipitation

Previously, we showed that functional traits and gene expression patterns as measured
in a common garden with older plants (1.5 year old plants vs 2 months-old plants in this
study) correlated with source site climate. Although using fewer populations in this
study, we observed similar patterns in those correlations. For example, in both studies
we observe associations between gene networks and minimum winter temperature,
elevation and mean annual precipitation. We also found that the climate variable that

has the most correlations with the gene networks to be elevation/minimum winter
temperature (2 variables highly correlated, Pearson's R = -0.91, **Table S2**), which was
also true for the common garden study. Although Protea populations throughout the
CFR should experience drought at least occasionally if not frequently, the drought
metrics used in this study are not as strong predictors of trait variation as
temperature/elevation is under the tested conditions (drought and control in a
greenhouse and a common garden). Even if our selection of leaf traits was biased
away from those likely to reflect drought adaptations (which we consider unlikely), this
could not be the case for gene expression networks, which include all major groupings
of coexpressed genes detected. Gene expression networks should therefore reflect an
unbiased pattern in plant responses, and these networks also had a high number of
correlations with home-site temperature or elevation.

Possible explanations for the dominance in temperature/elevation correlations
are (1) we have not identified the most influential metrics of drought stress in the CFR,
such that we fail to detect these associations; or (2) temperature-gradients across the
CFR have shaped trait evolution in *Protea repens* more strongly than rainfall-gradients
recently. This could have arisen because *P. repens* plants experience drought less
frequently than expected due to rapidly growing roots that reach and remain in the water
table despite water shortages at the surface. It could also be because drought is an
inconstant selection pressure, relative to temperature. Populations in the CFR
experience drought periodically, and although some of the populations we studied
experienced a decrease in precipitation over the past 115 years, this effect is not as
strong as the increase in temperature over the same period (**Figure 6**, Mason & Jury
1997; Kruger & Shongwe 2004). Compared to a periodic drought cycle that varies
widely in strength, this temperature increase potentially imposes a more stable and
continuous stress and thus a stronger selection force. Additionally, the fact that
precipitation patterns are affected in a more variable way compared to more stable
increases in temperature might make it harder to detect patterns of selection in
response to precipitation changes across climate gradients even under drought
conditions as in our experiments. For example, a recent meta-analysis of 21 plant traits
from 25354 species concluded that temperature is a stronger driver of plant trait
evolution compared to precipitation (Moles *et al.* 2014).

Carbohydrate reserves play a role in survival

Surprisingly, we found a positive correlation between survival under drought and precipitation; plants from wetter sites survived longer under drought. Populations from wetter sites also showed higher stomatal conductance and shoot starch indicating higher photosynthetic rates. The fact that populations from wetter sites surviving better under drought is counterintuitive. However, in a sudden drought onset, like the one imposed in this experiment, carbohydrate reserves might be a decisive factor in survival, especially in earlier stages of development, and thus if plants from wetter sites have built up larger reserves, they may survive longer (O'Brien *et al.* 2014).

Both climatic heterogeneity and favorable environmental conditions shape plasticity evolution

Here, we also tested if plasticity is favored and thus selected for in heterogeneous and/or milder climatic conditions. A heterogeneous environment in theory should promote plasticity evolution as this can allow for a better match between the phenotype and the changing environment, potentially surpassing cost of plasticity. Similarly, in mild environments, plasticity can arise through a relaxed selection in which a mismatch between the phenotype and the climate does not impose strong fitness costs. We found evidence for both of these patterns for the functional traits and the gene expression patterns. These results support that plasticity can evolve under both milder and heterogeneous conditions. However in this study, we have more evidence for growth plasticity evolving under milder conditions than heterogeneous environments, i.e. higher plasticity for growth and GN5 (cytoskeleton organization) in populations from lower elevation/warmer sites (representing milder climates). Although heat stress can be detrimental, in the CFR cold stress may impose a stronger selective pressure for growth (Wilson *et al.* 2015). In fact, in a study of 56 Proteaceae species looking at climate and plant trait correlations, Mitchell et al, showed the most strongly supported evolutionary correlations to be between winter minimum temperature (next to mean annual temperature and elevation) and plant height (Mitchell *et al.* 2018). They also note that this negative effect of elevation on plant height might be coupled potentially with shallower soils in the high elevation sites (Campbell & Werger 1988). These combined effects might create harsher environments which increase the cost of mismatch between phenotype and the environment, thus decreasing fitness of more plastic individuals. We also find higher plasticity in wetter sites for GN13 and leaf area, and again these sites represent milder climates in this drought-prone region.

Similar to trait means, we also find more associations between temperature

related variables and trait plasticity, once again suggesting that temperature may create

a stronger or more consistent selective force compared to precipitation in this system.

Interestingly, the only correlations we observe for heterogeneity are also related to

temperature. Both plasticity in conductance and pigment content increases with

increasing heterogeneity. In environments where temperature change across seasons

is high, higher plasticity in stomatal conductance (thus photosynthesis and transpiration

rates) might decrease the phenotype-environment mismatch and increase plant fitness,

thus be selected for in these environments. Likewise, adjusting levels of anthocyanins

through plasticity in response to temperature changes might effectively protect plants

against abiotic stressors, thus promoting selection on higher plasticity.

CONCLUSION

Here, we show that both phenotypic traits and gene expression, and their plasticity can

be under selection along a climate gradient. However, detecting associations between

these traits and the environmental variables can be a challenging task as these

associations are context-dependent, and thus can be detected only when they are

measured under certain conditions. We also found that trait variation can be explained

better with temperature related-variables rather than precipitation related variables,

potentially due to the fact that the recent temperature increases associated with climate

change impose a stronger and more continuous selective pressure compared to the

more variable changes in precipitation. This study also provides insights into the drivers

of plasticity evolution: we found more evidence for selection for plasticity in mild

environments (or selection against plasticity in harsher environments) compared to

directional selection in heterogeneous environments. We also demonstrate that

integration of gene expression variables into studies of climate trait associations proves

valuable for detecting molecular mechanisms involved in selection and by representing

a comprehensive palette of traits that are shaped by both neutral and selective

evolutionary forces.

ACKNOWLEDGEMENTS

We thank Chris Adams, Megan Varvaro and Sarah Soorya for their assistance with

plant care and the drought experiments, Tony Rebelo, Guy Midgley and Louise Nurrish

for their assistance with seed collection. We also thank the Blackman Lab and Carl

Schlichting for their suggestions to an earlier version of the manuscript. This work was

supported by Dimensions of Biodiversity awarded to Carl Schlichting, Cynthia S. Jones

and Kent E. Holsinger (award no: 1046328) and Andrew M. Latimer (award no:

1045985).

REFERENCES

Ackerly DD, Dudley SA, Sultan SE *et al.* (2000) The evolution of plant ecophysiological

traits: Recent advances and future directions: new research addresses natural

selection, genetic constraints, and the adaptive evolution of plant ecophysiological

traits. *Bioscience*, **50**, 979–995.

Akman M, Bhikharie AV, McLean EH *et al.* (2012) Wait or escape? Contrasting

submergence tolerance strategies of Rorippa amphibia, Rorippa sylvestris and their

hybrid. *Annals of botany*, **109**, 1263–1276.

Akman M, Carlson JE, Holsinger KE, Latimer AM (2016) Transcriptome sequencing

reveals population differentiation in gene expression linked to functional traits and

environmental gradients in the South African shrub Protea repens. *The New*

phytologist, **210**, 295–309.

Allen CD, Macalady AK, Chenchouni H *et al.* (2010) A global overview of drought and

heat-induced tree mortality reveals emerging climate change risks for forests. *Forest*

ecology and management, **259**, 660–684.

Allsopp N, Colville JF, Verboom GA (2014) *Fynbos: Ecology, Evolution, and*

Conservation of a Megadiverse Region. Oxford University Press.

Alpert P, Simms EL (2002) The relative advantages of plasticity and fixity in different

environments: when is it good for a plant to adjust? *Evolutionary ecology*, **16**, 285–

297.

Anderson JT, Gezon ZJ (2015) Plasticity in functional traits in the context of climate

change: a case study of the subalpine forb Boechera stricta (Brassicaceae). *Global*

change biology, **21**, 1689–1703.

Aubin I, Munson AD, Cardou F *et al.* (2016) Traits to stay, traits to move: a review of

functional traits to assess sensitivity and adaptive capacity of temperate and boreal

trees to climate change. *Environmental Review*, **24**, 164–186.

Barshis DJ, Ladner JT, Oliver TA *et al.* (2013) Genomic basis for coral resilience to

climate change. *Proceedings of the National Academy of Sciences*, **110**, 1387–1392.

Bates D, Maechler M, Bolker B (2012) lme4: Linear mixed-effects models using S4

classes.

Baythavong BS, Stanton ML (2010) Characterizing selection on phenotypic plasticity in
response to natural environmental heterogeneity. *Evolution; international journal of*
organic evolution, **64**, 2904–2920.

Bhaskara GB, Wen T-N, Nguyen TT, Verslues PE (2017) Protein Phosphatase 2Cs and
Microtubule-Associated Stress Protein 1 Control Microtubule Stability, Plant Growth,
and Drought Response. *The Plant cell*, **29**, 169–191.

Bita CE, Gerats T (2013) Plant tolerance to high temperature in a changing environment:
scientific fundamentals and production of heat stress-tolerant crops. *Frontiers in plant*
science, **4**, 273.

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina
Sequence Data. *Bioinformatics* .

Bradshaw AD, Hardwick K (1989) Evolution and stress—genotypic and phenotypic
components. *Biological journal of the Linnean Society. Linnean Society of London*, **37**,
137–155.

Campbell BM, Werger MJA (1988) Plant Form in the Mountains of the Cape, South
Africa. *The Journal of ecology*, **76**, 637–653.

Carins Murphy MR, Jordan GJ, Brodribb TJ (2014) Acclimation to humidity modifies the
link between leaf size and the density of veins and stomata. *Plant, cell & environment*,
37, 124–131.

Carlson JE, Adams CA, Holsinger KE (2016) Intraspecific variation in stomatal traits, leaf
traits and physiology reflects adaptation along aridity gradients in a South African
shrub. *Annals of botany*, **117**, 195–207.

Carlson JE, Holsinger KE (2012) Developmental Plasticity in <italic>Protea</italic> as an
Evolutionary Response to Environmental Clines in the Cape Floristic Region. *PloS*
one, **7**, e52035.

Carlson JE, Holsinger KE, Prunier R (2011) Plant responses to climate in the Cape
Floristic region of South Africa: Evidence for adaptive differentiation in the Proteaceae.
Evolution; international journal of organic evolution, **65**, 108–124.

Chen B, Feder ME, Kang L (2018) Evolution of heat-shock protein expression underlying
adaptive responses to environmental stress. *Molecular ecology*.

Chevin L-M, Lande R, Mace GM (2010) Adaptation, plasticity, and extinction in a
changing environment: towards a predictive theory. *PLoS biology*, **8**, e1000357.

Colicchio JM, Monnahan PJ, Kelly JK, Hileman LC (2015) Gene expression plasticity
resulting from parental leaf damage in Mimulus guttatus. *The New phytologist*, **205**,
894–906.

Cowling RM (2009) From winter wet to winter dry: Fynbos diversity and evolution along
the west–east gradient. *South African journal of botany: official journal of the South*
African Association of Botanists = Suid-Afrikaanse tydskrif vir plantkunde: amptelike
tydskrif van die Suid-Afrikaanse Genootskap van Plantkundiges, **2**, 386–387.
Cuenca Cambronero M, Beasley J, Kissane S, Orsini L (2018) Evolution of thermal
tolerance in multifarious environments. *Molecular ecology*, **27**, 4529–4541.
Daac LP (2011) ASTER L1B. *Sioux Falls, South Dakota: USGS/Earth Resources*
Observation and Science (EROS) Center.
Dayan DI, Crawford DL, Oleksiak MF (2015) Phenotypic plasticity in gene expression
contributes to divergence of locally adapted populations of Fundulus heteroclitus.
Molecular ecology, **24**, 3345–3359.
Dudley SA, Schmitt J (1996) Testing the Adaptive Plasticity Hypothesis: Density-
Dependent Selection on Manipulated Stem Length in Impatiens capensis. *The*
American naturalist, **147**, 445–465.
Duputié A, Rutschmann A, Ronce O, Chuine I (2015) Phenological plasticity will not help
all species adapt to climate change. *Global change biology*, **21**, 3062–3073.
Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z (2009) GOrilla: a tool for discovery
and visualization of enriched GO terms in ranked gene lists. *BMC bioinformatics*, **10**,
48.
Etterson JR (2004) Evolutionary potential of Chamaecrista fasciculata in relation to
climate change. I. Clinal patterns of selection along an environmental gradient in the
great plains. *Evolution; international journal of organic evolution*, **58**, 1446–1458.
Forsman A (2015) Rethinking phenotypic plasticity and its consequences for individuals,
populations and species. *Heredity*, **115**, 276–284.
Franks PJ, Beerling DJ (2009) Maximum leaf conductance driven by CO2 effects on
stomatal size and density over geologic time. *Proceedings of the National Academy of*
Sciences of the United States of America, **106**, 10343–10347.
Fu L, Niu B, Zhu Z, Wu S, Li W (2012) CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* , **28**, 3150–3152.
Geber MA, Griffen LR (2003) Inheritance and Natural Selection on Functional Traits.
International journal of plant sciences, **164**, S21–S42.
Ghalambor CK, Hoke KL, Ruell EW *et al.* (2015) Non-adaptive plasticity potentiates rapid
adaptive evolution of gene expression in nature. *Nature*, **525**, 372–375.
Granados-Cifuentes C, Bellantuono AJ, Ridgway T, Hoegh-Guldberg O, Rodriguez-
Lanetty M (2013) High natural gene expression variation in the reef-building coral

Acropora millepora: potential for acclimative and adaptive plasticity. *BMC genomics*, **14**, 1–12.

Grime JP (1994) The role of plasticity in exploiting environmental heterogeneity. *Exploitation of environmental heterogeneity by plants: ecophysiological processes above-and belowground*, **1**, 19.

Gugger S, Kesselring H, Stöcklin J, Hamann E (2015) Lower plasticity exhibited by high-versus mid-elevation species in their phenological responses to manipulated temperature and drought. *Annals of botany*, **116**, 953–962.

Hannon GJ Fastx-toolkit. 2010. *GNU Affero General Public License*, **706**.

Hasanuzzaman M, Nahar K, Alam MM, Roychowdhury R, Fujita M (2013) Physiological, biochemical, and molecular mechanisms of heat stress tolerance in plants. *International journal of molecular sciences*, **14**, 9643–9684.

Hatier J-HB, Gould KS (2009) Anthocyanin Function in Vegetative Organs. In: *Anthocyanins: Biosynthesis, Functions, and Applications* (eds Winefield C, Davies K, Gould K), pp. 1–19. Springer New York, New York, NY.

Haugen R, Steffes L, Wolf J *et al.* (2008) Evolution of drought tolerance and defense: dependence of tradeoffs on mechanism, environment and defense switching. *Oikos* , **117**, 231–244.

Heschel MS, Evankow A, Wolfson KB, Carlson JE, Holsinger KE (2014) Drought Response Diversification in African Protea Species. *International journal of plant sciences*, **175**, 442–449.

Hewitson BC, Crane RG (2006) Consensus between GCM climate change projections with empirical downscaling: precipitation downscaling over South Africa. *International Journal of Climatology*, **26**, 1315–1337.

Jacobs BS, Latimer AM (2012) Analyzing reaction norm variation in the field vs. greenhouse: Comparing studies of plasticity and its adaptive value in two species of Erodium. *Perspectives in plant ecology, evolution and systematics*, **14**, 325–334.

Kempel A, Schädler M, Chrobock T, Fischer M, van Kleunen M (2011) Tradeoffs associated with constitutive and induced plant resistance against herbivory. *Proceedings of the National Academy of Sciences of the United States of America*, **108**, 5685–5689.

Kruger AC, Shongwe S (2004) Temperature trends in South Africa: 1960--2003. *International Journal of Climatology*, **24**, 1929–1945.

Langfelder P, Horvath S (2008) WGCNA: an R package for weighted correlation network analysis. *BMC bioinformatics*, **9**, 559.

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nature*
methods, **9**, 357–359.

Lasky JR, Des Marais DL, Lowry DB *et al.* (2014) Natural Variation in Abiotic Stress
Responsive Gene Expression and Local Adaptation to Climate in Arabidopsis thaliana.
Molecular biology and evolution, **31**, 2283–2296.

Latimer A, Silander JA Jr, Rebelo AG, Midgley GF (2009) Experimental biogeography:
the role of environmental gradients in high geographic diversity in Cape Proteaceae.
Oecologia, **160**, 151–162.

Lázaro-Nogal A, Matesanz S, Godoy A *et al.* (2015) Environmental heterogeneity leads
to higher plasticity in dry-edge populations of a semi-arid Chilean shrub: insights into
climate change responses. *The Journal of ecology*, **103**, 338–350.

Ledón-Rettig CC, Pfennig DW, Chunco AJ, Dworkin I (2014) Cryptic genetic variation in
natural populations: a predictive framework. *Integrative and comparative biology*, **54**,
783–793.

Loehle C (1998) Height growth rate tradeoffs determine northern and southern range
limits for trees. *Journal of biogeography*, **25**, 735–742.

Maia R, Eliason CM, Bitton P-P, Doucet SM, Shawkey MD (2013) pavo : an R package
for the analysis, visualization and organization of spectral data (A Tatem, Ed,).
Methods in ecology and evolution / British Ecological Society, **I**.

Mason SJ, Jury MR (1997) Climatic variability and change over southern Africa: a
reflection on underlying processes. *Progress in Physical Geography: Earth and*
Environment, **21**, 23–50.

Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R (2010) Reactive oxygen species
homeostasis and signalling during drought and salinity stresses. *Plant, cell &*
environment, **33**, 453–467.

Mitchell N, Carlson JE, Holsinger KE (2018) Correlated evolution between climate and
suites of traits along a fast-slow continuum in the radiation of Protea. *Ecology and*
evolution, **8**, 1853–1866.

Moles AT, Perkins SE, Laffan SW *et al.* (2014) Which is a better predictor of plant traits:
temperature or precipitation? (A Helm, Ed,). *Journal of vegetation science: official*
organ of the International Association for Vegetation Science, **25**, 1167–1180.

Morris MRJ, Richard R, Leder EH *et al.* (2014) Gene expression plasticity evolves in
response to colonization of freshwater lakes in threespine stickleback. *Molecular*
ecology, **23**, 3226–3240.

Muñoz NJ, Farrell AP, Heath JW, Neff BD (2014) Adaptive potential of a Pacific salmon
challenged by climate change. *Nature climate change*, **5**, 163.

Nicotra AB, Atkin OK, Bonser SP *et al.* (2010) Plant phenotypic plasticity in a changing
climate. *Trends in plant science*, **15**, 684–692.

Nicotra AB, Segal DL, Hoyle GL *et al.* (2015) Adaptive plasticity and epigenetic variation
in response to warming in an Alpine plant. *Ecology and evolution*, **5**, 634–647.

O'Brien MJ, Leuzinger S, Philipson CD, Tay J, Hector A (2014) Drought survival of
tropical tree seedlings enhanced by non-structural carbohydrate levels. *Nature climate*
change, **4**, 710.

Oliveros JC (2016) Venny. An interactive tool for comparing lists with Venn's diagrams.
2007--2015.

Olszyk D, Wise C, VanEss E, Tingey D (1998) Elevated temperature but not elevated
CO2 affects long-term patterns of stem diameter and height of Douglas-fir seedlings.
Canadian journal of forest research. Journal canadien de la recherche forestiere, **28**,
1046–1054.

Pachter L (2011) Models for transcript quantification from RNA-Seq. *arXiv [q-bio.GN]*.

Palacio-López K, Beckage B, Scheiner S, Molofsky J (2015) The ubiquity of phenotypic
plasticity in plants: a synthesis. *Ecology and evolution*, **5**, 3389–3400.

Pearce DW, Millard S, Bray DF, Rood SB (2006) Stomatal characteristics of riparian
poplar species in a semi-arid environment. *Tree physiology*, **26**, 211–218.

Peppe DJ, Royer DL, Cariglino B *et al.* (2011) Sensitivity of leaf size and shape to
climate: global patterns and paleoclimatic applications. *The New phytologist*, **190**,
724–739.

Pratt JD, Mooney KA (2013) Clinal adaptation and adaptive plasticity in Artemisia
californica: implications for the response of a foundation species to predicted climate
change. *Global change biology*, **19**, 2454–2466.

Prunier R, Holsinger KE, Carlson JE (2012) The effect of historical legacy on adaptation:
do closely related species respond to the environment in the same way? *Journal of*
evolutionary biology, **25**, 1636–1649.

Ramachandra Reddy A, Chaitanya KV, Vivekanandan M (2004) Drought-induced
responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of*
plant physiology, **161**, 1189–1202.

Rebelo A (2001) Saso proteas: a field guide to the proteas of Southern Africa.

Sauquet H, Weston PH, Anderson CL *et al.* (2009) Contrasted patterns of

hyperdiversification in Mediterranean hotspots. *Proceedings of the National Academy*

of Sciences, **106**, 221–225.

Scheiner SM (2013) The genetics of phenotypic plasticity. XII. Temporal and spatial

heterogeneity. *Ecology and evolution*, **3**, 4596–4609.

Schlichting CD, Pigliucci M (1995) Gene regulation, quantitative genetics and the

evolution of reaction norms. *Evolutionary ecology*, **9**, 154–168.

Schneider RF, Li Y, Meyer A, Gunter HM (2014) Regulatory gene networks that shape

the development of adaptive phenotypic plasticity in a cichlid fish. *Molecular ecology*,

23, 4511–4526.

Schulze RE (1997) *South African atlas of agrohydrology and-climatology*. Water

Research Commission, Pretoria, South Africa.

Schulze RE (2007) *South African atlas of climatology and agrohydrology: WRC Report*

1489/1/06. Water Resource Commission.

Sgrò CM, Terblanche JS, Hoffmann AA (2016) What Can Plasticity Contribute to Insect

Responses to Climate Change? *Annual review of*.

Skelton RP, Midgley JJ, Nyaga JM, Johnson SD, Cramer MD (2012) Is leaf pubescence

of Cape Proteaceae a xeromorphic or radiation-protective trait? *Australian journal of*

botany, **60**, 104–113.

Stenström A, Jónsdóttir IS, Augner M (2002) Genetic and environmental effects on

morphology in clonal sedges in the Eurasian Arctic. *American journal of botany*, **89**,

1410–1421.

Steyn WJ, Wand SJE, Jacobs G, Rosecrance RC, Roberts SC (2009) Evidence for a

photoprotective function of low-temperature-induced anthocyanin accumulation in

apple and pear peel. *Physiologia plantarum*, **136**, 461–472.

Stoks R, Govaert L, Pauwels K, Jansen B, De Meester L (2016) Resurrecting

complexity: the interplay of plasticity and rapid evolution in the multiple trait response

to strong changes in predation pressure in the water flea Daphnia magna. *Ecology*

letters, **19**, 180–190.

Sultan SE (2003) Phenotypic plasticity in plants: a case study in ecological development.

Evolution & development, **5**, 25–33.

Team RC (2014) R: A language and environment for statistical computing. Vienna,

Austria: R Foundation for Statistical Computing; 2014.

Valladares F, Matesanz S, Guilhaumon F *et al.* (2014) The effects of phenotypic
plasticity and local adaptation on forecasts of species range shifts under climate
change. *Ecology letters*, **17**, 1351–1364.

Van Dyken JD, Wade MJ (2010) The genetic signature of conditional expression.
Genetics, **184**, 557–570.

Van Kleunen M, Fischer M (2005) Constraints on the evolution of adaptive phenotypic
plasticity in plants. *The New phytologist*, **166**, 49–60.

Van Wilgen BW, Forsyth GG, De Klerk H (2010) Fire management in Mediterranean-
climate shrublands: a case study from the Cape fynbos, South Africa. *Journal of*
Applied.

Vitasse Y, Hoch G, Randin CF *et al.* (2013) Elevational adaptation and plasticity in
seedling phenology of temperate deciduous tree species. *Oecologia*, **171**, 663–678.

Wadgymar SM, Mactavish RM, Anderson JT (2018) Transgenerational and Within-
Generation Plasticity in Response to Climate Change: Insights from a Manipulative
Field Experiment across an Elevational Gradient. *The American naturalist*, **192**, 698–
714.

Wang H, Fan W, Li H *et al.* (2013) Functional characterization of dihydroflavonol-4-
reductase in anthocyanin biosynthesis of purple sweet potato underlies the direct
evidence of anthocyanins function against abiotic stresses. *PloS one*, **8**, e78484.

Wennersten L, Forsman A (2012) Population-level consequences of polymorphism,
plasticity and randomized phenotype switching: a review of predictions. *Biological*
reviews of the Cambridge Philosophical Society, **87**, 756–767.

West AG, Dawson TE, February EC (2012) Diverse functional responses to drought in a
Mediterranean-type shrubland in South Africa. *New*.

West-Eberhard MJ (2003) *Developmental Plasticity and Evolution*. Oxford University
Press.

Whitehead A (2012) Comparative genomics in ecological physiology: toward a more
nuanced understanding of acclimation and adaptation. *The Journal of experimental*
biology, **215**, 884–891.

Wilson AM, Latimer AM, Silander JA (2015) Climatic controls on ecosystem resilience:
Postfire regeneration in the Cape Floristic Region of South Africa. *Proceedings of the*
National Academy of Sciences, **112**, 9058–9063.

Young DJN, Stevens JT, Earles JM *et al.* (2017) Long-term climate and competition
explain forest mortality patterns under extreme drought. *Ecology letters*, **20**, 78–86.

Yu H, Chen X, Hong Y-Y *et al.* (2008) Activated expression of an Arabidopsis HD-

START protein confers drought tolerance with improved root system and reduced

stomatal density. *The Plant cell*, **20**, 1134–1151.

Zeng X-Q, Chow WS, Su L-J, Peng X-X, Peng C-L (2010) Protective effect of

supplemental anthocyanins on Arabidopsis leaves under high light. *Physiologia*

plantarum, **138**, 215–225.

Züst T, Rasmann S, Agrawal AA (2015) Growth--defense tradeoffs for two major anti-

herbivore traits of the common milkweed Asclepias syriaca. *Oikos* , **124**, 1404–1415.

Table 1. Results of linear mixed models testing functional trait and performance differences of *Protea repens* seedlings at 6 and 12 days into the dry down experiment. The model compares values between watered and unwatered treatments as well as among 8 wild-sourced populations and the population by treatment interaction. Significant values are bolded.

Response variable	Fixed effects	Fixed effects at 6 days				Fixed effects at 12 days			
		df	Chi-sq	P-value	q-val	df	Chi-sq	P-value	
Leaf area (cm^2)	Population	7	49.91	**<0.0001**	**<0.0001**	6	69.16	**<0.0001**	**<0.0001**
	Treatment	1	0.17	0.678	0.881	1	23.39	**<0.0001**	**<0.0001**
	Pop. × Trt.	7	8.61	0.282	0.423	6	13.3	0.039	0.050
SLA (cm^2/g) Log transform	Population	7	5.15	0.642	0.881	**6**	**15.35**	**0.018**	**0.027**
	Treatment	**1**	**12.13**	**0.0005**	**0.002**	1	73.14	**<0.0001**	**<0.0001**
	Pop. × Trt.	**7**	**19.23**	**0.008**	**0.027**	**6**	**16.71**	**0.010**	**0.017**
LWR	Population	7	67.48	**<0.0001**	**<0.0001**	6	78.55	**<0.0001**	**<0.0001**
	Treatment	1	0.06	0.814	0.881	1	1.86	0.173	0.210
	Pop. × Trt.	7	3.47	0.839	0.881	6	1.52	0.958	0.958
Stomatal density	Population	**7**	**17.46**	**0.015**	**0.040**	6	29.73	**<0.0001**	**0.0001**
	Treatment	1	3.07	0.080	0.192	**1**	**12.95**	**0.0003**	**0.0007**
	Pop. × Trt.	7	9.56	0.215	0.369	6	3.59	0.732	0.765
Stomatal pore size (µm)	Population	7	3.87	0.795	0.881	**6**	**21.92**	**0.001**	**0.002**
	Treatment	1	0.04	0.844	0.881	**1**	**27.68**	**<0.0001**	**<0.0001**
	Pop. × Trt.	7	3.05	0.881	0.881	**6**	**14.27**	**0.027**	**0.038**
Stem pigmentation	Population	7	34.13	**<0.0001**	**<0.0001**	7	38.51	**<0.0001**	**<0.0001**
	Treatment	1	1.85	0.174	0.336	1	0.42	0.515	0.579
	Pop. × Trt.	7	10.12	0.182	0.336	7	5.71	0.574	0.630
Stem height (cm)	Population	**7**	**105.34**	**<0.0001**	**<0.0001**	7	121.24	**<0.0001**	**<0.0001**
	Treatment	**1**	**5.9**	**0.015**	**0.040**	1	49.4	**<0.0001**	**<0.0001**
	Pop. × Trt.	7	4.1	0.768	0.881	7	6.86	0.443	0.511
Stomatal conductance	Population	7	0	0.254	0.406	**6**	**17.52**	**0.008**	**0.014**
	Treatment	**1**	**0**	**<0.0001**	**<0.0001**	**1**	**104.7**	**<0.0001**	**<0.0001**
	Pop. × Trt.	7	0	0.148	0.323	**6**	**23.43**	**0.0007**	**0.001**

Table 2. Results of linear mixed models comparing eigengene values (E0-E13) at 6 and 12 days into the dry down experiment for watered and unwatered plants of *P. repens*. Significant values are bolded.

	Fixed effects	Fixed effects at 6 days				Fixed effects at 12 days			
		df	Chi-sq	p-value	q-val	ddf	Chi-sq	p-value	q-val
E0	Population	**7**	**24.89**	**0.0008**	**0.0042**	**6**	**80.38**	**<0.0001**	**<0.0001**
	Treatment	1	2.05	0.152	0.281	1	0.77	0.380	0.479
	Pop. × Trt	7	5.72	0.573	0.655	6	8.51	0.203	0.294
E1	Population	**7**	**18.59**	**0.010**	**0.042**	**6**	**16.49**	**0.011**	**0.026**
	Treatment	**1**	**91.40**	**<0.0001**	**<0.0001**	**1**	**123.77**	**<0.0001**	**<0.0001**
	Pop. × Trt	7	11.30	0.126	0.252	6	6.71	0.348	0.457
E2	Population	7	8.91	0.259	0.363	6	13.87	0.031	0.0566
	Treatment	**1**	**13.01**	**0.0003**	**0.0018**	**1**	**50.11**	**<0.0001**	**<0.0001**
	Pop. × Trt	7	9.65	0.210	0.351	6	3.22	0.781	0.841
E3	Population	7	8.56	0.286	0.387	**6**	**18.70**	**0.005**	**0.013**
	Treatment	1	0.07	0.796	0.836	**1**	**9.88**	**0.00167**	**0.005**
	Pop. × Trt	7	9.30	0.232	0.351	6	1.47	0.962	0.962
E4	Population	7	16.16	0.024	0.078	**6**	**36.29**	**<0.0001**	**<0.0001**
	Treatment	**1**	**7.52**	**0.006**	**0.028**	**1**	**75.50**	**<0.0001**	**<0.0001**
	Pop. × Trt	7	6.04	0.536	0.643	6	3.80	0.704	0.799
E5	Population	7	12.91	0.074	0.177	**6**	**26.58**	**0.0002**	**0.0007**
	Treatment	**1**	**13.37**	**0.0003**	**0.0018**	**1**	**68.66**	**<0.0001**	**<0.0001**
	Pop. × Trt	7	9.32	0.231	0.351	6	2.90	0.821	0.841
E6	Population	7	12.84	0.076	0.177	6	11.76	0.067	0.113
	Treatment	**1**	**56.14**	**<0.0001**	**<0.0001**	**1**	**107.13**	**<0.0001**	**<0.0001**
	Pop. × Trt	7	9.00	0.253	0.363	6	7.72	0.260	0.352
E7	Population	7	15.34	0.032	0.096	**6**	**19.99**	**0.003**	**0.008**
	Treatment	**1**	**67.46**	**<0.0001**	**<0.0001**	**1**	**86.20**	**<0.0001**	**<0.0001**
	Pop. × Trt	7	10.67	0.154	0.281	**6**	**14.79**	**0.022**	**0.049**
E8	Population	7	15.11	0.035	0.098	6	3.69	0.719	0.799
	Treatment	1	0.65	0.420	0.540	1	1.49	0.222	0.311
	Pop. × Trt	7	9.96	0.191	0.334	6	4.10	0.663	0.796
E9	Population	7	12.56	0.084	0.186	**6**	**23.63**	**0.0006**	**0.0019**
	Treatment	**1**	**26.83**	**<0.0001**	**<0.0001**	**1**	**102.74**	**<0.0001**	**<0.0001**
	Pop. × Trt	7	6.21	0.515	0.636	6	13.97	0.030	0.057
E10	Population	7	1.76	0.972	0.972	6	10.58	0.102	0.165
	Treatment	1	0.04	0.840	0.860	1	0.13	0.723	0.799
	Pop. × Trt	7	5.46	0.604	0.655	6	14.12	0.028	0.056
E11	Population	7	13.34	0.064	0.168	6	12.34	0.055	0.096
	Treatment	**1**	**31.91**	**<0.0001**	**<0.0001**	**1**	**64.04**	**<0.0001**	**<0.0001**
	Pop. × Trt	7	9.27	0.234	0.351	6	6.33	0.388	0.479
E12	Population	7	7.05	0.424	0.540	6	9.38	0.154	0.231
	Treatment	1	2.75	0.097	0.204	1	0.06	0.808	0.841
	Pop. × Trt	7	5.53	0.596	0.655	6	9.41	0.152	0.231
E13	Population	7	16.84	0.018	0.063	6	14.14	0.028	0.056
	Treatment	1	0.26	0.608	0.655	**1**	**30.16**	**<0.0001**	**<0.0001**
	Pop. × Trt	**7**	**17.85**	**0.013**	**0.0496**	6	16.57	0.011	0.026

Table 3. Results of linear mixed models testing for differences in growth and carbohydrate storage between watered and unwatered *Protea repens* seedlings and among 8 populations at the end of the dry-down experiment (day 12). . Significant values are bolded.

Response variable	Fixed effects	df	Chi-sq	p-val	q-val
Soluble sugars mass (g) belowground	Population	7	55.57	**<0.0001**	**<0.0001**
	Treatment	1	6.33	**0.012**	**0.019**
	Pop. × Trt.	7	17.52	**0.014**	**0.022**
Soluble sugars mass (g) aboveground	Population	7	80.08	**<0.0001**	**<0.0001**
	Treatment	1	25.69	**<0.0001**	**<0.0001**
	Pop. × Trt.	7	24.27	**0.001**	**0.002**
Starch mass (g) belowground	Population	7	21.06	**0.004**	**0.008**
	Treatment	1	0.10	0.748	0.765
	Pop. × Trt.	7	4.57	0.712	0.763
Starch mass (g) aboveground	Population	7	57.40	**<0.0001**	**<0.0001**
	Treatment	1	60.10	**<0.0001**	**<0.0001**
	Pop. × Trt.	7	100.76	**<0.0001**	**<0.0001**
Growth rate (height on Day 12-6)	Population	7	34.85	**<0.0001**	**<0.0001**
	Treatment	1	194.34	**<0.0001**	**<0.0001**
	Pop. × Trt.	7	26.33	**0.0004**	**0.0009**
Root length (mm)	Population	6	9.36	0.154	0.192
	Treatment	1	6.66	**0.010**	**0.017**
	Pop. × Trt.	6	6.3	0.391	0.463
Stem pigment accumulation	Population	7	15.47	**0.031**	**0.041**
	Treatment	1	5.08	**0.024**	**0.035**
	Pop. × Trt.	7	15.58	**0.029**	**0.040**

Table 4. Significant effects from mixed models testing associations between climate variables and plasticity measures, for traits and eigengenes on day 6 and day 12.

Day	Plasticity measure (RDPI)	Climate variable	P-val.	Slope	Plasticity high in pops. from:
Day 6	GN5	Elevation	0.015	0.0005	lower elevation
Day 12	GN10	Mean annual precip.	0.035	0.0014	wettest and driest sites
Day 12	GN13	Mean annual precip.	0.213	-0.0014	wetter sites
Day 12	Growth	Elevation	0.014	-0.0005	lower elevation
Day 12	Growth	Summer max. temp.	0.016	0.0709	warmer sites
Day 12	Leaf area	Mean annual precip.	0.019	0.0006	wetter sites
Day 12	Conductance	Temperature range	0.007	0.1235	higher heterogeneity
Day 12	Pigment content	Temperature range	0.039	0.0474	higher heterogeneity
Day 12	Pigment content	Summer max. temp.	0.041	0.0477	coolest and warmest sites

FIGURES

Fig 1. Map of CFR and locations of sampling sites (A); Principal component analysis for the climatic variables used in this study

Fig 2. Multi-dimensional scaling (MDS) plots for gene expression patterns at day 6 (A) and day 12 (B); principal component analysis of measured traits at day 6 (C) and day 12 (D) into the drought treatment.

Fig 3. Number of differentially expressed genes for the 8 populations for after 6 and 12 days of the drought treatment (A). Note that population A is not sampled and does not have any DE genes at day 12. Number of differentially expressed genes among populations under control and drought conditions for day 6 and 12 separately (B).

Fig 4. Heatmap of standardized regression coefficients for climate variables as predictors and traits and gene networks (GN) as response variables for measurements (A) on day 6 or (B) day 12. For variables with significant interaction effects, each treatment was analyzed separately to estimate slopes and test their significance within treatments. Solid boxes indicate a significant assosication and the box is divided in cases where there is a significant treatment and climate effect, indicating that the climate association was significant only in one of the treatments. The upper half of the box represents the watered treatment and the lower half represents the dry treatment. Stars indicate significant correlations for either of the two treatments.

Fig 5. Response curves for the studied 8 populations under well-watered and drought conditions and the associations between plasticity and climate variables for growth (A), leaf area (B), stomatal conductance (C) and gene network 13 (D).

Fig 6. Mean annual precipitation (A) and mean annual temperature (B) averaged across the 8 population sites between 1901-2016.

Supplementary info

Table S1. Population source site climatic information

Table S2. Correlations between climatic variables

Table S3. WGCNA GO categories and highest gene network membership genes

Table S4. Up and down-regulated genes for all the populations under drought

Table S5. Gene network and trait correlations

Figure S1. Sampling scheme

Figure S2. WGCNA sample clustering

Figure S3. Functional traits response curves

Figure S4. Gene network response curves

Figure S5. Performance traits response curves

Figure S6. MDS plot with all samples for day 6 and day 12

Figure S7. Contigs that show a population x treatment effect, from figure 3