**Running title: Climate explains population divergence in plasticity**

**Climate explains population divergence in drought-induced plasticity of**

**functional traits and gene expression in a South African *Protea***

Melis Akman[1,2], Jane E. Carlson[3,4], Andrew M. Latimer[1]

[1] Department of Plant Sciences, UC Davis

[2] Department of Plant and Microbial Biology, UC Berkeley

[3] Department of Biology, Nicholls State University

[4] Gulf Coast Network Inventory and Monitoring Program, National Park Services

Corresponding author: Melis Akman

**Abstract**

Long term environmental variation often drives local adaptation and leads to trait differentiation across populations. Additionally, when traits change in an environment-dependent way through phenotypic plasticity, the genetic variation underlying plasticity will also be under selection. These processes could create a landscape of differentiation across populations in traits and their plasticity. Here, we studied drought responses in seedlings of a shrub species from the Cape Floristic Region, the common sugarbush (*Protea repens*). We measured morphological and physiological traits, and sequenced whole transcriptomes in 8 populations that represent both the climatic and the geographic distribution of this species. We found that there is substantial variation in how populations respond to drought, but we also observed common patterns such as reduced leaf size and leaf thickness, and upregulation of stress- and down-regulation of growth-related gene groups. Both high environmental heterogeneity and milder source site climates were associated with higher plasticity in various traits and co-expression gene networks. Associations between traits, trait plasticity, gene networks and the source site climate suggests that temperature may play a bigger role in shaping these patterns when compared to precipitation, in line with recent changes in the region due to climate change. We also found that traits respond to climatic variation in an environment dependent manner: some associations between traits and climate were apparent only under certain growing conditions. Together, our results uncover common responses of *P. repens* populations to drought, and climatic drivers of population differentiation in functional traits, gene expression and their plasticity.

 Keywords: phenotypic plasticity, climatic variation, transcriptomics, *Protea*, drought, climate change

## INTRODUCTION

Populations can persist in changing environmental conditions through three mechanisms: by tracking their climatic niche in space through migration or dispersal, by shifting their niche through adaptation, and/or by matching traits to the new environment through phenotypic plasticity (Chevin, Lande, and Mace 2010). For plants, plasticity can be the fastest response to environmental change, since it involves rapid adjustments of morphology, physiology and gene expression within a single generation. Like mean trait values, the form and magnitude of trait plasticity can be under selection and thus plasticity can also vary among populations along environmental gradients (Geber and Griffen 2003; Wennersten and Forsman 2012; Lázaro-Nogal et al. 2016, 2015; Wadgymar, Mactavish, and Anderson 2018). Previous studies have demonstrated that such population differences in plasticity might or might not have climatic drivers (Pratt and Mooney 2013; Lázaro-Nogal et al. 2015; Pohlman, Nicotra, and Murray 2005; Cochrane et al. 2015).

While plasticity can have strong consequences on functional traits (e.g. (Etterson 2004), it remains less clear how often plasticity confers environment-specific fitness benefits (Forsman 2015; Acasuso-Rivero et al. 2019). Plasticity has been directly shown to be adaptive: for example, Dudley and Schmitt (1996) showed that plastic stem elongation was beneficial at high but not low plant densities. Other studies have characterized patterns of plasticity that suggest adaptive plasticity (Sultan 2003; Nicotra et al. 2015; Baythavong 2011; Lázaro-Nogal et al. 2016; Cavender-Bares and Ramírez-Valiente 2017). Resurrection studies using historical and modern *Daphnia magna* populations showed evolution of plasticity for heat tolerance and fish predation

(Cuenca Cambronero et al. 2018; Stoks et al. 2016). In light of these studies and theoretical predictions (Grime 1994; Bradshaw and Hardwick 1989; Alpert and Simms 2002; Schlichting and Pigliucci 1995; West-Eberhard 2003; Ghalambor et al. 2015), we expect that selection should promote local adaptation in plasticity under two main circumstances. First, plants from environments with strong and relatively predictable forms of spatial or temporal heterogeneity should be more plastic than those from homogeneous environments (Ackerly et al. 2000; Van Kleunen and Fischer 2005; Baythavong and Stanton 2010; Scheiner 2013; Lázaro-Nogal et al. 2015; Bonamour et al. 2019; Chevin and Hoffmann 2017; Reed et al. 2010); but see Palacio-López et al. (2015). Second, plants from harsh climates are expected to be less plastic than those from milder climates (Vitasse et al. 2013; Gugger et al. 2015), because the fitness costs of mismatch between the phenotype and the environment should be greater in resource-poor conditions (Alpert and Simms 2002). Elevation contrasts provide one example of this latter pattern: for an alpine herb, *Wahlenbergia ceracea,* inhabiting high montane and alpine elevations, plants from colder, higher-elevation environments tend to have both lower and less plastic growth rates than plants from warmer sites (Nicotra et al. 2015).

The form and magnitude of functional plasticity which organisms express can be expected to influence the vulnerability of populations to environmental change. This study focuses on a widespread plant species from the Cape Floristic Region (CFR) of South Africa, a biodiversity hotspot (Cowling 2009), that is predicted to be strongly affected by climate change. Our study species, *Protea repens* (common sugar bush), is the most widespread species in this plant genus with 112 species, ~85% of which are endemic (Sauquet et al. 2009; Rebelo 2001). Climate models project warming up

to 3 °C by 2070 (Wilson, Latimer, and Silander 2015), and precipitation projections being more variable, most models project drying in the super-diverse western mountains (Wilson, Latimer, and Silander 2015; Hewitson and Crane 2006). Most Mediterranean-climate regions, including California and Southern Australia as well as the CFR, have recently experienced unprecedented droughts, which caused large mortality events in trees (Young et al. 2017; Allen et al. 2010) and shrubs (Valliere et al. 2017). Thus, understanding the patterns and effects of drought plasticity in functional traits is a vital issue for plant vulnerability in these semi-arid regions.

Plasticity in key plant traits, including stomatal density and conductance, growth rate, and leaf morphology, has been observed to vary among populations in long-lived woody plants (e.g. (Aubin et al. 2016) as well as herbaceous annuals (Jacobs and Latimer 2012). Likewise, along the environmental gradients *Protea* species inhabit, we have found evidence for local adaptation and environment-dependent benefits in several traits *(Carlson, Adams, and Holsinger 2016; Carlson, Holsinger, and Prunier 2011; Latimer et al. 2009; Mitchell, Carlson, and Holsinger 2018; Prunier, Holsinger, and Carlson 2012)*. Specifically, we found evidence for positive selection on stomatal density in populations from hotter and drier sites, with links to higher stomatal conductance in *P. repens (Carlson, Adams, and Holsinger 2016)*. Additionally, in the monophyletic group of white proteas, *P. subvestita* shows correlations between root length and rainfall seasonality. Biomass also decreases with increasing elevation in *P. mundii*, *P. punctata* and *P. lacticolor*, but shows the opposite trend in *P. mundii* (Prunier, Holsinger, and Carlson 2012). On a broader phylogenetic scale, we also found associations between temperature and plant size related traits such as leaf area and plant height (Mitchell, Carlson, and Holsinger 2018). There is also evidence for

differential developmental plasticity and drought plasticity in leaf traits of several other

*Protea* species (*Protea sect. Exsertae*: (Carlson and Holsinger 2012; Heschel et al.

2014).

Previously, we also showed that gene expression in populations grown in a

common garden show patterns of differentiation corresponding to climatic variables

such as elevation (a good proxy for temperature in CFR), minimum winter temperature

and mean annual precipitation in *P. repens* (Akman et al. 2016). These gene networks

also show correlations with key traits such as height, leaf area, stomatal density/size,

providing insights into the genes involved in climate driven differentiation.

Measurements of plasticity in gene expression, in which thousands of "phenotypes"

can be measured simultaneously, are also valuable for investigating the causes and

consequences of plasticity. Gene expression patterns can change within seconds of

a stimulus, long before larger biochemical shifts and morphological changes become

apparent. Gene expression patterns can also be linked to morphological and

physiological adjustments, potentially providing insights into the functional genetic

basis of adaptation and/or plasticity (Whitehead 2012; Akman et al. 2016) We note

that such links do not pinpoint the genetic locus associated with the phenotypic

variation, as loci influencing expression may lie in regulatory regions or in others genes

whose expression is upstream of the focal gene. Previous studies have revealed

patterns of adaptive plasticity in gene expression in fish (Dayan, Crawford, and

Oleksiak 2015; Morris et al. 2014; Schneider et al. 2014), plants (Lasky et al. 2014;

Colicchio et al. 2015) and corals (Barshis et al. 2013; Granados-Cifuentes et al. 2013).

These studies demonstrate that gene expression profiling can usefully complement

standard functional trait measurements to help understand the functional genetic basis and physiological significance of plasticity and how it varies across populations.

With this study, we aim to answer two questions: 1) How do populations of *P. repens* respond to drought, both in terms of functional traits and gene expression, together with their plasticity, and how do these responses vary across populations? and 2) Do the climatic conditions of the source sites covary with the mean values and plasticity of functional traits and gene expression levels? To do so, we imposed a severe, experimental drought on seedlings in a dry down experiment, and measured functional traits and gene expression patterns. Then, we tested for population differentiation in trait means and drought-response plasticity in functional traits and gene expression patterns, and evaluated associations between patterns of population differentiation and their potential climatic drivers.

## MATERIALS AND METHODS

**Species description**

*Protea repens* L. is a common evergreen shrub endemic to South Africa, with particularly high abundance in semi-arid fynbos of the CFR (Rebelo 2001). This species has narrow, sclerophyllous leaves and reaches 2-4 m in height. All viable and nonviable fruits remain within a woody, serotinous infructescence until the adult plant is killed by fire, which occurs at ~10-20 year intervals (with variation among geographic regions (Van Wilgen, Forsyth, and De Klerk 2010). Fruits disperse by tumbling, and seeds germinate during the next rainy season (May-Aug for much of the CFR). Due to the Mediterranean rainfall patterns in the western CFR, some western sites receive far more wintertime precipitation than do eastern population sites.

**Field sampling and climate data**

In 2011, we collected seeds from 8 wild populations of *P. repens* spanning its geographic distribution in the CFR (**Figure 1A** and **Table S1**). The populations encompass a range of precipitation levels and degrees of seasonal concentration -- some sites have strong winter rainfall seasonality, while others experience both winter precipitation and some summer monsoonal rain (Table S1). The elevation range was deliberately wide, from ~150m to ~1500m. The reason why the lower elevation, warmer sites are "milder" is that they have little or no frost in winter, while being only moderately dry. High-elevation sites, in contrast, receive snow in winter and have high winds and shorter growing seasons. There are hot, dry, low-elevation sites in the interior of the Cape region that are not mild, but these tend to be too dry to support our study species, and were not included in our sampling. From 40-50 plants per population (mean n= 45 plants), we collected 1-2 mature seed heads from the previous year's growth. Plants were sampled along transects through the population centre, with at least 1 m between each sampled plant for small populations (<100 plants), and 5 m for larger populations. Seed heads were allowed to air-dry until achenes were released. GPS coordinates of each site near the population center were used to extract elevation and six climate metrics for rainfall and temperature from GIS layers. A principal component analysis performed on these data confirmed that the sample sites cover most of the range of variation evenly in major environmental factors (**Figure 1B**).

The climate data layers were from the South African Atlas of Agrohydrology and Climatology (Roland Edgar Schulze 1997) with higher resolution for 4 and 5 below, and (R. E. Schulze 2007) for the rest, and were based on 50 year (collected between

1950-1999 when data was available from field stations) climate averages, extremes,

or variances. The climate layers used were (1) mean annual rainfall, (2) average daily

maximum temperature for summer (January), (3) average daily minimum temperature

for winter (July), (4) precipitation concentration (an seasonality indicator we refer to as

"rainfall seasonality"), measured as Markham's concentration index (Markham 1970),

which ranges from 0 to 100, with a value of 0 indicating perfectly even precipitation

across all months, and 100 indicating that all precipitation falls in a single month (5)

inter-annual rainfall variability, measured as the coefficient of variation in mean annual

rainfall across years, and (6) extreme temperature range within a year, measured as

the difference between summer maximum and winter minimum temperatures. Our

seventh variable was elevation, which was extracted separately from ASTER GDEM

(Daac 2011). Elevation is correlated strongly with winter minimum temperature, and

we interpret it primarily as an indicator of temperature regime, although some other

factors such as precipitation and wind speed are more weakly associated with

elevation in the region (**Table S2**). Most of these layers have been used in our earlier

work on *P. repens*, except for (5) and (6), which were added to complement rainfall

seasonality as additional measures of environmental heterogeneity. Soil analyses for

these source population locations were not available, but we standardized the general

soil type: populations occurred (as is typical for this species) on low-nutrient, coarse,

sandy soil derived primarily from quartzite sandstone parent material.

Only two pairs of climate variables were strongly correlated with each other:

winter minimum temperature with elevation (Pearson's R= -0.91) and summer

maximum temperature with annual temperature range (R=0.70; **Table S2**). Rainfall

seasonality in our dataset is positively associated with wintertime rainfall (total for Dec-

Feb, Pearson's R=0.80) and negatively associated with summertime rainfall (total for June-August, Pearson's R=-0.90). The 12-year gap between the last collection year of the climate variables and seed collection date might hinder some of the associations we detect in later sections. However, the age of the plants (9-23 years based on veld age through burn history when available) from which seeds were collected will potentially eliminate this gap. We predict that selection should have already acted on the maternal plants and pollen sources, and their adaptations, if any, should be reflected in the germplasm we performed our experiments on.

**Experimental setup and dry-down treatments**

In July 2014, we sowed seeds from 8 *Protea repens* populations on low nutrient mix (50% bark, 50% sand) in trays of 100 individual plugs 2x2 cm in size with a complete random design. Trays were placed in a long-day growth chamber (8 hours in dark at 8℃, 16 hours in light at 20℃). The trays were randomized every 3 days with each watering in the growth chambers. After 1.5 months, we transplanted germinated seedlings into small, deep pots (5 cm diameter x 17 cm deep) in trays of 50 each, in the same low nutrient mix. After transplanting, seedlings were rested for two weeks indoors under growth lights before being transferred to a single greenhouse bench at Nicholls State University Farm, Thibodaux, LA.

Two weeks post-greenhouse transfer and two weeks prior to the start of the dry-down experiment, we selected a total of 316 out of 419 seedlings (19-20 seedlings/population x 8 populations x 2 treatments) for uniform size for the experiment. We then reorganized the 2.5 month-old seedlings into trays containing 20-25 seedlings each, in a complete random design. In the experiment, seedlings were

assigned to one of two treatments: control, in which we applied regular watering

(~every 3 days, equal amounts of water was given to each plant) and drought, in which

we stopped watering entirely for 12 days. Because we had a variable number of

seedlings per maternal plant, we could not perfectly balance maternal lines across

populations and treatments. To the extent possible, however, we included at least one

seedling per maternal family per treatment; for 82% of the 83 maternal families, we

were able to include at least one seedling per maternal family in each treatment, and

for 51% of families, we included multiple seedlings per treatment (Table S3). The

drought experiment began in late October 2014, when greenhouse conditions were

warm but relatively dry, and thus most similar to summer conditions in the native range.

**Trait measurements and gene expression sampling**

At 6 days and 12 days into the drought experiment, plants were measured for

functional traits and leaves were sampled for transcriptomics. Our sampling design is

depicted in **Figure S1**. Functional traits that were measured at both timepoints

included plant height (cm), stomatal conductance, leaf area, leaf length:width ratio

(LWR), specific leaf area (fresh leaf area/dry leaf mass; SLA), stomatal density

(number per mm$^2$), stomatal pore length (mm) and stem pigmentation. Stem

pigmentation, measured as stem hue, was derived using Endler's segment

classification (Maia *et al.* 2013), based on the average of two reflectance spectra

readings (400-700 nm) taken at mid-stem (JAZ, Ocean Optics). On the segment

classification scale for hue, higher values represent redder stems that contain higher

levels of anthocyanin. Stomatal conductance was measured using a steady-state leaf

porometer (SC-1, Decagon) on one leaf per plant, with an additional adjacent leaf

measured simultaneously if the chamber was not filled by a single leaf due to small

leaf size.

All other leaf measurements were taken from the 2nd or 3rd youngest

expanding leaf from the stem apex. Immediately following conductance measurement,

the leaf was removed to measure leaf area, width, length and stomatal traits. Fresh

leaves were digitally scanned shortly after harvest at 400 dpi on a flatbed scanner.

After scanning, acrylic peels were taken from the adaxial leaf surface and placed on

cellophane, which were later viewed under a light microscope at 40x for stomatal

measurements (see Carlson *et al.* 2016). Although height and pigmentation were

measured on all plants on both timepoints (mean sample size per population=39), all

other traits were measured on only a subset of plants on each of these days, typically

in association with the transcriptomic data (10-20 individuals per population for each

day). Two additional variables were calculated by subtracting day 6 from day 12 values

on the same plant: pigment accumulation (day 12 hue - day 6 hue) and growth (day

12 height - day 6 height). A principle component analysis was performed on the traits

measured separately for day 6 and day 12. Details of sampling for gene expression

and carbohydrate analysis are included in Supplementary Materials and Methods.

Briefly, leaves were sampled at both time points for whole transcriptome sequencing,

and at day 12, the whole plants were freeze dried and extracted with 70% methanol,

and carbohydrates were measured with anthrone method (as described in (Akman et

al. 2012)).

**Gene expression analyses and bioinformatics**

See Supplementary Materials and Methods for transcriptome library preparation and

sequencing details. Raw reads were trimmed and demultiplexed with trimmomatic

(Bolger, Lohse, and Usadel 2014) and fastx toolkit (Hannon, n.d.). By using bowtie

(Langmead and Salzberg 2012), trimmed reads were mapped to a previously

published de-novo transcriptome of *P. repens* (Akman et al. 2016a) assembled by

using sequences of individuals that also include all of our focal populations. Before

reads were mapped to the transcriptome, a further clustering of the contigs were done

using cd-hit (Fu et al. 2012). Read counts, extracted using eXpress (Pachter 2011),

averaged 6.02 million reads per sample (ranged between 2.46-21.80 million). Contigs

with <1 counts per million (cpm) were excluded from the analyses. Differential gene

expression analyses were performed using edgeR. For each time point, differentially

expressed genes per population were revealed using pairwise comparisons using

negative binomial models. The common set of differentially expressed genes was

obtained by extracting the overlap of differentially expressed genes for each

population at day 12 (**Table S5**). We then used linear models in edgeR, including

sequencing lane as a fixed factor, to detect contigs that showed variation between

treatments and among populations (population x treatment). Multi-dimensional scaling

(MDS) analysis was performed for each time point to visualize general patterns of

gene expression for all the samples using the top 500 genes that explain the most

variation (**Figure 2A-B and Figure S6**).

Co-expression gene networks were computed using WGCNA (Langfelder and

Horvath 2008) with consensus clustering option for the 2 timepoints together.

Normalized and cpm-filtered read counts were transformed with "voom" in limma

package. A soft-threshold of 6 was used for clustering and 14 co-expressed gene

networks were constructed. The number of genes within each gene network varied

between 49 (GN13) to 7682 (GN1). The gene network "grey" (called GN0 in this study)

as constructed by WGCNA for genes that do not fit any gene network has 3124 genes.

A representative "eigen gene" value, which is the first principal component axis for all

genes within the network, was used for further analyses of climate and trait

associations. By using gene orthology of our genes with Arabidopsis (Akman et al.

2016a), we also tested for Gene Ontology (GO) enrichment for the 13 networks. This

analysis was performed in Gorilla (Eden et al. 2009), using the network genes as target

set and the whole gene set as the background. Please note that, when we refer to a

statistical test for a given gene network (GN), an association or a linear mixed model

test, we are referring to the eigengene value of that gene network. However, when we

mention an associated function, this is based on all the genes on that network and the

corresponding GO analyses. Contigs within each module, GO enrichments and the

genes with the highest module membership for a single network are provided in **Table
S4**.

**Statistical analyses**

*How do populations respond to drought and how do these responses vary?*

We tested whether drought significantly affected traits in *Protea repens* using linear

mixed models. We used a total of 23 plant metrics in our analyses, including 8 traits

on day 6, and 15 traits on day 12. We analyzed each trait on each day separately,

because preliminary analyses on combining both days revealed significant timepoint

effects for almost every response variable. Each model included the fixed effects of

treatment, population, and the population by treatment interaction (G x E) and random

effects of maternal line nested within population. In these models, the treatment-by-

population interaction term indicates whether there was differential plasticity among

populations, or in other words, that plants from different populations responded

differently to the treatment (G x E effect). These analyses were performed using the

lme4 package (Bates, Maechler, and Bolker 2012) in R 3.1 (Team 2014). To obtain p-

values for each fixed effect, we successively dropped one predictor from the full model

and used a chi-square test to compare AIC scores between models. Given the large

number of statistical tests that this entailed, we also adjusted p-values according to

estimated false discovery rates (FDR; R package "fdrtool", local method).

Gene expression analyses followed the same general approach as for traits.

As response variables, these analyses focused on eigengenes in the 14 gene

networks (GN) that were produced through the gene clustering analysis. Models for

gene networks included fixed effects of treatment, population, and their interaction,

random effects of maternal line nested within population, and a random effect of

sequencing lane. We also tested for genes that are differentially expressed within a

population. Commonly regulated genes for all populations are given in **Table S5**. We

also tested for genes differentially expressed among populations for both timepoints

and days separately.

*Do the climatic conditions of the source sites covary with traits, gene expression and*

*their plasticity?*

To test whether characteristics of the home climate predicted population

variation in traits and gene expression, we used mixed effect models. In these models,

traits and gene networks (GN) were response variables, the climate variables were

fixed effect terms, and maternal line nested in population was a random effect variable.

For testing the significance of the fixed factors, we calculated p-values based on chi-

square tests as explained above. FDR corrections were applied to p-values in two

groups: traits and gene networks. Day 6 and Day 12 FDR corrections were also

applied separately. For variables with significant climate and treatment interaction,

each treatment was also analyzed separately to estimate slopes and test their

significance within treatments.

We used another set of linear mixed models to test whether there were

significant associations between traits and gene expression patterns. We performed

these analyses with lme4 in R as above, with each pairwise comparison between a

trait and a gene network in its own model. Response variables were one of 8 (at day

6) or 15 (at day 12) traits and predictor variables were one of 14 eigen gene values for

a gene network. Random effects were sequencing lane, population and maternal line

nested within population. Correlations were tested separately for the two timepoints,

and p-values were calculated as explained previously. An FDR correction was applied

separately to the two time points (Table S6).

To characterize plasticity for individual traits and to quantify the direction and

magnitude of plasticity, we used maternal line plasticity indices. For each maternal

family, we calculated plasticity as the family mean trait value in the wet treatment

minus the family mean trait value in the drought treatment, divided by the overall family

mean trait value (for families with only one plant per treatment, we used the trait value

of that plant rather than a family mean). Note that this plasticity metric can take on

both positive and negative values: values close to 0 indicate low plasticity (little

difference between treatments), while large negative and large positive values both

indicate high plasticity.

We used mixed effect models to determine whether characteristics of the home

climate predicted cross-population variation in plasticity of traits and gene expression.

In our analyses, plasticity in traits and gene networks were response variables (these

are the above mentioned family means), the 7 climate variables were fixed effect

variables and population was included as a random effect. Each climate variable was

tested against each response variable in a separate model, with day 6 and day 12

measurements analyzed separately (lmer library). In order to do this, we used a subset

of gene network and trait pairs that already showed a climate x treatment effect (before

FDR-correction;**Table S7&8**). These included 5 and 11 tests for gene networks, and

2 and 15 for traits for day 6 and day 12, respectively.

**RESULTS**

**How do responses to drought vary among populations?**

*Drought responses are stronger in gene expression compared to traits*

Six days into the drought experiment, only three of the 8 traits differed significantly

between treatments: unwatered plants had shorter stems, lower stomatal conductance

and more variable SLA (**Table 1**). At the later stage, treatment effects became

pronounced for more traits measured (12 of 15): unwatered plants had lower stomatal

conductance, decreased height, slower growth in height (from day 6 to day 12), higher

SLA, smaller leaf area, denser stomata, smaller stomatal pore size, increasing

pigment accumulation, lower carbohydrates aboveground (but not belowground). In

the earlier stages of the stress, morphological and physiological traits did not show

responses as pronounced as those observed in gene expression patterns, potentially

due to time required for morphological changes to happen (**Figure 2**). In contrast to

most of the traits, the treatment effect was significant for over half of the gene networks

already at day 6 into the treatment. This trend of more pronounced gene expression

response was visible both in co-expression gene networks (**Table 3**, **Figure S2**) and

Multi-dimensional Scaling (MDS) analyses (**Figure 2A-B**), which indicates a lag

between the rapid adjustments in gene expression and trait onset. However, we

should also note that the MDS for gene expression is based on the top 500 genes that

explain the most variation, and we might not have measured morpho-physiological

traits that might also respond as quickly. Notably, population V (Baviaanskloof, a high-

elevation eastern site) did not show a treatment effect in traits at the later stage,

although the effect was quite pronounced in gene expression (**Figure 2B&D** and **3A**).

At day 6 into the treatment, unwatered plants had higher eigengene values for

gene networks GN 2, 4, 5, 6, and 7 and lower values for gene networks GN 1, 9 and

11, relative to watered plants (**Table 3**, **Figure S3**). These modules show gene

enrichment in GO categories for ATPase activity (GN2), oxidoreductase activity (GN4),

cytoskeleton organization (GN5), photorespiration (GN6), response to high light

intensity and temperature (GN7; note that p-value<0.0005, fdr-corrected p-value is not

significant), cell wall organization (GN1) and pyridine-containing compound metabolic

process (GN9). As the stress advanced after 12 days into the drought treatment,

watered and unwatered plants differed significantly for most traits and gene networks.

Significant treatment responses were detected for 10 of 14 gene networks and 12 of

15 traits. (**Tables 1-3**, **Figures S3-5**). In addition to the gene networks with treatment

effects at day 6, GN3 (tropism and carbohydrate metabolism related, and shows

associations with leaf area, height and aboveground soluble carbohydrates) and GN13

(homeostasis related) also had evident treatments effects at day 12, with GN3 having

lower values and GN13 higher values under drought.

*Drought up-regulates stress related genes and down-regulates growth related genes*

The number of genes differentially regulated between earlier and later stages of the

stress and among populations differed significantly (**Figure 3A&B**). For example,

population G (Garcia's Pass) does not have any genes differentially expressed at day

6, indicating either a lower level of stress or lack of accurate perception and timely

signal transduction of the environmental cues to gene expression phenotypes. In

contrast, population B (Bredasdorp) had several hundred differentially expressed

genes by day 6. All populations had at least several hundred differentially expressed

genes by day 12. Two populations (C-Cederberg and S-Swartberg) showed few

differentially expressed genes initially, but over 1500 differentially expressed genes by

day 12 (**Figure 3A**). Although the number of genes regulated upon drought stress was

highly variable, we found 110 and 63 genes to be consistently up- or down- regulated,

respectively, among populations (**Table S5**). The up-regulated genes include heat

shock proteins, auxin signaling and ABA catabolism genes, genes related to drought

stress and photosynthesis, and a leaf amylase; whereas down-regulated genes

included ARF genes (involved in cell elongation and division), and salt stress-, fatty

acid and amino acid biosynthesis-related genes.

*Population differentiation in stress response becomes more evident as the stress*

*advances*

Variation among populations is evident in most of the traits, 11 out of 15 traits were significantly different between the populations at day 6, and all traits except for root length showed a population effect at day 12 (**Table 1&2**). We also observed G x E effects (variation in population responses to drought) to a limited extent on day 6, and to a greater extent on day 12 (**Table 1**). At day 6, only SLA and gene network 13 (enriched for homeostasis GO term) showed variation in responses to drought across populations (**Table 1&3**). SLA was lower under drought for western populations (R, G, C and F), but it was higher under drought in the remaining 4 populations (V, A, B, S; **Table 1**, **Figure S3**). At day 12, populations showed variation in their response to drought for SLA, stomatal size, pigment accumulation, stomatal conductance, stem growth rate, starch biomass aboveground, sugar biomass above and belowground. In addition to GN13 at day 6, GN7 expression (linked to high light intensity and high temperature GO term) also showed a G x E effect at day 12 (**Table 3**).

For each treatment, we also tested for genes differentially expressed among populations (testing for population effect) and timepoints separately. This analysis reveals which genes are differentially expressed among populations, indicating population differentiation, within a treatment and a timepoint. With these analyses, we find 851 and 668 genes to be differentially regulated among populations for the control treatment, and 717 and 786 for drought treatment at day 6 and 12, respectively (FDR-corrected p-value<0.01). We also observed some overlap between these groups (**Figure 3B**); 45 of the genes being differentially expressed among populations in both of the treatments for both days. Interestingly, the majority of genes differentially expressed in a treatment did not overlap between the two timepoints and this holds true for both of the treatments (e.g. purple and pink in **Figure 3B**). Likewise, the genes

differentially expressed in the different treatments did not show overlap within a single

timepoint (e.g. purple and yellow in **Figure 3B**). These results show that together with

a strong treatment effect, there is a pronounced timepoint effect in our experiment for

gene expression (note that the plants were all harvested at the same time of the day

but 6 days apart).

*Most gene networks are associated with physio-morphological traits*

Testing for associations between gene networks and traits, we showed that 12 of the

14 gene networks show correlations with at least one trait (Table S6). For example,

GN 4 (oxidoreductase activity), 5 (cytoskeleton organization), 7 (response to high light

intensity and temperature), 9 (pyridine-containing compound metabolic process) and

13 (homeostasis) all show correlations with plant height at day 6. At day 12, these

same gene networks and an additional GN3 (tropism and carbohydrate metabolism-

related) are correlated with plant height. GN13 also shows correlation with leaf area

and stomatal conductance at day 6, and additionally SLA, stomatal size, aboveground

starch and soluble carbohydrates at day 12. GN4 and 5 show correlations with SLA

and leaf area at day 6, and above- and belowground soluble carbohydrates

additionally at day 12. Similarly, GN6 (photorespiration) is associated with leaf area,

stomatal conductance, aboveground starch and soluble carbohydrates and

belowground soluble carbohydrates at day 12. Additionally, GN7 shows correlations

with SLA at day 6, and GN12 (no GO enrichments) shows correlations with leaf area

at day 12. GN9 shows correlations with leaf area and SLA at both time points, and

with conductance at day 6 and above ground soluble carbohydrates and starch and

below ground soluble carbohydrates at day 12.

**Do the climatic conditions covary with traits, gene expression and their plasticity?**

*Gene networks and functional traits correlate with temperature*

In order to assess climatically driven trait and gene expression variation among populations, we tested for correlations between source-site climate variables and co-expression gene networks and functional traits. We found several gene networks that covary with climatic variables (**Figure 4**; boxed combinations). These include GN 3, 4, 5, 9 and 13 at day 6, and GN3, 4, 5, 6, 7, 10, 12 and 13 at day 12. Most of these gene networks showed strong correlations with elevation and the closely related variables winter minimum temperature and summer maximum temperature. We also observed correlations between mean annual precipitation and GN6, 10 and 13, stomatal conductance and starch content in shoots at day 12. Stem growth showed correlations with elevation, summer maximum and winter minimum temperature. Aboveground starch content co-varied with elevation, mean annual precipitation and intra-annual precipitation variation. Similarly, aboveground soluble carbohydrates showed associations with elevation.

We tested for climate by treatment interaction which revealed if associations between traits and the climate are significant for only one of the treatments. We found that gene expression patterns for GN7 at day 12 show a positive correlation with elevation (and minimum winter temperature) only for plants under drought conditions. Contrastingly, the correlations between GN12 and elevation, and GN13 and mean annual precipitation are significant only for watered plants. Similarly, for stem growth and aboveground starch, the correlations with elevation are significant only for watered

plants, and stomatal conductance with mean annual precipitation only under watered

conditions.

*Plasticity differentiation is associated with climate*

We tested if plasticity in traits and gene expression reflects signatures of climatic

effects. At day 6, plasticity in GN5 (related to cytoskeleton organization) is the only

gene network that show a correlation with a climatic variable: low elevation populations

showed higher plasticity in gene expression patterns of this gene network (**Table 4**).

We also observe higher plasticity in stomatal density in wetter sites. At day 12, we

found that plasticity in GN10 (related to translation) and GN13 (linked to homeostasis)

covaries with mean annual precipitation. For GN10, populations from both the driest

and the wettest sites show higher plasticity, and populations from wetter sites show

higher plasticity for GN13 (**Table 4**, **Figure 5D**). Additionally, plasticity in stem growth

correlates with both elevation and summer maximum temperature. High plasticity in

growth is observed in populations from low elevation and warmer sites (**Figure 5A**).

Similar to GN13, plasticity in leaf area shows correlations with mean annual

precipitation with higher plasticity in populations from wetter sites (**Figure 5B**).

Pigment accumulation shows correlations with temperature heterogeneity

(temperature range within years) (**Figure 5C**) and   maximum temperature. However,

pigment-related associations are driven by a single population in Cederberg, the most

northern population we studied. We should also note that this trend is driven by the

direction of plasticity in this population for pigment content, which is opposite to what

we observe in all the other populations:, plants from the Cederberg population have

higher pigment under drought and lower pigments when watered (Figure S3).

## DISCUSSION

Local adaptation can lead to trait divergence among populations along environmental gradients. Plasticity in traits, as any other phenotypic character, can also be under selection, creating matching patterns between plasticity differentiation and the source site climatic variation among populations. In this study, we show that functional traits and gene expression patterns, and their plasticity are associated with source-site climate in *Protea repens*, suggesting local adaptation in both trait means and their plasticity.

*Drought imposes a complex stress regulating many gene expression networks*

Drought has been a strong driver of plant evolution in the Cape Floristic Region and other Mediterranean climate zones (Allsopp, Colville, and Verboom 2014). When water availability is limited, as it typically is for several months during the summer in most of the region, plants experience a suite of complex stresses (West, Dawson, and February 2012; Ramachandra Reddy, Chaitanya, and Vivekanandan 2004) reflected in regulation of various pathways including ABA-dependent pathways, carbohydrate metabolism, ROS scavenging and photosynthesis under drought (Heschel et al. 2014; Miller et al. 2010). Accordingly, we observe differential gene expression in 10 out of 13 gene networks under drought conditions. For example, the genes with the highest gene network membership in GN5 (related to cytoskeleton organization) are downregulated under drought in all the populations, and this gene network also shows correlations with height, leaf area and aboveground carbohydrate content. Cytoskeleton organization related pathways were also shown to be highly hampered

during drought stress in Arabidopsis (Bhaskara et al. 2017), linking this process to

reduced growth. A general pattern similar to that of GN5 is seen in other gene networks

important for growth (such as GN1 related to cell wall organization and other

carbohydrate metabolism related gene networks). This "shut-off" of growth also

explains the shorter stems and smaller carbohydrate content aboveground under 12

days of drought for all populations. Additionally, the genes that were consistently up-

regulated among all the populations were genes involved in stress responses.

Together these results indicate a trade-off in which resources are channeled from

growth to stress acclimation. A similar trade-off has been observed across many plant

species for herbivory (Kempel et al. 2011; Züst, Rasmann, and Agrawal 2015), cold

stress (Loehle 1998) and drought (Haugen et al. 2008). We also detected reduced

stomatal conductance in all populations in the drought treatment. This indicates a

potential drop in photosynthesis, which might further contribute to the energy deficit

hampering growth. The only gene network, GN6 (enriched for photorespiration related

genes) that shows a correlation with conductance at the later stage of the stress is

also correlated with leaf area and carbohydrate content. Interestingly, we also find a

correlation between this gene network and mean annual precipitation suggesting local

adaptation to water availability via regulation of carbohydrate accumulation.

*Associations with the climate are context dependent*

When there are fitness consequences of a trait that is induced conditionally in

response to environmental stressors, with no fitness costs/benefits when those

stressors are absent, genetic variation that controls trait plasticity will be under

selection only under the inducing stressful environment (Van Dyken and Wade 2010;

Ledón-Rettig et al. 2014). Strikingly, we observed climatic correlations in gene

expression and traits that only exist in either one of the treatments (**Figure 4**). Gene network 7, for example, shows a correlation with elevation and winter minimum temperature only when the plants are under drought stress. Gene ontology enrichments in this module point to regulation by high light and temperature stress. A similar gene network in our previous study in a common garden with more mature plants (18 months old) was also enriched for GO terms related to high light intensity and temperature, and this module showed correlations with stem diameter and stomatal density, but surprisingly not with any climate variables (Akman et al. 2016b). GN7 in this study also shows significant correlations with height at both timepoints and SLA in the earlier timepoint (**Table S6**). The top members of this gene network with Arabidopsis homology are down-regulated in populations coming from higher elevation, one of these genes is 4CL3 (AT1G65060) involved in last step of phenylpropanoid pathway - the upstream pathway anthocyanins are derived from. We also found that plants from warmer sites (higher maximum summer temperature) have higher pigment accumulation (p-value =0.029, fdr-corrected value = 0.106). Previous studies showed a negative effect of heat stress on plant growth (Olszyk et al. 1998; Hasanuzzaman et al. 2013; Bita and Gerats 2013), and anthocyanin accumulation as a protective mechanism against heat, cold, oxidative stress and UV radiation (Hatier and Gould 2009; Steyn et al. 2009; Zeng et al. 2010; Wang et al. 2013). The higher amount of anthocyanins accumulated in populations originating from warmer sites in this study might enable a better protection against stressors such as reactive oxygen species, and thus lead to taller stature. In a parallel finding, under drought conditions, differentially expressed genes among populations were enriched for only oxidative

stress related genes, suggesting selection on expression of these genes along the

climate gradient.

Contrastingly, GN13 (homeostasis related; the best representative gene

showing homology with an Arabidopsis heat shock protein- AT5G59720) shows a

correlation with mean annual precipitation only in watered conditions. Heat shock

proteins are shown to be involved in responses to stresses such as heat, cold, UV

radiation, salt and drought (Chen, Feder, and Kang 2018). Under control conditions,

plants from wetter source sites express genes in this network at lower levels compared

to populations from drier sites, and show a bigger up-regulation under drought,

eventually reaching similar levels of expression for all populations. This gene network

also shows correlations with leaf area, SLA, stomatal pore size and aboveground

carbohydrate content. In plants from drier environments, higher constitutive

expression of these genes might be crucial for a prompt response to drought via

regulation of stomatal pore size, leaf size and thickness. However, plants from

populations that rarely experience drought might not benefit from constitutive

expression of these genes, creating a mismatch between costs and benefits.

*Minimum temperature and elevation show widespread associations with trends in*

*gene expression*

Previously, we showed that functional traits and gene expression patterns as

measured in a common garden with older plants (1.5 year old plants vs 2 months-old

plants in this study) correlated with source site climate (Akman *et al*., 2016). Although

using fewer populations in this study, we observed similar patterns in those

correlations. For example, in both studies we observe associations between gene

expression networks and minimum winter temperature, elevation and mean annual precipitation (note that minimum winter temperature and elevation are highly correlated: Pearson's R = -0.91, **Table S2)**. Taken together, these variables have several associations with gene network variation, as we also found in the previous study. Although *Protea* populations throughout the CFR experience drought at least occasionally, the drought metrics used in this study are not as strong predictors of trait variation as temperature/elevation is under the tested conditions (drought and control in a greenhouse and a common garden). Since gene expression networks include all major groupings of coexpressed genes, they reflect an unbiased pattern in plant responses, and in this case, this pattern indicates fairly widespread variation in gene expression across temperature/elevation gradients.

Why was temperature/elevation associated with population differences in gene expression more often than precipitation?  We can't say conclusively, but note that populations in the CFR experience drought periodically, and although some of the populations we studied experienced a decrease in precipitation over the past 115 years, this effect is not as strong as the increase in temperature over the same period (**Figure** 6, (Kruger and Shongwe 2004; Mason and Jury 1997). Both historically and during recent warming, temperature regimes potentially impose a more continuous selection force than precipitation patterns. Precipitation has  been more variable historically and recent climate change trends are variable across the region, and this variability might make it harder to detect patterns of selection in response to precipitation changes across climate gradients even if measurements are performed under drought conditions as in our experiments. These results also line up with a broader trend: a meta-analysis of 21 plant traits from 25354 species concluded that

temperature is a stronger driver of plant trait evolution compared to precipitation

(Moles et al. 2014).

*Carbohydrate reserves play a role in survival*

Surprisingly, we found a positive correlation between survival under drought and

precipitation; plants from wetter sites survived longer under drought. Populations from

wetter sites also showed higher stomatal conductance and shoot starch content,

indicating higher photosynthetic rates. The fact that populations from wetter sites

survived better under drought is counterintuitive. However, in a sudden drought onset,

like the one imposed in this experiment, carbohydrate reserves might be a decisive

factor in survival, especially in earlier stages of development, and thus if plants from

wetter sites have built up larger reserves, they may survive longer (O'Brien et al. 2014).

*Both climatic heterogeneity and favorable environmental conditions promote plasticity*

*evolution*

Here, we also tested if plasticity is favored and thus selected for in heterogeneous

and/or milder climatic conditions. A heterogeneous environment in theory should

promote plasticity evolution as this can allow for a better match between the phenotype

and the changing environment, potentially surpassing cost of plasticity. Similarly, in

mild environments, plasticity can arise through a relaxed selection in which a mismatch

between the phenotype and the climate does not impose strong fitness costs. We

found evidence for both of these patterns for the functional traits and the gene

expression patterns. However, in this study, we have more evidence for growth

plasticity evolving under milder conditions than heterogeneous environments, i.e.

higher plasticity for growth and GN5 (cytoskeleton organization) in populations from

lower elevation/warmer sites (representing milder climates). Although heat stress can

be detrimental, in the CFR cold stress may impose a stronger selective pressure for

growth (Wilson, Latimer, and Silander 2015). In fact, in a study of 56 Proteaceae

species looking at climate and plant trait correlations, Mitchell et al, showed the most

strongly supported evolutionary correlations to be between winter minimum

temperature (next to mean annual temperature and elevation) and plant height

(Mitchell, Carlson, and Holsinger 2018). Authors also note that this negative effect of

elevation on plant height might be coupled potentially with shallower soils in the high

elevation sites (Campbell and Werger 1988). These combined effects might create

harsher environments which increase the cost of mismatch between phenotype and

the environment, thus decreasing fitness of more plastic individuals. We also find

higher plasticity in wetter sites for GN13 (homeostasis related) and leaf area, and

again these sites represent milder climates in this drought-prone region. The only

correlation we observe for heterogeneity is also related to temperature: plasticity in

pigment accumulation increases with increasing heterogeneity. In environments

where temperature change across seasons is high, levels of anthocyanins might

effectively protect plants against abiotic stressors, thus promoting selection on higher

plasticity.

To conclude, our results suggest that both phenotypic traits and gene expression, and

their plasticity can be under selection along a climate gradient. However it is also

important to note that without studying fitness consequences of the climate, our results

do not prove that this variation is adaptive. We also showed that detecting associations

between these traits and the environmental variables can be a challenging task as

these associations are environment-dependent, and thus can be detected only when

they are measured under certain conditions. Additionally, we found that trait variation

can be explained better with temperature related-variables rather than precipitation

related variables we chose. This could arise from recent temperature increase

associated with climate change that can potentially impose a stronger and more

continuous selection pressure compared to the more variable changes in precipitation.

This study also provides insights into the drivers of plasticity evolution: we found more

evidence for higher plasticity in mild environments compared heterogeneous

environments. Although our results demonstrate population differentiation in traits and

their plasticity and their links to climate, we should also note that we found only 24/350

trait/gene expression and climate associations to be significant together with 9/33 for

plasticity. This relatively low number could either be due to the fact that the climate

variables we chose cannot fully represent the source site environments, that our

experimental design does not capture the genetic variation fully, or that drift has

affected population differentiation. In this regard, future work with long-term transplant

experiments in several common gardens and collection of genomics/transcriptomics

data would enable heritability and Fst/Qst calculations, and coupling these with fitness

measurements would conclusively uncover patterns of local adaptation and the

underlying genetic/gene expression mechanisms in this vulnerable species in the

CFR.

**ACKNOWLEDGEMENTS**

We thank Chris Adams, Megan Varvaro and Sarah Soorya for their assistance with

plant care and the drought experiments, Tony Rebelo, Guy Midgley and Louise Nurrish

for their assistance with seed collection. We thank Jessica Orozco and the Zwieniecki

Lab for their contribution in the carbohydrate analysis. We also thank the Blackman

Lab and Carl Schlichting for their suggestions on an earlier version of the manuscript.

Finally, we thank the reviewers for providing detailed constructive suggestions that

increased the quality of our manuscript. This work was supported by Dimensions of

Biodiversity awarded to Carl Schlichting, Cynthia S. Jones and Kent E. Holsinger

(award no: 1046328) and Andrew M. Latimer (award no: 1045985).

**REFERENCES**

Acasuso-Rivero, Cristina, Courtney J. Murren, Carl D. Schlichting, and Ulrich K.
Steiner. 2019. "Adaptive Phenotypic Plasticity for Life-History and Less Fitness-
Related Traits." *Proceedings. Biological Sciences / The Royal Society* 286 (1904):
20190653.
Ackerly, David D., Susan A. Dudley, Sonia E. Sultan, Johanna Schmitt, James S.
Coleman, C. Randall Linder, Darren R. Sandquist, et al. 2000. "The Evolution of
Plant Ecophysiological Traits: Recent Advances and Future Directions: New
Research Addresses Natural Selection, Genetic Constraints, and the Adaptive
Evolution of Plant Ecophysiological Traits." *Bioscience* 50 (11): 979–95.
Akman, Melis, Amit V. Bhikharie, Elizabeth H. McLean, Alex Boonman, Eric J. W.
Visser, M. Eric Schranz, and Peter H. van Tienderen. 2012. "Wait or Escape?
Contrasting Submergence Tolerance Strategies of Rorippa Amphibia, Rorippa
Sylvestris and Their Hybrid." *Annals of Botany* 109 (7): 1263–76.
Akman, Melis, Jane E. Carlson, Kent E. Holsinger, and Andrew M. Latimer. 2016.
"Transcriptome Sequencing Reveals Population Differentiation in Gene
Expression Linked to Functional Traits and Environmental Gradients in the South
African Shrub Protea Repens." *The New Phytologist* 210 (1): 295–309.
Allen, Craig D., Alison K. Macalady, Haroun Chenchouni, Dominique Bachelet, Nate
McDowell, Michel Vennetier, Thomas Kitzberger, et al. 2010. "A Global Overview
of Drought and Heat-Induced Tree Mortality Reveals Emerging Climate Change
Risks for Forests." *Forest Ecology and Management* 259 (4): 660–84.
Allsopp, Nicky, Jonathan F. Colville, and G. Anthony Verboom. 2014. *Fynbos:
Ecology, Evolution, and Conservation of a Megadiverse Region*. Oxford
University Press.
Alpert, Peter, and Ellen L. Simms. 2002. "The Relative Advantages of Plasticity and
Fixity in Different Environments: When Is It Good for a Plant to Adjust?"
*Evolutionary Ecology* 16 (3): 285–97.
Aubin, I., A. D. Munson, F. Cardou, P. J. Burton, N. Isabel, J. H. Pedlar, A. Paquette,
et al. 2016. "Traits to Stay, Traits to Move: A Review of Functional Traits to Assess
Sensitivity and Adaptive Capacity of Temperate and Boreal Trees to Climate
Change." *Environmental Review* 24 (2): 164–86.
Barshis, Daniel J., Jason T. Ladner, Thomas A. Oliver, François O. Seneca, Nikki

Traylor-Knowles, and Stephen R. Palumbi. 2013. "Genomic Basis for Coral Resilience to Climate Change." *Proceedings of the National Academy of Sciences* 110 (4): 1387–92.

Bates, Douglas, Martin Maechler, and Ben Bolker. 2012. "lme4: Linear Mixed-Effects Models Using S4 Classes."

Baythavong, Brooke S. 2011. "Linking the Spatial Scale of Environmental Variation and the Evolution of Phenotypic Plasticity: Selection Favors Adaptive Plasticity in Fine-Grained Environments." *The American Naturalist* 178 (1): 75–87.

Baythavong, Brooke S., and Maureen L. Stanton. 2010. "Characterizing Selection on Phenotypic Plasticity in Response to Natural Environmental Heterogeneity." *Evolution; International Journal of Organic Evolution* 64 (10): 2904–20.

Bhaskara, Govinal Badiger, Tuan-Nan Wen, Thao Thi Nguyen, and Paul E. Verslues. 2017. "Protein Phosphatase 2Cs and Microtubule-Associated Stress Protein 1 Control Microtubule Stability, Plant Growth, and Drought Response." *The Plant Cell* 29 (1): 169–91.

Bita, Craita E., and Tom Gerats. 2013. "Plant Tolerance to High Temperature in a Changing Environment: Scientific Fundamentals and Production of Heat Stress-Tolerant Crops." *Frontiers in Plant Science* 4 (July): 273.

Bolger, Anthony M., Marc Lohse, and Bjoern Usadel. 2014. "Trimmomatic: A Flexible Trimmer for Illumina Sequence Data." *Bioinformatics* , April. https://doi.org/10.1093/bioinformatics/btu170.

Bonamour, Suzanne, Luis-Miguel Chevin, Anne Charmantier, and Céline Teplitsky. 2019. "Phenotypic Plasticity in Response to Climate Change: The Importance of Cue Variation." *Philosophical Transactions of the Royal Society B: Biological Sciences*. https://doi.org/10.1098/rstb.2018.0178.

Bradshaw, A. D., and K. Hardwick. 1989. "Evolution and Stress—genotypic and Phenotypic Components." *Biological Journal of the Linnean Society. Linnean Society of London* 37 (1-2): 137–55.

Campbell, B. M., and M. J. A. Werger. 1988. "Plant Form in the Mountains of the Cape, South Africa." *The Journal of Ecology* 76 (3): 637–53.

Carlson, Jane E., Christopher A. Adams, and Kent E. Holsinger. 2016. "Intraspecific Variation in Stomatal Traits, Leaf Traits and Physiology Reflects Adaptation along Aridity Gradients in a South African Shrub." *Annals of Botany* 117 (1): 195–207.

Carlson, Jane E., and Kent E. Holsinger. 2012. "Developmental Plasticity in <italic>Protea</italic> as an Evolutionary Response to Environmental Clines in the Cape Floristic Region." *PloS One* 7 (12): e52035.

Carlson, Jane E., Kent E. Holsinger, and Rachel Prunier. 2011. "Plant Responses to Climate in the Cape Floristic Region of South Africa: Evidence for Adaptive Differentiation in the Proteaceae." *Evolution; International Journal of Organic Evolution* 65 (1): 108–24.

Cavender-Bares, Jeannine, and José Alberto Ramírez-Valiente. 2017. "Physiological Evidence from Common Garden Experiments for Local Adaptation and Adaptive Plasticity to Climate in American Live Oaks (Quercus Section Virentes): Implications for Conservation under Global Change." In *Oaks Physiological Ecology. Exploring the Functional Diversity of Genus Quercus L.*, 107–35. Springer.

Chen, Bing, Martin E. Feder, and Le Kang. 2018. "Evolution of Heat-Shock Protein Expression Underlying Adaptive Responses to Environmental Stress." *Molecular*

*Ecology*. https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.14769.

Chevin, Luis-Miguel, and Ary A. Hoffmann. 2017. "Evolution of Phenotypic Plasticity
in Extreme Environments." *Philosophical Transactions of the Royal Society of*
*London. Series B, Biological Sciences* 372 (1723).
https://doi.org/10.1098/rstb.2016.0138.

Chevin, Luis-Miguel, Russell Lande, and Georgina M. Mace. 2010. "Adaptation,
Plasticity, and Extinction in a Changing Environment: Towards a Predictive
Theory." *PLoS Biology* 8 (4): e1000357.

Cochrane, Anne, Gemma L. Hoyle, Colin J. Yates, Jeff Wood, and Adrienne B.
Nicotra. 2015. "The Phenotypic Response of Co-Occurring Banksia Species to
Warming and Drying." *Plant Ecology* 216 (1): 27–39.

Colicchio, Jack M., Patrick J. Monnahan, John K. Kelly, and Lena C. Hileman. 2015.
"Gene Expression Plasticity Resulting from Parental Leaf Damage in Mimulus
Guttatus." *The New Phytologist* 205 (2): 894–906.

Cowling, R. M. 2009. "From Winter Wet to Winter Dry: Fynbos Diversity and Evolution
along the West–east Gradient." *South African Journal of Botany: Official Journal*
*of the South African Association of Botanists = Suid-Afrikaanse Tydskrif Vir*
*Plantkunde: Amptelike Tydskrif van Die Suid-Afrikaanse Genootskap van*
*Plantkundiges* 2 (75): 386–87.

Cuenca Cambronero, Maria, Jordan Beasley, Stephen Kissane, and Luisa Orsini.
2018. "Evolution of Thermal Tolerance in Multifarious Environments." *Molecular*
*Ecology* 27 (22): 4529–41.

Daac, L. P. 2011. "ASTER L1B." *Sioux Falls, South Dakota: USGS/Earth Resources*
*Observation and Science (EROS) Center*.

Dayan, David I., Douglas L. Crawford, and Marjorie F. Oleksiak. 2015. "Phenotypic
Plasticity in Gene Expression Contributes to Divergence of Locally Adapted
Populations of Fundulus Heteroclitus." *Molecular Ecology* 24 (13): 3345–59.

Dudley, Susan A., and Johanna Schmitt. 1996. "Testing the Adaptive Plasticity
Hypothesis: Density-Dependent Selection on Manipulated Stem Length in
Impatiens Capensis." *The American Naturalist* 147 (3): 445–65.

Eden, Eran, Roy Navon, Israel Steinfeld, Doron Lipson, and Zohar Yakhini. 2009.
"GOrilla: A Tool for Discovery and Visualization of Enriched GO Terms in Ranked
Gene Lists." *BMC Bioinformatics* 10 (1): 48.

Etterson, Julie R. 2004. "Evolutionary Potential of Chamaecrista Fasciculata in
Relation to Climate Change. I. Clinal Patterns of Selection along an
Environmental Gradient in the Great Plains." *Evolution; International Journal of*
*Organic Evolution* 58 (7): 1446–58.

Forsman, A. 2015. "Rethinking Phenotypic Plasticity and Its Consequences for
Individuals, Populations and Species." *Heredity* 115 (4): 276–84.

Fu, Limin, Beifang Niu, Zhengwei Zhu, Sitao Wu, and Weizhong Li. 2012. "CD-HIT:
Accelerated for Clustering the next-Generation Sequencing Data." *Bioinformatics*
28 (23): 3150–52.

Geber, Monica A., and Lauren R. Griffen. 2003. "Inheritance and Natural Selection on
Functional Traits." *International Journal of Plant Sciences* 164 (S3): S21–42.

Ghalambor, Cameron K., Kim L. Hoke, Emily W. Ruell, Eva K. Fischer, David N.
Reznick, and Kimberly A. Hughes. 2015. "Non-Adaptive Plasticity Potentiates
Rapid Adaptive Evolution of Gene Expression in Nature." *Nature* 525 (7569): 372–
75.

Granados-Cifuentes, Camila, Anthony J. Bellantuono, Tyrone Ridgway, Ove Hoegh-Guldberg, and Mauricio Rodriguez-Lanetty. 2013. "High Natural Gene Expression Variation in the Reef-Building Coral Acropora Millepora: Potential for Acclimative and Adaptive Plasticity." *BMC Genomics* 14 (1): 1–12.

Grime, J. P. 1994. "The Role of Plasticity in Exploiting Environmental Heterogeneity." *Exploitation of Environmental Heterogeneity by Plants: Ecophysiological Processes above-and Belowground* 1: 19.

Gugger, Simona, Halil Kesselring, Jürg Stöcklin, and Elena Hamann. 2015. "Lower Plasticity Exhibited by High- versus Mid-Elevation Species in Their Phenological Responses to Manipulated Temperature and Drought." *Annals of Botany* 116 (6): 953–62.

Hannon, G. J. n.d. "Fastx-Toolkit. 2010." *GNU Affero General Public License* 706.

Hasanuzzaman, Mirza, Kamrun Nahar, Md Mahabub Alam, Rajib Roychowdhury, and Masayuki Fujita. 2013. "Physiological, Biochemical, and Molecular Mechanisms of Heat Stress Tolerance in Plants." *International Journal of Molecular Sciences* 14 (5): 9643–84.

Hatier, Jean-Hugues B., and Kevin S. Gould. 2009. "Anthocyanin Function in Vegetative Organs." In *Anthocyanins: Biosynthesis, Functions, and Applications*, edited by Chris Winefield, Kevin Davies, and Kevin Gould, 1–19. New York, NY: Springer New York.

Haugen, Riston, Lexi Steffes, Joy Wolf, Paul Brown, Steven Matzner, and David H. Siemens. 2008. "Evolution of Drought Tolerance and Defense: Dependence of Tradeoffs on Mechanism, Environment and Defense Switching." *Oikos* 117 (2): 231–44.

Heschel, M. Shane, Ann Evankow, Katherine Bibee Wolfson, Jane E. Carlson, and Kent E. Holsinger. 2014. "Drought Response Diversification in African Protea Species." *International Journal of Plant Sciences* 175 (4): 442–49.

Hewitson, B. C., and R. G. Crane. 2006. "Consensus between GCM Climate Change Projections with Empirical Downscaling: Precipitation Downscaling over South Africa." *International Journal of Climatology* 26 (10): 1315–37.

Jacobs, B. S., and A. M. Latimer. 2012. "Analyzing Reaction Norm Variation in the Field vs. Greenhouse: Comparing Studies of Plasticity and Its Adaptive Value in Two Species of Erodium." *Perspectives in Plant Ecology, Evolution and Systematics* 14 (5): 325–34.

Kempel, Anne, Martin Schädler, Thomas Chrobock, Markus Fischer, and Mark van Kleunen. 2011. "Tradeoffs Associated with Constitutive and Induced Plant Resistance against Herbivory." *Proceedings of the National Academy of Sciences of the United States of America* 108 (14): 5685–89.

Kruger, Andries C., and Stephen Shongwe. 2004. "Temperature Trends in South Africa: 1960--2003." *International Journal of Climatology* 24 (15): 1929–45.

Langfelder, Peter, and Steve Horvath. 2008. "WGCNA: An R Package for Weighted Correlation Network Analysis." *BMC Bioinformatics* 9 (1): 559.

Langmead, Ben, and Steven L. Salzberg. 2012. "Fast Gapped-Read Alignment with Bowtie 2." *Nature Methods* 9 (4): 357–59.

Lasky, Jesse R., David L. Des Marais, David B. Lowry, Inna Povolotskaya, John K. McKay, James H. Richards, Timothy H. Keitt, and Thomas E. Juenger. 2014. "Natural Variation in Abiotic Stress Responsive Gene Expression and Local Adaptation to Climate in Arabidopsis Thaliana." *Molecular Biology and Evolution*

31 (9): 2283–96.

Latimer, Andrewm, J. A. Silander Jr., A. G. Rebelo, and G. F. Midgley. 2009. "Experimental Biogeography: The Role of Environmental Gradients in High Geographic Diversity in Cape Proteaceae." *Oecologia* 160 (1): 151–62.

Lázaro-Nogal, Ana, Silvia Matesanz, Alice Godoy, Fernanda Pérez-Trautman, Ernesto Gianoli, and Fernando Valladares. 2015. "Environmental Heterogeneity Leads to Higher Plasticity in Dry-Edge Populations of a Semi-Arid Chilean Shrub: Insights into Climate Change Responses." *The Journal of Ecology* 103 (2): 338–50.

Lázaro-Nogal, Ana, Silvia Matesanz, Lea Hallik, Alisa Krasnova, Anna Traveset, and Fernando Valladares. 2016. "Population Differentiation in a Mediterranean Relict Shrub: The Potential Role of Local Adaptation for Coping with Climate Change." *Oecologia* 180 (4): 1075–90.

Ledón-Rettig, Cris C., David W. Pfennig, Amanda J. Chunco, and Ian Dworkin. 2014. "Cryptic Genetic Variation in Natural Populations: A Predictive Framework." *Integrative and Comparative Biology* 54 (5): 783–93.

Loehle, Craig. 1998. "Height Growth Rate Tradeoffs Determine Northern and Southern Range Limits for Trees." *Journal of Biogeography* 25 (4): 735–42.

Markham, Charles G. 1970. "SEASONALITY OF PRECIPITATION IN THE UNITED STATES." *Annals of the Association of American Geographers. Association of American Geographers* 60 (3): 593–97.

Mason, S. J., and M. R. Jury. 1997. "Climatic Variability and Change over Southern Africa: A Reflection on Underlying Processes." *Progress in Physical Geography: Earth and Environment* 21 (1): 23–50.

Miller, Gad, Nobuhiro Suzuki, Sultan Ciftci-Yilmaz, and Ron Mittler. 2010. "Reactive Oxygen Species Homeostasis and Signalling during Drought and Salinity Stresses." *Plant, Cell & Environment* 33 (4): 453–67.

Mitchell, Nora, Jane E. Carlson, and Kent E. Holsinger. 2018. "Correlated Evolution between Climate and Suites of Traits along a Fast-Slow Continuum in the Radiation of Protea." *Ecology and Evolution* 8 (3): 1853–66.

Moles, Angela T., Sarah E. Perkins, Shawn W. Laffan, Habacuc Flores-Moreno, Monica Awasthy, Marianne L. Tindall, Lawren Sack, et al. 2014. "Which Is a Better Predictor of Plant Traits: Temperature or Precipitation?" Edited by Aveliina Helm. *Journal of Vegetation Science: Official Organ of the International Association for Vegetation Science* 25 (5): 1167–80.

Morris, Matthew R. J., Romain Richard, Erica H. Leder, Rowan D. H. Barrett, Nadia Aubin-Horth, and Sean M. Rogers. 2014. "Gene Expression Plasticity Evolves in Response to Colonization of Freshwater Lakes in Threespine Stickleback." *Molecular Ecology* 23 (13): 3226–40.

Nicotra, Adrienne B., Deborah L. Segal, Gemma L. Hoyle, Aaron W. Schrey, Koen J. F. Verhoeven, and Christina L. Richards. 2015. "Adaptive Plasticity and Epigenetic Variation in Response to Warming in an Alpine Plant." *Ecology and Evolution* 5 (3): 634–47.

O'Brien, Michael J., Sebastian Leuzinger, Christopher D. Philipson, John Tay, and Andy Hector. 2014. "Drought Survival of Tropical Tree Seedlings Enhanced by Non-Structural Carbohydrate Levels." *Nature Climate Change* 4 (June): 710.

Olszyk, D., Claudia Wise, Erica VanEss, and David Tingey. 1998. "Elevated Temperature but Not Elevated CO2 Affects Long-Term Patterns of Stem Diameter

and Height of Douglas-Fir Seedlings." *Canadian Journal of Forest Research. Journal Canadien de La Recherche Forestiere* 28 (7): 1046–54.

Pachter, Lior. 2011. "Models for Transcript Quantification from RNA-Seq." *arXiv [q-bio.GN]*. arXiv. http://arxiv.org/abs/1104.3889.

Palacio-López, Kattia, Brian Beckage, Samuel Scheiner, and Jane Molofsky. 2015. "The Ubiquity of Phenotypic Plasticity in Plants: A Synthesis." *Ecology and Evolution* 5 (16): 3389–3400.

Pohlman, Catherine L., Adrienne B. Nicotra, and Brad R. Murray. 2005. "Geographic Range Size, Seedling Ecophysiology and Phenotypic Plasticity in Australian Acacia Species." *Journal of Biogeography* 32 (2): 341–51.

Pratt, Jessica D., and Kailen A. Mooney. 2013. "Clinal Adaptation and Adaptive Plasticity in Artemisia Californica: Implications for the Response of a Foundation Species to Predicted Climate Change." *Global Change Biology* 19 (8): 2454–66.

Prunier, R., K. E. Holsinger, and J. E. Carlson. 2012. "The Effect of Historical Legacy on Adaptation: Do Closely Related Species Respond to the Environment in the Same Way?" *Journal of Evolutionary Biology* 25 (8): 1636–49.

Ramachandra Reddy, Attipalli, Kolluru Viswanatha Chaitanya, and Munusamy Vivekanandan. 2004. "Drought-Induced Responses of Photosynthesis and Antioxidant Metabolism in Higher Plants." *Journal of Plant Physiology* 161 (11): 1189–1202.

Rebelo, A. 2001. "Saso Proteas: A Field Guide to the Proteas of Southern Africa." http://kbd.kew.org/kbd/detailedresult.do?id=352313.

Reed, Thomas E., Robin S. Waples, Daniel E. Schindler, Jeffrey J. Hard, and Michael T. Kinnison. 2010. "Phenotypic Plasticity and Population Viability: The Importance of Environmental Predictability." *Proceedings. Biological Sciences / The Royal Society* 277 (1699): 3391–3400.

Sauquet, Hervé, Peter H. Weston, Cajsa Lisa Anderson, Nigel P. Barker, David J. Cantrill, Austin R. Mast, and Vincent Savolainen. 2009. "Contrasted Patterns of Hyperdiversification in Mediterranean Hotspots." *Proceedings of the National Academy of Sciences* 106 (1): 221–25.

Scheiner, Samuel M. 2013. "The Genetics of Phenotypic Plasticity. XII. Temporal and Spatial Heterogeneity." *Ecology and Evolution* 3 (13): 4596–4609.

Schlichting, Carl D., and Massimo Pigliucci. 1995. "Gene Regulation, Quantitative Genetics and the Evolution of Reaction Norms." *Evolutionary Ecology* 9 (2): 154–68.

Schneider, Ralf F., Yuanhao Li, Axel Meyer, and Helen M. Gunter. 2014. "Regulatory Gene Networks That Shape the Development of Adaptive Phenotypic Plasticity in a Cichlid Fish." *Molecular Ecology* 23 (18): 4511–26.

Schulze, R. E. 2007. "South African Atlas of Climatology and Agrohydrology: WRC Report 1489/1/06." Water Resource Commission.

Schulze, Roland Edgar. 1997. "South African Atlas of Agrohydrology and-Climatology." 1868452719. Pretoria, South Africa: Water Research Commission.

Steyn, Willem J., Stephanie J. E. Wand, Gerard Jacobs, Richard C. Rosecrance, and Stephanie C. Roberts. 2009. "Evidence for a Photoprotective Function of Low-Temperature-Induced Anthocyanin Accumulation in Apple and Pear Peel." *Physiologia Plantarum* 136 (4): 461–72.

Stoks, Robby, Lynn Govaert, Kevin Pauwels, Bastiaan Jansen, and Luc De Meester. 2016. "Resurrecting Complexity: The Interplay of Plasticity and Rapid Evolution

in the Multiple Trait Response to Strong Changes in Predation Pressure in the Water Flea Daphnia Magna." *Ecology Letters* 19 (2): 180–90.

Sultan, Sonia E. 2003. "Phenotypic Plasticity in Plants: A Case Study in Ecological Development." *Evolution & Development* 5 (1): 25–33.

Team, R. Core. 2014. "R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2014."

Valliere, Justin M., Irina C. Irvine, Louis Santiago, and Edith B. Allen. 2017. "High N, Dry: Experimental Nitrogen Deposition Exacerbates Native Shrub Loss and Nonnative Plant Invasion during Extreme Drought." *Global Change Biology* 23 (10): 4333–45.

Van Dyken, J. David, and Michael J. Wade. 2010. "The Genetic Signature of Conditional Expression." *Genetics* 184 (2): 557–70.

Van Kleunen, Mark, and Markus Fischer. 2005. "Constraints on the Evolution of Adaptive Phenotypic Plasticity in Plants." *The New Phytologist* 166 (1): 49–60.

Van Wilgen, B. W., G. G. Forsyth, and H. De Klerk. 2010. "Fire Management in Mediterranean-climate Shrublands: A Case Study from the Cape Fynbos, South Africa." *Journal of Applied*. https://agupubs.onlinelibrary.wiley.com/doi/full/10.1111/j.1365-2664.2010.01800.x.

Vitasse, Yann, Günter Hoch, Christophe F. Randin, Armando Lenz, Chris Kollas, J. F. Scheepens, and Christian Körner. 2013. "Elevational Adaptation and Plasticity in Seedling Phenology of Temperate Deciduous Tree Species." *Oecologia* 171 (3): 663–78.

Wadgymar, Susana M., Rachel M. Mactavish, and Jill T. Anderson. 2018. "Transgenerational and Within-Generation Plasticity in Response to Climate Change: Insights from a Manipulative Field Experiment across an Elevational Gradient." *The American Naturalist* 192 (6): 698–714.

Wang, Hongxia, Weijuan Fan, Hong Li, Jun Yang, Jirong Huang, and Peng Zhang. 2013. "Functional Characterization of Dihydroflavonol-4-Reductase in Anthocyanin Biosynthesis of Purple Sweet Potato Underlies the Direct Evidence of Anthocyanins Function against Abiotic Stresses." *PloS One* 8 (11): e78484.

Wennersten, Lena, and Anders Forsman. 2012. "Population-Level Consequences of Polymorphism, Plasticity and Randomized Phenotype Switching: A Review of Predictions." *Biological Reviews of the Cambridge Philosophical Society* 87 (3): 756–67.

West, A. G., T. E. Dawson, and E. C. February. 2012. "Diverse Functional Responses to Drought in a Mediterranean-type Shrubland in South Africa." *New*. https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1469-8137.2012.04170.x.

West-Eberhard, Mary Jane. 2003. *Developmental Plasticity and Evolution*. Oxford University Press.

Whitehead, Andrew. 2012. "Comparative Genomics in Ecological Physiology: Toward a More Nuanced Understanding of Acclimation and Adaptation." *The Journal of Experimental Biology* 215 (6): 884–91.

Wilson, Adam M., Andrew M. Latimer, and John A. Silander. 2015. "Climatic Controls on Ecosystem Resilience: Postfire Regeneration in the Cape Floristic Region of South Africa." *Proceedings of the National Academy of Sciences* 112 (29): 9058–63.

Young, Derek J. N., Jens T. Stevens, J. Mason Earles, Jeffrey Moore, Adam Ellis, Amy

L. Jirka, and Andrew M. Latimer. 2017. "Long-Term Climate and Competition
Explain Forest Mortality Patterns under Extreme Drought." *Ecology Letters* 20 (1):
78–86.
Zeng, Xue-Qin, Wah Soon Chow, Li-Juan Su, Xin-Xiang Peng, and Chang-Lian Peng.
2010. "Protective Effect of Supplemental Anthocyanins on Arabidopsis Leaves
under High Light." *Physiologia Plantarum* 138 (2): 215–25.
Züst, Tobias, Sergio Rasmann, and Anurag A. Agrawal. 2015. "Growth--Defense
Tradeoffs for Two Major Anti-Herbivore Traits of the Common Milkweed Asclepias
Syriaca." *Oikos* 124 (10): 1404–15.

**Data Accessibility**

All sequencing data will be made available at NCBI SRA. Bioinformatics and R scripts

will be made available upon acceptance of the manuscript on Akman's github. Climate

data, phenotypic measurement and gene expression counts will be made available in

Dryad.

**Author contributions**

Akman, Carlson and Latimer designed the research. Akman and Carlson performed

the dry down experiments. Akman prepared sequencing libraries and performed gene

expression analyses. Akman, Carlson and Latimer performed climate association

analyses and wrote the manuscript.

1061

## FIGURES

Fig 1. Map of CFR and locations of sampling sites (A); Principal component analysis for the climatic variables used in this study

Fig 2. Multi-dimensional scaling (MDS) plots for gene expression patterns at day 6 (A) and day 12 (B); principal component analysis of measured traits at day 6 (C) and day 12 (D) into the drought treatment.

Fig 3. Number of differentially expressed genes for the 8 populations for after 6 and 12 days of the drought treatment (A). Note that population A is not sampled at day 12, and population G does not have any differentially expressed genes between the treatments at day 6. Number of differentially expressed genes among populations under control and drought conditions for day 6 and 12 separately (B). For example, the intersecting ellipse between Day6 control and DAY6 drought indicates number of genes that are differentially expressed among populations, regardless of the treatment.

Fig 4. Heatmap of standardized regression coefficients for climate variables as predictors and traits and gene networks (GN) as response variables for measurements (A) on day 6 or (B) day 12. For variables with significant interaction effects, each treatment was analyzed separately to estimate slopes and test their significance within treatments. Solid boxes indicate a significant association and the box is divided in cases where there is a significant treatment and climate effect, indicating that the climate association was significant only in one of the treatments. The upper half of the box represents the watered treatment and the lower half represents the dry treatment. Stars indicate significant correlations for either of the two treatments.

Fig 5. Response curves for the studied 8 populations under well-watered and drought conditions and the associations between plasticity and climate variables for growth (A), leaf area (B) and gene network 13 (C).

Fig 6. Mean annual precipitation (A) and mean annual temperature (B) averaged across the 8 population sites between 1901-2016.

## TABLES

Table 1. Results of linear mixed models testing trait differences of *Protea repens* seedlings at 6 and 12 days into the dry down experiment. The model compares values between watered and unwatered treatments as well as among 8 wild-sourced populations and the population by treatment interaction. Significant values are bolded.

Table 2. Results of linear mixed models testing for differences in growth and carbohydrate storage between watered and unwatered *Protea repens* seedlings and among 8 populations at the end of the dry-down experiment (day 12). Significant values are bolded.

Table 3. Results of linear mixed models comparing eigengene values of the gene
networks (GN0-GN13) at 6 and 12 days into the dry down experiment for watered and
unwatered plants of *P. repens*. Significant values are bolded.

Table 4. Significant effects from mixed models testing associations between climate
variables and plasticity measures, for traits and eigengenes of gene networks (GN) on
day 6 and day 12.

## SUPPLEMENTARY INFORMATION

**Table 1**. Results of linear mixed models testing functional trait and performance differences of *Protea repens* seedlings at 6 and 12 days into the dry down experiment. The model compares values between watered and unwatered treatments as well as among 8 wild-sourced populations and the population by treatment interaction. Significant values are bolded.

| Response variable | Fixed effects | day 6 | | | | day 12 | | | |
|---|---|---|---|---|---|---|---|---|---|
| | | df | Chi-sq | P-value | q-val | df | Chi-sq | p-value | q-value |
| Leaf area (cm$^2$) | Population | 7 | 49.91 | **<0.0001** | **<0.0001** | 6 | 69.16 | **<0.0001** | **<0.0001** |
| | Treatment | 1 | 0.17 | 0.678 | 0.881 | 1 | 23.39 | **<0.0001** | **<0.0001** |
| | Pop. × Trt. | 7 | 8.61 | 0.282 | 0.423 | 6 | 13.3 | 0.039 | 0.050 |
| SLA (cm$^2$/g) Log transform | Population | 7 | 5.15 | 0.642 | 0.881 | **6** | **15.35** | **0.018** | **0.027** |
| | Treatment | **1** | **12.13** | **0.0005** | **0.002** | 1 | 73.14 | **<0.0001** | **<0.0001** |
| | Pop. × Trt. | **7** | **19.23** | **0.008** | **0.027** | **6** | **16.71** | **0.010** | **0.017** |
| LWR | Population | 7 | 67.48 | **<0.0001** | **<0.0001** | 6 | 78.55 | **<0.0001** | **<0.0001** |
| | Treatment | 1 | 0.06 | 0.814 | 0.881 | 1 | 1.86 | 0.173 | 0.210 |
| | Pop. × Trt. | 7 | 3.47 | 0.839 | 0.881 | 6 | 1.52 | 0.958 | 0.958 |
| Stomatal density | Population | **7** | **17.46** | **0.015** | **0.040** | **6** | **29.73** | **<0.0001** | **0.0001** |
| | Treatment | 1 | 3.07 | 0.080 | 0.192 | **1** | **12.95** | **0.0003** | **0.0007** |
| | Pop. × Trt. | 7 | 9.56 | 0.215 | 0.369 | 6 | 3.59 | 0.732 | 0.765 |
| Stomatal pore size (um) | Population | 7 | 3.87 | 0.795 | 0.881 | **6** | **21.92** | **0.001** | **0.002** |
| | Treatment | 1 | 0.04 | 0.844 | 0.881 | **1** | **27.68** | **<0.0001** | **<0.0001** |
| | Pop. × Trt. | 7 | 3.05 | 0.881 | 0.881 | **6** | **14.27** | **0.027** | **0.038** |
| Stem pigmentation | Population | 7 | 34.13 | **<0.0001** | **<0.0001** | 7 | 38.51 | **<0.0001** | **<0.0001** |
| | Treatment | 1 | 1.85 | 0.174 | 0.336 | 1 | 0.42 | 0.515 | 0.579 |
| | Pop. × Trt. | 7 | 10.12 | 0.182 | 0.336 | 7 | 5.71 | 0.574 | 0.630 |
| Stem height (cm) | Population | **7** | **105.34** | **<0.0001** | **<0.0001** | 7 | 121.24 | **<0.0001** | **<0.0001** |
| | Treatment | **1** | **5.9** | **0.015** | **0.040** | 1 | 49.4 | **<0.0001** | **<0.0001** |
| | Pop. × Trt. | 7 | 4.1 | 0.768 | 0.881 | 7 | 6.86 | 0.443 | 0.511 |
| Stomatal conductance | Population | 7 | 0 | 0.254 | 0.406 | **6** | **17.52** | **0.008** | **0.014** |
| | Treatment | **1** | **0** | **<0.0001** | **<0.0001** | **1** | **104.7** | **<0.0001** | **<0.0001** |
| | Pop. × Trt. | 7 | 0 | 0.148 | 0.323 | **6** | **23.43** | **0.0007** | **0.001** |

**Table 2**. Results of linear mixed models testing for differences in growth, carbohydrate storage, root length and pigment accumulation between watered and unwatered *Protea repens* seedlings and among 8 populations at the end of the dry-down experiment (day 12). Significant values are bolded.

| Response variable | Fixed effects | df | Chi-sq | p-val | q-val |
|---|---|---|---|---|---|
| Soluble sugars mass (g) belowground | Population | **7** | **55.57** | **<0.0001** | **<0.0001** |
| | Treatment | **1** | **6.33** | **0.012** | **0.019** |
| | Pop. × Trt. | **7** | **17.52** | **0.014** | **0.022** |
| Soluble sugars mass (g) aboveground | Population | **7** | **80.08** | **<0.0001** | **<0.0001** |
| | Treatment | **1** | **25.69** | **<0.0001** | **<0.0001** |
| | Pop. × Trt. | **7** | **24.27** | **0.001** | **0.002** |
| Starch mass (g) belowground | Population | **7** | **21.06** | **0.004** | **0.008** |
| | Treatment | 1 | 0.10 | 0.748 | 0.765 |
| | Pop. × Trt. | 7 | 4.57 | 0.712 | 0.763 |
| Starch mass (g) aboveground | Population | **7** | **57.40** | **<0.0001** | **<0.0001** |
| | Treatment | **1** | **60.10** | **<0.0001** | **<0.0001** |
| | Pop. × Trt. | **7** | **100.76** | **<0.0001** | **<0.0001** |
| Growth rate (height on Day 12-6) | Population | **7** | **34.85** | **<0.0001** | **<0.0001** |
| | Treatment | **1** | **194.34** | **<0.0001** | **<0.0001** |
| | Pop. × Trt. | **7** | **26.33** | **0.0004** | **0.0009** |
| Root length (mm) | Population | 6 | 9.36 | 0.154 | 0.192 |
| | Treatment | **1** | **6.66** | **0.010** | **0.017** |
| | Pop. × Trt. | 6 | 6.3 | 0.391 | 0.463 |
| Stem pigment accumulation | Population | **7** | **15.47** | **0.031** | **0.041** |
| | Treatment | **1** | **5.08** | **0.024** | **0.035** |
| | Pop. × Trt. | **7** | **15.58** | **0.029** | **0.040** |

**Table 3**. Results of linear mixed models comparing eigengene values (GN0-GN13) at 6 and 12 days into the dry down experiment for watered and unwatered plants of *P. repens*. Significant values are bolded.

|  | Fixed effects | Fixed effects at 6 days | | | | Fixed effects at 12 days | | | |
|---|---|---|---|---|---|---|---|---|---|
|  |  | df | Chi-sq | p-value | q-val | ddf | Chi-sq | p-value | q-val |
| GN0 | Population | **7** | **24.89** | **0.0008** | **0.0042** | **6** | **80.38** | **<0.0001** | **<0.0001** |
|  | Treatment | 1 | 2.05 | 0.152 | 0.281 | 1 | 0.77 | 0.380 | 0.479 |
|  | Pop. × Trt. | 7 | 5.72 | 0.573 | 0.655 | 6 | 8.51 | 0.203 | 0.294 |
| GN1 | Population | **7** | **18.59** | **0.010** | **0.042** | **6** | **16.49** | **0.011** | **0.026** |
|  | Treatment | **1** | **91.40** | **<0.0001** | **<0.0001** | **1** | **123.77** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | 7 | 11.30 | 0.126 | 0.252 | 6 | 6.71 | 0.348 | 0.457 |
| GN2 | Population | 7 | 8.91 | 0.259 | 0.363 | 6 | 13.87 | 0.031 | 0.0566 |
|  | Treatment | **1** | **13.01** | **0.0003** | **0.0018** | **1** | **50.11** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | 7 | 9.65 | 0.210 | 0.351 | 6 | 3.22 | 0.781 | 0.841 |
| GN3 | Population | 7 | 8.56 | 0.286 | 0.387 | **6** | **18.70** | **0.005** | **0.013** |
|  | Treatment | 1 | 0.07 | 0.796 | 0.836 | **1** | **9.88** | **0.00167** | **0.005** |
|  | Pop. × Trt. | 7 | 9.30 | 0.232 | 0.351 | 6 | 1.47 | 0.962 | 0.962 |
| GN4 | Population | 7 | 16.16 | 0.024 | 0.078 | **6** | **36.29** | **<0.0001** | **<0.0001** |
|  | Treatment | **1** | **7.52** | **0.006** | **0.028** | **1** | **75.50** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | 7 | 6.04 | 0.536 | 0.643 | 6 | 3.80 | 0.704 | 0.799 |
| GN5 | Population | 7 | 12.91 | 0.074 | 0.177 | **6** | **26.58** | **0.0002** | **0.0007** |
|  | Treatment | **1** | **13.37** | **0.0003** | **0.0018** | **1** | **68.66** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | 7 | 9.32 | 0.231 | 0.351 | 6 | 2.90 | 0.821 | 0.841 |
| GN6 | Population | 7 | 12.84 | 0.076 | 0.177 | 6 | 11.76 | 0.067 | 0.113 |
|  | Treatment | **1** | **56.14** | **<0.0001** | **<0.0001** | **1** | **107.13** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | 7 | 9.00 | 0.253 | 0.363 | 6 | 7.72 | 0.260 | 0.352 |
| GN7 | Population | 7 | 15.34 | 0.032 | 0.096 | **6** | **19.99** | **0.003** | **0.008** |
|  | Treatment | **1** | **67.46** | **<0.0001** | **<0.0001** | **1** | **86.20** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | 7 | 10.67 | 0.154 | 0.281 | **6** | **14.79** | **0.022** | **0.049** |
| GN8 | Population | 7 | 15.11 | 0.035 | 0.098 | 6 | 3.69 | 0.719 | 0.799 |
|  | Treatment | 1 | 0.65 | 0.420 | 0.540 | 1 | 1.49 | 0.222 | 0.311 |
|  | Pop. × Trt. | 7 | 9.96 | 0.191 | 0.334 | 6 | 4.10 | 0.663 | 0.796 |
| GN9 | Population | 7 | 12.56 | 0.084 | 0.186 | **6** | **23.63** | **0.0006** | **0.0019** |
|  | Treatment | **1** | **26.83** | **<0.0001** | **<0.0001** | **1** | **102.74** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | 7 | 6.21 | 0.515 | 0.636 | 6 | 13.97 | 0.030 | 0.057 |
| GN10 | Population | 7 | 1.76 | 0.972 | 0.972 | 6 | 10.58 | 0.102 | 0.165 |
|  | Treatment | 1 | 0.04 | 0.840 | 0.860 | 1 | 0.13 | 0.723 | 0.799 |
|  | Pop. × Trt. | 7 | 5.46 | 0.604 | 0.655 | 6 | 14.12 | 0.028 | 0.056 |
| GN11 | Population | 7 | 13.34 | 0.064 | 0.168 | 6 | 12.34 | 0.055 | 0.096 |
|  | Treatment | **1** | **31.91** | **<0.0001** | **<0.0001** | **1** | **64.04** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | 7 | 9.27 | 0.234 | 0.351 | 6 | 6.33 | 0.388 | 0.479 |
| GN12 | Population | 7 | 7.05 | 0.424 | 0.540 | 6 | 9.38 | 0.154 | 0.231 |
|  | Treatment | 1 | 2.75 | 0.097 | 0.204 | 1 | 0.06 | 0.808 | 0.841 |
|  | Pop. × Trt. | 7 | 5.53 | 0.596 | 0.655 | 6 | 9.41 | 0.152 | 0.231 |
| GN13 | Population | 7 | 16.84 | 0.018 | 0.063 | 6 | 14.14 | 0.028 | 0.056 |
|  | Treatment | 1 | 0.26 | 0.608 | 0.655 | **1** | **30.16** | **<0.0001** | **<0.0001** |
|  | Pop. × Trt. | **7** | **17.85** | **0.013** | **0.0496** | **6** | **16.57** | **0.011** | **0.026** |

**Table 4.** Significant effects from mixed models testing associations between climate variables and plasticity measures, for traits and eigengenes on day 6 and day 12.

| Day | Plasticity measure | Climate variable | p-value | Plasticity high in pops. from: |
|---|---|---|---|---|
| Day 6 | GN5 | Elevation | 0.015 | lower elevation |
| Day 6 | Stomatal density | Mean annual precip. | 0.008 | wetter sites |
| Day 12 | GN10 | Mean annual precip. | 0.035 | wettest and driest sites |
| Day 12 | GN13 | Mean annual precip. | 0.021 | wetter sites |
| Day 12 | Growth | Elevation | 0.014 | lower elevation |
| Day 12 | Growth | Summer max. temp. | 0.016 | warmer sites |
| Day 12 | Leaf area | Mean annual precip. | 0.019 | wetter sites |
| Day 12 | Pigment accumulation | Temperature range | 0.039 | higher heterogeneity |
| Day 12 | Pigment accumulation | Summer max. temp. | 0.041 | coolest and warmest sites |