Split sex-ratio and variability in aggressive behavior in desert ants

Shani Inbar[#] and Eyal Privman

Department of Evolutionary and Environmental Biology, Institute of Evolution, University of Haifa, Haifa

3498838, Israel

[#] Corresponding author: shaniinbar@gmail.com

Abstract

Several ant species display split sex-ratio distributions, in which colonies specialize in the

production of either male or female sexuals. Here, we report split-sex ratio distributions in desert

ants of the *Cataglyphis niger* species complex. We also show that male- and female-producing

colonies display elevated aggression toward conspecifics, compared to colonies without

sexuals, and that encounters between workers from female- and male-producing colonies tend

to be the most aggressive. Moreover, elevated aggression of workers from sexual-producing

colonies overrides the previously reported 'nasty-neighbor' effect and persists even toward ants

from distant nests. These results suggest that the need to protect the substantial energetic

investment in sexuals leads to elevated aggression levels and alters aggressive behavior

patterns.

Introduction

Bimodal sex-ratio distribution

Trivers and Hare's seminal theoretical work on haplodiploid social insects (ants, bees, and

wasps) showed that relatedness asymmetries between colony members can lead to different

predicted sex ratios when queens or workers control reproduction (Trivers and Hare 1976). In

colonies with singly-mated queens, queens and workers should favor 1:1 and 3:1 ratios of

gynes to drones, respectively. Studies from singly-mated species revealed frequent female-

biased sex ratios indicating that workers often prevail (Trivers and Hare 1976, Herbers 1984,

Nonacs 1986). Conversely, the queen may thwart worker control by specializing in producing

either gynes or drones, so that workers are unable to manipulate the sex ratio by selectively

eliminating males. This phenomenon is known as split-sex ratio (Grafen 1986, Boomsma and

Grafen 1990, Boomsma 1991, Boomsma and Grafen 1991, Chan and Bourke 1994, Pamilo and

Seppä 1994, Sundström 1994, Ratnieks and Boomsma 1997, Queller and Strassmann 1998,

Chapuisat and Keller 1999, Mehdiabadi et al. 2003, Bourke 2005)

When queens are multiply mated, colonies are expected to vary in relatedness asymmetries

depending on the number of mates. Boomsma and Grafen (1990, 1991) argued that the

variation in relatedness asymmetries among colonies promotes the evolution of split sex ratios

(Boomsma and Grafen 1990, 1991). Furthermore, according to these studies, colonies

composed of highly related females are predicted to produce female sexuals, while colonies

with lower relatedness will produce males. Here, we report split sex-ratios in colonies founded

by multiply-mated queens, and show that male and female producing colonies differ in their

level of aggression.

Aggression in conspecific encounters

In *C. niger*, as in other social insect species, workers discriminate between nestmates and non-

nestmates according to their cuticular hydrocarbons profiles and non-nestmates interactions are

typically highly agonistic (Soroker et al. 1994, Soroker et al. 1995b, Soroker et al. 1995a, Lahav

et al. 1999). Aggressive behavior between non-nestmates depends on genetic relatedness

(Stuart and Herbers 2000, van Zweden and d'Ettorre 2010), environmental cues like nest

material, climate, and queen odors (Jutsum et al. 1979, Carlin and Hölldobler 1986, Crosland

1989, Chen and Nonacs 2000), diet (Silverman and Liang 2001, Buczkowski et al. 2005, Grover

et al. 2007), worker caste, task, and age (Nowbahari and Lenoir 1989, Wagner et al. 1998), past

experience (Newey et al. 2010a, Van Wilgenburg et al. 2010, Esponda and Gordon 2015), etc.

Studies in different species reported variable effects of colony size on aggression toward non-

nestmates, including both positive and negative correlation (Crosland 1990, Stuart 1991). These

findings suggest that various factors such as life history and reproductive strategy determine

whether increased aggression is beneficial for young and small colonies vs. old and large

colonies.

The cost of sexuals

The production of sexuals in ants is highly energy-consuming. Gynes and drones are typically

larger in size than the average worker. Unlike workers, both gynes and drones need to develop

flight muscles, and store sufficient energy reserves for flight. Gynes also accumulate fat and

protein in order to decrease, or altogether avoid, foraging during colony founding (Keller and

Passera 1988, 1989). Such an energetic investment by the colony is likely to be protected and

may result, we hypothesized, in elevated worker aggression between the emergence of sexuals

and their departure for mating. This hypothesis is supported by evidence indicating seasonal

variation in aggression levels between workers (Ichinose 1991, D'Ettorre et al. 2004, Brandt et

al. 2005, Katzerke et al. 2006, Thurin and Aron 2008) which is determined by both intrinsic (e.g.

worker density, relatedness within the nest) and extrinsic factors (e.g. temperature, availability

and quality of food) (Thurin and Aron 2008). Additional evidence also indicated that some ant

species have periodic variation in cuticular hydrocarbon profiles (Vander Meer et al. 1989,

Nielsen et al. 1999, Liu et al. 2001), which may alter the response toward non-nestmates. We

hypothesized that the reproductive phase of the colony is an important intrinsic factor affecting

worker response threshold to various types of threats and that the presence of sexuals in the

colony affects intercolonial acceptance threshold.

The distance between neighboring colonies

Studies suggest that ants can discern whether an ant they encounter comes from a distant nest,

or a close one, and that it affects the displayed level of aggression. In *C. niger*, workers tend to

be more aggressive toward conspecifics from close nests and exhibit lesser efforts to engage in

agonistic behavior when they encounter ants from distant nests, a phenomenon known as the

'nasty neighbor' effect (Saar et al. 2014). This phenomenon is also known in other *Cataglyphis*

species (Knaden and Wehner 2003), as well as in other ant genera (Sanada-Morimura et al.

2003, Boulay et al. 2007, Thomas et al. 2007, Wilgenburg 2007, Newey et al. 2010b, Roux et al.

2013). It should be mentioned that, in some species, the opposite effect was also reported, a

phenomenon known as the 'dear neighbor' effect (Langen et al. 2000). However, these studies

did not investigate the 'nasty/dear neighbor' effects in the context of the reproductive phase of

the colony and the presence of sexuals. Another objective of this study, therefore, was to

determine whether the 'nasty neighbor' effect persists, or is amplified, during the sexual phase

of *C. niger* colonies, when there is a greater resource investment at risk.

Methods

Sampling

Ants were sampled in April 2016 from Betzet beach on the northern Israeli coastline (from

N33.05162, E35.10245 to N33.07868, E35.10705). This population was previously described as

C. drusus (Eyer et al. 2017), but our recent species delimitation study raised the question of

whether *C. drusus* is separate species or is it the same species as *C. niger*, because these

populations are not differentiated by their nuclear genomic DNA (Reiner-Brodetzki et al.).

Colonies of this population are monogyne (headed by a single queen), polyandrous (queens are

multiply mated) and monodomous (single nest per colony) (Eyer et al. 2017). Spring is the

reproductive season in this species, and sexuals, i.e., gynes (winged, unmated queens) and

males can be found in nests in early spring. Fifty nests along a transect of 3.1km were dug (one

nest every 80m on average) and 30 workers were collected from each nest. Workers were

collected from both the entrance of the nests and from inside the nests for a heterogeneous

sample. Samples from each nest were kept in a 30x20x10cm box and given water and honey-

water. All behavioral assays were conducted within a week after collection. In the 50 colonies

sampled, we found gynes in 14 colonies and drones in 7 colonies. None of the colonies had

both gynes and males (Fig. 1). Any sexuals found were collected and frozen on the same

evening of collection so that workers were left in the boxes without sexuals.

Fig. 1: Map of 50 sampled colonies along a transect in Betzet beach (from N33.05162, E35.10245 to

N33.07868, E35.10705). Blue squared M indicate colonies where drones were found. 'Yellow ladies'

indicate colonies were gynes were found.

Behavioral assays

All test encounters were dyadic (two workers per encounter) and, in each, workers from either

female (F), male (M), or non-sexual (N) colonies were introduced to each other and the level of

aggression between them was scored (i.e., six types of encounters all together: FF, MF, MM,

NM, NF, NN). Five encounters from each pair of selected colonies were introduced and scored.

All encounters were conducted in a neutral arena (9 cm diameter, lined with a filter paper that

was changed after each test). Before each test, workers were acclimatized to the arena for 2

min by placing them on the arena surface covered in separate glass tubes. The test began by

removing the tubes, and aggressive behavior was recorded for 2 min. Aggression was scored

according to the commonly practiced method (Errard and Hefetz 1997) using the following

behavioral acts: 1 point - antennation; 2- mandibular threat; 3- short biting with jumping; 4-

biting; 5- spraying of formic acid. The aggression index was calculated as follows:

$$\frac{\sum_{i=1}^{n} A_i * t_i}{T}$$

where A_i and t_i are the aggression score and duration of each act respectively, and T is the total

interaction time (120 sec). Aggression scores were not normally distributed for some types of

encounters (Shapiro-Wilk test p-values: FF- 0.064, MF- 0.125, MM- 0.005, NF- 0.01, NM- 0.001,

NN- 0.042), and thus encounter types were compared using the non-parametric Kruskal-Wallis

test.

Results

In a sample of 50 colonies, gynes were found in 14 colonies and drones were found in 7

colonies (Fig. 1). No colonies had both gynes and drones. Interactions between workers from

female producing colonies and male producing colonies (MF) were significantly more

aggressive than any of the interactions which involved workers from colonies that did not

produce sexuals (Fig. 2; Kruskal-Wallis test *p* < 0.0001 followed by Dunn's multiple pairwise

comparisons corrected for multiple testing (Benjamini and Hochberg 1995): *p* = 0.001, 0.003,

and 0.001, for the pairwise comparisons of FM with NF, NM and NN, respectively). FF

encounters were also marginally significantly more aggressive in comparison to all type of

interactions which involved N colonies (NF, NM and NN *p* = 0.055, 0.055, 0.003, respectively).

Fig. 2: Distribution of aggression scores for interactions between colonies that produced, or did not produce,

sexuals. Aggression scores for interactions between workers from female-producing colonies (F), male-producing

colonies (M) and colonies in which no sexuals were found (N). Means are indicated by red + signs and medians are

indicated by red horizontal lines. Significant differences are marked with asterisks.

For the analysis of the effect of the distance between nests (Fig. 3A), all interactions between

workers from sexuals-producing colonies were grouped together (i.e., FF+MF+MM = S-S),

interactions between sexuals-producing colonies and colonies where no sexuals were found

were grouped together (i.e., NM+NF = S-NS), and a third group consisted of interaction only

between workers from colonies where no sexuals were found (NN = NS-NS). Significant

negative correlation was found between aggressive behavior and the distance between the

nests of the participating parties in encounters between non-sexual colonies (NS-NS; Fig. 3B;

Spearman's r=-0.39, p=0.001). This result is in line with previous reports of the 'nasty neighbor'

effect in *C. niger* (Saar et al. 2014). Encounters between sexual colonies (S-S) also showed

significant negative correlation (Fig. 3C; r=-0.261, p=0.005). However, encounters between

sexuals and non-sexuals (S-NS) did not exhibit correlation between aggression and distance

(Fig. 3D; r=0.055, p=0.491). In contrast to non-sexual colonies, encounters between sexual

colonies persisted in high aggression scores for longer distances, that is, ants remain

aggressive even when they encountered ants from nests up to 2km away. Similarly, high

aggression was also observed in S-NS interactions at large distances.

Fig. 3: Aggression scores according to distance in three types of interactions (sexuals and sexuals,

167 non-sexuals and sexuals, non-sexuals and non-sexuals). (A) Scatter plot of observed aggression scores,

168 up to 400m between participating nests. (B) Aggression score by distance intervals in NS-NS interactions

169 (C) Aggression score by distance intervals in S-S interactions. (D) Aggression score by distance intervals

170 in NS-S interactions. Red + marks indicate mean scores.

Discussion

We report a survey of a *Cataglyphis* population during the reproductive season. Our results

reveal split-sex ratios in this population. Behavioral assays show that the presence of sexuals in

colonies affects both magnitude and patterns of aggressive behavior toward non-nestmates.

Workers from female and male producing colonies are expected to protect their energy

investment in sexuals, rich with lipid and protein (Keller and Passera 1988, 1989). Our results

show that, indeed, the presence of sexuals leads to elevated levels of aggression. Furthermore,

it appears that aggression levels are highest when workers from female producing colonies

interact with workers form male producing colonies. This may be a result of differences in CHCs

between drones and gynes (Cuvillier-Hot et al. 2001, Smith et al. 2014), which may be

incorporated in the CHC profiles of the workers, thus making the identification of a non-

nestmate quicker and eliciting a greater aggressive response. Another explanation for increased

aggression levels between workers from female and male producing colonies may be

differences in intra-colony relatedness. Crozier suggested that individuals with rare labels

should be rejected more often than individuals with common labels (Crozier 1986, Crozier 1987,

1988, Elgar and Crozier 1989) and that individuals from high-diversity colonies should be

rejected more frequently than individuals from low-diversity colonies. And indeed, there are

reports that colonies with greater genetic relatedness tend to be more aggressive toward

colonies of lesser genetic relatedness (Tsutsui et al. 2003). Since colonies with greater genetic

relatedness may also tend to produce gynes (Boomsma and Grafen 1990, 1991, Sundström

1994, Evans 1995, Bourke 2015) this may explain the elevated levels of aggression we

observed in FM interactions.

It is also expected that, in the presence of sexuals, workers would invest greater efforts in

protecting their energetic investment in them and exhibit greater aggression response toward

intruders in general. In this study, in addition to estimating the magnitude of aggressive

responses, we, also, examined whether the presence of sexuals alters the pattern of aggressive

behavior. Our results show that, while colonies that did not produce sexuals exhibit a clear

'nasty neighbour' effect, as previously reported (Saar et al. 2014), colonies with gynes or males

persist in aggressive behavior even when they encounter ants from distant nests. Although S-S

interactions do have a significant negative correlation between distance and aggressiveness,

aggression levels decrease only at ~500m from the nest. S-NS interaction, on the other hand,

do not exhibit any correlation between distance and aggressiveness and ants persist in

aggressive behavior for even longer distances. This result may suggest, again, that risk

assessment in the presence of sexuals play a role in deciding to invest in aggressive behavior.

Acknowledgement

E.P. was supported by Israel Science Foundation Grants no. 646/15, 2140/15, and 2155/15 and

US-Israel Binational Science Foundation Grant no. 2013408.

References

Benjamini, Y., and Y.J.J.o.t.r.s.s.S.B. Hochberg (1995) Controlling the false discovery rate: a
213   practical and powerful approach to multiple testing. 289-300.
214   **Boomsma, J.** (1991) Adaptive colony sex ratios in primitively eusocial bees. *Trends in Ecology*
215   *& Evolution* 6, 92-95.
216   **Boomsma, J., and A. Grafen** (1990) Intraspecific variation in ant sex ratios and the Trivers-
217   Hare hypothesis. *Evolution* 44, 1026-1034.
218   **Boomsma, J., and A. Grafen** (1991) Colony-level sex ratio selection in the eusocial
219   Hymenoptera. *Journal of evolutionary Biology* 4, 383-407.
220   **Boulay, R., X. Cerdá, T. Simon, M. Roldan, and A. Hefetz** (2007) Intraspecific competition in
221   the ant Camponotus cruentatus: should we expect the 'dear enemy'effect? *Animal*
222   *Behaviour* 74, 985-993.
223   **Bourke, A.F.** (Year) Genetics, relatedness and social behaviour in insect societies. p. 1 in
224   Proceedings of the SYMPOSIUM-ROYAL ENTOMOLOGICAL SOCIETY OF LONDON,
225   2005.
226   **Bourke, A.F.** (2015) Sex investment ratios in eusocial Hymenoptera support inclusive fitness
227   theory. *Journal of evolutionary Biology* 28, 2106-2111.
228   **Brandt, M., J. Heinze, T. Schmitt, and S. Foitzik** (2005) A chemical level in the coevolutionary
229   arms race between an ant social parasite and its hosts. *Journal of evolutionary Biology*
230   18, 576-586.
231   **Buczkowski, G., R. Kumar, S.L. Suib, and J. Silverman** (2005) Diet-related modification of
232   cuticular hydrocarbon profiles of the Argentine ant, Linepithema humile, diminishes
233   intercolony aggression. *Journal of Chemical Ecology* 31, 829-843.
234   **Carlin, N.F., and B. Hölldobler** (1986) The kin recognition system of carpenter ants
235   (Camponotus spp.). *Behavioral Ecology and Sociobiology* 19, 123-134.
236   **Chan, G.L., and A.F. Bourke** (1994) Split sex ratios in a multiple-queen ant population. *Proc.*
237   *R. Soc. Lond. B* 258, 261-266.
238   **Chapuisat, M., and L. Keller** (1999) Testing kin selection with sex allocation data in eusocial
239   Hymenoptera. *Heredity* 82, 473.
240   **Chen, J.S., and P. Nonacs** (2000) Nestmate recognition and intraspecific aggression based on
241   environmental cues in Argentine ants (Hymenoptera: Formicidae). *Annals of the*
242   *Entomological Society of America* 93, 1333-1337.

243   **Crosland, M.W.** (1989) Kin recognition in the ant Rhytidoponera confusa I. Environmental
244   odour. *Animal Behaviour* 37, 912-919.
245   **Crosland, M.W.** (1990) The influence of the queen, colony size and worker ovarian
246   development on nestmate recognition in the ant Rhytidoponera confusa. *Animal*
247   *Behaviour* 39, 413-425.
248   **Crozier, R.** (1986) Genetic clonal recognition abilities in marine invertebrates must be
249   maintained by selection for something else. *Evolution* 40, 1100-1101.
250   Crozier, R.H. (1987) Genetic aspects of kin recognition: concepts, models, and synthesis pp.
251   55–73 in D. J. C. Fletcher and C. D. Michener. (Eds.) Kin Recognition in Animals.
252   Chichester, Wiley.
253   Crozier, R.H. (1988) Kin Recognition Using Innate Labels: A Central Role for Piggybacking? in
254   R. K. Grosberg, D. Hedgecock and K. Nelson. (Eds.) Invertebrate Historecognition.
255   Boston, MA, Springer.
256   **Cuvillier-Hot, V., M. Cobb, C. Malosse, and C. Peeters** (2001) Sex, age and ovarian activity
257   affect cuticular hydrocarbons in Diacamma ceylonense, a queenless ant. *Journal of*
258   *Insect Physiology* 47, 485-493.
259   **D'Ettorre, P., E. Brunner, T. Wenseleers, and J. Heinze** (2004) Knowing your enemies:
260   seasonal dynamics of host–social parasite recognition. *Naturwissenschaften* 91, 594-
261   597.
262   **Elgar, M.A., and R.H. Crozier** (1989) Animal allorecognition systems: how to get to know
263   yourself. *Trends in Ecology & Evolution* 4, 288-289.
264   **Errard, C., and A. Hefetz** (1997) Label familiarity and discriminatory ability of ants reared in
265   mixed groups. *Insectes sociaux* 44, 189-198.
266   **Esponda, F., and D.M. Gordon** (2015) Distributed nestmate recognition in ants. *Proc. R. Soc.*
267   *B* 282, 20142838.
268   **Evans, J.D.** (1995) Relatedness threshold for the production of female sexuals in colonies of a
269   polygynous ant, Myrmica tahoensis, as revealed by microsatellite DNA analysis.
270   *Proceedings of the National Academy of Sciences* 92, 6514-6517.
271   **Eyer, P., R. Seltzer, T. Reiner-Brodetzki, and A. Hefetz** (2017) An integrative approach to
272   untangling species delimitation in the Cataglyphis bicolor desert ant complex in Israel.
273   *Molecular phylogenetics and evolution* 115, 128-139.
274   **Grafen, A.** (1986) Split sex ratios and the evolutionary origins of eusociality. *Journal of*
275   *theoretical biology* 122, 95-121.
276   **Grover, C.D., A.D. Kay, J.A. Monson, T.C. Marsh, and D.A. Holway** (2007) Linking nutrition
277   and behavioural dominance: carbohydrate scarcity limits aggression and activity in
278   Argentine ants. *Proceedings of the Royal Society of London B: Biological Sciences* 274,
279   2951-2957.
280   **Herbers, J.M.** (1984) Queen-worker conflict and eusocial evolution in a polygynous ant species.
281   *Evolution* 38, 631-643.
282   **Ichinose, K.** (1991) Seasonal variation in nestmate recognition in Paratrechina flavipes (Smith)
283   worker ants (Hymenoptera: Formicidae). *Animal Behaviour* 41, 1-6.
284   **Jutsum, A., T. Saunders, and J. Cherrett** (1979) Intraspecific aggression is the leaf-cutting ant
285   Acromyrmex octospinosus. *Animal Behaviour* 27, 839-844.
286   **Katzerke, A., P. Neumann, C.W. Pirk, P. Bliss, and R.F. Moritz** (2006) Seasonal nestmate
287   recognition in the ant Formica exsecta. *Behavioral Ecology and Sociobiology* 61, 143-
288   150.
289   **Keller, L., and L. Passera** (1988) Energy investment in gynes of the Argentine ant Iridomyrmex
290   humilis (Mayr) in relation to the mode of colony founding in ants (Hymenoptera:
291   Formicidae). *International Journal of Invertebrate Reproduction and Development* 13,
292   31-38.

Keller, L., and L. Passera (1989) Size and fat content of gynes in relation to the mode of
colony founding in ants (Hymenoptera; Formicidae). *Oecologia* 80, 236-240.
Knaden, M., and R. Wehner (2003) Nest defense and conspecific enemy recognition in the
desert ant Cataglyphis fortis. *Journal of Insect Behavior* 16, 717-730.
Lahav, S., V. Soroker, A. Hefetz, and R.K. Vander Meer (1999) Direct behavioral evidence for
hydrocarbons as ant recognition discriminators. *Naturwissenschaften* 86, 246-249.
Langen, T.A., F. Tripet, and P. Nonacs (2000) The red and the black: habituation and the
dear-enemy phenomenon in two desert Pheidole ants. *Behavioral Ecology and*
Sociobiology 48, 285-292.
Liu, Z., A. Bagnères, S. Yamane, Q. Wang, and J. Kojima (2001) Intra-colony, inter-colony
and seasonal variations of cuticular hydrocarbon profiles in Formica japonica
(Hymenoptera, Formicidae). *Insectes sociaux* 48, 342-346.
Mehdiabadi, N.J., H.K. Reeve, and U.G. Mueller (2003) Queens versus workers: sex-ratio
conflict in eusocial Hymenoptera. *Trends in Ecology & Evolution* 18, 88-93.
Newey, P.S., S.K. Robson, and R.H. Crozier (2010a) Know thine enemy: why some weaver
ants do but others do not. *Behavioral Ecology* 21, 381-386.
Newey, P.S., S.K. Robson, and R.H. Crozier (2010b) Weaver ants Oecophylla smaragdina
encounter nasty neighbors rather than dear enemies. *Ecology* 91, 2366-2372.
Nielsen, J., J.J. Boomsma, N. Oldham, H. Petersen, and E. Morgan (1999) Colony-level and
season-specific variation in cuticular hydrocarbon profiles of individual workers in the ant
Formica truncorum. *Insectes sociaux* 46, 58-65.
Nonacs, P. (1986) Sex-ratio determination within colonies of ants. *Evolution* 40, 199-204.
Nowbahari, E., and A. Lenoir (1989) Age-related changes in aggression in ant Cataglyphis
cursor (Hymenoptera, Formicidae): influence on intercolonial relationships. *Behavioural*
processes 18, 173-181.
Pamilo, P., and P. Seppä (1994) Reproductive competition and conflicts in colonies of the ant
Formica sanguinea. *Animal Behaviour* 48, 1201-1206.
Queller, D.C., and J.E. Strassmann (1998) Kin selection and social insects. *Bioscience* 48,
165-175.
Ratnieks, F.L., and J.J. Boomsma (1997) On the robustness of split sex ratio predictions in
social Hymenoptera. *Journal of theoretical biology* 185, 423-439.
Reiner-Brodetzki, T., S. Inbar, P. Cohen, A. Hefetz, and E. Privman Social polymorphism in
Cataglyphis niger. *In preperation,*
Roux, O., V. Rossi, R. Céréghino, A. Compin, J.-M. Martin, and A. Dejean (2013) How to
coexist with fire ants: the roles of behaviour and cuticular compounds. *Behavioural*
processes 98, 51-57.
Saar, M., L. Leniaud, S. Aron, and A. Hefetz (2014) At the brink of supercoloniality: genetic,
behavioral, and chemical assessments of population structure of the desert ant
Cataglyphis niger. *Frontiers in ecology and evolution* 2, 13.
Sanada-Morimura, S., M. Minai, M. Yokoyama, T. Hirota, T. Satoh, and Y. Obara (2003)
Encounter-induced hostility to neighbors in the ant Pristomyrmex pungens. *Behavioral*
Ecology 14, 713-718.
Silverman, J., and D. Liang (2001) Colony disassociation following diet partitioning in a
unicolonial ant. *Naturwissenschaften* 88, 73-77.
Smith, A.A., W. Vanderpool, J.G. Millar, L.M. Hanks, and A.V. Suarez (2014) Conserved
male-specific cuticular hydrocarbon patterns in the trap-jaw ant Odontomachus
brunneus. *Chemoecology* 24, 29-34.
Soroker, V., C. Vienne, and A. Hefetz (1995a) Hydrocarbon dynamics within and between
nestmates inCataglyphis niger (Hymenoptera: Formicidae). *Journal of Chemical Ecology*
21, 365-378.

Soroker, V., C. Vienne, A. Hefetz, and E. Nowbahari (1994) The postpharyngeal gland as a
"Gestalt" organ for nestmate recognition in the antCataglyphis niger.
Naturwissenschaften 81, 510-513.
Soroker, V., A. Hefetz, M. Cojocaru, J. Billen, S. Franke, and W. Francke (1995b) Structural
and chemical ontogeny of the postpharyngeal gland in the desert ant Cataglyphis niger.
Physiological Entomology 20, 323-329.
Stuart, R.J. (1991) Nestmate recognition in leptothoracine ants: testing for effects of queen
number, colony size and species of intruder. *Animal Behaviour* 42, 277-284.
Stuart, R.J., and J.M. Herbers (2000) Nest mate recognition in ants with complex colonies:
within-and between-population variation. *Behavioral Ecology* 11, 676-685.
Sundström, L. (1994) Sex ratio bias, relatedness asymmetry and queen mating frequency in
ants. *Nature* 367, 266.
Thomas, M., C. Payne-Makrisa, A. Suarez, N. Tsutsui, and D. Holway (2007) Contact
between supercolonies elevates aggression in Argentine ants. *Insectes sociaux* 54, 225-
233.
Thurin, N., and S. Aron (2008) Seasonal nestmate recognition in the polydomous ant
Plagiolepis pygmaea. *Animal Behaviour* 75, 1023-1030.
Trivers, R.L., and H. Hare (1976) Haploidploidy and the evolution of the social insect. *Science*
191, 249-263.
Tsutsui, N.D., A.V. Suarez, and R.K. Grosberg (2003) Genetic diversity, asymmetrical
aggression, and recognition in a widespread invasive species. *Proceedings of the*
National Academy of Sciences 100, 1078-1083.
Van Wilgenburg, E., J. Clémencet, and N.D. Tsutsui (2010) Experience influences
aggressive behaviour in the Argentine ant. *Biology Letters* 6, 152-155.
van Zweden, J.S., and P. d'Ettorre (2010) Nestmate recognition in social insects and the role
of hydrocarbons. *Insect hydrocarbons: biology, biochemistry and chemical ecology* 11,
222-243.
Vander Meer, R.K., D. Saliwanchik, and B. Lavine (1989) Temporal changes in colony
cuticular hydrocarbon patterns ofSolenopsis invicta. *Journal of Chemical Ecology* 15,
2115-2125.
Wagner, D., M.J. Brown, P. Broun, W. Cuevas, L.E. Moses, D.L. Chao, and D.M. Gordon
(1998) Task-related differences in the cuticular hydrocarbon composition of harvester
ants, Pogonomyrmex barbatus. *Journal of Chemical Ecology* 24, 2021-2037.
Wilgenburg, E.v. (2007) The influence of relatedness, neighbourhood and overall distance on
colony mate recognition in a polydomous ant. *Ethology* 113, 1185-1191.

378