An agent-based framework for improving wildlife disease surveillance: A case

study of chronic wasting disease in Missouri white-tailed deer.

Aniruddha V. Belsare[1#*], Matthew E. Gompper[1], Barbara Keller[2], Jason Sumners[2],

Lonnie Hansen[1], and Joshua J. Millspaugh[3]

[1] School of Natural Resources, University of Missouri, Columbia, Missouri, United

States of America

[2] Missouri Department of Conservation, 3500 East Gans Road, Columbia, Missouri,

United States of America

[3] W.A. Franke College of Forestry and Conservation, University of Montana, Missoula,

Montana, United States of America

[#] Current Affiliation: Center for Modeling Complex Interactions, University of Idaho,

Moscow, Idaho, United States of America

*Corresponding author

Email: abelsare@uidaho.edu

Abstract

Epidemiological surveillance for important wildlife diseases often relies on samples obtained from hunter-harvested animals. A problem, however, is that although convenient and cost-effective, hunter-harvest samples are not representative of the population due to heterogeneities in disease distribution and biased sampling. We developed an agent-based modeling framework that simulates spatial and demographic heterogeneities in disease distribution as well as hunter harvest sampling, thereby facilitating population-specific sample size calculations for prompt detection of disease. Application of this framework is illustrated using the example of chronic wasting disease (CWD) surveillance in Missouri's white-tailed deer (*Odocoileus virginianus*) population. We show how confidence in detecting CWD is grossly overestimated under the unrealistic, but standard, assumptions that sampling effort and disease are randomly and independently distributed. We then provide adjusted sample size recommendations based on more realistic assumptions. Wildlife agencies can use these open-access models to design their CWD surveillance. Furthermore, these models can be readily adapted to other regions and other wildlife disease systems.

Keywords: agent-based models; chronic wasting disease; sample size; surveillance; white-tailed deer; wildlife diseases.

1. Introduction

Surveillance of wildlife populations is an essential tool for effective management of diseases that pose conservation, economic or public health threats. Samples for wildlife disease surveillance are frequently obtained from hunter-harvested animals (e.g. [1–4]). This approach is practical and cost-effective as samples are obtained from a large geographic area and sampling costs are dramatically reduced. In recent decades, wildlife agencies across USA and Canada have implemented harvest-based chronic wasting disease (CWD) surveillance in cervid populations [5,6]. CWD is an emerging prion disease of cervids, including white-tailed deer (*Odocoileus virginianus*), mule deer (*O. hemionus*) and elk (*Cervus elaphus*). Currently reported from 26 states and provinces in North America, the continuing spread of CWD is a serious long-term threat to the health of free-ranging cervid populations [7–10].

For emerging wildlife diseases like CWD, surveillance programs are aimed at prompt detection in populations where the disease has not been previously documented. Epidemiological data generated from sampling the population of interest is used to infer disease status of the population (presence/absence, prevalence, distribution). Reliable inferences can be made only if sufficient number of representative samples are obtained. Sample sizes for epidemiological surveillance are often determined from standardized tables (or online programs) that use hypergeometric distribution (sampling without replacement) for finite populations or a binomial distribution (sampling with replacement) for large popultions (>100000), under the assumption that samples are representative of the population of interest [5,11,12].

However, hunter-harvested samples will rarely be representative of the population and

may result in biased estimates and erroneous inferences about disease presence and

distribution [6,13]. Biases in hunter-harvested samples can arise from disease

distribution heterogeneities in the population (infected individuals are often clustered

rather than being uniformly or randomly distributed through a population) and factors

like harvest regulations, hunter selectivity, variable land access, and uneven hunter

distribution. Wildlife agencies need to consider such biases and adapt their disease

surveillance strategies accordingly. This issue is particularly important when the disease

prevalence rates are very low and yet the potential for impact on wildlife, domestic

animals, or public health is high.

The white-tailed deer is one of Missouri's most valuable natural resources, and the

continuing spread of CWD poses a serious threat to the health of Missouri's free-

ranging deer populations. Logistical and financial constraints limit the ability of mangers

to implement strict probability-based sampling schemes, hence CWD surveillance

continues to rely on hunter-harvested samples. Statewide surveillance of hunter-

harvested deer for CWD was initiated by the Missouri Department of Conservation in

2002 [14], but the first case of CWD in free-ranging deer was detected in 2012. As of

December 2017, more than 76,600 free-ranging deer have been tested and 43 found

positive for CWD in Missouri (0.06% of sampled deer) [15–17]. In 2016-17 alone, the

state tested >25,600 deer and detected nine positive cases in five counties (0.04% of

sampled deer). These numbers illustrate both the daunting challenge involved with

surveillance, and the level of concern over the disease in the state.

CWD occurs at a low prevalence in most areas, and has a clustered distribution with

focal areas of infection near the point of introduction [18–20]. Three considerations are

therefore vital in the context of CWD surveillance: (1) the very low prevalence of

disease when detection matters most, (2) the non-random distribution of diseased

individuals, and (3) the non-random nature of samples derived from hunter harvested

animals. Agent-based models (ABMs, also called Individual-based models) are well-

suited to solving these problems. ABMs are formulated using relevant information about

individuals, their behavior and the landscape they occupy, and therefore can readily

incorporate non-random and heterogeneous aspects of the population, the disease, and

harvest. Moreover, ABMs have been used to inform policy in many fields including

natural resource management, epidemiology and public health.

We have developed an agent-based modeling framework that incorporates real-world

heterogeneities arising due to disease distribution on the landscape and sampling

biases in hunter-harvest, and provides a rigorous, quantitative framework for informing

disease surveillance strategies. Our ABM framework has two constituent models: (1)

MO*Ov*POP (MissOuri *Odocoileus virginianus* POPulation simulation) simulates a

realistic deer population for the region of interest and (2) MO*Ov*POP*surveillance*

distributes the disease in the population in a realistic manner and then samples the

population for disease through hunter-harvest. Combining these two models allows

managers to iteratively estimate population-specific sample size for detection of disease

with a high-level of confidence. Here, we illustrate the use of this framework to derive

optimum sample sizes for efficient and reliable CWD surveillance in Missouri's white-

tailed deer populations. Though we focus on CWD in white-tailed deer populations of

Missouri, this model-based framework can be readily adapted to other regions, and

other host-pathogen systems.

2. Background information and modeling considerations

The purpose of the first model, MO*Ov*POP, is to generate a snapshot of a target deer

population that is as realistic as possible. In the model, deer are treated as individual

agents that occupy habitat patches; they interact with each other and the environment.

The habitat patches are based on forest cover data for the area of interest. In the

Midwest North America, deer thrive well in landscapes where forest cover and

agricultural food are juxtaposed, but less so in heavily forested or predominantly

agricultural areas [21,22]. Therefore, contiguous patches in the model landscape with

average forest cover between 25% and 75% are designated as deer habitat.

The model is initialized using demographic attributes derived from field data,

specifically, density, sex-ratio, age composition, reproduction, mortality (both hunting

and non-hunting). Social organization, group dynamics and dispersal are explicitly

incorporated in the model as these processes influence interactions among individual

deer and underpin the heterogeneity in the population. These heterogeneities should be

accounted for when designing surveillance for a disease like CWD.

White-tailed deer are social animals exhibiting an intermediate level of sociality, typically

occurring in small, relatively stable groups of adult females and their recent offspring

(doe social groups), loose bachelor groups of adult males, or as solitary individuals

(male and female) [23,24]. Doe social group size ranges between 2 and 12 [25,26], but

smaller group sizes (less than 8) are commonly seen in Missouri, and bachelor group

size typically ranges between 2 and 5 [27–29], but larger groups of up to 8 are

occasionally seen in Missouri (L. Hansen, personal observation). Our modeled deer

population maintains the group sizes in this suggested range throughout the simulation.

Furthermore, the pattern and strength of social affiliations in white-tailed deer

populations fluctuate temporally. For instance, pregnant females seek isolation during

the fawning season and become aggressive towards other deer including group

members [30,31]. Similarly, bachelor groups break up and males are solitary during the

breeding season [32]. Pre-breeding interactions (sparring matches, flehmen behavior or

tending behavior) and mating also influence the pattern of interactions between deer.

We have accounted for such temporal fluctuations that influence group dynamics in this

model.

Dispersal is an important component of white-tailed deer ecology, and can create

opportunities for the spread of pathogens like CWD [33].Though dispersal behavior is

predominantly exhibited by yearling bucks [34,35], female dispersal also occurs, albeit

at lower rates [36]. Yearling buck dispersal can occur during the parturition period, cued

by the intersexual aggression of pregnant females, or during the rut period, cued by the

intrasexual aggression by adult males [37]. Agonistic behavior by pregnant does just

before parturition is thought to be the reason for juvenile female dispersal [38]. Every

year of the model run, yearling buck and juvenile female dispersal is scheduled before

parturition in the fifth month. Additionally, yearling buck dispersal also occurs before

rutting activity in the 11^{th} month every year. Dispersal rates and distances for yearling

bucks and juvenile females are extrapolated from published literature (details provided

in Parameterization and Calibration section). Similarly, parameters influencing deer

reproduction and harvest have been extrapolated from the multi-year harvest data

collected by MDC.

We simulate the population forward in time until it achieves an equilibrium where the

age and sex composition and population growth are all stable.

The second model, MO*OvPOPsurveillance*, imports the heterogeneous, model-

generated deer population and simulates disease distribution and sampling. The

assumptions underpinning non-random harvest and sampling processes in this model

are derived from the hunter and landowner survey data collected by MDC. Prompt

detection in populations where CWD has been recently introduced is vital for controlling

its spread. Low prevalence scenarios are therefore particularly important in the context

of CWD surveillance. MO*OvPOPsurveillance* can simulate very low prevalence (0.1%)

as well as age-sex specific prevalence scenarios in the model deer population, thus

facilitating sample size calculations in the real-world context. This allows us to address

the central goal of determining the sample strategy for detecting CWD with high

confidence.

3. Model description

3.1 Purpose

The purpose of the model-based framework is to provide a decision-making context for

collection and analysis of harvest based epidemiological surveillance data. The

framework is adapted for chronic wasting disease (CWD) surveillance of Missouri white-

tailed deer populations.

3.2 State variables and scales

Entities: MO*Ov*POP and MO*Ov*POP*surveillance* have two entities: patches and deer.

Irrespective of the region selected for simulation, each patch in the model landscape

represents one square mile. Deer are modeled as individuals occupying the patches.

State variables: Patches located on the edges of the model landscape are designated

as border patches (*border*). Each patch is characterized by its percent forest cover

(*forest-percent*) and the mean forest-percent (*dfp*). The mean forest-percent is required

in the male yearling dispersal submodel and is calculated by averaging forest cover of a

patch and its immediate neighbors. During the setup, patches are assessed for deer

occupancy (*do*) based on percent forest cover. Patch variable *dh* (deer habitat) is used

for grouping of contiguous deer occupancy patches, thus facilitating initial distribution of

deer in the model landscape.

Each patch in MO*Ov*POP*surveillance* has two additional patch variables: *add* (average

deer density) takes the value of average number of deer occurring on the patch and its

immediate neighbors (8 patches), and *hp* designates the relative harvest pressure for a

patch.

Each deer has eight state variables, which define individual characteristics that include

age, sex, group membership and status. Deer in MO*Ov*POP*surveillance* have three

additional state variables that designate CWD infection status (*cwd*), harvest status

(*marked-for-harvest*?) and sampling status (*marked-for-testing?*).

Patch and deer state variables are listed in Table 1.

Spatial scales: MO*Ov*POP has a spatially-explicit landscape that can be set up for

individual counties or CWD management zones using geographic information (forest

cover data) for the selected region. We have used miles rather than kilometers as the

measurement unit because of the past and current norms of the region and its

management agencies, and the related need to make the results immediately applicable

to those same agencies.

Temporal scale: MO*Ov*POP has a monthly time step, and the deer population is

simulated for a period of 25 years. MO*Ov*POP*surveillance* simulates harvest and

sampling using a pre-harvest snapshot of MOOvPOP-generated deer population. The

model therefore runs for just one time-step (one month).

3.3 Process overview and scheduling

MO*Ov*POP and MO*Ov*POP*surveillance* were both developed in NetLogo 6.0, a

software platform for implementing agent-based models [39]. Model description is

provided following the ODD (Overview, Design concepts, Details) protocol for individual-

based models [40,41]. Model codes are available via website repository "Open ABM

CoMSES Computational Model Library" [42,43].

Processes: Processes included in this model are all related to deer: individual growth

(aging), male and female yearling dispersal, bachelor group formation, doe social group

formation, fawning, hunting and non-hunting mortality.

MO*Ov*POP*surveillance* implements three processes: individual growth (deer age by one

month), non-hunting mortality and hunting mortality with CWD testing.

Schedule: Individual growth (increase in age by one month) is scheduled at the

beginning of each time step and is followed by non-hunting mortality. We classify deer

in four age-classes: young fawns (up to 6 months old), older fawns (7 to 12 months old),

yearlings (13 to 24 months old) and adults (25 months or older). Processes of dispersal

(male yearling/female yearling), group formation (bachelor group/ doe social group), and

hunting mortality which only occur during a specific time-step (month) are executed after

deer-growth and non-hunting mortality. Fawning is scheduled in the fifth month every

year and is preceded by yearling buck and juvenile female dispersal. Yearling buck

dispersal is also scheduled in in the 11th month every year. Census and plotting graphs

(observer actions) are scheduled at the end of each time step. The month counter

resets after every 12 months.

In MO*Ov*POP*surveillance*, a proportion of harvested deer from each age-sex class

(specified by percent harvest tested) is queried for CWD+ status, and results are

returned without error. The order of processes occurring in MO*Ov*POP and

MO*Ov*POP*surveillance* is illustrated in Figure 1.

3.4 Design concepts

Emergence: Age and sex structure of the model deer population, as well as the deer

distribution pattern, emerge from the model.

Adaptation: Adaptation is modeled in MO*Ov*POP in two processes: male dispersal and

female group dynamics. Yearling buck dispersal distances are derived from percent

forest cover of the individual's home range [44,45]. Female group size dynamics (group-

fission or new group formation) are based on the current group size and membership.

Sensing: Agents (deer) are modeled to 'sense' their environment (patch variables or

state variables of other agents) before making some behavioral decisions. Yearling

bucks perceive the percent forest cover of their home range and determine dispersal

distance (described in Submodels: yearling male dispersal). Doe social group members

can sense current group size, group leaders can sense solitary female deer on their

patch, and solitary female deer can sense the number of doe social groups and group

leaders in their neighborhood (own patch and eight neighboring patches, nine square

mile area).

Interaction: Deer group dynamics and sociality is included in this model to implicitly

simulate within- and between- group interactions in the deer population.

Stochasticity: Each individual deer in the model is subject to a mortality probability

during a time step. The mortality probability is derived from the mortality rate (natural or

hunting). Male and female group leaders are selected from a set of potential candidates

during the setup, or when leaders have to be replaced due to mortality events. In

MO*OvPOPsurveillance*, CWD+ status is assigned randomly during the set up.

Observation: MO*Ov*POP has a graphical display of deer distribution in the landscape.

Deer abundance and distribution are updated as the model executes. Additionally, three

graphical displays are included: one plots deer abundance versus time, and the other

two plot frequency distribution of bachelor and doe social group sizes respectively.

Monitors display number of deer in each age-sex-class. Pre-harvest and post-harvest

population abundance by sex-age-class is recorded in an output file for each year of

model run.

MO*Ov*POP*surveillance* interface has monitors for the following outputs: total number of

adult deer (male and female) in the population, number of CWD+ deer in the population,

number of CWD+ deer in the harvest, total sample size as well as age-sex group wise

sample distribution, and CWD+ deer in the sample. For scenarios with clustering of

CWD, the graphical display highlights area where CWD cases are clustered in the

model landscape. Additionally, following information is recorded in an output file for

each year of the model run: number of fawns, yearlings and adults harvested

(separately for both sexes), number of CWD+ deer in the hunter harvest, number of

CWD+ deer detected in the sample, CWD detection probability and observed (sample)

prevalence.

3.5 Initialization

MO*Ov*POP is initialized with forest cover data and post-harvest deer density for the

desired sampling region. We have converted forest cover data (United States

Geological Survey 1992 National Land Cover Data) for selected sampling regions in

Missouri to a forest percentage grid of 1 square mile patches to facilitate import in

NetLogo. MO*Ov*POP*surveillance* is initialized by importing model-generated pre-harvest

deer population data (abundance, sex-age composition and distribution in the

landscape) for the sampling region under consideration.

3.6 Input data

The only model input is forest cover data for the selected sampling region.

3.7 Submodels

3.7.1 Individual growth

This submodel is executed at the beginning of every time step. All deer in the model

landscape update their state variable *aim* (age in months) by one month.

3.7.2 Deer census

Post-harvest census is scheduled in the 1st month (one time step after annual harvest),

and pre-harvest census in the 11th month (one time step before the annual harvest) of

each year. Number of male and female deer in each of the three age categories is

reported separately.

3.7.3 Deer group dynamics and sociality

We include two processes, *bachelor-group-formation* and *doe-group-size-regulator*,

which facilitate simulation of group dynamics as well as within- and between-group

contact patterns in the model deer populations.

Bachelor-group-formation

Adult male deer are solitary during breeding season, but otherwise form temporary

bachelor groups of nonrelated individuals [32]. During the first month every year

immediately after post-harvest census, potential number of bachelor groups in the deer

population is calculated based on total number of adult and yearling males in the

population and the parameter *mean-bachelor-group-size*.

$$Number\ of\ bachelor\ groups = \frac{yearling\ males + adult\ males}{mean-bachelor-group-size}$$

Surviving bachelor group leaders from the previous year maintain their leader status. If

the potential number of bachelor groups exceed the available number of group leaders,

appropriate number of bucks older than 32 months are randomly selected and

designated as potential bachelor group leaders. The leaders then form bachelor groups

by first setting their potential group size (using the parameter *mean-bachelor-group-*

size), and then recruiting available adult bucks from patches within a 1.5 mile radius;

surviving group members from the previous year are recruited before new members.

doe-group-size-regulator

After the fawning season (month = 5), doe social groups regulate their group size using

the parameter *doe-social-group-size-regulator*. Groups with membership approaching or

exceeding the value set by *doe-social-group-size-regulator* undergo fission. Up to two

female group members (adults or yearling) lose their group affiliation and become

solitary along with their fawns. A deer is considered a member of a doe social group

when its state variable *groid* has the group leader's ID number, and the other state

variable *gr* has a value of -1. After fawning (month = 5), designated leaders of doe

social groups with four or less members increase their group size by seeking solitary

females in a 1.5 mile radius and adding up to two females along with their new-born

fawns to the group. After any birth, death or dispersal event, group size is updated using

the process *review-group-dynamics*.

3.7.4 Yearling dispersal

The proportion of dispersing yearlings is set using two parameters: *yearling-male-*

dispersal-rate and *yearling-female-dispersal-rate*. Dispersing individuals travel the

calculated dispersal distance (described below) as an equivalent number of patches in a

random direction. We assume that the number of individuals dispersing out of the model

landscape is equal to the number of individuals dispersing into the model landscape.

Therefore, at any point during dispersal, if a deer moves past the edge of the model

landscape (world wraps horizontally as well as vertically), it reappears on the opposite

edge as a different deer (its state variable *momid* is changed to 0).

Yearling male dispersal

Dispersal distances for yearling bucks are modeled using percent forest cover, as

suggested by Diefenbach et al. [44]. Average forest cover for a patch and its immediate

neighbors is first calculated for each non-border patch and the value is stored as a

patch variable (*dfp*). Border patches set their *dfp* equal to that of one of their non-border

neighboring patches. Mean dispersal distance is predicted using the equation from [45]

333
$$\bar{x} = 35.07 - 48.14 \, dfp$$

334   where *dfp* is the patch variable representing mean percent forest of the patch and its

335   neighbors. Variance of dispersal distance is predicted using the equation from [44]

336
$$\log_e(s^2) = a + b\bar{x}$$

where $a = 3.51$ (SE = 0.597) and $b = 0.77$ (SE = 0.025). Dispersal distance is obtained

from a log-normal distribution using the predicted mean dispersal distance and

predicted variance of dispersal distance [44]. If a male yearling reaches a non-deer

occupancy patch after dispersal, it is transferred to the nearest deer occupancy patch.

After dispersal, the dispersing individual's state variable *mgroid* takes a value of -1.

Yearling female dispersal

Dispersal distance for dispersing juvenile female is derived from a random distribution

using parameters *mean-female-dispersal-distance* and *stddev-dispersal-distance*. If a

dispersing individual reaches a non-deer occupancy patch after dispersal, it is

transferred to the nearest deer occupancy patch. Dispersing yearling females change

their state variables *groid* and *gr* to -1 and -2 respectively.

3.7.5 Fawning

A proportion of female yearlings aged 13 months (determined by the parameter

juvenile-pregnancy-rate), and a proportion of adult female deer (determined by the

parameter *adult-pregnancy-rate*) are randomly selected to produce fawns (using 'hatch-

deer' to create new deer). Juvenile deer give birth to one fawn and adult deer give birth

to twins (MDC data). Sex ratio at birth is set at 1:1 [46]. Fawns inherit two state

variables from their dam: *groid* (female social group identifier) and *gr* (group size).

Additionally, male fawns have the state variable *mgroid* set to -2.

3.7.6 Deer mortality

If a female group leader dies (hunting or natural mortality), leadership is either a)

transferred to another adult female in the same group (new leader's state variable *gl*

changes from 0 to 1; members change their state variable *groid* to the new leader's ID

('who number'); b) if no adult female member exists in the group, members join other

small group (group size <= 3) on the same patch (change their state variable *groid* to

the new group leader's ID); or c) the group members become solitary (change their

state variables *gr* and *groid* to -2 and -1. If a bachelor group leader dies due to natural

mortality, leadership is transferred to one of the surviving group members.

Non-hunting mortality

The probability of a deer dying of natural or other non-hunting related causes during

every time step is determined by age- and sex- specific monthly mortality rates [47,48].

Irrespective of these rates, old deer (>240 months) have an overall high probability of

dying (0.8) during a time step. Fawns are functional ruminants at two months of age

[27], and therefore can possibly survive the death of their mother. We assume that

fawns less than two months old do not survive if their mother dies. If members of a doe

social group die during a time step, the group leader's state variable *gr* (accounting for

the group size) is decreased accordingly. If there are no members remaining in a group,

the leader becomes a solitary deer (state variables *fgroid* and *gr* changed to -1 and -2

respectively).

Hunting mortality

Hunting mortality is scheduled one time step after the rut period. Deer surviving the

monthly non-hunting mortality execute the hunting mortality submodel. Number of deer

harvested is specified by age- and sex- specific hunting mortality rates derived from

hunter-harvest data collected by the MDC.

In MO*OvPOPsurveillance*, sampling for CWD can be simulated as a random process

with equal probability of harvest for all deer in the model landscape, and random

sampling of the harvested deer for CWD testing. Harvest and sampling can also be

simulated as a non-random process. The non-random harvest scenario assigns

differential harvest probabilities to patches in the model landscape. Twenty-five percent

of the total patches occupied by deer are randomly selected and designated as no-

harvest patches, 15% as high harvest patches and remaining as low harvest patches.

Patches with most deer (average deer density) are selected as high harvest patches, as

50% of the total annual harvest is set to occur on high harvest patches. This differential

harvest scheme is extrapolated from the hunter and landowner survey data collected by

MDC.

Non-random sampling of harvested deer is simulated by assigning differential

probabilities for sampling. For every 8000 deer in the population, one high harvest patch

is designated as a location for a sampling station. Proximity to a sampling station

increases the probability of a harvested deer to be included in the subset that is tested

for CWD.

The following counters for each age class-sex category are updated after the harvest:

total deer harvested, total CWD+ deer harvested, total deer tested, total CWD+ deer

tested.

3.8 Parameterization and simulation experiments

3.8.1 Parameterization and Calibration

During initial setup, four parameters define the abundance and the structure of deer

population: *post_harvest_density*, *sexratio*, *adultprop* and *yearlingprop*. The proportion

of fawns in the initial population is calculated by subtracting the sum of *adultprop* and

yearlingprop from 1 (proportion of fawns = 1 – [*adultprop* + *yearlingprop*]).

Population dynamics of the model deer population is defined by two sets of age-sex-

specific parameters, *hunting mortality rates* and *non-hunting mortality rates*. Hunting

mortality rates are annual, while non-hunting mortality rates are monthly, and derived

from harvest data collected by MDC (Table 2). To facilitate use of the model in other

contexts, the interface has a chooser *recommended_parameter_values* which resets all

parameter sliders on the interface to values calibrated for sampling regions in Missouri.

The recommended parameter values are estimated from peer-reviewed literature,

derived from field-based surveys, or are based on expert opinions (Table 3).

Group sizes in the model are regulated using reporters, *mean-bachelor-group-size* and

doe-group-size-regulator. Values for *mean-bachelor-group-size* and *doe-group-size-*

regulator were calibrated so that the model group sizes are in agreement with the

references and expert opinion (Table 3).

Body mass attained during the breeding season appears to be a strong determinant of a

fawn's ability to breed [49]. The percent of female fawns that breed is influenced by the

population's level of nutrition. The reporter, *juvenile-pregnancy-rate*, was set at 20%,

based on data from an ongoing deer study in Missouri (Jon McRoberts, personal

communication). The proportion of adult females that breed, *adult-pregnancy-rate*, was

set at 80% (calibrated using MDC data available here:

https://extension2.missouri.edu/g9479).

Dispersal rates for yearling bucks range between 46 and 80% [45] but predictive

relationships are poorly understood [44]. We have set the yearling buck dispersal rate at

46% for simulations described in this paper. Similarly, based on observations from west-

central Illinois, the juvenile female dispersal rate is set at 22% [50]. Mean dispersal

distance for juvenile females was set at 11 miles – this value was extrapolated for a

post-harvest deer density of ~25 per forested km^2 from a logistic regression model

based on meta-analysis of juvenile female dispersal data [36].

MO*Ov*POP*surveillance* has 12 additional parameters that define the distribution of CWD

cases in the model landscape, nature of sampling (random or non-random) and

sampling intensity. Four sliders on the interface (*fawn-prevalence*, *yearling-prevalence*,

adult-prevalence, and *m-f-prev-ratio*) are used to specify age-sex class wise distribution

of CWD prevalence. The chooser *cwd_distribution* specifies the distribution of CWD+

deer: CWD+ deer are either randomly distributed in the model deer population

(*random_dist*) or are limited in distribution on a group of contiguous patches

(*clustered_dist*). In this model, clustering of CWD cases on the model landscape is best

simulated for very low prevalence scenarios when infected individuals are clustered

near the point of introduction [18–20]. For prevalence of ≤1%, *clustered_dist* distributes

CWD+ deer on a cluster of patches with an area ≤ 5% of the total area occupied by deer

in the model landscape. Another chooser, '*sampling*' selects the sampling process,

random or non-random sampling (described under *Hunting mortality*). Sample size for

each age-sex class are set using sliders on the interface (*%fawn-male-harvest-tested*,

%fawn-female-harvest-tested, *%yearling-male-harvest-tested*, *%yearling-female-*

harvest-tested, *%adult-male-harvest-tested* and *%adult-female-harvest-tested*).

4. Model evaluation

4.1 Evaluation of MO*Ov*POP performance

We initialized MO*Ov*POP with GIS coverage data (percent forest cover) for Franklin

County in eastern Missouri (Figure 2). Franklin County is of interest because of recent

detections of CWD-positive deer, and had the largest opening weekend deer harvest

(n=2041) in Missouri during the 2016-2017 deer harvest season [17]. Deer harvest data

(2016-17) collected by MDC was used to estimate the post-harvest density for Franklin

County (23 deer per forested square mile), age composition (35% fawns, 25% yearlings

and 40% adults), M:F ratio (1:1.2) and age-sex mortality parameters (Table 2). As per

MDC's data, the deer population in this region is very stable, therefore we simulated

deer population for Franklin County until a relatively stationary abundance was reached

(average population growth rate λ = 1). Annual population growth rate was calculated

using pre-harvest abundance.

The deer populations simulated for Franklin County using MO*Ov*POP reached a

relatively stationary pre-harvest abundance of ~24700 after 17 years (Figure 3). We

conservatively simulated the model for 25 years before exporting the pre-harvest

population snapshot for Franklin County. Franklin County deer population estimate prior

to the 2017 harvest season matched the simulated deer population abundance (B.

Keller, personal communication). Similarly, the age and sex structure of the model deer

population closely matched the harvest-data derived structure (Figure 4). Doe social

group sizes ranged between two and 11 individuals, and bachelor group size ranged

between two and eight individuals (Figure 5).

4.2 Evaluation of MO*Ov*POP *surveillance* performance

MO*Ov*POP*surveillance* was initialized using MO*Ov*POP generated pre-harvest deer

population for Franklin County. We evaluated MO*Ov*POP*surveillance* by comparing

model output (detection probabilities) with empirical data from standardized sample size

tables. Standardized tables provide sample sizes for representative samples, assuming

that disease is distributed randomly, and samples are obtained using a random process.

Sample size tables and online calculators like EpiTools (Sergeant, ESG, 2018. Epitools

epidemiological calculators. Ausvet Pty Ltd. Available at: http://epitools.ausvet.com.au)

use an approximation of the hypergeometric distribution to calculate sample sizes

necessary for desired detection probability (~confidence in detecting disease) if the

disease occurs at the specified prevalence in the target population [11].

We simulated random distribution of diseased individuals (four levels of prevalence:

0.5%, 1%, 2% and 5%) in a population of ~4000 deer (= pre-harvest abundance of adult

male deer in Franklin County). We specifically selected a simple scenario for model

evaluation, but MO*Ov*POP*surveillance* has the capacity to simulate age-sex specific

CWD prevalence in the model deer population, and thus can be used to determine

sample sizes for all age classes. Deer harvest and sampling of harvested individuals

were also simulated as random processes. For each prevalence level, sample sizes

with high detection probabilities (90%, 95% and 99%) were obtained from standardized

sample size tables and used here for model evaluation. We undertook 100 iterations of

the model for each prevalence-sample size scenario, and calculated detection

probability as the number of iterations in which at least one diseased deer was

detected. Detection probabilities derived from MO*Ov*POP*surveillance* simulations were

in close agreement with those obtained from sample size tables and EpiTools

epidemiological calculator (Table 4).

4.3 Sensitivity analysis

We evaluated how MO*Ov*POP*surveillance* derived detection probability responded to a)

change in CWD clustering pattern and b) change in non-random sampling process.

CWD prevalence in adult deer was set at 1% and sample size at 50% of the adult

harvest (~ 1460). The baseline scenario did not have any clustering (randomly

distributed CWD cases), and harvest as well as sampling of harvested deer were

simulated as random processes. For the 'Alternate' scenario, 100% of the CWD cases

were clustered on contiguous patches ~ ≤ 5% of the total area occupied by deer in the

model landscape, and non-random harvest was simulated by designating 15% of deer

habitat patches as high harvest pressure patches. Half of the total deer harvest (50%)

occurs on high harvest pressure patches. Sampling of harvested deer was also

simulated as a non-random process and is influenced by proximity to a sampling

station. Figures 6 and 7 illustrate CWD distribution (random and clustered) and

sampling (random and non-random), respectively. We further explored scenarios with

one cluster of infected individuals ('cluster 98%', 'cluster 96%', 'cluster 94%', 'cluster

92% and 'cluster 90%') and few infected cases (2 – 10%) randomly distributed in the

population. Harvest as well as sampling were simulated as non-random processes

similar to the alternate scenario. Finally, we explored scenarios where non-random

harvest was simulated by designating 18 - 30% of deer habitat patches as high harvest

pressure patches where 50% harvest occurred ('non-random sampling 18', 'non-random

sampling 21', 'non-random sampling 24', 'non-random sampling 27', 'non-random

sampling 30'). In these scenarios, clustering of CWD cases was similar to the alternate

scenario.

For each scenario, detection probabilities were derived from 10 replicates of 100 model

iterations (Figure 8). Without any clustering of CWD cases and random sampling, the

baseline scenario consistently yielded high detection probabilities (100%). Detection

probability decreased to ~76% with the alternate scenario, where 100% of CWD cases

were clustered and 15% of deer habitat patches had high harvest pressure (non-

random sampling). Detection probability was sensitive to changes in the number of

infected individuals outside the cluster. For the Franklin County deer population with 1%

CWD prevalence in adult deer and a sample size of ~1460, the confidence in detecting

CWD increased from 76% to 90% when 8-10 infected deer occurred outside the cluster.

This is particularly relevant for very low prevalence situations, for instance, where CWD

is recently introduced and all the cases are clustered around the point of infection. The

sample size threshold of detecting CWD confidently would increase later in the course

of the epizootic after CWD cases spread to areas outside the cluster. Changing the non-

random harvest pattern did not have much effect on the detection probability.

5. Model Application

We now illustrate how the models can be used to support CWD surveillance.

Simulations were designed to estimate the confidence in detecting CWD (detection

probability) for a range of prevalence - sample size combinations. We used five

hypothesized prevalence levels in adult deer of Franklin County (0.2%, 0.4%, 0.6%,

0.8% and 1%). Sample sizes were simulated as percent of the total adult deer

harvested (10%, 20%, 30%,40% and 50%). The adult deer harvest in our model

(~3000) approximates the adult harvest during 2016 Fall Firearm harvest in Franklin

County (MDC Deer Harvest Summary 2016-2017). For each hypothesized prevalence -

sample size combination, we simulated baseline and alternate scenarios (described

under Sensitivity analysis). We undertook ten replicates of 100 iterations of the model

for each prevalence-sample size and scenario combination.

Under the baseline scenario, 17 of 25 prevalence-sample size combinations had high

detection probabilities (≥95%) (Figure 9). Though CWD detection probability increased

with sample size and CWD prevalence, the relationship is not linear, and even under

such ideal assumptions the likelihood of reaching the 95% detection threshold was low

for the lowest prevalence rates (typical of the low prevalence rates of CWD that

currently are thought to occur in Missouri) and at the lowest sampling efforts. Thus,

under ideal assumptions, model-derived sample sizes for confident detection of CWD in

Franklin County deer population range between 1189 (CWD detection threshold set at

0.2% prevalence) and 297 (CWD detection threshold set at 1% prevalence). There was

a strong increase in sample sizes necessary for >95% detection probability for each

scenario as prevalence rates decrease. For example, necessary sampling efforts

double as prevalence declines from 1% to 0.6% and increases 3 fold - 5 fold at lower

prevalence.

The alternate scenario with spatial clustering of CWD and biased hunter harvest had

strong impacts on CWD detection probabilities. None of the tested sample sizes under

the alternate scenario had high detection probabilities. Even with 1% prevalence, the

largest sample size that we tested (50% of the adult harvest for Franklin County = ~

1500) had a detection probability of 78%. Detection probabilities further decreased as

prevalence decreased, and for 0.2% prevalence it dropped to 58%. We explored

increasing prevalence levels under the alternate scenario to determine the prevalence

threshold at which the sample size ~ 50% of adult harvest (~1500 deer tested) has a

high detection probability. This sample size approximates total deer sampled from

Franklin County during 2016 hunting season and represents the upper limit of sample

size under realistic conditions. Under non-realistic assumptions, a sample size of ~1500

should be able to confidently detect CWD when the adult prevalence is less than 0.2%

(<20 infected adults in the population). Our model explorations indicate that under

realistic assumptions, this sample size can confidently detect CWD (at least one CWD+

deer in the sample) when the adult population prevalence is more than 5% (>500

infected adults in the population).

The results of these simulation experiments (scenario-detection probability data) can be

used to determine CWD detection thresholds, calculate optimum sample sizes to

confidently detect CWD in target populations, and design a sustainable CWD

surveillance strategy.

6. Discussion

Sampling is a fundamental component of epidemiological surveillance as sample data

are used to infer about the presence or absence of a disease in the target population.

Surveillance of important wildlife diseases like avian influenza, bovine tuberculosis,

brucellosis, West Nile virus, and chronic wasting disease often rely upon samples

obtained from hunter-harvested individuals [1,3,4,19,51–55]. But hunter harvest

sampling does not approximate random sampling or any other probabilistic sampling

method, and such samples are not representative of the population. Nonetheless,

wildlife management agencies and researchers often calculate sample size (and

associated confidence level) necessary for disease detection in a target population

under the assumption that harvest-based sampling approximates probabilistic sampling

[14,52,55–57]. Such an approach may result in misleading and biased inferences about

the presence or absence of a disease in the target population.

Using computer simulation and simplified assumptions (uniform deer density across a

30 square mile landscape, and CWD cases clustered in the center of the landscape),

Samuel et al. [5] demonstrated that non-random sampling decreases the CWD

detection probability. Similarly, Nusser et al. [13] used a simulated white-tailed deer

population for one county in Iowa, USA to compare probability-based and alternative

sampling strategies, and their results similarly demonstrated that sample sizes

necessary for confidently detecting CWD are underestimated if assumptions of random

CWD distribution and random sampling are used for sample size calculations.

As recommended by Nusser et al. [13], we have developed a model-based tool that can

incorporate real world heterogeneities and biases, and can be applied to different

regions / deer management units to guide collection and analysis of CWD surveillance

data that relies on non-probabilistic methods like harvest-based sampling. An important

feature of our framework is the ability to simulate age-sex specific scenarios and

interventions as relevant individual host characteristics (age, sex, group membership)

and behaviors (dispersal, grouping behavior) have been incorporated in the model

programs. Further, the model programs are designed to be user friendly; the interface

sliders and choices allow users to update model assumptions based on their current

best knowledge of the system, and thereby employ an adaptive management approach

for defensible decision making while planning and implementing CWD surveillance

activities.

Results of the simulation experiments described in this paper (Table 5) show that

model-derived sample size requirement for detecting CWD with ≥ 95% confidence are

lower for the ecologically unrealistic baseline scenario that assumes random sampling

and random distribution of infected animals, compared to the alternate scenario that

incorporates realistic biases in sampling and disease distribution. For example, if the

objective of surveillance in Franklin County is to detect CWD with high confidence (≥

95%) when 1% (or more) of adult deer are infected, then under the baseline scenario

(random CWD distribution and random sampling) the model-derived sample size is ~

297 (10% of adult deer harvest for Franklin County). Sample size tables (based on

hypergeometric distribution, and assumption of random sampling) also recommend a

sample size of ~300 from a host population (~adult deer population in Franklin County)

with disease prevalence of 1% if the objective is disease detection with high confidence

(95%) [5,11]. But under the alternate scenario with 1% prevalence, sample size required

for detecting CWD with high confidence (≥95%) will be larger than 1500 (~50% adult

deer harvest for Franklin County). Unrealistic assumptions result in underestimation of

sample sizes due to the failure to account for spatial structure of the deer population,

CWD and harvest.

The importance of determining sample size targets using realistic assumptions cannot

be overemphasized, especially if the main objective of surveillance is to detect the

disease in areas where it has not been found yet so that control measures may be

rapidly implemented. As CWD is known to occur at a low prevalence near the point of

introduction during the early stages of the epizootic, sample size targets should be

determined using a low prevalence detection threshold [18–20]. The trade-off between

sample size and CWD detection threshold is an important consideration: the lower the

detection threshold is set, the larger the required sample size to confidently detect CWD

in the target population. But sampling effort also has to be considered in the context of

sustainability of surveillance efforts, as resources to support sampling efforts are limited

and subject to waning public interest. On the other hand, the very purpose of

surveillance is defeated if the detection threshold is set at a higher prevalence level to

facilitate use of a lower sample size. Our modeling framework can be used to optimize

surveillance strategies by 1) applying the model to smaller units (for example, township

units) to ensure reasonable sample size, low detection threshold, and meaningful

inferences about CWD occurrence and distribution; and 2) developing a 'risk map'

based on sampling history and other factors like proximity to areas where CWD has

been detected, captive facilities, geographic features, etc. to prioritize areas for

sampling.

Sampling history for a region (previous sampling effort and outcomes) can be useful for

making valid inferences about the presence or absence of the disease in the local

population. In particular, negative surveillance outcomes (no infected individuals

detected in the sample) can be used to infer the upper limit for the number of infected

individuals that could be present in the population. But such inferences would be

misleading if the sampling method assumes random sampling when surveillance

activities are based on non-random sampling. For instance, if no CWD-infected deer are

detected in a random sample of ~1189 adult deer harvested from Franklin County, then

the inferred upper limit of CWD prevalence in Franklin County can be estimated at 0.2%

(~ 20 infected adult deer) (Table 5). But under realistic assumptions (clustered CWD

distribution and non-random sampling), not finding an infected deer in a sample of

~1189 harvested deer from Franklin County would translate to an upper limit of CWD

prevalence of > 5% (> 500 infected deer).

Importantly, our model-based framework supports an adaptive management approach.

Surveillance strategies for target populations can be iteratively improved by updating

the models with current information on host demography, harvest patterns and disease

prevalence. Furthermore, the model landscape can readily incorporate locale-specific

geographic features to better simulate host, harvest and disease distribution. Using the

example of CWD surveillance in white-tailed deer of Missouri, we have illustrated how

the model-based framework can guide collection and analysis of surveillance data that

relies on non-probabilistic methods like harvest-based sampling. The broader

framework can be readily adapted for surveillance of CWD in white-tailed deer

populations of other regions, or for surveillance of other important diseases of white-

tailed deer like bovine tuberculosis and epizootic hemorrhagic disease. The framework

can also be adapted to other host-pathogen systems and used for informed-decision

making while planning and implementing disease surveillance in wild and free-ranging

species.

Acknowledgments

We thank and acknowledge D. Corcoran Barrios, J. Batten, T. Bixler, E. Flinn, V.
Grimm, J. McRoberts, C. Miller and C. Scroggins for useful suggestions. Support for this
study came from the Missouri Department of Conservation. AB was also partly
supported by NSF Award P20GM104420.

References

1.   Drew L, Jessup A, Burr A, Franti CE. Serologic survey for brucellosis in feral

swine, wild ruminants, and black bear of California, 1977 to 1989. J Wildl Dis.

1992;28: 355–363. doi:10.7589/0090-3558-28.3.355

2.   Muñoz PM, Boadella M, Arnal M, de Miguel MJ, Revilla M, Martínez D, et al.

Spatial distribution and risk factors of brucellosis in Iberian wild ungulates. BMC

Infect Dis. 2010;10: 46. doi:10.1186/1471-2334-10-46

3.   Dusek RJ, Bortner JB, DeLiberto TJ, Hoskins J, Franson JC, Bales BD, et al.

Surveillance for high pathogenicity avian influenza virus in wild birds in the Pacific

Flyway of the United States, 2006-2007. Avian Dis. 2009;53: 222–230.

doi:10.1637/8854.1

4.   Dusek RJ, Hagen CA, Franson JC, Budeau DA, Hofmeister EK. Utilizing hunter

harvest effort to survey for wildlife disease: A case study of West Nile virus in

greater sage-grouse. Wildl Soc Bull. 2014;38: 721–727. doi:10.1002/wsb.472

5.   Samuel MD, Joly DO, Wild MA, Wright SD, Otis DL, Werge RW, et al.

Surveillance strategies for detecting chronic wasting disease in free-ranging deer

and elk: results of a CWD surveillance workshop, 10-12 December 2002. United

States Geol Surv Natl Wildl Heal Center, Madison, Wisconsin, USA. Madison,

Wisconsin, USA; 2003; 41.

6.   Rees EE, Merrill EH, Bollinger TK, Hwang Y Ten, Pybus MJ, Coltman DW.

Targeting the detection of chronic wasting disease using the hunter harvest during

early phases of an outbreak in Saskatchewan, Canada. Prev Vet Med. Elsevier

B.V.; 2012;104: 149–159. doi:10.1016/j.prevetmed.2011.10.016

7. Haley NJ, Hoover EA. Chronic wasting disease of cervids: Current knowledge and

future perspectives. Annu Rev Anim Biosci. 2015;3: 305–325.

doi:10.1146/annurev-animal-022114-111001

8. Samuel MD, Storm DJ. Chronic wasting disease in white-tailed deer: infection,

mortality and implications for heterogeneous transmission. Ecology. 2016;97:

3195–3205.

9. Gross JE, Miller MW. Chronic wasting disease in mule deer: disease dynamics

and control. J Wildl Manage. 2001;65: 205–215. doi:10.2307/3802899

10. Edmunds DR, Kauffman MJ, Schumaker BA, Lindzey FG, Cook WE, Kreeger TJ,

et al. Chronic wasting disease drives population decline of white-tailed deer. PLoS

One. 2016;11: e0161127. doi:10.1371/journal.pone.0161127

11. Cannon RM, Roe RT. Livestock disease surveys. A field manual for veterinarians.

Bureau of Resource Science, Department of Primary Industry. Aust. Govt. Publ.

Serv. Canberra; 1982.

12. Fosgate GT. Practical sample size calculations for surveillance and diagnostic

interventions. J Vet Diagnostic Investig. 2009;21: 3–14.

doi:10.1177/104063870902100102

13. Nusser SM, Clark WR, Otis DL, Huang L. Sampling considerations for disease

surveillance in wildlife populations. J Wildl Manage. 2008;72: 52–60.

doi:10.2193/2007-317

14.   Beringer J, Hansen L, Millspaugh J, Meyer T. A statewide surveillance effort for

detecting chronic wasting disease in wild white-tailed deer in Missouri. Wildl Soc

Bull. 2003;31: 873–881. Available: http://www.jstor.org/stable/3784613

15.   ProMED-Mail. Chronic wasting disease, cervid - USA (02):(MISSOURI). ProMed-

mail 20120125.1020977. 2012.

16.   ProMED-Mail. Chronic wasting disease, cervid - USA (22): (Missouri). ProMed-

mail 20151217.3870576. 2015.

17.   CWD surveillance and monitoring in Missouri [Internet]. [cited 21 Dec 2017].

Available: https://huntfish.mdc.mo.gov/hunting-trapping/wildlife-diseases/chronic-

wasting-disease-cwd/cwd-surveillance-and-monitoring

18.   Walsh DP. Enhanced surveillance strategies for detecting and monitoring chronic

wasting disease in free-ranging cervids: [Internet]. U.S. Geological Survey Open-

File Report. 2012. Available: http://pubs.usgs.gov/of/2012/1036/

19.   Joly DO, Samuel MD, Langenberg JA, Blanchong JA, Batha CA, Rolley RE, et al.

Spatial epidemiology of chronic wasting disease in Wisconsin white-tailed deer. J

Wildl Dis. 2006;42: 578–588. doi:10.7589/0090-3558-42.3.578

20.   Osnas EE, Heisey DM, Rolley RE, Samuel MD. Spatial and temporal patterns of

chronic wasting disease: Fine-scale mapping of a wildlife epidemic in Wisconsin.

Ecol Appl. 2009;19: 1311–1322. doi:10.1890/08-0578.1

21.   VerCauteren KC, Hygnstrom SE. White-tailed deer. In: Wishart D, editor.

Encyclopedia of the Great Plains. Lincoln, NE: University of Nebraska Press;

2004. pp. 642–643.

22.   Walter WD, VerCauteren KC, Campa H, Clark WR, Fischer JW, Hygnstrom SE, et

al. Regional assessment on influence of landscape configuration and connectivity

on range size of white-tailed deer. Landsc Ecol. 2010;24: 1405–1420.

doi:10.1007/s10980-009-9374-4

23.   Ozoga J. Whitetail intrigue. Krause, Iola, Wisconsin USA; 2000.

24.   Monteith KL, Sexton CL, Jenks JA, Terry Bowyer R. Evaluation of techniques for

categorizing group membership of white-tailed deer. J Wildl Manage. 2007;71:

1712–1716.

25.   Nelson ME, Mech LD. Dispersal in female white-tailed deer. J Mammal. 1992;73:

891–894.

26.   Miller BF, DeYoung R., Campbell TA, Laseter BR, Ford WM, Miller KV. Fine-scale

genetic and social structuring in a central Appalachian white-tailed deer herd. J

Mammal. 2010;93: 681–689.

27.   Marchinton RL, Hirth D. Behavior. In: Halls LK, editor. White-tailed deer: Ecology

and management. Harrisburg, PA: Stackpole Books; 1984. pp. 129–168.

28.   Smith, W P. Odocoileus virginianus. Mamm Species. 1991;388: 1–13.

29.   Miller KV, Muller LI, Demarias S. White-tailed deer (Odocoileus virginianus). In:

Feldhamer GA, Thompson BC, Chapman JA, editors. Wild mammals of North

America: Biology, management, and conservation. 2nd ed. Baltimore, Maryland:

Johns Hopkins University Press; 2003. pp. 906–930.

30. Ozoga J., Verme L., Bienz C. Parturition behavior and territoriality in white-tailed

deer: Impacts on neonatal mortality. J Wildl Manage. 1982;46: 1–11.

31. Schwede G, Hendrichs H, McShea W. Social and spatial organization of female

white-tailed deer, Odocoileus virginianus, during the fawning period. Anim Behav.

1993;45: 1007–1017.

32. Hirth D. Social behavor of white-tailed deer in relation to habitat. Wildl Monogr.

1977;53: 1–55.

33. Cullingham CI, Merrill EH, Pybus MJ, Bollinger TK, Wilson GA, Coltman DW.

Broad and fine-scale genetic analysis of white-tailed deer populations: Estimating

the relative risk of chronic wasting disease spread. Evol Appl. 2011;4: 116–131.

doi:10.1111/j.1752-4571.2010.00142.x

34. Nelson ME. Natal dispersal and gene flow in white-tailed deer in northeastern

Minnesota. J Mammal. 1993;74: 316–322.

35. Purdue JR, Smith MH, Patton JC. Female philopatry and extreme spatial genetic

heterogeneity in white-tailed deer. J Mammal. 2000;81: 179–185.

36. Lutz CL, Diefenbach DR, Rosenberry CS. Population density influences dispersal

in female white-tailed deer. J Mammal. 2015;96: 494–501.

37. Long ES, Diefenbach DR, Rosenberry CS, Wallingford BD. Multiple proximate

and ultimate causes of natal dispersal in white-tailed deer. Behav Ecol. 2008;19:

1235–1242.

38.   Lutz CL, Diefenbach DR, Rosenberry CS. Proximate influences on female

dispersal in white-tailed deer. J Wildl Manage. 2016;80: 1218–1226.

39.   Wilensky U. NetLogo [Internet]. Center for Connected Learning and Computer-

Based Modeling, Northwestern University, Evanston, IL; 1999. Available:

http://ccl.northwestern.edu/netlogo/.

40.   Grimm V, Berger U, Bastiansen F, Eliassen S, Ginot V, Giske J, et al. A standard

protocol for describing individual-based and agent-based models. Ecol Modell.

2006;198: 115–126.

41.   Grimm V, Berger U, DeAngelis DL, Polhill JG, Giske J, Railsback SF. The ODD

protocol: A review and first update. Ecol Modell. 2010;221: 2760–2768.

doi:10.1016/j.ecolmodel.2010.08.019

42.   Belsare A V., Gompper ME, Millspaugh JJ. MOOvPOP (Version 8). In: CoMSES

Computational Model Library [Internet]. 2017. Available:

https://www.openabm.org/model/5585/version/8

43.   Belsare A V., Gompper ME, Millspaugh JJ. MOOvPOPsurveillance. In: CoMSES

Computational Model Library [Internet]. 2017. Available:

https://www.openabm.org/model/5576/version/4

44.   Diefenbach DR, Long ES, Rosenberry CS, Wallingford BD, Smith DR. Modeling

distribution of dispersal distances in male white-tailed deer. J Wildl Manage.

2008;72: 1296–1303. doi:http://dx.doi.org/10.2193/2007-436

45.   Long ES, Diefenbach DR, Rosenberry CS, Wallingford BD, Grund MD. Forest

cover influences dispersal distance of white-tailed deer. J Mammal. 2005;86:

623–629. doi:http://dx.doi.org/10.1644/1545-1542(2005)86[623:FCIDDO]2.0.CO;2

46. Ditchkoff SS. Anatomy and Physiology. In: Hewitt DG, editor. Biology and

management of white-tailed deer. Boca Raton, FL: CRC Press; 2011. pp. 43–73.

47. Hiller TL, Henry Campa I, Winterstein SR. Survival and space use of fawn white-

tailed deer in southern Michigan. Am Midl Nat. 2008;159: 403–412.

48. Van Deelen TR, Campa Iii H, Haufler JB, Thompson PD. Mortality patterns of

white-tailed deer in Michigan's Upper Peninsula. J Wildl Manage. 1997;61: 903–

910.

49. Gaillard JM, Festa-Blanchet M, Yoccoz N., Loison A, Toigo C. Temporal variation

in fitness components and population dynamics of large herbivores. Annu Rev

Ecol Syst. 2000;31: 367–393.

50. Nixon CM, Mankin PC, Etter DR, Hansen LP, Brewer PA, Chelsvig JE, et al.

White-tailed deer dispersal behavior in an agricultural environment. Am Midl Nat.

2007;157: 212–220.

51. Bevins SN, Dusek RJ, White CL, Gidlewski T, Bodenstein B, Mansfield KG, et al.

Widespread detection of highly pathogenic H5 influenza viruses in wild birds from

the Pacific Flyway of the United States. Sci Rep. 2016;6: 28980.

doi:10.1038/srep28980

52. Schöning JM, Cerny N, Prohaska S, Wittenbrink MM, Smith NH, Bloemberg G, et

al. Surveillance of bovine tuberculosis and risk estimation of a future reservoir

formation in wildlife in Switzerland and Liechtenstein. Gordon S V., editor. PLoS

One. 2013;8: e54253. doi:10.1371/journal.pone.0054253

53. O'Brien DJ, Schmitt SM, Berry DE, Fitzgerald SD, Lyon TJ, Vanneste JR, et al.

Estimating the true prevalence of Mycobacterium bovis in free-ranging elk in

Michigan. J Wildl Dis. 2008;44: 802–810.

54. Rees EE, Merrill EH, Bollinger TK, Hwang Y Ten, Pybus MJ, Coltman DW.

Targeting the detection of chronic wasting disease using the hunter harvest during

early phases of an outbreak in Saskatchewan, Canada. Prev Vet Med. Elsevier

B.V.; 2012;104: 149–159. doi:10.1016/j.prevetmed.2011.10.016

55. Diefenbach DR, Rosenberry CS, Boyd RC. Efficacy of detecting chronic wasting

disease via sampling hunter-killed white-tailed deer. Wildl Soc Bull. 2004;32: 267–

272.

56. Schmitt SM, Fitzgerald SD, Cooley TM, Bruning-Fann CS, Sullivan L, Berry D, et

al. Bovine tuberculosis in free-ranging white-tailed deer from Michigan. J Wildl

Dis. 1997;33: 749–758.

57. Bevins SN, Dusek RJ, White CL, Gidlewski T, Bodenstein B, Mansfield KG, et al.

Widespread detection of highly pathogenic H5 influenza viruses in wild birds from

the Pacific Flyway of the United States. Sci Rep. 2016;6: 28980.

doi:10.1038/srep28980

Table 1. State variables of agents included in MO*Ov*POP and MO*Ov*POP*surveillance*.

All state variables except the deer state variable 'aim' are unitless.

Agent	Variable	Description
Patch	forest-percent	forest cover expressed as a proportion
	border	patches on the edge of the model landscape have border = 1, other patches have border = 0
	dfp	mean forest-percent for a patch and its immediate neighbors
	do	deer occupancy; 1 if deer occur on a patch, 0 otherwise
	dh	deer habitat; ≥ 1 if a patch or a group of contiguous patches qualifies as deer habitat, otherwise < 1
	add	average deer density calculated for a patch and its immediate neighbors (8 patches)
	hp	designates harvest pressure on the patch: 0 no harvest, 1 low harvest and 2 high harvest
Deer	sex	1 if male, 2 if female
	aim	age in months
	momid	dam's id number
	gl	1 if a doe social group leader, 0 otherwise
	ml	1 if a bachelor group leader, 0 otherwise
	groid	≥ 0 if member of a doe social group, -1 if a solitary female, 0 if a male deer
	gr	≥ 1 if a group leader, -1 if member of a doe social group, otherwise -2
	mgroid	0 if a female deer, -2 if a male fawn, group leader id if member of a bachelor group
	cwd	CWD infection status of a deer: 0 uninfected and 1 infected
	marked-for-harvest?	deer marked for harvest
	marked-for-testing?	deer marked for sampling (CWD testing)

851 Table 2. Age- and sex-specific mortality parameter values used in MO*Ov*POP and
852 MO*Ov*POP*surveillance*. These values are derived from harvest data collected by the
853 Missouri Department of Conservation.

Age-Sex Class	Non-hunting mortality [a]		Hunting mortality [b]	
	Parameter	Value	Parameter	Value
male fawns (0 – 6 months)	*mf6nhm*	0.055	*mf6hm*	0
female fawns (0 – 6 months)	*ff6nhm*	0.055	*ff6hm*	0
male fawns (7 – 12 months)	*mf12nhm*	0.05	*mf12hm*	0.05
female fawns (7 – 12 months)	*ff12nhm*	0.05	*ff12hm*	0.02
male yearlings (13 – 24 months)	*mynhm*	0.01	*myhm*	0.25
female yearlings (13 – 24 months)	*fynhm*	0.00	*fyhm*	0.15
male adults (> 25 months)	*manhm*	0.01	*mahm*	0.40
female adults (> 25 months)	*fanhm*	0.02	*fahm*	0.20

854 [a] per month; [b] per year

855 Table 3. Parameter values used for initial population setup and simulation of white-tailed
856 deer behavior. Calibrated values are indicated by an asterisk.

Parameter	Description	Value	Reference
Population setup and distribution			
post_harvest_density	Initial deer density (per forested sq. mile)	23	MDC
sexratio	Male: female ratio in the population	1:1.2*	MDC
adultprop	Adult proportion (≥ 25 months) in the population	0.4*	MDC
yearlingprop	Yearling proportion in the population	0.25*	MDC
min-forestcover-percent	Minimum percent forest cover of a patch for deer occupancy	0.25*	[21,22]
max-forestcover-percent	Maximum percent forest cover of a patch for deer occupancy	0.75*	[21,22]
Behavior			
mean-bachelor-group-size	mean (± sd) number of adult bucks in a bachelor group	2 - 5*	[27–29], L.Hansen, pers.obs
doe-group-size-regulator	doe social group size above which it undergoes fission	6*	[25,26], L.Hansen, pers.obs
juvenile-pregnancy-rate	proportion of female fawns that reproduce	0.2	J. McRoberts, pers.comm.
adult-pregnancy-rate	proportion of adult females that reproduce	0.8*	MDC
yearling-male-dispersal-rate	proportion of dispersing yearling males	0.46*	[44,45]
yearling-female-dispersal-rate	proportion of dispersing yearling females	0.22*	[50]
mean-female-dispersal-distance	mean dispersal distance for yearling females (miles)	11*	[36]
stddev-dispersal-distance	standard deviation for the mean dispersal distance of yearling females (miles)	4*	[36]

859 Table 4. MO *OvPOP surveillance* derived detection probabilities for hypothesized CWD
860 prevalence-sample size scenarios for a deer population size ~4000 assuming random
861 CWD distribution and random sampling. Model-derived detection probabilities closely
862 matched those derived from hypergeometric distribution.

Population prevalence	Sample Size	Detection probability	
		Sample size table*	MO *OvPOP surveillance*
0.5%	434	90%	90%
	550	95%	95%
	821	99%	97%
1%	223	90%	89%
	288	95%	96%
	433	99%	100%
2%	113	90%	90%
	146	95%	96%
	222	99%	97%
5%	45	90%	90%
	58	95%	94%
	89	99%	99%

865 *Approximation of the hypergeometric distribution.

866 Table 5. Sample sizes with detection probabilities for Franklin County contrasted for the
867 baseline (random CWD distribution, random harvest and sampling) and alternate
868 (clustered CWD distribution, non-random harvest and sampling) scenarios.

	Prevalence				
Scenarios	**0.2%**	**0.4%**	**0.6%**	**0.8%**	**1%**
Random CWD distribution, random sampling	1189 (≥95%)	891 (≥95%)	594 (≥95%)	297 (≥95%)	297 (≥95%)
Clustered CWD distribution, non-random sampling	1486 (59%)	1463 (64%)	1463 (75%)	1463 (75%)	1463 (78%)

871 Fig 1. Schedule of processes in MO*Ov*POP and MO*Ov*POP*surveillance*. The
872 rectangular boxes represent actions performed by deer ('agents'), and the shaded
873 boxes represent actions performed by the model at the population scale.

875

Fig 2. Map of Missouri with county outlines. Franklin County is marked with an asterisk.

Fig 3. Population growth rate (λ) for the Franklin County, Missouri, deer population
simulated using MO*Ov*POP. The population reached a relatively stationary abundance
after 17 years (vertical line), when λ converged to 1.

Fig 4. Age composition of the simulated Franklin County deer population. After the 5th

year, age and sex composition of the population remained stable and in agreement with

the derived values for Franklin County.

Fig 5. Group size distribution snapshot for bachelor and doe social groups in the
simulated Franklin County deer population. Group size distributions were in agreement
with references and expert opinions throughout the model run.

Fig 6. Random (A) or clustered (B) distribution of infected deer in
MO*OvPOPsurveillance* landscape representing Franklin County, Missouri. Infected deer
(red) are shown on the model landscape, and yellow colored patches in (B) represent
the disease cluster.

(A)             (B)

Fig 7. (A) MO *OvPOP* *surveillance* landscape representing Franklin County with uniform
harvest pressure. (B) The same landscape with non-uniform harvest pressure to
simulate non-random sampling. Blue color indicates patches where deer are not
harvested, yellow indicates low harvest pressure and orange indicates high harvest
pressure. Patches labelled 'SS' indicate sampling stations.

(A)   (B)

Fig 8. Model-derived detection probabilities for Franklin County deer population with
different CWD clustering and/or harvesting (sampling) patterns. CWD prevalence was
simulated at 1% (adult deer only) and the sample size was set at ~1460 (= 50% adult
harvest). Non-random sampling scenarios indicate the proportion of high harvest
pressure patches in the model landscape and cluster scenarios indicate the proportion
of CWD cases occurring within the cluster. Baseline scenario had no clustering of CWD
cases and uniform harvest pressure, while alternate scenario had 100% of CWD cases
within the cluster and 15% of the patches designated as high harvest patches. Shaded
area indicates high detection probabilities (≥ 95%).

Fig 9. Model-derived detection probabilities for a range of sample sizes (10% to 50% of
adult deer harvest) and CWD prevalence levels (0.2% to 1%). The baseline scenario
assumes random distribution of infected individuals in the population, random harvest
and random sampling. The alternate scenario assumes clustered distribution of infected
individuals, non-random harvest and non-random sampling. Shaded area indicates high
detection probabilities (≥ 95%).