Biospeckle laser digital image processing for quantitative and statistical evaluation of the activity of Ciprofloxacin on *Escherichia coli* K-12

Hilda Cristina Grassi[1], Ana Velásquez[2*], Olga Mercedes Belandria[1], María Lorena Lobo-Sulbarán[3], Jesús E. Andrades-Grassi[4], Humberto Cabrera[5,6], Efrén D. J. Andrades[1]

[1] Facultad de Farmacia y Bioanálisis, Universidad de Los Andes, Mérida, Venezuela

[2] Centro Multidisciplinario de Ciencias, Instituto Venezolano de Investigaciones Científicas, Mérida, Venezuela

[3] Cátedra de Farmacología y Terapéutica, Departamento de Biopatología, Facultad de Odontología, Universidad de Los Andes, Mérida, Venezuela

[4] Departamento de Ordenación de Cuencas, Facultad de Ciencias Forestales y Ambientales, Universidad de Los Andes Mérida, Venezuela

[5] Abdus Salam International Centre for Theoretical Physics (ICTP), Trieste, Italy

[6] Istituto Nazionale di Fisica Nucleare, Trieste, Italy

*Corresponding Author
avelasqu@ivic.gob.ve

Abstract

Antibiotic susceptibility testing is a necessary step prior to the treatment of clinical infections. A major concern is the time required to obtain a fast and reliable result. The aim of this work is to use Biospeckle laser in a 15min assay for an antimicrobial susceptibility test of Ciprofloxacin in serial two-fold dilutions on *Escherichia coli* K-12 using Venereal Disease Research Laboratory (VDRL) plates. Analysis of images by video edition is performed on a quantitatively selected region of interest, and processed with ImageJ-ImageDP; and by the construction of time series and analysis with either statistical diagnostics tests or Autoregressive Integrated Moving Average (ARIMA) models. Antimicrobial susceptibility tests are also performed for the purpose of quantitative comparison, showing a profile that is comparable to the result obtained with ImageJ-ImageDP processing after 15min of antibiotic action. Only the time series of the least affected bacteria (low Ciprofloxacin concentration) behaves in an expected manner, being non-independent and mainly non-linear, non-normal, and heteroscedastic. The most affected bacteria (higher Ciprofloxacin concentration) are non-independent and tend to be linear, normal and heteroscedastic. Adjustment to a linear regression identifies both, the culture medium without bacteria and the most affected bacteria, normality identifies the most affected bacteria and heteroscedasticity-homoscedasticity distinguishes the presence-absence of bacteria, respectively. ARIMA models $(1,1,1)(1,0,1)_{11}$ and $(4,1,1)(1,1,1)_{11}$ fit the time series of the most affected bacteria while the latter also fits the culture medium without bacteria. The time series of the least affected bacteria are identified by a $(7,1,2)(1,0,1)_{11}$ model. The non-linear, non-normal and heteroscedastic behavior of this group is probably responsible for its adjustment to a model with a relatively high parameter. The four methods: diagnostic statistical tests, fitting of ARIMA models, ImageJ-ImageDP and antimicrobial susceptibility tests, show similar results, being able to distinguish among the groups of assays with bacteria and Ciprofloxacin below and above the Minimal Inhibitory Concentration.

Author Summary

Biospeckle laser patterns occur when a dynamic surface is illuminated. This research describes its application to the activity of *Escherichia coli* bacteria and the effect of 18 different concentrations of an antibiotic (Ciprofloxacin) in a 15min assay, using VDRL plates where the sample has a relatively small volume and is flat shaped. The assay is performed on an anti-vibration table in a dark room with a laser that sequentially illuminates each of the wells of the plate. A camera takes short 30sec videos with approximately 750 frames and sends them to a computer where image processing takes place. In order to select a segment of 80 successive frames to analyze, the region with the higher variation was identified, punched out and edited as a "flip-book animation" with a program named ImageJ and processed with another program named ImageDP that takes the difference between successive frames and is able to describe the speed with which the Biospeckle dots move, expecting to show that the antibiotic affects the bacteria by changing the speed with which they move. Also in each video, within the region of higher differential activity, a pixel was selected to construct a time series which is the successive value of that pixel in 253 frames, representing a recording of 8sec. This was analyzed with two statistical methods: diagnostic statistics tests and ARIMA models, both of which try to demonstrate how the results are organized. All the results, the speed of the dots in the "flip-book

94 animation" and the structure of the data of the time series, were comparable to those
95 obtained with traditional antimicrobial susceptibility tests with the same bacteria and
96 antibiotic.
97
98

Introduction
Antibiotic susceptibility testing is a necessary step prior to the treatment of clinical
infections. A major concern is the time required to obtain a fast and reliable result
especially in the case of acute bacterial infections for which it is necessary to determine the
appropriate antibiotic that must be used. Established assays such as broth and agar dilution,
and disk diffusion are the accepted techniques [1] but new approaches are needed in order
to reduce the required time and the sample volume. Another issue is to search for the
possibility of testing direct clinical samples reducing the assay requirements for handling
and treatment of the microbial contaminated fluid as well as bacterial isolation and
enrichment. Fast antimicrobial susceptibility testing has been achieved by diverse methods,
among these are the molecular techniques [2,3] biosensors [4] fluorescence detection [5] ,
as well as the adaptation of new techniques such as Raman Spectroscopy [6]. Therefore it is
desirable to perform an exploratory assay which could eventually set the basis for a fast
antibiogram.
113
Biospeckle Laser is a non-invasive technique and refers to a pattern that occurs when a
laser beam illuminates a dynamic surface [7], such as a microorganism contaminated
sample which has a movement of its surface that causes the wave fronts to interfere and
produce a pattern of moving dots [8]. This technique has been used in different fields of
research as well as at macroscopic and microscopic levels for a variety of applications [9,
10,11]. Usually this dynamic phenomenon is caught in a video for image processing.
Diverse approaches and descriptors have been proposed that enable the information of the
videos to be transformed in a number that represents the activity of the sample [12].
However, this is a multivariate phenomenon that is affected by the size of the organism to
be detected, the speed and the direction of its movement [13], the type of medium or
environment in which the organism is contained, etc.
125
The aim of the present work is to adapt the Biospeckle laser technique to antimicrobial
susceptibility testing for Ciprofloxacin on *Escherichia coli* K-12. Ciprofloxacin is a broad-
spectrum antimicrobial carboxyfluoroquinoline which selectively poisons preferentially the
GyrA subunit of gyrase in Gram-negative bacteria and the ParC subunit of Topo IV in
Gram-positive bacteria, with generation of chromosome breaks, which culminate in cell
death [14,15]. An assay using VDRL plates that had already been used for testing
Benznidazol on *Trypanosoma cruzi* [16], was adapted to *E. coli*. However, due to the type
of microorganism, its size, motility and cell density, the processing of the images was
performed on a quantitatively selected region of interest (ROI) which was analyzed through
different approaches, such as video edition-processing and descriptive statistics; and time
series construction and analysis with either statistical diagnostics tests or ARIMA models.
A time series data set consists of observations on a variable or several variables over time
which can be used for repeated-measures analysis or model adjustment [17]. Interpretation
of Biospeckle images by time series analysis has already been proposed [18]. The selection

of a ROI resulted in an important step because it allowed the consideration of different
activity-level regions, as previously reported [19,20]. In this work a low Biospeckle
intensity (I) region with a high differential bacterial activity was identified which was
suitable for further analysis. Antimicrobial susceptibility tests were also performed for the
purpose of quantitative and statistic comparison.
145
146

Materials and Methods

Antibiotic, Bacteria and culture conditions.
Escherichia coli K-12 (*HfrH b_1^-*) [21] designated as strain 322 in the Laboratory Collection,
was used in all the assays. For the Biospeckle and microbiological techniques, bacteria
were grown overnight in Mueller-Hinton Broth (MHB) Himedia and prior to use it was
adjusted to 0.5 McFarland standard which corresponds to approximately 1.5E+08
bacteria/mL (Absorbance 0.08-0.1 at λ_{625nm} in a Spectronic 20) [22]. The antibiotic used
was Ciprofloxacin (BACIPRO^{TM} BIOGALENIC), i.v. infusion, 200mg/100mL. The leaflet
indicates that it contains NaCl 0.9% (354 mg Na/100mL). This antibiotic was used in two-
fold serial dilutions in sterile NaCl 0.9%.
158

VDRL Plate Assay design for Biospeckle laser.
The assay was designed as previously described [16]. Ciprofloxacin was serially diluted in
saline (0.9% sterile NaCl) . 95 μL of either Mueller-Hinton Broth (18 odd numbered wells)
or bacterial suspension in Mueller-Hinton Broth (18 even numbered wells) were distributed
in an alternating manner in wells of VDRL plates. At t=0, five μL of the antibiotic dilution
were added to two contiguous wells, one without bacteria and one with bacteria, so that
only the even numbered wells contained bacteria and both contained antibiotic at the same
concentration. The plates were left still to avoid vibrations and to allow 15 minutes for the
action of the drug at room temperature (20-25°C). Also 4 control assays were performed.
168
Table 1 shows the composition of each well. After 15 minutes, each well of the VDRL
plate was processed under the experimental setup to obtain a video for each well. Four
control wells (videos) were performed: Control A (95μL MHB + 5μL NaCl 0.9%); Control
B (95μL with 1.43E+07 bacteria in MHB + 5μL NaCl 0.9%); Control C (100μL MHB);
Control D (100μL with 1.50E+07 bacteria in MHB). The dimensions of the VDRL Plates
were as specified by the supplier and previously reported; the mean internal diameter of
each well was 12.7+/- 0.1 mm and the height of 100μL in the well, was 0.75mm [16]. As it
is described further on, the laser beam has a diameter of 10mm on the surface. With these
dimensions the calculated volume of the illuminated liquid is 60μL and the number of
bacteria present in this volume is approximately 8.6E+06, which is an appropriate number
for evaluating the activity of the Biospeckle pattern.

Table 1. VDRL Plate Assay design for Biospeckle laser.

VIDEOS (Wells)	CIPROFLOXACIN µg/mL in well
1 and 2	7.63E-04
3 and 4	1.53E-03
5 and 6	3.05E-03
7 and 8	6.10E-03
9 and 10	1.22E-02
11 and 12	2.44E-02
13 and 14	4.88E-02
15 and 16	9.77E-02
17 and 18	1.95E-01
19 and 20	3.91E-01
21 and 22	7.81E-01
23 and 24	1.56E+00
25 and 26	3.13E+00
27 and 28	6.25E+00
29 and 30	1.25E+01
31 and 32	2.50E+01
33 and 34	5.00E+01
35 and 36	1.00E+02

Table 1. Concentration of Ciprofloxacin in the wells of the VDRL plates for Biospeckle laser. Odd numbered videos contained 95µL of MHB and 5µL of the antibiotic. Even numbered wells contained 95µL of bacteria (approximately 1.43E+07 total cells) in MHB and 5µL of the antibiotic. The same number was assigned to the well and to the video.

Experimental setup.

The experimental setup was the same as previously described [16] and is shown in Fig 1. Briefly, the setup rests on top of an anti-vibration table. The Biospeckle laser imaging system has a 1mW He-Ne unpolarized laser operating at 632.8nm. The laser which is located at a distance of 50cm is coupled to a convex divergent lens and the beam forms a 10mm spot in the individual wells of the VDRL plate, with an angle of incidence of 72°. Since the wells have an internal diameter of 12.7+/- 0.1mm, the beam illuminates the center of the well of the VDRL plate which rests on an opaque black cardboard, avoiding sparkles and edge effects. A CCD camera (Thorlabs USB.2, 30fps) is located at 30cm from the sample and is connected to a PC which records 0.5-1min videos. The resolution of the camera is 320×240, pixel size 6.45µm, speckle size aprox. 21.6µm. The speckle video data is sent to the computer for video and image processing. In order to visualize a significant number of bacteria per pixel, a relatively low resolution was chosen for the camera. Thus it was possible to apply statistical diagnostic tests and models on the time series of a selected pixel and descriptive statistics on the region of interest (ROI).

Fig 1. Schematic diagram of the speckle experimental setup. Experimental setup, with data of the setup and the VDRL well, L laser; C CCD camera; PC computer; l1 lens of the laser; l2 lens of the camera; AT anti-vibration table; VP VDRL plate; SW sample well; OP object's plane;W width; HL height of the liquid; HW height of the well. The diagram is not drawn to scale. [16].

Acquisition and handling of image data.

Videos. 40 videos were obtained. For each video in gray scale, the frames were downloaded using a frame-by-frame mode in the image processing unit. Videos contained over 750 frames (a recording of 30-60 seconds), each one with dimensions x=320, y=240 pixels, and each frame having an exposure time of approximately 0.033 sec. Regardless of the length of the video, the processed frames were always the same for each experiment and were taken starting on frame 100 which corresponds to 3.33 sec after the beginning of the video for repeated measures and time series and at frame 500 which corresponds to 16.5 sec, for video edition and processing. There are 4 control videos without antibiotic and 36 videos, each one having a dilution of the antibiotic, 18 of them are assays without bacteria and 18 of them are assays with bacteria. Although each video was processed individually, the results are sometimes combined in groups of low, intermediate and high antibiotic concentration. The Minimal Inhibitory Concentration (MIC) is ≤1 µg/mL according to CLSI [1], and 0.488 µg/mL as obtained in this work, which is close to the concentration of videos 19 and 20 and is usually included in the intermediate concentration group so as to differentiate the antibiotic effect of the high and low concentration groups.

Selection of the Region of Interest (ROI). Video 1, which was taken from the well that contained culture medium with the smallest amount of added antibiotic, without bacteria, and Video 2 which was taken from the well that contained culture medium, bacteria and the same amount of added antibiotic, were used to construct a spatial-temporal speckle matrix that would be a guide to find the region with the maximal differential activity and would be selected as the region of interest (ROI). In each video, using the program ImageJ and starting with the raster images, each one being a systematic division of the illuminated space, a segment of 11 lines was taken (y=115 to y=125, both inclusive) from 11 successive frames which include a temporal resolution of 0.37 seconds (from 3.33sec to 3.70 sec of each video). Each line spanned from x=0 to x=319. Since this was performed in 11 successive speckle images (frames 100-110), for each "x", there were 121 values (11 "y" values from 11 frames). These 121 values were used to obtain the Standard Deviation (SD) for each one of the 320 "x" values of Video 1 and of Video 2. Fig 2 shows a diagram of the selection of the ROI. Then, the 320 SD values of each video were used to obtain 320 values of Qx_i, as follows:

Equation 1

$$Qx_i = \frac{SDx_i\,(v2)}{SDx_i\,(v1)}$$

Where SDx_i is the Standard Deviation in each "x", i spans from 0 to 319, and $v1$ and $v2$
refer to Video 1 and Video 2, respectively. This yields 320 values for Qx_i, each one derived
from a spatial temporal speckle matrix of 121 values from each video.

The region with the highest values for Qx_i, was further examined in order to select the ROI.

Fig 2. Selection of the region of interest (ROI). Schematic representation of the
quantitative selection of the ROI.

Image-ROI video edition and processing by the method of Generalized
Differences. In each of the 40 videos, the region of interest (ROI) was extracted from 80
successive frames. A region with coordinates $x282$-$x311$; $y105$-$y135$, was extracted which
resulted in a square matrix with 30x30 pixels, from the 80 frames. Then, using the program
ImageJ (a public domain Java image processing program developed at NIH, USA), a new
video was constructed with the 80 selected regions, generating an uncompressed digital
"flip book animation" (Fig 3), at a frame rate of 7fps. Each new edited video was processed
with ImageDP, according to previous studies [16], obtaining a value of A which is a
measure of the Biospeckle activity of the sample, as captured in each video. Briefly, Eq (2)
was developed as an algorithm of averages that analyzed all the frames in one video
sequence. A is a measure of mean intensity (I), being expressed as an averaged mean
intensity by the method of Generalized Differences:

Equation 2

$$A = \sum_{n=1}^{N} \langle |I_n - I_{n+1}| \rangle \Big/ N - 1$$

where $|.|$ is the absolute value, $<.>$ is the mean value, I is intensity in gray scale and N
represents the total number of frames analyzed, which in this case is 80. The mean intensity
of the absolute value of a subtraction pixel by pixel of whole successive images is averaged
to obtain A, when a video has 80 images, there will be 79 mean differences (N-1). With this
algorithm, successive images that were constructed with a square matrix of 30x30 pixels
are subtracted, transformed to the absolute value of the difference, the mean for the
absolute values of the intensities of the differences is obtained and finally the means are
averaged. It should be noted that with this algorithm, the final number corresponds to the
average of means. In the case of Eq (2), this was achieved with a free software program,
Image Delta Processor (named as ImageDP), written for this purpose in Java. ImageDP is
available on line at http://bigjocker.com/biospeckle/ and it can be used according to the
instructions of the ImageDP Manual (S1 Text). Thus for each assay, a video was
constructed with ImageJ and processed with ImageDP. The 40 values of A obtained were
used to fit a linear regression and also combined in groups of low, intermediate and high
antibiotic content as described above.

301
Fig 3. Image-ROI video edition and processing. Schematic representation of the selection
303 and punching out of a ROI, construction of a new video with ImageJ and processing with
304 ImageDP.
305
Selection of the pixel for the time series. The region with the highest values for
Qx_i, was further examined in order to select a value of *"x"*, which would define one of the
coordinates for the pixel to be selected. The other coordinate was chosen as $y=120$ because
it represents the middle of the y-scale of the frame and also of the constructed matrix
($y=115$ to $y=125$). The chosen pixel was ($x=303$, $y=120$) and it was used to develop the
time series for each of the 40 videos. In each one of the 40 videos, this pixel lays within the
region with the least intensity (I), making it possible to detect the Qx_{303}, as one of the
highest among all the Qx_i values.
314
Construction of the time series. SAGA (System for Automated Geoscientific
Analysis) was used for the construction of a time series of each of the 40 videos. The
"Shapes" Module was used (Shapes points, Convert table to points and Shapes Grid, Add
grid values to points) of the Coordinate ($x303,y120$). Each time series had a length of 253
frames which corresponded with a time period of approximately 8 seconds. The time series
were processed individually with the statistics tools.
321

Statistics tools applied on the time series for repeated-measures analysis.

324
Normality test: Shapiro Wilk. For the Shapiro Wilk Normality test, the H_0 states that
the population is normally distributed [23]. If the p-value is less than 0.05, the H_0 is
rejected and there is evidence that the data are not from a normally distributed population,
accepting the alternative hypothesis (H_1). This test was performed in R and R commander.
Since this test is sensitive to the number of values it was performed with the first 50 values
of the time series.
331
Linear Regression. In this case the H_0 states that there is no significant linear
correlation between the dependent and the independent variable, p-value greater than 0.05.
If the p-value is less than 0.05, H_0 is rejected; hence there is a significant relationship
between the variables in the linear regression model. However, linear regressions are based
on assumptions of homoscedasticity and independence that will be tested below. This test
was performed in R and R commander.
338
339 **Homoscedasticity / Heteroscedasticity: Breusch-Pagan.** In this test the H_0
340 states that there is homoscedasticity, p-value greater than 0.05; therefore, a p-value less
341 than 0.05 is indicative of heteroscedasticity [24]. The test was performed in R and R
342 commander.
343
344 **Independence / Autocorrelation: Durbin-Watson test.** The H_0 (p-value greater
345 than 0.05) states that the errors are serially uncorrelated against the H_1 (p-values less than

346 0.05) that they follow a first order autoregressive process. Furthermore, for the D statistic
347 "small" values or values close to 0, are indicative of positive correlation [25,26]. This test
348 was performed in R and R commander.
349

350 **Autocorrelation: ACF and PACF.** For a given time series, the ACF
351 (Autocorrelation Function), the PACF (Partial Autocorrelation Function) as well as the run
352 sequence plot of the data as a function of time, were obtained [27]. This test was performed
353 in R and R commander.
354

Auto Regressive Integrated Moving Average (ARIMA) Models. The
$(p,d,q)(P,D,Q)_m$ terms of the ARIMA models were selected according to the Box and
Jenkins methodology [27]. Briefly, for a given time series, the run sequence plot is
examined to see if it is stationary and if it has trend and seasonality. The ACF and PACF
are examined to evaluate if the stochastic process is random and independent (white noise).
If the process is not white noise, the ACF and PACF functions are evaluated in order to
identify the type of temporal autocorrelation, as follows: the ACF is examined to see if the
data is random, if there is autocorrelation and auto regression and if it does not represent
white noise, also the shape of the decay of the ACF is indicative of the type of model that
should be chosen; the PACF is examined to see the lags that show statistical significance.
The first-difference trend and seasonal transformations are performed in order to obtain
second order stationarity. Usually for a time series that has trend and seasonality the first-
difference run sequence plot will have a mean of the differenced data around zero. Also, the
significant lags of the PACF of the original data will indicate the "AR" (p) term while the
significant lags of the ACF of the differenced data will indicate the "MA" (q) term. Trend
(d) was included starting with the lowest possible value and seasonality $(P,D,Q)_m$ was
selected by a quantitative approach, with the sub index term "m" (the number of periods in
each season) being 11. Although each video was processed individually, the tendency of
the results is sometimes analyzed in groups as described above in the *Videos* epigraph and
in reference to the MIC.
375

376 # Antimicrobial susceptibility testing.
377 Antimicrobial testing methods were performed in order to confirm the susceptibility of the
378 strain to the concentration range of Ciprofloxacin, under the conditions used.   The MIC
379 value was recorded as the lowest concentration of the assayed antimicrobial agent that
380 inhibits the growth of the microorganism.
381

382 **Broth Macro-dilution.** This test was prepared in tubes with 1mL MHB containing
383 serial two-fold dilutions of Ciprofloxacin and 1mL of MHB containing bacteria at 0.5 Mc
384 Farland Standard. The microorganism was allowed to grow for 24 hours at 36°C then the
385 Absorbance at 625nm was taken.
386

Agar dilution spot test. The bacterial strain was grown in MHB up to 0.5 Mc Farland
Standard and plated on the surface of Mueller-Hinton Agar (MHA), Becton, Dickinson and
Company. Spots of 5μL serial two-fold dilutions of Ciprofloxacin were placed on the
surface [28]. The plates were incubated for 24 hours at 36°C and inhibition of growth was
evaluated, measuring the diameter of the halo under the magnifying microscope.

RESULTS

The present paper aims at applying the Biospeckle methodology already designed [16] to a
bacterial culture in the wells of VDRL plates, with the purpose of performing an
exploratory assay which could eventually set the basis for a fast antibiogram.
398

Selection of the Region of Interest (ROI). Developing the Biospeckle
methodology with bacteria has to take into account the size and the density of the
microorganism, as well as the motility and Brownian motion. Moreover, the assumption of
homogeneity cannot be applied directly to Biospeckle laser monitoring of microorganisms
in a medium [29]. Since there are variations of homogeneity in the illuminated sample,
particularly in a culture medium with cells, it was necessary to select a region of the frames
where the activity of the microorganism and the effect of the antibiotic are properly
revealed. It should be noted that the laser illuminated the plate with an angle of incidence of
72° in order to obtain different intensity regions within each image. The same illumination
was applied to all the samples.
409

A line at the middle of the frame (y120) was selected for all the values of "x" (x0-x319). Fig
4a and Fig 4b show the intensity (I) for all the pixels in that line, for frame "100", taken
from both, Video 1 (culture medium with the lowest added antibiotic concentration, without
bacteria) and Video 2 (culture medium with the lowest added antibiotic concentration, with
bacteria) respectively, with both showing a similar profile and dispersion pattern, the
double arrow points at the region x282-x311 at the middle of frame "100" (y120), which
was selected as a high activity region as is shown later. The usual Biospeckle practice is to
take the region with the mean intensity for analysis. However, as Fig 4a and Fig 4b show,
in this case the mean intensity region also has the highest slope, implying that it is a region
with a high variation. Nevertheless, all the regions (from pixel x=0 to pixel x=319) were
evaluated in order to find the pixels where the presence of bacteria was best revealed. Fig
5a shows the surface plot for the segment x0-x319;y115-y125 of Video 1 (without bacteria)
and Fig 5b shows the same area of Video 2 (with bacteria).
423

Fig 4. Intensity (I) of pixels at line y120 of Videos 1 and 2. Intensity of pixels at line x0-
319 y120 of
A. Video 1 which was taken from an assay that contained culture medium (MHB)
without bacteria (WOB) and the least amount of Ciprofloxacin (7.63E-04 μg/mL in
the well), and
B. Video 2 which was taken from an assay that contained culture medium (MHB) with
bacteria (WB) and the least amount of Ciprofloxacin (7.63E-04 μg/mL in the well).
432

434 **Fig 5. Surface plot of a segment of a Video.** Surface plot of the segment x0-x319 y115-
435 y125 of
436   A. Video 1 (culture medium (MHB) without bacteria and the least amount of
437   Ciprofloxacin, 7.63E-04 μg/mL in the well) and

438   B. Video 2 (culture medium (MHB) with bacteria and the least amount of
439   Ciprofloxacin, 7.63E-04 µg/mL in the well).

440
441

For the selection of the ROI, the data of Video 1 and Video 2 were used applying Equation
1, as described in Materials and Methods (*Selection of the Region of Interest (ROI)* and Fig
2). Fig 6 shows the result of this spatial-temporal analysis in a plot with the x-axis crossing
at a value of 1 in order to distinguish the areas where the SD of Video 2 (with bacteria) is
higher than the SD of Video 1 (without bacteria). A region of higher differential Biospeckle
activity with coordinates $x282$-$x311;y115$-$y125$ is shown between the arrows. The purpose
of this spatial-temporal analysis was to select the ROI through a quantitative method,
avoiding qualitative approaches.

450
451

Fig 6. Spatial temporal analysis of the SD. Application of Equation 1 to the data of Video
1 (MHB without bacteria and Ciprofloxacin 7.63E-04 µg/mL) and Video 2 (MHB with
bacteria and Ciprofloxacin 7.63E-04 µg/mL). The SD was taken for each column of pixels
in "*x*" for $y120$-115 in 11 frames. Equation 1 was applied and the quotient of
Video2/Video1 is plotted with the x-axis crossing at a value of 1 in order to distinguish the
areas where the SD of Video 2 (with bacteria) is higher than the SD of Video 1 (without
bacteria).

Generalized differences as descriptor of Biospeckle activity of the edited videos.
Having identified the area of high differential activity, a segment of 30x30 pixels ($x282$-
$x311; y105$-135) was extracted from each video. Figs 7 and Fig 8 show the selected areas
for Videos 1 and 2, respectively. It should be noted that although this selected area has the
lowest intensity (*I*) value within the frame, it has a higher differential SD, suggesting that it
is in this region that the presence of bacteria and the effect of the antibiotic could be
detected. Hence in each case (Videos 1-36) a new edited video was constructed as shown in
Fig 3, with the segment of maximal differential activity. For this purpose, the region of
interest (ROI) was extracted from 80 successive frames, a video was constructed and each
new video was processed with ImageDP, according to previous studies [16], obtaining a
value of *A* which is a measure of the Biospeckle activity of the sample, as captured in the
edited video. S1 Data shows the values of *A* for the videos with and without bacteria,
however videos 31/32 to 35/36 were not included due to a distortion caused by the higher
antibiotic concentrations. This type of behavior was also observed in [16]. Fig 9 shows the
linear regression for the values of *A* taken from S1 Data, as a function of the logarithm of
the concentration of Ciprofloxacine. Although the curve with bacteria shows a clear
tendency to decrease as the antibiotic concentration rises, it also shows a low determination
coefficient with a high dispersion of the points.

478
479

480   **Fig 7. Surface plot of the selected ROI of Video 1.** Surface plot of the segment $x282$-$x311$
481    $y115$-$y125$of Video 1 (culture medium (MHB) without bacteria and the least amount of
482   Ciprofloxacin, 7.63E-04 µg/mL in the well).

483
484

Fig 8. Surface plot of the selected ROI of Video 2. Surface plot of the segment $x282$-$x311$ $y115$-$y125$ of Video 2 (culture medium (MHB) without bacteria and the least amount of Ciprofloxacin, 7.63E-04 µg/mL in the well).

Fig 9. **Linear regression for the values of A as a function of the logarithm of the concentration of Ciprofloxacin.** WB, with bacteria; WOB without bacteria.

Due to the dispersion of the data, the videos with and without bacteria, were organized in three groups, low (7.6E-04 to 1.2E-02 µg/mL), intermediate (2.4E-02 to 3.9E-01 µg/mL) and high concentration (7.8E-01 to 1.25E+01 µg/mL) of Ciprofloxacin. It should be noted that for the purpose of comparison, all groups have the same number of videos (5 videos each group). However, within each group the range of the antibiotic concentration may have up to two orders of magnitude. According to CLSI, the MIC for Ciprofloxacin on *E. coli* is ≤1µg/mL [1]. In this work the MIC of Ciprofloxacin on the used strain was determined at 0.488 µg/mL by two methods, Broth Macrodilution and Agar Dilution as shown below. This is close to the antibiotic concentration of videos 19 and 20 (0.391 µg/mL), both included in the intermediate group.   S1 Data contains the the data of the edited videos. Odd and even numbered videos 31 to 36 were excluded from this analysis because the high antibiotic concentration caused a distortion in the Biospeckle pattern. This effect had already been described when benznidazol was the test drug on *Trypanosoma cruzi* [16]. In each group, the Biospeckle activity was obtained with ImageDP as an A value, then the the mean and the SD of the groups, were calculated and are shown in Fig. 10. As Fig 10 shows, the Biospeckle activity of the videos without bacteria is similar in the three groups, with a slight tendency to increase at the higher antibiotic concentrations. In the group of videos with bacteria, the Biospeckle activity is higher at the lower antibiotic concentrations and drops as the concentration increases. The Biospeckle activity of the bacteria at the highest concentrations of antibiotic approaches the Biospeckle activity of the culture medium without bacteria and the same antibiotic concentrations.

Fig 10. Activity of the Biospeckle pattern of the edited videos. Each video was edited by punching out the ROI and constructing a new video with ImageJ, which was processed with ImageDP, obtaining in each case, a value of A which represents a measure of the activity of the Biospeckle pattern. The results are organized in groups according to the concentration of Ciprofloxacin, as follows, low (7.6E-04 to 1.2E-02 µg/mL), intermediate (2.4E-02 to 3.9E-01 µg/mL) and high (7.8E-01 to 1.25E+01 µg/mL) concentration of Ciprofloxacin; WOB without bacteria, WB with bacteria.

All this indicates that, if the ROI is properly selected, and new videos are constructed with ImageJ and analyzed with ImageDP, it is possible to distinguish the videos with bacteria from those without bacteria and the effect of the antibiotic. Moreover, if the videos are analyzed in separate groups, above and below the MIC, they show a difference in the activity of the Biospeckle pattern with different values, when the antibiotic is present at concentrations above and below the MIC.

Diagnostic statistical tests for repeated-measures analysis (linear regression model, normality, homoscedasticity and independence of the time series of each video). In an attempt to improve the detection of the presence of bacteria and the effect of the antibiotic, a time series was extracted within the selected ROI of each of 38 videos (odd numbered videos 1-35; even numbered videos 2-36 and control videos A and B). The coordinates were chosen from a high differential activity pixel within the selected ROI. In all cases the time series was extracted at pixel $x303;y120$. Statistic tests were performed for each time series.

Figs 11- 14 show the results of diagnostic statistical tests, for the time series of each video and two control videos (see S2 Data). In this case each test was performed on the individual time series and the results were considered in groups for the purpose of analysis and discussion.

In the assays with culture medium, bacteria and low antibiotic concentrations (7.6E-04 to 2.4E-02, even numbered videos 2-12) below the MIC, the p-value for a linear regression is greater than 0.05 in 6 out of 7 assays, so there is not a significant relationship between the dependent and independent variables, in a linear regression model; there is non-normality (Shapiro-Wilk test) in 5 out of 7 assays, there is heteroscedasticity (according to the Breusch-Pagan test) in 6 out of 7 assays and positive autocorrelation (Durbin-Watson test) in all of them. Therefore, the presence of bacteria with the least antibiotic effect, promotes a Biospeckle pattern in the time series that has positive autocorrelation and is mainly heteroscedastic, non-linear with a non-normal distribution of the data. In this case, the non-linear regression is compatible with the heteroscedasticity and non-independence (autocorrelation).

In assays with culture medium, bacteria and the antibiotic that approaches the MIC from the lower concentrations (4.88E-02 to 7.81E-01, even numbered videos 14 to 22) and includes it, there is a mixed behavior which suggests a transient condition between the low and the high concentration groups, in the linear regression and homo/heteroscedasticity, with non-normal behavior and positive autocorrelation in all of the assays.

In assays with culture medium, bacteria and the antibiotic at concentrations above the MIC (1.56E+00 to 1E+02, even numbered videos 24 to 36), in 5 out of 7 time series, there is a significant relationship between the dependent and independent variables in a linear regression model, being similar to the culture medium without bacteria and with antibiotic and contrary to the low concentration group (bacteria with low antibiotic concentrations). Furthermore, this high concentration group shows mainly heteroscedasticity (5 out of 7) and a normal behavior (5 out of 7) with all of the time series having positive autocorrelation. This is an unexpected combination of results because the assumptions for a linear regression are not satisfied.

Considering all the assays with culture medium with different concentrations of Ciprofloxacin (7.6E-04 to 1.0E+02, odd numbered videos 1-35) without bacteria, in 13 out of 19 time series the p-value for a linear regression is less than 0.05 so there is a significant

relationship between the dependent and independent variables. In 14 of the 19 videos, there is homoscedasticity according to the Breusch-Pagan test and in all of them there is a non-normal distribution as indicated in the Shapiro-Wilk test and positive autocorrelation with the Durbin-Watson test.   Therefore, the time series of the culture medium without bacteria and different concentrations of the antibiotic have a Biospeckle pattern that have a non-normal distribution with positive autocorrelation and are mainly homoscedastic and linear. All of the time series that follow a linear regression are expected to be homoscedastic and independent according to the assumptions.

Fig 11. Shapiro-Wilk Normality Test. This test was performed on the time series of all of the videos with the values of pixel $x303;y120$ of the first 50 frames (frames 100-150) because it is sensitive to the number of input values. Note: black indicates that the H_0 is accepted and there is normal distribution of the data; white means that the H_1 is accepted and the data are not normally distributed.

Fig 12. Adjustment to a linear regression. This test was performed on the time series of all of the videos on the values of pixel $x303;y120$ over 253 frames (frames 100-353). Note: black indicates that the H_0 is accepted and there is no relation among the variables; white means that the H_1 is accepted and there is a linear relation among the dependent and independent variables.

Fig 13. Breusch-Pagan homoscedasticity / heteroscedasticity test. This test was performed on the time series of all of the videos on the values of pixel $x303;y120$ over 253 frames (frames 100-353). Note: black means that the H0 is accepted and there is homoscedasticity; white means the H_1 is accepted and there is heteroscedasticity.

Fig 14. Durbin-Watson test for independence. This test was performed on the time series of all of the videos on the values of pixel $x303;y120$ over 253 frames (frames 100-353). In all cases the p-values are less than 0.05, rejecting the H_0 which states that the errors are serially uncorrelated and accepting the H_1 which states that the errors follow a first order autoregressive process. Small values of the D statistic indicate that successive error terms are close in value to one another, or positively correlated. Note: gray means there is positive autocorrelation, which was found in all the time series.

Autocorrelation Function (ACF), Partial Autocorrelation Function (PACF) and Selection of ARIMA models. Since the results of the Durbin-Watson test (Fig 14) showed that all the time series were non-independent with positive autocorrelation, the ACF and PACF were performed for each one of the 40 time series, confirming the autocorrelation. As an example, the run sequence plot, ACF and PACF for the time series of Video 1 (culture medium with the lowest added antibiotic concentration, without bacteria) and Video 2 (culture medium with the lowest added antibiotic concentration, with bacteria), are shown in S3 Data. Model adjustment was individually

625 made in each of the 40 videos but the discussion is performed considering the groups as
626 above.
627

628 Controls A (culture medium with NaCl 0.9%, without bacteria) and C (culture medium
629 without bacteria and without NaCl) were used as an orientation for model selection. Once it
630 was determined that there is autocorrelation, an ARIMA model was explored for the time
631 series of pixel x303;y120 of Control A, according to Box-Jenkins approach. A process to
632 obtain the best ARIMA $(p, d, q)(P,D,Q)_m$ model was worked out as follows.
633

634 The time series was examined for Control A. The run sequence plot of the time series
635 shows that it is non-stationary (Fig 15), the ACF (Fig 16) shows that the data does not
636 represent white noise, the data are not random, the observations are related to adjacent
637 observations and there is a strong autocorrelation and auto regression. The PACF function
638 (Fig 17) shows statistical significance for lags 1 and 2.
639

641 **Fig 15. Run sequence plot of Control A** (culture medium with NaCl 0.9%, without
642 bacteria).
643

645 **Fig 16. Autocorrelation Function (ACF) of Control A** (culture medium with NaCl 0.9%,
646 without bacteria).
647

649 **Fig 17. Partial Autocorrelation Function (PACF) of Control A** (culture medium with
650 NaCl 0.9%, without bacteria).
651

653 A first-difference trend transformation was performed. The run sequence plot of the
654 differenced data (Fig 18) shows that the mean of the differenced data is around zero, with
655 the differenced result less autocorrelated than the original data. The ACF of the differenced
656 data (Fig 19) and the PACF (Fig 20) both with a 95% confidence band show that only the
657 autocorrelation at lag 1 and possibly at lag 2 are significant, suggesting that the differenced
658 data are stationary.
659

661 **Fig 18. Run sequence plot of first-difference trend transformation of Control A**
662 (culture medium with NaCl 0.9%, without bacteria).
663

665 **Fig 19. Autocorrelation Function (ACF) of first-difference trend transformation of**
666 **Control A** (culture medium with NaCl 0.9%, without bacteria).
667

669 **Fig 20. Partial Autocorrelation Function (PACF) of first-difference trend**
670 **transformation of Control A** (culture medium with NaCl 0.9%, without bacteria).
671

Similarly, a first-difference seasonal transformation was performed, showing stationarity (Figs 21-23).

Fig 21. Run sequence plot of first-difference seasonal transformation of Control A (culture medium with NaCl 0.9%, without bacteria).

Fig 22. Autocorrelation Function (ACF) of first-difference seasonal transformation of Control A (culture medium with NaCl 0.9%, without bacteria).

Fig 23. Partial Autocorrelation Function (PACF) of first-difference seasonal transformation of Control A (culture medium with NaCl 0.9%, without bacteria).

Since the ACF plot of the original data shows that the autocorrelations are very strong and positive and decay very slowly, the time series was identified as an "AR(p) signature". The PACF shows significance at lags 1 and 2 indicating that this is possibly an Auto-regressive, AR(2) signature. Since the ACF of the differenced data shows significance at lag 1 and possibly at lag 2, a Moving Average, MA(2) model is suggested.

Furthermore, these examinations indicate that the time series has trend and seasonality. Therefore, a term for trend (d) as well as the terms for seasonality $(P,D,Q)_{11}$ were included. It should be noted that this rationale was used as a general orientation to select a model that could fit the largest number of videos obtained either with or without bacteria. Therefore, a $(2,1,2)(2,1,0)_{11}$ model was thought to be close enough for this purpose and was chosen as the first selected model which has the further advantage of having been widely used. Furthermore, a similar analysis of the time series for Videos 1 and 2 give the same orientation (see S3 Data).

A similar process was carried out with Control C. The results are shown in Figs 24 to Fig 32. The run sequence plot (Fig 24) shows non-stationarity. The ACF (Fig 25) shows an "AR(p) signature" for which an AR(4) signature was suggested from the PACF (Fig 26), and an MA(1) was suggested from the ACF plot of the differenced data. Since stationarity is achieved with the differenced data, trend and seasonality were included in the model. The AR(4) can also be observed from the PACF of the differenced data. Taking into account that the most suitable ARIMA models are those with the lower number of parameters possible, a term (1,1,1) was assayed for variations in seasonality. The best fit was approached with a model $(4,1,1)(1,1,1)_{11}$. Again, this rational was thought to be close enough for the purpose of having a starting criterion to guide towards finding the most suitable models.

Fig 24. Run sequence plot of Control C (culture medium without bacteria and without NaCl).

Fig 25. Autocorrelation Function (ACF) of Control C (culture medium without bacteria and without NaCl).

Fig 26. Partial Autocorrelation Function (PACF) of Control C (culture medium without bacteria and without NaCl).

Fig 27. Run sequence plot of first-difference trend transformation of Control C (culture medium without bacteria and without NaCl).

Fig 28. Autocorrelation Function (ACF) of first-difference trend transformation of Control C (culture medium without bacteria and without NaCl).

Fig 29. Partial Autocorrelation Function (PACF) of first-difference trend transformation of Control C (culture medium without bacteria and without NaCl).

Fig 30. Run sequence plot of first-difference seasonal transformation of Control C (culture medium without bacteria and without NaCl).

Fig 31. Autocorrelation Function (ACF) of first-difference seasonal transformation of Control C (culture medium without bacteria and without NaCl).

Fig 32. Partial Autocorrelation Function (PACF) of first-difference seasonal transformation of Control C (culture medium without bacteria and without NaCl).

The different models were evaluated on the individual time series and the results are shown individually and also considered in terms of bacteria in the presence of Ciprofloxacin at low concentration (7.6E-04 to 2.4E-02, even numbered videos 2-12), intermediate concentration (4.88E-02 to 7.81E-01, even numbered videos 14 to 22) and high concentration (1.56E+00 to 1E+02, even numbered videos 24 to 36) of the antibiotic and the culture medium with all the concentrations of Ciprofloxacin (7.6E-04 to 1.0E+02, odd numbered videos 1-35).

Fig 33 shows the results of applying these two models ($(2,1,2)(2,1,0)_{11}$ and $(4,1,1)(1,1,1)_{11}$) to the 38 videos with and without bacteria and varying amounts of Ciprofloxacin. Black boxes symbolize that the model fits the time series, with coefficients that are <1 and with residuals that behave as white noise. Both models tend to fit better the time series for the odd numbered videos (that contain culture medium without bacteria with varying amounts of antibiotic), than the even numbered videos with bacteria and the low and intermediate amounts of antibiotic. Also, the time series from videos with bacteria and the highest amount of antibiotic, adjust to both models as would be expected if the antibiotic tends to affect the bacteria, behaving like the culture medium with antibiotic and without bacteria.

766 Therefore, fitting models $(2,1,2)(2,1,0)_{11}$ and $(4,1,1)(1,1,1)_{11}$ on the time series, it is
767 possible to distinguish among the conditions with and without bacteria and among
768 antibiotic concentrations below and above the MIC.
769
770
771 **Fig 33. Adjustment of ARIMA models $(2,1,2)(2,1,0)_{11}$ and $(4,1,1)(1,1,1)_{11}$.** Each one of
772 these models was performed with each one of the time series that are indicated above the
773 boxes. Black boxes indicates that the model adjusts to the time series with 253 values;
774 white boxes indicates that the model does not adjust to the time series.
775
776 In order to be able to further distinguish the groups of assays, several models were tested
777 separately to find the best fit for the culture medium without bacteria, the affected bacteria
778 and the unaffected bacteria, increasing and decreasing the parameters of the ARIMA
779 models.
780
Increasing the parameters of the ARIMA models. Since the purpose of this
work is to distinguish the presence and absence of bacteria and the effect of the antibiotic,
variations of the model were performed so as to obtain the best fit with the different
conditions. Therefore, in an attempt to detect the bacteria that have been least affected by
the antibiotic and are probably the most active, alternative models were assayed. Since
these time series have an "AR(p) signature", different models with increasing values of (p)
were tested, starting from AR(2), MA(2). Fig 34 shows the best fitted model for this
purpose, $(7,1,2)(1,0,1)_{11}$, adjusting to the time series of bacteria in the presence of the least
amount of antibiotic. It should be noted that in the case of the even numbered videos (with
bacteria) a tendency that fades off as the antibiotic concentration rises from low to
intermediate is seen, and it is not an "all or nothing" result, again resembling a transient
condition (see Figs 12 and 13).
793
794
795 **Fig 34. Adjustment of ARIMA model $(7,1,2)(1,0,1)_{11}$.** The model was performed with
796 each one of the time series that are indicated above the boxes. Black boxes indicates that
797 the model adjusts to the time series with 253 values; white boxes indicates that the model
798 does not adjust to the time series.
799
800
Decreasing the parameters of the ARIMA model. Since it is recommended that
the most suitable ARIMA models are those with the lower number of parameters, a
decrease in the numbers of the model was assayed, with variations in seasonality. Thus, the
model **(1,1,1)** was assayed, combined with the seasonal parameters $(P,D,Q)_{11}$ that were
permutations of 1 and 0 (0,0,0); (0,0,1); (0,1,0); **(0,1,1)**; **(1,0,0)**; **(1,0,1)**; (1,1,0); **(1,1,1)**. Fig
35 shows only the results for 4 selected models which are highlighted in this text. All of
them are able to detect the bacteria affected by the higher antibiotic concentrations. The
other combinations (not highlighted) do not adjust completely to the time series.
809
810
811 **Fig 35. Adjustment of ARIMA models $(1,1,1)(1,0,1)_{11}$; $(1,1,1)(0,1,1)_{11}$; $(1,1,1)(1,1,1)_{11}$**
812 **and $(1,1,1)(1,0,0)_{11}$.** Each one of these models was performed with each one of the time

series that are indicated above the boxes. Black boxes indicates that the model adjusts to the time series with 253 values; white boxes indicates that the model does not adjust to the time series.

Model (1,1,1)(1,0,0) does not fit any of the assays of culture medium without bacteria and model (1,1,1)(1,0,1) fits only one time series in this group. These models, especially the latter, are able to detect the bacteria affected by the higher antibiotic concentrations. Furthermore, model (1,1,1)(1,0,1) has a pattern that is similar to the result obtained with Shapiro-Wilk Normality test (see Fig 11). In general, the first 3 models tend to have an adjustment with the time series of assays with bacteria and the highest antibiotic concentration. In other words, these models have the tendency to distinguish the bacteria that are affected by the concentrations of Ciprofloxacin that are above the MIC that also show a normal distribution of the data.

Antimicrobial susceptibility testing. In order to test the susceptibility of the *E. coli* K-12 strain HfrH to the Ciprofloxacin i.v. infusion used in this work, classic broth and agar dilution methods were performed to determine the lowest concentration of the assayed antimicrobial agent (MIC) that, under defined test conditions, inhibits the growth of the bacterium. The broth macro-dilution method showed a MIC value of 0.488μg/mL (S4 Data) which corresponds with a concentration that approaches the assays of Videos 19 and 20 (0.39 μg/mL). Furthermore, Fig 36 shows the results of the mean and SD of the growth of the bacteria as determined by its Absorbance at 625nm arranged in three groups of low, intermediate and high antibiotic concentration that are similar to those of Fig 10. The low concentration group goes from 1.53E-02 to 2.44E-01, the intermediate concentration group goes from 4.88E-01 to 7.81E00 (including the MIC) and the high concentration group goes from 1.56E+01 to 2.50E+02.The results obtained with agar dilution indicate (see S4 Data) that 5 μg of Ciprofloxacin have an inhibition diameter that is ≥21mm as required in the CLSI standards [1] for a susceptible strain. Furthermore, in this agar dilution test [28], a concentration of 0.488 μg/mL, which again is close to the assays of Videos 19 and 20 (0.39 μg/mL) of the Biospeckle technique, is the minimal concentration that promotes a precise inhibition halo which can be detected and measured under the magnifying microscope. All this indicates that Ciprofloxacin has the expected anti-microbial activity on the selected strain of *E. coli* K-12 and that the antimicrobial susceptibility tests show results that may be comparable to those obtained with Biospeckle under different analysis methods.

Fig 36. Absorbance at λ_{625nm} of the macro-broth dilution test. The macro-broth dilution test was carried out as described in Materials and Methods with bacteria and varying concentrations of Ciprofloxacin, obtaining in each case, a value of Absorbance which represents a measure of the growth of the microorganism. The y axis expresses the CFU/mL. The results of the assays with bacteria in MHB and varying antibiotic concentrations are organized in groups that are similar as the groups of Fig 10 because, due to the experimental design, variations in concentrations are obtained, however, the same groups below and above the MIC are presented, as follows, Group (1) assays with Ciprofloxacin concentration from 3.81E-03 to 3.05E-02; Group (2) assays with Ciprofloxacin concentration from 6.10E-02 to 9.77E-01; and Group (3) assays with Ciprofloxacin concentrations from 1.95E+00 to 1.25E+02.

DISCUSSION

An assay designed to develop dynamic laser speckle or Biospeckle aims at providing a digital signal which is caught in a video. Biologic materials [29] as well as the illumination in the setup tend to cause inhomogeneities that affect the digital processing of the images. Furthermore, subsurface inhomogeneities have been reported and revealed with laser speckle [30]. These inhomogeneities could interfere if the whole photogram is used for calculations, obscuring the real effect that is to be revealed. Therefore, the selection of a ROI is an important step prior to the analysis of the results. Braga *et al* [29] have addressed the problem of inhomogeneities by selecting pixels at random, whereas Meschino *et al* [18] have performed a Self-Organizing Map based on Artificial Neural Networks. According to those authors, care must be taken so as to select a representative area for calculation. Therefore, in the present work a quantitative approach was developed. The rational indicated that areas of the photogram with high illumination should be avoided because small changes could be obscured within the high intensity (I) peaks. Areas of high variation should also be avoided because the variable background could make the activity of the bacteria undetectable, due to their size, their motility and the high cell density usually required. In this context and with the idea of performing statistical analysis, a low resolution of the camera was selected. The quantitative analysis of the SD of 11 pixels in 11 photograms in terms of a video with bacteria and a video without bacteria, both with the lowest Ciprofloxacin concentration, revealed an appropriate approach to find the ROI which is located in the area of low intensity (I).

After having identified the ROI, three approaches were taken which focus on the evolution of intensity (I) as a function of the time domain. For this purpose the behavior for a length of time of either an area of 900 pixels or of the time series of individual pixels, was examined. In the first approach, after quantitatively identifying the ROI, the segment was punched out of each frame and all the extracted segments were used to edit a new video with 80 segmented frames which corresponds to a time period of over 2 seconds. The number of frames was chosen from previous studies [16] in which it was determined that 80 frames was an appropriate length for ImageDP processing. With the combination of ImageJ for video construction and ImageDP for video processing (Fig 2), it was possible to obtain a linear tendency which shows that the mean intensity (A) of the Biospeckle pattern of the bacteria decreases as the antibiotic concentration increases. However, since there is a high dispersion of the points, the analysis was improved organizing the results within groups above and below the MIC. For the other two approaches, a pixel with a high differential activity was chosen within the selected ROI, for the extraction of a time series with a length of over 8 seconds and then analyzing each one either through diagnostic statistical tests or by adjusting an ARIMA model for each time series.
The diagnostic statistical tests were chosen in reference to linear regressions. In other words, a linear regression model and three of its principal assumptions (normality, homoscedasticity and independence) were tested on the time series as a repeated-measures analysis. Only the time series of the least affected bacteria (low Ciprofloxacin concentration) behaved in an expected manner, being non independent and mostly non-linear, non-normal and heteroscedastic. On the other hand, the time series of the most

affected bacteria (higher Ciprofloxacin concentration) are non-independent and tend to be
linear, normal and heteroscedastic; and the time series of the culture medium without
bacteria (all Ciprofloxacin concentrations) are non-independent and non-normal and tend to
be linear and homoscedastic. Therefore, adjustment to a linear regression identifies both,
the culture medium and the most affected bacteria, normality identifies the most affected
bacteria, and heteroscedasticity-homoscedasticity distinguishes the presence-absence of
bacteria, respectively.
913
Through the use of ARIMA models, we explored the relationship between the activities of
the Biospeckle pattern of *E. coli* K-12 as it becomes increasingly affected by Ciprofloxacin.
Thus, the activity of the Biospeckle pattern was modeled as a univariate ARIMA with a
seasonal component which indicates that the values not only depend on their own past
values and random errors, but also they depend on the values of a previous cycle. Based on
the ACF and PACF we fitted several ARIMA models with varying AR and MA orders. In
the fitting process, the AR and MA coefficients were estimated using the log-likelihood
method. The residuals were further inspected for autocorrelation through ACF and PACF.
Models with autocorrelated residuals or with coefficients >1, were discarded. ARIMA
models were adjusted showing that a $(1,1,1)(1,0,1)_{11}$ model identifies the most affected
bacteria (higher antibiotic concentration); a $(4,1,1)(1,1,1)_{11}$ model is able to identify the
time series of the most affected bacteria (higher antibiotic concentration) and of the culture
medium (with varying antibiotic concentrations), indicating that the bacteria, as they
become affected by increasing concentrations of the antibiotic, tend to behave in the same
manner as the culture medium. The time series of the least affected bacteria (lowest
antibiotic concentration) are identified by a $(7,1,2)(1,0,1)_{11}$ model. The non-linear, non-
normal and heteroscedastic behavior of this group is probably responsible for its adjustment
to a model with a relatively high parameter. Thus, with the three methods it is possible to
distinguish the presence and absence of bacteria and also bacteria with antibiotic
concentrations below and above the MIC.
934
Bacteria move with a run-and-tumble navigation. During runs the organism moves
approximately straight and during tumbles, or reorientations at zero speed, the organism
explores other directions [31]. This strategy is affected by the presence of positive and
negative feedbacks. In the presence of a gradient it has been described that there is a non-
normal dynamical structure of the positive feedback and a quadratic non-linearity in the
speed along the gradient [13]. This is in agreement with the results of this work in the case
of the time series of bacteria at low concentrations of Ciprofloxacin. Moreover, in *Bacillus*
subtilis, a motile Gram positive bacterium, it has been shown that under sub-lethal
antibiotic concentrations, the statistics of collectively swarming cells, experiments a
transition from normal to anomalous [32]. The authors conclude that the transition is due to
changes in the properties of the bacterial motion and the formation of a motility-defective
sub-population that self-segregates into regions. This phenomenon suggests a new strategy
bacteria employ to fight antibiotic stress. In the present work this phenomenon could
represent the transient condition of sub-lethal concentrations of antibiotic ranging from
4.88E-02μg/mL to 7.81E-01 μg/mL. This group is flanked on one side by the very low sub-
lethal antibiotic concentrations (7.63E-04μg/mL to 2.44E-02μg/mL) and on the other side
by the high lethal antibiotic concentrations (1.56 μg/mL to 1.00E+02μg/mL). The effect of
Ciprofloxacin at these concentration ranges was confirmed in the antimicrobial

susceptibility tests performed on the bacterial strain by macro-broth and agar dilution
assays by growth inhibition and by the size of the diameter of the inhibition halo
respectively, as established by CLSI [1]. Moreover, the results of these microbiological
tests show a profile that is comparable to the result obtained with the images after 15min of
antibiotic action and with ImageJ-ImageDP processing. In this work, a combination of a
low intensity (I) of the selected region (ROI) and a low resolution of the camera were
adequate to manage the data with several methods. This approach was appropriate for this
case in which the microorganism is a bacterium which has a small size, high motility and is
used at a high cell density. Moreover, all the performed analysis lead to comparable results
possibly due to the quantitative method used for the selection of the ROI, on which the
edition of the new video and the selection of a time series were performed. This allowed
both, the edition and quantification with a "flip-book animation" extracted at the ROI and
the selection of a pixel for the time series construction for diagnostic statistics. The four
evaluation methods used: diagnostic statistical tests, fitting of ARIMA models, ImageJ-
ImageDP processing and antimicrobial susceptibility tests, show similar results, being able
to distinguish among the groups of assays with bacteria and Ciprofloxacin below and above
the MIC. This result opens a possible way to perform a fast method to evaluate antibiotic
susceptibility in control as well as in clinical samples.
971
972

CONCLUSIONS
A Biospeckle laser assay was designed for the evaluation of Ciprofloxacin susceptibility of
a strain of *E. coli* K-12. Digital image processing was performed by three methods that
were comparable to classical antimicrobial susceptibility tests in statistical terms.
Diagnostic statistical tests, fitting of ARIMA models and ImageJ-ImageDP processing
showed results that were comparable to antimicrobial susceptibility tests. This Biospeckle
laser method is a possible starting point for testing body fluids and other samples and it
opens a possible way to perform a fast method to evaluate antibiotic susceptibility.
981

982 # Supporting Information
983
984 **S1 DATA.** Data of the edited videos using ImageJ–ImageDP
985
986 **S2 DATA.** Diagnostic statistical tests (linear regression model, normality,
987 homoscedasticity and independence) of the time series for each video
988
989 **S3 DATA.** Run sequence plot, ACF and PACF for the time series of Video 1 and Video 2.
990
991 **S4 DATA.** Data of the antimicrobial susceptibility tests
992
993 **S1 TEXT.** ImageDP Manual for online processing
994

995 # Acknowledgements
996 We thank Centro Multidisciplinario de Ciencias, Instituto Venezolano de Investigaciones

Científicas, Mérida (IVIC-Mérida), Venezuela. The advice of Francisco A. Andrades-Grassi is gratefully acknowledged.

References

1. CLSI. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fifth Informational Supplement. CLSI document M100-S25. Wayne, PA: Clinical and Laboratory Standards Institute; 2015

2. Waldeisen JR, Wang T, Mitra D, Lee LP (2011) A Real-Time PCR Antibiogram for Drug-Resistant Sepsis. PLoS ONE 6(12): e28528. https://doi.org/10.1371/journal.pone.0028528

3. Hansen WL, Beuving J, Verbon A, Wolffs PF (2012) One-day Workflow Scheme for Bacterial Pathogen Detection and Antimicrobial Resistance Testing from Blood Cultures. J. Vis. Exp. (65), e3254, DOI : 10.3791/3254

4. Liu T, Lu Y, Gau V, Liao JC, Wong PK (2014) Rapid Antimicrobial Susceptibility Testing with Electrokinetics Enhanced Biosensors for Diagnosis of Acute Bacterial Infections. Annals of Biomedical Engineering 42(11) 2314–2321. DOI: 10.1007/s10439-014-1040-6

5. Mohan R, Mukherjee A, Sevgen SE, Sanpitakseree Ch, Lee J, Schroeder ChM, Kenis PA (2013) A multiplexed microfluidic platform for rapid antibiotic susceptibility testing. Biosensors and Bioelectronics 49, 118-125. http://dx.doi.org/10.1016/j.bios.2013.04.046

6. Liu1 Ch-Y, Han Y-Y, Shih P-H, Lian W-N, Wang H-H, Lin Ch-H, Hsueh P-R, Wang J-K, Wang Y-L (2016) Rapid bacterial antibiotic susceptibility test based on simple surface-enhanced Raman spectroscopic biomarkers. Scientific Reports | 6:23375 | DOI: 10.1038/srep23375. www.nature.com/scientificreports/

7. Martí-López L, Cabrera H, Martínez-Celorio RA, González-Peña R (2010) Temporal difference method for processing dynamic speckle patterns. Optics Communications 283, 4972–4977.

8. Sendra H, Murialdo S, Passoni L (2007) Dynamic laser speckle to detect motile bacterial response of Pseudomonas aeruginosa. Journal of Physics: Conference Series 90 (2007) 012064 doi:10.1088/1742-6596/90/1/012064

9. González-Peña RJ, Braga Jr. RA, Cibrián RM, Salvador-Palmer R, Gil-Benso R, San Miguel T (2014) Monitoring of the action of drugs in melanoma cells by dynamic laser speckle. Journal of Biomedical Optics 19(5), 057008. doi: 10.1117/1.JBO.19.5.057008

10. Ramírez-Miquet EE, Cabrera H, Grassi HC, Andrades EdJ, Otero I, Rodríguez D, Darias JG (2017) Digital imaging information technology for biospeckle activity

assessment relative to bacteria and parasites. Lasers Med Sci doi: 10.1007/s10103-017-2256-0

11. Cojoc D, Finaurini S, Livshits P, Gur, E, Shapira A, Mico V, Zalevsky Z (2012) Toward fast malaria detection by secondary speckle sensing microscopy. Biomedical Optics Express 3(5)993.

12. Sendra GH, DaiPra AL, Passoni LI, Arizaga R, Rabal HJ, Trivi M. (2010) Biospeckle descriptors: a performance comparison. Proc. of SPIE 7387, 73871K · doi: 10.1117/12.870682

13. Long J, Zucker SW, Emonet T (2017) Feedback between motion and sensation provides nonlinear boost in run-and-tumble navigation. PLoS Comput Biol 13(3): e1005429. doi:10.1371/journal.pcbi.1005429

14. Drlica K, Zhao X (1997) DNA Gyrase, Topoisomerase IV, and the 4-Quinolones. Microbiology and Molecular Biology Reviews 61(3)377–392

15. Pommier Y, Leo E, Zhang HL, Marchand C (2010) DNA Topoisomerases and Their Poisoning by Anticancer and Antibacterial Drugs. Chemistry & Biology 170. DOI 10.1016/j.chembiol.2010.04.012

16. Grassi HC, García LC, Lobo-Sulbarán ML, Velaásquez A, Andrades-Grassi FA, Cabrera H, Andrades-Grassi JE, Andrades EDJ (2016) Quantitative Laser Biospeckle Method for the Evaluation of the Activity of *Trypanosoma cruzi* Using VDRL Plates and Digital Analysis. PLoS Negl Trop Dis 10(12): e0005169. doi:10.1371/journal. pntd.0005169

17. Wooldridge JM (2013) Econometrics, A Modern Approach, 5[th] Edition. South-Western, Cengage Learning,USA. Library of Congress Control Number: 2012945120, ISBN-13: 978-1-111-53104-1, ISBN-10: 1-111-53104-8

18. Meschino G, Murialdo S, Passoni L, Rabal H, Trivi M (2010) Biospeckle Image Stack Process based on Artificial Neural Networks. Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Conference. DOI: 10.1109/IEMBS.2010.5627620

19. Passoni LI, Dai Pra A, Rabal H, Arizaga R (2005) Dynamic speckle processing using wavelets based entropy. Optics Communications 246(1-3):219-228. DOI: 10.1016/j.optcom.2004.10.054

20. Zdunek A, Adamiak A, Pieczywek PM, Kurenda A (2014) The biospeckle method for the investigation of agricultural crops: A review. Optics and Lasers in Engineering 52(2014)276–285. http://dx.doi.org/10.1016/j.optlaseng.2013.06.017

21. Hayes W (1953) The mechanism of genetic recombination in Escherichia coli. Cold Spring Harb. Symp. Quant. Biol. 18,75-93. doi:10.1101/SQB.1953.018.01.016

22. Isenberg HD (1992) Clinical Microbiology Procedures Handbook. American Society for Microbiology, USA. ISBN 1555810381, 9781555810382

23. Shapiro SS, Wilk MB (1965) An analysis of variance test for normality (complete samples). Biometrika 52(3–4)591–611. doi:10.1093/biomet/52.3-4.591

24. Breusch, T.S. and Pagan, A.R. (1979) A Simple Test for Heteroscedasticity and
Random Coefficient Variation. Econometrica, 47(5)1287-1294.
http://dx.doi.org/10.2307/1911963
25. Durbin, J.; Watson, G. S. (1950). "Testing for Serial Correlation in Least Squares
Regression, I". Biometrika. 37 (3–4): 409–428. doi:10.1093/biomet/37.3-4.409
26. Durbin, J.; Watson, G. S. (1951). "Testing for Serial Correlation in Least Squares
Regression, II". Biometrika. 38 (1–2): 159–179. doi:10.1093/biomet/38.1-2.159
27. Box GEP, Jenkins G (1976). Time Series Analysis: Forecasting and Control.
Holden-Day, USA.
28. Balouiri M, Sadiki M, Ibnsouda SK (2016) Methods for in vitro evaluating
antimicrobial activity: A review. Journal of Pharmaceutical Analysis 6 (2016)71–
79. http://dx.doi.org/10.1016/j.jpha.2015.11.005
29. Braga Jr. RA, González-Peña RJ, Campos Viana D, Pujaico Rivera F (2017)
Dynamic laser speckle analyzed considering inhomogeneities in the biological
sample. Journal of Biomedical Optics 22(4), 045010. 1083-3668/2017
30. Nothdurft R, Yao G (2005) Imaging obscured subsurface inhomogeneity using laser
speckle. Optics Express 13(25)10034.
31. Berg HC, Brown DA (1972) Chemotaxis in Escherichia coli analysed by three-
dimensional tracking. Nature 239(5374)500 – 504. doi:10.1038/239500a0
32. Benisty S, Ben-Jacob E, Ariel G, Be'er A (2015) Antibiotic-Induced Anomalous
Statistics of Collective Bacterial Swarming. Phys. Rev. Lett. 114, 018105. DOI:
10.1103/PhysRevLett.114.018105
1100
1101

Shapiro-Wilk Normality Test

Figure

Linear Regression

A	1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35		B	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36
☐	☐	■	■	☐	☐	☐	☐	☐	■	☐	☐	■	☐	☐	■	☐	■	☐		■	■	■	■	■	☐	■	☐	☐	☐	■	■	■	☐	☐	■	☐	☐	☐

Figure

Breush-Pagan Homoscedasticity Test

Figure

Durbin – Watson Independence Test

Figure

(212)(210)	A	1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35		B	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36
	☐	☐	■	■	☐	■	☐	■	☐	☐	■	■	☐	☐	☐	☐	☐	■	☐		☐	■	☐	☐	☐	☐	☐	■	■	☐	☐	☐	☐	☐	■	■	■	☐	■
	A	1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35		B	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36
(411)(111)	■	☐	■	■	■	■	☐	■	■	☐	☐	■	☐	■	☐	■	☐	■	☐		☐	☐	☐	☐	☐	■	☐	☐	☐	☐	☐	☐	☐	☐	☐	■	■	■	■

Figure

(712)(101)	A	1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35		B	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36
	■	■	□	□	□	□	□	■	□	□	□	□	□	□	■	■	■	□	□		■	■	■	□	■	■	□	□	■	□	■	□	□	□	□	□	□	□	□

Figure

(111)(101)

A	1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35
	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□	■	□	□

B	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36
	□	■	□	□	□	■	□	□	■	□	□	□	□	■	■	■	□	■	■

(111)(011)

A	1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35
	□	□	□	■	□	□	□	■	□	□	□	■	■	□	□	□	■	□	□

B	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36
	□	□	□	□	□	■	□	□	□	□	□	□	□	□	■	□	■	□

(111)(111)

A	1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35
□	□	■	■	□	■	□	■	■	□	□	■	□	□	□	□	□	■	□

B	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36
□	□	□	□	□	■	□	□	□	□	□	□	□	□	□	■	□	■	■

(111)(100)

A	1	3	5	7	9	11	13	15	17	19	21	23	25	27	29	31	33	35
	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□

B	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36
	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□	□	■

Figure

Figure

Figure

ImageJ

ImageDP

A

Figure

A

B

Figure

A

B

Figure

Figure

Figure

Figure

Figure

Figure

Index

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

I

Figure

Figure

Figure

Figure