1

Temporal and spatial selectivity of hippocampal connectivity with sensorimotor cortex during
individual finger movements

Michael J. Miller, Douglas D. Burman

short title: Spatio-temporal properties of hippocampal-sensorimotor connectivity

Department of Radiology, NorthShore University HealthSystem,

Evanston, Illinois, United States of America

Email: MMiller4@northshore.org

Abstract

Cognitive control refers to brain processes involved in regulating behavior according to internal goals or plans. This study examines whether hippocampal connectivity with sensorimotor cortex during paced movements shows a pattern of spatial and temporal selectivity required for cognitive control. Functional magnetic resonance imaging activity was recorded from thirteen right-handed subjects during a paced, non-mnemonic (repetitive tapping) motor task. Direct and inverse connectivity in sensorimotor cortex were examined from psychophysiological interactions (PPI) from hippocampal seed activity during two sets of analyses: the first identified motor interactions relative to rest, whereas the second identified interactions in motor activity between fingers. Finger representations identified in a previous study were used to evaluate patterns of temporal and spatial selectivity in hippocampal connectivity. Changes in the magnitude of connectivity were identified within the sensorimotor representations of the first (index) through third (ring) fingers across time periods when each finger moved; at each finger representation, hippocampal connectivity was greatest when the represented finger was moving, reflecting temporal selectivity for the timing of finger movements. Similarly, the seeds associated with each finger representation differed in their magnitude of connectivity for adjacent finger representations, reflecting spatial selectivity for the moving finger. The patterns of spatial and temporal selectivity of connectivity during volitional movements in this study meets the criteria for cognitive control adapted from oculomotor studies.

Introduction

To move purposefully, one or more signals within our brain direct the motor system to carry out the intended movement. This is the essence of *cognitive control*, the process by which goals or

plans influence behavior. Although various regions of prefrontal cortex have been broadly

implicated in cognitive control (1-5), the source of cognitive influence on the motor system is

still subject to speculation.

Requirements for cognitive control of movements have been demonstrated in oculomotor

studies, particularly those investigating the neural basis of reading (6-8) and other purposive eye

movements (9-11). Such studies emphasize three components: 1) neural activity evident during

conditions that require cognition; 2) spatial specificity, reflecting the spatial goal of the eye

movement (i.e., target location); and 3) temporal specificity, which determines when the

movement occurs.

The frontal eye field (FEF) is involved specifically in voluntary saccades, both in human and

non-human primates (11, 12). Recording activity from nerve cells is generally avoided in

humans, but FEF neurons in non-human primates reflect all three of these properties (13). These

three properties are also evident from deficits following FEF lesions; for example, impairments

in the accuracy and latency for a variety of volitional, but not reflexive eye movements are

observed in both humans (11, 14-20) and non-human primates (21-24).   In humans, lesions of

the FEF and parietal eye field (PEF) differentially affect the selection of saccade targets and their

timing (25); in monkeys, lesions of either area produce modest impairments in the accuracy and

initiation of saccades, whereas combined lesions produce profound impairments (26). These

studies demonstrate that the FEF, singly or in combination with PEF, is critically involved in the

generation of volitional eye movements, providing the timing of purposive eye movements and

their target location.

Analogous to the FEF in the oculomotor system (27), the primary motor cortex is necessary for

volitional movements of individual fingers (28, 29), with short fiber tracts that connect

postcentral with precentral regions of sensorimotor cortex (SMC) providing sensory feedback

required for accurate performance (30). Neural activity in the FEF and sensorimotor cortex both

reflect cognitive input. FEF properties that reflect cognitive input includes activity that predicts

two sequential saccades (31-33), target selection (34-37), covert attention (38-42), and changes

during learning (27, 43). Similarly, cognitive input to SMC is evident from neural changes

during motor learning, both in its response properties (44-49) and changes in connectivity (50-

52). Although cognitive input to both areas can be inferred, direct evidence for the source (or

sources) of this input is sparse. Indirect evidence from neural properties and lesion effects has

implicated the supplementary eye field (SEF) (53, 54), basal ganglia (55), and dorsolateral

prefrontal cortex (56) in the cognitive control of eye movements, and the basal ganglia (57),

cingulate (58), and prefrontal cortex (1, 59-63) for the cognitive control of skeletomuscular

movements.

From this, it is tempting to speculate that the prefrontal cortex regulates finger movements. The

prefrontal cortex has traditionally been associated with cognitive control, yet its connectivity to

the SMC is indirect via the dorsal premotor cortex (2, 64, 65). Prefrontal areas are functionally

coupled to premotor areas when learning movement sequences but not during repetitive tapping

(66, 67), suggesting prefrontal cortex could be involved in cognitive control when learning

complex movement sequences but not during simple movements. In its role, premotor cortex is

analogous to the SEF (27), which is specialized for learning eye movement sequences (27, 68-

70) and resolving conflicts between potential saccade targets (71, 72). Because prefrontal

connectivity to FEF and SMC is indirect and limited to complex movements, cognitive control of

simpler movements mediated specifically through FEF and SMC may arise from a different

source. Thus, a relationship between neural activity and task complexity is unlikely when

studying cognitive control in SMC, even though such a relationship has been used effectively to

study cognitive control elsewhere (4, 73, 74).

The control of finger movements is more complex than eye movements for at least two reasons.

First, the spatial reference for finger movements is different. Whereas eye movements are

encoded from the current position of the eyes within the orbit (retinocentric space), finger

movements are based upon the current position of the body (body-centered space); thus, the

combination of finger muscles required to achieve a goal differs with different body positions.

Second, the load on eye muscles is constant, whereas muscles moving the fingers may require

different forces when moving objects of different resistance. When a task requires the initiation

of finger movements from a set position with minimal load, however, requirements for cognitive

control are the same as for eye movements: the neural mechanism must be active during

volitional movements that require *cognition*, while specifying the timing and goal of the

movement. The goal under these conditions reflects the body part that moves (e.g., hand or

finger), requiring *cortical specificity* for neural activity within the SMC hand representation.

An earlier study examined hippocampal-SMC connectivity within the framework of volitional

movements (75); the current study considers whether this connectivity shows spatial and

temporal selectivity, necessary to specify which finger moves and when. Connectivity associated

with movements of individual fingers is shown to be organized topographically, with

connectivity in the representation of the moving finger differing from adjacent fingers (spatial

selectivity); furthermore, connectivity within each finger representation is greatest when the

finger moves, dropping when movement stops (temporal selectivity). Combined with the earlier

study, hippocampal-SMC connectivity thus fulfills the three conditions of selectivity required for

cognitive control. After considering alternative interpretations, we conclude that hippocampal

connectivity is consistent with cognitive control of motor function, and describe experiments that

could further elucidate this functionality.

112   Materials and methods

113   Subjects

Thirteen right-handed adults from the Chicago metropolitan area participated in the study (ages

24–59, mean=42.3, five females). Experimental procedures were explained to each subject

before obtaining written consent; consent procedures complied with the Code of Ethics set forth

in the Declaration of Helsinki, and were approved by the Institutional Review Board at the

NorthShore University HealthSystem / Research Institute. Consented subjects had no history of

the following: a previous concussion, psychiatric illness, learning disability, attention deficit

disorder, abnormal physical or cognitive development that would affect educational performance,

central neurological disease (or structural pathology), or neurosurgical intervention.

Immediately prior to their fMRI tests, subjects were trained on one cycle of each experimental

task.

125   Experimental tasks

The tasks were described in detail elsewhere (75). Briefly, the visual/motor task used in this

study comprised 6 cycles of a specified sequence of visual and motor conditions. Motor

conditions were preceded by an instruction screen, accompanied by a metronome ticking at 2

beats per second. During the sequence learning task, subjects rehearsed a remembered sequence

of button presses, tapping in synchrony with the metronome; this was followed by a visual

condition, which required fixation at the center without finger movements while a circular

checkerboard flashed at 4Hz. This initial presentation of the visual task ("visA") was followed

by a block of repetitive tapping, subjects repeatedly tapped the same finger on both hands,

corresponding to the button identified on-screen; the index finger ('F1') tapped first, and the

finger increased incrementally every 4s (eight taps). After tapping movements from all four

fingers (F1-F4 during time periods T1-T4, respectively), the visual condition was repeated

("visB").

This report uses two sets of psychophysiological (PPI) connectivity for selectivity analyses; the

first used the visA task condition as baseline (comparing activity during movement and rest),

whereas the second used the T4 period (comparing activity between movement of different

fingers).

MRI data acquisition

Images were acquired using a 12-channel head coil in a 3 Tesla Siemens scanner (Verio). Visual

stimuli projected onto a screen (Avotec Silent Vision) were viewed via a mirror attached to the

head coil, and behavioral responses were recorded by Eprime (Psychology Software Tools, Inc.)

from an optical response box (Current Designs, Philadelphia, PA). Blood-oxygen level

dependent (BOLD) functional images were acquired using the echo planar imaging (EPI)

method, using the following parameters: time of echo (TE) = 25ms, flip angle = 90°, matrix size

= 64 x 64, field of view = 22cm, slice thickness = 3mm, number of slices = 32; time of repetition

(TR) = 2000ms. The number of repetitions for the visual/motor condition was 182. A structural

T1 weighted 3D image (TR = 1600ms, TE = 3.46ms, flip angle = 9 °, matrix size = 256 x 256,

field of view = 22cm, slice thickness = 1mm, number of slices = 144) was acquired in the same

orientation as the functional images.

fMRI data processing

SPM12 software (http://www.fil.ion.ucl.ac.uk/spm) was used to process and analyze data, applying

procedures described elsewhere (75,76). Briefly, images were spatially aligned to the first

volume to correct for small movements, sinc interpolation minimized timing-errors between

slices, and functional images were coregistered to the anatomical image and normalized to the

standard T1 Montreal Neurological Institute (MNI) template, then smoothed with a 10mm

isotropic Gaussian kernel. During analysis, conditions of interest were specified for sequence

learning, visual, and each of the four fingers moving during the repetitive tapping block; global

normalization scaled the mean intensity of each brain volume to a common value to correct for

whole brain differences over time.

The representations for F1 (index finger) through F3 (ring finger) were adapted from an earlier

study (77).

Psychophysiological interactions (PPI)

Preprocessing

Connectivity analysis was carried out using psychophysiological interactions (75), modified to

account for individual variability in connectivity (76). Each voxel in the left and right

hippocampus of the normalized brain was sampled, as delimited by the aal atlas in the WFU

PickAtlas toolbox (http://fmri.wfubmc.edu/software/PickAtlas).

To examine temporal and spatial selectivity, two sets of connectivity analyses were carried out,

differing only in the interaction term used to identify task-specific activity. In the first analysis,

an interaction term specified a greater effect of seed activity during movement of a single finger

than during the visual condition; this identified the effect of movement relative to rest (e.g.,

T1>visA, T1 specifying the time period for tapping the first finger). In the second analysis, the

interaction term specified a greater effect of seed activity during movement of one finger relative

to another finger (e.g., T1>T4). Each set of analyses used a constant time period for the baseline

comparison (visA and T4, respectively), so that differences in connectivity between finger

representations or time periods would not result from different baselines. Following adjustments

for regional differences in timing, a regression analysis identified the magnitude of the BOLD

signal in SMC correlated with the PPI interaction term.

Seed selection

In this study, "functional seeds" were selected (see Burman 2019), i.e., the hippocampal voxel

generating the greatest movement-related connectivity anywhere within SMC. Functional seeds

were selected both for high values of direct connectivity within SMC ("MAX") and for low

values, which reflect inverse connectivity ("MIN"). For both direct and inverse connectivity, a

voxel seed was selected from both the left and right hippocampus, and a conjunction (global)

analysis was used to the magnitude of connectivity.

As noted elsewhere (75), the locus of maximal connectivity associated with functional seeds

often lay outside the finger representations and could occur in either hemisphere. Seeds were *not*

selected based on measurements within a specific finger representation.

Finger topography and spatio/temporal selectivity

A previous study used a variant of our paced repetitive tapping task to identify individual finger

representations (77). This study used high-resolution functional images (1mm isovoxels),

identifying a finger representation for the right hand as the left SMC region where activation

during movement of one finger exceeded the mean activation during movement of the remaining

fingers. The finger representation for the little finger (F4 as labeled in the current study) could

not be mapped, perhaps because some subjects rotated their wrist rather than moving their little

finger to press the key. The high-resolution activation map for each finger representation from

the earlier study was superimposed on a low-resolution (4mm isovoxel) image used in the

current study; low-resolution coordinates for each finger representation are recorded in Table 1.

Activation maps were flipped along the x-axis to map finger representations in the right

hemisphere. Because high-resolution activation did not coincide with the boundaries of the low-

resolution voxels, a few voxels in the current study were associated with more than one finger

representation.

Table 1. Finger representations in left and right sensorimotor cortex. Coordinates are shown based
on 4mm isovoxels for the normalized brain; voxels that appear in more than one finger representation are
shaded in colors to show the dual-finger representation.

Left postcentral	F1	F2	F3
	(-46,-24,54)	(-34,-32,58)	(-34,-32,54)
	(-50,-24,54)	(-34,-32,62)	(-34,-32,58)
	(-42,-24,58)	(-34,-36,62)	(-34,-36,58)
Left precentral			
	(-38,-24,58)	(-30,-32,58)	(-34,-24,58)
	(-42,-20,58)	(-30,-28,62)	(-34,-28,58)
Right postcentral			
	(50,-28,54)	(34,-32,62)	(42,-32,54)

	(42,-28,58)	(34,-36,62)	(38,-32,58)
	(46,-28,58)	(34,-36,66)	(42,-32,58)
	(38,-32,58)	(34,-32,58)	(38,-32,62)
	(42,-32,58)	(38,-32,58)	(34,-32,58)
Right precentral			
	(38,-20,54)	(34,-24,58)	(34,-24,54)
	(42,-20,54)	(38,-24,58)	(34,-20,54)
	(38,-20,58)	(34,-24,54)	(38,-24,54)
	(38,-24,54)		

With the voxels identified for each finger representation (F1-F3), the amplitude of connectivity
from each functional seed (S1-S3) was identified and plotted for each time period (T1-T4).
Paired t-tests from each voxel within a finger representation across time periods provided a
measure of *temporal selectivity* from a given seed; this identified significant differences in the
magnitude of connectivity between two time periods, corresponding to the movement of different
fingers. Similarly, paired t-tests between finger representations from the same seed provided a
measure of *spatial selectivity* within a time period; this identified significant differences in the
magnitude of connectivity among finger representations during each time period, when only one
finger was moving. In this spatial analysis, the same number of voxels near the central
representation of each finger were statistically compared, avoiding those voxels common to both
fingers.

The absolute value of beta estimates for connectivity varied widely between subjects and
analyses, reflecting subject variability in several factors tangential to the analysis of interest (e.g.,

subject variability in spatiotemporal offsets used to best fit SMC activity to the PPI interaction

term, plus variability in response to the baseline condition). Results are reported as percentage

changes in connectivity from the median amplitude across all conditions; the median amplitude

was calculated for each voxel within a finger representation, with the total range specified

separately for temporal and spatial selectivity analyses. A previous study showed connectivity

changes selectively within the SMC hand representation (75); percentage changes in connectivity

measures from the midline thus reflect the relative influence of seed activity on a finger

representation during each condition.

Results

Temporal selectivity of connectivity within a finger representation

Temporal selectivity within a finger representation was demonstrated when the magnitude of

connectivity was greatest during the time period that the represented finger was moving.

Evidence for temporal selectivity for functional seeds is shown in Figure 1 for both the L and R

postcentral gyrus. In the L postcentral gyrus, the PPI interaction term compared seed activity

during finger movements vs. the preceding visual rest condition; significant connectivity was *not*

observed when comparing seed activity during movement of one finger relative to the fourth

finger. In the R postcentral gyrus, the reverse was true; significant connectivity was observed

only when the PPI interaction term compared seed activity during movement of one finger

relative to activity during movement of the fourth finger.

Across all four time periods (T1-T4), Fig 1A shows the hippocampal connectivity maximum

(MAX, solid line) and minimum (MIN, dashed line) for each functional seed (S1-S3) at its

corresponding finger representation (F1-F3). In each hemisphere, MAX connectivity for a seed

was greatest during the movement of its corresponding finger (e.g., S1 in F1 during T1);

typically, the magnitude of MIN connectivity below the mean (inverse connectivity) was greatest

during this same time period.

Hippocampal connectivity at a finger representation was compared between time periods using

paired t-tests (Fig 1A, inset). In both hemispheres, connectivity during the time period where the

represented finger was moving (demarcated by gold ovals) consistently differed from the control

condition (gray ovals). Significant differences with the time period immediately before or after

the movement of a given finger typically involved MIN connectivity (black bars extending one

time segment from a gold oval). Significant differences for MAX connectivity (gray bars) was

observed, however, at F3 in both hemispheres between T3 and T4, the control condition, and

both MAX and MIN connectivity (striped bar) were observed at F1 in the right hemisphere

between T1 and T2. MAX connectivity, however, more often produced changes spanning

multiple time periods.

The MAX and MIN hippocampal connectivity values (Fig 1B) and the MAX-MIN spread (Fig

1C) were compared for each time period. The spread was greatest for each seed/representation

area during the movement of its corresponding finger.

Figure 1. **Temporal selectivity of hippocampal connectivity with SMC. A**: For hippocampal

functional seeds S1-S3, maximum and minimum connectivity with finger representation areas

F1-F3, respectively, in the L and R postcentral gyrus are shown at time periods corresponding to

the movement of fingers 1-4 (T1-T4, respectively). Gold arrows indicate the time period of

movement for the fingers corresponding to F1-F3. Note that both maximum and minimum

connectivity were greatest in each representation area during movement of its corresponding

finger. Task-related connectivity in the postcentral gyrus was calculated relative to either the

visual resting state VisA (L) or the movement of finger 4 (R). Error bars indicate the SEM.

Insets: Paired t-test was used to identify significant differences in direct (gray) and inverse

(black) connectivity among all time periods (T1-T4) at each finger representation F1-F3 in the L

and R postcentral gyrus. Gold ovals indicate the period of movement for the fingers

corresponding to areas F1-F3; silver ovals indicate the control condition for each side. In each

finger representation, significant connectivity differences were typically linked to the period of

movement for its represented finger and to the control state. **B**: Maximum and minimum

hippocampal connectivity are compared for S1/F1 (blue) S2/F2 (red) and S3/F3 (green) at each

time period T1-T4. During each period of movement, both the maximum and minimum

connectivity were greatest in the moving finger's representation. **C**: The Max-Min spread of

connectivity is compared at each time period for S1/F1 (blue), S2/F2 (red), and S3/F3 (green).

For each finger representation, the spread was greatest during the period of the represented

finger's movement.

Spatial selectivity across finger representations

In Figs 2-3, we examined the magnitude of connectivity within a specified time period for each

seed (S1-S3), as well as differences in connectivity between finger representations. Fig 2 shows

the connectivity analysis for the left postcentral gyrus, which used rest as the baseline; Fig 3

shows the same analysis for the right postcentral gyrus, which used the period of movement for

F4 as the baseline (T4).

Fig 2A compares MAX and MIN connectivity within finger representations F1-F3 in the L

postcentral gyrus during each time period (from T1 through T3). MAX and MIN seeds in F1 and

F3 all showed significant connectivity in each time period, whereas significant connectivity in F2

was only observed for the S2 MAX seed during T2. The representation of the moving finger

(i.e., F1 in T1, F2 in T2 and F3 in T3) did not necessarily show the greatest magnitude of MAX

or MIN connectivity.

For each time period, Fig 2B shows differences in connectivity between finger representations;

all significant differences involved inverse connectivity. The finger representation corresponding

to the moving finger is demarcated by the gold bar. Hippocampal connectivity with the

representation of the moving finger was significantly different from one or more neighboring

fingers; connectivity differences between the stationary fingers were less consistent. Figure 2C

compares the MAX-MIN spread at each finger representation across time periods. No consistent

pattern in this spread was observed for any seed, either between time periods or between finger

representations.

Figure 2. **Spatial selectivity of hippocampal connectivity with L postcentral gyrus.**

Movement-related connectivity was calculated relative to the rest condition VisA. **A**: Magnitude

of connectivity in finger representations F1-F3 for functional seeds during time periods T1-T3;

both direct (MAX) and inverse connectivity values (MIN) are shown for each seed. MAX and

MIN connectivity were both significant for F1 and F3, regardless of time period, whereas

connectivity with F2 was significant only for the S2-MAX seed during T2. Error bars indicate

the SEM, asterisks indicate significant connectivity after family-wise error correction ($p<0.05$).

B: Paired t-test identified significant differences between finger representation during a time

period; gold bars indicate the representation area corresponding to the finger moving during each

time period. Significant differences were observed in the magnitude of inverse connectivity

(black) between the representations of the moving finger and an adjacent finger. Differences in

inverse connectivity were sometimes observed between other finger representations, but not

direct connectivity (gray). **C**: Max-Min spread of connectivity was compared for each seed

across time periods and finger representations; solid bars highlight the time period of movement

for each finger representation. No consistent relationship was evident between the Max-Min

spread and the timing of the finger movement associated with a finger representation.

The pattern of MAX and MIN hippocampal connectivity for the R postcentral gyrus differed

from the L postcentral gyrus (see Fig 3). Significant connectivity in each time period (Fig 3A)

was present only in the finger representation for the moving finger. During T1 and T2, both

MAX and MIN connectivity were significant when the represented finger was moving; during

T3, only MIN connectivity (plus MAX for S1) was significant for F3.

Figure 3. **Spatial selectivity of hippocampal connectivity with R postcentral gyrus.** Task-

related connectivity was calculated relative to T4, the time period for moving finger 4; other

conventions as in Fig 2. **A**: Maximum and minimum connectivity of hippocampal functional

seeds. MAX and MIN connectivity were significant and larger within each finger's

representation when the represented finger was moving. **B**: Paired t-test identified significant

differences in direct (gray) and inverse (black) connectivity between finger representations;

striped bars indicate significant differences for both. Significant differences in connectivity were

observed between the representations of the moving finger and an adjacent finger; during T1 and

T3, significant differences were also observed between the non-moving fingers. Spatial

differences in direct connectivity (silver) predominated during T1, whereas differences in inverse

connectivity (black) predominated during T3. **C**: Max-Min spread of connectivity was compared

for each seed across time periods and finger representations. All seeds provided the greatest

MAX-MIN spread in connectivity during a single time period, regardless of the finger

representation.

During T1, only the S3 seed produced significant differences in F1 connectivity from that of

other finger representations (Fig 3B). During T2 and T3, however, direct and/or inverse

connectivity of the moving finger was significantly different from those of neighboring finger

representations for all seeds.   Also, the MAX-MIN connectivity spread was invariably highest

in each finger representation when the represented finger was moving (Fig 3C).

Topography and differentiation between fingers

In previous sections, T4 was used as the baseline for all connectivity analyses in the R

sensorimotor cortex, which reflected differences in seed activity when different fingers are

moving. Using visA as the baseline, by contrast, reflected differences in seed activity when a

finger is moving vs. rest. This raises a question about the spatial resolution of hippocampal

connectivity; can it differentiate between the movement of adjacent fingers?

Fig 4 shows the time course, extent, and topography of hippocampal connectivity delineated by

differential seed activity from movement of adjacent fingers. The time course of the connectivity

signal reflects the time course of the adjacent fingers compared; the time course when comparing

activity from a finger's movement with the preceding finger (top row) is earlier than for the same

finger compared with the subsequent finger movement (bottom row). Although the amplitude of

the connectivity signal may differ, the time course is similar from the left hippocampal seed

(solid line) and the right hippocampal seed (dashed line). Connectivity associated with

movement of each finger shows a topographical arrangement, with connectivity from F1 and F2

located inferior within postcentral gyrus to F4.

Figure 4. **Hippocampal connectivity with R postcentral gyrus based on movements of adjacent fingers.** Connectivity was calculated using functional seeds from the L (solid line) and R hippocampus (dashed line); functional activation across all time periods (dotted line) allows timing comparisons. Connectivity increases from the L and R hippocampal seeds followed the same time course, but connectivity derived from seed activity during movement of one finger vs. the preceding finger (top row) appeared earlier than connectivity derived from movement of the same finger compared to the successive finger (bottom row). Spatial overlap from the top and bottom analyses identified connectivity associated with a specific finger; connectivity associated

382 with the fingers was organized topographically (middle row of images). F1=yellow, F2=red, and

383 F4=blue.

385 The location, size, and magnitude of finger-specific connectivity is summarized in Table 2.

386 **Table 2. Location of finger-specific connectivity in right SMC.**

Finger/task interaction	Hemisphere (SMC)	Region	Cluster size	Z-score (peak)	Peak Coordinates
F1>F2	right	precentral	9	4.36	(38,-12,66)
		postcentral	18	4.05	(38,-32,50)
F2>F1	right	postcentral	22	5.05	(50,-24,50)
		postcentral			(38,-32,62)
		precentral			(50,-16,46)
F2>F3	right	postcentral	22	4.41	(42,-28,54)
		postcentral		4.21	(34,-32,58)
F3>F2	right	postcentral	58	5.11	(50,-24,54)
		precentral		5.03	(34,-12,66)
		precentral		4.69	(46,-16,50)
F3>F4	right	precentral	58	4.01	(34,-16,66)
		postcentral		3.61	(50,-24,54)
		precentral		3.59	(38,-16,46)

		precentral		4.44	(50,-12,46)
F4>F3	right	postcentral	71	4.08	(42,-28,58)
		precentral		4.07	(38,-12,58)

387 Connectivity clusters in the right SMC from movement of individual fingers, surviving an intensity
388 threshold of p=0.05 with a family-wise error correction.

Discussion

To evaluate the possible role of the hippocampus in cognitive control, this study sought to

demonstrate spatial and temporal selectivity of hippocampal connectivity with the SMC during a

sensorimotor task.

Previously we examined hippocampal-SMC connectivity during mnemonic vs. non-mnemonic

finger movements and found connectivity increases during both tasks, suggesting a hippocampal

involvement in volitional control (75). As noted in the introduction, cognitive control of a

movement requires neural activity during volitional movements plus temporal and spatial

selectivity.

Temporal selectivity within the finger representations

401 Temporal selectivity for finger movements was identified as differences in the magnitude of

402 connectivity across time periods within a given finger representation area. Consistent with this

403 definition, maximal connectivity at each finger representation occurred when the finger was

404 moving, with reduced connectivity during other time periods.   A "spill over" effect was also

405 evident, in which the time periods adjacent to the period corresponding to the moving finger

exhibited elevated connectivity, likely because adjacent fingers do not move with complete

independence.

Temporal selectivity was observed in both pre- and postcentral finger representations. This is

expected, as motor and sensory systems operate jointly with movements closely tied to sensory

feedback.

Spatial selectivity

The topography of hippocampal connectivity with SMC was consistent with our previous study

(77) and those of others (78-82), which showed overlapping activation between nearby fingers.

Representations for later fingers (e.g., F4) were located dorsal to earlier fingers (F1-F3). The

topographic organization of connectivity associated with individual finger movements followed

this pattern in the postcentral gyrus of both hemispheres; although connectivity was present, no

topography was apparent in the precentral gyrus.

During a single time period, spatial selectivity reflected differences in the magnitude of

connectivity between finger representations. Within each time period, significant differences in

spatial connectivity were observed; however, the L and R hemispheres exhibited different

patterns of MAX and MIN connectivity, as will be summarized below. The functional relevance

for many of these differences cannot be explained until we understand the specific behavioral

effects mediated by MAX (positive) and MIN (inverse) connectivity.

In the L postcentral gyrus, the visual rest condition was required as a baseline to demonstrate

significant effects. Significant MAX and MIN connectivity was observed for F1 and F2 during

each time period, whereas F2 only showed greater MAX connectivity in its representation area

during movement. The amplitude of MIN connectivity in the representation of the moving finger

invariably differed from that of the adjacent finger, whereas no differences between fingers were

observed for MAX connectivity. In the R hemisphere, by contrast, T4 was required as the

baseline; furthermore, significant MAX and MIN connectivity changes were observed only in

the representation of the moving finger. The magnitude of both MIN and MAX in the moving

finger differed from those of an adjacent finger.

The complex spatial distribution of connectivity changes across finger representations in both

hemispheres is not surprising, considering the multiple motor processes that occur during a

single time period. Moving vs. waiting, activation of flexors vs. extensors, and movement

readiness vs. actual movement are some of the opposing factors that can potentially result from

direct vs. inverse connectivity. Fingers do not necessarily move independently of their

neighboring fingers; moreover, deliberate movement of one finger may also involve the active

suppression of stationary fingers.

In the L hemisphere, connectivity was measured relative to a baseline visual resting condition;

the interaction term for this connectivity analysis provides a signal simply to "move this finger,"

suggesting direct control of an individual finger independent of others. Connectivity in the R

hemisphere, on the other hand, required an interaction term that explicitly signaled, "move this

finger but not that one," suggesting an interactive process between the control of different

fingers. Because all subjects in this study were right-handed and volitional movements of the

right hand are controlled by the left SMC, these findings suggest hippocampal influences on

sensorimotor activity provide better independence of individual finger movements in the

dominant hand.

Our analysis further demonstrated that differential connectivity was resolvable between adjacent

fingers, although elevated connectivity for the moving finger could "spill over" into an adjacent

finger representation. Significant differences in connectivity were observed during T1, for

example, between the representations for F2 and F3, as well as between F1 and F2. A similar

spillover effect has been observed for fMRI activation during finger movements, where

movement-related activation extends into the representation of adjacent fingers (78-82).

Hippocampal properties are consistent with cognitive control

Because the hippocampus is preferentially involved in conscious memories and environmental

interactions (83, 84), studies of hippocampal interactions with the motor system have

traditionally been focused on its role in explicit learning and memory consolidation.

Hippocampal connectivity with the striatum increases during explicit motor learning and

consolidation (85-87); consolidation after rehearsal is further reflected through increased motor

activation following sleep in both the primary motor area and the hippocampus (85, 88). Our

previous study showed that hippocampal connectivity to SMC occurs during paced repetitive

tapping movements as well as sequence learning, and subjects' anticipatory responses indicate

volitional control of these movements (75). These findings demonstrate hippocampal

involvement in movements and motor learning under explicit cognitive control; by contrast, the

hippocampus is inactive during implicit learning of motor sequences (89).

The second requirement for cognitive control is spatial selectivity. The hippocampus is sensitive

to spatial location and navigation, with spatial functioning localized to the posterior

hippocampus (90, 91), including space cells that encode a spatial map (92-96). Because the

spatial location of the hand on the response pad was fixed in the current study, the spatial

location of a button press corresponded to a SMC finger representation, and posterior structural

seeds generated connectivity restricted to the SMC hand representation. Furthermore, spatial

selectivity was observed between finger representations, as the magnitude of connectivity for the

moving finger differed from that of the surrounding fingers from the representation of the finger

currently moving. Thus, spatial properties of the hippocampus provided the selectivity in motor

response required for cognitive control.

The third requirement for cognitive control is temporal selectivity. The hippocampus responds

differentially to sequences of events that differ in order (97-100), and to the intervals between

stimuli within a sequence (101-103). In the current study, cognitive awareness of the temporal

intervals between metronome beats provided the basis for anticipatory behavioral responses.

Furthermore, temporal selectivity was observed within each finger representation, such that

connectivity was greatest when the represented finger was moving.

Thus, connectivity with SMC during motor tasks is consistent with known hippocampal

properties and meets the three suggested criteria for the cognitive control of movements. This

does not preclude an additional role in cognitive control of SMC from other areas. Prefrontal

cortex, for example, has been suggested to play an indirect role through its connectivity with

dorsal premotor cortex (2), although functional coupling of prefrontal with premotor areas is

limited to movement sequences (66). During paced repetitive tapping, at least, hippocampal

connectivity provides the best explanation for the cognitive control of SMC.

Cognitive control vs. other theories of hippocampal function

There is currently no consensus about the functional significance of the spatial and temporal

properties of the hippocampus. Some have suggested the hippocampus has cognitive functions

beyond its traditional roles (104-106), whereas others suggest its diverse properties merely

reflect the varied components of episodic and long-term declarative memory (92, 93, 107).

Although hippocampal connectivity with SMC meets the requirements for cognitive control,

could the results be better explained through its other known functions?

Memory of the pacing interval, plus associations between the numerical onscreen display and the

corresponding fingers, were required for accurate performance during motor tasks in this study;

these modest memory requirements might arguably require hippocampal input. Neither spatial

nor temporal selectivity, however, is required to access this mnemonic information. Spatial

selectivity is unnecessary because the mapping between numerals and fingers never changed,

whereas temporal selectivity is unnecessary because widespread cortical rhythms (such as theta)

could serve to time events. The observed pattern of spatial and temporal selectivity is *required*

for cognitive control, however, to specify the finger to be moved and when.

An alternative explanation is that hippocampal connectivity in this study represents a form of

sensorimotor processing, facilitating the transformation of sensory signals into motor output

(108-111). This interpretation could explain why its topography and spatial selectivity were

most prominent in the postcentral gyrus. Even without hippocampal intervention, however,

extensive connections between pre- and postcentral gyrus provides sensory feedback (112), so

aside from cognitive control, the role of hippocampal sensorimotor processing in SMC is

unclear. Furthermore, facilitation of sensorimotor processing does not require temporal

specificity. Given the directional influence of the hippocampus on SMC (implicit in PPI

analysis), a hippocampal role in sensorimotor processing begs the question: what directs the

hippocampus to provide relevant spatio-temporal information to SMC as needed for task

performance? Considering the tight relationship between neural activity in the hippocampal

seeds and SMC (shown in Burman 2019, Fig 2), an additional control signal in SMC seems

unlikely.

This does not exclude the possibility of other sources of cognitive control under different

conditions. By analogy with the oculomotor system, damage to a single structure (e.g., FEF or

SMC) can produce deficits without eliminating all function (eye movements or hand movements)

due to the preservation of other structures with overlapping functions (SEF / PEF, premotor

cortex / basal ganglia). Similarly, executive functions show deficits following damage to

prefrontal areas, yet are not eliminated (113-115). Hippocampal dysfunction in Alzheimer's

impairs volitional motor control (116), but with overlapping systems for cognitive control, it

would not eliminate volitional movements.

Implications and directions for future research

This study provides evidence that the hippocampus is involved in the cognitive control of

volitional finger movements. Whether this represents a new function, or represents another

component of a single cognitive function, is unclear. These alternative possibilities suggest two

broad experimental questions for further investigation.

First, how extensive is the hippocampal role in cognitive control during volitional movements?

Based on the criteria for cognitive control, hippocampal connectivity should be lost when

"mindless" finger movements are generated while simultaneously performing a mentally-

challenging task; connectivity also needs to be demonstrated during volitional movements of

other body parts. In addition, a fixed spatial relationship between fingers and button presses in

the current study allowed cortical specificity for finger movements to provide a measure of

spatial selectivity; it's unclear how the hippocampal pattern of cortical / spatial selectivity would

be affected by changes in the spatial target of finger movements.

Second, is the hippocampal role in cognitive control limited to volitional movements? The

hippocampus is critically involved in the formation and recall of episodic and declarative

memories, thus fulfilling the first criterion for cognitive control (preferentially active during

volitional states). Whether hippocampal interactions with cortical areas meet the remaining

criteria for cognitive control over other mental functions is unclear, although theoretical accounts

of memory function suggest they might. When spatial and temporal selectivity of hippocampal

connectivity with cortical areas is necessary to perform cognitive tasks, a role of the

hippocampus in cognitive control is implied. The role of the hippocampus in cognitive control

may be evaluated through experimental manipulations that affect cognitive awareness and

performance, evaluating effects of these variables on the connectivity of the hippocampus with

brain areas directly involved in task performance. Comparing effects of these variables on

hippocampal and prefrontal connectivity may help identify conditions where these regions are

either complementary or synergistic in the cognitive control of behavior.

Conclusions

The hippocampus showed task-specific connectivity with SMC during paced movement tasks,

meeting the criteria for cortical control adapted from studies of cognition in the oculomotor

system: associated with volitional movements, restricted to the representations of the moving

fingers, and selective for the timing of the movements. Still present in a motor task without

demonstrable learning or memory effects, this pattern of connectivity is consistent with known

hippocampal properties, yet cannot be explained by its established functions. A role for the

hippocampus in cognitive control is indicated; a collaborative role with prefrontal cortex is

suggested from their close functional relationship across tasks and neurological conditions.

References

1. Cieslik EC, Zilles K, Caspers S, Roski C, Kellermann TS, Jakobs O, et al. Is there "one"☐ DLPFC in cognitive action control? Evidence for heterogeneity from co-activation-based parcellation. Cerebral cortex 2012:bhs256.

2. Cole MW, Schneider W. The cognitive control network: integrated cortical regions with dissociable functions. Neuroimage 2007;37(1):343-360.

3. Ochsner KN, Ray RD, Cooper JC, Robertson ER, Chopra S, Gabrieli JD, et al. For better or for worse: neural systems supporting the cognitive down- and up-regulation of negative emotion. Neuroimage 2004;23(2):483-99.

4. Rajah MN, Ames B, D'Esposito M. Prefrontal contributions to domain-general executive control processes during temporal context retrieval. Neuropsychologia 2008;46(4):1088-103.

5. Ridderinkhof KR, Ullsperger M, Crone EA, Nieuwenhuis S. The role of the medial frontal cortex in cognitive control. Science 2004;306(5695):443-7.

6. Liversedge SP, Findlay JM. Saccadic eye movements and cognition. Trends in Cognitive Sciences 2000;4(1):6-14.

7. Rayner K, Liversedge SP, White SJ, Vergilino-Perez D, Mail E, rayner@psych.umass.edu. Reading disappearing text: Cognitive control of eye movements. Psychological Science 2003;14(4) Jul 2003:United Kingdom http://www.

8. Pollatsek A, Rayner K. Eye movements and lexical access in reading. In: Comprehension processes in reading: Routledge; 2012. p. 165-186.

9. Hayhoe M, Ballard D. Eye movements in natural behavior. Trends in cognitive sciences 2005;9(4):188-194.

593   10.   Hutton SB. Cognitive control of saccadic eye movements. Brain and cognition

594   2008;68(3):327-340.

595   11.   McDowell JE, Dyckman KA, Austin BP, Clementz BA. Neurophysiology and

596   neuroanatomy of reflexive and volitional saccades: evidence from studies of humans.

597   Brain & Cognition 2008;68(3):255-70.

598   12.   Vernet M, Quentin R, Chanes L, Mitsumasu A, Valero-Cabré A. Frontal eye field, where

599   art thou? Anatomy, function, and non-invasive manipulation of frontal regions involved

600   in eye movements and associated cognitive operations. Frontiers in integrative

601   neuroscience 2014;8:66.

602   13.   Bruce CJ, Goldberg ME. Primate frontal eye fields: I. Single neurons discharging before

603   saccades. J. Neurophysiol. 1985;53:603-635.

604   14.   Guitton D, Buchtel HA, Douglas RM. Frontal lobe lesions in man cause difficulties in

605   suppressing reflexive glances and in generating goal-directed saccades. Exp. Brain Res.

606   1985;58:455-472.

607   15.   Keating EG. Frontal eye field lesions impair predictive and visually-guided pursuit eye

608   movements. Exp. Brain Res. 1991;86:311-323.

609   16.   Henik A, Rafal R, Rhodes D. Endogenously generated and visually guided saccades after

610   lesions of the human frontal eye fields. Journal of Cognitive Neuroscience

611   1994;6(4):400-411.

612   17.   Munoz DP, Everling S. Look away: the anti-saccade task and the voluntary control of eye

613   movement. Nature Reviews Neuroscience 2004;5(3):218.

18.   Rivaud S, Müri RM, Gaymard B, Vermersch AI, Pierrot-Deseilligny C. Eye movement
disorders after frontal eye field lesions in humans. Experimental Brain Research
1994;102(1):110-120.
19.   Stuyven E, Van der Goten K, Vandierendonck A, Claeys K, Crevits L. The effect of
cognitive load on saccadic eye movements. Acta psychologica 2000;104(1):69-85.
20.   Van der Stigchel S, Van Koningsbruggen M, Nijboer TCW, List A, Rafal RD. The role of
the frontal eye fields in the oculomotor inhibition of reflexive saccades: Evidence from
lesion patients. Neuropsychologia 2012;50(1):198-203.
21.   Gaymard B, Ploner CJ, Rivaud-Pechoux S, Pierrot-Deseilligny C. The frontal eye field is
involved in spatial short-term memory but not in reflexive saccade inhibition.
Experimental Brain Research 1999;129(2):288-301.
22.   Schiller PH, True SD, Conway JL. Deficits in eye movements following frontal eye field
and superior colliculus ablations. J. Neurophysiol. 1980;44:1175-1189.
23.   Schiller PH, Chou Ih. The effects of frontal eye field and dorsomedial frontal cortex
lesions on visually guided eye movements. Nature neuroscience 1998;1(3):248.
24.   Tehovnik EJ, Sommer MA, Chou IH, Slocum WM, Schiller PH. Eye fields in the frontal
lobes of primates. Brain Research Reviews 2000;32(2-3):413-448.
25.   Braun D, Weber H, Mergner TH, Schulte-Mönting J. Saccadic reaction times in patients
with frontal and parietal lesions. Brain 1992;115(5):1359-1386.
26.   Lynch JC. Saccade initiation and latency deficits after combined lesions of the frontal and
posterior eye fields in monkeys. Journal of Neurophysiology 1992;68(5):1913-1916.
27.   Chen LL, Wise SP. Supplementary eye field contrasted with the frontal eye field during
acquisition of conditional oculomotor associations. J. Neurophysiol. 1995;73:1122-1134.

28.   Lang CE, Schieber MH. Differential impairment of individuated finger movements in

humans after damage to the motor cortex or the corticospinal tract. Journal of

neurophysiology 2003;90(2):1160-1170.

29.   Lang CE, Schieber MH. Reduced muscle selectivity during individuated finger

movements in humans after damage to the motor cortex or corticospinal tract. Journal of

neurophysiology 2004;91(4):1722-1733.

30.   Catani M, Dell'Acqua F, Vergani F, Malik F, Hodge H, Roy P, et al. Short frontal lobe

connections of the human brain. Cortex 2012;48(2):273-291.

31.   Colby CL, Goldberg ME. The updating of the representation of visual space in parietal

cortex by intended eye movements. Science 1992;255(5040):90-92.

32.   Phillips AN, Segraves MA. Predictive Activity in Macaque Frontal Eye Field Neurons

During Natural Scene Searching. J Neurophysiol 2009;103:1238-1252.

33.   Umeno MM, Goldberg ME. Spatial processing in the monkey frontal eye field. I.

Predictive visual responses. Journal of neurophysiology 1997;78(3):1373-1383.

34.   Burman DD, Segraves MA. Primate frontal eye field activity during natural scanning eye

movements. Journal of Neurophysiology 1994;71(3):1266-1271.

35.   Hanes DP, Patterson Ii WF, Schall JD. Role of Frontal Eye Fields in Countermanding

Saccades: Visual, Movement, and Fixation Activity. 1998.

36.   Schall JD, Hanes DP. Neural basis of saccade target selection in frontal eye field during

visual search. Nature 1993;366(6454):467.

37.   Schall JD, Purcell BA, Heitz RP, Logan GD, Palmeri TJ. Neural mechanisms of saccade

target selection: gated accumulator model of the visual-motor cascade. European Journal

of Neuroscience 2011;33(11):1991-2002.

38. Garg A, Schwartz D, Stevens AA. Orienting auditory spatial attention engages frontal eye

fields and medial occipital cortex in congenitally blind humans. Neuropsychologia

2007;45(10):2307-21.

39. Heitz RP, Cohen JY, Woodman GF, Schall JD. Neural correlates of correct and errant

attentional selection revealed through N2pc and frontal eye field activity. Journal of

neurophysiology 2010;104(5):2433-2441.

40. Moore T, Fallah M. Microstimulation of the frontal eye field and its effects on covert

spatial attention. Journal of neurophysiology 2004;91(1):152-162.

41. Thompson KG, Bichot NP. A visual salience map in the primate frontal eye field.

Progress in brain research 2005;147:249-262.

42. Wardak C, Ibos G, Duhamel J-R, Olivier E. Contribution of the monkey frontal eye field

to covert visual attention. Journal of Neuroscience 2006;26(16):4228-4235.

43. Little DM, Klein R, Shobat DM, McClure ED, Thulborn KR. Changing patterns of brain

activation during category learning revealed by functional MRI. Cognitive Brain

Research 2004;22(1):84-93.

44. Floyer-Lea A, Wylezinska M, Kincses T, Matthews PM. Rapid modulation of GABA

concentration in human sensorimotor cortex during motor learning. Journal of

neurophysiology 2006;95(3):1639-1644.

45. Muellbacher W, Ziemann U, Wissel J, Dang N, Kofler M, Facchini S, et al. Early

consolidation in human primary motor cortex. Nature 2002;415(6872):640.

46. Rosenkranz K, Kacar A, Rothwell JC. Differential modulation of motor cortical plasticity

and excitability in early and late phases of human motor learning. Journal of

Neuroscience 2007;27(44):12058-12066.

47.   Sanes JN, Donoghue JP. Plasticity and primary motor cortex. Annual Review of

684   Neuroscience 2000;23:393-415.

685   48.   Pollok B, Latz D, Krause V, Butz M, Schnitzler A. Changes of motor-cortical oscillations

686   associated with motor learning. Neuroscience 2014;275:47-53.

687   49.   Ostry DJ, Gribble PL. Sensory Plasticity in Human Motor Learning. Trends in

688   Neurosciences 2016;39(2):115.

689   50.   Ma L, Wang B, Narayana S, Hazeltine E, Chen X, Robin DA, et al. Changes in regional

690   activity are accompanied with changes in inter-regional connectivity during 4 weeks

691   motor learning. Brain research 2010;1318:64-76.

692   51.   Steele CJ, Penhune VB. Specific increases within global decreases: a functional magnetic

693   resonance imaging investigation of five days of motor sequence learning. The Journal of

694   neuroscience 2010;30(24):8332-8341.

695   52.   Sun FT, Miller LM, Rao AA, D'Esposito M. Functional connectivity of cortical networks

696   involved in bimanual motor sequence learning. Cerebral cortex 2006;17(5):1227-1234.

697   53.   Stuphorn V, Schall JD. Executive control of countermanding saccades by the

698   supplementary eye field. Nature neuroscience 2006;9(7):925.

699   54.   Stuphorn V, Brown JW, Schall JD. Role of supplementary eye field in saccade initiation:

700   executive, not direct, control. Journal of neurophysiology 2009;103(2):801-816.

701   55.   Hikosaka O, Takikawa Y, Kawagoe R. Role of the basal ganglia in the control of

702   purposive saccadic eye movements. Physiological reviews 2000;80(3):953-978.

703   56.   Pierrot-Deseilligny C, Milea D, Muri RM. Eye movement control by the cerebral cortex.

704   Current Opinion in Neurology 2004;17:17-25.

57. Graybiel AM. The basal ganglia: learning new tricks and loving it. Current opinion in

neurobiology 2005;15(6):638-644.

58. Hoffstaedter F, Grefkes C, Caspers S, Roski C, Palomero-Gallagher N, Laird AR, et al.

The role of anterior midcingulate cortex in cognitive motor control. Human brain

mapping 2014;35(6):2741-2753.

59. Azuar C, Reyes P, Slachevsky A, Volle E, Kinkingnehun S, Kouneiher Fdr, et al. Testing

the model of caudo-rostral organization of cognitive control in the human with frontal

lesions. Neuroimage 2014;84:1053-1060.

60. Badre D. Cognitive control, hierarchy, and the rostro-caudal organization of the frontal

lobes. Trends in cognitive sciences 2008;12(5):193-200.

61. Cole MW, Yarkoni T, Repovš G, Anticevic A, Braver TS. Global connectivity of

prefrontal cortex predicts cognitive control and intelligence. The Journal of Neuroscience

2012;32(26):8988-8999.

62. Koechlin E, Ody C, Kouneiher Fdr. The architecture of cognitive control in the human

prefrontal cortex. Science 2003;302(5648):1181-1185.

63. Ridderinkhof KR, Van Den Wildenberg WPM, Segalowitz SJ, Carter CS. Neurocognitive

mechanisms of cognitive control: the role of prefrontal cortex in action selection,

response inhibition, performance monitoring, and reward-based learning. Brain and

cognition 2004;56(2):129-140.

64. Chouinard PA, Paus TÅ. The primary motor and premotor areas of the human cerebral

cortex. The neuroscientist 2006;12(2):143-152.

65. Koch G, Franca M, Del Olmo MF, Cheeran B, Milton R, Sauco MA, et al. Time course of
functional connectivity between dorsal premotor and contralateral motor cortex during
movement selection. The Journal of neuroscience 2006;26(28):7452-7459.
66. Abe M, Hanakawa T. Functional coupling underlying motor and cognitive functions of
the dorsal premotor cortex. Behavioural Brain Research 2009;198(1):13-23.
67. Kantak SS, Stinear JW, Buch ER, Cohen LG. Rewiring the brain: potential role of the
premotor cortex in motor control, learning, and recovery of function following brain
injury. Neurorehabilitation and neural repair 2012;26(3):282-292.
68. Chen LL, Wise SP. Neuronal activity in the supplementary eye field during acquisition of
conditional oculomotor associations. J. Neurophysiol. 1995;73:1101-1121.
69. Gaymard B, Pierrot-Deseilligny C, Rivaud S. Impairment of sequences of memory-
guided saccades after supplementary motor area lesions. Annals of Neurology 1990.
70. Isoda M, Tanji J. Cellular activity in the supplementary eye field during sequential
performance of multiple saccades. Journal of neurophysiology 2002;88(6):3541-3545.
71. Husain M, Parton A, Hodgson TL, Mort D, Rees G. Self-control during response conflict
by human supplementary eye field. Nature neuroscience 2003;6(2):117.
72. Parton A, Nachev P, Hodgson TL, Mort D, Thomas D, Ordidge R, et al. Role of the
human supplementary eye field in the control of saccadic eye movements.
Neuropsychologia 2007;45(5):997-1008.
73. Harvey P-O, Fossati P, Pochon J-B, Levy R, LeBastard G, Lehericy S, et al. Cognitive
control and brain resources in major depression: an fMRI study using the n-back task.
Neuroimage 2005;26(3):860-869.

74.   Lavie N. Attention, distraction, and cognitive control under load. Current Directions in

749   Psychological Science 2010;19(3):143-148.

750   75.   Burman DD. Hippocampal connectivity with sensorimotor cortex during volitional finger

751   movements: Laterality and relationship to motor learning. PLoS One

752   2019;14(9):e0222064.

753   76.   Burman DD. Hippocampal Influences on Movement, Sensory and Language Processing:

754   A Role in Cognitive Control? In: Hippocampus – Cytoarchitecture and Diseases: InTech

755   Open; 2021. p. 57-79.

756   77.   Gitelman DR, Penny WD, Ashburner J, Friston KJ. Modeling regional and

757   psychophysiologic interactions in fMRI: the importance of hemodynamic deconvolution.

758   Neuroimage 2003;19(1):200-207.

759   78.   Burman DD, Minas T, Bolger DJ, Booth JR. Age, sex, and verbal abilities affect location

760   of linguistic connectivity in ventral visual pathway. Brain and language 2013;124(2):184-

761   193.

762   79.   Burman DD, Lie-Nemeth T, Brandfonbrener AG, Parisi T, Meyer JR. Altered Finger

763   Representations in Sensorimotor Cortex of Musicians with Focal Dystonia: Precentral

764   Cortex. Brain Imaging and Behavior 2009;3(1):10-23.

765   80.   Beisteiner R, Windischberger C, Lanzenberger R, Edward V, Cunnington R, Erdler M, et

766   al. Finger somatotopy in human motor cortex.[see comment]. Neuroimage 2001;13(6 Pt

767   1):1016-26.

768   81.   Dechent P, Frahm J. Functional somatotopy of finger representations in human primary

769   motor cortex. Human Brain Mapping 2003;18(4):272-83.

770   82.   Hlustik P, Solodkin A, Gullapalli RP, Noll DC, Small SL. Somatotopy in human primary

771   motor and somatosensory hand representations revisited. Cerebral Cortex

772   2001;11(4):312-21.

773   83.   Kleinschmidt A, Nitschke MF, Frahm J. Somatotopy in the human motor cortex hand

774   area. A high-resolution functional MRI study. European Journal of Neuroscience

775   1997;9(10):2178-86.

776   84.   Volkmann J, Schnitzler A, Witte OW, Freund H. Handedness and asymmetry of hand

777   representation in human motor cortex. Journal of Neurophysiology 1998;79(4):2149-54.

778   85.   Behrendt R-P. Hippocampus and consciousness. Reviews in the Neurosciences

779   2013;24(3):239-266.

780   86.   Huijgen J, Samson S. The hippocampus: a central node in a large-scale brain network for

781   memory. Revue neurologique 2015;171(3):204-216.

782   87.   Albouy G, King BR, Maquet P, Doyon J. Hippocampus and striatum: Dynamics and

783   interaction during acquisition and sleep-related motor sequence memory consolidation.

784   Hippocampus 2013;23(11):985-1004.

785   88.   Albouy G, Fogel S, King BR, Laventure S, Benali H, Karni A, et al. Maintaining vs.

786   enhancing motor sequence memories: Respective roles of striatal and hippocampal

787   systems. Neuroimage 2015;108:423-434.

788   89.   Fernandez-Seara MA, Aznarez-Sanado M, Mengual E, Loayza FR, Pastor MA.

789   Continuous performance of a novel motor sequence leads to highly correlated striatal and

790   hippocampal perfusion increases. Neuroimage 2009;47(4):1797-1808.

791   90.   Walker MP, Stickgold R, Alsop D, Gaab N, Schlaug G. Sleep-dependent motor memory

792   plasticity in the human brain. Neuroscience 2005;133(4):911-917.

91.   Rose M, Haider H, Salari N, Buchel C. Functional dissociation of hippocampal

mechanism during implicit learning based on the domain of associations. J. Neurosci.

2011;31:13739-13745.

92.   Strange BA, Witter MP, Lein ES, Moser EI. Functional organization of the hippocampal

longitudinal axis. Nature Reviews Neuroscience 2014;15(10):655.

93.   Travis SG, Huang Y, Fujiwara E, Radomski A, Olsen F, Carter R, et al. High field

structural MRI reveals specific episodic memory correlates in the subfields of the

hippocampus. Neuropsychologia 2014;53:233-245.

94.   Eichenbaum H. On the Integration of Space, Time, and Memory. Neuron

2017;95(5):1007.

95.   Eichenbaum H. The role of the hippocampus in navigation is memory. Journal of

neurophysiology 2017;117(4):1785.

96.   Ekstrom AD, Ranganath C. Space, time, and episodic memory: the hippocampus is all

over the cognitive map. Hippocampus 2017.

97.   Libby LA, Hannula DE, Ranganath C. Medial temporal lobe coding of item and spatial

information during relational binding in working memory. J. Neurosci. 2014;34:14233-

14242.

98.   Rolls ET. Pattern separation, completion, and categorisation in the hippocampus and

neocortex. Neurobiology of learning and memory 2016;129:4-28.

99.   Davachi L, DuBrow S. How the hippocampus preserves order: the role of prediction and

context. Trends in cognitive sciences 2015;19(2):92-99.

100. Ekstrom AD, Copara MS, Isham EA, Wang W-c, Yonelinas AP. Dissociable networks

involved in spatial and temporal order source retrieval. Neuroimage 2011;56(3):1803-

1813.

101. Paz R, Gelbard-Sagiv H, Mukamel R, Harel M, Malach R, Fried I. A neural substrate in

the human hippocampus for linking successive events. Proceedings of the National

Academy of Sciences 2010;107(13):6046-6051.

102. Warburton EC, Brown MW. Findings from animals concerning when interactions

between perirhinal cortex, hippocampus and medial prefrontal cortex are necessary for

recognition memory. Neuropsychologia 2010;48(8):2262-2272.

103. Barnett AJ, O'neil EB, Watson HC, Lee ACH. The human hippocampus is sensitive to the

durations of events and intervals within a sequence. Neuropsychologia 2014;64:1-12.

104. MacDonald CJ, Fortin NJ, Sakata S, Meck WH. Retrospective and prospective views on

the role of the hippocampus in interval timing and memory for elapsed time. Timing &

Time Perception 2014;2(1):51-61.

105. MacDonald CJ. Prospective and retrospective duration memory in the hippocampus: is

time in the foreground or background? Phil. Trans. R. Soc. B 2014;369(1637):20120463.

106. Behrendt R-P. Consciousness, memory, and hallucinations. In: Hallucinations: Springer;

2013. p. 17-31.

107. Olsen RK, Moses SN, Riggs L, Ryan JD. The hippocampus supports multiple cognitive

processes through relational binding and comparison. Frontiers in human neuroscience

2012;6:146.

108.   Poch C, Campo P. Neocortical-hippocampal dynamics of working memory in healthy and

diseased brain states based on functional connectivity. Frontiers in human neuroscience

2012;6.

109.   Moscovitch M, Cabeza R, Winocur G, Nadel L. Episodic memory and beyond: the

hippocampus and neocortex in transformation. Annual review of psychology

2016;67:105-134.

110.   Bast T, Feldon J. Hippocampal modulation of sensorimotor processes. Progress in

neurobiology 2003;70(4):319-345.

111.   Cruikshank LC, Singhal A, Caplan JB. Theta oscillations reflect a putative neural

mechanism for human sensorimotor integration. Journal of Neurophysiology 2012:jn.

00893.2010.

112.   Dypvik AT, Bland BH. Functional connectivity between the red nucleus and the

hippocampus supports the role of hippocampal formation in sensorimotor integration.

Journal of neurophysiology 2004;92(4):2040-2050.

113.   Scheidt RA, Zimbelman JL, Salowitz NMG, Suminski AJ, Mosier KM, Houk J, et al.

Remembering forward: neural correlates of memory and prediction in human motor

adaptation. Neuroimage 2012;59(1):582-600.

114.   Catani M, Dell'Acqua F, Vergani F, Malik F, Hodge H, Roy P, et al. Short frontal lobe

connections of the human brain. cortex 2012;48(2):273-291.

115.   Alvarez JA, Emory E. Executive function and the frontal lobes: a meta-analytic review.

Neuropsychology review 2006;16(1):17-42.

116.   Barbey AK, Colom R, Solomon J, Krueger F, Forbes C, Grafman J. An integrative

857   architecture for general intelligence and executive function revealed by lesion mapping.

858   Brain 2012;135(4):1154-1164.

859