***Syntrophus* Conductive Pili Demonstrate that Common Hydrogen-Donating Syntrophs can**

have a Direct Electron Transfer Option

David J.F. Walker[1,2], Kelly P. Nevin[1], Dawn E. Holmes[1,3], Amelia-Elena Rotaru[1,4], Joy E.

Ward[1], Trevor L. Woodard[1], Jiaxin Zhu[5], Toshiyuki Ueki[1], Stephen S. Nonnenmann[2,5], Michael

J. McInerney[6], and Derek R. Lovley[1,2*]

[1]Department of Microbiology, University of Massachusetts-Amherst, Amherst, MA, USA

[2]Institute for Applied Life Sciences, University of Massachusetts-Amherst

[3]Department of Physical and Biological Science, Western New England University, Springfield,

MA, USA

[4]Department of Biology, University of Southern Denmark, Odense, Denmark

[5]Department of Mechanical and Industrial Engineering, University of Massachusetts-Amherst

[6]Department of Microbiology and Plant Biology, University of Oklahoma, Norman, OK, USA

*Corresponding author: dlovley@microbio.umass.edu

Abstract

Syntrophic interspecies electron exchange is essential for the stable functioning of diverse

anaerobic microbial communities. Hydrogen/formate interspecies electron transfer (HFIT), in

which H_2 and/or formate function as diffusible electron carriers, has been considered to be the

primary mechanism for electron sharing because most common syntrophs were thought to lack

biochemical components, such as electrically conductive pili (e-pili), necessary for direct

interspecies electron transfer (DIET). Here we report that *Syntrophus aciditrophicus*, one of the

most intensively studied microbial models for HFIT, produces e-pili and can grow via DIET.

Pilin genes likely to yield e-pili were found in other genera of hydrogen/formate-producing

syntrophs. The finding that DIET is a likely option for diverse syntrophs that are abundant in

many anaerobic environments necessitates a reexamination of the paradigm that HFIT is the

predominant mechanism for syntrophic electron exchange within anaerobic microbial

communities of biogeochemical and practical significance.

Introduction

H_2/formate interspecies electron transfer (HFIT) and direct interspecies electron transfer

(DIET) are mechanisms for the electron-donating partner in syntrophic consortia to dispose of

electrons released from the oxidation of key intermediates (organic acids, alcohols, aromatics)

during the anaerobic degradation of complex organic matter [1-5]. The relative proportion of

electron flux through DIET or HFIT can influence the speed of interspecies electron transfer, the

stability of anaerobic microbial communities, and their ability to adapt to environmental change

[4,6,7]. However, there are no accurate methods for measuring the rates at which H_2 and formate are

transferred between microbes or for quantifying interspecies electrical currents in complex

communities. Therefore, the importance of HFIT or DIET in microbial communities is typically

inferred from the composition of the microbial community.

It has previously been considered that DIET is primarily restricted to environments in

which *Geobacter* species are abundant [8-12], because *Geobacter* species are the only microbes in

pure culture that have been definitively shown to function as electron-donating partners for DIET

[9,13-17]. Studies with *Geobacter metallireducens* have suggested that electrically conductive pili

(e-pili) are an important conduit for electron transfer from the electron-donating partner [9,14,17,18].

It was initially considered that only short pilin monomers (ca. 60 amino acids), such as those

found in *G. metallireducens* and *G. sulfurreducens*, could assemble into e-pili [19]. However, it

was subsequently found that larger pilin monomers (> 100 amino aids), phylogenetically distinct

from *Geobacter* pilins, can yield e-pili and that e-pili have independently evolved multiple times

[20]. This raises the possibility that some microorganisms not previously known to be capable of

extracellular electron transfer may express e-pili to enable DIET.

For example, *Syntrophus aciditrophicus* is one of the most intensively studied pure

culture models for HFIT [21-24]. It was previously concluded that *S. aciditrophicus* would be

unlikely to participate in DIET [24,25] because it lacks a gene encoding a protein homologous to the

Geobacter pilin monomer (PilA) that assembles into electrically conductive pili (e-pili) [26].

Here we report that *S. aciditrophicus* expresses e-pili and is capable of growing via

DIET. These results, and analysis of the pilin genes of other common syntrophs, indicate that the

capacity for DIET should be considered as an option for microorganisms known to grow via

HFIT and suggest that DIET may be more widespread than previously considered.

Results and Discussion

Transmission electron microscopy of *S. aciditrophicus* revealed filaments with a

morphology typical of type IVa pili (Fig. 1a). A complement of genes for protein components

required for type IV pilus assembly (*pilA, pilD, pilM, pilN, pilO, pilP, pilQ*) and transcriptional

control of pilus expression (*pilS, pilR*) are also present in the genome (Supplementary Fig. 1).

One gene, SYN_00814 encodes a peptide with a N-terminal domain characteristic of PilA, the

pilin monomer for Type IVa pili found in other microorganisms (Fig. 1b). This includes a short

leader peptide (13 amino acids) that is cleaved by PilD at the G|FTLIE recognition site and a

highly conserved, hydrophobic, pilin assembly interface that facilitates pilin polymerization into

pili [27,28].

The amino acid sequence of the putative PilA fits the empirical criteria [20] for a pilin

monomer likely to yield e-pili: 1) aromatic amino acids are located in the key positions required

for conductivity in *G. sulfurreducens* e-pili; 2) the abundance of aromatic amino acids (10.9 % of

amino acids) is above the minimum threshold of 9 % thought to be necessary for high e-pili

conductivity; and 3) no large gaps (> 40 amino acids) that lack aromatic amino acids (Fig. 1b).

Fig.1. Pili protruding from *S. aciditrophicus* and the putative *S. aciditrophicus* PilA pilin

monomer peptide identified from the genome sequence. (a) Transmission electron

micrograph showing multiple pili protruding from *S. aciditrophicus* (pili highlighted with black

80 arrows). (**b**) Key characteristics of the predicted amino acid sequence of the *S. aciditrophicus*

81 pilin monomer. The PilA monomer of *G. sulfurreducens* that yields conductive pili is shown for

82 comparison. Aromatic amino acid abundance is calculated as a percentage of the mature peptide.

84   Low culture densities of *S. aciditrophicus* prevented harvesting sufficient quantities of

85 pili to measure pili conductance on the electrode arrays previously employed for the study of

86 other e-pili [20]. Therefore, the method initially employed to document the presence of e-pili in *G.*

87 *sulfurreducens* [29] was adapted [30] as an alternative approach. Cultures of *S. aciditrophicus* were

88 directly drop-cast on highly ordered pyrolytic graphite (HOPG), washed with deionized water,

89 and examined with atomic force microscopy. Topographic imaging, (Fig. 2a), indicated that the

90 distance from the surface of the HOPG to the top of the pili (Fig. 2b) was 4.0 ± 0.7 nm (n=27;

91 nine different locations on three separate pili). The 4 nm diameter of the *S. aciditrophicus* pili is

92 larger than the 3 nm reported for the e-pili of *Geobacter* species [31-33], but thinner than the ca. 5-6

93 nm diameter of the intensively studied type IVa pili of *Psedumonas* and *Neisseria* species [34-36].

94   Conductive imaging revealed that the pili were electrically conductive (Fig. 2b, c). Point-

95 mode current-voltage (I-V) spectroscopy (Fig. 2d; Supplementary Fig. 2) yielded an Ohmic-like

96 response with a conductance of 6.1 ± 1.2 nS (mean $\pm$ standard deviation; n=9) that was similar to

97 the previously reported [30] conductance of 4.5 ± 0.3 nS for wild-type *G. sulfurreducens* pili. As

98 previously reported [30], the pili from *G. sulfurreducens* strain Aro-5, which lack key aromatic

99 amino acids required for high conductivity [31,33,37,38], had much lower $(0.004 \pm 0.002$ nS)

100 conductance (Fig. 2d).

Fig. 2. Conducting tip atomic force microscopy demonstrated that _Syntrophus_ _aciditrophicus_ pili are electrically conductive. (**a**) Contact mode topographic imaging of pili on highly ordered pyrolytic (HOPG). Red line designates the cross section examined in (b). (**b**) Topographic analysis of the height/diameter of an individual pilus and corresponding current measurements (100 mV differential between the tip and the HOPG) across the pilus cross-section. (**c**) Current response of the pili shown in (a) with an applied 100 mV differential. (**d**) Current-voltage analysis of individual pili of _S. aciditrophicus_ (blue data points), wild-type _G. sulfurreducens_ (black data points) and the Aro-5 strain of _G. sulfurreducens_ (green data points). Current-voltage spectroscopy is shown for one pilus of each type and are representative of

analysis of three distinct locations on three separate pili of each type. Additional scans of *S.*

aciditrophicus pili available in Supplementary Fig. 2. The *G. sulfurreducens* wild-type and

strain Aro-5 data are from reference [30].

The lack of tools for genetic manipulation of *S. aciditrophicus* limited further functional

analysis of the putative PilA gene predicted to yield its e-pili in the native organism. Therefore,

the gene was heterologously expressed in *G. sulfurreducens*, replacing the native *G.*

sulfurreducens pilA with an approach that has successfully yielded *G. sulfurreducens* strains that

express a diversity of both highly conductive and poorly conductive heterologous pili [20,32,39-41].

This new *G. sulfurreducens* strain, designated strain SP (for *Syntrophus* pili), expressed abundant

pili (Fig. 3a). The heterologously expressed pili had a diameter (4.1 ± 0.4 nm) and conductance

(5.9 ± 0.7 nS) nearly identical to the native pili expressed by *S. aciditrophicus* (Fig. 3b). *G.*

sulfurreducens strain SP produced electrical current densities comparable to the control strain

expressing the *G. sulfurreducens* wild-type *pilA* (Fig. 3c). Such high current densities are only

possible when *G. sulfurreducens* expresses e-pili [20,37]. As previously reported [20,37], *G.*

sulfurreducens strain Aro-5, with its poorly conductive pili, produced much lower current

densities (Fig. 3c). Networks of pili sheared from the electrode-grown biofilm of strain SP,

purified, and drop cast on electrode arrays as previously described [20], had a conductance

comparable to *G. sulfurreducens* wild-type pili networks (Fig. 3d). These results further

demonstrated that the *S. aciditrophicus* PilA gene yields a pilin monomer that can assemble into

e-pili.

Fig. 3. Physical, electrical, and functional, analysis of the putative *Syntrophus aciditrophicus*

PilA via heterologous expression in *Geobacter sulfurreducens*. (**a**) Transmission electron

micrograph of pili expression in the strain of *G. sulfurreducens* in which the native *pilA* was

replaced with the *S. aciditrophicus pilA*. Examples of the location of pili is designated with

white arrows. (**b**) Current-voltage profile for three separate locations on three pili expressed by *S.*

aciditrophicus pili (data aggregated from individual plots in Fig.1d and Supplementary Fig. S2)

and for three separate locations on three pili expressed in the SP strain of *G. sulfurreducens* in

which the wild-type *pilA* was replaced with the *S. aciditrophicus pilA* (individual plots shown in

Supplentary Fig. 3). (**c**) Current production by *G. sulfurreducens* with wild-type PilA; the

synthetic Aro-5 PilA designed to yield poorly conductive pili; or *S. aciditrophicus* PilA. (**d**)

Four-probe conductance (mean $\pm$ standard deviation n=9) of films of purified pili from strains of

G. sulfurreducens expressing; *S. aciditrophicus* PilA compared with previously reported [20]

conductances of pili derived from wild-type and Aro-5 strains of *G. sulfurreducens.*

The presence of e-pili in *S. aciditrophicus* suggested it might be capable of establishing

an electrical connection for DIET. To evaluate this, *S. aciditrophicus* was grown in co-culture

with *G. sulfurreducens,* the microbe most intensively studied as an electron-accepting partner for

DIET [4]. *G. sulfurreducens* can also function as a H_2- and formate-consuming partner for HFIT [42].

Therefore, even if *S. aciditrophicus* was unable to grow via DIET it would be expected to grow

in co-culture with *G. sulfurreducens* via HFIT, providing a positive control to ensure that the

culture medium was appropriate for co-culture growth. The electron donor was benzoate, a

substrate that *S. aciditrophicus* can metabolize, but *G. sulfurreducens* cannot. The electron

acceptor was fumarate, an electron acceptor that only *G. sulfurreducens* can utilize. In the

presence of a H_2/formate-consuming partner, *S. aciditrophicus* metabolizes benzoate to acetate

with the production of either H_2:

$C_7H_5O_2^- + 7\ H_2O \rightarrow 3CH_3COO^- + HCO_3^- + 3\ H^+ + 3\ H_2$

or formate:

$C_7H_5O_2^- + 2\ HCO_3^- + 4\ H2O \rightarrow 3\ CH_3COO^- + 3\ H^+ + 3\ HCOO^-$

With electron transfer via DIET the relevant reaction is:

$C_7H_5O_2^- + 7\ H_2O \rightarrow 3CH_3COO^- + HCO_3^- + 9\ H^+ + 6\ e^-$

In addition to oxidizing H_2 and formate, or consuming electrons released during DIET, *G.*

sulfurreducens can also metabolize acetate, making the overall reaction expected for the

oxidation of benzoate with the reduction of fumarate to succinate in the co-culture:

$C_7H_5O_2^- + 15\ ^-OOC\text{-}CH\text{=}CH\text{-}COO^- + 19\ H_2O \rightarrow 7\ HCO_3^- + 15\ ^-OOC\text{-}CH_2\text{-}CH_2\text{-}COO^- + 6\ H^+$

S. aciditrophicus/G. sulfurreducens co-cultures grew with repeated sub-culturing and,

within the experimental error, exhibited the expected stoichiometry of benzoate consumption and

succinate production (Fig. 4a). In these co-cultures HFIT, DIET, or a combination of the two,

were feasible. Therefore, to eliminate the possibility of HFIT, *S. aciditrophicus* was co-cultured

with the previously described strain of *G. sulfurreducens* [18], designated here as *G.*

*sulfurreducens*HF, that cannot utilize H_2 or formate because the genes for the uptake hydrogenase

and formate dehydrogenase were deleted. The *S. aciditrophicus/ G. sulfurreducens*HF co-culture

had a longer initial lag period, but then metabolized benzoate with the reduction of fumarate

almost as fast as the co-culture with wild-type *G. sulfurreducens* (Fig. 4b). These results

demonstrate that *S. aciditrophicus* can grow via DIET.

DIET requires physical electrical contacts between the electron-donating and electron-

accepting partner, whereas contact is not required for HFIT [4,5]. In some co-cultures the

requirement for contact manifests as visible aggregates [13], but other DIET co-cultures produce

small, relatively fragile aggregates [14]. There were no visible aggregates in the *S.*

aciditrophicus/G. sulfurreducens co-cultures. However, more small clumps were seen in

scanning electron micrographs of co-cultures in which *G. sulfurreducens*HF was the electron-

accepting partner than co-cultures with wild-type *G. sulfurreducens* provided as the electron-

accepting partner (Fig. 4 d,e). This observation is consistent with the need for contact between

electron-donating and electron-accepting partners participating in DIET. In contrast, HFIT was

an option for the co-cultures established with wild-type *G. sulfurreducens* and aggregation is not

required when *G. sulfurreducens* grows via HFIT [42].

Granular activated carbon (GAC) can greatly reduce the initial lag time in establishing

DIET-based co-cultures because both partners attach to GAC, which functions as an electrically

conductive conduit [14,43]. GAC substantially reduced the lag time of the *S. aciditrophicus/ G.*

*sulfurreducens*HF co-cultures (Fig. 4c). As previously observed for other co-cultures in which

GAC promoted DIET, cells from the *S. aciditrophicus/ G. sulfurreducens*HF co-culture heavily

colonized the GAC (Fig. 4e), consistent with a GAC conduit for DIET.

Fig. 4. Co-cultures of *Syntrophus aciditrophicus* and *Geobacter sulfurreducens* grow via

DIET. Metabolism with *S. aciditrophicus* co-cultured with (**a**) wild-type *G. sulfurreducens*, (**b**)

*G. sulfurreducens*HF, which is unable to use H_2 or formate. (**c**) *G. sulfurreducens*HF with granular

activated carbon (GAC) amendment. No acetate or formate was detected in any of the co-

cultures. GAC interfered with determination of benzoate, which is not shown for GAC-amended

cultures. Scanning electron micrographs (SEM) of cells collected on filters from co-cultures with

(**d**) *G. sulfurreducens* wild-type or (**e**) *G. sulfurreducens* strain$_{HF}$ demonstrating greater

aggregation in co-cultures with *G. sulfurreducens*$_{HF}$ in which DIET was the only option for

interspecies electron exchange. (**f**) SEM of cells on GAC from co-culture of *S. aciditrophicus*

with *G. sulfurreducens* strain$_{HF}$. Circles-benzoate; squares-succinate. Data are the mean and

standard deviation of triplicate cultures.

In addition to e-pili, multi-heme outer-surface *c*-type cytochromes appear to play an

important role in DIET in *Geobacter* species [4,5]. However, the S. *aciditrophicus* genome encodes

only a few putative *c*-type cytochromes [24] and cytochromes were not readily apparent in heme-

stained preparations of cell protein (Supplementary Fig. 4). Not all microbes require

cytochromes for effective electron transport to the outer cell surface [44]. More detailed

examination of the role of e-pili and other *S. aciditrophicus* components in DIET will require the

development of methods for genetic manipulation of this microorganism.

S. aciditrophicus is the first isolate outside the genus *Geobacter* found to grow via DIET

and it is the first syntroph shown to have the option to grow via HFIT or DIET. Other

Syntrophus species also have pilin genes with the empirical criteria, detailed above, of aromatic

amino acid placement and abundance likely to yield e-pili (Fig. 5). The pilins of other diverse

genera of syntrophic microorganisms known to grow via HFIT in defined co-cultures also meet

these criteria (Fig. 5). Establishing conditions that favor DIET often enrich for microbes in these

genera [6]. For example, in enrichment cultures specifically designed to promote the metabolism of

propionate or butyrate via DIET, *Smithella* (propionate enrichment) or *Syntrophomonas*

(butyrate enrichment) species were the most abundant bacteria [45-47]. The greater energetic

demands [48] required for synthesizing the abundant aromatic amino acids that are required for e-

pili conductivity suggests that e-pili provide a strong selective advantage under some

environmental conditions. Conferring the capacity for DIET is a likely explanation.

Fig. 5. A diversity of syntrophs known to produce H_2 and/or formate as an interspecies

electron carrier encode for pilin monomers likely to yield electrically conductive pili.

Position of aromatic amino acids designated in magenta and the largest gap that lacks aromatic

amino acids designated in green. The full protein sequence alignment for each pilin can be found

in Supplementary Fig. 5. Each of the pilins meets the empirical criteria derived from previous

studies [20] of: 1) aromatic amino acids located in the key positions required for conductivity in *G.*

sulfurreducens e-pili; 2) the abundance of aromatic amino acids greater than 9%; and 3) no

aromatic-free gaps of greater than 40 amino acids.

For decades, interactions within complex anaerobic microbial communities have been

interpreted through the lens of HFIT, but the available data does not rule out the possibility of

DIET, which was not known to be an option at the time. For example, estimated H_2 turnover

rates in anaerobic digesters, rice paddy soils, and aquatic sediments were consistently less than

10 % of the independently determined rate of methane production derived from the reduction of

carbon dioxide to methane [49-51], a result that is consistent with DIET providing most of the

electrons for carbon dioxide reduction. The mismatch between measured H_2 turnover rates and

the concept of H_2 as a primary interspecies electron carrier was rationalized with the suggestion

that there was a separate pool of H_2 within closely juxatapositioned assemblages of H_2 producers

and H_2 consumers. However, the existence of two distinct pools of H_2 that did not equilibrate

over time via diffusion, a seemingly physical impossibility, was never verified. Attempts to

accurately measure formate fluxes have also been problematic [52] and there is no method for

directly measuring DIET-based electron fluxes in complex communities.

A fresh perspective and new analytical tools will be required to resolve the relative

importance of HFIT and DIET in diverse anaerobic microbial communities. Just as electron-

accepting partners have different gene expression patterns depending on whether they are

participating in HFIT or DIET [18,53], it may be possible to determine whether electron-donating

syntrophs are engaged in DIET or HFIT from metatranscriptomic analysis of anaerobic

communities.

METHODS

Bacterial strains, plasmids and culture conditions

Syntrophus aciditrophicus, G. sulfurreducens wild-type, *G. sulfurreducens* strain Aro-5,

and *G. sulfurreducens*HF were obtained from our laboratory culture collections. A strain of *G.*

sulfurreducens expressing *pilA* from *S. aciditrophicus* rather than native *G. sulfurreducens pilA*

was constructed as previously described [20]. *S. aciditrophicus* and *G. sulfurreducens* strains were

routinely grown under strict anaerobic conditions at 30 °C in previously described [54] defined,

bicarbonate-buffered medium with N_2:CO_2 (80:20) as the gas phase. For *S. aciditrophicus,* the

medium was amended with crotonate (20 mM) and for *G. sulfurreducens* strains, acetate (10

mM) was the electron donor and fumarate (40 mM) was the electron acceptor. The presence of *c*-

type cytochromes in whole cell lysates was evaluated with heme-staining of proteins separated

on denaturing polyacrylamide gels as previously described [17]. *G. sulfurreducens* strains were

grown with a graphite electrode as the electron acceptor as previously described [55].

Cocultures were established in 10 ml of culture medium [54] in anaerobic pressure tubes

with benzoate (1 mM) as the electron donor and fumarate (40 mM) as the electron acceptor with

cysteine (2 mM) and sulfide (1 mM) added as reducing agents. When noted, cultures were

amended with granular activated carbon (0.25 g; 8-20 mesh). Benzoate, acetate, and succinate

were analyzed with high-performance liquid chromatography as previously described [16].

Previously described methods were employed for transmission electron microscopy [21] and

scanning electron microscopy [26].

Pili conductivity

A 100 µl sample of cultures was drop cast onto highly oriented pyrolytic graphite

(HOPG). After 10 min, the HOPG was washed twice with 100 µl of deionized water, blotted dry

to remove excess water, and dried for 12 hours at 24 °C in a desiccator. Samples were

equilibrated with atmospheric humidity for at least 2 hours and then examined with an Oxford

Instruments Cypher ES Environmental AFM in ORCA electrical mode equipped with a Pt/Ir-

coated Arrow-ContPT tip with a 0.2 N/m force constant (NanoWorld AG, Neuchâtel,

Switzerland). Pili were located in contact mode, with a set point of 0.002 V and a scan rate of 1.5

Hz. For conductive imaging, the grounded tip, attached to a transimpedance amplifier, served as

a translatable top electrode to locally detect the current response of the individual pili to a 100

mV bias applied to the HOPG substrate. Individual pili conductivity was further characterized by

lightly pressing the AFM tip (set point 0.002 V) to the top of the pili and applying a

quadruplicate amplitude of ±0.6 V voltage sweep at a frequency of 0.99 Hz, receiving ca. 8,000

points of reference per measurement. Pili expressed in *G. sulfurreducens* were further analyzed

with four probe conductivity measurements on films of pili purified from cells as previously

described [20].

References
294

295   1   Stams, A. J. & Plugge, C. M. Electron transfer in syntrophic communities of anaerobic
296   bacteria and archaea. *Nature Rev. Microbiol.* **7**, 568-577 (2009).
297   2   McInerney, M. J., Sieber, J. R. & Gunsalus, R. P. Syntrophy in anaerobic global carbon
298   cycles. *Curr Opin. Biotechnol.* **20**, 623-632 (2009).
299   3   Kouzuma, A., Kato, S. & Wantanabe, K. Microbial interspecies interactions: recent
300   findings in syntrophic consortia. *Front, Microbiol.* **6**, 477 (2015).
301   4   Lovley, D. R. Syntrophy goes electric: direct interspecies electron transfer. *Ann. Rev.*
302   *Microbiol.* **71**, 643-664 (2017).
303   5   Lovley, D. R. Happy together: microbial communities that hook up to swap electrons.
304   *ISME J.* **11**, 327-336 (2017).
305   6   Martins, G., Salvador, A. F., Pereira, L. & Alves, M. M. Methane production and
306   conductive materials: a critical review. *Environ. Sci. Technol.* **52**, 10241-10253 (2018).
307   7   Park, J.-H., Kang, H.-J., Park, K.-H. & Park, H.-D. Direct interspecies electron transfer
308   via conductive materials: A perspective for anaerobic digestion applications. *Bioresource*
309   *Technol.* **254**, 300-311 (2018).
310   8   Morita, M. *et al.* Potential for direct interspecies electron transfer in methanogenic
311   wastewater digester aggregates. *mBio* **2**, e00159-11 (2011).
312   9   Rotaru, A.-E. *et al.* A new model for electron flow during anaerobic digestion: direct
313   interspecies electron transfer to Methanosaeta for the reduction of carbon dioxide to
314   methane. *Energy & Environ. Sci.* **7**, 408-415 (2014).
315   10   Shrestha, P. M. *et al.* Correlation between microbial community and granule conductivity
316   in anaerobic bioreactors for brewery wastewater treatment. *Bioresource Tech.* **174**, 306-
317   310 (2014).
318   11   Ha, P. T. *et al.* Syntrophic anaerobic photosynthesis via direct interspecies electron
319   transfer. *Nature Comm.* **7**, 13924 (2017).
320   12   Holmes, D. E. *et al.* Metatranscriptomic evidence for direct interspecies electron transfer
321   between Geobacter and Methanothrix species in rice paddy soils. *App.l Environ.*
322   *Microbiol.* **83**, e00223-00217 (2017).
323   13   Summers, Z. M. *et al.* Direct exchange of electrons within aggregates of an evolved
324   syntrophic co-culture of anaerobic bacteria. *Science* **330**, 1413-1415 (2010).

14 Rotaru, A.-E., Shrestha, P. M., Liu, F., Nevin, K. P. & Lovley, D. R. Direct interspecies
electron transfer during syntrophic growth of Geobacter metallireducens and
Methanosarcina barkeri on ethanol. *Appl. Environ. Microbiol.* **80**, 4599-4605 (2014).
15 Rotaru, A.-E., Woodard, T. L., Nevin, K. P. & Lovley, D. R. Link between capacity for
current production and syntrophic growth in Geobacter species. *Front. Microbiol.* **6**, 744
(2015).
16 Wang, L.-Y., Nevin, K. P., Woodard, T. L., Mu, B.-Z. & Lovley, D. R. Expanding the
diet for DIET: electron donors supporting direct interspecies electron transfer (DIET) in
defined co-cultures. *Front. Microbiol.* **7**, 236 (2016).
17 Ueki, T. *et al.* Geobacter strains expressing poorly conductive pili reveal constraints on
direct interspecies electron transfer mechanisms. *mBio* **9**, e01273-18 (2018).
18 Shrestha, P. M. *et al.* Transcriptomic and genetic analysis of direct interspecies electron
transfer. *Appl. Environ. Microbiol.* **79**, 2397-2404 (2013).
19 Holmes, D. E., Dang, Y., Walker, D. J. F. & Lovley, D. R. The electrically conductive
pili of Geobacter species are a recently evolved feature for extracellular electron transfer.
Microbial Genomics **2**, doi: 10.1099/mgen.1090.000072 (2016).
20 Walker, D. J. F. *et al.* Electrically conductive pili from genes of phylogenetically diverse
microorganisms. *ISME J.* **12**, 48-58 (2018).
21 Hopkins, B. T., McInerney, M. J. & Warikoo, V. Anaerobic syntrophic benzoate
degradation: evidence for a threshold and isolation of a syntrophic benzoate degrader.
Appl. Environ. Microbiol. **61**, 526-530 (1995).
22 Bradley, E. J., Bhupathiraju, V. K., Tanner, R. S., Woese, C. R. & McInerney, M. J.
Syntrophus aciditrophicus sp. nov., a new anaerobic bacterium that degrades fatty acids
and benzoate in syntrophic association with hydrogen-using microorganisms. *Arch.*
Microbiol. **171**, 107-114 (1999).
23 McInerney, M. J. *et al.* The genome of Syntrophus aciditrophicus: life at the
thermodynamic limit of microbial growth. *Proc Natl Acad Sci U S A* **104**, 7600-7605
(2007).
24 Sieber, J. R., Le, H. M. & McInerney, M. J. The importance of hydrogen and formate
transfer for syntrophic fatty, aromatic and alicyclic metabolism. *Environ. Microbiol.* **16**,
177-188 (2014).
25 Sieber, J. R., McInerney, M. J. & Gunsalus, R. P. Genomic insights into syntrophy: the
paradigm for anaerobic metabolic cooperation. *Ann. Rev. Microbiol.* **66**, 429-452 (2012).
26 Lovley, D. R. Electrically conductive pili: biological function and potential applications
in electronics. *Curr. Opin. Electrochem.* **4**, 190-198 (2017).
27 Craig, L. & Li, J. Type IV pili: paradoxes in form and function. *Curr. Opin. Struct. Biol.*
18, 267-277 (2008).
28 Hospenthal, M. K., Costa, T. R. D. & Waksman, G. A comprehensive guide to pilus
biogenesis in Gram-negative bacteria. *Nature Rev. Microbiol.* **15**, 365-379 (2017).
29 Reguera, G. *et al.* Extracellular electron transfer via microbial nanowires. *Nature* **435**,
1098-1101 (2005).
30 Walker, D. J. F. *et al.* The archaellum of Methanospirillum hungatei is electrically
conductive. *mBio* **10**, (in press) (2019).
31 Adhikari, R. Y., Malvankar, N. S., Tuominen, M. T. & Lovley, D. R. Conductivity of
individual *Geobacter* pili. *RSC Adv.* **6**, 8354-8357 (2016).

32   Tan, Y. *et al.* Expressing the Geobacter metallireducens PilA in Geobacter sulfurreducens
yields pili with exceptional conductivity. *mBio* **8**, e02203-16 (2017).
33   Malvankar, N. S., Yalcin, S. E., Tuominen, M. T. & Lovley, D. R. Visualization of
charge propagation along individual pili proteins using ambient electrostatic force
microscopy. *Nature Nanotechnol.* **9**, 1012-1017 (2014).
34   Craig, L. *et al.* Type IV pilus structure by cryo-electron microscopy and crystallography:
implications for pilus assembly and functions. *Molecular Cell* **23**, 651-662 (2006).
35   Kolappan, S. *et al.* Structure of the Neisseria meningitidis Type IV pilus. *Nature*
Commun. **7**, 13015 (2016).
36   Wang, F. *et al.* Cryoelectron microscopy reconstructions of the Pseudomonas aeruginosa
and Neisseria gonorrhoeae type IV pili at sub-nanometer resolution. *Structure* **25**,
14231435 (2017).
37   Vargas, M. *et al.* Aromatic amino acids required for pili conductivity and long-range
extracellular electron transport in Geobacter sulfurreducens *mBio* **4**, e00105-13. (2013).
38   Malvankar, N. S. *et al.* Structural basis for metallic-like conductivity in microbial
nanowires. *mBio* **6**, e00084-15 (2015).
39   Liu, X. *et al.* A Geobacter sulfurreducens strain expressing Pseudomonas aeruginosa type
IV pili localizes OmcS on pili but Is deficient in Fe(III) oxide reduction and current
production. *Appl. Environ. Microbiol.* **80**, 1219-1224 (2014).
40   Tan, Y. *et al.* The low conductivity of Geobacter uraniireducens pili suggests a diversity
of extracellular electron transfer mechanisms in the genus Geobacter. *Front. Microbiol.*
7, 980 (2016).
41   Tan, Y. *et al.* Synthetic biological protein nanowires with high conductivity. *Small* **12**,
4481–4485 (2016).
42   Rotaru, A.-E. *et al.* Interspecies electron transfer via H2 and formate rather than direct
electrical connections in co-cultures of Pelobacter carbinolicus and Geobacter
sulfurreducens. *Appl. Environ. Microbiol.* **78**, 7645-7651 (2012).
43   Liu, F. *et al.* Promoting direct interspecies electron transfer with activated carbon. *Energy*
Environ. Sci. **5**, 8982-8989 (2012).
44   Light, S. H. *et al.* A flavin-based extracellular electron transfer mechanism in diverse
Gram-positive bacteria. *Nature* **562**, 140-144 (2018).
45   Zhao, Z. *et al.* Communities stimulated with ethanol to perform direct interspecies
electron transfer for syntrophic metabolism of propionate and butyrate. *Wat. Res.* **102**,
475-484 (2016).
46   Zhang, W., Zhang, J. & Lu, Y. Stimulation of carbon nanomaterials on syntrophic
oxidation of butyrate in sediment enrichments and a defined coculture. *Sci. Rep.* **8**, 12185
(2018).
47   Zhao, Z., Li, Y., Yu, Q. & Zhang, Y. Ferroferric oxide triggered possible direct
interspecies electron transfer between Syntrophomonas and Methanosaeta to enhance
waste activated sludge anaerobic digestion. *Bioresource Technology* **250** (2018).
48   Akashi, H. & Gojobori, T. Metabolic efficiency and amino acid composition in the
proteomes of Escherichia coli and Bacillus subtilis. *Proc Natl Acad Sci U S A* **99**, 3695–
3700 (2002).
49   Conrad, R., Phelps, T. J. & Zeikus, J. G. Gas metabolism evidence in support of
juxtaposition of hydrogen-producing and methanogenic bacteria in sewage sludge and
lake sediments. *Appl. Environ. Microbiol.* **50**, 595-601 (1985).

50   Conrad, R. & Babbel, M. Effect of dilution on methanogenesis, hydrogen turnover and
interspecies hydrogen transfer in anoxic paddy soil. *FEMS* **62**, 21-28 (1989).
51   Conrad, R., Mayer, H.-P. & Wust, M. Temporal change of gas metabolism by hydrogen-
syntrophic methanogenic bacterial associations in anoxic paddy soil. *FEMS Microbiol.*
Lett. **62**, 265-273 (1989).
52   Lovley, D. R. & Klug, M. J. Intermediary metabolism of organic carbon in the sediment
of a eutrophic lake. *Appl. Environ. Microbiol* **43**, 552-560 (1982).
53   Holmes, D. E. *et al.* Electron and proton flux for carbon dioxide reduction in
Methanosarcina barkeri during direct interspecies electron transfer *Fronti. Microbiol.* **9**,
3109 (2018).
54   Coppi, M. V., Leang, C., Sandler, S. J. & Lovley, D. R. Development of a genetic system
for *Geobacter sulfurreducens*. *Appl. Environ. Microbiol.* **67**, 3180-3187 (2001).
55   Nevin, K. P. *et al.* Anode biofilm transcriptomics reveals outer surface components
essential for high current power production in *Geobacter sulfurreducens* fuel cells. *PLoS*
ONE **4**, e5628 (2009).
431