TMS-evoked EEG potentials from prefrontal and parietal cortex: reliability, site specificity, and effects of NMDA receptor blockade

Nigel C. Rogasch[1], Carl Zipser[2], Ghazaleh Darmani[2], Tuomas P. Mutanen[3], Mana Biabani[1], Christoph Zrenner[2], Debora Desideri[2], Paolo Belardinelli[2], Florian Müller-Dahlhaus[2,4]*, Ulf Ziemann[2]*

*Equal contribution

1. Brain and Mental Health Research Hub, Monash Institute of Cognitive and Clinical Neurosciences and Monash Biomedical Imaging, Monash University, Melbourne, Australia.
2. Department of Neurology and Stroke, and Hertie Institute for Clinical Brain Research, Eberhard Karls University of Tübingen, Tübingen, Germany.
3. Centre for Cognitive Neuroimaging, Institute of Neuroscience and Psychology, University of Glasgow, Glasgow, United Kingdom
4. Department of Psychiatry and Psychotherapy, University Medical Center Mainz, Mainz, Germany.

Corresponding author
Nigel C. Rogasch
Monash University
Building 220, Clayton Campus
770 Blackburn Rd.
Clayton, VIC, 3800
Australia
nigel.rogasch@monash.edu

Abbreviations: EEG, electroencephalography; GABA, γ-aminobutyric acid; NMDA, n-methyl-d-aspartate; PAR, parietal cortex; PFC, prefrontal cortex; RMT, resting motor threshold; TEP, TMS-evoked EEG potential; TMS, transcranial magnetic stimulation.

ABSTRACT

Background: Measuring the brain's response to transcranial magnetic stimulation (TMS) with electroencephalography (EEG) offers a unique insight into the local cortical circuits and networks activated following stimulation, particularly in non-motor regions where less is known about TMS physiology. However, the mechanisms underlying TMS-evoked EEG potentials (TEPs) remain largely unknown.

Objective: We assessed TEP reliability, site-specificity, and sensitivity to changes in excitatory neurotransmission mediated by n-methyl-d-aspartate (NMDA) receptors following stimulation of non-motor regions.

Methods: In fourteen male volunteers, resting EEG and TEPs from prefrontal (PFC) and parietal (PAR) cortex were measured before and after administration of either dextromethorphan (an NMDA receptor antagonist) or placebo across two sessions separated by at least a week in a double-blinded pseudo-randomised crossover design.

Results: At baseline, TEPs showed lower within- than between-subject variability for both stimulation sites across sessions, demonstrating the reliability of non-motor TEPs within individuals. There were differences in amplitude between PFC and PAR TEPs across a wide time range (15-250 ms), however the signals were correlated after ~80 ms, suggesting that early peaks reflect site-specific activity, whereas late peaks reflect activity patterns less dependent on the stimulated sites. TEPs were not altered following dextromethorphan compared to placebo, however low frequency resting oscillations were reduced in power.

Conclusions: Our findings suggest that TEPs from PFC and PAR: 1) are reliable within and variable between individuals; 2) reflect stimulation site specific activity across early (<80 ms) time points; and 3) are not sensitive to changes in NMDA receptor-mediated neurotransmission.

Keywords: TMS-EEG, dextromethorphan, NMDA receptor, excitability, prefrontal cortex, parietal cortex

INTRODUCTION

Transcranial magnetic stimulation (TMS) is a brain stimulation method capable of non-invasively activating cortical neurons across the scalp in humans via electromagnetic induction [1]. A single TMS pulse evokes a series of time-locked peaks and troughs in electroencephalographic (EEG) recordings of brain activity [2], which are commonly known as TMS-evoked EEG potentials (TEPs). TEPs are reliable within and between sessions [3–5], are sensitive to changes in TMS parameters such as intensity [4,6] and pulse shape [7], and differ depending on the cortical site stimulated [4,8,9]. In addition, TEPs are sensitive to changes in cortical properties resulting from differing brain states [10–12], plasticity-inducing brain stimulation paradigms [13,14], and brain disorders [15,16]. As such, TMS-EEG is emerging as a powerful method for investigating cortical dynamics in health and disease.

Despite the recent uptake of TMS-EEG within the brain stimulation field, it remains largely unclear what physiological properties underlie the size, shape and distribution of TEPs, thereby limiting their interpretability. Current hypotheses suggest that TEPs primarily reflect fluctuations in cortical excitability resulting from excitatory and inhibitory neurotransmission at the site of stimulation, as well as the propagation of activation through cortical networks following TMS [17]. In support, pharmacological agonists of inhibitory neurotransmission mediated by fast activating γ-aminobutyric acid (GABA)-A receptors given at sub-anaesthetic doses increase the amplitude of early TEPs (e.g. N45) following motor cortex stimulation [18–20], and reduce the propagation of activity following premotor and parietal cortex stimulation at anaesthetic doses [21,22]. Agonists of slow acting GABA-B receptors at sub-anaesthetic doses, on the other hand, increase the amplitude of latter peaks (e.g. N100) following motor cortex stimulation [18]. Although evidence for the sensitivity of single-pulse TEPs to inhibitory neurotransmission is growing, the effect of excitatory neurotransmission on TEPs is less clear. Several studies have linked early motor TEPs between 15-40 ms after TMS with fluctuations in cortical excitability

measured via motor-evoked potentials [6,23–25]. However, TEPs following single-pulse TMS of premotor or parietal cortex are unaffected following anaesthetic doses of ketamine, an n-methyl-d-aspartate (NMDA) receptor antagonist [22]. To date, no studies have assessed the sensitivity of single-pulse TEPs to changes in NMDA receptor mediated neurotransmission while individuals are conscious.

The primary aim of this study was to investigate the contribution of NMDA receptor-mediated neurotransmission to the generation of TEPs following single-pulse TMS in conscious, healthy adults. We measured TEPs following prefrontal (PFC) and parietal (PAR) stimulation before and after a sub-anaesthetic dose of dextromethorphan, an NMDA receptor antagonist, or a placebo in a double-blinded pseudo-randomized crossover design. We hypothesised that early (15-40 ms) TEPs would be reduced following dextromethorphan, but not placebo. Given that TMS-EEG is a relatively new method, particularly outside of motor cortex, we also took advantage of the repeated session experimental design to: 1) characterise within- and between-subject reliability/variability in the spatiotemporal profile of TEPs; and 2) compare differences and similarities between TEPs following stimulation of different sites. As there is currently no consensus on the best way to process TMS-EEG data, we also assessed the impact of different cleaning pipelines on the study outcomes.

MATERIALS AND METHODS

Participants

Fifteen right-handed male participants were recruited for the study. Data from one participant was removed due to a fault in the TMS noise-masking in one condition, leaving a total of fourteen participants (mean age ± S.D. = 28.7± 5 years, range = 21-39 years; see supplementary methods for exclusion criteria). The experimental procedures were approved by the local ethics committee of the Eberhard-Karls-University Medical Faculty, Tübingen (protocol 526/2014BO1), and all participants provided signed, informed consent in accordance with the latest version of the Declaration of Helsinki.

Experimental design

Participants underwent a pseudo-randomised, placebo-controlled, double-blind cross-over experiment to assess the effects of dextromethorphan on TEPs resulting from PFC and PAR stimulation. Dextromethorphan is a non-competitive antagonist of the glutamatergic NMDA receptor, but also interacts with serotonin transporters, sigma-1 receptors, and α3β4 nicotinic acetylcholine receptors [28]. Prior to the experimental sessions, all participants underwent a T1-weighted magnetic resonance imaging (MRI) scan of their brain for use in TMS neuronavigation and EEG electrode position digitisation (supplementary methods). Participants then attended two experimental sessions at least one week apart. During testing, participants were seated comfortably in a chair with hands resting on a pillow in their lap. Baseline measures included: systolic and diastolic blood pressure, resting motor threshold (RMT), two 4 min periods of resting EEG (eyes open and closed; measured to assess the impact of dextromethorphan on resting oscillations), and TEPs following stimulation of PFC and PAR. Following baseline measures, participants ingested either 120 mg of dextromethorphan or placebo (session order pseudorandomised across subjects). After 60 min, blood pressure, resting EEG, and TEP measures were repeated. 60 min was chosen based on dextromethorphan pharmacokinetics,

with blood plasma levels peaking ~60-120 min after drug ingestion [29]. Blood pressure was measured again at the end of the experimental session.

EEG

EEG was recorded from 62 TMS-compatible, c-ring slit electrodes (EASYCAP, Germany) using a TMS-compatible EEG amplifier (BrainAmp DC, BrainProducts GmbH, Germany). During eyes open resting EEG, participants were asked to look at a fixation cross and blink as normal. During eyes closed, participants were asked to close their eyes and avoid going to sleep.

TMS

For TEPs, two sites were stimulated using monophasic TMS (Magstim company, UK): left superior frontal gyrus (PFC; MNI coordinates: -20, 35, 55) and left superior parietal lobule (PAR; -20, -65, 65) (see supplementary methods for TMS details). We deliberately chose sites close to the midline to minimise TMS activation of scalp/facial muscles [30,31]. The TMS coil position was determined and monitored throughout the experiment using frameless stereotaxic neuronavigation co-localised to individual T1-weighted MR scans (TMS Navigator, Localite GmbH, Germany). Coil angle was positioned so that the coil handle ran perpendicular to the underlying gyrus with the handle pointing laterally. As there are currently no standardised methods for determining TMS intensity in non-motor regions, TMS intensity was set to 100% of RMT for each site (supplementary methods). 150 TMS pulses were delivered at a rate of 0.2 Hz ± 25% jitter for each site and the order of sites was randomised at each measurement point. Participants were asked to look at a fixation cross during stimulation and blink as normal. Muscle activity and excessive eye movement were monitored by an experimenter throughout the session and fed back to the participant via a tap on the shoulder if too high.

EEG analyses

Analyses were performed in MATLAB r2017a (MathWorks Inc.) using EEGLAB (v14.1.1) [32], TESA (v0.1.0) [33], FieldTrip (v20170815) [34], Brainstorm (v20180108) [35], and FreeSurfer (v5.3) [36,37] toolboxes, and custom code. All code is available at: (https://github.com/nigelrogasch/DXM_TMS-EEG_paper).

TMS-EEG: As we were interested in early TEP peaks, we developed a novel TMS-EEG cleaning pipeline including two analysis methods designed to recover early TMS-evoked activity (< 45 ms) from TMS-related artifacts; the source-estimate-utilizing noise-discarding (SOUND) [38] algorithm and signal-space projection source-informed reconstruction (SSP-SIR) [39]. See supplementary methods for details and table S1 for number of trials, channels and components removed.

As there is currently no consensus on the best pipeline for cleaning TMS-EEG data, we re-cleaned the data using a pipeline we have used previously [40] (supplementary methods; table S2) and repeated the analyses to assess whether the cleaning procedure impacted the outcomes of the study.

In addition to scalp analysis, we also applied source estimation using two different methods, dipole fitting and minimum-norm estimation (MNE) [41], to assess which cortical regions most likely explained the EEG scalp data. See supplementary methods for details.

Resting EEG: Eyes open and eyes closed resting EEG were cleaned using identical pipelines. See supplementary methods for details and table S3 for number of segments, channels and components removed.

Statistics

TEP variability and comparisons between sites: To assess TEP variability between sessions, absolute TEP amplitude differences averaged over electrodes and time points (15-500 ms) were calculated and compared between baseline TEPs within-subjects (dextromethorphan vs placebo session; 14 comparisons) and between-subjects (91 comparisons) using Mann-Whitney U tests. The analysis was repeated with data collapsed across time and electrodes. To assess differences in TEPs following PFC and PAR stimulation, baseline TEPs were compared between stimulation sites for each condition using cluster-based permutation statistics (cluster threshold: $p < 0.05$ dependent t-test; cluster alpha<0.05 two-tailed; randomisation=5000; time included: 15-250 ms). To assess similarities between stimulation sites, Spearman's correlations were performed on TEP amplitudes across electrodes (scalp) and vertices (source) for each time point, converted to z scores, and compared with baseline measures using Mann-Whitney U tests.

Effects of dextromethorphan on TEPs and resting oscillations: Cluster-based permutation statistics were used to compare changes in TEP amplitude and resting oscillations across time following dextromethorphan and placebo administration, and between conditions by comparing post values subtracted from pre values (cluster threshold: $p < 0.05$ dependent t-test; cluster alpha<0.05 two-tailed; randomisation=5000). TEP analyses included a broad time range (i.e. no *a priori* assumptions about peak times; 15-250 ms), and at six peaks evident following PFC and PAR stimulation (cluster alpha<0.008; Bonferroni corrected; supplementary methods). Data from TEP peaks were also compared using Bayes Factor (BF) analysis to assess evidence for

the null hypothesis that changes in peak amplitudes did not differ following dextromethorphan or placebo (JASP v0.8.1.2; Cauchy prior=0.07; $BF_{01}>3$ taken as moderate evidence; supplementary methods). For resting oscillations, data were averaged into five canonical oscillation bands prior to cluster-based analysis: delta (1-3 Hz); theta (4-7 Hz); alpha (8-12 Hz); beta (13-29 Hz); and gamma (30-45 Hz) (cluster alpha<0.01; Bonferroni corrected).

RESULTS

All experimental procedures were generally well tolerated, with several individuals reporting mild dizziness and one individual nausea following dextromethorphan. These side effects did not affect the subjects' capacity to fully comply with study requirements. There was no difference in RMT at baseline between drug conditions (dextromethorphan=48.4 ± 8% maximum stimulator output, MSO; placebo=47.8 ± 8% MSO; p=0.09).

Within- and between-subject TEP variability

We first compared within- and between-subject differences in baseline TEPs across conditions to assess TEP reliability/variability. Differences in TEPs across sessions within individuals were lower than differences between individuals for both stimulation sites (PFC, $p=4.7 \times 10^{-6}$; PAR, $p=1.1 \times 10^{-4}$; fig. 1A,B). Across space, lower within-subject than between-subject TEP differences were observed across the majority of electrodes for both stimulation sites (fig. 1C,D), whereas across time, lower within-subject TEP differences were strongest between ~30-513 ms following PFC stimulation and ~25-323 ms following PAR stimulation (fig. 1E,F). These findings suggest that the spatiotemporal profile of TEPs are reliable across sessions within individuals, but show variability between individuals (see fig. 2 for an example in 5 subjects).

Figure 1: Within- and between-subject variability in baseline TEPs. *A-B) Mean absolute differences in baseline TEPs (15-500 ms, all electrodes) between the dextromethorphan and placebo condition within- and between-subjects following prefrontal cortex (PFC; A) and parietal cortex (PAR; B) stimulation. * indicates p<0.05 (Mann-Whitney U test). C-D) Topoplots displaying z-scores (Mann-Whitney U tests) comparing within- and between-subject baseline TEP differences at individual electrodes (averaged across time between 15-500 ms) following*

PFC (C) and PAR (D) stimulation. Negative z-scores indicate within-subject TEP differences are less than between-subject TEP differences. White dots indicate p<0.05. E-F) Z-scores (Mann-Whitney U tests) comparing within- and between-subject TEP differences at individual time points (averaged across all electrodes) following PFC (E) and PAR (F) stimulation. Dotted black lines indicate the time of the TMS pulse. Dashed red lines indicate z = ±1.96. Solid red lines indicate p<0.05.

Figure 2: Example of within- and between-subject variability in individual participants.

The top row shows baseline TEPs from the dextromethorphan (blue lines, left topoplots) and placebo (red lines, right topoplots) sessions following PFC stimulation, and the bottom row following PAR stimulation, in five example participants (S5-S9). TEP line plots are taken from an electrode near the site of stimulation (indicated with white dot on topoplots; Fz for PFC stimulation; Pz for PAR stimulation), and TEP topoplots for a representative point in time (indicated with triangles in line plots; 70 ms for PFC; 75 ms for PAR). Both the shape and spatial distribution of the baseline TEPs are more similar within-subjects than they are between-subjects.

Baseline TEPs following PFC and PAR stimulation

Next we assessed the differences and similarities between TEPs following stimulation of different sites. As TEPs were reliable within individuals between sessions, we averaged across baseline conditions to maximise TEP signal strength. Figure 3 shows TEP waveforms and spatial maps following PFC and PAR stimulation averaged across all individuals. When comparing across a broad time window (15-250 ms), TEPs following PFC stimulation differed in amplitude compared with PAR stimulation across all time points (fig. 3C). Despite the amplitude differences, the spatial distribution of TEPs were highly correlated between stimulation sites after ~83 ms (fig. 4A), suggesting that late peaks may represent similar underlying cortical sources regardless the stimulated sites.

Figure 3: Comparison of baseline TEPs between stimulation sites. *Butterfly plots of grand average TEPs across all individuals following prefrontal (PFC; A) and parietal cortex (PAR; B) stimulation at baseline (averaged across conditions). The red dashed line represents the timing of the TMS pulse and the blue triangles the latencies plotted in C and D. C) Topoplots showing the grand average amplitude of TEPs at different time points following PFC (top row), and PAR stimulation (middle row). The bottom row shows t-statistics comparing the amplitude of PFC and PAR stimulation. White and black dots indicate significant negative and positive clusters (cluster-based permutation tests on 15-250 ms; 2 positive clusters [p=0.040, 81-142 ms;*

p=0.006, 148-250 ms]; 1 negative cluster [p=0.002, 15-192 ms]). D) Minimum-norm estimate source maps averaged across participants showing peak activity at each time point in C following PFC (top row) and PAR (bottom row) stimulation. Activity has been thresholded to 85% of maximum activity at each time point. The blue dot represents the target for PFC stimulation and the green dot the target for PAR stimulation.

Figure 4: Spatial correlations between prefrontal (PFC) and parietal (PAR) TEPs.

Spearman correlations comparing the relationship between PFC and PAR TEPs at the scalp (A) and source (B) level for each time point. The thick blue line represents the mean rho values across individuals, and the shaded bars the 95% confidence intervals. The thick red line indicates post stimulation time points where correlations are greater than at equivalent pre stimulation time points (p<0.05; Mann-Whitney U test). Note that rho values were converted to z for statistics, then back to rho for plotting.

To further explore the origin of early and late TEPs, we applied two different source estimation methods: dipole fitting and MNE. For early peaks, the location of the best fitting dipole tended to be closer to the site of stimulation compared to the non-target site (e.g. the PAR when the PFC was stimulated and *vice versa*; table 1). In contrast, the dipole locations corresponding to late peaks were closer to the PAR target regardless of stimulation site. For MNE, estimated source distributions were located close to the site of stimulation for early peaks (25-55 ms; fig. 3D), showed some overlap between stimulation sites for middle peaks (75, 110 ms), and were similar for late peaks (200 ms). Similar to the scalp data, MNE spatial distributions were highly correlated between PFC and PAR TEPs from ~129 ms to ~259 ms (fig. 4B). Taken together, these findings suggest that early TEP peaks reflect neural activation specific to the site of stimulation, whereas late peaks reflect common activation patterns, which differ in amplitude between stimulation sites.

Table 1: *Distance from TMS target sites to best-fitting dipoles at baseline.*

	Distance from target (mm)	Distance from non-target (mm)	Goodness of fit (GoF)	p-value
PFC (15-45 ms)	60 [18-133]	73 [40-103]	0.93 [0.81-0.99]	0.135
PFC (95-125 ms)	80 [24-129]	59 [20-110]	0.88 [0.68-0.99]	0.077
PFC (175-205 ms)	80 [37-123]	**51 [21-122]**	0.84 [0.69-0.99]	0.003
PAR (15-45 ms)	**49 [23-91]**	97 [68-130]	0.93 [0.69-0.99]	4.7×10^{-5}
PAR (95-125 ms)	**52 [22-94]**	87 [51-110]	0.90 [0.79-0.97]	1.5×10^{-4}
PAR (175-205 ms)	**44 [19-85]**	82 [42-130]	0.87 [0.72-0.99]	0.001

NB: *Values in column 1-3 represent the mean [range]. Bold numbers indicate which site was closest to the best fitting dipole (target vs. non-target; $p<0.05$, Mann-Whitney U test). PFC, prefrontal cortex; PAR, parietal cortex.*

Effect of dextromethorphan on TEPs

We next assessed whether dextromethorphan altered TEP amplitude. We could not find any differences in TEP amplitudes across time following either dextromethorphan or placebo for PFC stimulation (all p>0.05), whereas there was a change in PAR TEP amplitude following dextromethorphan (positive cluster, p=0.006, 126-207 ms; negative cluster, p=0.0132, 125-201 ms), but not following placebo (p>0.05). However, these changes were not aligned to TEP peaks (fig. S1) and we could not find any difference between conditions when directly comparing the change in TEP amplitudes following dextromethorphan and placebo for either stimulation site (all p>0.05; 15-250 ms; fig. 5), suggesting the changes observed following dextromethorphan with PAR stimulation were not robust. To ensure that the size of latter clusters was not biasing the analysis against smaller earlier clusters, we reran the analysis averaging across shorter time windows capturing the main TEP peaks, but could not detect any differences across time or between conditions (all p>0.05; Bonferroni corrected; fig. 6). We then ran Bayesian t-tests over ROIs for each peak (determined from baseline data) to assess evidence for the null hypothesis that changes in TEP amplitudes did not differ between conditions. For all comparisons, the BF_{01} was between 1-4, providing weak/moderate evidence that changes in TEP peak amplitude did not differ between dextromethorphan and placebo (table 2).

***Figure 5: TEPs from single electrodes following dextromethorphan (DXM) and placebo
(PBO).*** *A-B) TEPs measured from the Fz electrode following prefrontal cortex (PFC) stimulation
pre and post dextromethorphan and placebo administration. C-D) TEPs measured from the Pz
electrode pre and post dextromethorphan and placebo administration. Thick coloured lines
represent the group mean and shaded colour lines represent 95% confidence intervals.*

Figure 6: Comparison of changes in TEPs following dextromethorphan (DXM) and

placebo (PBO). *Topoplots showing changes in TEP amplitude at peak latencies following*

prefrontal (PFC; A) and parietal cortex (PAR; B) stimulation after dextromethorphan (top row)

and placebo (middle row). Topoplots showing t-statistics (within-subject t-tests) comparing TEP

changes between dextromethorphan and placebo are shown on the bottom row. No significant

differences were observed between conditions (cluster-based permutation tests).

Table 2: *Bayes factors comparing the change in TEP peak amplitude following dextromethorphan (DXM) vs. placebo (PBO).*

	DXM vs PBO	
TEP peaks	**PFC (BF_{01})**	**PAR (BF_{01})**
33, 25	**3.0**	1.3
43, 41	2.5	**3.7**
60, 54	**3.6**	2.6
77, 73	**3.7**	**3.6**
115, 112	**3.6**	1.9
184, 194	**3.4**	**3.0**

NB: *Values in column one represent the mean TEP peak latency for prefrontal cortex (PFC) and parietal cortex (PAR) stimulation respectively. Bold numbers indicate moderate evidence for no difference between conditions.*

Effect of processing pipeline on TEP results

As we used a novel TEP cleaning pipeline, we reran all of the analyses using a more conventional pipeline with two rounds of ICA [33,40]. As with pipeline one, we found low within-subject TEP differences between sessions, differences and similarities in amplitude between stimulation sites, and non-significant effects of dextromethorphan on TEPs using pipeline two (figs. S2-S7; tables S4-S5), indicating that the choice of cleaning pipeline does not impact the main conclusions of the study.

Effect of dextromethorphan on resting oscillations

In addition to TEPs, we also assessed whether dextromethorphan altered resting oscillations. We could not detect any differences in resting oscillations at baseline between sessions (all $p > 0.05$), suggesting that the spatio-spectral profile of oscillations was stable across sessions within individuals. Delta oscillatory power was reduced following dextromethorphan in the eyes open ($p=0.002$) and eyes closed ($p=0.009$) conditions, whereas beta oscillatory power was reduced following placebo in the eyes closed condition only ($p=2.0\times10^{-4}$). When comparing conditions, reductions in delta power tended to be larger following dextromethorphan than placebo for eyes open ($p=0.013$; fig. 7A), although this did not survive correction for multiple comparisons, whereas a reduction in theta power was larger following dextromethorphan than placebo for the eyes closed condition ($p=0.009$; Bonferroni-corrected; fig. 7B). We could not detect differences in oscillatory power changes between dextromethorphan and placebo for any other frequency band (all $p > 0.05$). Taken together, these findings suggest that dextromethorphan reduces power in low frequency oscillations (delta and theta) during resting states.

Figure 7: Comparison of changes in resting oscillations following dextromethorphan (DXM) and placebo (PBO). *Topoplots showing changes in oscillatory power in different frequency bands during eyes open (A) and eyes closed (B) resting conditions following dextromethorphan (top row) and placebo (middle row). Topoplots showing t-statistics (within-subject t-tests) comparing power changes between dextromethorphan and placebo are shown on the bottom row. White dots indicate significant clusters with Bonferroni correction and blue dots uncorrected clusters.*

Effect of dextromethorphan on blood pressure

There was a main effect of time on diastolic blood pressure ($F_{1,13}$=7.9, p=0.006), with pressure increasing across the experiment, however we could not detect any main effect of condition ($F_{1,13}$=0.2, p=0.71), or time × condition interaction ($F_{1,13}$=0.4, p=0.66). We could not detect any main effects of time ($F_{1,13}$=2.4, p=0.13) or condition ($F_{1,13}$=0.0, p=0.90), or a time × condition interaction ($F_{1,13}$=1.1, p=0.34) on systolic pressure, suggesting dextromethorphan and placebo did not have differential effects on blood pressure.

DISCUSSION

TEPs offer unique insight into the effects of TMS on local cortical circuits and networks, however the precise mechanisms reflected by TEPs remain largely unclear. In the current study, we have shown that TEPs from PFC and PAR stimulation are reliable within-individuals, but are variable between-individuals. When comparing TEPs from different stimulation sites, early TEPs (<50 ms) are localised to regions close to the site of stimulation, whereas late peaks (>80 ms) showed common activation patterns independent of the stimulated sites. We also provide weak/moderate evidence that TEPs are not altered by dextromethorphan, suggesting that TMS-evoked EEG responses following single-pulses applied to PFC and PAR are insensitive to changes in NMDA receptor mediated neurotransmission. Our findings confirm the reliability of TEPs for assessing the cortical response to TMS in regions outside the motor cortex, and provide a deeper understanding of the physiological mechanisms reflected by TEPs elicited by prefrontal and parietal cortex stimulation.

Reliability and variability of TEPs

TEPs are generally considered highly reliable within-individuals over short (e.g. hours) to moderate (e.g. weeks) time periods. Indeed, high test-retest reliability has been demonstrated for single-pulse TEPs following stimulation of motor [3,42], premotor [4], dorsolateral prefrontal [3,5], superior parietal [4], and visual cortices [4] using a variety of different metrics. We further confirm the reliability of TEPs following stimulation of superior frontal and superior parietal cortex using a simple metric which takes into account the entire spatio-temporal TEP profile by comparing the mean absolute difference in TEP amplitude within- and between-individuals across sessions (i.e. weeks). In addition to demonstrating within-individual reliability, this finding also suggests that TEPs are variable between individuals (e.g., fig. 2). The sources of this variability are likely multi-faceted, and could reflect individual differences in cortical physiology and anatomy (e.g. balance of excitation/inhibition, shape of gyri/sulci) and/or experimental

factors (e.g., success of sensory masking, relative stimulation intensity). Understanding what contributes to this between-individual variability in TEP profile will likely provide important insights into how individual physiology shapes the local and global cortical response to TMS from regions outside of the motor cortex.

Dependence of TEPs on stimulation site

Studies directly comparing TEPs following stimulation of different cortical sites have shown both differences and similarities in the local response profile and the cortical networks activated by TMS. For instance, the local oscillatory profile following TMS appears to differ along an anterior-posterior gradient, with frontal sites oscillating at higher frequencies than parietal and occipital sites [15,43]. Furthermore, stimulation of motor cortex results in larger TEPs than non-motor regions [8], with a unique oscillatory profile [9]. The broader cortical networks activated following TMS also differ depending on the stimulation site, even within stimulation of functionally-related regions [44].

Despite the differences in TEPs following stimulation of different cortical sites, several studies have reported similarities in TEPs regardless of the target site, especially at periods ~100 ms, and ~200 ms following stimulation [45,46]. These periods coincide with auditory-evoked potentials resulting from the TMS clicking noise, and bone-conducted sensory responses from coil vibration [47,48]. To minimise sensory contamination, noise-masking is typically provided during stimulation (e.g. white noise played through headphones) and/or foam is placed under the coil to minimise vibration [10,49,50]. Even with such measures, several recent studies have reported that TEPs are highly correlated with control conditions (e.g. TMS of the shoulder or electrical stimulation of the scalp) [51–53].

In the current study, we applied auditory masking, and stimulated sites close to the midline to minimise sensation resulting from the activation of scalp muscles with TMS. We found differences in TEP amplitude following stimulation of PFC and PAR at the scalp level across a broad time range (15-250 ms). However the spatial distribution of the TEPs were highly correlated between sites from ~80 ms onwards. Source estimation using two different methods (dipole fitting and MNE) suggested that the early TEP response (15-55 ms) reflected activity from regions close to the site of stimulation, whereas late TEP peaks reflected activity from partially or fully overlapping central regions regardless of stimulation site. Taken together, one possibility is that some of the late TMS responses measured with EEG reflects indirect activation of the cortex from sensory input, regardless of efforts to minimise TMS-evoked sensation and audition. Another possibility for explaining similarity of spatial distribution of late TEPs in the present study is that areas of the fronto-parietal network have been stimulated and that this may lead to common network activation at late time points.

Excitatory neurotransmission and TEPs

Pharmacological studies targeting inhibitory receptors have provided evidence that certain TEP peaks around 45 and 100 ms are sensitive to changes in GABAergic neurotransmission [18,19], whereas peaks at 30 ms, 45 ms and 180 ms are sensitive to anti-epileptic drugs targeting voltage-gated sodium channels [54,55]. However, the sensitivity of TEPs to changes in excitatory neurotransmission is less clear. Several lines of indirect evidence suggest that early TEP peaks between 15 to 40 ms may reflect excitatory neurotransmission. First, excitatory postsynaptic potentials generated by NMDA receptor activation peak at ~15-40 ms in rodents following electrical stimulation of the neocortex [56,57], latencies which are similar to early TEP peaks. Second, the amplitude of early TEP peaks in motor cortex (N15, P30) correlate with fluctuations in MEP amplitude (which reflect activation of the corticomotoneuronal system) [23], and show similar changes with TMS intensity [6], coil angle [6,24], and paired pulse paradigms

[25] to MEPs. Third, early TEP peaks increase in amplitude with time awake [58], consistent with changes in cortical excitability during wakefulness. Collectively, this body of evidence has led to the hypothesis that early TEP peaks may reflect excitatory postsynaptic potentials following TMS, possibly mediated by NMDA receptors [17].

We could not find any evidence that changes in TEPs differed following administration of the NMDA receptor antagonist dextromethorphan compared to placebo at any time point following stimulation of either site. The lack of change in TEPs following dextromethorphan was not impacted by our choice of statistical approach or the TEP processing pipeline. Although our sample was relatively small (n=14), Bayes factor analysis provided moderate evidence for the null hypothesis in 8 of the 12 peaks tested across sites, and weak evidence in the other peaks, suggesting that we were adequately powered to test our hypothesis. In line with our findings, TEPs following single-pulse TMS to premotor and parietal cortex are unaffected by anaesthetic doses of ketamine [22], another NMDA receptor antagonist, suggesting that single-pulse TEPs are insensitive to changes in NMDA receptor mediated neurotransmission. As NMDA receptors are dependent both on glutamatergic binding and depolarisation of the postsynaptic neuron, it is possible that a single TMS pulse is not sufficient to open NMDA receptors. Instead, paired-pulse TMS-EEG paradigms at intervals between 10-40 ms may be required to observe NMDA receptor mediated neurotransmission, similar to intracortical facilitation paradigms measured with MEPs in motor cortex [59].

A potential limitation of the current study is the dose of dextromethorphan provided (120 mg). High doses of dextromethorphan (>300 mg/70 kg) are required to produce hallucinations [60] and cognitive impairment [61], which are hallmarks of the effects of other NMDA receptor antagonists, such as ketamine and phencyclidine. As such, it is possible that the dose provided here was too low to provide adequate antagonism of the NMDA receptor. However, we did

observe reductions in low-frequency oscillations in the resting EEG, which are partially consistent with changes observed following sub-anesthetic, psychotomimetic doses of ketamine [62]. Furthermore, dextromethorphan at similar doses blocks plasticity assessed via changes in single-pulse TEP amplitude following paired associative stimulation of the DLPFC [63], a plasticity-inducing protocol dependent on glutamatergic neurotransmission mediated by NMDA receptors [64]. Another potential limitation is that we only tested TEPs at one intensity. The effect of certain drugs can impact TEPs in a way which is dependent on stimulation intensity [65]. As such, future studies assessing drug effects on TEPs should take into account a range of stimulation intensities.

CONCLUSIONS

Our findings provide evidence that single-pulse TEPs following stimulation of prefrontal and parietal cortex in conscious humans are not sensitive to changes in excitatory neurotransmission mediated by NMDA receptors. However, TEPs from these cortical regions are reliable within-individuals, and the early TEP peaks provide information specific to the site of stimulation, whereas late TEPs reflect activity less dependent on the stimulated sites. Future work using pharmacological agents targeting different excitatory and inhibitory receptor types is required to disentangle the physiological mechanisms contributing to early TEPs following TMS, and to test if these pharmacological effects are different when stimulating different cortical sites.

ACKNOWLEDGMENTS

The authors wish to thank Dr. Ben D. Fulcher for advice on analysis.

FUNDING

This work was supported by the National Health and Medical Research Council of Australia [grant number: GNT1072057], and the Deutscher Akademischer Austauschdienst (German Academic Exchange Service).

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

[1]   Barker AT, Jalinous R, Freeston IL. Non-invasive magnetic stimulation of human motor cortex. Lancet 1985;325:1106–7.

[2]   Ilmoniemi RJ, Virtanen J, Ruohonen J, Karhu J, Aronen HJ, Näätänen R, et al. Neuronal responses to magnetic stimulation reveal cortical reactivity and connectivity. Neuroreport 1997;8:3537–40.

[3]   Lioumis P, Kicić D, Savolainen P, Mäkelä JP, Kähkönen S. Reproducibility of TMS-Evoked EEG responses. Hum Brain Mapp 2009;30:1387–96.

[4]   Casarotto S, Romero Lauro LJ, Bellina V, Casali AG, Rosanova M, Pigorini A, et al. EEG responses to TMS are sensitive to changes in the perturbation parameters and repeatable over time. PLoS One 2010;5:e10281.

[5]   Kerwin LJ, Keller CJ, Wu W, Narayan M, Etkin A. Test-retest reliability of transcranial magnetic stimulation EEG evoked potentials. Brain Stimul 2018;11:536–44.

[6]   Komssi S, Kähkönen S, Ilmoniemi RJ. The effect of stimulus intensity on brain responses evoked by transcranial magnetic stimulation. Hum Brain Mapp 2004;21:154–64.

[7]   Casula EP, Rocchi L, Hannah R, Rothwell JC. Effects of pulse width, waveform and current direction in the cortex: A combined cTMS-EEG study. Brain Stimul 2018;11:1063–70.

[8]   Kähkönen S, Wilenius J, Komssi S, Ilmoniemi RJ. Distinct differences in cortical reactivity of motor and prefrontal cortices to magnetic stimulation. Clin Neurophysiol 2004;115:583–8.

[9]   Fecchio M, Pigorini A, Comanducci A, Sarasso S, Casarotto S, Premoli I, et al. The spectral features of EEG responses to transcranial magnetic stimulation of the primary motor cortex depend on the amplitude of the motor evoked potentials. PLoS One 2017;12:e0184910.

[10]   Massimini M, Ferrarelli F, Huber R, Esser SK, Singh H, Tononi G. Breakdown of cortical effective connectivity during sleep. Science 2005;309:2228–32.

[11]   Morishima Y, Akaishi R, Yamada Y, Okuda J, Toma K, Sakai K. Task-specific signal transmission from prefrontal cortex in visual selective attention. Nat Neurosci 2009;12:85–91.

[12]   Johnson JS, Kundu B, Casali AG, Postle BR. Task-dependent changes in cortical excitability and effective connectivity: a combined TMS-EEG study. J Neurophysiol 2012;107:2383–92.

[13]   Chung SW, Lewis BP, Rogasch NC, Saeki T, Thomson RH, Hoy KE, et al. Demonstration of short-term plasticity in the dorsolateral prefrontal cortex with theta burst stimulation: A TMS-EEG study. Clin Neurophysiol 2017;128:1117–26.

[14]   Hill AT, Rogasch NC, Fitzgerald PB, Hoy KE. Effects of prefrontal bipolar and high-definition transcranial direct current stimulation on cortical reactivity and working memory in healthy adults. Neuroimage 2017;152:142–57.

[15]   Ferrarelli F, Sarasso S, Guller Y, Riedner BA, Peterson MJ, Bellesi M, et al. Reduced natural oscillatory frequency of frontal thalamocortical circuits in schizophrenia. Arch Gen Psychiatry 2012;69:766–74.

[16]   Rosanova M, Gosseries O, Casarotto S, Boly M, Casali AG, Bruno M-A, et al. Recovery of cortical effective connectivity and recovery of consciousness in vegetative patients. Brain 2012;135:1308–20.

[17]   Rogasch NC, Fitzgerald PB. Assessing cortical network properties using TMS-EEG. Hum Brain Mapp 2013;34:1652–69.

[18]   Premoli I, Castellanos N, Rivolta D, Belardinelli P, Bajo R, Zipser C, et al. TMS-EEG signatures of GABAergic neurotransmission in the human cortex. J Neurosci 2014;34:5603–12.

[19]   Darmani G, Zipser CM, Böhmer GM, Deschet K, Müller-Dahlhaus F, Belardinelli P, et al. Effects of the Selective α5-GABAAR Antagonist S44819 on Excitability in the Human Brain:

A TMS-EMG and TMS-EEG Phase I Study. J Neurosci 2016;36:12312–20.

[20] Premoli I, Király J, Müller-Dahlhaus F, Zipser CM, Rossini P, Zrenner C, et al. Short-interval and long-interval intracortical inhibition of TMS-evoked EEG potentials. Brain Stimul 2018;11:818–27.

[21] Ferrarelli F, Massimini M, Sarasso S, Casali A, Riedner BA, Angelini G, et al. Breakdown in cortical effective connectivity during midazolam-induced loss of consciousness. Proc Natl Acad Sci U S A 2010;107:2681–6.

[22] Sarasso S, Boly M, Napolitani M, Gosseries O, Charland-Verville V, Casarotto S, et al. Consciousness and Complexity during Unresponsiveness Induced by Propofol, Xenon, and Ketamine. Curr Biol 2015;25:3099–105.

[23] Mäki H, Ilmoniemi RJ. The relationship between peripheral and early cortical activation induced by transcranial magnetic stimulation. Neurosci Lett 2010;478:24–8.

[24] Bonato C, Miniussi C, Rossini PM. Transcranial magnetic stimulation and cortical evoked potentials: a TMS/EEG co-registration study. Clin Neurophysiol 2006;117:1699–707.

[25] Rogasch NC, Daskalakis ZJ, Fitzgerald PB. Mechanisms underlying long-interval cortical inhibition in the human motor cortex: a TMS-EEG study. J Neurophysiol 2013;109:89–98.

[26] Smith MJ, Keel JC, Greenberg BD, Adams LF, Schmidt PJ, Rubinow DA, et al. Menstrual cycle effects on cortical excitability. Neurology 1999;53:2069–2069.

[27] Rossi S, Hallett M, Rossini PM, Pascual-Leone A, Safety of TMS Consensus Group. Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. Clin Neurophysiol 2009;120:2008–39.

[28] Taylor CP, Traynelis SF, Siffert J, Pope LE, Matsumoto RR. Pharmacology of dextromethorphan: Relevance to dextromethorphan/quinidine (Nuedexta®) clinical use. Pharmacol Ther 2016;164:170–82.

[29] Kazis A, Kimiskidis V, Niopas I. Pharmacokinetics of dextromethorphan and dextrorphan in epileptic patients. Acta Neurol Scand 1996;93:94–8.

[30] Rogasch NC, Thomson RH, Daskalakis ZJ, Fitzgerald PB. Short-Latency Artifacts Associated with Concurrent TMS–EEG. Brain Stimul 2013;6:868–76.

[31] Mutanen T, Mäki H, Ilmoniemi RJ. The effect of stimulus parameters on TMS-EEG muscle artifacts. Brain Stimul 2013;6:371–6.

[32] Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. J Neurosci Methods 2004;134:9–21.

[33] Rogasch NC, Sullivan C, Thomson RH, Rose NS, Bailey NW, Fitzgerald PB, et al. Analysing concurrent transcranial magnetic stimulation and electroencephalographic data: A review and introduction to the open-source TESA software. Neuroimage 2017;147:934–51.

[34] Oostenveld R, Fries P, Maris E, Schoffelen J-M. FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell Neurosci 2011;2011:156869.

[35] Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM. Brainstorm: a user-friendly application for MEG/EEG analysis. Comput Intell Neurosci 2011;2011:879716.

[36] Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. Neuroimage 1999;9:179–94.

[37] Fischl B, Sereno MI, Dale AM. Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. Neuroimage 1999;9:195–207.

[38] Mutanen TP, Metsomaa J, Liljander S, Ilmoniemi RJ. Automatic and robust noise suppression in EEG and MEG: The SOUND algorithm. Neuroimage 2018;166:135–51.

[39] Mutanen TP, Kukkonen M, Nieminen JO, Stenroos M, Sarvas J, Ilmoniemi RJ. Recovering TMS-evoked EEG responses masked by muscle artifacts. Neuroimage 2016;139:157–66.

[40] Rogasch NC, Thomson RH, Farzan F, Fitzgibbon BM, Bailey NW, Hernandez-Pavon JC, et

al. Removing artefacts from TMS-EEG recordings using independent component analysis: importance for assessing prefrontal and motor cortex network properties. Neuroimage 2014;101:425–39.

[41] Hämäläinen MS, Ilmoniemi RJ. Interpreting magnetic fields of the brain: minimum norm estimates. Med Biol Eng Comput 1994;32:35–42.

[42] Premoli I, Rivolta D, Espenhahn S, Castellanos N, Belardinelli P, Ziemann U, et al. Characterization of GABAB-receptor mediated neurotransmission in the human cortex by paired-pulse TMS-EEG. Neuroimage 2014;103:152–62.

[43] Rosanova M, Casali A, Bellina V, Resta F, Mariotti M, Massimini M. Natural frequencies of human corticothalamic circuits. J Neurosci 2009;29:7679–85.

[44] Garcia JO, Grossman ED, Srinivasan R. Evoked potentials in large-scale cortical networks elicited by TMS of the visual cortex. J Neurophysiol 2011;106:1734–46.

[45] Du X, Choa F-S, Summerfelt A, Rowland LM, Chiappelli J, Kochunov P, et al. N100 as a generic cortical electrophysiological marker based on decomposition of TMS-evoked potentials across five anatomic locations. Exp Brain Res 2017;235:69–81.

[46] Harquel S, Bacle T, Beynel L, Marendaz C, Chauvin A, David O. Mapping dynamical properties of cortical microcircuits using robotized TMS and EEG: Towards functional cytoarchitectonics. Neuroimage 2016;135:115–24.

[47] Nikouline V, Ruohonen J, Ilmoniemi RJ. The role of the coil click in TMS assessed with simultaneous EEG. Clin Neurophysiol 1999;110:1325–8.

[48] Tiitinen H, Virtanen J, Ilmoniemi RJ, Kamppuri J, Ollikainen M, Ruohonen J, et al. Separation of contamination caused by coil clicks from responses elicited by transcranial magnetic stimulation. Clin Neurophysiol 1999;110:982–5.

[49] ter Braack EM, de Vos CC, van Putten MJAM. Masking the Auditory Evoked Potential in TMS-EEG: A Comparison of Various Methods. Brain Topogr 2015;28:520–8.

[50] Ilmoniemi RJ, Hernandez-Pavon JC, Makela NN, Metsomaa J, Mutanen TP, Stenroos M, et al. Dealing with artifacts in TMS-evoked EEG. Conf Proc IEEE Eng Med Biol Soc 2015;2015:230–3.

[51] Conde V, Tomasevic L, Akopian I, Stanek K, Saturnino GB, Thielscher A, et al. The non-transcranial TMS-evoked potential is an inherent source of ambiguity in TMS-EEG studies. Neuroimage 2018. doi:10.1016/j.neuroimage.2018.10.052.

[52] Herring JD, Thut G, Jensen O, Bergmann TO. Attention Modulates TMS-Locked Alpha Oscillations in the Visual Cortex. J Neurosci 2015;35:14435–47.

[53] Gordon PC, Desideri D, Belardinelli P, Zrenner C, Ziemann U. Comparison of cortical EEG responses to realistic sham versus real TMS of human motor cortex. Brain Stimul 2018;11:1322–30.

[54] Premoli I, Biondi A, Carlesso S, Rivolta D, Richardson MP. Lamotrigine and levetiracetam exert a similar modulation of TMS-evoked EEG potentials. Epilepsia 2017;58:42–50.

[55] Darmani G, Bergmann TO, Zipser C, Baur D, Müller- Dahlhaus F, Ziemann U. Effects of antiepileptic drugs on cortical excitability in humans. A TMS-EMG and TMS-EEG study. Hum Brain Mapp 2018.

[56] Sutor B, Hablitz JJ. EPSPs in rat neocortical neurons in vitro. I. Electrophysiological evidence for two distinct EPSPs. J Neurophysiol 1989;61:607–20.

[57] Sutor B, Hablitz JJ. EPSPs in rat neocortical neurons in vitro. II. Involvement of N-methyl-D-aspartate receptors in the generation of EPSPs. J Neurophysiol 1989;61:621–34.

[58] Huber R, Mäki H, Rosanova M, Casarotto S, Canali P, Casali AG, et al. Human cortical excitability increases with time awake. Cereb Cortex 2013;23:332–8.

[59] Ziemann U, Chen R, Cohen LG, Hallett M. Dextromethorphan decreases the excitability of the human motor cortex. Neurology 1998;51:1320–4.

[60] Reissig CJ, Carter LP, Johnson MW, Mintzer MZ, Klinedinst MA, Griffiths RR. High doses of dextromethorphan, an NMDA antagonist, produce effects similar to classic

hallucinogens. Psychopharmacology 2012;223:1–15.

[61] Carter LP, Reissig CJ, Johnson MW, Klinedinst MA, Griffiths RR, Mintzer MZ. Acute cognitive effects of high doses of dextromethorphan relative to triazolam in humans. Drug Alcohol Depend 2013;128:206–13.

[62] de la Salle S, Choueiry J, Shah D, Bowers H, McIntosh J, Ilivitsky V, et al. Effects of Ketamine on Resting-State EEG Activity and Their Relationship to Perceptual/Dissociative Symptoms in Healthy Humans. Front Pharmacol 2016;7:348.

[63] Salavati B, Daskalakis ZJ, Zomorrodi R, Blumberger DM, Chen R, Pollock BG, et al. Pharmacological Modulation of Long-Term Potentiation-Like Activity in the Dorsolateral Prefrontal Cortex. Front Hum Neurosci 2018;12:155.

[64] Stefan K, Kunesch E, Benecke R, Cohen LG, Classen J. Mechanisms of enhancement of human motor cortex excitability induced by interventional paired associative stimulation. J Physiol 2002;543:699–708.

[65] Premoli I, Costantini A, Rivolta D, Biondi A, Richardson MP. The Effect of Lamotrigine and Levetiracetam on TMS-Evoked EEG Responses Depends on Stimulation Intensity. Front Neurosci 2017;11:585.