8   # Dynamic properties of internal noise
9   # probed by modulating binocular rivalry
10
11
12
13 *Short title*: Internal noise in binocular rivalry
14
15
16

17 *Authors*:   Daniel H. Baker[a] (Orcid ID: 0000-0002-0161-443X)
18   Bruno Richard[a,b] (Orcid ID: 0000-0003-4297-2492)
19
20
21
22 *Author affiliations*:
23   a) Department of Psychology, University of York, Heslington, York, United Kingdom,
24   YO10 5DD
25   b) Department of Mathematics and Computer Science, Rutgers University – Newark,
26   Newark, New Jersey, USA, 07102
27
28
29
30 *Corresponding author:* Daniel H Baker, Department of Psychology, University of York,
31 Heslington, York, United Kingdom, YO10 5DD. Email: daniel.baker@york.ac.uk

32
33
34

35 *Keywords*: binocular rivalry; double pass consistency; internal noise; computational
36 modelling; bistable stimuli.

37
38

39
40
41
42

43   *Abstract*
44
Neural systems are inherently noisy, and this noise can affect our perception from moment
to moment. This is particularly apparent in binocular rivalry, where our perception of
competing stimuli shown to the left and right eyes alternates over time in a seemingly random
fashion. We investigated internal noise using binocular rivalry by modulating rivalling stimuli
using dynamic sequences of external noise of various rates and amplitudes. As well as
measuring the effect on dominance durations, we repeated each external noise sequence
twice, and assessed the consistency of percepts across repetitions. External noise
modulations with standard deviations above 4% contrast increased consistency scores above
baseline, and were most effective at 1/8Hz. A computational model of rivalry in which internal
noise has a $1/f$ (pink) temporal amplitude spectrum, and a standard deviation of 16%,
provided the best account of our data, and was able to correctly predict perception in
additional conditions. Our novel technique provides detailed estimates of the dynamic
properties of internal noise during binocular rivalry, and by extension the stochastic processes
that drive our perception and other types of spontaneous brain activity.

64   *Significance statement*
65
Although our perception of the world appears constant, sensory representations are variable
because of the 'noisy' nature of biological neurons. Here we used a binocular rivalry
paradigm, in which conflicting images are shown to the two eyes, to probe the properties of
this internal variability. Using a novel paradigm in which the contrasts of rivalling stimuli are
modulated by two independent external noise streams, we infer the amplitude and character
of this internal noise. The temporal amplitude spectrum of the noise has a 1/f spectrum,
similar to that of natural visual input, and consistent with the idea that the visual system
evolved to match its environment.

Introduction

Despite appearing constant, our sensory perception fluctuates from moment to moment because of the non-deterministic nature of biological neurons. This 'internal noise' operates at multiple timescales, and affects our decisions about sensory information. Internal noise is particularly apparent in bistable phenomena such as binocular rivalry, in which our perception of conflicting images shown to the two eyes fluctuates over time in a stochastic fashion. Because phenomena like rivalry make otherwise invisible processes available to conscious perception, they provide a useful tool for probing the properties of internal noise.

In a typical rivalry experiment, participants view sine wave grating patterns with orthogonal orientations in the left and right eyes. They are asked to report which stimulus they perceive at each moment in time by continuously pressing a response button that corresponds to the perceived orientation. Histograms of the durations for which each image remains dominant typically have positive skew, corresponding to a gamma distribution (or a normal distribution on logarithmic axes). Computational models of rivalry (e.g. 1–4) have successfully explained the precise pattern of percepts reported by assuming the presence of three key processes: inhibition between neurons representing the two stimuli, adaptation to the dominant stimulus, and noise. Inhibitory properties have been investigated using dichoptic masking paradigms (5–7) and by varying the properties of rivalling stimuli (8–11), and there is direct evidence of adaptation during a period of dominance (12). However, comparatively little is known about the precise properties of the noise, as there have been few attempts to investigate it directly.

One exception is a study that aimed to influence rivalry alternations by periodically changing the contrast of the rivalling stimuli. Kim, Grabowecky and Suzuki (1) used a square wave temporal modulation to alter the contrast of rivalling stimuli in antiphase (i.e. one stimulus increased in contrast and the other decreased at the same time) at a range of temporal frequencies from 0.28Hz to 2.48Hz. This manipulation caused a peak in the histogram of dominance durations at the half period of the modulation frequency. The increase was greatest when the half period was 600ms, a duration corresponding to the peak of the histogram for unmodulated rivalry using the same stimuli. Furthermore, there were additional peaks at odd integer harmonics of the modulation frequency. The authors consider this to be evidence of a stochastic resonance effect, and support this with a computational model of rivalry alternations.

Here we modify this approach by modulating the contrast of rivalling stimuli using two independent dynamic noise sequences instead of square wave modulations (see Figure 1). As well as measuring the effect on dominance durations, this design allows us to reverse correlate the participant's reported percept with the timecourse of the external noise. In addition, we can use the same noise sequences multiple times, and measure the consistency of the participants' percepts in a dynamic version of the 'double pass' paradigm (13, 14). If the external noise sequences were entirely determining perception, responses should be identical across the two repetitions. On the other hand, if the external noise sequences have no influence on perception then the similarity of responses will be determined by internal noise, and response consistency will be that expected by chance. By manipulating the variance and temporal frequency content of the noise sequences, we can investigate

122 properties of the internal noise that influences rivalry alternations. We interpret the results
123 with reference to an established computational model of rivalry proposed by Wilson (3, 4)
124 (see Figure 2), to which we add different types of internal noise.
125

126
Figure 1: Methodological details. Panel (a) shows example stimuli with conflicting orientations, surrounded by a
Voronoi texture to aid binocular fusion. Panel (b) shows example waveforms used to modulate stimulus
contrasts at the five temporal frequencies used in the experiment. Panel (c) shows example trial timecourses for
two repetitions of an unmodulated condition (left) and a modulated condition (right). Red (green) regions in the
lower two plots indicate periods of time when the left (right) eye's stimulus was perceived. Note that in the
example on the right, percepts closely followed the physical contrast with a slight lag.
133

Figure 2. Model details. (a) Model diagram of the two competing units. Each receives as input an independent white noise stream, bandpass filtered at one of five different temporal frequencies (see Methods). The minimum rivalry model (4) defines the oscillatory behaviour of rivalry between two units with self-adaptation and mutual inhibition. We include additive internal independent monocular noise in our model, marked by the (+) symbol. (b) Examples of the five different internal noise spectral slopes ($\alpha = 0 - 2.0$) of the model for the left (green) and right (red) responding units. Noise streams with steeper slopes have an increased relative amplitude of low temporal frequencies relative to high, which leads to slower changes in the noise amplitude. (c) Example oscillatory behaviour of the model for a given trial (60s). The colours represent the responses of the left (green) and right (red) responding units.

Results

External noise strongly modulates binocular rivalry alternations

In the absence of any noise modulations, binocular rivalry produced a typical histogram of dominance durations with a positive skew (see grey curve in Figure 3ai), and a mean of 2.7 seconds. A 5x5 repeated measures ANOVA indicated that the mean dominance duration depended on both temporal frequency ($F(4,16)=34.43$, $p<0.001$, $\eta_p^2=0.90$) and modulation contrast ($F(4,16)=8.15$, $p<0.01$, $\eta_p^2=0.67$), and also showed that the two variables interacted ($F(16,64)=18.01$, $p<0.001$, $\eta_p^2=0.82$). The histograms in Figure 3a show that at low temporal frequencies, strong contrast modulation resulted in slightly more long-duration percepts (an

156 increase in positive skew), whereas at high temporal frequencies the peak of the histogram
157 shifted leftwards. These patterns were reflected in both the change in means (Figure 3b) and
158 also the shift in the autocorrelation functions (Figure 3c), such that high temporal frequencies
159 (e.g. the purple curve) had a shorter lag than long ones (e.g. the red curve). The functions in
160 Figure 3b begin to diverge at a contrast of around 4%.
161
We also cross-correlated the noise time course (difference between left and right eye
contrasts for the 16% contrast modulation conditions pooled across all temporal frequencies)
with the participants' responses (Figure 3d). This revealed a mean response lag of 583ms,
somewhat faster than estimates from previous studies (8). The mean cross-correlation
coefficient at this time point was 0.35, indicating that a substantial proportion of the variance
in participant percepts was predictable from the changes in stimulus contrast. Functions for
individual participants are shown by the thin traces in Figure 3d, and are similar to the mean.
169

170
Figure 3: Traditional rivalry measures for all conditions, averaged (or pooled) across all participants (N=5). Panel
(a) shows histograms of pooled dominance durations at five temporal frequencies (i-v) and a range of contrast
levels (standard deviations of 0 – 16%, increasing down each plot). The grey histogram, duplicated in each plot,
shows the baseline condition with no contrast modulation. For low temporal frequency, high contrast
modulations, there were more very long dominance periods (the positive skew of the red histogram increases).
For high temporal frequency, high contrast modulations there were more short dominance periods, and the
histograms shifted left. Panel (b) shows mean dominance durations for all conditions, plotted as a function of
modulation contrast. The grey horizontal line shows the baseline (no modulation) condition. Error bars (and
dotted lines) show ±1SE across participants. Panel (c) shows autocorrelation functions averaged across
participants for the baseline condition (grey curve) and the highest contrast modulation at each temporal
frequency (curves, see panels a,b for colour legends). Panel (d) shows the cross correlation between the
difference in noise modulations at the highest modulation contrast, averaged across all modulation frequencies.
The thin grey lines denote individual participants and the thick black line is the average.
184
Next, we calculated the consistency of responses across pairs of presentations of identical
noise streams. In the absence of any noise modulation, the mean consistency was slightly
above the expected baseline of 0.5, having a value of 0.53 (horizontal grey lines in Figure 4).
The most likely explanation for this is that slight eye dominances or biases towards one or

other stimulus will increase the consistency across repetitions, however the effect is very
small. For conditions where the stimulus contrast was modulated, a 5x5 repeated measures
ANOVA indicated that the response consistency depended on both temporal frequency
(F(4,16)=9.90, p<0.001, η_p^2=0.71) and modulation contrast (F(4,16)=28.81, p<0.001,
η_p^2=0.88), as well as the interaction between the two variables (F(16,64)=3.55, p<0.001,
η_p^2=0.47). These effects are shown in Figure 4, which plots the same data twice as a function
of either modulation contrast (Fig 4a) or temporal frequency (Fig 4b). The general trends are
that consistency increases with contrast, and at each contrast is strongest for the 1/8Hz
temporal frequency (shown in green). The maximum consistency was 0.72, for the 1/8Hz, 16%
contrast condition, which is particularly noteworthy given that this temporal frequency had
the weakest influence on dominance durations (see green points in Figure 3b). Consistency
exceeded baseline for the 1/8Hz condition at around 4% modulation contrast (green
diamonds in Figure 4).
202

203
Figure 4: Response consistency across two passes through the experiment. The same data are plotted in both
panels, as a function of modulation contrast (a) or temporal frequency (b). In each panel, the thick grey line
represents the baseline (no modulation) condition, colours represent different temporal frequencies, and
symbol types represent different contrasts. All data points are averaged across participants, with error bars
indicating ±1SE of the mean. The dashed horizontal line at y=0.5 indicates a theoretical baseline in the absence
of any response bias or eye dominance effects.
210

211   *A computational model with pink internal noise describes the human results*
212

We first investigated how the amplitude of internal noise, and the spectral slope (α), affected
model behaviour. We selected a single stimulus condition (stimulus noise frequency of 1/8Hz
and amplitude of 16%) and ran the model with a range of internal noise contrast levels (SD =
1 − 64%) at five different spectral slopes (α = 0 − 2). The results of our simulations on
dominance duration and response consistency are shown in Figure 5a(i-v), with the
equivalent human data plotted in green for comparison. For all spectral slopes, as internal
noise contrast increased it more strongly affected rivalry alternations. This is shown by the
change in dominance duration (Figure 5b; increases for steep slopes and decreases for
shallow slopes), and response consistency (Figure 5c), which decreased as responses became
increasingly dominated by internal noise.

Figure 5: Summary of model behaviour for internal noise amplitude and spectral slope estimation. (a) The histograms of dominance durations for each spectral slope (α = 0.0 – 2.0) and contrast (SD = 1% – 64%) values. Within each subplot, the uppermost (green shaded) histogram is the equivalent human data for a stimulus temporal frequency of 1/8Hz and a contrast modulation of SD = 16%. The solid vertical green line marks the average dominance duration for human observers. Histograms below show model dominance duration distributions for each internal noise contrast value. (b) Average dominance durations of the model for each spectral slope (coloured lines). The green line and shaded area mark human average dominance duration and ±1SE of the mean, respectively. Average dominance duration was affected by internal noise once its contrast reached 4%. Noise with steeper slopes (α = 1.5-2.0) increased mean dominance duration as a function of noise contrast, while noise with shallower slopes decreased mean dominance duration. (c) Response consistency decreased as a function of internal noise contrast for all spectral slopes. The green line and shaded area mark human observer average consistency and ±1SE of the mean, respectively. For all α>0, response consistency reached human levels at an internal noise contrast of 16%.

We can use the joint dominance durations and consistency scores to rule out several types of internal noise. White internal noise (α = 0) is not viable because there is no internal noise level for which both durations and consistency are close to human levels. Internal noise with steep amplitude slopes (α > 1) produces sensible consistency scores, but dominance durations become too long. This leaves slopes of α = 0.5 and α = 1, for which an internal noise contrast of around 16% gives a good approximation to the human data. We performed full simulations for all noise spectral slopes with this contrast. A slope of α = 1 was the best predictor of the human data, so these simulations are discussed in the main text, with simulations of other spectral slopes are presented in Supplementary Figures S1 and S2.

The histograms of dominance durations, mean dominance duration and response consistency of the model simulations for all 26 stimulus conditions are shown in Figure 6. The model replicated the pattern of human data shown in Figures 3 & 4 remarkably well. The histograms of dominance durations of the model (Figure 6a i-v) show similar trends to those of human observers (Figure 3a). Slow modulation frequencies (1/16Hz and 1/8Hz) increased positive skew at high modulation contrasts (Figure 6a i-ii), while higher modulation frequencies shifted the peak of the dominance duration histograms leftwards as modulation contrast increased. The shifts in the histograms is reflected in the mean dominance durations of the model (Figure 6b), just as with human observers. Similarly, response consistency (Figure 6c, d) increased

when stimulus noise contrast reached 4% and was highest for each contrast at a temporal frequency of 1/8Hz. While human response consistency was quite bandpass (peaking at 1/8Hz and dropping quickly for faster frequencies), the model exhibited slightly broader tuning at high stimulus noise contrast. This may be due to the other parameters of the model that were fixed prior to our simulations, or it could imply additional physiological constraints such as bandpass temporal filters on the input, or response lag.

Figure 6: Summary of modelling results. (a) Histograms of dominance durations of the model with pink (α = 1) internal noise of 16% contrast for each stimulus temporal frequency (i-v) and contrast SD. The solid line colour serves as a legend for the stimulus noise temporal frequency (red = 1/16Hz, green = 1/8Hz, blue = 1/4Hz, yellow = 1/2Hz, purple = 1Hz). The histogram marked in grey represents baseline dominance durations with no contrast modulation. (b) The mean dominance durations of the histograms in (a). Marker colour represents the modulation temporal frequency, while the x-axis gives the modulation contrast. The grey line marks the baseline dominance duration of the model (3.18s), slightly slower than that of the human data. (c-d) Model response consistency plotted in the same manner as Figure 4. In (c), marker colour indicates the modulation temporal frequency while the x-axis indicates the modulation contrast. For all stimulus frequencies, response consistency increased according to modulation contrast, greatest when the stimulus temporal frequency was 1/8Hz. (d) Identical data but plotted with modulation temporal frequency as the x-axis. The grey line (c,d) marks response consistency at baseline with no external noise fed to the model (0.49).

The model predicts consistency with antiphase noise sequences

We next explored whether the model could predict performance in novel conditions. Inspired by Kim et al. (1), we designed a further condition in which the noise modulations were in antiphase across the eyes (i.e. a contrast increase in one eye matched with an equal contrast decrease in the other). We chose a temporal frequency of 1/8Hz, and tested four of our original participants. With no free parameters, the model described above made a strong prediction about performance in this condition (see Figure 7a), namely that double pass consistency should be reliably increased for the antiphase noise (brown curve in Figure 7a), compared to the equivalent condition from the main experiment with two independent streams of external noise (green curve in Figure 7a). This prediction was borne out empirically, as shown in Figure 7b. We note that dominance duration histograms from our human participants (and therefore mean dominance durations) remained relatively unaffected by

this manipulation (see Figure 7c), consistent with performance with independent noise streams (Figure 3aii).

Figure 7: Summary of further conditions testing antiphase modulation and monocular rivalry. Panel (a) shows response consistency predictions of the model for the independent (green) and antiphase (brown) external noise (modulation temporal frequency = 1/8hz). Panel (b) shows the human response consistency for the same conditions as (a). Panel (c) shows histograms of human dominance durations in the same format as Figure 3a, with the unmodulated rivalry condition shown at the top in grey. Panel (d) shows the response consistency of the model when the oscillatory mechanism is removed and modulations are driven by internal and external noise only (cyan) versus the response consistency for the main model (green). Panel (e) shows human response consistency to the 'monocular rivalry' condition (cyan) compared with that of the main experiment (green). Panel (f) shows human dominance duration histograms for the 'monocular rivalry' condition. Error bars and dotted lines show ±1SE across participants (N=4; for the conditions from the main experiment, we omitted results from the participant who did not complete the additional conditions when constructing this figure).

We also tested a condition in which we presented both stimuli to both eyes as a plaid, and modulated the contrast of the components. Just as in the main experiment, we asked participants to report which component appeared higher in contrast at each moment in time, though there was no binocular rivalry. This 'monocular rivalry' condition also produced greater consistency scores than the equivalent condition from the main experiment (see Figure 7e), and demonstrates that the technique can be used to dynamically monitor perception even in the absence of interocular competition. The distributions of dominance durations were rather broader for low contrast modulations, but narrowed at higher contrasts (see Figure 7f).

We reasoned that one way to model this condition might be to remove the rivalry mechanism from the model, leaving only the combination of internal and external noise to determine

dominance at each moment. The predictions for this arrangement are shown by the cyan symbols in Figure 7d, and involve markedly *lower* consistency scores than both the model and empirical binocular rivalry conditions (green curves in Figures 7d,e), and also the monocular rivalry data itself (cyan data in Figure 7e). Clearly then, monocular rivalry does still involve an alternation process, but the increased empirical consistency scores in this condition suggest that the alternating mechanism is more strongly driven by the external noise modulations than during binocular rivalry.

Discussion

Using a combination of psychophysical experiments and computational modelling, we infer that the source of internal noise relevant to perceptual alternations during binocular rivalry has an amplitude spectrum of 1/*f*, and a standard deviation of around 16%. Our method facilitates these inferences because it uses a double pass design, in which an external noise sequence is repeated twice, under the assumption that internal noise will be different on each pass. Although the double pass method has been used previously for briefly presented stimuli (13, 15), this is the first time (to our knowledge) it has been used in a dynamic paradigm. We now discuss details of the rivalry model, relevance to other work on noise in binocular vision, and broader implications for our understanding of internal noise in the brain.

Model variants and alternative models of rivalry

In the course of developing the model, we also considered several variants using same architecture that were either less successful or less plausible. One variant was a model in which a single source of internal noise was added to both channels. In this arrangement, the internal noise was less effective, because it increased or decreased the response in both channels by the same amount, and so did not materially influence the competition between channels. Another variant placed the internal noise sources outside of the gain control equation (i.e. added after eqn 1 rather than appearing on the numerator and denominator). Although moving internal noise later is consistent with the assumptions of a family of popular computational models of early binocular vision (16, 17 see next section), this was less successful than our main model because internal noise levels sufficient to influence consistency had too large an effect on dominance durations. This rendered the dynamic properties of the model moot, with rivalry percepts being largely determined by the internal noise streams.

We also tested alternative values of the main parameters in the rivalry model. These altered model behaviour in the unmodulated baseline condition much as described in previous work (4), but had relatively minimal effects on dominance durations and consistency scores with strongly noise-modulated stimuli, where rivalry alternations depend more on the interplay of internal and external noise than on adaptation and inhibition. We anticipate that other rivalry models with architectures related to that of Wilson (3, 4) could be modified in a similar way as described here to achieve comparable effects, but have not tested this assumption.

Related work on rivalry

As mentioned above, Kim et al. (1) modulated the contrast of rivalling stimuli periodically in
antiphase at a range of temporal frequencies. They implement three computational models
to account for their results, each of which has random walk (i.e. brown) noise with a spectral
slope of $1/f^2$, but report obtaining similar results with white noise for their experimental
conditions. Furthermore, one of the models they implement is a version of the Wilson (3)
model considered here, but they report the best performance when the internal noise is
added to the adaptation differential equation (see Methods), rather than the rivalling units.
In our simulations, we found similar effects on the dominance duration distributions for
internal noise placed either in the main equation or adaptation equation (not shown here).
However, placing internal noise in the adaptation differential equation resulted in response
consistency that was not tuned to modulation frequency (i.e., flat). We suspect that Kim et
al.'s paradigm did not afford sufficient constraints to distinguish between the two very
different internal noise types or the locus of internal noise.
378
Other models that have incorporated a stochastic component include the model of Lehky (2)
which also used random walk (brown) noise, Kalarickal and Marshall (18) who used additive
uniformly distributed (effectively white) noise, and Stollenwerk and Bode (19) who used
temporally white noise that was correlated across space. However none of these studies
report testing other types of internal noise, nor were their experimental conditions sufficient
to offer meaningful constraints on the internal noise properties. As far as we are aware, this
is the first study that has compared internal noise of different amplitudes and spectral
properties with strong predictions from empirical results.
387
Baker & Graf (8) explored binocular rivalry using broadband pink noise stimuli that also varied
dynamically in time. By testing factorial combinations of temporal amplitude spectra across
the two eyes, they showed that stimuli with $1/f$ temporal amplitude spectra tended to
dominate over stimuli with different spectral slopes (the same was also true of static stimuli
with a $1/f$ spatial amplitude spectrum). Whilst these results do not directly imply anything
about the properties of internal noise, they are consistent with the idea that the visual system
is optimised for stimuli encountered in the natural world, which are typically $1/f$ in both space
and time (e.g. 20–24). Our findings here imply that as well as having a preference for external
stimuli with naturalistic properties, the internal structure of the visual system might itself
have evolved to emulate these temporal constraints (21, 25–27).
398
399 *Internal noise in binocular vision and throughout the brain*
400
Early models of binocular signal combination attributed the $\sqrt{2}$ improvement in contrast
sensitivity for fusible stimuli viewed binocularly vs monocularly to the pooling of independent
monocular noise sources (28). However this model assumes that during monocular
presentation, the noise in the unstimulated eye can be ignored, which is implausible in a well-
designed experiment (16). Contemporary binocular models of contrast detection and
discrimination assume noise that is late and additive, occurring at a point beyond binocular
signal combination (17). It is generally assumed that this late source of noise is the
combination of multiple noise generators at successive stages of processing, though relatively
little is known about their precise characteristics. However a small number of studies have
investigated this issue, as we now summarise.
411

Pardhan & Rose (29) added binocular external noise during a monocular or binocular
detection task and found that binocular summation decreased at high levels of external noise,
and that equivalent input noise (the minimum external noise level required to influence
thresholds) was higher for monocular than binocular targets. One interpretation of these
results is that the effective internal noise is greater for monocularly presented stimuli (see
also 30). However, the type of external noise that they used was broadband white pixel noise,
which can also cause substantial gain control suppression (see 15), potentially confounding
the effects of increased variance. These results are therefore relatively inconclusive regarding
sources of internal noise in binocular vision.
421

Recently, Ding & Levi (31) have demonstrated that the inclusion of early (monocular)
multiplicative noise in gain control models can account for some subtle features of binocular
contrast discrimination performance. It has also been suggested that monocular noise might
be increased in the affected eye of individuals with amblyopia (32). Finally, we have recently
shown (33) using a contrast discrimination paradigm that EEG and MEG data are consistent
with both an early (~100ms post stimulus onset) noise source in low level visual areas, and a
later noise source in more frontal and parietal brain areas, both of which affect perceptual
decisions. All of these results are therefore consistent with an early monocular source of
internal noise, as included in our model, but do not preclude the addition of later sources of
noise which we do not consider here.
432

Regarding noise more generally, surprisingly few studies have addressed the spectral and
distribution properties of internal noise using psychophysical methods. The default
assumption is typically that internal noise is Gaussian (owing to Central Limit Theorem) and
white. However, Neri (34) concluded that internal noise had a Laplacian distribution, and
other psychophysical work has assumed Poisson processes for internal noise (35), based on
single cell recordings (36). Noise with a pink amplitude spectrum typically retains a Gaussian
distribution, though in principle non-Gaussian distributions (such as Laplacian or Poisson
distributions) could also be altered to have a pink spectrum. Although we are unaware of any
other psychophysical studies attempting to estimate the spectral characteristics of internal
noise, we note that measurements of spontaneous neural activity using ECoG and fMRI also
have fractal properties, and a slope of approximately $1/f$ in visual areas (37).
444

445   *Conclusions*
446

Using a novel dynamic double pass paradigm with binocular rivalry, we measured how
alternation rates and response consistency were affected by different types and amounts of
external noise. The results were consistent with a computational model of rivalry in which
internal noise was independent in each monocular channel. We conclude that internal noise
relevant to rivalry has an amplitude spectrum of $1/f$, and a standard deviation of around 16%.
We anticipate that future studies might use temporally sensitive neuroimaging techniques
such as EEG and MEG to further investigate these sources of internal noise.
454

455   **Methods**
456

457   *Participants*
458

459 The main experiment was completed by five psychophysically experienced observers, who
460 provided written informed consent. A control experiment was completed by four of the same
461 observers. All observers had no known abnormalities of binocular vision, and wore their
462 standard optical correction if required. Procedures were approved by the Ethics Committee
463 of the Department of Psychology at the University of York.
464
465 *Apparatus and stimuli*
466
Stimuli were sinusoidal grating patches with a spatial frequency of 1c/deg, subtending two
degrees of visual angle, and ramped in contrast by a cosine function over a further ¼ degree.
The gratings shown to the left and right eyes had orthogonal orientations (±45 degrees) which
were assigned randomly on each trial (see Figure 1a for examples). The mean Michelson
contrast of the gratings was 50%, but this was modulated by dynamic noise streams of various
centre frequencies (1/16 Hz to 1Hz) and standard deviations (1% to 16% Michelson contrast).
The noise streams were constructed by bandpass filtering white noise at the required
frequency using a one octave bandpass filter (see Figure 1b). In the main experiment, the
noise streams used to modulate the contrast of each eye were independent.
476
Stimuli were displayed on a ViewPixx 3D display (VPixx Ltd., Canada), driven by an Apple
Macintosh computer. The monitor operated with 16 bits of greyscale luminance resolution
(M16 mode) and was gamma corrected using a Minolta LS110 photometer. Independent
stimulation of the left and right eyes was achieved using stereo shutter glasses (NVidia 3D
Vision), synchronised with the monitor refresh rate of 120Hz via an infra-red signal. To
promote good vergence and binocular alignment, each stimulus was surrounded by a static
high contrast greyscale Voronoi texture (squares of 14 x 14 degrees, with a 7 degree diameter
disc in the centre set to mean luminance) that was identical in both eyes (see Figure 1a). A
different texture was presented on each trial, selected at random from a set of 99 pre-
generated textures.
487
488 *Procedure*
489
Participants sat in a darkened room and viewed the display from a distance of 57cm. Stimuli
were presented for 60 seconds per trial, with condition order determined at random.
Participants were instructed to indicate using a two-button mouse which of the two grating
stimuli they perceived at each moment in time by holding down one or other button. If both
stimuli were perceived, they were instructed to choose the stimulus that was most visible (i.e.
that took up the largest part of the image), or to hold down both buttons if they were equally
salient. At the end of each trial, there was a minimum blank interval of three seconds, with
the following trial initiated by the participant.
498
Each of the 26 conditions (5 contrasts * 5 temporal frequencies + 1 baseline) was repeated 5
times by each observer using unique noise sequences in each repetition, and then a further 5
times using the same noise sequences as in the first pass. This resulted in 260 trials (4.3 hours
of rivalry data) per participant, which were completed across multiple sessions (each typically
lasting 20-30 minutes) over several days. Raw data are available online at: http://dx.doi.org/
10.6084/m9.figshare.7262201
505

Modelling

There are multiple models that have been successful at capturing the oscillatory behaviour of dominant percepts in binocular rivalry (1–4, 38). While they vary in complexity, all include two key characteristics: inhibition between units responding to the left and right monocular stimuli, and self-adaptation. These guarantee that only one unit will be active at a given moment, and that over time, the active unit will decrease its firing rate sufficiently to allow the suppressed unit to be released from inhibition. Apart from a few exceptions (1, 2, 18, 19, 39), most computational investigations of binocular rivalry have focused on deterministic implementations of their models to investigate how suppression and self-adaptation contribute to oscillations in perceptual dominance. It is, however, fairly straightforward to adapt these models of rivalry to include an additive noise term and directly probe the properties (i.e., amplitude and spectral qualities) of internal noise. Here, we probe the properties of internal noise with the minimum rivalry model of Wilson (3, 4).

The minimum rivalry model defines the response of a single unit by two differential equations (equation 1 and equation 2), which incorporate stimulus excitation (*L/R*), self-excitation ($\varepsilon = 0.2$), competitive inhibition ($\omega = 3.5$), self-adaptation (*H*), and here, an additive internal noise term (N). For the unit responding to stimuli presented to the left eye (E_L), the response term is

$$\tau \frac{dE_L}{dt} = -E_L + \frac{M\left[L - \omega E_R + \epsilon E_L + gH_L + N_L\right]_+}{1 + \left[L - \omega E_R + \epsilon E_L + gH_L + N_L\right]_+^{0.8}} \quad (1)$$

and self-adaptation is

$$\tau_h \frac{dH_L}{dt} = -H_L + E_L \quad (2)$$

which is identical for activity in the right eye (E_R), but with the subscripts switched. The constants *M* and *g* serve to scale the response gain and adaptation strength and were set to values of 1.0 and 3.0, respectively. The excitatory (τ) and hyperpolarizing (τ_h) time constants of equation 1 and equation 2 were set to 15ms and 4000ms respectively. All model parameters were fixed in our simulations. Internal noise was additive and independently generated for each eye. As previous studies have already investigated the locus of internal noise with this particular model (1), we chose here to only conduct model simulations with noise added to the unit response equation (equation 1). Note that as the stimulus input to the model is identical to that of the psychophysical experiment (see Figure 2a), we use a contrast gain control variant of the Minimum rivalry model (4) to account for any differences in contrast between eyes. This also means that the noise term is added to both the numerator and denominator of equation 1.

We probed the spectral characteristics of internal noise by injecting the model with
broadband noise patterns ($1/f^\alpha$) generated at one of five different spectral slopes[*], where α =
[0, 0.5, 1.0, 1.5, 2.0] (see Figure 2b). Noise patterns were generated in the Fourier domain by
first creating a flat (α = 0) amplitude spectrum and then multiplying the amplitude coefficient
at each frequency by f^α. The phase of each frequency component was assigned a random
value between $-\pi$ and π. Two different phase spectra were generated in order to create two
independent noise streams (N_L and N_R) with the same amplitude spectrum. These were
rendered in the temporal domain by taking the inverse Fourier transform and adding them to
the left and right units separately.
553
Perceptual switches were implemented as a winner-take-all rule: the dominance of a percept
was defined by the magnitude of $E_{L/R}$ at any given moment in time (if $E_L > E_R$, E_L is dominant;
see Figure 2c) Finally, all model simulations were conducted in MATLAB (version R2017a)
using ODE45 to solve the 4 differential equations that define the response of each unit and
their self-adaptation over 60 seconds (i.e. the duration of a trial in the psychophysical
experiment). We simulated binocular rivalry twice – with different internal noise samples but
the same external noise sequences – for each stimulus noise condition in order to calculate
the response consistency of the model. This was repeated 1000 times, and the model outputs
(dominance duration and response consistency) were averaged across repetitions.
563
564
565 **Acknowledgements**
566

---

[*] We also conducted simulations with bandpass filtered internal noise streams with the same frequency as that of the stimulus sequences, in addition to the broadband internal noise simulations. Response consistency was high for all stimulus conditions, which suggests that this type of internal noise is incapable of modulating model responses beyond that of the external noise sequences. As these results do not offer any additional insight to the characteristics of internal noise, we do not show them here.

References

1. Kim Y-J, Grabowecky M, Suzuki S (2006) Stochastic resonance in binocular rivalry. *Vision Res* 46(3):392–406.

2. Lehky SR (1988) An astable multivibrator model of binocular rivalry. *Perception* 17(2):215–228.

3. Wilson HR (2003) Computational evidence for a rivalry hierarchy in vision. *Proc Natl Acad Sci U S A* 100(24):14499–14503.

4. Wilson HR (2007) Minimal physiological conditions for binocular rivalry and rivalry memory. *Vision Res* 47(21):2741–2750.

5. Baker DH, Meese TS (2007) Binocular contrast interactions: dichoptic masking is not a single process. *Vision Res* 47(24):3096–3107.

6. Legge GE (1979) Spatial frequency masking in human vision: binocular interactions. *J Opt Soc Am* 69(6):838–847.

7. Meese TS, Baker DH (2009) Cross-orientation masking is speed invariant between ocular pathways but speed dependent within them. *J Vis* 9(5):2.

8. Baker DH, Graf EW (2009) Natural images dominate in binocular rivalry. *Proc Natl Acad Sci U S A* 106(13):5436–5441.

9. Baker DH, Graf EW (2009) On the relation between dichoptic masking and binocular rivalry. *Vision Res* 49(4):451–459.

10. Stuit SM, Cass J, Paffen CLE, Alais D (2009) Orientation-tuned suppression in binocular rivalry reveals general and specific components of rivalry suppression. *J Vis* 9(11):17.

11. Stuit SM, Paffen CLE, van der Smagt MJ, Verstraten FAJ (2011) Suppressed images selectively affect the dominant percept during binocular rivalry. *J Vis* 11(10):7.

12. Alais D, Cass J, O'Shea RP, Blake R (2010) Visual sensitivity underlying changes in visual consciousness. *Curr Biol CB* 20(15):1362–1367.

13. Burgess AE, Colborne B (1988) Visual signal detection. IV. Observer inconsistency. *J Opt Soc Am A* 5(4):617–627.

14. Green DM (1964) Consistency of auditory detection judgements. *Psychol Rev* 71:392–407.

15. Baker DH, Meese TS (2012) Zero-dimensional noise: the best mask you never saw. *J Vis* 12(10):20.

16. Legge GE (1984) Binocular contrast summation--II. Quadratic summation. *Vision Res* 24(4):385–394.

17. Meese TS, Georgeson MA, Baker DH (2006) Binocular contrast vision at and above
threshold. *J Vis* 6(11):1224–1243.

18. Kalarickal GJ, Marshall JA (2000) Neural model of temporal and stochastic properties
of binocular rivalry. *Neurocomputing* 32–33:843–853.

19. Stollenwerk L, Bode M (2003) Lateral neural model of binocular rivalry. *Neural
Comput* 15(12):2863–2882.

20. Dong DW, Atick JJ (1995) Statistics of natural time-varying images. *Netw Comput
Neural Syst* 6(3):345–358.

21. Field DJ (1987) Relations between the statistics of natural images and the response
properties of cortical cells. *J Opt Soc Am A* 4(12):2379–2394.

22. Geisler WS (2008) Visual perception and the statistical properties of natural scenes.
Annu Rev Psychol 59:167–192.

23. Simoncelli EP, Olshausen BA (2001) Natural image statistics and neural representation.
Annu Rev Neurosci 24:1193–1216.

24. Hansen B, Essock E (2005) Influence of scale and orientation on the visual perception
of natural scenes. *Vis Cogn* 12(6):1199–1234.

25. Haun AM, Peli E (2013) Perceived contrast in complex images. *J Vis* 13(13):3.

26. Schweinhart AM, Shafto P, Essock EA (2017) Distribution of content in recently-
viewed scenes whitens perception. *J Vis* 17(3):8.

27. Schwartz O, Simoncelli EP (2001) Natural signal statistics and sensory gain control. *Nat
Neurosci* 4(8):819–825.

28. Campbell FW, Green DG (1965) Monocular versus binocular visual acuity. *Nature*
208(5006):191–192.

29. Pardhan S, Rose D (1999) Binocular and monocular detection of Gabor patches in
binocular two-dimensional noise. *Perception* 28(2):203–215.

30. Anderson PA, Movshon JA (1989) Binocular combination of contrast signals. *Vision
Res* 29(9):1115–1132.

31. Ding J, Levi DM (2016) Binocular contrast discrimination needs monocular
multiplicative noise. *J Vis* 16(5):12.

32. Baker DH, Meese TS, Hess RF (2008) Contrast masking in strabismic amblyopia:
attenuation, noise, interocular suppression and binocular summation. *Vision Res*
48(15):1625–1640.

33. Vilidaite G, Marsh E, Baker D (2018) Internal noise in contrast discrimination
propagates forwards from early visual cortex. doi:10.1101/364612.

34. Neri P (2013) The statistical distribution of noisy transmission in human sensors. *J Neural Eng* 10(1):016014.

35. May KA, Solomon JA (2015) Connecting psychophysical performance to neuronal response properties I: Discrimination of suprathreshold stimuli. *J Vis* 15(6):8.

36. Goris RLT, Movshon JA, Simoncelli EP (2014) Partitioning neuronal variability. *Nat Neurosci* 17(6):858–865.

37. He BJ, Zempel JM, Snyder AZ, Raichle ME (2010) The Temporal Structures and Functional Significance of Scale-free Brain Activity. *Neuron* 66(3):353–369.

38. Laing CR, Chow CC (2002) A spiking neuron model for binocular rivalry. *J Comput Neurosci* 12(1):39–53.

39. Brascamp JW, van Ee R, Noest AJ, Jacobs RHAH, van den Berg AV (2006) The time course of binocular rivalry reveals a fundamental role of noise. *J Vis* 6(11):1244–1256.

655
656
657

660
661   Figure S1. (a) Model dominance duration histograms for each of the five noise αs and stimulus condition as in
662   Figure 6a. The solid line colour indicates the stimulus temporal frequency while the fill colour marks the noise
663   α. The grey vertical line marks the mean dominance duration of the 0% modulation contrast condition. For very
664   steep slopes (α = 2) the mean exceeds the x axis limit (~10s). (b) The average dominance duration for each
665   model noise α as in Figure 6b. Note the different scale for the y axis with internal noise αs of 1.5 and 2.0.

666
667 Figure S2. Response consistency for all five internal noise α values investigated here. The left column charts
668 response consistency for each modulation contrast while the right column shows the same data replotted
669 according to modulation frequency as in Figure 6c and 6d.