Dynamically Evolving Cell Sizes During Early Development Enable Normal Gastrulation Movements In Zebrafish Embryos

Triveni Menon[1], Asfa Sabrin Borbora[1], Rahul Kumar[1] and Sreelaja Nair[1*]

[1]Department of Biological Sciences, Tata Institute of Fundamental Research, Homi Bhabha Road, Colaba, Mumbai, 400005, India.

[*]Corresponding author: Sreelaja Nair, email: s.nair@tifr.res.in

Short title: Cell migration and cell size heterogeneity in embryos

The authors declare that they have no conflict of interest.

ABSTRACT

Current knowledge of the mechanisms of cell migration is based on differentiated cells in culture where it is known that the actomyosin machinery drives migration via dynamic interactions with the extracellular matrix and adhesion complexes. However, unlike differentiated cells, cells in early metazoan embryos must also dynamically change cell sizes as they migrate. The relevance of cell size to cell migration and embryonic development is not known. Here we investigate this phenomena in zebrafish embryos, a model system in which reductive cell divisions causes cell sizes to decrease naturally over time as cells migrate collectively to sculpt the embryonic body plan. We show that cell size reduction during early development follows power-law scaling. Because mutations that can perturb cell sizes so early in development do not exist, we generate haploid and tetraploid zebrafish embryos and show that cell sizes in such embryos are smaller and larger than the diploid norm, respectively. Cells in embryos made of smaller or larger than normal cells migrate sub-optimally, leading to gastrulation defects. Multiple lines of evidence suggest that the observed defects originate from altered cell size rather than from pleotropic effects of altered ploidy. This interpretation is strengthened by the result wherein restoring cell sizes to normal diploid-like values rescues gastrulation defects. Live imaging of chimeric embryos where haploid/tetraploid cells are introduced into diploid embryos reveal the cell-autonomous nature of the migration defects. Additionally, aberrant intracellular actin dynamics with respect to the vectorial direction of motion suggests a cellular mechanism behind the migration defects. Taken together, early reductive cell divisions potentially allow dynamic, stage-specific cell size norms to emerge, which enables efficient collective cell migration to correctly position cells in space and time to shape an amorphous ball of blastoderm into an embryo.

Keywords: Zebrafish, cell size, gastrulation, cell migration

INTRODUCTION

Cell migration is a complex interplay between actomyosin contractility in migrating

cells, the molecular nature of the communication between cells and their environments,

the physical properties of the extracellular environment and physical contacts between

neighboring cells (Lee & Losert, 2019, Martin & Lewis, 1992, Norden & Lecaudey, 2019,

Petrie, Gavara et al., 2012, Schaks, Giannone et al., 2019, Trepat, Wasserman et al.,

2009). Cell migration is also an integral feature of both natural and disease conditions,

such as embryonic development, cancer metastasis and wound healing. Though the

molecular and cellular mechanisms that drive cell migration are known, how sizes of cells

within a migrating collective would influence migration is not known. This is because the

molecular and cellular basis of cell migration has been largely gleaned from cells in culture

or from differentiated cells later in embryonic development (Lee & Losert, 2019, Norden &

Lecaudey, 2019, Petrie et al., 2012, Schaks et al., 2019, Trepat et al., 2009).

Cell size is a fundamental feature of biological systems, which varies across

organisms and across cell types within a multicellular organism (Arendt, 2007, Levy &

Heald, 2015). In unicellular eukaryotes such as yeast, optimal cell size is an integrated

property that emerges as a balance between cell growth and cell division (Umen, 2005). In

metazoans, cell size must be viewed minimally in two diverse contexts: adult homeostasis

and embryonic development. During homeostasis, organs in the adult animal and the

organism as a whole, strives to maintain an optimal organ/tissue size by balancing cell

proliferation with cell growth. This balancing act during homeostasis is heavily influenced

by autonomous and non-autonomous cues of growth factor signaling (Artiles, Anastasia et

al., 2009, Conlon & Raff, 1999, Edgar, 2006, Jorgensen, Nishikawa et al., 2002,

Jorgensen & Tyers, 2004, Nurse, 1985, Tumaneng, Russell et al., 2012). The phenomena

of cell growth and division can be coupled to or uncoupled from each other in diverse

developmental and homeostasis contexts (Dong, Feldmann et al., 2007, Qu, Weiss et al.,

2004, Su & O'Farrell, 1998). In general, blocking cell division leads to cell growth, whereas

blocking cell growth elicits a cell division arrest (Jorgensen & Tyers, 2004). An exception

to this general rule is the early developmental phase of all metazoan embryos wherein the

absence of cell growth does not evoke a cell division arrest. On the contrary, early

embryonic development in all metazoans is characterized by a visually striking

phenomenon of progressive reduction in cell size precisely because cell division is

uncoupled from cell growth. This allows newly fertilized single-cell zygotes to generate

increasing numbers of progressively smaller daughter cells to form a multicellular embryo.

Thus, the inherent nature of early cell divisions in metazoan embryos generates an ever-

changing cell size landscape, on a temporal scale of minutes to hours depending on the
species.

Experimental analysis in metazoan embryos reveals that when cell size is
constantly changing during reductive cell divisions, sizes of intracellular macromolecular
assemblies such as the nucleus and the mitotic spindle scales with cell size (Good, Vahey
et al., 2013, Levy & Heald, 2010, Loughlin, Wilbur et al., 2011). Thus, embryos must have
mechanisms to sense cell sizes early in development, even when cell size is a constantly
evolving parameter with each subsequent division. Cell size control has been studied with
respect to whole body size of an organism and, is a complex interplay of cell autonomous
and non-autonomous events, which are organism, organ or context specific (Henery, Bard
et al., 1992, Lloyd, 2013). In general, individual cell sizes in adult animals remain
comparable despite dramatic variations in body size, suggesting that cell numbers, rather
than cell size is modulated to achieve a body size (Conlon & Raff, 1999, Day & Lawrence,
2000). This idea has been experimentally tested in developing mammalian and amphibian
embryos. Increasing or decreasing blastomere numbers in mouse embryos or fusing two
8-cell mouse embryos into one results in embryos of the normal size, an outcome that is
achieved by regulation of cell proliferation during development (Lewis & Rossant, 1982,
Rossant, 1976, Tarkowski, 1959). Similarly, tetraploid amphibian embryos are made of
larger and fewer cells at early blastula stages, but adjust cell numbers to attain normal
larval sizes eventually (Conlon & Raff, 1999, Day & Lawrence, 2000, Gibeaux, Miller et al.,
2018). In these experiments, after the initial manipulations, embryo size was the end-point
readout and the effect on cell size during early development remained unaddressed.
Thus, the fundamental question of the importance of cell sizes in early embryonic
development for subsequent normal embryogenesis has remained open. Interestingly,
despite extensive mutagenesis screens there have been no mutants in animal models in
which early cell sizes during reductive cell divisions are shown to alter. Therefore, it also
remains unclear whether cell size norms exist at all during early reductive divisions.

We investigated the relevance of cell sizes during the reductive cell division phase
on vertebrate embryonic patterning using the rapidly dividing zebrafish embryo as the
experimental paradigm (Kimmel, Ballard et al., 1995). In addition to nutrient and growth
factor signaling, genome size is known to determine cell size (Cavalier-Smith, 2005, Lee,
Davidson et al., 2009, Orr-Weaver, 2015, Otto, 2007). Therefore, in the absence of
mutants, to obtain embryos with cell sizes smaller or larger than the normal diploid
embryos, we generated haploid and tetraploid zebrafish embryos. We additionally also
altered cell sizes in diploid embryos by transient pharmacological treatments. These

independent approaches allowed the experimental generation of zebrafish blastoderms
made of cells that were smaller or larger cells than the developmental stage-specific cell
size norm. Such embryos developed epiboly and gastrulation defects and were
morphologically abnormal. Comparative transcriptome analysis showed that the global
transcriptomes as well as levels of candidate transcripts of the planar cell polarity (PCP)
pathway remained unperturbed. Further, cell size deviations towards smaller and larger
than the norm *both* elicited similar defects in migration and gastrulation indicating that
mechanisms beyond transcriptional alterations potentially drive emergence of the
phenotypes. Importantly, epiboly and gastrulation defects in haploid embryos, which are
composed of cells smaller than the norm, could be rescued by experimentally increasing
cell sizes before initiation of gastrulation. We find that smaller or larger than normal cells
migrate sub-optimally during gastrulation, follow abnormally tortuous migration paths, and
exhibit gross defects in intracellular actin localization as they migrate. Thus, the inability of
sub-optimally sized cells to migrate efficiently may explain the epiboly and gastrulation
defects that emerge when cell sizes in the zebrafish blastoderm deviates from the norm.
Our study suggests that a developmental stage-specific cell size norm exists, and cells
sizes are dynamically set with each round of cell division to produce a cell size landscape
that ensures optimal cell migration leading to normal embryonic development.
158
159
160 **RESULTS**
161
Reductive cell divisions lead to power law reduction of cell sizes in the zebrafish
blastoderm
Observations of embryonic development show that early cell divisions in newly
fertilized zygotes result in daughter cells that are progressive smaller in size. However, no
studies have quantified the actual sizes of cells during this phase in any metazoan
embryo. We therefore began our investigations by characterizing cell sizes in normal
diploid zebrafish embryos (Fig. 1, Supplementary Fig. 1A, D, Supplementary Fig. 2A). We
chose the early blastoderm at 64-cell stage (2 hours post fertilization hpf), late blastoderm
at 1k-cell (3 hpf) and sphere stages (4 hpf), early gastrula at shield stage (6 hpf) and early
larva (24 hpf). Cell sizes decreased progressively during successive stages of embryonic
development (Fig. 1, Supplementary Fig. 1A, D and Supplementary Fig. 2A). We
quantified cell areas from the deep layer of the blastoderm using the BioVoxxel toolbox in
ImageJ (Brocher, 2014) (Fig. 1G-J). The average cell size in a zebrafish embryo

decreased ~200 fold during the first 24 hours of development (Fig. 1K, O). Average cell

areas decreased by ~28 fold during the first 6 hours of development, with a decrease of

~7 fold occurring between 1k-cell and shield stages (Fig. 1K-O). Regression analysis on a

log-log plot revealed a linear reduction of the cell areas between 64-cell to shield stages,

thus suggesting a power law reduction in cell size (Size ~Time^N), where N = -3 is the

calculated slope (Fig. 1P, Q, Supplementary Fig. 4A). To obtain a better understanding of

how cell sizes evolve dynamically during development, we binned the cells as small,

medium and large based on cell areas as follows: at any stage of development, medium

size cells were defined as cells with areas +/- 10% of the average area for that stage of

development. Cells with areas less than this bin were categorized small and those with

areas more than this bin were categorized as large (Supplementary Fig. 3A-D). This size

categorization allowed us to estimate potential changes in proportions of cells within a

specific size category across developmental stages. Interestingly, despite the

heterogeneity in cell sizes, the relative proportions of cells within a particular size category

remained comparable across developmental stages (Fig. 1R).

Cell sizes in haploid and tetraploid zebrafish blastoderms deviate from the normal

developmental stage-specific values

DNA content is known to affect cell sizes in varied contexts (Cavalier-Smith, 2005,

Gregory, 2001, Lee et al., 2009, Otto, 2007). Experimentally generated haploid and

tetraploid zebrafish embryos undergo normal rounds of early reductive divisions (Menon &

Nair, 2018). 1k-cell, sphere and shield stages are reliably identifiable landmark stages

during zebrafish development. These stages encompass developmental time points

before initiation of cell migration (1k-cell), initiation of epiboly movements (sphere stage)

and gastrulation movements (shield stage). Thus, potential cell size deviations from the

norm at these stages could uncover the role of cell size in cell migration in developing

embryos. Comparative analysis of cell sizes in haploids and tetraploids to diploid cell sizes

at 1k-cell, sphere and shield stages revealed that cell sizes in haploid and tetraploid

embryos were smaller and larger, respectively (Fig. 2A-C, Supplementary Fig. 1B, C, E, F,

Supplementary Fig. 2B, C). During the first 6 hours of development, the average cell area

in haploid embryos decreased by ~36 fold and that in tetraploids decreased by ~41 fold,

with a decrease of ~7-8 fold occurring between 1k-cell and shield stages (Fig. 2A-C).

Binning cell size values from haploid and tetraploid embryos according to normal size

values from diploid embryos showed that the small and large category of cells significantly

deviated from the cell size norms across developmental stages (Supplementary Fig. 3A-

D). Analysis of the proportions of cell size categories in haploids and tetraploid embryos
revealed an over-representation of small cells in haploids and large cells in tetraploids,
respectively (Fig. 2D-H). Regression analysis of reduction in cell area using log-log plots
showed that cell sizes continued to decrease linearly suggesting power law reduction in
cell size with development in haploids and tetraploids as well (Supplementary Fig. 4A-I).
215

Haploid and tetraploid embryos develop epiboly and dorsal convergence defects

Reductive cell divisions are followed by collective cell migration during gastrulation,
which rearranges cells in space over time to transform a collection of cells into an embryo.
Collective cell behaviors are influenced by cell sizes in several different contexts,
particularly during migration (Leal-Egana, Letort et al., 2017, Matsiaka, Penington et al.,
2018). Cell size deviations in haploids and tetraploids occurred before gastrulation,
allowing us to study the consequences of deviation from cell size norms on gastrulation. In
diploid embryos, analysis of embryonic development in live embryos after sphere stages
showed that diploid embryos underwent well-characterized collective cell behaviors such
as migration over the yolk ball from the sphere stage (epiboly), followed by involution at
the margin and anterior migration at the shield stage (initiation of gastrulation), and
subsequent dorsal convergence to thicken and elongate the body axis on one side by the
bud stage (end of gastrulation) (Fig. 3A-D). In haploids and tetraploids, initiation and
progression of epiboly, formation of the dorsal shield and anterior migration were delayed
(Fig. 3A-C). By the end of gastrulation, haploid and tetraploid embryos had a
comparatively shorter body axis (Fig. 3D), which was also evident at 24 hpf
(Supplementary Fig. 5A, C, E).

Whole embryo RNA in situ hybridization to check for defects in anterior migration
and dorsal convergence confirmed aberrant cell migration during gastrulation. At 7 hpf, the
prechordal plate mesoderm cells at the anterior express *gsc,* and *foxa2* marks the
posteriorly juxtaposed axial mesoderm (Fig. 3E). In haploids, the domains of *gsc* and
foxa2 expression remained intermingled and the *foxa2* expressing axial mesoderm cells
were found in a shorter domain, indicating a failure in anterior migration of prechordal
plate cells and elongation of axial mesoderm (Fig. 3F). In tetraploids, though there were
no overt defects in *gsc* and *foxa2* expression domains, in general *gsc* expression seemed
dampened (Fig. 3G). After completion of gastrulation at 10 hpf, the extent of dorsal
convergence of cells can be assessed by the distance between the bilateral rows of
ectodermal cells marked by *dlx3b,* across the dorsal midline (Fig. 3H). In both haploids
and tetraploids, the rows of ectoderm cells were located further away from the midline,

indicating a delay in or failure to converge dorsally (Fig. 3I, J). Thus, haploid and tetraploid
embryos began undergoing delayed epiboly, and displayed a range of cell migration
defects during gastrulation (Fig. 3K).

Early transcriptomes of haploids and tetraploids and PCP pathway transcripts remain largely unperturbed

The experimental system used in this study to uncover the relevance of cell size in
normal embryogenesis was alteration of ploidy in zebrafish embryos. This experimental
paradigm was chosen primarily because there are no zebrafish mutants with smaller or
larger cells during 1k-cell to shield stages that would allow us to study the consequence of
altering cell sizes before and during gastrulation on overall embryonic development. DNA
content has been shown to affect timing of zygotic genome activation (ZGA), which occurs
at 2.5 hpf in zebrafish (Kane & Kimmel, 1993, Newport & Kirschner, 1982a, Newport &
Kirschner, 1982b). To prove that gene expression levels in haploids and tetraploids
remained comparable to diploids, we performed whole transcriptome analysis at ~3.5 hpf.
Whole transcriptome comparisons showed that at ~3.5 hpf, majority of genes were not
mis-expressed in response to ploidy alteration (Fig. 4A, B). 21 genes in haploids and 33
genes in tetraploids were mis-expressed (Fig. 4C and Supplementary Table 1).
Interestingly, only 6 genes are mis-regulated in common between haploids and
tetraploids, yet both kinds of embryos manifest similar defects later in development
(discussed in later sections), suggesting that the developmental defects do not arise from
perturbed ploidy *per se.* Dorsal convergence, which is compromised in both haploids and
tetraploids, is most pronounced during late gastrula stages (Kimmel et al., 1995). A
molecular pathway known to drive collective cell migration in diverse cellular and
evolutionary contexts, including dorsal convergence is the Planar Cell Polarity (PCP)
pathway (Davey & Moens, 2017, Heisenberg, Tada et al., 2000, Topczewski, Sepich et
al., 2001). Quantitative real time PCR showed that transcript levels of 11 genes
representing diverse components of the PCP signaling pathway remained unaltered in
haploids and tetraploids at 8 hpf (Fig. 4D).

Drug-induced cell size increase in diploid blastoderm recapitulates gastrulation defects

277 Gene Ontology analysis of the 54 mis-expressed genes in haploids and tetraploids
278 showed that these genes controlled a diverse range of biological functions
279 (Supplementary Table 1). However, an unbiased approach to test whether deviation from

cell size norms during early development could result in gastrulation defects would be to recreate the cell size deviation phenomenon in diploids. A combinatorial treatment of Aphidicolin and Hydroxyurea (Aph+HU) has been used to effectively increase cell sizes in the zebrafish notochord (Liu, Sepich et al., 2017). To preclude the possibility that exposure to Aph+HU at early developmental stages was inducing DNA damage, we assayed for γH2AX foci in the nuclei after Aph+HU treatments. Diploid zebrafish embryos treated with Aph+HU from 1k-cell to shield stages showed negligible toxicity and DNA damage (Supplementary Fig. 6, 7). Average cell areas in Diploid Aph+HU embryos were larger than normal at shield stages, with large cells representing majority of the sample population (Fig. 5A-D, Supplementary Fig. 3E). By bud stages, diploid Aph+HU embryos had shorter body axis, which resembled the phenotypes seen in haploids and tetraploids (Fig. 5E, F, compare to Fig. 3D). Analysis of the extent of dorsal convergence at 12 hpf showed that, the *dlx3b* expressing ectoderm was found further away from the midline, a phenotype that also resembled the *dlx3b* expression in haploids and tetraploids (Fig. 5H, compare to Fig. 3I, J). Thus, in a ploidy-independent experimental paradigm, increasing cell sizes in diploid embryos between 1k-cell to shield stages also resulted in gastrulation defects similar to those seen in haploid and tetraploid embryos.

Drug-induced cell size increase in haploid blastoderms rescues gastrulation defects

If cell size deviations from the norm during 1k-cell to shield stages does indeed cause gastrulation defects, then restoring cell sizes to the norm may rescue the developmental defects. Cell size increase upon Aph+HU treatment allowed us to test this experimentally in haploid embryos, wherein cell sizes are smaller than the norm between 1k-cell and shield stages. Aph+HU treatment of haploid embryos from 1k-cell to shield stages (3 hour exposure) as was done for diploid embryos resulted in cell extrusions during gastrulation indicating that haploid embryos were fragile. A shorter Aph+HU exposure regime from 1k-cell to dome stage (1.5 hour exposure) was therefore used for haploid embryos. Haploid embryos treated with Aph+HU from 1k-cell to dome stages showed an increase in cell sizes to diploid-like values by the shield stage (Fig. 5I-K, Supplementary Fig. 3F). Rescuing the cell areas in haploid Aph+HU to diploid-like values by shield stage also restored the proportions of cells within different size categories to diploid-like proportions (Fig. 5L). Importantly, gastrulation defects were also partially rescued in haploid Aph+HU embryos (Fig. 5M, N). Analysis of dorsal convergence in

314   haploid Aph+HU embryos clearly showed that the ectoderm marked by *dlx3b* was closer

315   to the midline indicating a rescue in dorsal convergence (Fig. 5O, P).

Smaller or larger than normal cells migrate sub-optimally during gastrulation

To assay cell migration in embryos with smaller or larger size cells than the norm,

fluorescently labeled cell clones were generated in diploid, haploid and tetraploid embryos

and migration of individual cells were tracked during gastrulation (Supplementary Movies

1-3). Quantification of cell migration parameters revealed that cells in haploids and

tetraploids covered less distance (total path length travelled) and were also less displaced

(net change in position from beginning to end of imaging) in comparison to cells in diploid

embryos (Fig. 6A, B). The observed deficiencies in distance covered and displacement

could arise because cells move slower and/or took more tortuous paths during migration.

In comparison to cells in diploid embryos, cells in haploids and tetraploids did indeed take

significantly more tortuous paths during migration (Fig. 6C). The average speed of cells in

tetraploids was comparable to cells in diploids, whereas the average speed of cells in

haploids tended to be slightly slower (Fig. 6D).

An advantage of zebrafish as a model organism is the ease of generating chimeric

embryos to assay experimental outcomes when cells are provided developmental milieus

different from their native environments. We next tracked cell migration during gastrulation

in chimeric diploid embryos in which small groups of differentially labeled diploid:haploid

(red, green) or diploid:tetraploid (red, green) cells were co-transplanted. Cells were

transplanted into diploid host embryos at sphere stage, just prior to initiation of epiboly,

and tracked live by timelapse imaging till the end of gastrulation. Though the initial

positions of the normal-small (diploid:haploid) and normal-large (diploid:tetraploid) cells

were the same at sphere stages (Fig. 6E, H), during gastrulation smaller and larger cells

were found farther laterally in the embryo in comparison to co-transplanted normal size

cells (Fig. 6F, I and Supplementary Movies 4, 5). By the end of gastrulation, smaller and

larger cells were still displaced farther laterally and were consistently found further away

from the midline in comparison to the co-transplanted normal cells (Fig. 6G, J and

Supplementary Movies 4, 5). Quantification of cell migration parameters of co-

transplanted normal-small and normal-large cells revealed that small (haploid) and large

(tetraploid) cells traveled less distance (Fig. 6K), were displaced less (Fig. 6L) and took

significantly more tortuous paths (Fig. 6M), in comparison to normal (diploid) cells. The

average speed of large cells was comparable to normal cells whereas the average speed

of small cells was significantly slower than normal size cells (Fig. 6N).

Smaller or larger than normal cells show altered actin dynamics

Smaller or larger than normal cells may migrate aberrantly due to altered actin dynamics during collective cell migration. Therefore, actin dynamics during gastrulation was assayed by generating mosaic diploid embryos in which diploid (normal), haploid (small) or diploid Aph+HU (large) cells were transplanted at sphere stages and imaged live during gastrulation (Supplementary Movies 6-14). The transplanted cells were fluorescently labeled for cell membrane and actin. In normal (diploid) cells, an actin patch formed at the back of the cell, which rearranged into smaller dynamic membrane punctae and then reassembled as a patch at the back of the cell. To better understand the dynamic actin localizations, azimuthal distribution of actin patches within single cells was analyzed with respect to the direction of migration. This analysis revealed that actin localized to 3 of the 4 quadrants in a cell, leaving one quadrant, in the direction of migration, largely free of actin during migration (Fig. 7A-B, Supplementary Movies 6-8). Similar analysis of the positions of actin in migrating small cells (haploid) revealed several larger actin patches in all four quadrants of the cell, which moved randomly without disassembly during migration (Fig. 7C-D, Supplementary Movies 9-11). In large cells (diploid Aph+HU), a very large patch of actin formed at the front of the cell and remained at the front without disassembling during migration in addition to the actin punctae that localized to all four quadrants during migration (Fig. 7E-F, Supplementary Movies 12-14).

DISCUSSION

A quantitative analysis of the evolution of progressive reduction in cell sizes during development revealed that early zebrafish embryos are composed of heterogeneously sized cells, which exist in certain proportions. We do not know how these proportions are set in the developing embryo and the relevance of the proportions of cell size categories to normal development. The quantifications of cell areas from a sample cell population within the deep cell layer also revealed that cell size reduction leading up to gastrulation is linear, following power law behaviour. The cellular and molecular mechanisms that regulate cell size dynamics during early embryonic development are not known in any metazoan. Because alternate methodologies to alter cell sizes during the early reductive cell division phase of development are unavailable we used haploid and tetraploid zebrafish embryos together with pharmacological perturbations to alter cell sizes. Altered ploidy zebrafish embryos are advantageous for this study because unlike mouse haploid and tetraploid embryos, which are early embryonic lethal, zebrafish haploids and tetraploids are late larval lethal (Eakin, Hadjantonakis et al., 2005, Henery et al., 1992,

Kaufman & Gardner, 1974, Koizumi & Fukuta, 1995, Modlinski, 1975). As seen in multiple

cell types in both plants and animals (Henery et al., 1992, Orr-Weaver, 2015), we found

that cell size in zebrafish haploids and tetraploids correlated with the ploidy. Such

deviation in cell size was uniquely advantageous to this study as it allowed assessment of

the consequence of deviating from a cell size norm early in development on subsequent

embryogenesis in an endogenously diploid vertebrate. In haploids and tetraploids absolute

cell sizes as well as proportions of cells across size categories deviated from that in

diploids at comparable developmental stages. Haploid and tetraploid zebrafish embryos

were delayed in development during late blastula and early gastrula stages. Morphological

defects exacerbated with time and culminated in dorsal convergence defects by the end of

gastrulation. This study does not distinguish the possibility of developmental defects due

to cell size deviations versus due to perturbations in the proportions of size categories. We

interpret the developmental defects as arising from a combination of the deviations from

the normal cell size and proportion norm for a developmental stage.

Comparative transcriptomics in haploids and tetraploids just after ZGA revealed

that levels of majority of the transcripts remained comparable to diploids, suggesting no

gross transcriptional perturbations. This is also true with respect to ploidy in yeast

(Galitski, Saldanha et al., 1999). Interestingly, despite the fact that haploids and

tetraploids manifest similar gastrulation defects, mis-regulated transcripts between the two

conditions are largely unique. Therefore, independent transcriptional cascades may go

awry in haploids and tetraploids and yet cause similar aberrant developmental outcomes.

Alternatively, transcription independent mechanisms may regulate the evolution of cell

size dynamics during early development. A complementary, ploidy-independent alteration

of cell sizes in diploid embryos by transient pharmacological treatments support the

interpretation that independent of gene expression changes, deviations in cell size from

the norm during early stages can cause consistent developmental abnormalities. It would

be quite unlikely that alteration in ploidy and pharmacological perturbations both result in

cell size deviation from the norm and gastrulation defects due to general toxic or

pleiotropic effects. Such an interpretation is strengthened by a phenotypic rescue of

haploids wherein gastrulation defects were mitigated when cell sizes were restored to

normal-like sizes at shield stages. It is again highly unlikely that two separate and

independent sets of potential pleiotropic effects (arising from altered ploidy and

pharmacological perturbations) would add up to rescue developmental defects in haploids.

The developmental defects we describe do not resemble those seen when apico-

basal polarity or cell adhesion is perturbed in zebrafish embryos (Kane, McFarland et al.,

2005, Sonawane, Carpio et al., 2005). Recent work in size reduced zebrafish blastoderms

showed that signaling gradients dynamically scale to readjust somite sizes to scale with

smaller embryo sizes (Ishimatsu, Hiscock et al., 2018). In *D. melanogaster*, morphogen

gradients scale in response to cell size change to maintain a normal wing size (Day &

Lawrence, 2000). In early zebrafish embryos, when cell sizes deviate from a

developmental stage-specific norm, perhaps cells perceive signaling gradients in the

blastoderm or early gastrula differently. However, regardless of sensing the deviation from

a cell size norm, embryos are unable to compensate for the suboptimal early cell sizes to

restore normal patterning.

Regulating cell size during early development when cell size is a constantly

changing parameter versus regulating cell size during homeostasis where the goal is to

maintain a certain cell size may require distinct mechanisms. It is possible that the

dynamic nature of cell size change during early development uses an equally dynamic

cytoskeletal interaction between astral microtubules and cortical actin to sense cell

boundaries and adjust cell size dynamically, especially in the rapidly dividing zebrafish

embryo. The spatial analysis of gene expression domains indicated compromised cell

migration when developmental stage-specific cell sizes deviated from diploid-like values.

Tracking cell migration during gastrulation in chimeric and mosaic embryos during

gastrulation revealed that smaller or larger cells migrated sub-optimally in comparison to

normal size cells and was likely to cause aberrant positioning of cells during gastrulation,

resulting in phenotypes such as compromised dorsal convergence. Additionally, actin

reorganization during migration was altered in cells that were smaller or larger than

normal, suggesting that cell size deviation from the norm affects cell migration at least

partially by altering actin dynamics. The mis-localization of actin into all four quadrants in

smaller or larger cells may drive cells to take tortuous paths during migration, making the

whole process of migration suboptimal. Taken together, live tracking of cell migration

during gastrulation in zebrafish indicates that cell migration parameters are cell

autonomous and may depend on the size of the cell.

It is intuitive that during early embryogenesis, a phase characterized by growing

numbers of progressively smaller cells, optimum cell sizes may be important for normal

development. However, in the absence of experimental efforts directed at altering cell

sizes during the earliest phases of development, this hypothesis remains an intuitive

conjecture at best. Reductive cell division after fertilization is an evolutionarily conserved

phenomenon in metazoans. Our study shows that zebrafish embryos made of smaller or

larger than normal cells develop abnormally, likely due to cell autonomous and possibly

size-dependent effects on migratory properties of cells, which affects gastrulation. In developing embryos, early reductive cell divisions potentially allow dynamic, developmental stage-specific cell size norms to emerge. Perhaps such a landscape of optimal cell sizes enables efficient collective cell migration to correctly position cells in space and time to shape the amorphous ball of cells in a blastoderm into a patterned embryo at the end of gastrulation.

MATERIALS AND METHODS

Zebrafish: Adult *Danio rerio* of the standard laboratory strains Tubingen and AB were reared at 28.5°C under 14 hour light and 10 hour dark conditions. To synchronize developmental stages, embryos were obtained by in vitro fertilizations (IVF) for all experiments and raised at 28.5°C. Diploid, gynogenetic haploid and tetraploid embryos were generated as described in (Menon & Nair, 2018, Menon & Nair, 2019).

Validation of ploidy: 24 hpf larvae were dechorionated and arrested at metaphase in 4mg/mL Colchicine (Sigma C9754) at 28^{O}C for 90 min in the dark. The larvae were treated with 1.1% sodium citrate monobasic and yolks were punctured in a span of 8 mins at room temperature (RT) followed by incubation on ice for an additional 8 mins. Deyolked embryos were fixed overnight in 3:1 Methanol:Acetic acid at 4^{O}C, transferred into 50% Acetic acid and minced thoroughly with forceps. A single cell suspension was generated by repeated trituration using 50 μl glass capillary pipettes and 2-3 drops were dropped from a height onto a glass slide pre-warmed to 65^{O}C. The slides were incubated at 65^{O}C for 60 mins and stained with DAPI (Invitrogen P36974) overnight at RT. Metaphase chromosomes were imaged on Zeiss Axio-Imager M2 and counted using Image J.

Developmental Staging of embryos: To stage-match embryos for comparative analysis IVF embryos at 1k-stage (3 hpf), high stage (3.5 hpf), dome stage (4.5 hpf), sphere stage (4 hpf) and shield stage (6 hpf), were staged according to the characteristic morphology of the embryo at these landmark stages. For 11 and 12 hpf fixations, the somite number was used as the criterion for stage matching embryos. 2 and 24 hpf embryos were staged by absolute time after fertilization.

Immunofluorescence and imaging: Embryos were fixed using 4% paraformaldehyde and immunolabelled with rabbit anti-β-catenin (Sigma C2206, 1:1000) or rabbit anti-γH2AX (Cell Signaling Technology 2577S, 1:1500) and donkey anti-rabbit Alexa 488 (Invitrogen A21202, 1:100) (Pelegri, Knaut et al., 1999). DNA was stained using DAPI. Semi-flat mounts of the blastoderms were prepared in 70% glycerol containing DABCO (Sigma D27802) and imaged on an Olympus FV1200. Images were processed and analyzed using ImageJ and figure panels were assembled using Adobe Photoshop. For 1k-cell, sphere and shield stage embryos, all four quadrants were imaged for cell size quantifications to cover majority of the blastoderm.

Cell size quantifications: To quantify the cell area of the deep layer cells, a single z-slice from the middle $2/3^{rd}$ of a β-catenin and DAPI labeled confocal z-stack was chosen from all images. This z-slice was segmented using the macro

http://imagej.1557.x6.nabble.com/Index-of-hexagonality-SOLVED-td5013442.html.

The segmented confocal z-slice was an image in which each cell was outlined and numbered. These outlined and numbered cells were overlaid on the original confocal image to verify the segmentation. Aberrantly segmented cells were eliminated or manually outlined and numbered. To obtain maximal cross-sectional area of a cell, areas were only calculated from cells in which the nucleus was clearly visible. The correctly segmented image was used in the Extended Particle Analyzer and Shape Descriptor Map plugins of the Biovoxxel Toolbox in ImageJ which generated area heat maps and an excel sheet with the cross-sectional area of each numbered cell. 3 to 4 embryos were analyzed for 2 and 24 hpf stages. All four quadrants from 3 embryos each were analyzed for 1k-cell, sphere and shield stage embryos. Area calculations were done using Microsoft Excel. Since the data was not normally distributed, statistical significances were calculated using Mann Whitney and Kruskal Wallis tests in Graphpad Prism. Violin plots were made using Chart Studio at https://plot.ly

Whole mount RNA in situ hybridization: Embryos fixed in 4% paraformaldehyde were processed for whole mount in situ hybridization (Thisse, Thisse et al., 1993). Digoxigenin or Fluorescein labeled anti-sense RNAs for the genes described were detected using anti-digoxigenin (Roche 16646820) and anti-fluorescein antibodies (Roche 17110000), using NBT (Roche 11383213001) and BCIP substrate (Roche 11383221001) or Sigma Fast (Sigma F4648). Stained embryos were post-fixed in 4% PFA and mounted in 100%

glycerol for imaging on Zeiss AxioImager M2. Images were processed and assembled

using ImageJ and Adobe Photoshop.

Transcriptome analysis: 50 pairs of adults of the AB strain were used exclusively to

generate diploid, haploid and tetraploid embryos for transcriptome sequencing. For one

biological replicate, embryos were obtained from ~15 pair matings and diploid, haploid

and tetraploid embryos were generated from the same clutch of embryos as described in

(Menon & Nair, 2018). At high stage, total RNA was extracted from ~120 embryos of each

ploidy by Trizol method (Ambion 15596018). RNA quality was assayed on a Bioanalyzer

(Agilent 2100). The RNA set in which RNA from diploid, haploid and tetraploid embryos

generated in the same experiment had RIN values >/= 8-9 was sent for whole

transcriptome sequencing. The entire experiment of generating all three ploidies and RNA

extraction was repeated two more times on different days to obtain a biological replicate of

three RNA samples for each ploidy. Ploidy was verified by metaphase chromosome

spreads at 24 hpf for each sample that was used for transcriptome sequencing. Strand

specific library was generated and 15-20 million 100 bp paired-end Illumina reads were

obtained on HiSeq 4000. The Illumina reads were processed and analyzed using HISAT2

and Cuffdiff at the Galaxy open source platform (https://usegalaxy.org/) to obtain

differentially expressed transcripts from the three biological replicates of transcriptomes of

each ploidy. R software was used to generate the Volcano plots and heatmap.

Quantitative RT-PCR: Diploid, haploid and tetraploid embryos were generated from the

same IVF clutch as was done for the transcriptome sequencing. Total RNA was extracted

from ~40 diploid, haploid, and tetraploid embryos using Trizol. 1µg of the RNA converted

into cDNA using the SSIII RT First Strand cDNA synthesis kit (Invitrogen 18080051). qRT-

PCR was performed using Kapa Sybr (Kapa Biosystems KK4601) reaction mix on a

Roche   LightCycle   96.   The   q-RTPCR   primers   used   were: *fz2*-F-

GAGTTTGCTTTGTGGGCCTG   and   *fz2*-R-GCAAGCAGAAATG AGGTGCC, *fz3a*-F-A

GAGCGACTGCTACAAGCTC   and   *fz3a*-R-GCTCGAGGGTACGACTCTTC,   *fz6*-F-AAGT

ACCTGATGACGCTGGC   and   *fz6*-R-CGCTGACCGCATCTTTCTTG,   *fz7b*-F-AGGATAG

TTTGTGCCTCGCC   and   *fz7b*-R-TTCCCGTACCGCCATCATTG,   *vangl2*-F-AGCCGCTT

CTACAATGTGGG   and   *vangl2*-R-TGGGAAGGTTGAGTAAAGCAGG,   *celsr1a*-F-TCC

ACGAGAACATCTGAACGG   and   *celsr1a*-R-TCCTGCTGGGTCAGATTGATG,   *celsr1b*-F-

ACCTGGATAACAACAGGCCG   and   *celsr1b*-R-TGTTCTCCAGACGAACGGTG,   *celsr2*-F-

CTTTCATCGAGGGCGGAGTC   and   *celsr2*-R-ACGAGTCACAGTCGTCAACC,   *celsr3*-F-

GCGTCTCCGAAGTGTCACC and *celsr3*-R-GATCGCTCCAGCTGCTTTTG, *dvl1a*-F-ACGATGATGCCAGCAGACTC and *dvl1a*-R-TTGTGGTCTTTCGGTCTCCG, *dvl2*-F-TCCTTACTCCACCCAGCCTC and *dvl2*-R-AACGCTTTTAGGGCCCCTTC, *dvl3a*-F-AGTTGCTTTGCTTTGCCGAG and *dvl3a*-R-GGATGGACTCATGCCGTAGG, *dvl3b*-F-GGGATCCAAACCCTCGTAGC and *dvl3b*-R-GGCTCATACCGTACACTGGG, *tbp*-F-ACCTCCTTTCGCTCAAGGTC and *tbp*-R-CCAGG AGGGACAAGCTGTTG. The fold-change of test transcripts was analyzed with respect to *tbp* transcript levels. The data was analyzed and graphically represented using Microsoft Excel.

Pharmacological treatments: Standardization experiments for Aphidocolin (Sigma A0781) and Hydroxyurea (Sigma H8627) (Aph+HU) treatment of zebrafish embryos during blastoderm stages is shown in Supplementary Fig. 6. Before addition of the drugs, embryos were transferred into 0.3% Danieau medium (17mM NaCl, 0.21mM KCl, 0.12mM MgSO$_4$.7H$_2$O, 0.18mM Ca(NO$_3$)$_2$.4H$_2$O and 1.5mM HEPES; pH 7.6) and dechorionated using Pronase (Sigma P6911). At 1k-cell stage, Danieau medium was replaced with ~10 ml of 50 µM Aphidicolin and 5 mM Hydroxyurea solution prepared in 4% DMSO. This concentration of Aph+HU was used for all experiments after standardization. Control embryos were treated with only 4% DMSO. The drug solution or DMSO was discarded at shield stage for diploids or at 4.5 hpf for haploids and for diploid embryos used as donors in the actin timelapses. Embryos were used as donors for timelapse imaging, fixed at shield stages or allowed to develop at 28.5^{O}C till live imaging at end of gastrulation and fixation at somite stages. Live images at shield stages were recorded using Zeiss AxioCam MRm camera; 10 hpf images were recorded using Zeiss AxioCam ICc1.

Analysis of DNA damage in drug treated embryos: Aph+HU treated embryos were fixed at shield stages using 4% PFA. For positive control of DNA damage, dechorionated diploid embryos were irradiated with 254nm UV for 10 min at 1k-cell stage and fixed for DNA damage analysis at shield stage. DNA damage was assessed using γH2AX immunolabeling as described above.

Differential cell labeling for live cell migration assays: For labeling small clones of cells in whole embryos, 0.5 nL of 1% Tetramethylrhodamine Dextran (Invitrogen D1817), was injected into one cell at the 16-32 cell stage in diploid, haploid or tetraploid embryos.

For co-transplanting cells to generate mosaic diploid embryos, donor diploid embryos were injected with 0.5 nL of 1% Tetramethylrhodamine Dextran and donor

haploid/tetraploid embryos were injected with 0.5 nL of 1% Fluorescein Dextran (Invitrogen D1820) at the one-cell stage. At sphere stage, a few labeled cells from diploid and haploid/tetraploid donors were co-transplanted into an unlabeled diploid host embryo also at sphere stage. The initial position of the transplanted donor cells in the host embryo was recorded on a Zeiss AxioImager M2 epifluorescence microscope.

For assaying actin localization during migration, donor diploid and haploid embryos were injected at the one-cell stage with CAAX-mCherry mRNA (Addgene) to visualize membrane protrusions (Hancock, Cadwallader et al., 1991) and Alexa488-Phalloidin (0.7x in DMSO, Thermo Fisher A12379) to label actin during gastrulation. In our experiments and as has been previously reported, the concentration of Phalloidin injected did not cause developmental abnormalities (Li, Webb et al., 2008). To obtain embryos with larger cell sizes, injected diploid embryos were treated with Aph+HU as described above. At sphere stages, a few cells were transplanted from donor diploid, haploid or diploid Aph+HU embryos into unlabeled diploid host embryos.

Timelapse imaging during gastrulation:

For all timelapse experiments, embryos were screened at the shield stage for the position of labeled/transplanted cells relative to the dorsal shield. An embryo in which the dorsal shield was laterally in focus relative to the position of the labeled/transplanted cells was mounted in 0.5% low melting agarose for live timelapse imaging on Zeiss LSM880 using a 10x objective. Timelapse was recorded from shield stages to end of gastrulation and assembled on ImageJ. Cell migration was tracked using the Manual Tracking plugin in ImageJ. Tracking data was collected from 35 cells each from four diploid, haploid and tetraploid clonal timelapses and 80 cells each from four diploid-haploid and diploid-tetraploid mosaic timelapses. Statistical significance was calculated using the Mann Whitney test in GraphPad Prism 5.0.

Timelapses of actin were also recorded similarly using a 40x objective to visualize cell membrane and actin punctae during migration. From one host embryo, during shield stages to end of gastrulation, four separate clusters of cells were imaged for ~30 mins each. Two embryos were imaged for each condition and the timelapses were assembled in ImageJ. To record the positions of actin punctae in a migrating cell, cells were viewed as circles partitioned into four quadrants and the angles subtended by the actin punctae from the center of the circle were measured in ImageJ for the duration of the timelapse. The angle values were used to make the actin position maps in Adobe Illustrator with respect to the direction of migration.

Acknowledgements: This work was supported by an India Alliance Wellcome Trust/Department of Biotechnology Intermediate Fellowship IA/I/13/2/501042 to SN, and by the Tata Institute of Fundamental Research, Mumbai. We wish to thank Akshina Mehta for help with RNA preparations for the transcriptome analysis. We thank SN lab members for inputs.

Author Contributions:

TM performed all experiments, compiled and analysed data. SN designed the study, performed experiments, data analysis and compilation with TM. TM, ASB and SN performed cell migration quantifications and analysis. TM and RK performed the qRT-PCR experiments. SN wrote the manuscript with comments from TM.

Animal use ethics statement: All animal husbandry and experiments were conducted in accordance with national and institutional animal use guidelines (approval number TIFR/IAEC/2015-3).

Data availability statement: RNA sequencing data sets generated will be deposited with GenBank upon publication and will be freely available for use.

REFERENCES

1. Arendt J (2007) Ecological correlates of body size in relation to cell size and cell number: patterns in flies, fish, fruits and foliage. Biol Rev Camb Philos Soc 82: 241-56

2. Artiles K, Anastasia S, McCusker D, Kellogg DR (2009) The Rts1 regulatory subunit of protein phosphatase 2A is required for control of G1 cyclin transcription and nutrient modulation of cell size. PLoS Genet 5: e1000727

3. Brocher J (2014) Qualitative and Quantitative Evaluation of Two New Histogram Limiting Binarization Algorithms IJIP 8 30-48

4. Cavalier-Smith T (2005) Economy, speed and size matter: evolutionary forces driving nuclear genome miniaturization and expansion. Ann Bot 95: 147-75

5. Conlon I, Raff M (1999) Size control in animal development. Cell 96: 235-44

6. Davey CF, Moens CB (2017) Planar cell polarity in moving cells: think globally, act locally. Development 144: 187-200

7. Day SJ, Lawrence PA (2000) Measuring dimensions: the regulation of size and shape.
Development 127: 2977-87

8. Dong J, Feldmann G, Huang J, Wu S, Zhang N, Comerford SA, Gayyed MF, Anders
RA, Maitra A, Pan D (2007) Elucidation of a universal size-control mechanism in
Drosophila and mammals. Cell 130: 1120-33

9. Eakin GS, Hadjantonakis AK, Papaioannou VE, Behringer RR (2005) Developmental
potential and behavior of tetraploid cells in the mouse embryo. Dev Biol 288: 150-9

10. Edgar BA (2006) How flies get their size: genetics meets physiology. Nat Rev Genet 7:
907-16

11. Galitski T, Saldanha AJ, Styles CA, Lander ES, Fink GR (1999) Ploidy regulation of
gene expression. Science 285: 251-4

12. Gibeaux R, Miller K, Acker R, Kwon T, Heald R (2018) Xenopus Hybrids Provide
Insight Into Cell and Organism Size Control. Front Physiol 9: 1758

13. Good MC, Vahey MD, Skandarajah A, Fletcher DA, Heald R (2013) Cytoplasmic
volume modulates spindle size during embryogenesis. Science 342: 856-60

14. Gregory TR (2001) Coincidence, coevolution, or causation? DNA content, cell size,
and the C-value enigma. Biol Rev Camb Philos Soc 76: 65-101

15. Hancock JF, Cadwallader K, Paterson H, Marshall CJ (1991) A CAAX or a CAAL motif
and a second signal are sufficient for plasma membrane targeting of ras proteins.
EMBO J 10: 4033-9

16. Heisenberg CP, Tada M, Rauch GJ, Saude L, Concha ML, Geisler R, Stemple DL,
Smith JC, Wilson SW (2000) Silberblick/Wnt11 mediates convergent extension
movements during zebrafish gastrulation. Nature 405: 76-81

17. Henery CC, Bard JB, Kaufman MH (1992) Tetraploidy in mice, embryonic cell number,
and the grain of the developmental map. Dev Biol 152: 233-41

18. Ishimatsu K, Hiscock TW, Collins ZM, Sari DWK, Lischer K, Richmond DL, Bessho Y,
Matsui T, Megason SG (2018) Size-reduced embryos reveal a gradient scaling-based
mechanism for zebrafish somite formation. Development 145

19. Jorgensen P, Nishikawa JL, Breitkreutz BJ, Tyers M (2002) Systematic identification of
pathways that couple cell growth and division in yeast. Science 297: 395-400

20. Jorgensen P, Tyers M (2004) How cells coordinate growth and division. Curr Biol 14:
R1014-27

21. Kane DA, Kimmel CB (1993) The zebrafish midblastula transition. Development 119:
447-56

22. Kane DA, McFarland KN, Warga RM (2005) Mutations in half baked/E-cadherin block
cell behaviors that are necessary for teleost epiboly. Development 132: 1105-16

23. Kaufman MH, Gardner RL (1974) Diploid and haploid mouse parthenogenetic
development following in vitro activation and embryo transfer. J Embryol Exp Morphol
31: 635-42

24. Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF (1995) Stages of
embryonic development of the zebrafish. Dev Dyn 203: 253-310

25. Koizumi N, Fukuta K (1995) Preimplantation development of tetraploid mouse embryo
produced by cytochalasin B. Experimental animals 44: 105-9

26. Leal-Egana A, Letort G, Martiel JL, Christ A, Vignaud T, Roelants C, Filhol O, Thery M
(2017) The size-speed-force relationship governs migratory cell response to
tumorigenic factors. Mol Biol Cell 28: 1612-1621

27. Lee HO, Davidson JM, Duronio RJ (2009) Endoreplication: polyploidy with purpose.
Genes Dev 23: 2461-77

28. Lee RM, Losert W (2019) Dynamics phenotyping across length and time scales in
collective cell migration. Semin Cell Dev Biol 93: 69-76

29. Levy DL, Heald R (2010) Nuclear size is regulated by importin alpha and Ntf2 in
Xenopus. Cell 143: 288-98

30. Levy DL, Heald R (2015) Biological Scaling Problems and Solutions in Amphibians.
Cold Spring Harb Perspect Biol 8: a019166

31. Lewis NE, Rossant J (1982) Mechanism of size regulation in mouse embryo
aggregates. J Embryol Exp Morphol 72: 169-81

32. Li WM, Webb SE, Chan CM, Miller AL (2008) Multiple roles of the furrow deepening
Ca2+ transient during cytokinesis in zebrafish embryos. Dev Biol 316: 228-48

33. Liu Y, Sepich DS, Solnica-Krezel L (2017) Stat3/Cdc25a-dependent cell proliferation
promotes embryonic axis extension during zebrafish gastrulation. PLoS Genet 13:
e1006564

34. Lloyd AC (2013) The regulation of cell size. Cell 154: 1194-205

35. Loughlin R, Wilbur JD, McNally FJ, Nedelec FJ, Heald R (2011) Katanin contributes to
interspecies spindle length scaling in Xenopus. Cell 147: 1397-407

36. Martin P, Lewis J (1992) Actin cables and epidermal movement in embryonic wound
healing. Nature 360: 179-83

37. Matsiaka OM, Penington CJ, Baker RE, Simpson MJ (2018) Discrete and Continuum
Approximations for Collective Cell Migration in a Scratch Assay with Cell Size
Dynamics. Bull Math Biol 80: 738-757

38. Menon T, Nair S (2018) Transient window of resilience during early development
minimizes teratogenic effects of heat in zebrafish embryos. Dev Dyn 247: 992-1004

39. Menon T, Nair S (2019) Experimental Manipulation of Ploidy in Zebrafish Embryos and
Its Application in Genetic Screens. Methods Mol Biol 1920: 111-128

40. Modlinski JA (1975) Haploid mouse embryos obtained by microsurgical removal of one
pronucleus. J Embryol Exp Morphol 33: 897-905

41. Newport J, Kirschner M (1982a) A major developmental transition in early Xenopus
embryos: I. characterization and timing of cellular changes at the midblastula stage.
Cell 30: 675-86

42. Newport J, Kirschner M (1982b) A major developmental transition in early Xenopus
embryos: II. Control of the onset of transcription. Cell 30: 687-96

43. Norden C, Lecaudey V (2019) Collective cell migration: general themes and new
paradigms. Curr Opin Genet Dev 57: 54-60

44. Nurse P (1985) The genetic control of cell volume. The Evolution of genome size: 185-
196

45. Orr-Weaver TL (2015) When bigger is better: the role of polyploidy in organogenesis.
Trends Genet 31: 307-15

46. Otto SP (2007) The evolutionary consequences of polyploidy. Cell 131: 452-62

47. Pelegri F, Knaut H, Maischein HM, Schulte-Merker S, Nusslein-Volhard C (1999) A
mutation in the zebrafish maternal-effect gene nebel affects furrow formation and vasa
RNA localization. Curr Biol 9: 1431-40

48. Petrie RJ, Gavara N, Chadwick RS, Yamada KM (2012) Nonpolarized signaling
reveals two distinct modes of 3D cell migration. J Cell Biol 197: 439-55

49. Qu Z, Weiss JN, MacLellan WR (2004) Coordination of cell growth and cell division: a
mathematical modeling study. J Cell Sci 117: 4199-207

50. Rossant J (1976) Postimplantation development of blastomeres isolated from 4- and 8-
cell mouse eggs. J Embryol Exp Morphol 36: 283-90

51. Schaks M, Giannone G, Rottner K (2019) Actin dynamics in cell migration. Essays
Biochem

52. Sonawane M, Carpio Y, Geisler R, Schwarz H, Maischein HM, Nuesslein-Volhard C
(2005) Zebrafish penner/lethal giant larvae 2 functions in hemidesmosome formation,
maintenance of cellular morphology and growth regulation in the developing basal
epidermis. Development 132: 3255-65

53. Su TT, O'Farrell PH (1998) Size control: cell proliferation does not equal growth. Curr
Biol 8: R687-9

54. Tarkowski AK (1959) Experiments on the development of isolated blastomers of
mouse eggs. Nature 184: 1286-7

55. Thisse C, Thisse B, Schilling TF, Postlethwait JH (1993) Structure of the zebrafish
snail1 gene and its expression in wild-type, spadetail and no tail mutant embryos.
Development 119: 1203-15

56. Topczewski J, Sepich DS, Myers DC, Walker C, Amores A, Lele Z, Hammerschmidt M,
Postlethwait J, Solnica-Krezel L (2001) The zebrafish glypican knypek controls cell
polarity during gastrulation movements of convergent extension. Dev Cell 1: 251-64

57. Trepat X, Wasserman MR, Angelini TE, Millet E, Weitz DA, Butler JP, Fredberg JJ
(2009) Physical forces during collective cell migration. Nature Physics 5: 426-430

58. Tumaneng K, Russell RC, Guan KL (2012) Organ size control by Hippo and TOR
pathways. Curr Biol 22: R368-79

59. Umen JG (2005) The elusive sizer. Curr Opin Cell Biol 17: 435-41

  FIGURES

801   **Figure 1**

Figure 1: Quantification of cell size reduction during early development in diploid zebrafish
embryos.

β-catenin immunolabels of zebrafish blastoderms (A-F). Animal view of whole embryo at 2 hpf (A).
One of the 4 quadrants imaged at 1k-cell (3 hpf, B), sphere stage (4 hpf, C) and shield stage (6
hpf, D) hpf. Left side of panel E is the mid-hindbrain boundary at 24 hpf and a higher magnification
of top side of E (F). Quantification methodology to obtain cross-sectional areas of cells (G-J). Cell
area distributions, dotted horizontal line within box plots indicate mean areas, n indicates total
number of correctly segmented cells that were used for area calculations from multiple embryos
(K-O). Cell size reduction plotted as log-log plots (P) and stages 2 to 6 hpf re-plotted to show
linear reduction (Q), black line is the total cell population. Proportions of cell size categories during
development (R).

Figure 2

Figure 2: Ploidy alters the cell size norm during late blastula and gastrula stages of zebrafish development.

Comparisons of cell area distributions in haploids and tetraploids to diploid normal values for a particular developmental stage, dotted horizontal line within box plots indicate mean area, n indicates total number of correctly segmented cells that were used for area calculations from multiple embryos (A-D). Asterisks denote statistical significance based on Mann Whitney (p ≤ 0.05). Proportions of cell size categories (F-J). (D) diploids, (H) haploids and (T) tetraploids.

Figure 3

Figure 3: Embryos composed of smaller or larger than normal cells manifest epiboly and gastrulation defects.

DIC images during epiboly and gastrulation (A-D) and gene expression analysis (E-K). For panels A-C, stages are noted according to diploid developmental stages and live appearance of haploids and tetraploids when diploids reached a particular developmental stage were recorded. Haploids and tetraploids initiate epiboly later than controls (A) and are delayed through gastrulation (B, C). For panels in D, the appearance of the tail-bud was taken as end of gastrulation and haploids and tetraploids were found to have a shorter body axis at this stage (D). RNA in situs for *foxa2* (blue) and *gsc* (red) at 7 hpf (E) and *dlx3b* (blue) and *myod1* (red) at 11 hpf (F). Quantification of gastrulation defects shown in panels F, I, G and J (K).

Figure 4

Figure 4: Transcriptional profiling and analysis of PCP pathway transcripts.

Most genes are expressed normally in haploids and tetraploids just after ZGA at 3.5 hpf (black dots in A, B). Few genes are mis-regulated (red dots in A, B), which are genes differentially expressed with q values ≤ 0.05, the False Discovery rate corrected p-values. Heatmap of FPKMs of genes differentially expressed by Log_2 fold change (C), arranged based on diploid expression level, decreasing from top to bottom. Quantitative RT-PCR at 8 hpf for PCP components revealed no significant change in transcript levels in comparison to diploids (D).

Figure 5

Figure 5: Developmental defects due alteration of stage-specific cell size norm can be recapitulated and rescued by transient pharmacological treatment.

Diploids treated with 4% DMSO (A, E, G) or Aph+HU (B, F, H). At shield stages, cell sizes in diploid Aph+HU (B) are larger than in DMSO treated embryos (A). Cell area distributions in diploids and diploid Aph+HU at shield stages, dotted horizontal line within box plots indicate mean area, n indicates total number of correctly segmented cells that were used for area calculations from multiple embryos (C). Asterisks denote statistical significance based on Mann Whitney (p ≤ 0.05). Proportions of cell size categories at shield stsge (D). Diploid Aph+HU embryos have shorter body axis at end of gastrulation (FvsE) and ectoderm cells (blue in H, n=21/45) are further apart (G, n=29/29). Haploid embryos treated with 4% DMSO (I, M, O) or Aph+HU (J, N, P). At shield stages, cell sizes in haploid Aph+HU (J) are larger than in DMSO treated haploids (I). Cell area distributions in diploids, haploids and haploid Aph+HU at shield stages, dotted horizontal line within box plots indicate mean area, n indicates total number of correctly segmented cells that were used for area calculations from multiple embryos (K). Asterisks denote statistical significance based on Mann Whitney (p ≤ 0.05). Black and red asterisks indicate statistical significance in comparison to diploids and haploids, respectively. Proportions of cell size categories at shield stage (L). Bilaterally placed ectoderm cells (blue in P, n=21/27) are closer (O, n=16/26), indicating rescue of dorsal convergence.

Figure 6

Figure 6: Smaller or larger than normal cells migrate sub-optimally during gastrulation.

Tracking of cell migration in clonally labeled embryos (A-D) and in chimeric embryos (E-N).

Graphs show total distance travelled (A), net displacement during migration (B), tortuosity (C) and

average speed (D) of clonally labeled cells in diploids (D), haploids (H) and tetraploids (T) from

shield to bud stages of gastrulation. Diploid (red) cells co-transplanted with haploid or tetraploid

(green) cells into diploid hosts at sphere stage (4hpf, E, H) and imaged through gastrulation (F, G,

I, J). Graphs show total distance travelled (K), net displacement during gastrulation (L) tortuosity

(M) and average speed (N) of diploid (D, red), haploid (H, green) and tetraploid (T, green) cells.

Asterisks denote statistical significance based on Mann Whitney ($p \leq 0.05$).

Figure 7

Figure 7: Smaller or larger than normal cells mis-localize actin during migration.

Dynamic actin localization changes in migrating cells (A-F). Panels A, C and E are higher magnification still frames of one cell during migration from Supplementary Movies 6, 9 and 12, respectively. Actin localization in migrating cells over time shown as lines drawn at angles from the center of a cell, black arrowhead indicates the direction of migration in the movie.

Figure 8

Figure 8: A landscape of optimal cell sizes enables efficient collective cell migration for normal development

Haploid (A), diploid (B) and tetraploid (C) embryos with smaller cells in haploids (box in A), normal cell sizes in diploids (box in B) and larger cells in tetraploids (box in C). Diploid embryos (F) made of small (orange), medium (blue) and large (purple) cells in normal proportions, have stage-specific normal size ranges and gastrulate normally. Haploid and Tetraploid embryos deviate from the stage-specific cell size norm, have more small (D, orange) or large (H, purple) cells, and manifest epiboly and dorsal convergence defects during gastrulation (D, H). Aph+HU increases cell size (rectangular panels above E and G). In diploids, this cell size increase recapitulates in diploids gastrulation defects (G), while in haploids, this partially rescues gastrulation defects (E).