RAPID WHOLE CELL IMAGING REVEALS A CALCIUM-APPL1-DYNEIN NEXUS THAT REGULATES COHORT TRAFFICKING OF STIMULATED EGF RECEPTORS

York H M[1], Patil, A[1], Moorthi U K[1], Kaur A[1], Bhowmik A[1], Hyde G J[3], Gandhi H[1], Gaus K[1,4] and Arumugam S[1,2,4] *

[1]Monash Biomedicine Discovery Institute, Faculty of Medicine, Nursing and Health Sciences, Monash University, Clayton/Melbourne, VIC 3800, Australia
European Molecular Biological Laboratory Australia (EMBL Australia), Monash University, Clayton/ Melbourne, VIC 3800, Australia
[2]Independent scholar, Sydney, Australia
[3]Single Molecule Science, University of New South wales, Sydney, Australia.
[4]ARC Centre of Excellence in Advanced Molecular Imaging, UNSW, Sydney, Australia

*to whom correspondence should be addressed: Senthil.arumugam@monash.edu

ABSTRACT

Multicellular life processes such as proliferation and differentiation depend on cell
surface signaling receptors that bind ligands generally referred to as growth factors.
Recently, it has emerged that the endosomal system provides rich signal processing
capabilities for responses elicited by these factors [1-3]. At the single cell level,
endosomal trafficking becomes a critical component of signal processing, as
exemplified by the epidermal growth factor (EGF) receptors of the receptor tyrosine
kinase family. EGFRs, once activated by EGF, are robustly trafficked to the
phosphatase-enriched peri-nuclear region (PNR), where they are dephosphorylated
[4-8]. However, the details of the mechanisms regulating the movements of
stimulated EGFR in time and space, i.e., towards the PNR, are not known. What
endosomal regulators provide specificity to EGFR? Do modifications to the receptor
upon stimulation regulate its trafficking? To understand the events leading to EGFR
translocation, and especially the early endosomal dynamics that immediately follow
EGFR internalization, requires the real-time, long-term, whole-cell imaging of multiple
elements. Here, exploiting the advantages of lattice light-sheet microscopy [9], we
show that the binding of EGF by its receptor, EGFR, triggers a transient calcium
increase that peaks by 30 s, causing the desorption of APPL1 from pre-existing
endosomes within one minute, the rebinding of liberated APPL1 to EGFR within three
minutes, and the dynein-dependent translocation of APPL1-EGF-bearing endosomes
to the PNR within five minutes. The novel, cell spanning, fast acting network that we
reveal integrates a cascade of events dedicated to the cohort movement of activated
EGFR receptors. Our findings support the intriguing proposal that certain endosomal
pathways have shed some of the stochastic strategies of traditional trafficking, and
have evolved behaviors whose predictability is better suited to signaling [10, 11].
Work presented here demonstrates that our whole cell imaging approach can be a
powerful tool in revealing critical transient interactions in key cellular processes such
as receptor trafficking.

INTRODUCTION

The endocytic system is critical to a cell's ability to faithfully transduce signals and to
process their extracellular environment. At the plasma membrane, EGF stimulation
results in multiple parallel processes — transient increases in Ca^{2+} [12, 13], rapid
reorganization of actin filaments [14] and formation of new clathrin-coated pits (CCPs)
[15]. EGFR has been compared to transferrin in many studies as an example of a

ligand-induced, rather than a constitutive receptor, system[16, 17]. Stimulated EGFRs initiate new CCPs that are distinct from transferrin receptor containing CCPs [15, 18, 19]. Post internalization, the intracellular itineraries of the two receptors are distinct with stimulated EGFR localizing to late endosomes and transferrin receptors recycled back through the early endosomes marked by Rab5[16].

Rab5 effectors such as EEA1 and APPL1 and 2 have been shown to be involved in the pre-early endosomal steps of endosomal maturation [20]. APPL1 can bind directly to Rab5 [21] as well as to a lipid bilayer via a PH BAR domain [22]. APPL1 has been also shown to bind the cytosolic tail of various receptors, including the adiponectin receptor [23], the nerve growth factor receptor [24] and EGFR [20, 25] via its PTB domain. Following endocytosis, EGFR has been shown to enter a population of endosomes marked by APPL1 [10, 20, 26]. It has also been demonstrated that endosomes bearing a subset of clathrin dependent cargoes, including EGFR, are more mobile and mature faster post internalization [27], compared to those that carry transferrin. To better understand the apparent central role of APPL1 in regulating EGFR, its relation to the observed rapid dynamics of EGF bearing endosomes, and more generally, the molecular events and processes that lead to intracellular divergence of EGFR and transferrin receptors, we measure the dynamics of the newly formed EGF bearing endosomes as well as APPL1 within the first ten minutes of stimulation by EGF. Capitalizing on the rapid volumetric imaging capabilities of Lattice Light-Sheet Microscopy (LLSM), we visualized the first events of internalization of EGF stimulated receptors and discovered that 1. APPL1 relocates from pre-existing endosomes onto EGFR puncta and 2. APPL1 binding to EGFR results in a cohort, dynein mediated movement of EGFR towards the peri-nuclear region of the cell where the endosomal maturation and signal quenching of EGFRs occur. The presented novel behavior of EGF bearing endosomes is a significant detail in understanding signal processing by live cells that has not been described elsewhere to our knowledge.

RESULTS

APPL1 responds to EGF stimulation by binding immediately to EGFR bearing endosomes and display enhanced retrograde motility.

To investigate the whole cell distribution of APPL1 on receptor bearing endosomes following internalization of fluorescently labeled EGF or transferrin in living HeLa cells transfected with APPL1-eGFP, we took advantage of the more rapid volumetric imaging and decreased photo-bleaching offered by a lattice light-sheet microscope [9] (LLSM). Fast multi-color, whole-cell imaging revealed the temporal dynamics of all fluorescently labeled endosomal populations. To capture the first cargo bearing

endosomes post-internalization with a time resolution of 4 s for an entire volume of a
cell, we devised a setup that allowed us to inject EGF or transferrin as a temporal
pulse during image acquisition (Supplementary Fig. 1 and Supplementary Movie 1).
50 µL of 100 nM fluorescently labeled ligand was injected near the imaging region,
followed by about 200-300 µL of imaging medium. Thus, in single acquisition
sequences of up to 15 min, we could follow the initial binding of ligands to the
receptors at the plasma membrane, the appearance of early endosomes formed
directly by endocytosis of the ligands, and any of their further movements within the
cell; additionally, APPL1 was tracked using APPL1-GFP. We confirmed that the EGF-
bearing endosomes are not macropinosomes and are likely derived from clathrin
mediated endocytosis (Supplementary Fig. 2).   To independently track the
coordinates of endosomes labeled with APPL1-eGFP, EGF or transferrin ligands, we
performed 2-color (APPL1-GFP and Alexa 647 labeled transferrin or EGF) imaging of
whole cells at 4 s per volume (Supplementary Movies 1,2). From the independently
tracked coordinates of APPL1-GFP as compared to those of the two other ligands,
we quantified the number of spatio-temporally correlated tracks (Methods,
Supplementary Fig. 3). Both transferrin and EGF appeared as diffuse background
fluorescence immediately following injection, and subsequently appeared as
punctate structures on the plasma membrane (Fig. 1a,b). In the case of transferrin,
the punctate structures slowly acquired low levels of APPL1, with an average delay of
100 s (Fig. 1b,g) (Supplementary Movie 2). In contrast, EGF immediately colocalized
with APPL1 at the cell-periphery (Fig. 1a,d)), as indicated by the fraction of EGF tracks
positive for APPL1 (Fig. 1d) and track towards the peri-nuclear region (Supplementary
Movie 1).
25
To further analyze how APPL1 binding affects EGF bearing endosomes, we
developed an analysis workflow that selected the APPL1 tracks that co-tracked with
EGF (Fig. 1c). The overall motility characteristics of these tracks were then determined
using mean square mean square displacement analysis [14]. In general, upon addition
of EGF, the fraction of APPL1-EGF tracks exhibiting directed motion increased slightly
while those with constrained motion decreased (Fig. 1e,f). However, by 5 min, of all
APPL1 vesicles that showed directed motion, over 80% were also found to be positive
for EGF (compare pink/blue curves, Fig. 1e, and columns, Fig. 1f), indicating that
APPL1-EGF-bearing endosomes display a strong tendency towards being more
mobile (Fig. 1 e,f). In contrast, by 5 min after the addition of transferrin, the APPL1-
bearing endosomes most likely to show directed movements were those that lacked
transferrin (Fig. 1h,i). EGF stimulation also resulted in APPL1-bearing endosomes

exhibiting longer tracks, and higher maximum velocities, compared to unstimulated

cells (Fig. 2a).

EGF bearing APPL1 positive endosomes utilize dynein to move

To allow tracking at higher resolution we combined LLSM with micropatterning. On

micropatterned coverslips with 5 μm lines, cells acquired an elongated shape

(Methods and Supplementary Fig. 4). We predicted that the elongated shape of the

cells would accentuate the directionality of endosomal movements, which, in the case

of EGF-bearing APPL1 endosomes, are expected to occur in a PNR-directed,

retrograde manner (Fig. 2b). We positioned the lattice light sheet such that an oblique

cross-section illuminated the nucleus and adjacent PNR of each cell and acquired

images with 200 ms of exposure per frame (Supplementary Movies 3, 4). The number

of retrograde movements of APPL1 endosomes in wild-type cells treated with EGF

was significantly higher than in unstimulated cells (Fig. 2c,e and Supplementary

Movies 3, 4) indicating that EGF stimulation can cause a switch in APPL1 dynamics.

While APPL1 is associated with endosomes that exist prior to EGF-stimulation, once

it becomes attached to endosomes newly formed by the internalization of EGF, those

endosomes are much more likely to move in a retrograde manner. These observations

led us to ask three questions: What motors are engaged to enable retrograde

movement? What causes APPL1 to switch from pre-existing endosomes to newly

generated EGF-bearing endosomes? How does APPL1 localize to EGF bearing

endosomes?

Given the observed retrograde motility of EGF-APPL1-bearing endosomes, we

suspected that the main minus-end directed motor, dynein, might be involved.

Dynein has also been implicated in the translocation of EGF to the juxta-nuclear

region in earlier studies [28]. We expressed a fluorescently labeled version of p150

217 – 548 tagged with dsRed, which sequesters dynein, thus inhibiting dynein-

dynactin-based motility [29]. This allowed us to select dsRed-labeled cells, and then

assay the motility of their EGF and/or APPL1-bearing endosomes during the first 10

min post EGF addition (Supplementary Movie 5). Interestingly, while the total set of

APPL1-bearing endosomes did not show a significant change in motility, in the subset

that co-tracked with an EGF signal, motility was inhibited (Fig. 2f), with a majority of

the tracks showing constrained motion (>60%) and only about 15 % showing directed

motion (Fig. 2f). We also quantified the extent of accumulation of endosomes at the

PNR by measuring the distance from the centroid, for all endosomal populations

(Methods and Fig. 2g). In contrast to controls, no accumulation of EGF vesicles in the

PNR was observed when cells expressed p150 217 – 548 or were treated with another

dynein inhibitor, ciliobrevin (Fig. 2h). These results indicate that dynein is the major
motor protein involved in the translocation of these endosomes. In agreement with
this, the EGF-bearing endosomes exhibited a significantly increased fraction of
constrained movements under dynein inhibition, in contrast to the large fraction of
directed movements seen in untreated cells (Fig. 2i). Therefore, independent
experiments, that perturb dynein in distinct ways, both suggest that the accumulation
of EGF-APPL1-bearing endosomes in the PNR, via minus-end directed translocation,
requires the recruitment of dynein.
9
APPL1 mediated translocation to the perinuclear region is necessary for efficient
maturation of EGF bearing endosomes.
Maturation of APPL1 positive endosomes have been shown to occur through a
process of conversion, where one protein is shed off the surface of the endosome,
and a new endosomal marker is acquired. APPL1 endosomes convert into EEA1
positive endosomes. This has also been demonstrated for EGFR containing
endosomes. We first recapitulated the maturation of EGF bearing endosomes (Fig.
3a-d). With time, the number of EGF tracks that acquired EEA1 increased with time.
Within 5 minutes, a steady fraction of EGF-EEA1 double positive endosome is
achieved (Fig. 3d). As previously described, conversion processes where APPL1 is
shed off and EEA1 is acquired were also observed (Fig. 3c). Inhibiting retrograde
motility by treatment of cells with Ciliobrevin slowed down APPL1 to EEA1 maturation
of EGF bearing endosomes (Fig. 3d). The localization of EGF-APPL1 double positive
endosomes therefore, is necessary to maintain the flux of EGF bearing endosomes
into EGF-EEA1 double positive endosomes through conversion.
25
APPL1 desorption is dependent on calcium wave evoked by EGF stimulation
To investigate the observed recruitment of APPL1, from pre-existing to newly formed
EGF-bearing endosomes, we analyzed the whole-cell dynamics of APPL1. Upon
stimulation with 100 nM EGF, the APPL1 signal was lost from the endosomes of the
PNR (Fig. 4a,b and Supplementary Movie 6, 7) suggesting global redistribution of
APPL1. We used Imaris to segment out the PNR to quantify the loss of APPL1 signal
from the endosomes. We chose the endosomes at the PNR for this measurement as
they are relatively immobile within the time frame of APPL1 desorption, as compared
to the peripheral endosomes (Supplementary Movie 8). We found that PNR
endosomes lose their APPL1 signal completely or partially when stimulated,
respectively, with 100 nM EGF or 20 nM EGF (Fig. 4c). In contrast, upon transferrin
stimulation, we did not observe any APPL1 redistribution. EGF stimulation has been
shown to elicit Ca^{2+} signals through phospholipase cγ (PLCγ) and phospholipase A2

(PLA2) [12, 13]. We speculated that the transient increase in cytosolic Ca^{2+} evoked by
EGF binding could cause the loss of APPL1 from the endosomes. Previous studies of
other proteins that, like APPL1, contain a pleckstrin homology (PH) domain, have
shown that elevated intracellular Ca^{2+} prevented membrane binding of such proteins
through the formation of Ca-phosphoinositide complexes [30]. To determine if
elevated Ca^{2+} is responsible for the unbinding of APPL1, we imaged the dynamic
distribution of APPL1 GFP and the Ca^{2+} indicator R-GECO, after using 100 nM
ionomycin to induce a Ca^{2+} influx (Fig. 4d and Supplementary Movie 9) [31]. Upon
ionomycin addition, APPL1 dissociated from endosomes within 40 s of the R-GECO
signal peak (Fig. 4d). In all cases of ionomycin-mediated desorption, we observed
that the endosomes became transiently immobile, after which APPL1 desorbed from
them before subsequently rebinding to endosomes. We attribute the freezing of
motile endosomes upon an increase in cytosolic Ca^{2+} to 'Calcium-mediated Actin
Reset' (CaAR), which results in a dense meshwork of actin [32]. We also measured the
time of R-GECO peaks with respect to EGF binding, and APPL1 desorption with
respect to EGF binding (Fig. 4e and Supplementary Movie 10). We found the R-
GECO signal peaked about 30 s post EGF stimulation, and APPL1 desorption
occurred in the next 30 s (Fig. 4e).
19
APPL1 binds to EGFR through its PTB domain
It has been previously reported that APPL1 can bind directly to phosphorylated EGFR
through its PTB domain. To test whether this interaction plays a role in early stage
trafficking of EGFR, cells expressing WT-APPL1 were treated with 10 µM erlotinib for
1 h, to inhibit EGFR cross-phosphorylation [33], and imaged following EGF
stimulation (Fig. 5a). APPL1 localization in these cells was extremely rare and
transient. To investigate whether the PTB domain of APPL1 is specifically involved in
the binding to EGFR, as suggested elsewhere [25], we expressed a mutant of APPL1
with the PTB domain deleted (APPL1-ΔPTB) [34]. We found that APPL1-ΔPTB did not
localize to EGF-bearing endosomes and the signals were cytoplasmic. Furthermore,
in both experiments, in cells either expressing APPL1-ΔPTB or treated with erlotinib,
the peri-nuclear localization and directed motion of EGF-bearing endosomes were
strikingly impaired, as demonstrated by the abrogation of the decrease in mean
centroid to endosomal distances (Fig. 5 b,c). Some mobile endosomes were
observed, possibly attributable to kinesins involved in EGFR trafficking [35]. In the
case of cells expressing APPL1-ΔPTB, the impaired localization of EGF endosomes in
the PNR strongly suggests that this deletion mutant functions as a dominant negative.
The experiments together suggest that APPL1 is an intermediary between
phosphorylated EGFRs and dynein and is essential for perinuclear accumulation.

DISCUSSION

A mechanism for cohort transport of signaling EGFRs

Based on the aforementioned experiments that provide evidence of endosomal dynamics and protein redistributions, in Fig. 6 we present a model of the proposed events involving EGF, EGFR, Ca^{2+}, APPL1, and dynein from before, and up to 5 min after, EGF stimulation. Before exposure to EGF, EGFR exists in its monomeric state [36], APPL1 is bound to pre-existing endosomes, and Ca^{2+} levels are low. Within 30 s of EGF-EGFR binding, EGFR dimerizes and is phosphorylated, endocytosis generates the first EGF-bearing vesicles, and Ca^{2+} levels begin to rise. Rising Ca^{2+} levels result in the desorption of APPL1 from pre-existing endosomes presumably by blocking the interaction of the BAR and PH domains of APPL1 and endosomal phosphoinositides, a type of block reported for other PH-domain containing proteins, including Akt [30]. Extracellular EGF concentration controls the degree of Ca^{2+} elevation [13] and in turn, the extent of APPL1 desorption, thereby resulting in a transient availability of cytosolic APPL1 proportional to the EGF concentration as demonstrated by the dependence of APPL1 desorption on EGF concentration. The resulting transient increase in cytosolic APPL1 occurs within approximately 60 s post EGF-binding, within which time EGFR phosphorylation continues in parallel. Previously, EGFR phosphorylation dynamics, measured using a ratio-metric sensor based on EGFR - ECFP and PTB-YFP, revealed that phosphorylation occurred in less than a minute after EGF addition [37]. This is consistent with the dynamics of APPL1 localization to EGFR bearing endosomes, through the PTB domain, observed in our experiments.

APPL1 as a multifunctional endosomal signaling adaptor protein

The multiple interaction sites provided by APPL1's BAR, PH and PTB domains are central to the model. The PTB domain of APPL1 is at the C-terminus and is structurally similar to the PTB domain of Shc [22], which also binds EGFR [38]. The PTB domain of Shc recognizes NPX(phosphor)Y and has been found to bind tighter to phosphorylated peptides than to phosphatidylinositol [39] and may explain the preference of APPL1 binding to phosphorylated EGFR than the membranes though phosphoinositides. Our experiments with APPL1-ΔPTB, where APPL1 binding to EGF bearing endosomes was impaired, support the fact that the PTB domain of APPL1 binds directly to phosphorylated EGFR. Once APPL1 has bound to EGFR, a form of yet unknown interaction appears to engage dynein, and EGF-bearing endosomes are rapidly shunted to the PNR. Our results suggest that dynein acts downstream to

APPL1, in that its activity occurs after APPL1 recruitment, and when APPL1 was blocked from binding to EGFR (by erlotinib or APPL1-ΔPTB), no retrograde motility was seen. Dynein has previously been reported to be required for the translocation of EGF towards the nucleus [28]. While direct interactions between dynein and APPL1 have not been reported, thirteen key phosphorylation sites have been identified on APPL1, and their functions are yet to be fully elucidated [40]. How APPL1 coupled to EGFR promotes dynein engagement is unknown, however it is possible that APPL1 phosphorylation sites are involved. APPL1 phosphorylation has previously been implicated in regulating the recycling of activated GPCRs [41]. Collectively these studies highlight the importance of APPL1 phosphorylation and the need for further investigation of the multiple roles of APPL1 in regulating endosomes.

Endosomal trafficking and signaling

What could be the physiological significance of such rapid trafficking of the EGFR receptors? Endocytosis and trafficking of signaling receptors regulate signal transduction by providing specificity in space and time [3, 36]. Endosomal trafficking allows signaling receptors to be localized at specific regions of the cell or to be channeled, in a timed fashion, towards attenuation sites in multi-vesicular bodies or lysosomes [6]. A recent report by demonstrates that EGFR signaling is controlled by protein tyrosine phosphatases (PTPs) associated with ER, that are enriched in the PNR [8]. The ER associated PTPs (PTP1B and TCPTP) enable spatially restricted activity and efficiently de-phosphorylate the EGFR when EGFR localizes to the PNR [4, 8]. Our results also agree with the distinct rapid motility previously reported for EGF in contrast to transferrin [27], and reveal the APPL1 mediated engagement of dynein as a mechanism for these movements. This study reveals that APPL1 is central to this dynein dependent rapid transport, as well as to the peri-nuclear accumulation of stimulated EGFRs and provides a mechanism distinct from that used by constitutive receptors like transferrin. A recent study also reported APPL1 displaying fast retrograde movement in axonal transport hinting that the dynein-based motility that we describe may be crucial in different cell types in different contexts [42].

Vesicular transport is an important logistical support in the signal processing machinery of the cell. As mentioned above, localization of receptors at the PNR, that results in dephosphorylation and hence signal attenuation is an important step in the signal processing. It is likely that not just the localization, but the rate of localization from the periphery of the cell to the PNR will determine the signal sensing. In a scenario where the receptors depend on the steady state dynamics of the endosomal dynamics, the arrival times of the receptors at the PNR may be broadly distributed. However, as described in this study, if the receptor depends on a specific activated

1 mechanism where a cohort movement of receptors ensue, the arrival times may be
2 restricted to a narrow distribution.
3
CONCLUSION
Our findings also highlight the sophisticated organization of the endosomal system's
interaction matrix, where transient interactions driven by specific biochemistry take
place in timescales of minutes. Our approach reveals a previously unknown EGFR-
APPL1-dynein nexus and highlights how live cell imaging-based studies can unravel
transient, conditional interactions, capturing simultaneous multiple processes
triggered by a single protein. Such transient interactions are perhaps not discernible
in ensemble approaches where temporal resolution, and details of endosomal
specificity and dynamics, are lost.
13

14 ## Methods
15

Cell lines
HeLa and RPE1 (ATCC) cells were incubated at 37°C in 5% CO_2 in high glucose
Dulbecco's modified eagle media (DMEM) (Life Technologies), supplemented with
10% fetal bovine serum (FBS) and 1% penicillin and streptomycin (Life Technologies).
Cells were seeded at a density of 200,000 per well in a 6-well plate containing 5 mm
glass coverslips.
22

Live-cell imaging
Cells were imaged using a lattice light sheet microscope (3i, Denver, Colarado, USA).
Excitation was achieved using 488, 560 and 640 nm diode lasers MPB
communications) at 1-5% AOTF transmittance through an excitation objective
(Special Optics 28.6x 0.7 NA 3.74-mm immersion lens) and is detected via a Nikon
CFI Apo LWD 25x 1.1 NA water immersion lens with a 2.5x tube lens. Live cells were
imaged in 8 mL of 37°C-heated DMEM and were acquired with 2x Hamamatsu Orca
Flash 4.0 V2 sCMOS cameras using Latticescope. 600 μL of 100 nM Alexa647 labelled
EGF or transferrin, or 600 μL of 100 μg/μL dextran-fluorescein was added mid imaging
using a custom syringe-sample holder contraption that allowed precise injection of
the desired volume of fluorescently labelled ligands, followed by imaging media. The
fluorescently labelled ligand containing media were kept separated from the imaging
media by an air bubble (Supplementary Fig. 1). Similarly, non-fluorescent 600 μL of
100 nM ionomycin was added where indicated. Cells were serum starved in high
glucose DMEM medium 4 h prior to EGF addition.
38

Plasmids and transfection

Cells were transfected with pEGFPC1-human APPL1, a gift from Pietro De Camilli (Addgene plasmid #22198) (26), GFP-APPL1-ΔPTB, a gift from Donna Webb (Addgene plasmid #59768) (34), DsRed-p150 217-548, a gift from Trina Schroer (Addgene plasmid #51146) (29), EEA1 TagRFP-T (Addgene plasmid #42635), a gift from Silvia Corvera. Cells were transfected with a total of 1 µg DNA (plasmid of interest – 0.2 µg, blank DNA – 0.8 µg) for single protein expression or (plasmids of interest – 0.2 µg + 0.2 µg, blank DNA – 0.6 µg) using lipofectamine 3000 (Thermo Fisher Scientific) or the Neon electroporator (Invitrogen) at 1200 mV 20 ms, 2 pulses for RPE1 and 1005 mV 35 ms, 2 pulses for HeLa. HeLa cells were ensured for mild expression by the following: The transfection mix consisted of APPL1 or EEA1 plasmids adding up to 20 % of the total DNA amount for transfection, with the rest consisting of blank DNA. Secondly, it has been reported that overexpression of APPL1 or EEA1 results in colocalization of APPL1 and EEA1 on Rab5 endosomes. We optimized this concentration by screening for this artifact, where we see no overlap of APPL1 and EEA1. Thirdly, APPL1 overexpression impairs EGFR internalization. However, in all our experiments, EGF/EGFR complex is trafficked efficiently to the peri-nuclear region.

Micropatterning

Coverslips were sonicated with 70% ethanol for 30 min and plasma cleaned for 5 min. Coverslips were incubated with 1 mg/mL poly(L-lysine)-poly(ethylene-glycol) (PLL-g-PEG) in 1x phosphate buffered saline (PBS) at 4°C overnight. Coverslips were then placed on a chromium mask with 5 µm line patterns and illuminated for 5 min by an ultraviolet lamp. The coverslips were then incubated with 20 µg/mL fibronectin (Thermo Fisher Scientific) in 1x PBS with 0.02% Tween 20 and 0.04% glycerol for 1 h at room temperature. Following which cells were plated onto the slips and allowed to take the shape desired over 4 h (Supplementary Fig. 4).

Drug addition

Cells were incubated with ciliobrevin (CB) or dimethyl sulfoxide (DMSO) at a final concentration of 200 nM in 8 mL DMEM medium for 5 min before and during imaging. Cells were similarly treated with 100 nM final concentration of phorbol12-myristate-13-acetate (PMA) or PBS; and 10 µM erlotinib or DMSO control for 1 h prior to imaging as indicated.

Co-tracking analysis:

Imaris 9.2.1 (Bitplane) was used to detect and track vesicles via the spot detection

feature following Gaussian filtering 0.1 um. The track coordinates were exported and

analyzed using custom codes developed in-house. Tracks which followed the same

trajectory were filtered by the following routine: The errors in the measurement

between the two channels as acquired by the two cameras were obtained by diffusing

Tetraspeck beads (T7279, Thermo Fischer Scientific). This estimation of error

provided the minimum radius for analyzing the 'co-localization' of spots belonging to

two distinct trajectories in two different channels. The spots were considered

colocalizing if they were within a sphere defined by the minimum radius: and a time

filter of 10 consecutive frames, where p = x, y, z coordinates as a function of time for

one channel and c for another (Supplementary Fig. 3). The effective radius of co-

localization was set to be 500 nm to account for the sequential imaging and any

spatial segregation within a single endosome, based on the measurements from

diffusing beads. Next, once all the colocalizing spots were identified, using the

trajectory ID of the spots, they were filtered using a minimum time window of 3 frames

(12 s). We discarded any tracks that co-tracked for less than 12 s as they could be due

to transient interactions. Tables detailing the number of co-tracks identified with time

were exported, averaged for many samples and plotted using ORIGIN.

Motility analysis

The track coordinates were exported from Imaris. MSD analysis was performed as

described in previous studies [43, 44]. Briefly, the identified trajectories were used to

generate MSD plots. The MSD curves were fitted with the following equation: $MSD =$

$< R >^2 = 4Dt^\alpha + 2\sigma^2$, where D is the diffusion coefficient, t is the lag-time, α, the scaling

exponent, and σ is the measurement error measured on the LLSM using beads

adsorbed onto glass, fixed as 35 nm for the fits. The trajectories were characterized

according to the scaling exponent α, retrieved from the fits. Tracks with an

exponential value of <0.4 were termed as constrained, those with an exponential

value of 0.4< α <1.2 termed diffusive and those with a value >1.2 were said to

undergo directed motion.

Peri-nuclear localization (distance from centroid)

Endosomal distributions within the cell were quantified by calculating the centroid of

all endosomal localizations for a given time point. The Euclidian distances of each

endosome from this centroid were calculated. These centroid-to-endosome distances

were pooled, and ensemble statistical analysis was performed. Peri-nuclear

accumulation resulted in lower distances between the centroid and each endosomal

localization.

Author contribution

SA devised and directed the study, HY, AK, SA performed LLSM experiments. HY, AP, SA analyzed data. AB performed the micropatterning, AK, UK, HG contributed to molecular biology, SA, HY and GH wrote the manuscript.

Acknowledgements

The authors would like to thank Profs. Joe Howard, Marino Zerial for engaging discussions and Drs. Vaishnavi Ananthanarayanan, Angika Basant and Robert Weatheritt for valuable comments on the manuscript. This work was supported by ARC LIEF, LE150100163 awarded to Prof. Katharina Gaus.

1
Fig. 1. EGF rapidly colocalizes with APPL1 and is actively co-trafficked to the
perinuclear region.
(a) Cells transfected with APPL1-eGFP (green) were imaged using LLSM, during which
time EGF-A647 (magenta) was pulse injected. The inserts correspond to APPL1 and
EGFA647 from left to right for the denoted time points. Rectangles indicate zoomed
versions for each time point. Time is in seconds. Arrows indicate APPL1 and EGFA647
colocalization. Scale bar = 10 μm. (b) Similar experiment with fluorescently labeled
transferrin. No colocalization was observed. Scale bar = 15 μm. (c) Schematic of
tracking and analysis workflow. All identified tracks, or tracks filtered on the basis of
co-trafficking by presence of both channels within a determined radius sphere at each

time point, were selected for ensemble MSD analysis. Based on the MSD analysis of
the tracks, each track was characterized as constrained, diffusive or undergoing
directed motion (Methods) and the tracks colored correspondingly (bottom) or
exported for grouped statistical analysis. (d) Graph of fraction of cargo tracks with
time (seconds) following EGF addition to HeLa cells transfected with APPL1 EGFP.
Graphs show all the EGF tracks and the fraction of EGF tracks positive for APPL1.
Error bars indicate standard deviation (n = 13 cells). (e) Percentages of APPL1 tracks
categorized as constrained (yellow), diffusive (green) and directed (magenta) motions
as a function of time by MSD analysis in a single cell. In addition, the blue trace shows
the subset of APPL1 tracks positive for EGF that displayed directed motion,
demonstrating that most APPL1 tracks with directed motions were also positive for
the EGF. Error bars correspond to the standard deviation. (f) Percentage of APPL1
tracks undergoing constrained (yellow), diffusive (green), directed (magenta) motions
and directed motions of EGF bearing APPL1 endosomes (blue) that were grouped in
time as pre-cargo addition, 0-5 mins and 5-15 mins post addition based on example
data presented in (e). Error bars correspond to the standard deviation where
applicable (n = 9 cells). (g) Graph of the fraction of cargo tracks with time (seconds)
following transferrin addition to HeLa cells transfected with APPL1 EGFP. Graphs
show all the transferrin tracks and the fraction of transferrin tracks positive for APPL1.
Error bars indicate standard deviation (n = 8 cells). (h) Percentages of APPL1 tracks
categorized as constrained (yellow), diffusive (green) and directed (magenta) motions
as a function of time by MSD analysis in a single cell. In addition, the graphs show the
subset of APPL1 tracks positive for transferrin that display directed motion (grey),
demonstrating that only a subset of APPL1 tracks showed the presence of transferrin
in the initial time points post addition. Error bars correspond to the standard
deviation. (i) Percentages of APPL1 tracks undergoing constrained (yellow), diffusive
(green), directed (magenta) motions and directed motions of transferrin bearing
APPL1 endosomes (grey) that are grouped in time as pre-cargo addition, 0-5 mins
and 5-15 mins post addition based on example data presented in (e). Error bars
correspond to the standard deviation where applicable (n = 7 cells).

Fig. 2. Dynein is responsible for rapid EGF-APPL1 trafficking to the perinuclear

region.

(a) Scatter plot of calculated trajectory lengths in microns against maximum velocity

in um/s of APPL1 endosomes in unstimulated cells (black) and EGF-bearing APPL1

endosomes 2-7 mins following injection (magenta). Line of best fit plotted following

linear regression. (b) Schematic of single plane lattice illumination (dotted red line) of

cells micropatterned in 5 μm patterns (blue). This elongation of the cells accentuates

retrograde motility towards the PNR as indicated by the arrow. (c) Percentages of

APPL1 endosomes which underwent anterograde (grey), retrograde (red) or no net

movement (white) in micropatterned HeLa cells pre-EGF stimulation, 0-5 mins and

beyond 10 mins post-EGF stimulation. Error bars indicate standard deviation. (d)

Scatter plots of the distance in microns of EGF endosomes from a selected centroid in HeLa cells, stimulated with 100 nM EGF, at 0-2 mins and 5-7 mins post addition. The grey plot represents the total amount of EGF bearing endosomes and the blue plots indicate APPL1 positive endosomes bearing EGF. The inner box of the box plot represents the standard deviation, the inner bar the median and the dot the mean, the 'x' represents the counts within 1-99% of the sample and the horizontal bars the range. (e) Kymographs of APPL1 endosomes in HeLa cells pre-EGF stimulation (left) and 0-5mins post-EGF stimulation (right). The top graphs show a zoomed-out view, scale bar = 5 µm y-axis, 5 seconds x-axis. The overlaid lines show the area of the kymograph corresponding to the perinuclear region as seen in the oblique slice insert. The bottom graphs show the area of the dotted box zoomed 5x, '+' indicates plus-end directed motions '-' indicates minus-end directed motions. (f) Percentages of EGF bearing endosome tracks which showed confined (yellow), diffusive (green) and directed (magenta) motion in HeLa cells 5-15 mins post stimulation with 100 nM EGF. The cells were either uninhibited, treated with 50 µM ciliobrevin or transfected with DsRed p150 cc. The error bars represent standard deviation. (g) Schematic of endosomal distribution calculation. Euclidian distances between endosomal positions (circles) and the centroid of all the positions (squares) were calculated within a time interval of 2 mins immediately post EGF stimulation (0 – 2 mins) and 5 mins later (5 – 7 mins). (h) Scatter plots of the endosomal distances from centroid in microns of EGF endosomes in HeLa cells stimulated with 100 nM EGF, at 0-2 mins and 5-7 mins post addition. The dotted box and whisker plots represent cells treated with 50 µM ciliobrevin and the stripped plots cells transfected with DsRed p150 cc. The inner box of the box plot represents the standard deviation, the inner bar the median and the dot the mean, the 'x' represents the counts within 1-99% of the sample and the horizontal bars the range. (i) A bar graph of the percentage of APPL1 (right) and APPL1 + EGF (left) bearing endosome tracks which show confined (yellow), diffusive (green) and directed (magenta) motion in HeLa cells treated with 50 µM ciliobrevin, 5-15 mins post stimulation with 100 nM EGF. The error bars represent standard deviation.

1
Fig. 3. APPL1-EGF endosomes mature by acquiring EEA1 and is dependent on its
retrograde motility and localization to PNR. (a) Representative maximum intensity
projections of 100 mM EGF stimulation of HeLa cells expressing APPL1-eGFP and
EEA1 TagRFP-T (Magenta). Hollow arrows point at few examples of APPL1 and EGF
colocalization and the solid arrows point at APPL1-EEA-EGF1 triple or EEA1-EGF
double positive endosomes. Scale bar = 15 µm (b) A zoomed view of APPL1-EEA1-
EGF triple positive endosomes in the boxed region in (a). Scale bar = 0.5 µm (c) An
EGF-APPL1 double positive endosome undergoing maturation by conversion to EGF-
EEA1 positive endosome. Scale bar = 0.2 µm (d) Graph of fraction EGF tracks positive
for APPL1 and EEA1. Error bars indicate standard deviation (n = 6 cells). (e) Same plot
as (d) for cells treated with 50 µM Ciliobrevin. Double lines on the X-axis indicate
breaks in time.
14
15
16

Fig. 4. EGF mediated Ca²⁺ influx causes transient APPL1 desorption.

(a) Time-lapse montage of cross-sections across a single cell displaying desorption of

APPL1-eGFP following addition of 100 nM EGF. Scale bar = 4 μm. **(b)** Example image

depicting segmentation of APPL1 eGFP signal in the PNR using Imaris and its loss

upon EGF stimulation. Scale bar: 5 μm. **(c)** Change in relative intensity of APPL1-GFP

in the perinuclear region of HeLa cells following addition of an EGF pulse of either

100 nM (blue) or 20 nM (green). Time is recorded in seconds following EGF binding,

error bars represent standard deviation. **(d)** Maximum intensity projection of images

of APPL1-eGFP (top, green in Merge) and R-GECO (middle, magenta in Merge) in

HeLa cells stimulated with 100 nM pulse of ionomycin. Time recorded in seconds

from ionomycin addition. Scale bar = 10 μm. Inserts correspond to zoomed sections

(unbroken and dotted squares respectively) at each time point, scale bar = 1 μm. **(e)**

Timeline of relationships between EGF binding, R-GECO peaks and APPL1 desorption. Measured R-GECO fluorescence peak and APPL1 desorption (grey dots), each dot represents a single experiment, with R-GECO peaks overlaid to mean of EGF induced R-GECO peaks. 100 nM EGF was used to determine both the time of APPL1 desorption (blue) and R-GECO peak (orange).

Fig. 6. Summary figure of proposed mechanism of APPL1-mediated shunt
trafficking of activated EGFR.
In steady state cells, APPL1 (yellow semi-circles) is dispersed throughout the cell. EGF
binding leads to dimerization and phosphorylation of EGFR and an increase in
intracellular Ca^{2+} (blue background). The increase in intracellular Ca^{2+} impairs APPL1
PH binding to phosphoinositide causing APPL1 desorption. The increased cytosolic
APPL1 results in APPL1 binding to activated EGFR via a PTB domain. APPL1 positive
EGF bearing endosomes undergo dynein (purple figure) mediated motility to the
perinuclear region which is rich in ER.

References

1. Di Fiore, P.P. & De Camilli, P. Endocytosis and signaling. an inseparable partnership. *Cell* **106**, 1-4 (2001).

2. Sorkin, A. & von Zastrow, M. Endocytosis and signalling: intertwining molecular networks. *Nat Rev Mol Cell Biol* **10**, 609-622 (2009).

3. Neefjes, J., Jongsma, M.M.L. & Berlin, I. Stop or Go? Endosome Positioning in the Establishment of Compartment Architecture, Dynamics, and Function. *Trends Cell Biol* **27**, 580-594 (2017).

4. Mattila, E. *et al.* Negative regulation of EGFR signalling through integrin-alpha1beta1-mediated activation of protein tyrosine phosphatase TCPTP. *Nat Cell Biol* **7**, 78-85 (2005).

5. Yudushkin, I.A. *et al.* Live-cell imaging of enzyme-substrate interaction reveals spatial regulation of PTP1B. *Science* **315**, 115-119 (2007).

6. Baumdick, M. *et al.* EGF-dependent re-routing of vesicular recycling switches spontaneous phosphorylation suppression to EGFR signaling. *Elife* **4** (2015).

7. Jongsma, M.L. *et al.* An ER-Associated Pathway Defines Endosomal Architecture for Controlled Cargo Transport. *Cell* **166**, 152-166 (2016).

8. Stanoev, A. *et al.* Interdependence between EGFR and Phosphatases Spatially Established by Vesicular Dynamics Generates a Growth Factor Sensing and Responding Network. *Cell Syst* **7**, 295-309 e211 (2018).

9. Chen, B.C. *et al.* Lattice light-sheet microscopy: imaging molecules to embryos at high spatiotemporal resolution. *Science* **346**, 1257998 (2014).

10. Kalaidzidis, I. *et al.* APPL endosomes are not obligatory endocytic intermediates but act as stable cargo-sorting compartments. *J Cell Biol* **211**, 123-144 (2015).

11. Villasenor, R., Nonaka, H., Del Conte-Zerial, P., Kalaidzidis, Y. & Zerial, M. Regulation of EGFR signal transduction by analogue-to-digital conversion in endosomes. *Elife* **4** (2015).

12. Peppelenbosch, M.P., Tertoolen, L.G., den Hertog, J. & de Laat, S.W. Epidermal growth factor activates calcium channels by phospholipase A2/5-lipoxygenase-mediated leukotriene C4 production. *Cell* **69**, 295-303 (1992).

13. Marqueze-Pouey, B., Mailfert, S., Rouger, V., Goaillard, J.M. & Marguet, D. Physiological epidermal growth factor concentrations activate high affinity receptors to elicit calcium oscillations. *PLoS One* **9**, e106803 (2014).

14. Rijken, P.J., Hage, W.J., van Bergen en Henegouwen, P.M., Verkleij, A.J. & Boonstra, J. Epidermal growth factor induces rapid reorganization of the actin microfilament system in human A431 cells. *J Cell Sci* **100 (Pt 3)**, 491-499 (1991).

15. Johannessen, L.E., Pedersen, N.M., Pedersen, K.W., Madshus, I.H. & Stang, E. Activation of the epidermal growth factor (EGF) receptor induces formation of EGF receptor- and Grb2-containing clathrin-coated pits. *Mol Cell Biol* **26**, 389-401 (2006).

16. Dickson, R.B., Hanover, J.A., Willingham, M.C. & Pastan, I. Prelysosomal divergence of transferrin and epidermal growth factor during receptor-mediated endocytosis. *Biochemistry* **22**, 5667-5674 (1983).

17. Hopkins, C.R., Miller, K. & Beardmore, J.M. Receptor-mediated endocytosis of transferrin and epidermal growth factor receptors: a comparison of constitutive and ligand-induced uptake. *J Cell Sci Suppl* **3**, 173-186 (1985).

18. Warren, R.A., Green, F.A. & Enns, C.A. Saturation of the endocytic pathway for the transferrin receptor does not affect the endocytosis of the epidermal growth factor receptor. *J Biol Chem* **272**, 2116-2121 (1997).

19.   Stang, E. *et al.* Cbl-dependent ubiquitination is required for progression of EGF receptors into clathrin-coated pits. *Mol Biol Cell* **15**, 3591-3604 (2004).

20.   Miaczynska, M. *et al.* APPL proteins link Rab5 to nuclear signal transduction via an endosomal compartment. *Cell* **116**, 445-456 (2004).

21.   Zhu, G. *et al.* Structure of the APPL1 BAR-PH domain and characterization of its interaction with Rab5. *EMBO J* **26**, 3484-3493 (2007).

22.   Li, J., Mao, X., Dong, L.Q., Liu, F. & Tong, L. Crystal structures of the BAR-PH and PTB domains of human APPL1. *Structure* **15**, 525-533 (2007).

23.   Mao, X. *et al.* APPL1 binds to adiponectin receptors and mediates adiponectin signalling and function. *Nat Cell Biol* **8**, 516-523 (2006).

24.   Lin, D.C. *et al.* APPL1 associates with TrkA and GIPC1 and is required for nerve growth factor-mediated signal transduction. *Mol Cell Biol* **26**, 8928-8941 (2006).

25.   Lee, J.R. *et al.* Adaptor protein containing PH domain, PTB domain and leucine zipper (APPL1) regulates the protein level of EGFR by modulating its trafficking. *Biochem Biophys Res Commun* **415**, 206-211 (2011).

26.   Zoncu, R. *et al.* A phosphoinositide switch controls the maturation and signaling properties of APPL endosomes. *Cell* **136**, 1110-1121 (2009).

27.   Lakadamyali, M., Rust, M.J. & Zhuang, X. Ligands for clathrin-mediated endocytosis are differentially sorted into distinct populations of early endosomes. *Cell* **124**, 997-1009 (2006).

28.   Driskell, O.J., Mironov, A., Allan, V.J. & Woodman, P.G. Dynein is required for receptor sorting and the morphogenesis of early endosomes. *Nat Cell Biol* **9**, 113-120 (2007).

29.   Quintyne, N.J. & Schroer, T.A. Distinct cell cycle-dependent roles for dynactin and dynein at centrosomes. *J Cell Biol* **159**, 245-254 (2002).

30.   Kang, J.K. *et al.* Increased intracellular Ca(2+) concentrations prevent membrane localization of PH domains through the formation of Ca(2+)-phosphoinositides. *Proc Natl Acad Sci U S A* **114**, 11926-11931 (2017).

31.   Dana, H. *et al.* Sensitive red protein calcium indicators for imaging neural activity. *Elife* **5** (2016).

32.   Wales, P. *et al.* Calcium-mediated actin reset (CaAR) mediates acute cell adaptations. *Elife* **5** (2016).

33.   Ng, S.S., Tsao, M.S., Nicklee, T. & Hedley, D.W. Effects of the epidermal growth factor receptor inhibitor OSI-774, Tarceva, on downstream signaling pathways and apoptosis in human pancreatic adenocarcinoma. *Mol Cancer Ther* **1**, 777-783 (2002).

34.   Broussard, J.A. *et al.* The endosomal adaptor protein APPL1 impairs the turnover of leading edge adhesions to regulate cell migration. *Mol Biol Cell* **23**, 1486-1499 (2012).

35.   Hoepfner, S. *et al.* Modulation of receptor recycling and degradation by the endosomal kinesin KIF16B. *Cell* **121**, 437-450 (2005).

36.   Bakker, J., Spits, M., Neefjes, J. & Berlin, I. The EGFR odyssey - from activation to destruction in space and time. *J Cell Sci* **130**, 4087-4096 (2017).

37.   Offterdinger, M., Georget, V., Girod, A. & Bastiaens, P.I. Imaging phosphorylation dynamics of the epidermal growth factor receptor. *J Biol Chem* **279**, 36972-36981 (2004).

38.   Cantor, A.J., Shah, N.H. & Kuriyan, J. Deep mutational analysis reveals functional trade-offs in the sequences of EGFR autophosphorylation sites. *Proc Natl Acad Sci U S A* **115**, E7303-E7312 (2018).

39.   Smith, M.J., Hardy, W.R., Murphy, J.M., Jones, N. & Pawson, T. Screening for PTB domain binding partners and ligand specificity using proteome-derived NPXY peptide arrays. *Mol Cell Biol* **26**, 8461-8474 (2006).

40.   Gant-Branum, R.L., Broussard, J.A., Mahsut, A., Webb, D.J. & McLean, J.A. Identification of phosphorylation sites within the signaling adaptor APPL1 by mass spectrometry. *J Proteome Res* **9**, 1541-1548 (2010).

41.   Sposini, S. *et al.* Integration of GPCR Signaling and Sorting from Very Early Endosomes via Opposing APPL1 Mechanisms. *Cell Rep* **21**, 2855-2867 (2017).

42.   Goto-Silva, L., McShanee, M.P., Salinas, S., Kalaidzidis, Y., Schiavo, G., Zerial, M., Biorxiv, 2018.

43.   Qian, H., Sheetz, M.P. & Elson, E.L. Single particle tracking. Analysis of diffusion and flow in two-dimensional systems. *Biophys J* **60**, 910-921 (1991).

44.   Zajac, A.L., Goldman, Y.E., Holzbaur, E.L. & Ostap, E.M. Local cytoskeletal and organelle interactions impact molecular-motor- driven early endosomal trafficking. *Curr Biol* **23**, 1173-1180 (2013).