RAPID WHOLE CELL IMAGING REVEALS AN APPL1-DYNEIN NEXUS THAT REGULATES STIMULATED EGFR TRAFFICKING

York H M[1], Kaur A[1], Patil, A[1], Bhowmik A[1], Moorthi U K[1], Hyde G J[2], Gandhi H[1], Gaus K[1,3] and Arumugam S[1,3] *

[1]EMBL Australia Node in Single Molecule Science, School of Medical Sciences, University of New South Wales, Sydney 2052, Australia
[2]Independent scholar, Sydney, Australia
[3]ARC Centre of Excellence in Advanced Molecular Imaging, UNSW, Sydney, Australia

*to whom correspondence should be addressed: senthil.arumugam@unsw.edu.au

ABSTRACT

Multicellular life processes such as proliferation and differentiation depend on cell surface signaling receptors that bind ligands generally referred to as growth factors. Recently, it has emerged that the endosomal system provides rich signal processing capabilities for responses elicited by these factors [1-3]. At the single cell level, endosomal trafficking becomes a critical component of signal processing, as exemplified by the epidermal growth factor (EGF) receptors of the receptor tyrosine kinase family. EGFRs, once activated by EGF, are robustly trafficked to the phosphatase-enriched peri-nuclear region (PNR), where they are dephosphorylated [4-8]. However, the details of the mechanisms regulating the movements of stimulated EGFR in time and space, i.e., towards the PNR, are not known. What endosomal regulators provide specificity to EGFR? Do modifications to the receptor upon stimulation regulate its trafficking? To understand the events leading to EGFR translocation, and especially the early endosomal dynamics that immediately follow EGFR internalization, requires the real-time, long-term, whole-cell imaging of multiple elements. Here, exploiting the advantages of lattice light-sheet microscopy [9], we show that the binding of EGF by its receptor, EGFR, triggers a transient calcium increase that peaks by 30 s, causing the desorption of Adaptor protein, phosphotyrosine interacting with PH domain and leucine zipper 1 (APPL1) from pre-existing endosomes within one minute, the rebinding of liberated APPL1 to EGFR within three minutes, and the dynein-dependent translocation of APPL1-EGF-bearing endosomes to the PNR within five minutes. The novel, cell spanning, fast acting network that we reveal integrates a cascade of events dedicated to the cohort movement of activated EGFR receptors. Our findings support the intriguing proposal that certain endosomal pathways have shed some of the stochastic strategies of traditional trafficking, and have evolved behaviors whose predictability is better suited to signaling [10, 11]. Work presented here demonstrates that our whole cell imaging approach can be a powerful tool in revealing critical transient interactions in key cellular processes such as receptor trafficking.

MAIN

The endocytic system is critical to a cell's ability to faithfully transduce signals and to process their extracellular environment. At the plasma membrane, EGF stimulation results in multiple parallel processes — transient increases in Ca^{2+} [12, 13], rapid reorganization of actin filaments [14] and formation of new clathrin-coated pits (CCPs) [15]. EGFR has been compared to transferrin in many studies as an example of a ligand-induced, rather than a constitutive

receptor, system[16, 17]. Stimulated EGFRs initiate new CCPs that are distinct from transferrin receptor containing CCPs [15, 18, 19]. Post internalization, the intracellular itineraries of the two receptors are distinct with stimulated EGFR localizing to late endosomes and transferrin receptors recycled back through the early endosomes marked by Rab5[16].

Rab5 effectors such as EEA1 and APPL1 and 2 have been shown to be involved in the pre-early endosomal steps of the endosomal matrix [20]. APPL1 can bind directly to Rab5 [21] as well as to a lipid bilayer via a BAR-PH domain [22] at its N-terminus. APPL1 has been also shown to bind the cytosolic tail of various receptors, including the adiponectin receptor [23], the nerve growth factor receptor [24] and EGFR [20, 25] via its phosphotyrosine-binding (PTB) domain at its C-terminus. Following endocytosis, EGFR has been shown to enter a population of endosomes marked by APPL1 [10, 20, 26]. It has also been demonstrated that endosomes bearing a subset of clathrin-dependent cargoes, including EGFR, are more mobile and mature faster post internalization, compared to those that carry transferrin [27]. To better understand the apparent central role of APPL1 in regulating EGFR, its relation to the observed rapid dynamics of EGF bearing endosomes, and more generally, the molecular events and processes that lead to intracellular divergence of EGFR and transferrin receptors, we measure the dynamics of the newly formed EGF bearing endosomes as well as APPL1 within the first ten minutes of stimulation by EGF.

To investigate the whole cell distribution of APPL1 on receptor bearing endosomes following internalization of fluorescently labeled EGF or transferrin in living HeLa cells transfected with APPL1-eGFP, we took advantage of the more rapid volumetric imaging and decreased photo-bleaching offered by a lattice light-sheet microscope [9] (LLSM). Fast multi-color, whole-cell imaging revealed the temporal dynamics of all fluorescently labeled endosomal populations. To capture the first cargo bearing endosomes post-internalization with a time resolution of 4 s for an entire volume of a cell, we devised a setup that allowed us to inject EGF or transferrin as a temporal pulse during image acquisition (Supplementary Fig. 1 and Supplementary Movie 1). 50 µL of 100 nM fluorescently labeled ligand was injected near the imaging region, followed by about 200-300 µL of imaging medium. Thus, in single acquisition sequences of up to 15 min, we could follow the initial binding of ligands to the receptors at the plasma membrane, the appearance of early endosomes formed directly by endocytosis of the ligands, and any of their further movements within the cell; additionally, APPL1 was tracked using APPL1-EGFP. We confirmed that the EGF-bearing endosomes are not macropinosomes and are likely derived from clathrin mediated endocytosis (Supplementary Fig. 2). To independently track the coordinates of endosomes labeled with APPL1-eGFP, EGF or transferrin ligands, we performed 2-color (APPL1-EGFP and Alexa 647 labeled transferrin or EGF) imaging of whole cells at 4 s per volume (Supplementary Movies 2,3). From the independently tracked coordinates of APPL1-EGFP as compared to those of the two other ligands, we quantified the number of spatio-temporally correlated tracks (Methods, Supplementary Fig. 3). Both transferrin and EGF appeared as diffuse background fluorescence immediately following injection, and subsequently appeared as punctate structures on the plasma membrane (Fig. 1a,b). In the case of transferrin, the punctate structures slowly acquired low levels of APPL1, with an average delay of 100 s (Fig. 1b,g) (Supplementary Movie 2_. In contrast, EGF immediately colocalized with APPL1 at the cell-periphery (Fig. 1a,g), as indicated by the fraction of EGF tracks positive for APPL1 (Fig. 1d) and track towards the peri-nuclear region (Supplementary Movie 3).

To further analyze how APPL1 binding affects EGF bearing endosomes, we developed an
analysis workflow that selected the APPL1 tracks that co-tracked with EGF (Fig. 1c). The overall
motility characteristics of these tracks were then determined using mean square displacement
analysis [14]. In general, upon addition of EGF, the fraction of APPL1-EGF tracks exhibiting
directed motion increased slightly while those with constrained motion decreased (Fig. 1e,f).
However, by 5 min, of all APPL1 vesicles that showed directed motion, over 80% were also
found to be positive for EGF (compare pink/blue curves, Fig. 1e, and columns, Fig. 1f), indicating
that APPL1-EGF-bearing endosomes display a strong tendency towards being more mobile (Fig.
1 e,f). In contrast, by 5 min after the addition of transferrin, the APPL1-bearing endosomes most
likely to show directed movements were those that lacked transferrin (Fig. 1h,i). EGF stimulation
also resulted in APPL1-bearing endosomes exhibiting longer tracks, and higher maximum
velocities, compared to unstimulated cells (Fig. 2a).
13
To follow retrograde motilities distinctly from overall endosomal movements, we combined LLSM
with micropatterning. On micropatterned coverslips with 5 μm lines, cells acquired an elongated
shape (Methods and Supplementary Fig. 4). We predicted that the elongated shape of the cells
would accentuate the directionality of endosomal movements, which, in the case of EGF-bearing
APPL1 endosomes, are expected to occur in a PNR-directed, retrograde manner (Fig. 2b,
Supplementary Movie 4). We positioned the lattice light sheet such that an oblique cross-section
illuminated the nucleus and adjacent PNR of each cell and acquired images with 200 ms of
exposure per frame (Supplementary Movies 5,6). The number of retrograde movements of
APPL1 endosomes in wild-type cells treated with EGF was significantly higher than in
unstimulated cells (Fig. 2c,e and Supplementary Movies 5, 6) indicating that EGF stimulation can
cause a switch in APPL1 dynamics. While APPL1 is associated with endosomes that exist prior to
EGF-stimulation, once it becomes attached to endosomes newly formed by the internalization of
EGF, those endosomes are much more likely to move in a retrograde manner. These observations
led us to ask three questions: What motors are engaged to enable retrograde movement? What
causes APPL1 to switch from pre-existing endosomes to newly generated EGF-bearing
endosomes? How does APPL1 localize to EGF bearing endosomes?
30
Given the observed retrograde motility of EGF-APPL1-bearing endosomes, we suspected that
the main minus-end directed motor, dynein, might be involved. Dynein has also been implicated
in the translocation of EGF to the juxta-nuclear region in earlier studies [28]. We expressed a
fluorescently labeled version of $p150\ 217 - 548$ tagged with dsRed, which sequesters dynein,
thus inhibiting dynein-dynactin-based motility [29]. This allowed us to select dsRed-labeled cells,
and then assay the motility of their EGF and/or APPL1-bearing endosomes during the first 10
min post EGF addition (Supplementary Movie 7). Interestingly, while the total set of APPL1-
bearing endosomes did not show a significant change in motility, in the subset that co-tracked
with an EGF signal, motility was inhibited (Fig. 2d), with a majority of the tracks showing
constrained motion (>60%) and only about 15 % showing directed motion (Fig. 2f). We also
quantified the extent of accumulation of endosomes at the PNR by measuring the distance from
the centroid, for all endosomal populations (Methods and Fig. 2g). In contrast to controls, no
accumulation of EGF vesicles in the PNR was observed when cells expressed $p150\ 217 - 548$
or were treated with another dynein inhibitor, ciliobrevin (Fig. 2h). These results indicate that
dynein is the major motor protein involved in the translocation of these endosomes. In agreement
with this, the EGF-bearing endosomes exhibited a significantly increased fraction of constrained
movements under dynein inhibition, in contrast to the large fraction of directed movements seen
in untreated cells (Fig. 2i). Therefore, independent experiments, that perturb dynein in distinct

ways, both suggest that the accumulation of EGF-APPL1-bearing endosomes in the PNR, via minus-end directed translocation, requires the recruitment of dynein.

To investigate the observed recruitment of APPL1, from pre-existing to newly formed EGF-bearing endosomes, we analyzed the whole-cell dynamics of APPL1. Upon stimulation with 100 nM EGF, the APPL1 signal was lost from the endosomes of the PNR (Fig. 3a and Supplementary Movie 8,9) suggesting global redistribution of APPL1. We used Imaris to segment out the PNR to quantify the loss of APPL1 signal from the endosomes. We chose the endosomes at the PNR for this measurement as they are relatively immobile within the time frame of APPL1 desorption, as compared to the peripheral endosomes (Supplementary Movie 10,11). We found that PNR endosomes lose their APPL1 signal completely or partially when stimulated, respectively, with 100 nM EGF or 20 nM EGF (Fig. 3b). In contrast, upon transferrin stimulation, we did not observe any APPL1 redistribution. EGF stimulation has been shown to elicit Ca^{2+} signals through phospholipase c γ (PLC γ) and phospholipase A2 (PLA2) [12, 13]. We speculated that the transient increase in cytosolic Ca^{2+} evoked by EGF binding could cause the loss of APPL1 from the endosomes. Previous studies of other proteins that, like APPL1, contain a pleckstrin homology (PH) domain, have shown that elevated intracellular Ca^{2+} prevented membrane binding of such proteins through the formation of Ca-phosphoinositide complexes [30]. To determine if elevated Ca^{2+} is responsible for the unbinding of APPL1, we imaged the dynamic distribution of APPL1 GFP and the Ca^{2+}indicator R-GECO, after using 100 nM ionomycin to induce a Ca^{2+} influx (Fig. 3c and Supplementary Movie 12) [31]. Upon ionomycin addition, APPL1 dissociated from endosomes within 40 s of the R-GECO signal peak (Fig. 3d). In all cases of ionomycin-mediated desorption, we observed that the endosomes became transiently immobile, after which APPL1 desorbed from them before subsequently rebinding to endosomes. We attribute the freezing of motile endosomes upon an increase in cytosolic Ca^{2+} to 'Calcium-mediated Actin Reset' (CaAR), which results in a dense meshwork of actin [32]. We also measured the time of R-GECO peaks with respect to EGF binding, and APPL1 desorption with respect to EGF binding (Fig. 3e and Supplementary Movie 13). We found the R-GECO signal peaked about 30 s post EGF stimulation, and APPL1 desorption occurred in the next 30 s (Fig. 3e).

It has been previously reported that APPL1 can bind directly to phosphorylated EGFR through its PTB domain. To test whether this interaction plays a role in early stage trafficking of EGFR, cells expressing WT-APPL1 were treated with 10 µM erlotinib for 1 h, to inhibit EGFR cross-phosphorylation [33], and imaged following EGF stimulation (Fig. 4a). APPL1 localization in these cells was extremely rare and transient (Supplementary Movie 14). To investigate whether the PTB domain of APPL1 is specifically involved in the binding to EGFR, as suggested elsewhere [25], we expressed a mutant of APPL1 with the PTB domain deleted (APPL1-ΔPTB) [34]. We found that APPL1-ΔPTB did not localize to EGF-bearing endosomes and the signals were cytoplasmic. Furthermore, in both experiments, in cells either expressing APPL1-ΔPTB or treated with erlotinib, the peri-nuclear localization and directed motion of EGF-bearing endosomes were strikingly impaired, as demonstrated by the abrogation of the decrease in mean centroid to endosomal distances (Fig. 4b,c and Supplementary Movie 15). Some mobile endosomes were observed, possibly attributable to kinesins involved in EGFR trafficking [35]. In the case of cells expressing APPL1-ΔPTB, the impaired localization of EGF endosomes in the PNR strongly suggests that this deletion mutant functions as a dominant negative. The experiments together suggest that APPL1 is an intermediary between phosphorylated EGFRs and dynein and is essential for perinuclear accumulation.

Based on the aforementioned experiments that provide evidence of endosomal dynamics and protein redistributions, in Fig. 5 we present a model of the proposed events involving EGF, EGFR, Ca^{2+}, APPL1, and dynein from before, and up to 5 min after, EGF stimulation. Before exposure to EGF, EGFR exists in its monomeric state [36], APPL1 is bound to pre-existing endosomes, and Ca^{2+} levels are low. Within 30 s of EGF-EGFR binding, EGFR dimerizes and is phosphorylated, endocytosis generates the first EGF-bearing vesicles, and Ca^{2+} levels begin to rise. Rising Ca^{2+} levels result in the desorption of APPL1 from pre-existing endosomes presumably by blocking the interaction of the BAR and PH domains of APPL1 and endosomal phosphoinositides, a type of block reported for other PH-domain containing proteins, including Akt [30]. Extracellular EGF concentration controls the degree of Ca^{2+} elevation [13] and in turn, the extent of APPL1 desorption, thereby resulting in a transient availability of cytosolic APPL1 proportional to the EGF concentration as demonstrated by the dependence of APPL1 desorption on EGF concentration. The resulting transient increase in cytosolic APPL1 occurs within approximately 60 s post EGF-binding, within which time EGFR phosphorylation continues in parallel. Previously, EGFR phosphorylation dynamics, measured using a ratio-metric sensor based on EGFR - ECFP and PTB-YFP, revealed that phosphorylation occurred in less than a minute after EGF addition [37]. This is consistent with the dynamics of APPL1 localization to EGFR bearing endosomes, through the PTB domain, observed in our experiments.

The multiple interaction sites provided by APPL1's BAR, PH and PTB domains are central to the model. The PTB domain of APPL1 is at the C-terminus and is structurally similar to the PTB domain of Shc [22], which also binds EGFR [38]. The PTB domain of Shc recognizes NPX(phosphor)Y and has been found to bind tighter to phosphorylated peptides than to phosphatidylinositol [39] and may explain the preference of APPL1 binding to phosphorylated EGFR than the membranes though phosphoinositides. Our experiments with APPL1-ΔPTB, where APPL1 binding to EGF bearing endosomes was impaired, support the fact that the PTB domain of APPL1 binds directly to phosphorylated EGFR. Once APPL1 has bound to EGFR, a form of yet unknown interaction appears to engage dynein, and EGF-bearing endosomes are rapidly shunted to the PNR. Our results suggest that dynein acts downstream to APPL1, in that its activity occurs after APPL1 recruitment, and when APPL1 was blocked from binding to EGFR (by erlotinib or APPL1-ΔPTB), no retrograde motility was seen. Dynein has previously been reported to be required for the translocation of EGF towards the nucleus [28]. While direct interactions between dynein and APPL1 have not been reported, thirteen key phosphorylation sites have been identified on APPL1, and their functions are yet to be fully elucidated [40]. How APPL1 coupled to EGFR promotes dynein engagement is unknown, however it is possible that APPL1 phosphorylation sites are involved. APPL1 phosphorylation has previously been implicated in regulating the recycling of activated GPCRs [41]. Collectively these studies highlight the importance of APPL1 phosphorylation and the need for further investigation of the multiple roles of APPL1 in regulating endosomes.

What could be the physiological significance of such rapid trafficking of the EGFR receptors? Endocytosis and trafficking of signaling receptors regulate signal transduction by providing specificity in space and time [3, 36]. Endosomal trafficking allows signaling receptors to be localized at specific regions of the cell or to be channeled, in a timed fashion, towards attenuation sites in multi-vesicular bodies or lysosomes [6]. A recent report by demonstrates that EGFR signaling is controlled by protein tyrosine phosphatases (PTPs) associated with ER, that are enriched in the PNR [8]. The ER associated PTPs (PTP1B and TCPTP) enable spatially

restricted activity and efficiently de-phosphorylate the EGFR when EGFR localizes to the PNR [4, 8]. Our results also agree with the distinct rapid motility previously reported for EGF in contrast to transferrin [27], and reveal the APPL1 mediated engagement of dynein as a mechanism for these movements. This study reveals that APPL1 is central to this dynein dependent rapid transport, as well as to the peri-nuclear accumulation of stimulated EGFRs and provides a mechanism distinct from that used by constitutive receptors like transferrin.

Our findings also highlight the sophisticated organization of the endosomal system's interaction matrix, where transient interactions driven by specific biochemistry take place in timescales of minutes. Our approach reveals a previously unknown EGFR-APPL1-dynein nexus and highlights how live cell imaging-based studies can unravel transient, conditional interactions, capturing simultaneous multiple processes triggered by a single protein. Such transient interactions are perhaps not discernible in ensemble approaches where temporal resolution, and details of endosomal specificity and dynamics, are lost.

Methods

Cell lines
HeLa and RPE1 (ATCC) cells were incubated at 37°C in 5% CO_2 in high glucose Dulbecco's modified eagle media (DMEM) (Life Technologies), supplemented with 10% fetal bovine serum (FBS) and 1% penicillin and streptomycin (Life Technologies). Cells were seeded at a density of 200,000 per well in a 6-well plate containing 5 mm glass coverslips. Cells were transfected with 0.3-1 µg DNA using lipofectamine 3000 (Thermo Fisher Scientific) or the Neon electroporator (Invtrogen) at 1200 mV 20 ms, 2 pulses for RPE1 and 1005 mV 35 ms, 2 pulses for HeLa.

Live-cell imaging
Cells were imaged using a lattice light sheet microscope (3i, Denver, Colorado, USA). Excitation was achieved using 488, 560 and 640 nm diode lasers MPB communications) at 1-5% AOTF transmittance through an excitation objective (Special Optics 28.6x 0.7 NA 3.74-mm immersion lens) and is detected via a Nikon CFI Apo LWD 25x 1.1 NA water immersion lens with a 2.5x tube lens. Live cells were imaged in 8 mL of 37°C-heated DMEM and were acquired with 2x Hamamatsu Orca Flash 4.0 V2 sCMOS cameras using Latticescope. 600 µL of 100 nM Alexa647 labelled EGF or transferrin, or 600 µL of 100 µg/µL dextran-fluorescein was added mid imaging using a custom syringe-sample holder contraption that allowed precise injection of the desired volume of fluorescently labelled ligands, followed by imaging media. The fluorescently labelled ligand containing media were kept separated from the imaging media by an air bubble (Supplementary Fig. 1). Similarly, non-fluorescent 600 µL of 100 nM ionomycin was added where indicated. Cells were serum starved in high glucose DMEM medium 4 h prior to EGF addition.

Plasmids
Cells were transfected with 0.3 µg pEGFPC1-human APPL1, a gift from Pietro De Camilli (Addgene plasmid #22198) (26), 0.3 µg GFP-APPL1-ΔPTB, a gift from Donna Webb (Addgene plasmid #59768) (34), 0.7 µg DsRed-p150 217-548, a gift from Trina Schroer (Addgene plasmid #51146) (29).

Micropatterning

Coverslips were sonicated with 70% ethanol for 30 min and plasma cleaned for 5 min.

Coverslips were incubated with 1 mg/mL poly(L-lysine)-poly(ethylene-glycol) (PLL-g-PEG) in 1x

phosphate buffered saline (PBS) at 4°C overnight. Coverslips were then placed on a chromium

mask with 5 μm line patterns and illuminated for 5 min by an ultraviolet lamp. The coverslips

were then incubated with 20 μg/mL fibronectin (Thermo Fisher Scientific) in 1x PBS with 0.02%

Tween 20 and 0.04% glycerol for 1 h at room temperature. Following which cells were plated

onto the slips and allowed to take the shape desired over 4 h (Supplementary Fig. 4).

Drug addition

Cells were incubated with ciliobrevin (CB) or dimethyl sulfoxide (DMSO) at a final concentration

of 200 nM in 8 mL DMEM medium for 5 min before and during imaging. Cells were similarly

treated with 100 nM final concentration of phorbol12-myristate-13-acetate (PMA) or PBS; and

10 μM erlotinib or DMSO control for 1 h prior to imaging as indicated.

Co-tracking analysis:

Imaris 9.2.1 (Bitplane) was used to detect and track vesicles via the spot detection feature

following Gaussian filtering 0.1 um. The track coordinates were exported and analyzed using

custom codes developed in-house. Tracks which followed the same trajectory were filtered by

the following routine: The errors in the measurement between the two channels as acquired by

the two cameras were obtained by diffusing Tetraspeck beads (T7279, Thermo Fischer

Scientific). This estimation of error provided the minimum radius for analyzing the 'co-

localization' of spots belonging to two distinct trajectories in two different channels. The spots

were considered colocalizing if they were within a sphere defined by the minimum radius: and

a time filter of 10 consecutive frames, where p = x, y, z coordinates as a function of time for

one channel and c for another (Supplementary Fig. 3). The effective radius of co-localization

was set to be 500 nm to account for the sequential imaging and any spatial segregation within

a single endosome, based on the measurements from diffusing beads. Next, once all the

colocalizing spots were identified, using the trajectory ID of the spots, they were filtered using

a minimum time window of 3 frames (12 s). We discarded any tracks that co-tracked for less

than 12 s as they could be due to transient interactions. Tables detailing the number of co-tracks

identified with time were exported, averaged for many samples and plotted using ORIGIN.

Motility analysis

The track coordinates were exported from Imaris. MSD analysis was performed as described

in previous studies [42, 43]. Briefly, the identified trajectories were used to generate MSD plots.

The MSD curves were fitted with the following equation: $MSD = <R>^2 = 4Dt^{\alpha} + 2\sigma^2$, where D

is the diffusion coefficient, t is the lag-time, α, the scaling exponent, and σ is the measurement

error measured on the LLSM using beads adsorbed onto glass, fixed as 35 nm for the fits. The

trajectories were characterized according to the scaling exponent α, retrieved from the fits.

Tracks with an exponential value of <0.4 were termed as constrained, those with an exponential

value of 0.4< α <1.2 termed diffusive and those with a value >1.2 were said to undergo

directed motion.

Peri-nuclear localization (distance from centroid)

Endosomal distributions within the cell were quantified by calculating the centroid of all
endosomal localizations for a given time point. The Euclidian distances of each endosome from
this centroid were calculated. These centroid-to-endosome distances were pooled, and ensemble
statistical analysis was performed. Peri-nuclear accumulation resulted in lower distances
between the centroid and each endosomal localization.
6

Author contribution

SA devised and directed the study, HY, AK, SA performed LLSM experiments. HY, AP, SA analyzed data. AB performed the micropatterning, AK, UK, HG contributed to molecular biology, SA, HY and GH wrote the manuscript.

Acknowledgements

The authors would like to thank Profs. Joe Howard, Marino Zerial for engaging discussions and Drs. Vaishnavi Ananthanarayanan, Angika Basant and Robert Weatheritt for valuable comments on the manuscript. This work was supported by ARC LIEF, LE150100163 awarded to Prof. Katharina Gaus.

Fig. 1. EGF rapidly colocalizes with APPL1 and is actively co-trafficked to the perinuclear
region.
(a) Cells transfected with APPL1-eGFP (green) were imaged using LLSM, during which time EGF-
A647 (magenta) was pulse injected. The inserts correspond to APPL1 and EGFA647 from left
to right for the denoted time points. Rectangles indicate zoomed versions for each time point.
Time is in seconds. Arrows indicate APPL1 and EGFA647 colocalization. Scale bar = 10 µm. **(b)**
Similar experiment with fluorescently labeled transferrin. No colocalization was observed. Scale
bar = 15 µm. **(c)** Schematic of tracking and analysis workflow. All identified tracks, or tracks
filtered on the basis of co-trafficking by presence of both channels within a determined radius
sphere at each time point, were selected for ensemble MSD analysis. Based on the MSD analysis
of the tracks, each track was characterized as constrained, diffusive or undergoing directed
motion (Methods) and the tracks colored correspondingly (bottom) or exported for grouped
statistical analysis. **(d)** Graph of fraction of cargo tracks with time (seconds) following EGF

addition to HeLa cells transfected with APPL1 EGFP. Graphs show all the EGF tracks and the fraction of EGF tracks positive for APPL1. Error bars indicate standard deviation (n = 13 cells). **(e)** Percentages of APPL1 tracks categorized as constrained (yellow), diffusive (green) and directed (magenta) motions as a function of time by MSD analysis in a single cell. In addition, the blue trace shows the subset of APPL1 tracks positive for EGF that displayed directed motion, demonstrating that most APPL1 tracks with directed motions were also positive for the EGF. Error bars correspond to the standard deviation. **(f)** Percentage of APPL1 tracks undergoing constrained (yellow), diffusive (green), directed (magenta) motions and directed motions of EGF bearing APPL1 endosomes (blue) that were grouped in time as pre-cargo addition, 0-5 mins and 5-15 mins post addition based on example data presented in (e). Error bars correspond to the standard deviation where applicable (n = 9 cells). **(g)** Graph of the fraction of cargo tracks with time (seconds) following transferrin addition to HeLa cells transfected with APPL1 EGFP. Graphs show all the transferrin tracks and the fraction of transferrin tracks positive for APPL1. Error bars indicate standard deviation (n = 8 cells). **(h)** Percentages of APPL1 tracks categorized as constrained (yellow), diffusive (green) and directed (magenta) motions as a function of time by MSD analysis in a single cell. In addition, the graphs show the subset of APPL1 tracks positive for transferrin that display directed motion (grey), demonstrating that only a subset of APPL1 tracks showed the presence of transferrin in the initial time points post addition. Error bars correspond to the standard deviation. **(i)** Percentages of APPL1 tracks undergoing constrained (yellow), diffusive (green), directed (magenta) motions and directed motions of transferrin bearing APPL1 endosomes (grey) that are grouped in time as pre-cargo addition, 0-5 mins and 5-15 mins post addition based on example data presented in (e). Error bars correspond to the standard deviation where applicable (n = 7 cells).

Fig. 2. Dynein is responsible for rapid EGF-APPL1 trafficking to the perinuclear region.

(a) Scatter plot of calculated trajectory lengths in microns against maximum velocity in μm/s of

APPL1 endosomes in unstimulated cells (black) and EGF-bearing APPL1 endosomes 2-7 mins

following injection (magenta). Line of best fit plotted following linear regression. **(b)** Schematic

of single plane lattice illumination (dotted red line) of cells micropatterned in 5 μm patterns

(blue). This elongation of the cells accentuates retrograde motility towards the PNR as indicated

by the arrow. **(c)** Percentages of APPL1 endosomes which underwent anterograde (grey),

retrograde (red) or no net movement (white) in micropatterned HeLa cells pre-EGF stimulation,

0-5 mins and beyond 10 mins post-EGF stimulation. Error bars indicate standard deviation. **(d)**

Scatter plots of the distance in microns of EGF endosomes from a selected centroid in HeLa cells,

stimulated with 100 nM EGF, at 0-2 mins and 5-7 mins post addition. The grey plot represents

the total amount of EGF bearing endosomes and the blue plots indicate APPL1 positive

endosomes bearing EGF. The inner box of the box plot represents the standard deviation, the

inner bar the median and the dot the mean, the 'x' represents the counts within 1-99% of the

sample and the horizontal bars the range. **(e)** Kymographs of APPL1 endosomes in HeLa cells
pre-EGF stimulation (left) and 0-5mins post-EGF stimulation (right). The top graphs show a
zoomed-out view, scale bar = 5 µm y-axis, 5 seconds x-axis. The overlaid lines show the area
of the kymograph corresponding to the perinuclear region as seen in the oblique slice insert.
The bottom graphs show the area of the dotted box zoomed 5x, '+' indicates plus-end directed
motions '-' indicates minus-end directed motions. **(f)** Percentages of EGF bearing endosome
tracks which showed confined (yellow), diffusive (green) and directed (magenta) motion in HeLa
cells 5-15 mins post stimulation with 100 nM EGF. The cells were either uninhibited, treated with
50 µM ciliobrevin or transfected with DsRed p150 cc. The error bars represent standard
deviation. **(g)** Schematic of endosomal distribution calculation. Euclidian distances between
endosomal positions (circles) and the centroid of all the positions (squares) were calculated within
a time interval of 2 mins immediately post EGF stimulation ($0 - 2$ mins) and 5 mins later ($5 - 7$
mins). **(h)** Scatter plots of the endosomal distances from centroid in microns of EGF endosomes
in HeLa cells stimulated with 100 nM EGF, at 0-2 mins and 5-7 mins post addition. The dotted
box and whisker plots represent cells treated with 50 µM ciliobrevin and the stripped plots cells
transfected with DsRed p150 cc. The inner box of the box plot represents the standard
deviation, the inner bar the median and the dot the mean, the 'x' represents the counts within 1-
99% of the sample and the horizontal bars the range. **(i)** A bar graph of the percentage of
APPL1 (right) and APPL1 + EGF (left) bearing endosome tracks which show confined (yellow),
diffusive (green) and directed (magenta) motion in HeLa cells treated with 50 µM ciliobrevin,
5-15 mins post stimulation with 100 nM EGF. The error bars represent standard deviation.

Fig. 3. EGF mediated Ca²⁺ influx causes transient APPL1 desorption.

(a) Time-lapse montage of cross-sections across a single cell displaying desorption of APPL1-

eGFP following addition of 100 nM EGF. Scale bar = 4 µm. **(b)** Example image depicting

segmentation of APPL1 eGFP signal in the PNR using Imaris and its loss upon EGF stimulation.

Scale bar: 5 µm. **(c)** Change in relative intensity of APPL1-EGFP in the perinuclear region of

HeLa cells following addition of an EGF pulse of either 100 nM (blue) or 20 nM (green). Time

is recorded in seconds following EGF binding, error bars represent standard deviation. **(d)**

Maximum intensity projection of images of APPL1-eGFP (top, green in Merge) and R-GECO

(middle, magenta in Merge) in HeLa cells stimulated with 100 nM pulse of ionomycin. Time

recorded in seconds from ionomycin addition. Scale bar = 10 µm. Inserts correspond to zoomed

sections (unbroken and dotted squares respectively) at each time point, scale bar = 1 µm. **(e)**

Timeline of relationships between EGF binding, R-GECO peaks and APPL1 desorption.

Measured R-GECO fluorescence peak and APPL1 desorption (grey dots), each dot represents

a single experiment, with R-GECO peaks overlaid to mean of EGF induced R-GECO peaks. 100

nM EGF was used to determine both the time of APPL1 desorption (blue) and R-GECO peak (orange).

Fig. 4. APPL1 binds activated EGFR via its PTB domain to mediate minus-directed motility.
(a) Representative maximum intensity projection of LLSM imaging of WT-APPL1-EGFP (green)
in HeLa cells treated with erlotinib 5 min post 100 nM EGF-647 (magenta) stimulation. Scale
bar = 4 µm. Left and right panels indicate zoomed sections corresponding to unbroken and
dotted rectangles respectively, scale bar = 4 µm. **(b)** Percentage of EGF bearing endosome
tracks which showed confined (yellow), diffusive (green) or directed (magenta) motility in HeLa
cells transfected with APPL1-eGFP (no treatment), APPL1-ΔPTB eGFP or APPL1-eGFP and
treated with 10 µM erlotinib. Error bars correspond to the standard deviation. **(c)** Scatter plots
of EGF distance in microns from centroid in HeLa cells treated with 100 nM EGF at 0-2 mins and
5-7-mins post addition as described in Fig. 2g. The Hela cells were either transfected with
APPL1-EGFP (blank plot), APPL1-ΔPTB GFP (diagonal lined plot) or APPL1-EGFP and treated
with 10 µM erlotinib (crossed plot). The inner box of the box plot represents the standard
deviation, the inner bar the median and the dot the mean, the 'x' represents the counts within 1-
99% of the sample and the horizontal bars the range.
15
16

Fig. 5. Summary figure of proposed mechanism of APPL1-mediated shunt trafficking of
activated EGFR.
In steady state cells, APPL1 (yellow semi-circles) is dispersed throughout the cell. EGF binding
leads to dimerization and phosphorylation of EGFR and an increase in intracellular Ca^{2+} (blue
background). The increase in intracellular Ca^{2+} impairs APPL1 PH binding to phosphoinositide
causing APPL1 desorption. The increased cytosolic APPL1 results in APPL1 binding to activated
EGFR via a PTB domain. APPL1 positive EGF bearing endosomes undergo dynein (purple figure)
mediated motility to the perinuclear region which is rich in ER.
10
11
12
13

References

1.   Di Fiore, P.P. & De Camilli, P. Endocytosis and signaling. an inseparable partnership. *Cell* **106**, 1-4 (2001).

2.   Sorkin, A. & von Zastrow, M. Endocytosis and signalling: intertwining molecular networks. *Nat Rev Mol Cell Biol* **10**, 609-622 (2009).

3.   Neefjes, J., Jongsma, M.M.L. & Berlin, I. Stop or Go? Endosome Positioning in the Establishment of Compartment Architecture, Dynamics, and Function. *Trends Cell Biol* **27**, 580-594 (2017).

4.   Mattila, E. *et al.* Negative regulation of EGFR signalling through integrin-alpha1beta1-mediated activation of protein tyrosine phosphatase TCPTP. *Nat Cell Biol* **7**, 78-85 (2005).

5.   Yudushkin, I.A. *et al.* Live-cell imaging of enzyme-substrate interaction reveals spatial regulation of PTP1B. *Science* **315**, 115-119 (2007).

6.   Baumdick, M. *et al.* EGF-dependent re-routing of vesicular recycling switches spontaneous phosphorylation suppression to EGFR signaling. *Elife* **4** (2015).

7.   Jongsma, M.L. *et al.* An ER-Associated Pathway Defines Endosomal Architecture for Controlled Cargo Transport. *Cell* **166**, 152-166 (2016).

8.   Stanoev, A. *et al.* Interdependence between EGFR and Phosphatases Spatially Established by Vesicular Dynamics Generates a Growth Factor Sensing and Responding Network. *Cell Syst* **7**, 295-309 e211 (2018).

9.   Chen, B.C. *et al.* Lattice light-sheet microscopy: imaging molecules to embryos at high spatiotemporal resolution. *Science* **346**, 1257998 (2014).

10.   Kalaidzidis, I. *et al.* APPL endosomes are not obligatory endocytic intermediates but act as stable cargo-sorting compartments. *J Cell Biol* **211**, 123-144 (2015).

11.   Villasenor, R., Nonaka, H., Del Conte-Zerial, P., Kalaidzidis, Y. & Zerial, M. Regulation of EGFR signal transduction by analogue-to-digital conversion in endosomes. *Elife* **4** (2015).

12.   Peppelenbosch, M.P., Tertoolen, L.G., den Hertog, J. & de Laat, S.W. Epidermal growth factor activates calcium channels by phospholipase A2/5-lipoxygenase-mediated leukotriene C4 production. *Cell* **69**, 295-303 (1992).

13.   Marqueze-Pouey, B., Mailfert, S., Rouger, V., Goaillard, J.M. & Marguet, D. Physiological epidermal growth factor concentrations activate high affinity receptors to elicit calcium oscillations. *PLoS One* **9**, e106803 (2014).

14.   Rijken, P.J., Hage, W.J., van Bergen en Henegouwen, P.M., Verkleij, A.J. & Boonstra, J. Epidermal growth factor induces rapid reorganization of the actin microfilament system in human A431 cells. *J Cell Sci* **100 (Pt 3)**, 491-499 (1991).

15.   Johannessen, L.E., Pedersen, N.M., Pedersen, K.W., Madshus, I.H. & Stang, E. Activation of the epidermal growth factor (EGF) receptor induces formation of EGF receptor- and Grb2-containing clathrin-coated pits. *Mol Cell Biol* **26**, 389-401 (2006).

16.   Dickson, R.B., Hanover, J.A., Willingham, M.C. & Pastan, I. Prelysosomal divergence of transferrin and epidermal growth factor during receptor-mediated endocytosis. *Biochemistry* **22**, 5667-5674 (1983).

17.   Hopkins, C.R., Miller, K. & Beardmore, J.M. Receptor-mediated endocytosis of transferrin and epidermal growth factor receptors: a comparison of constitutive and ligand-induced uptake. *J Cell Sci Suppl* **3**, 173-186 (1985).

18.   Warren, R.A., Green, F.A. & Enns, C.A. Saturation of the endocytic pathway for the transferrin receptor does not affect the endocytosis of the epidermal growth factor receptor. *J Biol Chem* **272**, 2116-2121 (1997).

19.   Stang, E. *et al.* Cbl-dependent ubiquitination is required for progression of EGF receptors into clathrin-coated pits. *Mol Biol Cell* **15**, 3591-3604 (2004).

20.   Miaczynska, M. *et al.* APPL proteins link Rab5 to nuclear signal transduction via an endosomal compartment. *Cell* **116**, 445-456 (2004).

21.   Zhu, G. *et al.* Structure of the APPL1 BAR-PH domain and characterization of its interaction with Rab5. *EMBO J* **26**, 3484-3493 (2007).

22.   Li, J., Mao, X., Dong, L.Q., Liu, F. & Tong, L. Crystal structures of the BAR-PH and PTB domains of human APPL1. *Structure* **15**, 525-533 (2007).

23.   Mao, X. *et al.* APPL1 binds to adiponectin receptors and mediates adiponectin signalling and function. *Nat Cell Biol* **8**, 516-523 (2006).

24.   Lin, D.C. *et al.* APPL1 associates with TrkA and GIPC1 and is required for nerve growth factor-mediated signal transduction. *Mol Cell Biol* **26**, 8928-8941 (2006).

25.   Lee, J.R. *et al.* Adaptor protein containing PH domain, PTB domain and leucine zipper (APPL1) regulates the protein level of EGFR by modulating its trafficking. *Biochem Biophys Res Commun* **415**, 206-211 (2011).

26.   Zoncu, R. *et al.* A phosphoinositide switch controls the maturation and signaling properties of APPL endosomes. *Cell* **136**, 1110-1121 (2009).

27.   Lakadamyali, M., Rust, M.J. & Zhuang, X. Ligands for clathrin-mediated endocytosis are differentially sorted into distinct populations of early endosomes. *Cell* **124**, 997-1009 (2006).

28.   Driskell, O.J., Mironov, A., Allan, V.J. & Woodman, P.G. Dynein is required for receptor sorting and the morphogenesis of early endosomes. *Nat Cell Biol* **9**, 113-120 (2007).

29.   Quintyne, N.J. & Schroer, T.A. Distinct cell cycle-dependent roles for dynactin and dynein at centrosomes. *J Cell Biol* **159**, 245-254 (2002).

30.   Kang, J.K. *et al.* Increased intracellular Ca(2+) concentrations prevent membrane localization of PH domains through the formation of Ca(2+)-phosphoinositides. *Proc Natl Acad Sci U S A* **114**, 11926-11931 (2017).

31.   Dana, H. *et al.* Sensitive red protein calcium indicators for imaging neural activity. *Elife* **5** (2016).

32.   Wales, P. *et al.* Calcium-mediated actin reset (CaAR) mediates acute cell adaptations. *Elife* **5** (2016).

33.   Ng, S.S., Tsao, M.S., Nicklee, T. & Hedley, D.W. Effects of the epidermal growth factor receptor inhibitor OSI-774, Tarceva, on downstream signaling pathways and apoptosis in human pancreatic adenocarcinoma. *Mol Cancer Ther* **1**, 777-783 (2002).

34.   Broussard, J.A. *et al.* The endosomal adaptor protein APPL1 impairs the turnover of leading edge adhesions to regulate cell migration. *Mol Biol Cell* **23**, 1486-1499 (2012).

35.   Hoepfner, S. *et al.* Modulation of receptor recycling and degradation by the endosomal kinesin KIF16B. *Cell* **121**, 437-450 (2005).

36.   Bakker, J., Spits, M., Neefjes, J. & Berlin, I. The EGFR odyssey - from activation to destruction in space and time. *J Cell Sci* **130**, 4087-4096 (2017).

37.   Offterdinger, M., Georget, V., Girod, A. & Bastiaens, P.I. Imaging phosphorylation dynamics of the epidermal growth factor receptor. *J Biol Chem* **279**, 36972-36981 (2004).

38.   Cantor, A.J., Shah, N.H. & Kuriyan, J. Deep mutational analysis reveals functional trade-offs in the sequences of EGFR autophosphorylation sites. *Proc Natl Acad Sci U S A* **115**, E7303-E7312 (2018).

39.   Smith, M.J., Hardy, W.R., Murphy, J.M., Jones, N. & Pawson, T. Screening for PTB domain binding partners and ligand specificity using proteome-derived NPXY peptide arrays. *Mol Cell Biol* **26**, 8461-8474 (2006).

40.   Gant-Branum, R.L., Broussard, J.A., Mahsut, A., Webb, D.J. & McLean, J.A. Identification of phosphorylation sites within the signaling adaptor APPL1 by mass spectrometry. *J Proteome Res* **9**, 1541-1548 (2010).

41.   Sposini, S. *et al.* Integration of GPCR Signaling and Sorting from Very Early Endosomes via Opposing APPL1 Mechanisms. *Cell Rep* **21**, 2855-2867 (2017).

42.   Qian, H., Sheetz, M.P. & Elson, E.L. Single particle tracking. Analysis of diffusion and flow in two-dimensional systems. *Biophys J* **60**, 910-921 (1991).

43.   Zajac, A.L., Goldman, Y.E., Holzbaur, E.L. & Ostap, E.M. Local cytoskeletal and organelle interactions impact molecular-motor- driven early endosomal trafficking. *Curr Biol* **23**, 1173-1180 (2013).