Hfq CLASH uncovers sRNA-target interaction networks involved in adaptation to nutrient availability

Ira A. Iosub[1], Marta Marchioretto[2], Brandon Sy[4], Stuart McKellar[1], Karen J. Nieken[3], Rob W. van Nues[3], Jai J. Tree[4], Gabriella Viero[2] and Sander Granneman[1*]

Affiliations:

[1]Centre for Synthetic and Systems Biology, University of Edinburgh, Edinburgh EH9 3BF, UK.

[2]Institute of Biophysics, CNR Unit at Trento, Italy.

[3]Institute of Cell Biology, University of Edinburgh, Edinburgh EH9 3FF, UK.

[4]School of Biotechnology and Biomolecular Sciences, University of New South Wales, Sydney 2052, NSW, Australia.

*To whom correspondence should be addressed:
Sander Granneman
e-mail: sgrannem@ed.ac.uk
Tel: +44 131 6519082

Abstract

By shaping gene expression profiles, small RNAs (sRNAs) enable bacteria to very efficiently adapt to constant changes in their environment. To better understand how *Escherichia coli* acclimatizes to changes in nutrient availability, we performed UV cross-linking, ligation and sequencing of hybrids (CLASH) to uncover sRNA-target interactions. Strikingly, we uncovered hundreds of novel Hfq-mediated sRNA-target interactions at specific growth stages, involving many novel 3'UTR-derived sRNAs and a plethora of sRNA-sRNA interactions. We discovered sRNA-target interaction networks that play a role in adaptation to changes in nutrient availability. We characterized a novel 3'UTR-derived sRNA (MdoR), which is part of a regulatory cascade that enhances maltose uptake by (a) inactivating repressive pathways that block the accumulation of maltose transporters and (b) by reducing the flux of general porins to the outer membrane. Our work provides striking examples of how bacteria utilize sRNAs to integrate multiple regulatory pathways to enhance nutrient stress adaptation.

Microorganisms are renowned for their ability to adapt to environmental changes by
rapidly rewiring their gene expression program. These responses are mediated through
integrated transcriptional and post-transcriptional networks. Control at the transcriptional level
dictates which genes are expressed (Balleza et al., 2009; Martínez-Antonio et al., 2008) and
is well-characterised in *Escherichia coli*. Post-transcriptional regulation is key for controlling
adaptive responses. By using riboregulators and RNA-binding proteins (RBPs), cells can
efficiently integrate multiple pathways and incorporate additional signals into regulatory
circuits. *E. coli* employs many post-transcriptional regulators, including small regulatory RNAs
(sRNAs (Waters and Storz, 2009)), *cis*-acting RNAs (Kortmann and Narberhaus, 2012), and
RNA binding proteins (RBPs) (Holmqvist and Vogel, 2018). The sRNAs are the largest class
of bacterial regulators, which work in tandem with RBPs to regulate their RNA targets (Storz
et al., 2011; Waters and Storz, 2009). The base-pairing interactions are often mediated by
RNA chaperones such as Hfq and ProQ, which help to anneal or stabilize the sRNA and
sRNA-target duplex (Smirnov et al., 2017, 2016; Updegrove et al., 2016). Small RNAs can
repress or stimulate translation and transcription, as well as control mRNA stability
(Sedlyarova et al., 2016; Updegrove et al., 2016; Vogel and Luisi, 2011; Waters and Storz,
2009).

During growth in rich media, *E. coli* are exposed to continuously changing conditions,
such as fluctuations in nutrient availability, pH and osmolarity. Consequently, *E. coli* elicit
complex responses that result in physiological and behavioural changes such as envelope
composition remodelling, quorum sensing, nutrient scavenging, swarming and biofilm
formation. Even subtle changes in the growth conditions can trigger rapid adaptive responses.
Accordingly, each stage of the growth curve is characterised by different physiological states
driven by activation of different transcriptional and post-transcriptional networks. Moreover,
growth phase dependency of virulence and pathogenic behaviour has been demonstrated in
both Gram-positive and Gram-negative bacteria. In some cases a particular growth stage is
non-permissive for the induction of virulence (Mäder et al., 2016; Mouali et al., 2018). Although
the exponential and stationary phases have been characterised in detail(Navarro Llorens et
al., 2010; Pletnev et al., 2015), little is known about the transition between these two phases.
During this transition, the cell population starts to scavenge alternative carbon sources, which
requires rapid remodelling of their transcriptome (Baev et al., 2006a, 2006b; Sezonov et al.,
2007).

To understand sRNA-mediated adaptive responses, detailed knowledge of the
underlying post-transcriptional circuits is required. In *E. coli*, hundreds of sRNAs have been
discovered, but only a fraction of these have been characterised. A key step to unravel the
roles of sRNAs in regulating adaptive responses is to identify their targets. To tackle this
globally, high-throughput methods have been developed that have uncovered a plethora of

sRNA-target interactions, many more than previously anticipated (Han et al., 2016; Hör et al.,

2018; Hör and Vogel, 2017; Lalaouna et al., 2015; Melamed et al., 2016; Waters et al., 2016).

To uncover sRNA-target RNA interaction dynamics that take place during the entry

into stationary phase, we applied UV cross-linking, ligation and sequencing of hybrids

(CLASH) (Helwak et al., 2013; Kudla et al., 2011) on *E. coli*. First, we demonstrate that the

highly stringent purification steps make CLASH a robust method for direct mapping of Hfq-

mediated sRNA-target interactions in *E. coli*. This enabled us to significantly expand on the

sRNA-target interaction repertoire found by RNase E CLASH (Waters et al., 2016) and RIL-

seq (Melamed et al., 2016), and we show that Hfq CLASH can generate very reliable results.

Using CLASH we identified many potentially novel 3'UTR-derived sRNAs, confirming that this

class of sRNAs (Chao et al., 2012, 2017; Chao and Vogel, 2016; Miyakoshi et al., 2015a) is

highly prevalent.

Next, we focussed our analyses on interactions that were specifically recovered during

the transition phase where we identified a surprisingly large number of interactions, including

sRNA-sRNA interactions. Our data suggests that during the transition stage, ArcZ represses

CyaR levels, thereby indirectly controlling genes involving nutrient uptake during the transition

phase. We also characterized a novel 3'UTR-derived sRNA, which we refer to as MdoR (*mal*-

dependent OMP repressor). Unlike the majority of bacterial sRNAs, MdoR is transiently

expressed during the transition phase. We demonstrate that MdoR is a degradation

intermediate of the *malG* 3'UTR, the last transcript of the *malEFG* polycistron that encodes

components of the maltose transport system. We show that MdoR directly downregulates

several mRNAs encoding major porins and suppresses the envelope stress response

controlled by σ^E. We propose that MdoR is part of a regulatory network that, during the

transition phase, promotes accumulation of high affinity maltose transporters in the outer

membrane by repressing competing pathways.

Results

Hfq CLASH in *E. coli*.

To unravel the post-transcriptional networks that underlie the transition between

exponential and stationary growth phases in *E. coli*, we performed CLASH (Helwak et al.,

2013; Kudla et al., 2011) using Hfq as bait (Figure 1A). To generate high quality Hfq CLASH

data made a number of improvements to the original protocol used for RNase E CLASH

(Waters et al., 2016).

Figure 1. Hfq CLASH experiments at different growth phases in *E. coli*.

(**A**) Overview of the critical experimental steps for obtaining the Hfq CLASH data. *E. coli* cells expressing an HTF (His6-TEV-3xFLAG)-tagged Hfq(Jai J. Tree et al., 2014) were grown in LB and an equal number of cells were harvested at different optical densities (OD_{600}). Hfq binds to sRNA-target RNA duplexes, and RNA ends that are in close proximity are ligated together. After removal of the protein, cDNA libraries are prepared and sequenced. The single reads can be used to map Hfq-RNA interactions, whereas the chimeric reads can be traced to sRNA-target interactions. (**B**) A growth curve of the cultures used for the Hfq CLASH experiments, with OD_{600} at which cells were cross-linked indicated by circles, and each growth stage is indicated above the plot. The results show the mean and standard deviations of two biological replicates. Source data are provided as a Source Data file. (**C**) Cultures at the same OD_{600} cross-linked and harvested by filtration were analysed by Hfq CLASH, RNA-seq and Western blotting to detect Hfq.

Our Hfq CLASH protocol has several advantages over the related RIL-seq method (see Materials and Methods and Discussion). As negative controls, replicate CLASH experiments were performed on the untagged parental strain. When combined, the control samples had ~10 times lower number of single-mapping reads and contained only 297 unique chimeric reads, compared to the over 70.000 chimeras identified in the tagged Hfq data. This demonstrates that the purification method produced very low background levels.

Cell samples from seven different optical densities were subjected to Hfq CLASH.
Based on the growth curve analysis shown in Figure 1B, we categorized OD_{600} densities 0.4
and 0.8 as exponential growth phase, 1.2, 1.8, 2.4 as the transition phase from exponential to
stationary, and 3.0 and 4.0 as early stationary phase. To complement the CLASH data, RNA-
seq and Western blot analysis was performed on UV-irradiated cells to quantify steady state
RNA and Hfq protein levels, respectively (Figure 1C, Figure 1-figure supplement 1,
Supplementary Table 1). Western blot analyses revealed that Hfq levels gradually increased
during growth, however, when normalized to the levels of the chaperone GroEL, the increase
was modest (Figure 1-figure supplement 1A-B). To determine the cross-linking efficiency, Hfq-
RNA complexes immobilized on nickel beads were radiolabelled, resolved on NuPAGE gels
and detected by autoradiography. The data show that the recovery of Hfq and radioactive
signal was comparable at each optical density studied (Figure 1-figure supplement 1C).
Comparison of normalized read counts of replicate CLASH and RNA-seq experiments showed
that the results were highly reproducible (Figure 1-figure supplement 2).
145

Hfq binds to the transcriptome in a growth-stage dependent manner.
Meta-analyses of the Hfq CLASH sequencing data revealed that the distribution of Hfq
binding across mRNAs was very similar at each growth stage. We observed the expected Hfq
enrichment at the 5'UTRs and at the 3'UTRs at each growth stage (see Figure 1-figure
supplement 3A and 3B for examples). After identifying significantly enriched Hfq binding peaks
(FDR <= 0.05; see Methods for details) we used the genomic coordinates of these peaks to
search for Hfq binding motifs in mRNAs. The most enriched k-mer included poly-U stretches
(Figure 1-figure supplement 3C) that resemble the poly-U tracts characteristic to Rho-
independent terminators found at the end of many bacterial transcripts (Wilson and Hippel,
1995), and confirms the motif uncovered in CLIP-seq studies in *Salmonella* (Holmqvist et al.,
2016).
157
158

Figure 2. Hfq binding to sRNAs and sRNA steady state levels in _E. coli_ do not always correlate highly.

(**A**) Heatmaps showing the changes in sRNA steady-states (Left) and Hfq cross-linking (Right) relative to OD_{600} 0.4. The clustered RNA-seq data were generated by k-means clustering using the fold-changes of normalized counts (log_2 of transcripts per million (TPM)) relative to OD_{600} 0.4. A blue shade indicates a reduction in levels compared to OD_{600} 0.4, whereas a dark orange shade indicates an increase. The cluster assignment is indicated at the left, and the growth stage is indicated as OD_{600} units below each heatmap. Note that ssrS and ffs encode the cytoplasmic RNAs 6S, a regulator of RNA polymerase, and 4.5S, the signal recognition particle RNA. EyeA is an uncharacterized sRNA mapped by(Sætrom et al., 2005). (**B**) Global correlation of between Hfq binding and steady state RNA levels

increases at higher cell densities. The heatmap shows the changes in Pearson R correlation between
Hfq binding and RNA expression (normalised as in (a)), for sRNAs (top) and all transcripts (bottom).
(**C**) Assessment of correlation between changes in expression and Hfq binding profiles for individual
sRNAs. The y-axis shows the gene name of the sRNAs and the Pearson coefficient indicating the
correlation between Hfq binding and steady state levels for each sRNA is shown on the x-axis.

Given the established role of Hfq in sRNA stabilization and mediating sRNA-target
interactions, it was logical to assume that changes in Hfq binding would also be reflected in
changes in sRNA steady-state levels. This would imply that the Hfq binding data would show
a strong correlation with the RNA-seq data. To test this, we compared the Hfq cross-linking
data to the RNA-seq data. K-means clustering of the normalized data revealed 7 different
patterns of changes in normalised read counts in the Hfq cross-linking and total RNA-seq data
(Figure 2A). Reminiscent of recent work performed in *Salmonella* (Chao et al., 2012), most
sRNAs in *E. coli* appear to be preferentially expressed when the cells reach the transition and
stationary phase (Figure 2A). However, much to our surprise, the Hfq cross-linking profile did
not always follow the same trend (Figure 2A-C, Figure 2-figure supplement 1A). Globally,
changes in sRNA expression did not correlate strongly with the Hfq-binding profile (and *vice*
versa) (Fig 2B-C). The correlation between changes in sRNA expression levels versus
changes in Hfq cross-linking was particularly poor at lower cell density (OD_{600} 0.8: r = 0.10)
but gradually improved as the cells approach stationary phase (OD_{600} 4.0 r =0.54; Figure 2B
and Figure 2-figure supplement 1A). A very similar result was obtained when comparing all
Hfq-bound RNAs, including mRNAs (Figure 2B and Figure 2-figure supplement 1B). Striking
examples are MgrR and PsrD, which showed a strong anti-correlation between Hfq CLASH
and RNA-seq counts (Figure 2C) In the case of PsrD, Hfq binding showed a modest increase
during the growth phase (Figure 2A; right heat map, cluster 6), whereas sRNA levels steadily
decreased (Figure 2A; left heat map; cluster 2). PsrD/SraB has also been shown to bind ProQ
(Smirnov et al., 2017), which may explain why its accumulation does not correlate with Hfq
binding. On the other side of the spectrum, SdsR showed a very high positive Pearson
correlation (Figure 2), suggesting that its accumulation heavily relies on Hfq binding.
We conclude that the dynamics of sRNA expression and binding to Hfq are not always
highly correlated.

Hfq CLASH robustly detects RNA-RNA interactions.
To get a complete overview of the RNA-RNA interactions captured by Hfq CLASH, we
merged the data from the two biological replicates of CLASH growth phase experiments
(Supplementary Table 2.1). Overlapping paired-end reads were merged and unique chimeric
205

206 **Figure 3. Hfq CLASH detects RNA-RNA interactions in *E. coli*.**

(**A**) Intermolecular transcript combinations found in interactions captured by Hfq CLASH. Combination count of all uniquely annotated hybrids on genomic features. *tRNA-tRNA and rRNA-rRNA chimeras originating from different coding regions were removed. (**B**) Venn diagram showing the intersection between interactions from statistically filtered CLASH data from two biological replicates, recovered at three main growth stages: exponential (OD_{600} 0.4 and 0.8), transition (OD_{600} 1.2, 1.8, 2.4) and early stationary (OD_{600} 3.0 and 4.0). (**C**) Comparison of sRNA-mRNA interactions found in RIL-seq S-chimera data and Hfq CLASH data. (**D**) Same as in (**B**) but for sRNA-mRNA interactions. (**E**) (Top) Distribution of chimeras representing statistically filtered interactions, which uniquely map to mRNA genes. Overlapping fragments of the two individual parts of the chimeras were collapsed into single clusters followed by generation of distribution plots. Each gene was divided in 100 bins the number of clusters that map to each bin (hit density; y-axis) was calculated; (Bottom) For the distribution plot around the AUG, the gene length was normalized in 601 bins (x-axis) 5'-end overlap (-300) before the start of the coding sequence, and 300 bins downstream AUG (+300); the bins corresponding to the start codon are indicated with a pink line. (**F**) Enriched motifs in chimeras that uniquely overlap 5'UTRs and 3'UTRs; the logos were drawn using the top 20 K-mers.

reads were identified using the hyb pipeline (Travis et al., 2013). To select RNA-RNA interactions for further analysis, we applied a probabilistic analysis pipeline previously used for the analysis of RNA-RNA interactions in human cells (Sharma et al., 2016) and adapted it for the analyses of RNase E CLASH data (Waters et al., 2016). This pipeline tests the likelihood that observed interactions could have formed spuriously. Strikingly, 87% of the chimeric reads had a Benjamini-Hochberg adjusted p-value of 0.05 or less, indicating that it is highly unlikely that these interactions were generated by random ligation of RNA molecules. These analyses demonstrate the robustness of Hfq CLASH protocol. A complete overview of statistically significantly enriched chimeras is provided in Supplementary Table 2.2.

The distribution of combinations of transcript classes found in the statistically filtered chimeric reads indicates the sRNA-mRNA interactions as the most frequent recovered Hfq-mediated interaction type (~25%; Figure 3A). We suspect that this number might be higher, as about 16% of the chimeras contained sRNA and fragments that mapped to intergenic regions (Figure 3A). Manual inspection of several of these indicated that some of the intergenic sequences were located near genes for which the UTRs were unannotated or short. The vast majority of these interactions were growth stage-specific (Figure 3B). Hfq CLASH identified almost 2000 sRNA-mRNA interactions (Figure 3C; Supplementary Table 2). Around 20% of the interactions found with RIL-seq and ~21% (27 out of 126) of the experimentally verified interactions present in sRNATarbase3 (Supplementary Table 2.7) were recovered. These results suggest that while the CLASH data contained known and many novel interactions, the analyses clearly were not exhaustive.

Meta-analysis revealed that the majority of interactions were identified in the transition

phase (Figure 3D) and that the mRNA fragments found in chimeric reads were strongly

enriched in 5'UTRs peaking near the translational start codon (Figure 3E). The latter is

consistent with the canonical mode of translational inhibition by sRNAs (Bouvier et al., 2008)

and demonstrates the robustness of the data. Enrichment was also found in 3'UTRs of mRNAs

(Figure 3F). Motif analyses revealed a distinct sequence preference in 5'UTR and 3'UTR

binding sites (Figure 3G, Supplementary Table 3). The motifs enriched in the 5'UTRs chimeric

fragments are more consistent with Hfq binding to Shine Dalgarno-like $(ARN)_n$ sequences (Jai

J. Tree et al., 2014) and U-tracts, whereas the 3'UTR-containing chimera consensus motif

corresponds to poly-U transcription termination sites (Figure 3F and Supplementary Table 3).

Hfq CLASH predicts sRNA-sRNA interactions as a widespread layer of post-

transcriptional regulation.

We also uncovered a surprisingly large number of sRNA-sRNA interactions

(Supplementary Table 2.4), many of which were uniquely found in our Hfq CLASH data (Figure

4A). Many interactions were growth-stage specific and the sRNA-sRNA networks show

extensive rewiring across the exponential, transition and stationary phases (Figure 4-figure

supplement 1). The sRNA-sRNA network is dominated by several abundant sRNAs that

appear to act as hubs that have many interacting partners: ChiX, Spot42 (spf), ArcZ and GcvB.

In many cases the experimentally-validated sRNA seed sequences were found in the chimeric

reads, for both established and novel interactions. For example, the vast majority of ArcZ

sRNA-sRNA chimeras contained the known and well conserved seed sequence (Figure 4B,

Figure 4-figure supplement 2).

The sRNA-sRNA chimeras containing CyaR fragments were of particular interest, as

the sRNA is primarily expressed during the transition from late exponential to stationary phase

(De Lay and Gottesman, 2009). In the case of CyaR, the known seed sequence (De Lay and

Gottesman, 2009; Papenfort et al., 2008) as well as a conserved ~25 nt fragment in the 5'

region was found in chimeras (Figure 4-C; Figure 4-figure supplement 2). Similar seed

sequences were identified in CLASH experiments using RNase E as a bait (Waters et al.,

2016), suggesting that this region represents a *bona fide* interaction site. Notably, we identified

ArcZ-CyaR chimeras containing the seed sequence from both sRNAs (Figure 4-figure

supplement 2) and these were detected specifically in the transition phase (Figure 4B-C),

suggesting that these sRNAs could influence each other's activity. To validate these findings,

we used an *E. coli* plasmid-based assay that is routinely used to monitor sRNA-sRNA

interactions and expression of their target mRNAs (Melamed et al., 2016; Miyakoshi et al.,

2015b; Jai J. Tree et al., 2014). An advantage of this system is that each sRNA would be

uncoupled from the chromosomally encoded regulatory networks (that were thought to act

largely in a 1:1 stoichiometry) and to allow the specific effects of the sRNA-target RNA to be assessed (Miyakoshi et al., 2015b). Importantly, these sRNAs were induced during early exponential growth phase when the endogenous (processed) ArcZ and CyaR sRNAs are detectable at only very low levels (Figure 4-figure supplement 3B, lanes 1, 2, 5, 7). The qPCR data were subsequently normalized to the results obtained with a control scrambled sRNA to calculate fold changes in expression levels. Since it is difficult to predict directly from the CLASH data which sRNA in each pair acts as the decoy/sponge, we tested both directions. ArcZ overexpression not only decreased the expression of its mRNA targets (*tpx*, *sdaC*) by more than 50% but also that of CyaR (Figure 5D, panel I). Concomitantly, we observed a substantial increase in CyaR targets *nadE* and *yqaE* (Figure 4D, panel I). CyaR overexpression reduced the level of a direct mRNA target (*nadE*) by ~40% but it did not significantly alter the level of ArcZ or ArcZ mRNA targets (*tpx* and *sdaC*; Figure 4D, panel II). Notably, in this two-plasmid assay CyaR was not expressed at levels higher than ArcZ (Figure 4-figure supplement 3A, panel II). Therefore, it is possible that under the tested conditions the CyaR overexpression was not sufficient to see an effect on ArcZ. We find this unlikely as overexpression of CyaR also did not significantly affect endogenous ArcZ levels, which was ~80-fold less abundant than CyaR in this experiment (Figure 4-figure supplement 3A, panel III). The qPCR results were also confirmed by Northern blot analyses (Figure 4-figure supplement 3B, lanes 1-8), which also demonstrate that ArcZ processing was not affected upon CyaR overexpression. These results suggest that the regulation is unidirectional, reminiscent of what has been described Qrr3 in *Vibrio harveyi* (Feng et al., 2015). We conclude that ArcZ acts as a CyaR anti-sRNA and can trigger its degradation.

To provide additional support for direct interactions between these sRNAs, we generated mutations in the seed sequences of the sRNAs analysed here (Figure 4C). We found that two G to C nucleotide substitutions in ArcZ was sufficient to disrupt ArcZ downregulation of CyaR (Figure 4C-D; ArcZ 70-71 + CyaR). This regulation, however, was almost fully restored when complementary mutations were introduced in the CyaR region (Figure 5C-D; ArcZ 70-71 + CyaR 38-39). These data also demonstrate that it is very unlikely that the observed changes in CyaR levels were be the result of Hfq redistribution due to over-expression of ArcZ over-expression (Moon and Gottesman, 2011; Papenfort et al., 2009), as the ArcZ seed mutant stably accumulated (and therefore effectively binds Hfq), but did not affect CyaR levels. Unexpectedly, the wild-type ArcZ was also able to effectively suppress the CyaR seed mutant (Figure 5D; ArcZ + CyaR 38-39), possibly because ArcZ can still form stable base-pairing interactions with the CyaR mutant.

These results, together with the CLASH data, strongly support the notion that ArcZ and CyaR base-pair *in vivo*, resulting in degradation of CyaR but not *vice versa*.

Figure 4. sRNA-RNA interactions identified by CLASH are growth-stage specific.

(**A**) Hfq CLASH uncovers sRNA-sRNA networks: comparison between statistically filtered sRNA-sRNA

interactions in the Hfq CLASH data, RIL-seq S-chimeras (Melamed et al., 2016) (log and stationary)

and RNase E CLASH (Waters et al., 2016). Only core genome sRNAs were considered. Red coloured
sRNA-sRNA interactions have been characterized in more detail. (**B**) Heatmaps showing the read
density (log$_2$(chimera count)) of chimeric fragments mapping to ArcZ and CyaR. (Top) ArcZ regions
involved in sRNA-sRNA interactions. The location of the known ArcZ seed sequence is indicated above.
(Bottom) CyaR heatmaps that show all CyaR chimeras and CyaR-sRNA chimeras, respectively. The
location of the known CyaR seed sequence, as well as a new seed, is indicated above. (**C**) Base-pairing
interactions predicted from the ArcZ-CyaR chimeras using RNAcofold. The nucleotide substitutions for
experimental validation of direct base-pairing are shown as red or green residues. (**D**) SRNA-sRNA
interactions coordinate nutritional stress responses. ArcZ, CyaR were overexpressed and the levels of
their targets were monitored by RT-qPCR. The *tpx* and *sdaC* mRNAs are ArcZ mRNA targets. The
nadE and *yqaE* mRNAs are CyaR targets. The *dppA* mRNA is a GcvB target. Experiments were
performed in biological and technical triplicates; Error bars indicate the standard error of the mean
(SEM) of the three biological replicates. The dashed horizontal line indicates the level of the
overexpressed scrambled RNA. (**E**) ArcZ and CyaR directly interact. The sRNAs and mutants as in (**C**)
were ectopically co-expressed in *E. coli* and CyaR and CyaR 38-39 levels were quantified by RT-qPCR.
337
338

Hfq CLASH identifies novel sRNAs in untranslated regions
Two lines of evidence from our data indicate that many more mRNAs may be
harbouring sRNAs in their UTRs or be involved in base-pairing among themselves. First,
around 11% of the intermolecular chimeras mapped to mRNA-mRNA interactions (Figure 3A).
Secondly, we observed extensive binding of Hfq in 3'UTRs near transcriptional terminators
(Figure 1-figure supplement 3A, C), indicating that like in *Salmonella*, *E. coli* 3'UTRs may
harbour many functional sRNAs (Chao et al., 2017). We identified 122 3'UTR-containing
mRNA fragments that were involved in 550 interactions. Sixty-five of these interactions were
also identified in the RIL-seq S-chimeras data (Melamed et al., 2016). Eighteen of the 3'UTRs
were found as part of chimeras in the RIL-seq data, while 10 appeared stabilised upon
transient inactivation of RNase E performed in *Salmonella* (TIER-seq data (Chao et al., 2017));
Figure 6A, Supplementary Tables 2.5 and 2.6). Out of the 550 3'UTR-mRNA chimeric reads,
79 were 3'UTRs fused to 5'UTRs of mRNAs, suggesting that these may represent 3'UTR-
derived sRNAs that base-pair with 5'UTRs of mRNAs, a region frequently targeted by sRNAs
(Supplementary Table 2.6). Strikingly, 223 interactions contained 3'UTR fragment of *cpxP*, 51
of which were also found in the RIL-Seq data (Supplementary Table 2.6). In *Salmonella cpxP*
harbours the CpxQ sRNA (Chao and Vogel, 2016). Our analyses greatly increased the
number of potential CpxQ mRNA targets and show that the vast majority of CpxQ interactions
take place during the transition and stationary phases (Supplementary Table 2.6).

We identified six mRNA 3'UTRs that were uncovered in all three (Hfq CLASH, RIL-seq

and TIER-seq) datasets (Figure 5A), suggesting these likely contain sRNAs released from

3'UTRs by RNase E processing. Northern blot analyses confirmed the presence of sRNAs in

malG and *gadE* 3'UTRs (Figure 5B, Figure 5-figure supplement 1A). The latter was also

recently experimentally confirmed in the RIL-seq data and was annotated as GadF (Melamed

et al., 2016). Furthermore, significant Hfq cross-linking could be detected in the 3'UTRs of

these transcripts (Figure 5C). In addition, we could show that the 3'UTR of *ygaM*, which was

found in chimeric reads in our data, also likely harbours a ~100 nt sRNA (hereafter referred to

as YgaN; Figure 5-figure supplement 1A) and robust Hfq cross-linking could be detected in

this region (Figure 5B-C).

To substantiate these results, we analysed RNA-seq data from a study that used

Terminator 5'-Phosphate Dependent Exonuclease (TEX) to map transcription start sites

(TSS) of coding and non-coding RNAs in *E. coli* (Thomason et al., 2015). TEX degrades

processed transcripts that have 5' monophosphates, but not primary transcripts with 5'

triphosphates. Therefore, these data enabled us to determine whether (a) a TSS was detected

in the 3'UTR and whether these were generated by RNase-dependent processing (TEX

sensitive) or originated from an independent promoter (TEX insensitive). For 47 of the 122

predicted 3'UTR-derived sRNAs TEX data provided strong evidence for the presence of

sRNAs (Figure 5-figure supplement 1B-C, Supplementary Table 2.5 and see Data and Code

availability). The TEX data indicate that *ygaM* has (at least) two promoters, one of which is

located near the 3' end of the gene that we predict is the TSS for YgaN (Figure 5-figure

supplement 1B). Furthermore, we speculate that YgaN is processed by RNases. This is based

on the observation that multiple YgaN species were detected in the Northern blot analyses

(Figure 5-figure supplement 1A) and the TEX data indicate that shorter YgaN RNAs are

sensitive to TEX treatment (Figure 5-figure supplement 1B).

The majority of the sRNAs we analysed are more abundant at higher cell densities

(including GadF, YgaN and RybB; see Figure 5B, 2A). In sharp contrast, 3' UTR *malG* sRNA

was expressed very transiently and peaked at an OD_{600} of 1.8 (Figure 5B). We envisage that

the particularly transient expression of this sRNA may be correlated with a role in the adaptive

responses triggered during transition from exponential to stationary phases of growth. We

named it MdoR (*mal*-dependent OMP repressor) and characterized it in detail.

The steady state levels of *malG* and *malEFG* transcripts recapitulate the same

expression profile: both peak at OD_{600} of 1.8 and drop to very low levels at OD_{600} 2.4 (Figure

6A). Additionally, we identified shorter *malG* 3'UTR-containing fragments of intermediate

length between *malG* and MdoR that could be degradation intermediates (Figure 6B). The

detection of these intermediate species suggests that the *malEFG* primary transcript is

undergoing serial ribonucleolytic cleavage steps.

MdoR is a 104 nt sRNA that contains part of the *malG* coding sequence, including the

stop codon and the Rho-independent terminator (Figure 5-figure supplement 1A-B). Two lines

of evidence suggest that MdoR is generated via endonucleolytic cleavage: RNase E cleavage

was detected in the 3'UTR of *malG* in the *Salmonella* TIER-seq data (Chao et al., 2017).

Secondly, the TEX RNA-seq data supported the existence of a short RNA in the same region

that has a 5' monophosphate (Figure 5-figure supplement 1B-C). We verified the MdoR TEX

data by Northern blot analyses (Figure 6C). Consistent with the TEX RNA-seq data, MdoR

could not be detected in our RNA samples treated with TEX (Figure 6C, lanes 2 and 4),

confirming it bears a 5' monophosphate. Like MdoR, the positive control CpxQ (Chao and

Vogel, 2016) was degraded in the presence of TEX (Figure 6C). In contrast, RybB, an sRNA

with a 5' triphosphate transcribed from an independent promoter (Johansen et al., 2006;

Papenfort et al., 2006) was a poor substrate for the exonuclease (Figure 6C).

Figure 6. MdoR is a degradation product that emerges at the transition between exponential and stationary phase of growth.

(**A**) The *mal* regulon gene expression peaks at the transition between exponential and stationary phases of growth: the plot shows averages of $\log_2$(TPM) normalized RNA steady-state levels (y-axis); the x-axis indicates the cell densities (OD_{600}) at which samples were taken. (**B**) Northern blot using total RNA from *E. coli* harvested at different cell densities (OD_{600}) probed with an oligo antisense to *malG* 3'UTR; 23 S rRNA was used as the loading control; the identity of the bands is indicated at the left of the panel; horizontal bars indicate *malEFG* degradation intermediates. (**C**) MdoR is a degradation product: Northern blot using total RNA from cells at indicated OD_{600} with (+; lanes 2 and 4) or without (-; lanes 1 and 3) 5'-Phosphate-Dependent Exonuclease (TEX) treatment. The sRNAs RybB (5'ppp) and CpxQ (5'p) were used as negative and positive controls, respectively. The 5S rRNA is a loading control.

The text on the right of the blot indicates the phosphorylation state of the 5'-termini for each sRNA.

TEX treatment of the total RNA also reduced the levels of the longer intermediate

species as well as the full-length *malG*. These data support a mechanism by which the full-

length polycistronic RNA undergoes decay that is initiated at a site in the upstream *malEFG*

region. The distal gene *malE* is clipped off by the degradosome and selectively stabilized,

allowing it to be expressed at higher levels than other members of the operon (Newbury et al.,

1987). The *malG* 3'UTR, however, would be less susceptible to degradation as it is stabilized

by Hfq binding.

MdoR directly regulates the expression of major outer membrane porins and represses

the envelope stress response pathway

The MdoR chimeras frequently contained 5'UTR fragments of two mRNAs encoding

major porins, *ompC* and *ompA* (Figure 7A-C), which were also significantly enriched in the

RIL-seq data (Figure 8B). The Hfq CLASH data, however, also contained MdoR fragments

fused to a *cis*-encoded sRNA, OhsC (Figure 7B-C), suggesting that either MdoR controls its

expression by sponging/degradation, or *vice versa*). The most abundant and favourable

interactions of MdoR with mRNAs (*ompC* and *ompA*) appear to be utilizing roughly the same

region of the *malG* 3'UTR for base-pairing (Figure 7C), suggesting that the corresponding site

on the predicted sRNA may be a main, functional seed. The predicted interaction between

MdoR and *ompC* is unusually long and consists of two stems interrupted by a bulge,

suggesting that these two RNAs form a stable complex. Conservation analyses and *in silico*

target predictions (CopraRNA (Wright et al., 2014, 2013)) indicate that the seed sequence

predicted by CLASH is relatively well-conserved (Figure 7-figure supplement 1A), and could

be utilized for the regulation of multiple targets (Figure 7-figure supplement 1B).

To verify the MdoR CLASH data we pulse-overexpressed the sRNA from a plasmid-

borne arabinose inducible promoter followed by RNA sequencing (Figure 7D-E). To minimize

secondary changes in gene expression, cells were harvested after only 15 minutes of MdoR

induction. Note that induction was performed at OD_{600} = 0.4, when endogenous levels of MdoR

are very low. As a control we used cells harbouring an empty vector. Differential gene

expression analysis (DESeq2 (Love et al., 2014)), identified ~20 transcripts that were

significantly enriched in the control data compared to the MdoR overexpression data (Figure

7E-F; Supplementary Table 4). Thus, these transcripts are likely downregulated by MdoR *in*

vivo. This set of transcripts included the sigma factor *rpoE* (σ^E), which plays an important role

in controlling gene expression during stress responses, including envelope stress (Alba and

Gross, 2004; Bossi et al., 2008; De Las Peñas et al., 1997; Rhodius et al., 2006). The

observed reduction in mRNA levels of the anti-σ^E protein RseA can be explained by the fact

 that σ^E and RseA are encoded by the same operon.

Figure 7. **MdoR destabilizes *ompC* and *ompA* mRNAs and downregulates key members the σ^E regulon.**

(**A**) The three genome browser tracks show the distribution of target mRNA and sRNA fragments (*ompC, ompA* and OhsC) that were fused to MdoR fragments in chimeric reads. (**B**) MdoR-target interactions found in Hfq CLASH, RIL-seq S-chimera data (log and stationary phase) and RNase E CLASH data. (**C**) MdoR forms stable duplexes with the 5'UTR of porin-encoding mRNAs and the OhsC sRNA. *In silico* prediction (RNAcofold (Lorenz et al., 2011)) of hybrid structures derived from the most abundant MdoR chimeric reads with the indicated transcripts. The min. MFE is the minimum folding energy assigned by RNAcofold. (**D**) Pulse-overexpression of MdoR using L-arabinose. The empty pBAD plasmid served as a negative control. Samples were harvested 15-minutes after induction. (**E**) MdoR regulates expression of various mRNAs. DESeq2 analyses were performed on RNA-seq data from three biological replicates. Red points indicate differentially expressed transcripts. Transcripts with a $\log_2$ fold-change > 0 were enriched in the Control data; those with a $\log_2$ fold-change < 0 were enriched in the MdoR overexpression data. The annotated, enlarged red data points indicate several differentially expressed transcripts discussed in the text. (**F**) The MdoR seed region is important for target regulation. RT-qPCR analysis of several differentially expressed transcripts (gene names shown at the top of the plots), in the presence of sRNAs. The plasmid-borne sRNAs were induced for 15 minutes using L-arabinose. The sRNA names are indicated below each bar; the 'no sRNA' sample contains the empty plasmid as reference for fold-change calculations; *recA* was used as the internal reference gene; experiments were performed in technical triplicates; the standard error of the mean (SEM) of two biological replicates are reported as error bars.

Intriguingly, MdoR overexpression also reduced the levels of sRNAs RyeA and MicA, the latter of which depends on σ^E for its expression (Udekwu and Wagner, 2007). In *Salmonella*, MicA downregulates LamB, a high affinity maltose/maltodextrin transporter (Bossi and Figueroa-Bossi, 2007). Fragments of three mRNAs (*ompC, ompA* and *ptsH*) that were found in MdoR chimeric reads were also differentially expressed in the MdoR overexpression RNA-seq data, providing strong evidence that these are direct MdoR targets.

All the available data suggest that MdoR is part of a mixed coherent feed forward regulatory network (FFL) that enhances the uptake of maltose/maltodextrin by maltose transporters (see Discussion). By base-pairing with the 5'UTRs of the mRNAs it firstly reduces the flux of the more general porins such as OmpC and OmpA to the outer membrane. Secondly, we propose that MdoR stimulates the accumulation of the high-affinity maltose porin LamB in the OMP by suppressing the inhibitory σ^E pathway and MicA. To further test this model, we performed additional validation experiments. First, we confirmed the DESeq2 results for a number of the regulated genes (*ompC, rpoE*, *micA* and *ryeA*) by RT-qPCR (Figure 7F). We included an MdoR mutant in which seed sequence in stem 1 was changed into its complementary sequence (Figure 8-figure supplement 1A). As control we also included the RybB sRNA, which regulates *rpoE* and *ompC* expression (Gogol et al., 2011; Papenfort et al., 2006; Thompson et al., 2007). In all cases, the MdoR SM mutations reduced the negative

regulatory effect on target expression (Figure 7F).

To demonstrate direct target regulation *in vivo*, we employed a well-established
reporter system where an sRNA is co-expressed with a construct containing the mRNA target
region fused to the coding sequence of superfolder green fluorescent protein (sfGFP)
(Corcoran et al., 2012; Urban and Vogel, 2007) (Figure 8-figure supplement 1B). Fusions for
ompC, *ompA* and σ^E were constructed, but only the OmpC and OmpA-sfGFP reporters
produced stable fusions that could be analysed. We also included an MdoR sRNA seed
sequence mutant (MdoR SM) and an *ompC* mutant containing compensatory mutations
(OmpC SM; Figure 8-figure supplement 1A). Unfortunately, our OmpA-GFP reporter construct
that contained compensatory mutations in the target region did not generate a stable fusion.
Therefore, we were unable to use this reporter system to verify the MdoR-*ompA* interaction.
As positive controls we used the MicC and RybB sRNAs as they both regulate *E. coli ompC*
expression (Chen et al., 2004; Gogol et al., 2011). Fluorescence measurements confirmed
that levels of OmpC-sfGFP and OmpA-sfGFP fusions were significantly lower in cells
expressing MdoR (Figure 8-figure supplement 1C). Importantly, MdoR overexpression did not
change the expression of the GFP reporter itself (Figure 8-figure supplement 1C). Mutating
the MdoR seed region largely restored OmpA- and OmpC-sfGFP reporter levels to the levels
of the no sRNA negative control. The MdoR SM mutant was still able to partially suppress the
expression of the OmpC SM-sfGFP mutant, suggesting that base-pairing interactions between
these two mutants is less stable compared to the wild-type (Figure 8-figure supplement 1C).
The wild-type MdoR was also able to partially suppress the expression of the *ompC* SM
mutant. We suggest that the predicted base-pairing interactions between MdoR and *ompC* in
the second stem (Figure 8-figure supplement 1A) might be sufficient to partially suppress
ompC expression. Regardless, the data strongly imply that MdoR directly regulates *ompC*
expression.

To determine whether the changes in fluorescence signal correlate with changes in
reporter mRNA levels, we measured the expression of the GFP reporters by RT-qPCR. The
results were essentially identical to the GFP fluorescence measurements (Figure 8A);
Overexpression of the wild-type MdoR, but not seed mutant, reduced *ompC*-sfGFP mRNA
levels. The *ompC* seed mutation (SM) did not fully disrupt regulation by wild-type MdoR.
However, the MdoR mutant containing compensatory mutations (SM mutant) was able to
much better suppress the *ompC* SM mRNA levels, consistent with the idea that base-pairing
was (largely) restored. Next, we performed polysome profiling experiments to assess the level
of *ompC* translation upon overexpression of MdoR. Although MdoR overexpression did not
noticeably affect 70S and polysome levels (Figure 8-figure supplement 2A), we observed a
significant (~37%) reduction of *ompC* mRNA in the polysomal fractions, relative to the upper
fractions (Figure 8-figure supplement 2B). We conclude that MdoR can regulate *ompC*

expression at the post-transcriptional and translational level.

557
558

Figure 8. MdoR modulates expression of key factors involved in maltose intake.

(**A**) MdoR directly downregulates *ompC* mRNA through base-pairing interactions: RT-qPCR analyses

of the *ompC-sfGFP* and *ompC SM-sfGFP* fusions expression in the presence of MdoR. The bars

indicate the mean fold-change in expression relative to the no sRNA Control (horizontal dashed line).

Error bars indicate the standard error of the mean from two biological replicates. The significance of the

differences between the WT and MdoR SM was assessed with a two-tailed Student's t-test. (**B**)

Endogenous MdoR and *malG* expression is induced during growth on maltose, and *ompC* levels are

significantly lower in maltose compared to glucose. Total RNA extracted from exponentially growing

cells (OD_{600} 0.5) and MdoR, *malG* (Left), and MdoR direct (*ompC*) and indirect (MicA) targets was

quantified by RT-qPCR. The data were normalized to 5S rRNA levels. The bars indicate the mean fold-

change in expression relative to expression in cells growing in glucose (indicated on the plot with a

horizontal dashed line). Error bars indicate the standard error of the mean from two biological replicates.

(**C**) The mutant MdoR is ~50% less abundant than the wild-type. Cell growth and RT-qPCR analysis

of MdoR and *malG* expression was performed as in (b). The bars indicate the mean fold-change in

expression relative to the wild-type. Error bars indicate the standard error of the mean from two

biological replicates. (**D**) Increased MicA and decreased *lamB* levels in the MdoR seed mutant. P-values

were calculated using with a one-sample t-test. (**E**) MdoR seed sequences are important for RNase E

recruitment and MdoR biogenesis. Northern blot that compares MdoR and longer *malG*-3'UTR

containing fragments expression in wild-type *E. coli* and the MdoR seed mutant strain.

MdoR enhances maltoporin expression during maltose fermentation

To further substantiate these results, we next switched to a more controlled system to

investigate the effect of endogenous MdoR on its targets. To determine whether MdoR has a

role in adaptation to maltose-metabolising conditions, single overnight cultures grown in

glucose were split and (re)inoculated in fresh medium containing either glucose or maltose as

the sole carbon sources, and expression of several *mal* regulon genes and MdoR targets were

quantified. We show that MdoR and its parental transcript *malG* are almost undetectable

during growth in glucose, and highly expressed during growth in maltose (~35-fold increase,

Figure 8B). This is consistent with catabolite repression of the *mal* regulon by glucose, and its

induction my maltose (Boos and Shuman, 1998). Intriguingly, we observed that *ompC* mRNA

levels are overall significantly lower during growth in maltose, compared to glucose (Figure

8B). This suggests that porin expression is also regulated by nutrient source in *E. coli*.

Similarly, MicA, a repressor of LamB synthesis, has reduced levels in maltose compared to

glucose (Figure 8B). We next mutated the entire seed sequence of the chromosomal copy of

MdoR (both stem 1 and 2; Figure 8-figure supplement 1A) in the chromosomal context, to

completely disrupt base-pairing with *ompC* mRNA. Notably, the fully-processed mutant MdoR

sRNA is less abundant than the wild-type (Figure 9C) and longer (unprocessed) fragments

that contain upstream *malG* regions could be readily detected (Figure 9E). We speculate that

the mutation in the upstream MdoR sequence might have affected RNase E recruitment or

cleavage, impairing MdoR processing. The MdoR mutant strain also accumulated significantly

higher levels of MicA (Figure 8D) and less *lamB*. Levels of *ompC* mRNA levels were also

slightly elevated in the mutant in maltose relative to the parental strain, although in this case a similar increase was also observed in glucose.

Collectively, the data suggest a role for MdoR in enhancing the uptake of maltose when more favourable carbon sources become limiting.

Discussion

Microorganisms need to constantly adapt their transcriptional program to counteract changes in their environment, such as changes in temperature, cell density and nutrient availability. In bacteria, small RNAs (sRNAs) and their associated RNA-binding proteins are believed to play a key role in this process. By controlling translation and degradation rates of mRNAs upon stress imposition (Holmqvist and Wagner, 2017; Nitzan et al., 2017; Shimoni et al., 2007), they can regulate the kinetics of gene expression as well as suppress noisy signals (Beisel and Storz, 2011), enabling organisms to more efficiently adapt to environmental changes. A major challenge for bacteria is the transition from exponential growth to stationary phase, when the most favourable nutrients become limiting. To counteract this challenge, cells need to rapidly remodel their transcriptome to be able to efficiently metabolize alternative carbon sources. This transition is very dynamic and involves activation and repression of diverse metabolic pathways. However, it is unclear to what degree sRNAs contribute to this transition. The most useful piece of information would be to know what sRNAs are upregulated during this transition phase and to identify their RNA targets. This would help to uncover the regulatory networks that govern this adaptation, as well as provide a starting point for more detailed functional analyses on sRNAs predicted to play a key role in this process. For this purpose, we performed UV cross-linking, ligation and sequencing of hybrids (CLASH (Kudla et al., 2011)) to unravel the sRNA-target interactions during this transition. Using Hfq as a bait we uncovered thousands of unique sRNA-target interactions. Our data are consistent with previously published work (Melamed et al., 2016; Waters et al., 2016) but we also identified almost 1700 novel sRNA-mRNA interactions and over 100 novel sRNA-sRNA interactions. We experimentally validated several of the interactions found in our CLASH findings. We identified functional sRNA-sRNA interactions and describe a novel 3'UTR derived sRNA that plays a role in enhancing uptake of an alternative carbon source during the transition to stationary phase.

Figure 9. CLASH uncovers sRNA-target interaction networks that regulate adaptation to changes in nutrient availability.

(**A**) Model for MdoR biogenesis and its role in post-transcriptional regulation. During maltose (grey circles) utilization, MalT transcribes the *malEFG* operon that encodes components for maltose uptake. A fraction of the *malEFG* transcripts is degraded by RNase E, generating the 3'UTR-derived MdoR sRNA. MdoR base-pairs with *ompC* mRNA, leading to its decay. Additionally, MdoR downregulates expression of *ompA* and MicA that represses LamB synthesis. Therefore, as part of a feed-forward loop

(FFL), MdoR promotes accumulation of maltose specific porins, which facilitates maltodextrin uptake.
Increased production of LamB exerts additional pressure on the OM assembly machinery. This
extracytoplasmic stress, is suppressed by σ^E activation, that controls the expression of stress-
responsive genes and most of the assembly and insertion factors. But σ^E activation is not desirable in
these conditions for two reasons: its unbalanced activation has precarious effects on the cell, and would
produce high amounts of MicA, that would ultimately repress LamB production. For these reasons,
MdoR regulation includes a regulatory arm that mitigates excessive σ^E response through MdoR
repression of *rpoE* expression and prevention of σ^E activation. The latter is achieved by downregulation
of *omp* mRNAs, which indirectly relieves envelope stress. (**B**) sRNA-target networks regulating
peptide/amino acids and maltose/maltodextrin uptake. Boxed are the key transcription factors, sRNAs
are in bold and italicized names are mRNA targets. CRP-cAMP induces expression of CyaR, which is
repressed by ArcZ, specifically during the transition phase, a regulatory circuit that connects multiple
pathways related to the onset of stationary phase and biofilm formation to quorum sensing, cellular
adherence and the nutritional state of the cell. MalT transcribes *lamB* and, via *malEFG* transcription,
promotes MdoR accumulation. MdoR indirectly promotes LamB synthesis by repressing the opposing
σ^E pathway and its sRNA, MicA. Thus, MalT and MdoR jointly promote *lamB* expression, forming a
mixed coherent FFL.

Hfq CLASH

Our *S. cerevisiae* Cross-linking and cDNA analysis data (CRAC; (Granneman et al.,
2009)) showed that a percentage of the cDNAs were formed by intermolecular ligations of two
RNA fragments (chimeras) known to base pair *in vivo* (Kudla et al., 2011). These findings
prompted us to develop a refined protocol to enrich for sRNA-target chimeric reads using Hfq
as an obvious bait. The initial Hfq UV cross-linking data (CRAC; (Tree et al., 2014)) did not
yield sufficiently high numbers of chimeric reads to extract new biological insights. In line with
observations from other groups (Bandyra et al., 2012; Bruce et al., 2018; Morita et al., 2005),
it was proposed that duplexes formed by Hfq are rapidly transferred to the RNA degradosome
(Bandyra et al., 2012; Bruce et al., 2018; Morita et al., 2005). This can cause an extensive
reduction in the likelihood of capturing sRNA-target interactions with Hfq using CLASH
(Waters et al., 2016). However, a recent study demonstrated that Hfq can be used effectively
as a bait to enrich for sRNA-target duplexes under lower-stringency purification conditions
suggesting that sRNA-mRNA duplexes are sufficiently stable on Hfq during purification
(Melamed et al., 2016). This encouraged us to further optimize the CLASH method. We made
a number changes to the protocol that, when combined, enabled us to recover a large number
of sRNA-target chimeric reads (detailed in Materials and Methods). We shortened various
incubation steps to minimize RNA degradation and performed very long and stringent washes
after bead incubation steps to remove background binding of non-specific proteins and RNAs.
Crucially, we very carefully controlled the RNase digestion step that is used to trim the cross-

linked RNAs prior to making cDNA libraries, ensuring the recovery of longer chimeric RNA
fragments. The resulting cDNA libraries were paired-end sequenced to increase the recovery
of chimeric reads with high mapping scores from the raw sequencing data. These
modifications led to a substantial improvement in the recovery of chimeric reads (9.5%
compared to 0.001%. 0.71% were intermolecular chimeras).

Both RIL-seq and Hfq CLASH have advantages and disadvantages, but they are highly
complementary approaches. A major strength of CLASH, however, is that the purification
steps are performed under highly stringent and denaturing conditions. During the first FLAG
affinity purification steps the beads are extensively washed with high salt buffers and the
second Nickel affinity purification step is done under completely denaturing conditions (6M
guanidium hydrochloride). These stringent purification steps enable can significantly reduce
noise by strongly enriching for RNAs covalently cross-linked to the bait protein (Granneman
et al., 2009). Indeed, we show that Hfq CLASH can generate high quality RNA-RNA interaction
data with very low background: Only a few hundred chimeric reads were found in control
datasets, compared to the over 70.000 that co-purified with Hfq. The RIL-seq library
preparation protocol uses an rRNA depletion step to remove contaminating ribosomal RNA,
whereas for Hfq CLASH this is not necessary. Our library preparation protocol also includes
the use of random nucleotides in adapter sequences to remove potential PCR duplicates
("collapsing") from the data.

The very stringent purification conditions used in CLASH could, in some cases also be
a disadvantage as it completely relies on UV cross-linking to isolate directly bound RNAs. In
cases where protein-RNA cross-linking efficiencies are low (for example proteins that only
recognize double-stranded RNA), RIL-seq may be a better approach as it does not completely
rely on UV cross-linking (Melamed et al., 2016).

A large number of interactions were unique to both RIL-seq and Hfq CLASH datasets,
which we believe can be explained by a number of technical and experimental factors. The
denaturing purification conditions used with CLASH completely disrupts the Hfq hexamer
((Tree et al., 2014) and this work) and therefore during the adapter ligation reactions the RNA
ends are likely more accessible for ligation. In support of this, in the RIL-Seq data, the sRNAs
are mostly found in the second half of the chimeras (Melamed et al., 2016), in the Hfq CLASH
data we see sRNAs fragments roughly equally distributed in both sides (45% in left fragment
and 55% in right fragment). Indeed, it was proposed that in RIL-seq the 3' end of the sRNA is
buried in the hexamer and therefore not always accessible for ligation (Melamed et al., 2016).

For the RIL-seq experiments, the authors harvested the cells at 4ºC and they
resuspend the cells in ice-cold PBS prior to UV irradiation (Melamed et al., 2018, 2016), which
results in a cold-shock that can affect the sRNA-interactome as well as sRNA stability. We
cross-link actively growing cells in their growth medium and we UV irradiate our cells only for

several seconds using the Vari-X-linker we recently developed (van Nues et al., 2017). We use filtration devices to rapidly harvest our cells (less than 30 seconds) and the filtered cells are subsequently stored at -80ºC. We previously showed that filtration combined with short UV cross-linking times dramatically reduces noise introduced by the activation of the DNA damage response and significantly increased the recovery of short-lived RNA species (van Nues et al., 2017). We speculate that many of the interactions that are unique to our Hfq CLASH data represent short-lived RNA duplexes that are preferentially captured with our UV cross-linking and rapid cell filtration setup.

The 3'UTR derived sRNA MdoR functions in a mixed feed forward loop by suppressing opposing pathways

We found a large number of chimeras that represent over a hundred distinct intermolecular interactions between 3'UTRs and other mRNA regions, which implicate direct mRNA-mRNA communication. These interactions could have been formed by 3'UTR fragments that have been processed by RNase E (or other RNases) or by an sRNA located within the 3'UTR that transcribed from an internal promoter. These 3'UTR fragments are primarily described as decoys or sponges for other sRNAs (Miyakoshi et al., 2015b), but could act as *trans*-acting sRNAs as well (Chao and Vogel, 2016). Among the interactions in our data we found 79 interactions between 3'UTRs and 5'UTRs (Supplementary Table 2.6). We speculate that many of these represent novel 3'UTR-derived sRNAs that target 5'UTRs of mRNAs. Reanalysis of RNA-seq data from a study that globally mapped *E. coli* transcription start sites (TSS) (Thomason et al., 2015) identified 47 TSS in these 3'UTR, strongly suggesting that these 3'UTRs host sRNAs. The majority of these 3'UTR fragments appear to have 5' monophosphates. We also verified some of our findings by Northern blot analyses. Thus, we have potentially uncovered many novel 3'UTR-derived sRNAs.

Some of these predictions were validated by our follow up work (discussed below) and, while this work was in progress, also by others (Melamed et al., 2016). One of the 3'UTR-derived sRNAs we uncovered (MdoR) was of particular interest as it is only detected during the transition from late exponential to early stationary phase. A model of how MdoR contributes to maltose uptake is shown in Figure 9A.

The genetic structure and transcriptional regulation of the *mal* regulon are well understood. However, its post-transcriptional regulation has remained largely unexplored. Our work uncovered new links between the maltose uptake (*mal*) regulon, envelope stress-responses and membrane composition/assembly pathways. While initially cells metabolize more favourable carbon sources such as glucose, these are generally rapidly depleted and bacteria need to quickly switch to alternative carbon sources, such as maltose and maltodextrins. During maltose utilisation, the *malEFG* operon is transcribed by the MalT

transcription factor and the transcript is processed by RNase E as well as other degradosome

components (Khemici and Carpousis, 2004). Here we show that the MdoR sRNA is a product

of *malEFG* processing, which is protected from further degradation by Hfq.

MdoR is unique in a sense that it not only a 3'UTR derived sRNA that targets multiple

pathways, but also because it is part of a mixed coherent feed forward loop (FFL) that we

predict promotes maltose uptake via LamB. Efficient uptake of maltose/maltodextrin not only

requires the inner membrane transporters encoded by the *malEFG* operon, but also the high-

affinity maltose transporter LamB (Figure 9A), which cooperates with the inner membrane

proteins to import these carbon sources. *lamB* is significantly upregulated when cells start to

utilize alternative carbon sources. A major advantage of a coherent FFL is that it can

accelerate or delay responses to stimuli, such as changes in carbon source availability. In this

FFL, the key activator is the MalT transcription factor, which induces the expression of both

malEFG and *lamB* when cells start to consume maltose (Figure 9B). Expression of new OMPs,

however, needs to be carefully coordinated as any changes in the protein composition of the

OM, such as increased levels of LamB (or accumulation of misfolded LamB in the periplasm),

can lead to induction of the σ^E envelope stress response caused by the reorganisation of the

membrane when maltodextrins are utilised (Figure 9A)(Kenyon et al., 2005). We propose that

malEFG indirectly promotes LamB accumulation via the MdoR sRNA that (a) reduces the

levels of general porins (OmpC and OmpA) and (b), likely indirectly, dampens the σ^E stress

response as a result of OMP level reduction. Dampening σ^E expression important during

maltose uptake as it induces the expression of MicA, which directly downregulates *lamB*

translation (Bossi and Figueroa-Bossi, 2007). When ectopically expressed at high levels,

MdoR reduces mRNA levels of *ompC* and *ompA*, which may free up the resources enabling

efficient accumulation of LamB in the outer membrane. At physiological levels MdoR

significantly reduces MicA expression and we observed a ~20% increase in *lamB* mRNA

levels. *OmpC* mRNA levels were not substantially affected. While these changes appear

modest, it is important to take into consideration the very high abundance and intrinsic stability

of *lamB* and *ompC* mRNAs (minute-long half-lives). It is therefore expected that even a mild

reduction in their mRNA levels can profoundly relieve the pressure on the OMP translation

and assembly pathways (Guo et al., 2014). The net outcome of *mal* regulon transcription,

MdoR biogenesis and regulatory activity, is increased expression of high-affinity components

of maltose-specific transport (MalE and LamB).

MdoR is reminiscent of the *Vibrio cholerae* MicX sRNA that also mapped to the 3'

region of *malG* and regulates levels of outer membrane proteins (Davis and Waldor, 2007).

However, the MicX targets are different from MdoR. Furthermore, unlike MdoR, MicX is

produced from an independent promoter: Conditions that change the expression of the

upstream Mal operon in *V. cholerae* do not affect accumulation of MicX (Davis and Waldor,
2007), whereas the levels of MdoR strongly correlate with *malEFG* mRNA levels.
Nevertheless, both studies demonstrate that an sRNA resides in the 3'UTR of the *malG*
transcript and pinpoint a mechanism where the expression of multiple porins is regulated by
sRNAs encoded within transporter mRNAs.
797

CLASH reveals large sRNA-sRNA interaction networks.

Our analyses unearthed an unexpectedly large number (>100) of sRNA-sRNA
interactions. The majority of these interactions involved known sRNA seed sequences,
suggesting that these could represent *bona fide* interactions that prevent sRNAs from base-
pairing with their targets or result in the degradation of the sRNAs. About 40 sRNA-sRNA
interactions involving sRNAs from the core genome were detected in the RNase E CLASH
dataset, about a quarter of which were also detected in our Hfq data. The relatively low overlap
between the datasets is not surprising, given the differences in the growth conditions
(virulence inducing conditions for enterohaemorrhagic *E. coli* vs growth transitions for
commensal *E. coli* MG1655). Moreover, many of the sRNA-sRNA interactions recovered in
association with RNase E are likely duplexes in the process of being degraded. Many of the
sRNA-sRNA interactions unique to our Hfq data may represent interactions between anti-
sRNAs and seed sequences of sRNAs which may not necessarily involve recruitment/activity
of RNase E.
The chimeras containing CyaR fragments were of particular interest as CyaR is
preferentially expressed during the transition from late exponential to stationary phase (De
Lay and Gottesman, 2009; Wassarman et al., 2001) and may therefore play an important role
in adaptation to nutrient availability. We could show that overexpression of ArcZ, which base-
pairs with CyaR in our CLASH data, significantly reduced CyaR levels (Figure 9B). Using seed
mutants and compensatory mutations, we confirmed direct interactions between ArcZ and
CyaR and our data suggests that this interaction specifically takes place during the transition
phase. Interestingly in *Salmonella*, overexpression of ArcZ showed a dramatic reduction in
CyaR bound to Hfq and upregulation of CyaR targets, such as *nadE* (Papenfort et al., 2009),
suggesting that this activity is conserved between these two Gram-negative bacteria. A similar
type of asymmetric regulation has also been elegantly demonstrated for the Qrr3 sRNA of
Vibrio cholera (Feng et al., 2015). The fate of these sRNA-sRNA duplexes may depend on the
position of the interaction; It was shown that if sRNA-target RNA base-pair within a stabilizing
5' stem, the sRNA will be preferentially degraded (Feng et al., 2015). Consistent with this,
folding of the chimeric reads suggests that ArcZ preferentially base-pairs with CyaR at the 5'
end (Figure 4), which may alter secondary structures that normally help to stabilize the sRNA.

The biological role of ArcZ targeting CyaR is unclear, however, a possible function could be

to reduce noise in CyaR expression by preventing CyaR levels from overshooting during the

transition phase.

ArcZ and CyaR target mRNAs are associated with many different processes. Thus,

these interactions are expected to connect multiple pathways. For example, ArcZ regulation

of CyaR may connect adaptation to stationary phase/biofilm development (De Lay and

Gottesman, 2009; Monteiro et al., 2012) to quorum sensing and cellular adherence (De Lay

and Gottesman, 2009). CyaR expression is controlled by the global regulator Crp. Most of the

genes controlled by Crp are involved in transport and/or catabolism of amino acids or sugar.

Interestingly, ArcZ downregulates the *sdaCB* dicistron which encodes for proteins involved in

serine uptake and metabolism (Papenfort et al., 2009). This operon has been shown to be

regulated by Crp as well, suggesting that ArcZ can counteract the activity of Crp.

Materials and Methods

Bacterial strains and culture conditions

An overview of the bacterial strains used in this study is provided in Supplementary

Table 5. The *E. coli* MG1655, TOP10 or TOP10F' strains served as parental strains. The *E.*

coli K12 strain used for CLASH experiments, MG1655 *hfq*::HTF was previously reported (Tree

et al., 2014). Cells were grown in Lysogeny Broth (LB) or minimal medium with supplements

(1xM9 salts, 2 mM MgSO$_4$,0.1 mM CaCl$_2$, 0.03 mM thiamine, 0.2% carbon-source) at 37°C

under aerobic conditions with shaking at 200 rpm. The media were supplemented with

antibiotics where required at the following concentrations: ampicillin (Sigma, UK, A9518) - 100

µg/ml, chloramphenicol (Corning, US, C239RI) - 25 µg/ml, kanamycin (Gibco, US, 11815-024)

- 50 µg/ml. Where indicated, 0.2% glucose or maltose were used. For induction of sRNA

expression from plasmids, 1 mM IPTG, 200 nM anhydrotetracycline hydrochloride (Sigma,

1035708-25MG) or 0.2% L-arabinose (Sigma, A3256) were used.

Construction of sRNA expression plasmids

For the pulse-overexpression constructs, the sRNA gene of interest was cloned at the

transcriptional +1 site under P*ara* control by amplifying the pBAD+1 plasmid (Supplementary

Table 5) by inverse PCR using Q5 DNA Polymerase (NEB). The pBAD+1 template is derived

from pBAD*myc*HisA (Tree et al., 2014). The sRNA genes and seed mutants (SM) were

synthesized as ultramers (IDT; Supplementary Table 5), which served as the forward primers,

as described (Tree et al., 2014). The reverse primer (oligo pBAD+1_5P_rev) bears a

monophosphorylated 5'-end to allow blunt-end self-ligation. The PCR reaction was digested

with 10U DpnI (NEB) for 1h at 37°C and purified by ethanol precipitation. The sRNA-pBAD

linear PCR product was circularized by self-ligation, performed as above. Ligations were
transformed in DH5α competent cells. Positive transformants were screened by sequencing.
The control plasmid pBAD+1 was constructed similarly by self-ligation of the PCR product
generated from oligonucleotides pBAD+1_XbaI_fwd and pBAD+1_5P_rev. Small RNA
overexpression constructs derived from the pZA21MCS and pZE12*luc* (Expressys) were
generated identically, using the indicated ultramers in Supplementary Table 5 as forward
primers, and oligos pZA21MCS_5P_rev and pZE12_5P_rev as reverse primers, respectively,
and transformed in *E. coli* TOP10F'.

Construction of mRNA-superfolder GFP fusions

Supplementary Table 5 lists all the plasmids, gene fragments and primers used for
cloning procedures in this work. To construct constitutively expressed, in-frame mRNA-*sfGFP*
fusions for the fluorescence reporter studies, the 5'UTR, start codon and first ~5 codons of
target genes were cloned under the control of PLtetO-1 promoter in a pXG10-SF backbone
as previously described (Corcoran et al., 2012; Urban and Vogel, 2007). Derivatives of the
target–GFP fusion plasmids harbouring seed mutations (SM) were generated using synthetic
mutated gene-fragments (IDT, Belgium, Supplementary Table 5). To prepare the inserts, the
target region of mRNA of interest was either amplified by PCR from *E. coli* genomic DNA or
synthesized as g-blocks (IDT, Belgium) and cloned using NheI and NsiI restriction sites.
Transformants were screened by restriction digest analysis and verified by Sanger sequencing
(Edinburgh Genomics, Edinburgh, UK).

Hfq UV Cross-linking, Ligation and Analysis of Hybrids (Hfq-CLASH)

CLASH was performed essentially as described(Waters et al., 2016), with a number
of modifications including changes in incubation steps, cDNA library preparation, reaction
volumes and UV cross-linking. *E. coli* expressing the chromosomal Hfq-HTF were grown
overnight in LB at 37°C with shaking (200 rpm), diluted to starter OD_{600} 0.05 in fresh LB, and
re-grown with shaking at 37°C in 750 ml LB. A volume of culture equivalent to 80 OD_{600} per
ml was removed at the following cell-densities (OD_{600}): 0.4, 0.8, 1.2, 1.8, 2.4, 3.0 and 4.0, and
immediately subjected to UV (254 nm) irradiation for 22 seconds (~500 mJ/cm2) in the Vari-
X-linker (van Nues et al., 2017) (https://www.vari-x-link.com). Cells were harvested using a
rapid filtration device (van Nues et al., 2017) (https://www.vari-x-link.com) onto 0.45 μM
nitrocellulose filters (Sigma, UK, HAWP14250) and flash-frozen on the membrane in liquid
nitrogen. Membranes were washed with ~15 ml ice-cold phosphate-buffered saline (PBS),
and cells were harvested by centrifugation. Cell pellets were lysed by bead-beating in 1
volume per weight TN150 buffer (50mM Tris pH 8.0, 150 mM NaCl, 0.1% NP-40, 5 mM β-
mercaptoethanol) in the presence of protease inhibitors (Roche, A32965), and 3 volumes 0.1

mm Zirconia beads (Thistle Scientific, 11079101z), by performing 5 cycles of 1 minute
vortexing followed by 1-minute incubation on ice. One additional volume of TN150 buffer was
added. To reduce the viscosity of the lysate and remove contaminating DNA the lysate was
incubated with RQ1 DNase I (10U/ml Promega, M6101) for 30 minutes on ice. Two-additional
volumes of TN150 were added and mixed with the lysates by vortexing. The lysates were
centrifuged for 20 minutes at 4000 rpm at 4°C and subsequently clarified by a second
centrifugation step at 13.4 krpm, for 20 min at 4°C. Purification of the UV cross-linked Hfq-
HTF-RNA complexes and cDNA library preparation was performed as described (Granneman
et al., 2009). Cell lysates were incubated with 50 µl of pre-equilibrated M2 anti-FLAG beads
(Sigma, M8823-5ML) for 1-2 hours at 4°C. The anti-FLAG beads were washed three times 10
minutes with 2 ml TN1000 (50 mM Tris pH 7.5, 0.1% NP-40, 1M NaCl) and three times 10
minutes with TN150 without protease inhibitors (50 mM Tris pH 7.5, 0.1% NP-40, 150mM
NaCl). For TEV cleavage, the beads were resuspended in 250 µl of TN150 buffer (without
protease inhibitors) and incubated with home-made GST-TEV protease at room temperature
for 1.5 hours. The TEV eluates were then incubated with a fresh 1:100 dilution preparation of
RNaceIt (RNase A and T1 mixture; Agilent, 400720) for exactly 5 minutes at 37ºC, after which
they were mixed with 0.4g GuHCl (6M, Sigma, G3272-100G), NaCl (300mM), and Imidazole
(10mM, I202-25G). Note this needs to be carefully optimized to obtain high-quality cDNA
libraries. The samples were then transferred to 50 µl Nickel-NTA agarose beads (Qiagen,
30210), equilibrated with wash buffer 1 (6 M GuHCl, 0.1% NP-40, 300 mM NaCl, 50 mM Tris
pH 7.8, 10 mM Imidazole, 5 mM beta-mercaptoethanol). Binding was performed at 4ºC
overnight with rotation. The following day, the beads were transferred to Pierce SnapCap spin
columns (Thermo Fisher, 69725), washed 3 times with wash buffer 1 and 3 times with 1xPNK
buffer (10 mM MgCl$_2$, 50mM Tris pH 7.8, 0.1% NP-40, 5 mM beta-mercaptoethanol). The
washes were followed by on-column TSAP incubation (Thermosensitive alkaline
phosphatase, Promega, M9910) treatment for 1h at 37°C with 8 U of phosphatase in 60 µl of
1xPNK, in the presence of 80U RNasin (Promega, N2115). The beads were washed once
with 500 µl wash buffer 1 and three times with 500 µl 1xPNK buffer. To add 3'-linkers (App-
PE – Supplementary Table 5), the Nickel-NTA beads were incubated in 80 µl 3'-linker ligation
mix with (1 X PNK buffer, 1 µM 3'-adapter, 10% PEG8000, 30U Truncated T4 RNA ligase 2
K227Q (NEB, M0351L), 60U RNasin). The samples were incubated for 4 hours at 25°C. The
5'-ends of bound RNAs were radiolabelled with 30U T4 PNK (NEB, M0201L) and 3µl ^{32}P-
γATP (1.1µCi; Perkin Elmer, NEG502Z-500) in 1xPNK buffer for 40 min at 37°C, after which
ATP (Roche, 11140965001) was added to a final concentration of 1mM, and the incubation
prolonged for another 20 min to complete 5'-end phosphorylation. The resin was washed three
times with 500 µl wash buffer 1 and three times with equal volume of 1xPNK buffer. For on-

bead 5'-linker ligation, the beads were incubated 16h at 16°C in 1xPNK buffer with 40U T4

RNA ligase I (NEB, M0204L), and 1 µl 100 µM L5 adapter (Supplementary Table 5), in the

presence of 1mM ATP and 60U RNasin. The Nickel-NTA beads were washed three times with

wash buffer 1 and three times with buffer 2 (50 mM Tris–HCl pH 7.8, 50 mM NaCl, 10 mM

imidazole, 0.1% NP-40, 5 mM β-mercaptoethanol). The protein-RNA complexes were eluted

in two steps in new tubes with 200 µl of elution buffer (wash buffer 2 with 250 mM imidazole).

The protein-RNA complexes were precipitated on ice by adding TCA (T0699-100ML) to a final

concentration of 20%, followed by a 20-minute centrifugation at 4°C at 13.4 krpm. Pellets were

washed with 800 µl acetone, and air dried for a few minutes in the hood. The protein pellet

was resuspended and incubated at 65°C in 20 µl 1x NuPage loading buffer (Thermo Scientific,

NP0007), resolved on 4–12% NuPAGE gels (Thermo Scientific, NP0323PK2) and visualised

by autoradiography. The cross-linked proteins-RNA were cut directly from the gel and

incubated with 160 µg of Proteinase K (Roche, 3115801001) in 600 µl wash buffer 2

supplemented with 1% SDS and 5 mM EDTA at 55°C for 2-3 hours with mixing. The RNA was

subsequently extracted by phenol-chloroform extraction and ethanol precipitated. The RNA

pellet was directly resuspended in RT buffer and was transcribed in a single reaction with the

SuperScript IV system (Invitrogen, 18090010) according to manufacturer's instructions using

the PE_reverse oligo as primer. The cDNA was purified with the DNA Clean and Concentrator

5 kit (Zymo Research) and eluted in 11 µl DEPC water. Half of the cDNA (5 µl) was amplified

by PCR using Pfu Polymerase (Promega, M7745) with the cycling conditions (95°C for 2 min;

20-24 cycles: 95°C for 20s, 52°C for 30s and 72°C for 1 min; final extension of 72°C for 5 min).

The PCR primers are listed in Supplementary Table 5. PCR products were treated with 40U

Exonuclease 1 (NEB, M0293L) for 1 h at 37°C to remove free oligonucleotide and purified by

ethanol precipitation/ or the DNA Clean and Concentrator 5 kit (Zymo Research, D4003T).

Libraries were resolved on a 2% MetaPhor agarose (Lonza, LZ50181) gel and 175-300bp

fragments were gel-extracted with the MinElute kit (Qiagen, 28004) according to

manufacturer's instructions. All libraries were quantified on a 2100 Bionalyzer using the High-

Sensitivity DNA assay and a Qubit 4 (Thermo Scientific, Q33226). Individual libraries were

pooled based on concentration and barcode sequence identity. Paired-end sequencing (75

bp) was performed by Edinburgh Genomics on an Illumina HiSeq 4000 platform.

RNA-seq

E. coli MG1655 was cultured, UV-irradiated and harvested as described for the CLASH

procedure. Total RNA was extracted using the Guanidium thiocyanate phenol method. RNA

integrity was assessed with the Prokaryote Total RNA Nano assay on a 2100 Bioanalyzer

(Agilent, G2939BA). Sequencing libraries from two biological replicates were prepared by

NovoGene using the TruSeq library preparation protocol and 150bp paired-end sequencing
was performed on an Illumina NovaSeq 6000 system. This yielded ~7-8 million paired-end
reads per sample. For the overexpression analysis of MdoR, we generated RNA-seq libraries
using an in-house protocol. Genomic DNA was removed by incubating 10 µg of total RNA with
2U Turbo DNase (Thermo Scientific, AM2238) in a 50 µl final volume for 30 minutes at 37°C
in the presence of 10 U SuperaseIn RNase Inhibitor (Thermo Scientific, AM2694). RNA was
subsequently phenol-chloroform extracted and purified by ethanol-precipitation. Ribosomal
RNA was removed with the Ribo-Zero rRNA Removal Kit (Gram-Negative Bacteria; Illumina,
MRZGN126) according to the manufacturer's instructions. Successful rRNA depletion was
verified on the Agilent 2100 Bioanalyzer. The RNA was fragmented for 5 min at 95°C in the
presence of Superscript III buffer (Invitrogen) followed by a five-minute incubation on ice.
Reverse-transcription (RT) was performed with Superscript III (Thermo Scientific, 18080044)
in 20 µl reactions according to manufacturer's procedures using 250 ng of ribosomal RNA
depleted RNA and 2.5 µM random hexamers (PE_solexa_hexamer, oligo 73, Supplementary
Table 5). The RNA and free primers were degraded using 20U of Exonuclease I (NEB,
M0293L) and 50U RNaseIf (NEB, M0243S) and the cDNA was purified with the DNA Clean &
Concentrator 5 kit (Zymo Research). Ligation of the 5' adapter (P5_phospho_adapter, oligo
39) to the cDNA was performed using CircLigase II (Lucigen, CL9021K) for 6 hours at 60°C,
followed by a 10-minute inactivation at 80°C. The cDNA was purified with the DNA Clean &
Concentrator 5 kit. Half of the cDNA library was PCR amplified using Pfu polymerase
(Promega, M7745) using the P5 forward PCR oligonucleotide and barcoded BC reverse
oligonucleotides (200 nM; Supplementary Table 5; 95°C for 2 min, 95°C for 20s, 52°C for 30s
and 72°C for 1 min, and a final extension of 72°C for 5 min. 20 cycles of amplification). The
PCR products were treated with Exonuclease 1 (NEB, M0293L) for 1 h at 37°C and purified
by ethanol precipitation. Libraries were resolved on a 2% MetaPhor agarose gel 200-500 bp
fragments were gel-extracted using the MinElute kit. All libraries were quantified on a 2100
Bionalyzer using the High-Sensitivity DNA assay (Agilent, 5067-4627). Individual libraries
were pooled in equimolar amounts. Paired-end sequencing (75 bp) was performed by
Edinburgh Genomics on the Illumina HiSeq 4000 platform.
1003
Small RNA overexpression studies
Individual TOP10F' clones carrying pZA21 and pZE12-derived sRNA constructs and
control plasmids combinations (Supplementary Table 5) were cultured to OD_{600} 0.1 and
expression of sRNAs was induced with IPTG and anhydrotetracycline hydrochloride (Sigma,
I6758-1G and 1035708-25MG) for one hour. Cells were collected by centrifugation for 30
seconds at 14000 rpm, flash-frozen in liquid nitrogen and total RNA was isolated as above.

Genomic DNA was digested with Turbo DNase (Thermo Scientific, AM2238) , then the RNA

was purified with RNAClean XP beads (Beckman Coulter, A63987). Gene expression was

quantified by RT-qPCR (see below) using 10 ng total RNA as template, and expressed as fold

change relative to the reference sample containing pJV300 (Sittka et al., 2007) or pZA21.

For pulse-overexpression studies overnight MG1655 cultures containing pBAD::sRNA

and empty pBAD+1 control plasmids were inoculated in fresh LB-ampicillin medium at a

starting OD_{600} of 0.05 and grown aerobically at 37°C to OD_{600} 0.4. Pre-induction (0 min) and

post-induction samples were harvested. For induction, cultures were supplemented with L-

arabinose (Sigma, A3256) and rapidly collected by filtration and flash-frozen in liquid nitrogen

at the indicated time-points. RNA was extracted from three biological replicate time-series,

followed by RNA-seq library preparation, next generation sequencing and DESeq2 analysis

of differentially expressed genes.

1022

GFP reporter system to quantify sRNA effect on target expression

A two-plasmid system was used to express each sRNA, and mRNA-*sfGFP* fusions

(Corcoran et al., 2012; Urban and Vogel, 2007) with modifications. The sRNA and sfGFP-

fusion plasmids were co-transformed in *E. coli* TOP10 cells by electroporation and cells were

maintained on dual selection with ampicillin and chloramphenicol. In TOP10 cells, the mRNA-

sfGFP constructs are constitutively expressed, whereas sRNA expression requires L-

arabinose induction. The expression of sfGFP-fused targets in the presence or absence of

sRNAs was quantified at the protein level, by plate reader experiments and at the RNA level,

by RT-qPCR.

For the plate reader experiments, a single colony of bacterial strain harbouring a

sRNA-target-sfGFP combination was inoculated in a 96-well Flat Bottom Transparent

Polystyrene plate with lid (Thermo Scientific, 152038) and cultured overnight at 37°C in 100

µl LB supplemented with antibiotics and L-arabinose (Sigma, A3256) to induce expression of

sRNAs. Next day, each overnight inoculum was diluted 1:100 by serial dilution, in triplicate, in

LB with freshly prepared L-arabinose to a final volume of 100 µl. Cultures were grown in a 96-

well plate in an Infinite 200 Pro plate reader (Tecan) controlled by i-control software (Tecan)

for 192 cycles at 37°C with 1 min orbital shaking (4 mm amplitude) every 5th minute. To

monitor optical density over time, the following parameters were used: wavelength 600 nm,

bandwidth 9 nm. Fluorescence was monitored with excitation wavelength 480 nm, bandwidth

9 nm and emission wavelength 520 nm, bandwidth 20 nm. Measurements were recorded at

5-minute intervals, by top reading. Raw data was processed following guidance from previous

reports (Urban and Vogel, 2007). First, the range of linearity of increase of fluorescence with

OD_{600} was identified for all individual triplicates. Only the linearity range common to all

triplicates was considered for further analysis. For each set of triplicates, the mean
fluorescence was calculated at each OD_{600}. To correct for background and cell
autofluorescence, the mean fluorescence of a strain with plasmid pXG-0 was subtracted from
all strains with GFP plasmids at the equivalent OD_{600}. Ultimately, a curve was generated for
each sample, plotting the background-corrected fluorescence (GFP) versus OD_{600}. The
experiments were performed for three biological replicates, and mean values and standard
error of the means calculated for each strain.

1053

RT-qPCR
Total RNA (12.5µg) was treated with 2U of Turbo DNase (Thermo Scientific, AM2238)
for 1 hour at 37°C in a 10 µl reaction in the presence of 2U SuperaseIn RNase inhibitor
(Thermo Scientific, AM2694). The DNase was inactivated by 10 minutes incubation at 75°C.
Reverse transcription (RT) was performed in a single reaction for all target genes of interest
using a mix of gene-specific RT primers at 3.5 µM concentration each. After addition of 2.5 µl
RT primer mix, the RNA and primers were denatured at 70°C for 3 min, then snap chilled and
incubated on ice for 5 min. RT was performed for 1 hour at 55°C with SuperScript III (Thermo
Scientific, 18080051) using 5 µl of RNA-RT primers mix in 10 µl final volume (100 U
Superscript III, 2.5 mM DTT, 1xFS Buffer, 0.75 mM dNTPs) in the presence of 1U RNasin
(Promega, N2115). RT was followed by treatment with 5U RNase H for 30 min at 37°C to
remove the RNA from the RNA-cDNA duplexes. The cDNA was diluted 10-fold with DEPC
water. Quantitative PCR was performed on 50ng of DNAse I-treated total RNA using the
Brilliant III UltraFast SYBR Green QPCR Master Mix (Agilent, #600883) and the Luna
Universal One-Step RT-qPCR Kit (NEB, E3005E) according to manufacturer's instructions.
The qPCRs were run on a LightCycler 480 (Roche), and the specificity of the product was
assessed by generating melting curves, as follows: 65°C-60s, 95°C (0.11 ramp rate with 5
acquisitions per °C, continuous). The data analyses were performed with the IDEAS2.0
software, at default settings: Absolute Quantification/Fit Points for Cp determination and Melt
Curve Genotyping.
The qPCR efficiency of primer pairs was assessed by performing standard curves by serial
dilution of template RNA or genomic DNA. Negative controls such as -RT or no template
control were used throughout, and the qPCR for all samples was performed in technical
triplicate. Outliers from the samples with technical triplicate standard deviations of Cp > 0.3
were discarded from the analyses. To calculate the fold-change relative to the control, the 2^{-ddCp}
method was employed, using *recA* or 5S rRNA (*rrfD*) as the reference genes where
indicated. Experiments were performed for minimum two biological replicates, and the mean
fold-change and standard error of the mean were computed. Unless otherwise stated,

significance of the fold-change difference compared to the reference sample control (for which
fold-change =1) was tested with a one-sample t-test.
1084

Northern Blot analysis

Total RNA was extracted from cell lysates by GTC-Phenol extraction. For large RNA
fragments, 10 µg of total RNA was resolved on a 1.25% BPTE- gel (pH 7) and transferred to
a nylon membrane (HyBond N+, GEHealthcare, RPN1210B) by capillarity. For short RNA
fragments, 10 µg total RNA was separated on an 8% polyacrylamide TBE-Urea gel and
transferred to a nitrocellulose membrane by electroblotting for four hours at 50 V. Membranes
were pre-hybridised in 10 ml of UltraHyb Oligo Hyb (Thermo Scientific, AM8663) for one hour
and probed with ^{32}P-labeled DNA oligo at 42°C for 12-18 hours in a hybridization oven. The
sequences of the probes used for Northern blot detection are detailed in Supplementary Table
5. Membranes were washed twice with 2xSSC + 0.5% SDS solution for 10 minutes and
visualized using a Phosphor imaging screen and FujiFilm FLA-5100 Scanner (IP-S mode).
For detection of highly abundant species (5S rRNA) autoradiography was used for exposure.
1097

Western blot analyses

A fraction of the *E. coli* MG1655 Hfq::*htf* lysates used for the RNA-seq experiments
using strains cultured, cross-linked, harvested and lysed in identical conditions as the CLASH
experiments containing 40 µg protein was run on PAGE gels and transferred to a nitrocellulose
membrane. The membranes were blocked for one hour in blocking solution (5% non- fat milk
in PBST (1X phosphate saline buffer, 0.1% Tween-20). To detect Hfq-HTF protein, the
membrane was probed overnight at 4°C with the Rabbit anti-TAP polyclonal primary antibody
(Thermo Fisher, 1:5000 dilution in blocking solution), which recognizes an epitope at the
region between the TEV-cleavage site and His6. For the loading control we used a rabbit
polyclonal to GroEL primary antibody (Abcam, 1:150000 dilution, ab82592), for 2 hours at
room temperature. After 3x10 min PBST washes, the membranes were blotted for one hour
with a Goat anti-rabbit IgG H&L (IRDye 800) secondary antibody (Abcam, ab216773, 1:10000
in blocking solution) at room temperature. Finally, after three 10-minute PBST washes, the
blot was rinsed in PBS, and the proteins were visualised with a LI-COR (Odyssey CLx) using
the 800 nm channel and scan intensity 4. Image acquisition and quantifications were
performed with the Image Studio Software.
1114

Primer extension analysis

One microgram total RNA was reverse-transcribed using SuperScript III reverse
transcriptase (Thermo Scientific, 18080051) using ^{32}P-radiolabelled oligonucleotides as

primers (Supplementary Table 5). Primers were added to the RNA and annealing was
performed by heating the samples at 85°C for three minutes and then snap chilling them on
ice. The RT was performed for one hour at 45°C, followed by Exonuclease I and RNaseIf
(NEB M0293L and M0243S) (0.5 μl each) treatment for 30 minutes at 37°C. Reactions were
stopped by mixing with an equal volume of 2XRNA loading dye (NEB, B0363S), 2 minutes
incubation at 95°C and snap chilled. The sequencing ladders were prepared with Sequenase
v2.0 (Thermo Scientific, 70775Y200UN) according to specified instructions. Samples were
resolved on 6% PAA/8M TBE-urea gels and visualized using the FujiFilm FLA5100 scanner.
1126

Construction of the MdoR seed-mutant strain

To mutate the chromosomal copy of MdoR, we used the λRed system (Datsenko and
Wanner, 2000). We amplified the integration cassette from plasmid pKD4 with ultramers 895
and 896, containing homology regions to the coding sequence of *malG*, the desired MdoR
sequence and to the region immediately downstream of the Rho-independent terminator,
respectively. With this design, the scar after removal of the *Kan*[r] cassette was expected at a
site outside the MdoR/*malG* sequence. The PCR product was electroporated in *E. coli*
MG1566 strains carrying the pKD46 plasmid from which λRed recombinase was induced with
10 mM L-arabinose. Correct replacement of the MdoR seed sequence was screened by
colony PCR using primer pairs: 725 & 909 and 726 & 910 (Supplementary Table 5). The
antibiotic resistance cassette was removed from substitution mutants by FLP-recombinase
expressed constitutively from pE-FLP (St-Pierre et al., 2013). Successful allele replacement
was confirmed by Sanger sequencing.
1140

Polysome profiling analyses

Wild-type *E. coli* MG1655 containing empty pBAD plasmid and an isogenic strain
containing pBAD::MdoR were grown in LB until OD_{600} 0.4, then treated for 15 minutes with L-
arabinose (Sigma, A3256) to induce overexpression of MdoR, and cycloheximide (Sigma,
C4859-1ML) at a final concentration of 100 μg/ml for 3 minutes. 200 ml of cells were harvested
by rapid filtration and flash frozen. The cells were washed in ice-cold PBS supplemented with
100μg/ml cycloheximide.
Polysomal profiling was performed according to previously described protocols
(Bernabò et al., 2017; Lunelli et al., 2016) with minor changes in the lysis buffer (10 mM NaCl,
10 mM $MgCl_2$, 10 mM Tris-HCl (pH 7.5), 100 μg/ml cycloheximide, 1% (w/v) Na-Deoxycholate,
1U RQ DNAse I (Promega, M6101), 0.6U/mL RiboLock (Thermo Scientific, EO0381) in DEPC
water). Lysates were kept on ice for 30 min, centrifuged 3X at 15000 g for 10 min. The
supernatants were loaded on a linear 10%–30% [w/v] sucrose gradient and centrifuged for 4

hours using a SW41 rotor at 40000 rpm in a Beckman Optima XPN-100 Ultracentrifuge.
Fractions of 1 mL in volume were collected monitoring the absorbance at 254 nm with the UA-
6 UV/VIS detector (Teledyne Isco). Fractions from the entire gradient (total RNA) and from the
fractions corresponding to ribosomes (70S) and polysomes (polysomal RNA) were pooled
and RNA was purified by acid phenol–chloroform extraction according to (Tebaldi et al., 2012).
1159

Terminator™ 5′-PhosphateDependent Exonuclease treatment

Ten micrograms of total RNA extracted from cell-samples at OD_{600} 1.2 and 1.8 were
treated with 5'-Terminator Dependent Exonuclease (Lucigen, TER51020) as per manufacturer
instructions using Buffer A. The reaction was terminated by phenol extraction and ethanol
precipitation, and the RNA was loaded on 8% polyacrylamide-urea gels and transferred to
nylon membranes that were probed for MdoR, CpxQ, RybB and 5S rRNA (Supplementary
Table 5).
1167

Seed mutant studies

1169 Wild-type MG1655 and seed mutant strains were grown overnight in minimal medium
1170 with glucose. Next day, each starter culture was split and inoculated at OD_{600} 0.05 in fresh M9
1171 medium with glucose or maltose as the sole carbon source. Growth was monitored and cells
1172 were harvested at OD_{600} 0.5. Total RNA was extracted, and gene expression was quantified
1173 by RT-qPCR or Northern Blot.
1174

Computational analysis

Pre-processing of the raw sequencing data.

Raw sequencing reads in fastq files were processed using a pipeline developed by
Sander Granneman, which uses tools from the pyCRAC package (Webb et al., 2014). The
entire pipeline is available at https://bitbucket.org/sgrann/). The CRAC_pipeline_PE.py
pipeline first demultiplexes the data using pyBarcodeFilter.py and the in-read barcode
sequences found in the L5 5' adapters. Flexbar then trims the reads to remove 3'-adapter
sequences and poor-quality nucleotides (Phred score <23). Using the random nucleotide
information present in the L5 5'-adaptor sequences, the reads are then collapsed to remove
potential PCR duplicates. The reads were then mapped to the *E. coli* MG1655 genome using
Novoalign (www.novocraft.com). To determine to which genes the reads mapped to, we
generated an annotation file in the Gene Transfer Format (GTF). This file contains the start
and end positions of each gene on the chromosome as well as what genomic features (i.e.
sRNA, protein- coding, tRNA) it belongs to. To generate this file, we used the Rockhopper
software (Tjaden, 2015) on *E. coli* rRNA-depleted total RNA-seq data (generated by Christel
Sirocchi), a minimal GTF file obtained from ENSEMBL (without UTR information). The

resulting GTF file contained information not only on the coding sequences, but also complete
5' and 3' UTR coordinates. We then used pyReadCounters.py with Novoalign output files as
input and the GTF annotation file to count the total number of unique cDNAs that mapped to
each gene.
1195
Normalization steps.
To normalise the read count data generated with pyReadCounters.py and to correct
for differences in library depth between time-points, we calculated Transcripts Per Million
reads (TPM) for each gene. Briefly, for each time-point the raw counts for each gene was first
divided by the gene length and then divided by the sum of all the values for the genes in that
time-point to normalize for differences in library depth. The TPM values for each OD_{600} studied
were divided by the TPM values of the first sample (OD_{600} 0.4) and were then log_2-normalized.
The log_2-normalized fold-changes were used to compare RNA-seq and Hfq-cross-linking
profiles among samples, and to perform k-means clustering with the python sklearn.
cluster.KMeans class.
1206
Hfq-binding coverage plots.
For the analysis of the Hfq binding sites the pyCRAC package (Webb et al., 2014) was
used (versions. 1.3.2-1.4.4). The pyBinCollector tool was used to generate Hfq cross-linking
distribution plots over genomic features. First, PyCalculateFDRs.py was used to identify the
significantly enriched Hfq-binding peaks (minimum 10 reads, minimum 20 nucleotide
intervals). Next, pyBinCollector was used to normalize gene lengths by dividing their
sequences into 100 bins and calculate nucleotide densities for each bin. To generate the
distribution profile for all genes individually, we normalized the total number of read clusters
(assemblies of overlapping cDNA sequences) covering each nucleotide position by the total
number of clusters that cover the gene. Motif searches were performed with pyMotif.py using
the significantly enriched Hfq-binding peaks (FDR intervals). The 4-8 nucleotide k-mers with
Z-scores above the indicated threshold were used for making the motif logo with the k-mer
probability logo tool (Wu and Bartel, 2017) with the -ranked option (http://kplogo.wi.mit.edu/).
1220
Analysis of chimeric reads.
Chimeric reads were identified using the hyb package using default settings (Travis et
al., 2013) and further analysed using the pyCRAC package (Webb et al., 2014). To apply this
single-end specific pipeline to our paired-end sequencing data, we joined forward and reverse
reads using FLASH (Magoč and Salzberg, 2011), which merges overlapping paired reads into
a single read. These, as well as any remaining single reads, were then analysed using hyb.
The -anti option for the hyb pipeline was used to be able to use a genomic *E. coli* hyb database,

rather than a transcript database. Uniquely annotated hybrids (.ua.hyb) were used in
subsequent analyses. To visualise the hybrids in the genome browser, the .ua.hyb output files
were converted to the GTF format. To generate distribution plots for the genes to which the
chimeric reads mapped, the parts of the chimeras were clustered with pyClusterReads.py and
BEDtools (Quinlan and Hall, 2010) (intersectBed) was used to remove clusters that map to
multiple regions. To produce the coverage plots with pyBinCollector, each cluster was counted
only once, and the number of reads belonging to each cluster was ignored.

1235

sRNA density plots.

To visualize the nucleotide read density of sRNA-target pairs for a given sRNA, we
first merged the hyb datasets for all OD_{600} and biological replicates and filtered the interactions
that were found statistically significant in the unified dataset. For each sRNA-target pair in the
filtered dataset, the hit counts at each nucleotide position for all chimeras were summed. The
count data was log_2-normalised (actually log_2(Chimera count +1) to avoid NaN for nucleotide
positions with 0 hits when log-transforming the data).

1243

1244 *sRNA-sRNA network visualization.*

1245 Only the sRNA-sRNA chimeric reads representing statistically significant interactions
1246 in the merged CLASH dataset were considered. For each such interaction, chimera counts
1247 corresponding in either orientation were summed, log_2-transformed and visualized with the
1248 igraph Python package.

1249

1250 *Differential expression analyses*.

1251 For the differential expression analyses DESeq2 was used (Love et al., 2014). Three
1252 MdoR pulse-overexpression datasets were compared to three pBAD Control overexpression
1253 datasets. Only differentially expressed genes that had an adjusted p-value of 0.05 or lower
1254 were considered significant.

1255

1256 *Multiple sequence alignments and conservation analyses*.

1257 The homologous sequences of MdoR in other enterobacteria were retrieved by
1258 BLAST. JalView was used for the multiple sequence alignments, using the MAFFT algorithm
1259 (Waterhouse et al., 2009).

1260

1261 **Data and Code availability.**

1262 The next generation sequencing data have been deposited on the NCBI Gene
1263 Expression Omnibus (GEO) with accession number GSE123050. The python pyCRAC(Webb
1264 et al., 2014), kinetic-CRAC and GenomeBrowser software packages used for analysing the

data are available from https://bitbucket.org/sgrann (pyCRAC up to version 1.3.3), https://git.ecdf.ed.ac.uk/sgrannem/ and pypi (https://pypi.org/user/g_ronimo/). The hyb pipeline for identifying chimeric reads is available from https://github.com/gkudla/hyb. The FLASH algorithm for merging paired reads is available from https://github.com/dstreett/FLASH2. Bedgraph and Gene Transfer Format (GTF) generated from the analysis of the Hfq CLASH, RNA-seq and TEX RNA-seq data(Thomason et al., 2015) are available from the Granneman lab DataShare repository (https://datashare.is.ed.ac.uk/handle/10283/2915).

Acknowledgement

We are grateful to Lionello Bossi and Meriem El Karoui for their valuable feedback on the project and fruitful discussions. We thank Christel Sirocchi for help with preparing *E. coli* RNA-seq libraries, Erica de Leau for expert technical assistance and the members of the Granneman lab for critically reading the manuscript. We thank Jean-Marie Pages and Julia Vergalli for help the analyses of LamB levels and for fruitful discussions. This work was supported by grants from the Wellcome Trust (091549 to S.G and 102334 to I.A.I.), the Wellcome Trust Centre for Cell Biology core grant (092076), a Medical Research Council non Clinical Senior Research Fellowship (MR/R008205/1 to S.G.), the Australian National Health and Medical Research Council Project grants (GNT1067241 and GNT1139313 to J.J.T) and the Autonomous Province of Trento (Axonomix to G.V and M.M.). Sequencing of Hfq CLASH data was carried out by Edinburgh Genomics that is partly supported through core grants from NERC (R8/H10/56), MRC (MR/K001744/1) and BBSRC (BB/J004243/1).

Competing interest.

The authors declare no competing financial interest.

Materials & Correspondence

All requests for code, materials and reagents should be sent to Sander Granneman

(sgrannem@ed.ac.uk)

References

Alba BM, Gross CA. 2004. Regulation of the Escherichia coli ??E-dependent envelope stress response. *Mol Microbiol* **52**:613–619. doi:10.1111/j.1365-2958.2003.03982.x

Baev M V., Baev D, Jansco Radek A, Campbell JW. 2006a. Growth of Escherichia coli

MG1655 on LB medium: Monitoring utilization of amino acids, peptides, and
nucleotides with transcriptional microarrays. *Appl Microbiol Biotechnol* **71**:317–322.
doi:10.1007/s00253-005-0310-5

Baev M V., Baev D, Radek AJ, Campbell JW. 2006b. Growth of Escherichia coli MG1655 on
LB medium: Determining metabolic strategy with transcriptional microarrays. *Appl*
Microbiol Biotechnol **71**:323–328. doi:10.1007/s00253-006-0392-8

Balleza E, López-Bojorquez LN, Martínez-Antonio A, Resendis-Antonio O, Lozada-Chávez I,
Balderas-Martínez YI, Encarnación S, Collado-Vides J. 2009. Regulation by
transcription factors in bacteria: Beyond description. *FEMS Microbiol Rev* **33**:133–151.
doi:10.1111/j.1574-6976.2008.00145.x

Bandyra KJ, Said N, Pfeiffer V, Górna MW, Vogel J, Luisi BF. 2012. The Seed Region of a
Small RNA Drives the Controlled Destruction of the Target mRNA by the
Endoribonuclease RNase E. *Mol Cell* **47**:943–953. doi:10.1016/j.molcel.2012.07.015

Beisel CL, Storz G. 2011. The Base-Pairing RNA Spot 42 Participates in a Multioutput
Feedforward Loop to Help Enact Catabolite Repression in Escherichia coli. *Mol Cell*
41:286–297. doi:10.1016/j.molcel.2010.12.027

Bernabò P, Tebaldi T, Groen EJN, Lane FM, Perenthaler E, Mattedi F, Newbery HJ, Zhou H,
Zuccotti P, Potrich V, Shorrock HK, Muntoni F, Quattrone A, Gillingwater TH, Viero G.
2017. In Vivo Translatome Profiling in Spinal Muscular Atrophy Reveals a Role for SMN
Protein in Ribosome Biology. *Cell Rep*. doi:10.1016/j.celrep.2017.10.010

Boos W, Shuman H. 1998. Maltose / Maltodextrin System of Escherichia coli : Transport ,
Metabolism , and Regulation Maltose / Maltodextrin System of Escherichia coli :
Transport , Metabolism , and Regulation. *Microbiol Mol Biol Rev* **62**:204–229.

Bossi L, Figueroa-Bossi N. 2007. A small RNA downregulates LamB maltoporin in
Salmonella. *Mol Microbiol* **65**:799–810. doi:10.1111/j.1365-2958.2007.05829.x

Bossi L, Maloriol D, Figueroa-Bossi N. 2008. Porin biogenesis activates the σE response in
Salmonella hfq mutants. *Biochimie* **90**:1539–1544. doi:10.1016/j.biochi.2008.06.001

Bouvier M, Sharma CM, Mika F, Nierhaus KH, Vogel J. 2008. Small RNA binding to 5'
mRNA coding region inhibits translational initiation. *Mol Cell* **32**:827–837.

Bruce HA, Du D, Matak-Vinkovic D, Bandyra KJ, Broadhurst RW, Martin E, Sobott F,
Shkumatov A V., Luisi BF. 2018. Analysis of the natively unstructured RNA/protein-
recognition core in the Escherichia coli RNA degradosome and its interactions with
regulatory RNA/Hfq complexes. *Nucleic Acids Res*. doi:10.1093/nar/gkx1083

Chao Y, Li L, Girodat D, Forstner KU, Said N, Corcoran C, Smiga M, Papenfort K, Reinhardt
R, Wieden HJ, Luisi BF, Vogel J. 2017. *in vivo* cleavage map illuminates the central
cole of RNase E in coding and non-coding RNA pathways. *Mol Cell* **65**:39–51.
doi:10.1016/j.molcel.2016.11.002

Chao Y, Papenfort K, Reinhardt R, Sharma CM, Vogel J. 2012. An atlas of Hfq-bound
transcripts reveals 3' UTRs as a genomic reservoir of regulatory small RNAs. *EMBO J*
31:4005–4019.

Chao Y, Vogel J. 2016. A 3' UTR-Derived Small RNA Provides the Regulatory Noncoding
Arm of the Inner Membrane Stress Response. *Mol Cell* **61**:352–363.
doi:10.1016/j.molcel.2015.12.023

Chen S, Zhang A, Blyn LB, Storz G. 2004. MicC, a second small-RNA regulator of omp
protein expression in Escherichia coli. *J Bacteriol*. doi:10.1128/JB.186.20.6689-
6697.2004

Corcoran CP, Podkaminski D, Papenfort K, Urban JH, Hinton JCD, Vogel J. 2012.
Superfolder GFP reporters validate diverse new mRNA targets of the classic porin
regulator, MicF RNA. *Mol Microbiol* **84**:428–445.

Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**:6640–5.
doi:10.1073/pnas.120163297

Davis BM, Waldor MK. 2007. RNase E-dependent processing stabilizes MicX, a Vibrio
cholerae sRNA. *Mol Microbiol*. doi:10.1111/j.1365-2958.2007.05796.x

De Las Peñas A, Connolly L, Gross CA. 1997. The sigmaE-mediated response to

extracytoplasmic stress in Escherichia coli is transduced by RseA and RseB, two negative regulators of sigmaE. *Mol Microbiol* **24**:373–385. doi:9159523

De Lay N, Gottesman S. 2009. The crp-activated small noncoding regulatory RNA CyaR (RyeE) links nutritional status to group behavior. *J Bacteriol*. doi:10.1128/JB.01157-08

Feng L, Rutherford ST, Papenfort K, Bagert JD, Van Kessel JC, Tirrell DA, Wingreen NS, Bassler BL. 2015. A Qrr noncoding RNA deploys four different regulatory mechanisms to optimize quorum-sensing dynamics. *Cell*. doi:10.1016/j.cell.2014.11.051

Gogol EB, Rhodius VA, Papenfort K, Vogel J, Gross CA. 2011. Small RNAs endow a transcriptional activator with essential repressor functions for single-tier control of a global stress regulon. *Proc Natl Acad Sci U S A* **108**:12875–12880.

Granneman S, Kudla G, Petfalski E, Tollervey D. 2009. Identification of protein binding sites on U3 snoRNA and pre-rRNA by UV cross-linking and high-throughput analysis of cDNAs. *Proc Natl Acad Sci U S A* **106**:9613–9618. doi:10.1073/pnas.0901997106

Guo MS, Updegrove TB, Gogol EB, Shabalina SA, Gross CA, Storz G. 2014. MicL, a new σE-dependent sRNA, combats envelope stress by repressing synthesis of Lpp, the major outer membrane lipoprotein. *Genes Dev* **28**:1620–1634. doi:10.1101/gad.243485.114

Han K, Tjaden B, Lory S. 2016. GRIL-seq provides a method for identifying direct targets of bacterial small regulatory RNA by in vivo proximity ligation. *Nat Microbiol* **2**:16239.

Helwak A, Kudla G, Dudnakova T, Tollervey D. 2013. Mapping the human miRNA interactome by CLASH reveals frequent noncanonical binding. *Cell* **153**:654–665. doi:10.1016/j.cell.2013.03.043

Holmqvist E, Vogel J. 2018. RNA-binding proteins in bacteria. *Nat Rev Microbiol*. doi:10.1038/s41579-018-0049-5

Holmqvist E, Wagner EGH. 2017. Impact of bacterial sRNAs in stress responses. *Biochem Soc Trans*. doi:10.1042/BST20160363

Holmqvist E, Wright PR, Li L, Bischler T, Barquist L, Reinhardt R, Backofen R, Vogel J. 2016. Global RNA recognition patterns of post-transcriptional regulators Hfq and CsrA revealed by UV crosslinking in~vivo. *EMBO J* **35**:e201593360--1011.

Hör J, Gorski SA, Vogel J. 2018. Bacterial RNA Biology on a Genome Scale. *Mol Cell* 785–799. doi:10.1016/j.molcel.2017.12.023

Hör J, Vogel J. 2017. Global snapshots of bacterial RNA networks. *EMBO J* **36**:245–247. doi:10.15252/embj.201696072

Johansen J, Rasmussen AA, Overgaard M, Valentin-Hansen P. 2006. Conserved small non-coding RNAs that belong to the sigmaE regulon: role in down-regulation of outer membrane proteins. *J Mol Biol* **364**:1–8.

Kenyon WJ, Thomas SM, Johnson E, Pallen MJ, Spector MP. 2005. Shifts from glucose to certain secondary carbon-sources result in activation of the extracytoplasmic function sigma factor σE in Salmonella enterica serovar Typhimurium. *Microbiology* **151**:2373–2383. doi:10.1099/mic.0.27649-0

Khemici V, Carpousis AJ. 2004. The RNA degradosome and poly(A) polymerase of Escherichia coli are required in vivo for the degradation of small mRNA decay intermediates containing REP-stabilizers. *Mol Microbiol*. doi:10.1046/j.1365-2958.2003.03862.x

Kortmann J, Narberhaus F. 2012. Bacterial RNA thermometers: molecular zippers and switches. *Nat Rev Microbiol* **10**:255–265.

Kudla G, Granneman S, Hahn D, Beggs JD, Tollervey D. 2011. Cross-linking, ligation, and sequencing of hybrids reveals RNA-RNA interactions in yeast. *Proc Natl Acad Sci U S A* **108**:10010–10015. doi:10.1073/pnas.1017386108

Lalaouna D, Carrier M-C, Semsey S, Brouard J-S, Wang J, Wade JT, Massé E. 2015. A 3' External Transcribed Spacer in a tRNA Transcript Acts as a Sponge for Small RNAs to Prevent Transcriptional Noise. *Mol Cell* **58**:393–405.

Lorenz R, Bernhart SH, Höner Zu Siederdissen C, Tafer H, Flamm C, Stadler PF, Hofacker IL. 2011. ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**:26. doi:10.1186/1748-7188-6-26

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**:550.
Lunelli L, Bernabò P, Bolner A, Vaghi V, Marchioretto M, Viero G. 2016. Peering at Brain
Polysomes with Atomic Force Microscopy. *J Vis Exp*. doi:10.3791/53851
Mäder U, Nicolas P, Depke M, Pané-Farré J, Debarbouille M, van der Kooi-Pol MM, Guérin
C, Dérozier S, Hiron A, Jarmer H, Leduc A, Michalik S, Reilman E, Schaffer M, Schmidt
F, Bessières P, Noirot P, Hecker M, Msadek T, Völker U, van Dijl JM. 2016.
Staphylococcus aureus Transcriptome Architecture: From Laboratory to Infection-
Mimicking Conditions. *PLoS Genet* **12**:e1005962.
Magoč T, Salzberg SL. 2011. FLASH: Fast length adjustment of short reads to improve
genome assemblies. *Bioinformatics*. doi:10.1093/bioinformatics/btr507
Martínez-Antonio A, Janga SC, Thieffry D. 2008. Functional organisation of Escherichia coli
transcriptional regulatory network. *J Mol Biol* **381**:238–247.
doi:10.1016/j.jmb.2008.05.054
Melamed S, Faigenbaum-Romm R, Peer A, Reiss N, Shechter O, Bar A, Altuvia Y, Argaman
L, Margalit H. 2018. Mapping the small RNA interactome in bacteria using RIL-seq. *Nat
Protoc* **13**:1–33.
Melamed S, Peer A, Faigenbaum-Romm R, Gatt YEE, Reiss N, Bar A, Altuvia YYY,
Argaman L, Margalit H. 2016. Global Mapping of Small RNA-Target Interactions in
Bacteria. *Mol Cell* **63**:884–897. doi:10.1016/j.molcel.2016.07.026
Miyakoshi M, Chao Y, Vogel J. 2015a. Regulatory small RNAs from the 3' regions of
bacterial mRNAs. *Curr Opin Microbiol*. doi:10.1016/j.mib.2015.01.013
Miyakoshi M, Chao Y, Vogel J. 2015b. Cross talk between ABC transporter mRNAs via a
target mRNA-derived sponge of the GcvB small RNA. *EMBO J* **34**:e201490546--1492.
doi:10.15252/embj.201490546
Monteiro C, Papenfort K, Hentrich K, Ahmad I, Le Guyon S, Reimann R, Grantcharova N,
Römling U. 2012. Hfq and Hfq-dependent small RNAs are major contributors to
multicellular development in Salmonella enterica serovar typhimurium. *RNA Biol*.
doi:10.4161/rna.19682
Moon K, Gottesman S. 2011. Competition among Hfq-binding small RNAs in Escherichia
coli. *Mol Microbiol* **82**:1545–1562. doi:10.1111/j.1365-2958.2011.07907.x
Morita T, Maki K, Aiba H. 2005. RNase E-based ribonucleoprotein complexes: Mechanical
basis of mRNA destabilization mediated by bacterial noncoding RNAs. *Genes Dev*.
doi:10.1101/gad.1330405
Mouali Y El, Gaviria-Cantin T, Sá Nchez-Romero MA, Gibert M, Westermann AJ, Rg Vogel
J, Balsalobre C. 2018. CRP-cAMP mediates silencing of Salmonella virulence at the
post-transcriptional level 1–26. doi:10.1371/journal.pgen.1007401
Navarro Llorens JM, Tormo A, Martínez-García E. 2010. Stationary phase in gram-negative
bacteria. *FEMS Microbiol Rev* **34**:476–495. doi:10.1111/j.1574-6976.2010.00213.x
Newbury SF, Smith NH, Higgins CF. 1987. Differential mRNA stability controls relative gene
expression within a polycistronic operon. *Cell* **51**:1131–1143. doi:10.1016/0092-
8674(87)90599-X
Nitzan M, Rehani R, Margalit H. 2017. Integration of Bacterial Small RNAs in Regulatory
Networks. *Annu Rev Biophys* **46**:131–148. doi:10.1146/annurev-biophys-070816-
034058
Nues R van, Schweikert G, Leau E de, Selega A, Langford A, Franklin R, Iosub I,
Wadsworth P, Sanguinetti G, Granneman S, Van Nues R, Schweikert G, De Leau E,
Selega A, Langford A, Franklin R, Iosub I, Wadsworth P, Sanguinetti G, Granneman S,
Nues R van, Schweikert G, Leau E de, Selega A, Langford A, Franklin R, Iosub I,
Wadsworth P, Sanguinetti G, Granneman S, Van Nues R, Schweikert G, De Leau E,
Selega A, Langford A, Franklin R, Iosub I, Wadsworth P, Sanguinetti G, Granneman S.
2017. Kinetic CRAC uncovers a role for Nab3 in determining gene expression profiles
during stress. *Nat Commun* **8**:12. doi:10.1038/s41467-017-00025-5
Papenfort K, Pfeiffer V, Lucchini S, Sonawane A, Hinton JCD, Vogel J. 2008. Systematic
deletion of Salmonella small RNA genes identifies CyaR, a conserved CRP-dependent

riboregulator of OmpX synthesis. *Mol Microbiol*. doi:10.1111/j.1365-2958.2008.06189.x

Papenfort K, Pfeiffer V, Mika F, Lucchini S, Hinton JCD, Vogel J. 2006. σE-dependent small
RNAs of Salmonella respond to membrane stress by accelerating global omp mRNA
decay. *Mol Microbiol* **62**:1674–1688.

Papenfort K, Said N, Welsink T, Lucchini S, Hinton JCDD, Vogel J. 2009. Specific and
pleiotropic patterns of mRNA regulation by ArcZ, a conserved, Hfq-dependent small
RNA. *Mol Microbiol* **74**:139–158. doi:10.1111/j.1365-2958.2009.06857.x

Pletnev P, Osterman I, Sergiev P, Bogdanov A, Dontsova O. 2015. Survival guide:
Escherichia coli in the stationary phase. *Acta Naturae* **7**:22–33.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**:841–842.

Rhodius VA, Suh WC, Nonaka G, West J, Gross CA. 2006. Conserved and variable
functions of the σE stress response in related genomes. *PLoS Biol* **4**:0043–0059.
doi:10.1371/journal.pbio.0040002

Sætrom P, Sneve R, Kristiansen KI, Snøve O, Grünfeld T, Rognes T, Seeberg E. 2005.
Predicting non-coding RNA genes in Escherichia coli with boosted genetic
programming. *Nucleic Acids Res*. doi:10.1093/nar/gki644

Sedlyarova N, Shamovsky I, Bharati BK, Epshtein V, Chen J, Gottesman S, Schroeder R,
Nudler E. 2016. sRNA-Mediated Control of Transcription Termination in E.~coli. *Cell*
167:111--121.e13.

Sezonov G, Joseleau-Petit D, D'Ari R. 2007. Escherichia coli physiology in Luria-Bertani
broth. *J Bacteriol* **189**:8746–8749. doi:10.1128/JB.01368-07

Sharma E, Sterne-Weiler T, O'Hanlon D, Blencowe BJ. 2016. Global Mapping of Human
RNA-RNA Interactions. *Mol Cell* **62**:618–626. doi:10.1016/j.molcel.2016.04.030

Shimoni Y, Friedlander G, Hetzroni G, Niv G, Altuvia S, Biham O, Margalit H. 2007.
Regulation of gene expression by small non-coding RNAs: a quantitative view. *Mol Syst*
Biol **3**:138.

Sittka A, Pfeiffer V, Tedin K, Vogel J. 2007. The RNA chaperone Hfq is essential for the
virulence of Salmonella typhimurium. *Mol Microbiol*. doi:10.1111/j.1365-
2958.2006.05489.x

Smirnov A, Förstner KU, Holmqvist E, Otto A, Günster R, Becher D, Reinhardt R, Vogel J.
2016. Grad-seq guides the discovery of ProQ as a major small RNA-binding protein.
Proc Natl Acad Sci **113**:11591–11596. doi:10.1073/pnas.1609981113

Smirnov A, Wang C, Drewry LL, Vogel J. 2017. Molecular mechanism of mRNA repression
in *trans* by a ProQ-dependent small RNA. *EMBO J* **36**:1029–1045.
doi:10.15252/embj.201696127

St-Pierre F, Cui L, Priest DG, Endy D, Dodd IB, Shearwin KE. 2013. One-step cloning and
chromosomal integration of DNA. *ACS Synth Biol*. doi:10.1021/sb400021j

Storz G, Vogel J, Wassarman KM. 2011. Regulation by Small RNAs in Bacteria: Expanding
Frontiers. *Mol Cell* **43**:880–891. doi:10.1016/j.molcel.2011.08.022

Tebaldi T, Re A, Viero G, Pegoretti I, Passerini A, Blanzieri E, Quattrone A. 2012.
Widespread uncoupling between transcriptome and translatome variations after a
stimulus in mammalian cells. *BMC Genomics*. doi:10.1186/1471-2164-13-220

Thomason MK, Bischler T, Eisenbart SK, Förstner KU, Zhang A, Herbig A, Nieselt K,
Sharma CM, Storza G. 2015. Global transcriptional start site mapping using differential
RNA sequencing reveals novel antisense RNAs in Escherichia coli. *J Bacteriol*.
doi:10.1128/JB.02096-14

Thompson KM, Rhodius VA, Gottesman S. 2007. SigmaE regulates and is regulated by a
small RNA in Escherichia coli. *J Bacteriol* **189**:4243–4256.

Tjaden B. 2015. De novo assembly of bacterial transcriptomes from RNA-seq data. *Genome*
Biol **16**:1–10. doi:10.1186/s13059-014-0572-2

Travis AJ, Moody J, Helwak A, Tollervey D, Kudla G. 2013. Hyb: A bioinformatics pipeline
for the analysis of CLASH (crosslinking, ligation and sequencing of hybrids) data.
Methods.

Tree Jai J., Granneman S, McAteer SP, Tollervey D, Gally DL. 2014. Identification of

bacteriophage-encoded anti-sRNAs in pathogenic Escherichia coli. *Mol Cell* **55**:199–
213. doi:10.1016/j.molcel.2014.05.006
Tree J J, Granneman S, McAteer SP, Tollervey D, Gally DL. 2014. Identification of
bacteriophage-encoded anti-sRNAs in pathogenic Escherichia coli.
doi:10.1016/j.molcel.2014.05.006
Udekwu KI, Wagner EGH. 2007. Sigma E controls biogenesis of the antisense RNA MicA.
Nucleic Acids Res. doi:10.1093/nar/gkl1154
Updegrove TB, Zhang A, Storz G. 2016. Hfq: The flexible RNA matchmaker. *Curr Opin*
Microbiol **30**:133–138. doi:10.1016/j.mib.2016.02.003
Urban JH, Vogel J. 2007. Translational control and target recognition by Escherichia coli
small RNAs in vivo. *Nucleic Acids Res* **35**:1018–1037. doi:10.1093/nar/gkl1040
Vogel J, Luisi BF. 2011. Hfq and its constellation of RNA. *Nat Rev Microbiol* **9**:578–589.
Wassarman KM, Repoila F, Rosenow C, Storz G, Gottesman S. 2001. Identification of novel
small RNAs using comparative genomics and microarrays. *Genes Dev*.
doi:10.1101/gad.901001
Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2-A
multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**:1189–
1191. doi:10.1093/bioinformatics/btp033
Waters LS, Storz G. 2009. Regulatory RNAs in Bacteria. *Cell* **136**:615–628.
doi:10.1016/j.cell.2009.01.043
Waters SA, McAteer SP, Kudla G, Pang I, Deshpande NP, Amos TG, Leong KW, Wilkins
MR, Strugnell R, Gally DL, Tollervey D, Tree JJ. 2016. Small RNA interactome of
pathogenic E.~coli revealed through crosslinking of RNase E. *EMBO J* **36**:e201694639.
doi:10.15252/embj.201694639
Webb S, Hector RD, Kudla G, Granneman S. 2014. PAR-CLIP data indicate that Nrd1-
Nab3-dependent transcription termination regulates expression of hundreds of protein
coding genes in yeast. *Genome Biol* **15**:R8. doi:10.1186/gb-2014-15-1-r8
Wilson KS, Hippel PH Von. 1995. Transcription termination at intrinsic terminators: The role
of the RNA hairpin (Escherichia coli/RNA polymerase/rho-independent termination).
Biochemistry **92**:8793–8797. doi:10.1073/pnas.92.19.8793
Wright PR, Georg J, Mann M, Sorescu DA, Richter AS, Lott S, Kleinkauf R, Hess WR,
Backofen R. 2014. CopraRNA and IntaRNA: Predicting small RNA targets, networks
and interaction domains. *Nucleic Acids Res* **42**:119–123. doi:10.1093/nar/gku359
Wright PR, Richter AS, Papenfort K, Mann M, Vogel J, Hess WR, Backofen R, Georg J.
2013. Comparative genomics boosts target prediction for bacterial small RNAs. *Proc*
Natl Acad Sci **110**:E3487–E3496. doi:10.1073/pnas.1303248110
Wu X, Bartel DP. 2017. KpLogo: Positional k -mer analysis reveals hidden specificity in
biological sequences. *Nucleic Acids Res*. doi:10.1093/nar/gkx323
1557
1558
1559

Supplementary Figures and legends

Figure 1–figure supplement 1. Hfq expression and Hfq binding to RNAs at different cell densities in UV-irradiated *E. coli*.

(**A**) Western blot analyses of Hfq levels during various growth stages. Hfq-HTF was detected using an anti-TAP primary antibody, and a fluorescent secondary antibody. GroEL was used as a loading control.

(**B**) Quantification of Hfq levels from the Western blot result. The fluorescent signal for Hfq-HTF and GroEL was measured with the LI-COR from biological replicate experiments. The levels of Hfq were normalised to GroEL and expressed as fold-change relative to OD$_{600}$ 0.4.

(**C**) Hfq crosslinking to RNA is similar at each optical density. Autoradiogram showing the purified radioactively labelled Hfq-RNA complexes for each OD$_{600}$ after elution from the nickel beads. Source data for (**A**-**C**) are provided as a Source Data file.

Figure 1–figure supplement 2. RNAseq and Hfq CLASH replicate datasets are highly correlated.

(**A, B**) Scatter plots showing the distribution of $\log_2$ Transcripts Per Million (TPM) normalised read counts for Hfq CLASH (**A**) and RNA-seq (**B**) biological replicates. Pearson R coefficients describing the correlation between the two independent experiments at each OD_{600} are included.

Figure 1–figure supplement 3. Transcriptome-wide maps of Hfq binding to mRNA genes.

(**A**) Heatmaps showing the distribution of Hfq binding sites across all mRNA genes at OD_{600} 0.8 and 4.0. The genes are sorted by their sequence length (x-axis); the darker a nucleotide is, the more Hfq is crosslinked to it. To generate the heatmap, Hfq binding clusters were generated. A 5'-and 3'UTR length of 200bp was used.

(**B**) Hfq binds to poly-U tracks. Significant k-mers (4-8 nt in length) were identified using the pyMotif tool of the pyCRAC package(Webb et al., 2014) and the motif logo was generated using all k-mers with a Z-score > 3, with kpLogo(Wu and Bartel, 2017). (**C**) A more stringent selection of the genes used to generate the distribution of Hfq binding to the transcriptome: all genes with overlapping 5' or 3'UTRs were removed from the analysis to avoid 'duplicate' counting. For all remaining cDNAs, FDR intervals of minimum 20 nt were considered for distribution plotting. The interval length (with UTR flanks as in the GTF annotation file) for each gene was normalized over 100 bins (x-axis), and the fraction of hits in each bin was calculated (y-axis).

Figure 2–figure supplement 1. The sRNA and mRNA levels are not always strongly correlated with Hfq binding during the various stages of growth.
Scatter plots comparing changes in Hfq binding (x-axis) to RNA levels (y-axis) for the indicated OD_{600} for sRNAs (**A**) and all transcript classes (**B**) show that the correlation improves at higher OD_{600}. For each OD_{600}, the Transcripts Per Million (TPM) normalised read counts at each

density-point were divided by the OD$_{600}$ 0.4 data, and the resulting ratios were log$_2$-normalized. Pearson R coefficients describing the correlation between the two independent experiments at each OD$_{600}$ is also included for each plot.

Figure 4–figure supplement 1. sRNA-RNA interactions identified by CLASH are growth-stage specific. sRNA-sRNA network generated from the statistically significant CLASH

interactions from two biological replicates, recovered at three main growth stages: exponential (OD_{600} 0.4 and 0.8), transition (OD_{600} 1.2, 1.8, 2.4) and early stationary (OD_{600} 3.0 and 4.0). The thickness of the edges is proportional to the $\log_2$(unique chimera count for each interaction). Only sRNAs transcribed from independent promoters were included in the analysis.

Figure 4–figure supplement 2. Interactions between ArcZ, CyaR and GcvB are conserved. Alignments of ArcZ, CyaR and GcvB were compiled as previously described(van Nues et al., 2016). Names of the enteric bacteria from which the sequence was retrieved are given on the left. Indicated are possible stem-loops (brackets), seed regions (boxed in dashed lines) and their interactions with various sections of ArcZ, CyaR or GcvB (blue and purple bars) or with other sRNAs and mRNAs (black bars). The CyaR sequence indicated with a blue bar is predicted to interact with two regions in GcvB (see blue bars in GcvB alignment), including the second seed sequence. A second interaction (pink bars) involves the seed sequence regions of CyaR and GcvB.

Figure 4–figure supplement 3. ArcZ downregulates CyaR expression.
(**A**) ArcZ and CyaR were overexpressed from a plasmid-borne IPTG inducible promoter (pZE-ArcZ and pZE-CyaR) and the data were compared to data from cells carrying a scrambled RNA plasmid (pJV300). The co-expressed candidate target sRNAs (expressed from pZA-derived backbone) were induced with anhydrotetracycline hydrochloride (panels I and II). The bars indicate the mean fold-change in expression relative to the level of 5S rRNA (*rrfD*) in cells with the indicated vector. In panel III endogenous ArcZ levels were measured upon over-expression of CyaR. Error bars indicate the standard error of the mean from three biological replicates and three technical replicates per experiment.
(**B**) Northern blot analysis of ArcZ and CyaR. The cells containing both the empty pZA and pJV300 plasmids (lanes 1, 5, 9) do not express ArcZ and CyaR at detectable levels.

Figure 5–figure supplement 1. *YgaM, gadE* and *malG* contain sRNAs in their 3'UTRs.
(**A**) Validation of *malG* 3'UTR (MdoR), *ygaM* 3'UTR (YgaN) and *gadE* 3'UTR (GadF) sRNAs by Northern blot. Total RNA extracted from cells at the indicated optical densities (OD_{600}) was resolved on 8% PAA-UREA gels and subjected to Northern blotting using oligos that hybridize with the 3'UTR of the respective transcripts. The asterisk indicates cross-reactivity of the probe with the 5S rRNA. The locations of the 3'UTR-derived fragments are indicated. MdoR and YgaN are ~110nt, whereas GadF fragment is ~ 90nt.
(**B, C**) Analysis of (Terminator 5'-Phosphate Dependent Exonuclease (TEX) RNA-seq datasets(Thomason et al., 2015) indicates that YgaN has an independent promoter, while MdoR is a degradation product of the *malEFG* operon. Genome browser tracks showing the location and normalised reads of *ygaM* and *malG* fragments in the absence of TEX (–TEX) and in the presence of TEX (+TEX). The *ygaM* and putative YgaN promoters are indicated. Independently transcribed YgaN could be further processed by RNases, at the site marked with a dashed vertical line.

Figure 6–figure supplement 1. Conservation and target prediction analyses of MdoR.
(**A**) MdoR contains conserved and variable regions. Sequence conservation analysis of MdoR in several Gram-negative bacteria species (Mafft algorithm with defaults); the arrow indicates the 5'-end of MdoR; the *malG* stop codon and Rho-independent terminator sites are highlighted; the horizontal black lines indicate the base-pairing regions of MdoR with *ptsH* (top), OhsC, *ompA* and *ompC* (bottom) in *E. coli* as predicted by CLASH combined with *in silico* folding (RNACofold).
(**B**) MdoR is predicted to interact with its targets using two seed regions. Interaction regions within MdoR and top target mRNAs predicted by CopraRNA(Wright et al., 2014, 2013). Density plots showing the relative frequency of a specific MdoR (Left) or mRNA (Right) nucleotide position in all predicted sRNA-mRNA interactions with a p-value < 0.01 in all considered homologs. The vertical lines indicate local maxima; the aligned regions of the homologs are shown in grey, whereas the interacting regions are shown in arbitrary colors; only the top 20 representative clusters members are shown in the aligned regions, with the gene names indicated on the right.

Figure 8–figure supplement 1. Validation of MdoR-*ompC* interaction using GFP reporters.

(**A**) Design of the wild-type and mutant *ompC* constructs. The panel indicates the base-pairing region within the MdoR-*ompC* duplex that was mutated. We created an MdoR seed mutant (SM) and an *ompC* mutant in which base pairing with MdoR SM was restored.

(**B**) Plasmid system used for the reporter assay: in *E. coli* TOP10 cells, low-copy plasmids constitutively overexpress target 5'UTRs fused to sfGFP and medium-copy plasmids overexpress the full-length sRNAs upon induction with L-arabinose.

(**C**) MdoR downregulates expression of OmpC and OmpA sfGFP fusions. *In vivo* fluorescence measurements of OmpC, OmpC SM and OmpA sfGFP fusion proteins was measured using a Tecan plate reader system. As a negative control we included sfGFP alone in the presence or absence of sRNAs. The 'no sRNA' expressing strains contain the empty pBAD plasmid. The y-axis indicates fluorescence units (F.U.) reported by the plate reader. Experiments were performed in technical and biological triplicates; the fluorescence means and SEM of three biological replicates are reported. Source data for are provided as a Source Data file.

Figure 8–figure supplement 2. MdoR regulates *ompC* mRNA translation in *E. coli*.

(**A**) Cultures at OD_{600} 0.4 overexpressing MdoR or no sRNA for 15 minutes were subjected to polysome profiling. Profiles of the polysomal (2x70S) and subpolysomal fractions obtained for the empty plasmid control and MdoR overexpression samples.

(**B**) RT-qPCR analysis of the polysomal fractions: A 'total' fraction was obtained by mixing equal amounts/volumes of the polysomal and subpolysomal fractions and is representative of the cytosol/cell lysate content. Total RNA was extracted from all fractions (polysomal, subpolysomal and total). Expression of *ompC* in the polysomal fractions was quantified relative to the amount in 'total' fraction, normalized to *recA,* and calculated as fold-change relative to the control sample (y-axis). The experiments were performed in technical triplicates; the standard error of the mean (SEM) of three biological replicates fold changes are reported as error bars. Significance of the difference in *ompC* mRNA level in polysomes was assessed with a two-tailed Student's t-test. Source data for are provided as a Source Data file.

Supplementary References

Thomason MK, Bischler T, Eisenbart SK, Förstner KU, Zhang A, Herbig A, Nieselt K, Sharma CM, Storza G. 2015. Global transcriptional start site mapping using differential RNA sequencing reveals novel antisense RNAs in Escherichia coli. *J Bacteriol*. doi:10.1128/JB.02096-14

van Nues RW, Castro-Roa D, Yuzenkova Y, Zenkin N. 2016. Ribonucleoprotein particles of bacterial small non-coding RNA IsrA (IS61 or McaS) and its interaction with RNA polymerase core may link transcription to mRNA fate. *Nucleic Acids Res* **44**:2577–2592.

Webb S, Hector RD, Kudla G, Granneman S. 2014. PAR-CLIP data indicate that Nrd1-Nab3-dependent transcription termination regulates expression of hundreds of protein coding genes in yeast. *Genome Biol* **15**:R8. doi:10.1186/gb-2014-15-1-r8

Wright PR, Georg J, Mann M, Sorescu DA, Richter AS, Lott S, Kleinkauf R, Hess WR, Backofen R. 2014. CopraRNA and IntaRNA: Predicting small RNA targets, networks and interaction domains. *Nucleic Acids Res* **42**:119–123. doi:10.1093/nar/gku359

Wright PR, Richter AS, Papenfort K, Mann M, Vogel J, Hess WR, Backofen R, Georg J. 2013. Comparative genomics boosts target prediction for bacterial small RNAs. *Proc Natl Acad Sci* **110**:E3487–E3496. doi:10.1073/pnas.1303248110

Wu X, Bartel DP. 2017. KpLogo: Positional k -mer analysis reveals hidden specificity in biological sequences. *Nucleic Acids Res*. doi:10.1093/nar/gkx323