Multidimensional analysis of Gammaherpesvirus RNA expression reveals

unexpected heterogeneity of gene expression

Lauren M. Oko[1], Abigail K. Kimball[2], Rachael E. Kaspar[2], Ashley N. Knox[1], Carrie B.

Coleman[1], Rosemary Rochford[1], Tim Chang[3], Benjamin Alderete[4], Linda F. van Dyk[1]*,

Eric T. Clambey[2]*#

Affiliations:

[1] Department of Immunology and Microbiology, [2] Department of Anesthesiology,

University of Colorado Denver | Anschutz Medical Campus, Aurora, CO, 80045, USA

[3] MilliporeSigma, a business of Merck KGaA, Darmstadt, Germany (Seattle, WA)

[4] Luminex Corporation, Austin, TX, USA

* Co-Corresponding authors:

Eric T. Clambey, eric.clambey@ucdenver.edu, 303-724-7783 (phone)

Linda F. van Dyk, linda.vandyk@ucdenver.edu, 303-724-4207 (phone)

 # Lead author for MS correspondence

Short title: Single-cell heterogeneity of Gammaherpesvirus RNA expression

ABSTRACT

Virus-host interactions are frequently studied in bulk cell populations, obscuring

cell-to-cell variation. Here we investigate endogenous herpesvirus gene expression at

the single-cell level, combining a sensitive and robust fluorescent in situ hybridization

platform with multiparameter flow cytometry, to study the expression of

gammaherpesvirus non-coding RNAs (ncRNAs) during lytic replication, latent infection

and reactivation in vitro. This method allowed robust detection of viral ncRNAs of

murine gammaherpesvirus 68 (γHV68), Kaposi's sarcoma associated herpesvirus and

Epstein-Barr virus, revealing variable expression at the single-cell level. By quantifying

the inter-relationship of viral ncRNA, viral mRNA, viral protein and host mRNA

regulation during γHV68 infection, we find heterogeneous and asynchronous gene

expression during latency and reactivation, with reactivation from latency identified by a

distinct gene expression profile within rare cells. Further, during lytic replication with

γHV68, we find many cells have limited viral gene expression, with only a fraction of

cells showing robust gene expression, dynamic RNA localization, and progressive

infection. Lytic viral gene expression was enhanced in primary fibroblasts and by

conditions associated with enhanced viral replication, with multiple subpopulations of

cells present in even highly permissive infection conditions. These findings, powered by

single-cell analysis integrated with automated clustering algorithms, suggest inefficient

or abortive γHV infection in many cells, and identify substantial heterogeneity in viral

gene expression at the single-cell level.

AUTHOR SUMMARY

The gammaherpesviruses are a group of DNA tumor viruses that establish

lifelong infection. How these viruses infect and manipulate cells has frequently been

studied in bulk populations of cells. While these studies have been incredibly insightful,

there is limited understanding of how virus infection proceeds within a single cell. Here

we present a new approach to quantify gammaherpesvirus gene expression at the

single-cell level. This method allows us to detect cell-to-cell variation in the expression

of virus non-coding RNAs, an important and understudied class of RNAs which do not

encode for proteins. By examining multiple features of virus gene expression, this

method further reveals significant variation in infection between cells across multiple

stages of infection, even in conditions generally thought to be highly uniform. These

studies emphasize that gammaherpesvirus infection can be surprisingly heterogeneous

when viewed at the level of the individual cell. Because this approach can be broadly

applied across diverse viruses, this study affords new opportunities to understand the

complexity of virus infection within single cells.

INTRODUCTION

The *Herpesviridae* are a family of large dsDNA viruses that include multiple

prominent human and animal pathogens [1]. Although these viruses infect different cell

types, and are associated with diverse pathologies, they share conserved genes and

two fundamental phases of infection: lytic replication and latent infection [1]. Lytic

replication is characterized by a cascade of viral gene expression, active viral DNA

replication and the production of infectious virions. Conversely, latency is characterized

by limited viral gene expression and the absence of de novo viral replication. While

latent infection is a relatively quiescent form of infection, the herpesviruses can

reactivate from latency, to reinitiate lytic replication.

Among the herpesviruses, the gammaherpesviruses (γHV) are lymphotropic

viruses that include the human pathogens Epstein-Barr virus (EBV) [2] and Kaposi's

sarcoma associated herpesvirus (KSHV) [3]. Murine gammaherpesvirus 68 (γHV68, or

MHV-68; ICTV nomenclature *Murid herpesvirus 4*, MuHV-4), is a well-described small

animal model for the γHVs [4]. While these viruses establish a lifelong infection that is

often clinically inapparent, immune-suppressed individuals are particularly at risk for

γHV-associated malignancies [5].

Herpesvirus gene expression is extremely well-characterized in bulk populations.

Despite increasing evidence for single-cell heterogeneity in gene expression [6-8], there

remains limited understanding of herpesvirus infection at the single-cell level [9-12].

Here, we tracked endogenous viral and host RNAs using a sensitive, robust fluorescent

in situ hybridization assay combined with multiparameter flow cytometry (PrimeFlow^TM)

[13] to analyze the expression and inter-relationships of viral ncRNA, viral mRNA and

cellular mRNA at the single-cell level during γHV latency, reactivation and lytic

replication. These studies revealed unanticipated heterogeneity of infection,

emphasizing how single-cell analysis of virus infection can afford significant new

insights into the complexity of γHV infection.

RESULTS

Single-cell analysis of viral RNAs during lytic infection.

Traditional measurements of gene expression frequently rely on pooled cellular

material, obscuring intercellular variation in gene expression. To better define

expression of γHV RNAs at the single cell level, we employed the PrimeFlow™ RNA

assay [13] to study viral gene expression during murine gammaherpesvirus 68 (γHV68)

infection, a small animal γHV [4, 13]. This method is a highly sensitive, extremely

specific in situ hybridization assay, integrating Affymetrix-designed branched DNA

technology with single-cell analysis powered by multiparameter flow cytometry. This

method has been successfully used to detect both virus and host RNAs (e.g. in the

context of HIV infected individuals [13, 14]).

We first tested the ability of PrimeFlow™ to measure multiple viral RNAs during

lytic infection with γHV68, including small non-coding RNAs (tRNA-miRNA encoding

RNAs or TMERs [15]) and mRNAs. Mouse 3T12 fibroblasts were infected with a

multiplicity of infection (MOI=5 plaque forming units (PFU) of virus/cell). Under these

conditions, TMER-5, one of the eight γHV68 TMERs, and the γHV68 ORF73, were

readily detectable by conventional real-time PCR in γHV68-infected, but not mock-

infected, cultures (Fig. 1A, C). Parallel cultures were analyzed for RNA expression by

PrimeFlow™. Whereas mock-infected cells had no detectable expression of either the

γHV68 TMERs or ORF73, WT γHV68-infected fibroblasts had a prominent population of

TMER+ and ORF73+ cells, respectively (Fig. 1B and 1D). Infection of cells with a

TMER-deficient γHV68 (TMER-TKO [16]), in which TMER expression is ablated through

promoter disruption, revealed no detectable TMER expression (Fig. 1B), yet robust

ORF73 expression (Fig. 1D). Parallel studies revealed ready detection of ORF18,

another γHV68 gene product (Fig. 1E). These studies show that PrimeFlow^{TM} is a

sensitive, robust and specific method to detect both viral non-coding and messenger

RNAs during lytic infection, quantifying both the frequency of gene expression and

expression on a per cell basis.

Heterogeneous gene expression during γHV latency and reactivation from latency.

γHV latency is characterized by limited gene expression. We next

measured viral RNAs during latency and reactivation using the γHV68-infected

A20 HE2.1 cell line (A20.γHV68), a drug-selected latency model with restricted viral

gene expression that can reactivate following stimulation [17]. A20.γHV68 cells are

characterized by restricted viral gene expression, yet remain competent for reactivation

from latency and the production of infectious virions following chemical stimulation with

the phorbol ester, TPA [17, 18].

When we compared TMER expression between uninfected (parental, virus-

negative A20) and infected (A20.γHV68) cells by qRT-PCR, the viral ncRNA TMER-5

was readily detectable in A20.γHV68 cells above background signals in parental A20

cells, with minimal changes between untreated and chemically-stimulated conditions

(Fig. 2A). PrimeFlow^{TM} analysis of TMER expression in untreated A20.γHV68 cells

revealed that a majority of these cells expressed the TMERs, as defined by a positive

signal in samples subjected to the TMER probe relative to unstained cells (Fig. 2B).

Untreated A20.γHV68 cells contained a high frequency of cells expressing intermediate

levels of TMERs (i.e. TMER^{mid} cells), with a significant signal enrichment above

parental, virus-negative A20 cells (Fig. 2C). While the frequency of TMER^{mid} cells

remained relatively constant following treatment with TPA (compare "Untreated" versus

"TPA stimulated", Fig. 2C), TPA stimulated A20.γHV68 cultures also contained a small

fraction of cells with high levels of TMERs (i.e. TMER^{high} cells), not present in untreated

cultures (Fig. 2C-D). Chemical stimulation is known to result in variable penetrance of

reactivation in latently infected cell lines [17]. Based on this, we hypothesized that these

rare, TMER^{high} cells may represent a subset of cells that are undergoing reactivation

from latency.

To test this, we analyzed the properties of TMER^{mid} and TMER^{high} cells,

comparing viral protein expression in untreated and stimulated A20.γHV68 cells. We

analyzed: 1) a γHV68 expressed GFP-hygromycin resistance fusion protein

(HygroGFP), under the control of a heterologous viral promoter (the CMV immediate

early promoter) [17], and 2) the γHV68 regulator of complement activation (RCA), a viral

protein encoded by the γHV68 ORF4, an early-late transcript [19]. The vast majority of

TMER^{mid} cells were negative for HygroGFP and RCA (i.e. HygroGFP- RCA-),

regardless of whether the cells were present in untreated or stimulated cultures (Fig.

2E-F). Conversely, TMER^{high} cells, which were present at an increased frequency in

stimulated cultures, had a significantly increased frequency of HygroGFP+ cells with

induction of RCA protein+ cells in a subset of cells when compared to TMER^{mid} cells

present in either untreated or stimulated cultures (Fig. 2E-F). By using imaging flow

cytometry, we further analyzed the subcellular localization of TMERs in TMER^{mid} cells

compared with TMER^{high} RCA+ cells. TMERs were predominantly nuclear in both

TMER^{mid} and TMER^{high} RCA+ cells, as defined by co-localization with DAPI

fluorescence (Fig. 2G). These data demonstrate that the TMERs are expressed during

latency, and that following reactivation-inducing stimulation, TMERs are further induced

in a rare subset of cells which are characterized by increased viral transcription and

translation.

Detection of endogenous viral gene expression during KSHV latency and reactivation.

To extend these findings, we analyzed viral gene expression in the KSHV

infected B cell tumor line, BCBL-1, focused on detection of an abundant viral ncRNA,

the KSHV polyadenylated nuclear RNA (PAN, nut1, or T1.1) [20]. PAN RNA is known to

be highly inducible upon induction of reactivation in KSHV latently infected B cell

lymphoma cell lines [10, 20]. The frequency of PAN RNA+ cells was low in untreated

BCBL-1 cells, with ~1% of cells spontaneously expressing this ncNRA (Fig. 3A-B).

Despite the low frequency, this hybridization was clearly above background, as ncRNA

defined on the KSHV- and EBV-negative B cell lymphoma cell line BL41 [21, 22] (Fig.

3A-B). Upon stimulation of BCBL-1 cells with the reactivation-inducing stimuli TPA and

sodium butyrate, the frequency of PAN RNA+ cells significantly increased with

expression in ~25% of cells (Fig. 3A-B). Although stimulation of BCBL-1 cells

significantly increased the frequency of PAN RNA+ events compared to untreated

cultures, PAN RNA expression on an individual cell basis was comparable between

cells from untreated or stimulated cultures (Fig. 3C). As anticipated, stimulation of

BCBL-1 cells was associated with increased viral DNA, consistent with stimulated

cultures undergoing reactivation from latency (Supplemental Fig. 1).

We next analyzed the properties of BCBL-1 cells as a function of PAN RNA

expression. In untreated cells, PAN RNA+ or RNA- cells had comparable cell size

(define by forward scatter, FSC) and granularity (defined by side scatter, SSC). ORF73

RNA expression was low in untreated BCBL-1 samples, with signal intensity in PAN

RNA- cells close to the background fluorescence observed in unstained samples. PAN

RNA+ cells in untreated cultures had a modest increase in ORF73 RNA expression

relative to PAN RNA- cells (Fig. 3D-E). In stimulated BCBL-1 cultures, PAN RNA+ cells

had a modest decrease in cell size (defined by forward scatter) and a trend towards

reduced granularity (defined by side scatter) compared to PAN RNA- cells (Fig. 3F-G).

Stimulated BCBL-1 cultures also had an increased ORF73 signal when compared to

unstained samples (Fig. 3F), with PAN RNA+ cells again showing ~2-fold increase

compared to PAN RNA- cells (Fig. 3F-G). These data demonstrate robust detection of

PAN RNA by PrimeFlow™, and further identify PAN RNA expression in a subset of both

untreated and reactivation-induced BCBL-1 cells.

Detection of endogenous viral gene expression during EBV latency and reactivation.

EBV encodes two abundant non-coding RNAs, the EBV-encoded RNAs (EBERs)

EBER1 and EBER2. We tested the ability of the PrimeFlow™ method to detect EBER in

an EBV positive, Burkitt lymphoma type I latency cell line, Mutu I [23]. EBER expression

was detected in ~45% of Mutu I cells in either untreated or TPA stimulated conditions,

with EBER+ cells defined relative to background probe hybridization in the KSHV- and

EBV-negative BL41 cell line, a conservative measurement (Fig. 4A). TPA stimulated

Mutu I cells showed a modest, 2-fold increase in EBER expression on an individual cell

basis, relative to untreated EBER+ cells (Fig. 4B). Based on these data, EBER

expression in Mutu I cells appears to be constitutive, with stimulation under these

conditions resulting in minimal consequences on either the frequency or per-cell

expression of the EBERs.

To extend these findings, we further analyzed EBER expression in a panel of

LCLs and during in vitro infection of primary human B cells. EBER expression was

readily detected in LCL cultures, with EBER+ cells also found in a subset of human

primary B cells following in vitro EBV infection (Supplemental Fig. 2, 3). During EBV

infection of human primary B cells, EBER+ B cells had an increased cell size and

granularity relative to EBER- B cells, with increased expression of both CD69 and actin

mRNA, an activated cell phenotype (Supplemental Fig. 3D). These data demonstrate

cell to cell variation in EBER expression and suggest EBER expression as a potential

discriminator to investigate variability during EBV infection.

Single-cell analysis of actin mRNA degradation as a readout of virus-induced host

shutoff.

Many herpesviruses, including γHV68, EBV and KSHV, induce host shutoff

during lytic replication and reactivation from latency, a process characterized by

dramatic decreases in host mRNAs [24, 25]. Consistent with published reports [24],

qRT-PCR analysis of a cellular housekeeping gene, β-actin (Actb), showed reduced

actin mRNA in γHV68 lytically-infected fibroblasts by 18 hours pi (Fig. 5A). While mock-

infected samples had a uniformly positive population of actin RNA^{high} cells detectable by

PrimeFlow^{TM}, γHV68-infected fibroblast cultures demonstrated a bimodal distribution of

actin RNA^{high} and actin RNA^{low} cells (Fig. 5B). The actin RNA^{low} population had a

fluorescent signal that was only modestly above background fluorescence (defined by

the "No probe" sample), suggesting an all-or-none phenomenon in which cells either

had no change in actin RNA levels or had pronounced actin RNA degradation.

Simultaneous analysis of TMER and actin RNA expression revealed that actin RNA^{low}

cells were frequently TMER^{high}, with actin^{high} cells frequently TMER^{negative} at this time

(Fig. 5C).

To determine whether actin RNA regulation could also be observed during γHV68

latency and reactivation, we measured actin RNA levels in A20.γHV68 cells. Parental,

virus-negative A20 cells and A20.γHV68 cells had relatively comparable actin RNA

levels by qRT-PCR, in both untreated and stimulated cells (Fig. 5D). Given that host

shutoff is expected to primarily occur in rare, reactivating cells, we measured actin RNA

degradation relative to TMER expression by the PrimeFlow^{TM} method. Untreated

A20.γHV68 cultures had no discernable population of TMER+ actin RNA^{low} events,

whereas stimulated cultures were characterized by a rare population of TMER^{high} actin

RNA^{low} cells (Fig. 5E). We further compared actin RNA expression between TMER^{mid}

and TMER^{high} cells, in untreated versus stimulated cultures using our previously defined

subpopulations (Fig. 2). While TMER^{mid} cells from either untreated or stimulated

cultures were predominantly actin RNA+, TMER^{high} cells from stimulated cultures

showed a significant increased frequency of actin RNA^{low} events (Fig. 5F-G). These

studies reveal actin RNA as a sensitive indicator of virus-induced host shutoff, and

demonstrate this as an all-or-none phenomenon that can be readily queried at the

single-cell level.

Heterogeneity of gene expression during de novo lytic replication.

Next, we revisited our analysis of gene expression during de novo lytic infection

of fibroblasts, to examine co-expression relationships between viral ncRNA (TMERs),

viral mRNA (the γHV68 ORF73), viral protein (RCA protein) and cellular actin mRNA

degradation [19, 24]. Mouse 3T12 fibroblasts were infected with a multiplicity of infection

(MOI=5 plaque forming units of virus/cell), harvested 16 hpi and then subjected to the

PrimeFlow™ method.

To enable an unbiased, automated analysis of gene expression profiles in γHV68

lytically infected cells relative to mock infected cells, data were subjected to the

automated clustering algorithm X-shift [26], to identify potential subpopulations of cells

with heterogeneous gene expression in these cultures. By sampling 1,000 cells from

multiple mock- and virus-infected cultures, the X-shift algorithm consistently identified 7

major clusters of cells (Fig. 6A) defined by varying gene expression patterns. While

some of the clusters were exclusively found in mock-infected cultures, virus-infected

cultures contained three broad types of cell clusters: 1) cells, with no detectable

expression of either the TMERs or ORF73 and normal actin RNA, 2) fully infected cells,

with robust expression of the TMERs, ORF73, actin RNA downregulation and frequent

expression of the RCA protein, and 3) intermediate populations characterized by

variable expression of the TMERs and ORF73 (Fig. 6A).

To validate these findings using a more conventional method, we compared

TMER and ORF73 RNA co-expression on a biaxial plot. By comparing mock-infected,

WT-infected and TMER-TKO-infected cultures, this analysis revealed five populations of

gene expression (Fig. 6B), including cells with: 1) no detectable expression of either

viral RNA (TMER- ORF73-), 2) TMER+ ORF73- cells (bottom right quadrant), 3) TMER-

ORF73+ cells (upper left quadrant), 4) TMER^{low} ORF73^{low} cells (lower left edge of the

upper right quadrant), and 5) TMER^{high} ORF73^{high} cells (upper right quadrant). The

definition of TMER positive events was defined based on background fluorescent levels

observed in TMER-TKO infected cultures (Fig. 6B). These 5 populations were each

assigned a unique color for subsequent analysis (Fig. 6C).

We compared cellular phenotype and gene expression across these 5

populations. Analysis of TMERs, ORF73, actin RNA, RCA protein, cell size (forward

scatter), and granularity (side scatter) revealed multiple types of viral gene expression.

TMER- ORF73- cells (in black) had no evidence of viral gene expression, with no

detectable viral protein (RCA) or actin downregulation (Fig. 6D-E). Cells with low

expression of either the TMERs and/or ORF73 contained viral RNAs, but had minimal

expression of either viral protein or actin downregulation (Fig. 6D-E). In stark contrast,

cells that were TMER^{high} ORF73^{high} (in red, Fig. 6D-E) had multiple characteristics of

progressive virus infection including a prominent fraction of cells that expressed RCA

and/or had actin RNA downregulation. Further, TMER^{high} ORF73^{high} cells were

consistently smaller in cell size (defined by forward scatter, FSC) and higher in

granularity (defined by side scatter, SSC), a feature that was unique to this phenotype

(Fig. 6D-E).

TMER^{high} ORF73^{high} also demonstrated robust expression of an early gene

(ORF64) and a late gene (ORF18) (Supplemental Fig. 4A). While some TMER^{low}

ORF73^{low} cells expressed early and late genes, cells defined as TMER- 73- had no

discernable expression of either early or late genes (Supplemental Fig. 4A). These data

emphasize that a subset of 3T12 fibroblasts subjected to infection with an MOI=5 and

harvested at 16 hpi have negligible signs of viral gene expression using this method.

This is further illustrated by comparing mock and γHV68 infected cultures for TMER,

ORF73, ORF64 and ORF18 expression which demonstrates a subset of cells with

minimal expression across each of these viral genes (Supplemental Fig. 4B). One

potential explanation for this could be that TMER- ORF73- cells are not infected. To test

this, we sort purified cells based on TMER and ORF73 expression, followed by

quantitation of viral DNA. TMER^{high} ORF73^{high} cells and TMER+ 73- cells had

comparable levels of viral DNA, which were approximately ten-fold higher than viral

DNA levels in TMER- 73- cells (Supplemental Fig. 4C). Despite this moderate difference

in viral DNA, however, viral DNA levels in TMER- 73- cells were still significantly above

background signal present in mock infected cultures (Supplemental Fig. 4C). Another

potential explanation for this limited gene expression in TMER- 73- cells is that perhaps

they are in a distinct phase of the cell cycle relative to other cells. We found that there

was no sizable impact of cell cycle stage on whether cells were TMER- ORF73- or

TMER^{high} ORF73^{high} (Supplemental Fig. 4D). These data identify unexpected

heterogeneity during in vitro lytic infection, with TMER^{high} ORF73^{high} cells characterized

by robust viral transcription, TMER^{low} ORF73^{low} cells characterized by a lower frequency

of cells expressing early and late genes, and TMER- ORF73- cells containing viral DNA

with little to no detectable viral gene expression.

Given the heterogeneous patterns of RNA and protein expression among

lytically-infected cells, we next queried TMER subcellular localization as a function of

viral gene expression using imaging flow cytometry. While the majority of TMER+ cells

had a primarily nuclear TMER localization (defined by DAPI co-localization, as in [27]),

the frequency of cells with nuclear TMER localization was highest among TMER+

ORF73- cells and lowest among TMER+ ORF73+ RCA+ cells (Fig. 7A-B). These data

suggest that the TMERs can be localized in either the nucleus or cytoplasm during

γHV68 lytic replication, and that this localization is not strictly a function of magnitude of

gene expression.

Finally, we used tSNE, a dimensionality reduction algorithm, to better delineate

the relationship between TMER, ORF73, RCA protein and actin downregulation across

populations defined by variable TMER and ORF73 expression. In tSNE-based

visualization, events are subjected to dimensionality reduction, with all events plotted

according to the composite parameters tSNE1 and tSNE2. In each tSNE-based plot,

values from individual cells are depicted as individual dots on the plot. Cells with similar

expression profiles are visually clustered together using this algorithm (e.g. [28]).

Consistent with our histogram analysis (Fig. 6D-E), cells that were negative for TMER

and ORF73 and cells that expressed either TMERs or ORF73 were relatively uniform in

gene expression (Fig. 7C). In contrast, TMER^{high} ORF73^{high} cells expressed a wider

array of phenotypes, including both a predominant fraction of cells that were actin

RNA^{low} RCA+, and a distinct group of cells that were actin RNA+ RCA- (Fig. 7C).

Notably, RCA expression and actin degradation were inversely correlated, with very few

cells that expressed RCA also high for actin RNA. Actin RNA+ populations among

TMER^{high} ORF73^{high} cells were associated with larger cell size (Supplemental Fig. 5).

The diversity of phenotypes among TMER^{high} ORF73^{high} cells was confirmed by biaxial

gating of actin RNA versus RCA protein expression (Fig. 7D). In total, these data

indicate heterogeneous progression of lytic replication in vitro. While some cells have

robust viral mRNA and protein expression, additional cell subsets are characterized by

limited or divergent gene expression.

Lytic cycle viral gene expression is influenced by target cell type, changes over time,

and is modulated by conditions that alter lytic replication.

Our previous studies of γHV68 lytic replication used 3T12 fibroblasts, an

immortalized cell line. Mouse embryonic fibroblasts (MEFs) are primary cells that are

highly permissive for γHV68 infection, with 5 to 10-fold greater sensitivity to virus

infection than 3T12 fibroblasts [29]. We therefore compared viral gene expression

between 3T12 fibroblasts and MEFs, using MEFs derived either from wild-type or

IFNAR1 KO mice. All cells were infected with WT γHV68 (MOI=5) and analyzed at 16

hpi. Both WT and IFNAR KO MEFs showed enhanced viral gene expression relative to

infected 3T12 cells at 16 hpi, with infected MEF cultures characterized by an increased

frequency of events that expressed ORF73, TMERs, RCA and were actin^{low} (Fig. 8A).

Infected MEFs also had an increased frequency of ORF73+TMER+ cells and a

decreased frequency of ORF73-TMER- cells relative to 3T12 cells (Fig. 8B). When we

analyzed the cellular distribution of viral gene expression across all 16 possible

combinations of viral gene expression, resulting from different combinations of ORF73,

TMER, RCA and actin^{low} phenotypes, infected MEF cultures compared to infected 3T12

cultures had: i) fewer ORF73-TMER- cells, ii) an increased frequency of

ORF73+TMER+actin^{low} cells, iii) with the most frequent subset defined as cells with full

viral gene expression (i.e. ORF73+TMER+RCA+actin^{low}) (Fig. 8C). Viral gene

expression distribution between WT and IFNAR1 KO MEFs were generally comparable,

suggesting little to no contribution of type I IFN receptor mediated signaling as a

regulator of viral gene expression in MEFs. These data demonstrate that the diversity

and relative abundance of viral gene expression can be influenced by target cell type.

The increased viral gene expression observed in MEFs raised the possibility that

viral gene expression could be further modulated in 3T12 cells, either as a function of

time, multiplicity of infection or viral genotype. We therefore compared viral gene

expression in 3T12 fibroblasts, comparing: i) WT γHV68 infection with 5 PFU/cell, ii) C-

RTA γHV68 (a recombinant virus engineered to overexpress Rta, the immediate early

viral transactivator [30]) infection with 5 PFU/cell, and iii) WT infection using 100

PFU/cell. Cells were harvested at 6 and 16 hpi. When comparing WT infection at MOI=5

between 6 and 16 hpi, ORF73 expression peaked at 6 hpi with decreased expression

by 16 hpi, in contrast to TMER, RCA and actin^{low} phenotypes which were infrequent at 6

hpi and increased by 16 hpi (Fig. 8D). 3T12 cells subjected to WT infection at an

MOI=100 showed an increased expression of ORF73, TMER and RCA by 6 hpi, above

WT infection with an MOI=5. By 16 hpi, cultures infected with C-RTA and WT MOI=100

had increased frequencies of TMER+ and ORF73+TMER+ events relative to WT MOI=5

cultures (Fig. 8E).

The diversity and progression of viral gene expression was further illustrated by

analyzing the frequency of all 16 different combinations of possible gene expression

between these different virus conditions. While WT MOI=5 cultures showed a time-

dependent switch from ORF73+ events to TMER+, ORF73+TMER+, and

ORF73+TMER+RCA+actin^{low} events, the frequency of each of these populations was

typically below 10% of cultures (Fig. 8F). In contrast, cultures infected with C-RTA had

limited viral gene expression at 6 hpi, but showed enhanced progression to both

TMER+RCA+actin^{low} and ORF73+TMER+RCA+actin^{low} phenotypes relative to WT

infection MOI=5 (Fig. 8F). Cultures infected with WT virus at an MOI=100 showed rapid

induction of multiple phenotypes, including a sizable portion of cultures which were

ORF73+RCA+ and ORF73+TMER+RCA+ by 6 hpi, both populations which retained

actin RNA expression. By 16 hpi, cultures infected with WT virus at an MOI=100 had

sizable frequencies of TMER+RCA+actin^{low} and ORF73+TMER+RCA+actin^{low} events

(Fig. 8F). While C-RTA and MOI=100 infected cultures showed enhanced viral gene

expression, WT cultures treated with phosphonoacetic acid (PAA), an inhibitor of viral

DNA synthesis, showed impaired expression of TMER with a complete absence of

TMER^{high} events, negligible ORF73 expression and a pronounced inability to induce an

actin^{low} phenotype (Supplemental Fig. 6). These data demonstrate diversity of viral gene

expression that can occur during lytic cycle and provide direct demonstration that these

different gene expression profiles can be experimentally manipulated by either

enhancing or restraining lytic replication.

DISCUSSION

Herpesvirus gene expression has been historically analyzed in bulk cell

populations. These studies have provided an essential cornerstone to understanding

the transcriptional and translational capacity of the herpesviruses. Despite this, recent

studies on cellular and viral transcription from other systems have emphasized a high

degree of cell-to-cell variation in gene expression [6-11, 13, 14], something we have

further investigated here. By applying the PrimeFlow^{TM} methodology to measure

endogenous viral gene expression across multiple gammaherpesviruses, and multiple

stages of infection, we have gained critical new insights into the inter-relationships of

gene expression at the single-cell level.

A primary focus of the current study has been to analyze expression of γHV

ncRNAs. Although the TMERs, EBERs and PAN RNA all represent abundant γHV

ncRNAs, these ncRNAs are transcribed by distinct mechanisms: KSHV PAN is a highly-

inducible, RNA pol II-transcribed ncRNA [20], in contrast to the RNA pol III-transcribed

TMERs and EBERs [15, 31]. This differential regulation was mirrored in the expression

patterns we observed. Whereas TMERs and EBERs were detected in a large fraction of

latently infected cells, PAN RNA was expressed in a low frequency of latently infected

cells, with prominent induction following cell stimulation and the induction of

reactivation. The viral ncRNAs were efficiently detected, as might be predicted due to

their abundance. The viral ORF73 encodes a transcription factor that is expressed at a

far lower level and are also efficiently detected, demonstrating that rare mRNAs can be

measured coincidently with abundant RNAs and with proteins, with no modifications

required. A unique advantage of our current approach is the ability to measure the

frequency of ncRNA expressing cells and changes in expression within individual cells.

This has been particularly insightful for the identification of rare PAN RNA+ cells in

untreated BCBL-1 cells and a TMER[high] subpopulation of cells in reactivating

A20.γHV68 cells. Integrating this method with cell sorting will afford future opportunities

to investigate unique properties of these rare cell populations.

Among the viral ncRNAs measured, in-depth analysis of TMER expression

during γHV68 infection has revealed new insights into infection. In the context of

latency, the TMERs are constitutively expressed in many, but not all, latently infected

cells using the A20.γHV68 model. Further, stimulating these cells to undergo

reactivation has a minimal effect on the frequency of cells expressing intermediate

levels of TMERs (i.e. TMER[mid] cells), instead resulting in the appearance of a minor

population of TMER[high] cells. Notably, TMER[high] cells show additional features of lytic

cycle progression, including actin RNA degradation and RCA protein expression. Why

only some latently infected cells show the TMER[high] phenotype, and what regulates the

inducible expression of the RNA pol III-transcribed TMERs remain important questions

raised by this analysis.

Of the γHVs studied here, only γHV68 has a robust in vitro lytic replication

system. Our studies on γHV68 lytic replication revealed multiple unanticipated results.

First, our analysis identified heterogeneity of viral gene expression, stratified by

differential viral gene expression of the TMERs and ORF73. Strikingly, in cultures

subjected to infection with 5 PFU per cell, conditions generally considered to induce

synchronous lytic replication, there were many cells with limited viral gene expression,

expressing low levels of either the TMERs and/or ORF73, but lacking additional signs of

virus gene expression (i.e. actin RNA degradation or RCA protein expression). This

analysis further identified a reproducible subset of cells with negligible viral gene

expression, defined by a TMER- ORF73- phenotype. Curiously, these cells contain viral

DNA but did not have further evidence of viral gene expression. Whether these cells

represent an abortive state of infection, or if viral genes are expressed at levels below

our current limit of detection remains to be determined. Our studies further revealed that

only some viral RNA+ cells showed full progression of virus infection characterized by

robust viral gene expression, defined as TMER^{high} ORF73^{high} RCA+ actin RNA^{low}.

Though progression to full viral gene expression was enhanced by infection of primary

fibroblasts, infection with the C-RTA virus, or a significantly higher MOI, even in these

conditions there remained heterogeneity in viral gene expression. Whether

heterogeneity of viral gene expression persists when cells manifest cytopathic effect, a

process which occurs after 16 hpi, is not yet known. While there is precedence that

reactivation from latency in KSHV infection can be asynchronous [9], this heterogeneity

of viral gene expression during in vitro lytic replication was unanticipated and suggests

that lytic infection under these reductionist conditions is either asynchronous, abortive,

or inefficient. This heterogeneity of gene expression raises important questions

regarding the universality of the prototypical cascade of immediate early, early and late

gene expression that is widely accepted in the herpesvirus field and suggests additional

levels of complexity that may be obscured by bulk cell analysis. The molecular

mechanisms that are responsible for this heterogeneity still remain to be elucidated but

appear to be independent of cell cycle stage with little to no contribution of type I

interferon receptor mediated signaling based on relatively comparable gene expression

profiles in wild-type and IFNAR1 KO MEFs.

This method allows multiplexed analysis of single-cell gene expression, to both

directly measure viral RNAs and downstream consequences of gene expression

including viral protein production and host RNA degradation, secondary to protein

translation. This approach has notable advantages to conventional analyses of gene

expression: 1) it can measure endogenous viral gene expression (both mRNA and

ncRNA) in the absence of recombinant viruses or marker genes, and 2) it can rapidly

analyze gene and protein expression inter-relationships, across millions of cells,

providing unique complementary strengths to other single-cell methodologies (e.g.

single-cell RNA-seq). In the future, this method can be further integrated with additional

antibody-based reagents, to simultaneously query post-translational modifications (e.g.

protein phosphorylation) as a function of cell cycle stage. It is also notable that through

the use of imaging flow cytometry, it is possible to interrogate subcellular RNA and

protein localization throughout distinct stages of virus infection, studies revealing single

cell variability in subcellular RNA localization. We anticipate that this approach will have

widespread utility, from addressing fundamental questions about herpesvirus gene

expression using reductionist approaches, to a better delineation of replication or

reactivation defects in viral mutants, to the in vivo identification of viral gene expression

in primary infected samples.

The approach presented here provides a powerful complementary method to

other single cell methods, affording the opportunity to query a diverse set of

experimental manipulations in a relatively rapid manner. It is worthwhile to note,

however, some important considerations with this approach. First, these studies rely on

fluorescent measurements of probe hybridization using a flow cytometer. Though this

approach detects a range of viral and host RNAs, sensitivity of this method is influenced

by target RNA expression level, fluorophores used for the analysis and controls to

define probe specificity and limit of detection. Ideally, comparisons can be strengthened

by comparing isogenic conditions (e.g. comparing WT virus with a genetically deficient

virus, as done with the TMER deficient virus). In cases where non-isogenic conditions

are used (e.g. comparing across cell lines), variable background fluorescence may limit

the sensitivity of this method. For new users of this technology, we strongly suggest the

use of multiple controls, including a full-minus-one (i.e. FMO) control to accurately

define background and signal to noise ratio for targets of interest. We consider

multiparameter flow cytometry, as well as mass cytometry [32, 33], two increasingly

useful technologies to afford new insight into heterogeneity of virus infection and gene

expression that provide a complementary approach to other single cell technologies.

In total, these studies demonstrate the power of single-cell analysis of

herpesvirus gene expression. Our data emphasize the heterogeneity of γHV gene

expression at the single-cell level, even in conditions considered to result in uniform

infection. The factors that underlie this heterogeneity are currently unknown, but could

reflect either asynchronous or inefficient infection in many infected cells (e.g. in the

context of lytic infection). The existence of specific infected cell subsets, based on

heterogeneous gene expression, may identify new susceptibilities and points for

intervention during the course of virus infection. Whether this variation arises from viral

or cellular heterogeneity is a fundamental question for future research.

MATERIALS AND METHODS

<u>Viruses and tissue culture</u>. γHV68 viruses were derived from the γHV68 strain WUMS

(ATCC VR-1465) [34], using either bacterial artificial chromosome-derived wild-type

(WT) γHV68 or γHV68.TMER-Total KnockOut (TMER-TKO) [16], or virus derived by

homologous recombination, γHV68.C-RTA [30] . Virus stocks were passaged, grown,

and titered as previously described [16]. Mouse 3T12 fibroblasts (ATCC CCL-164) were

infected with a multiplicity of infection (MOI) of either 5 or 100 plaque forming units/cell,

analyzed 6-18 hpi. Primary mouse embryonic fibroblasts isolated from C57BL/6 mice

(B6 MEFs) or from B6 IFNAR1 KO mice (IFNAR KO MEFs, kindly provided by Dr.

Thomas E. Morrison at the University of Colorado) were cultured in DMEM with 10%

FBS, 1% Penicillin/Streptomycin and then either mock- or virus-infected (MOI=5)

harvested at 16 hpi. For experiments using PAA, PAA (Sigma-Aldrich, catalog #284270)

was added at a concentration of 200 μg/mL at time of virus infection and left on cultures

until the time of harvest. The parental, virus-negative A20 B cell lymphoma cell line was

obtained from ATCC (ATCC® TIB-208™) and cultured in RPMI 1640 with 10% FBS, 1%

Penicillin/Streptomycin, L-glutamine and 50 μM β-mercaptoethanol (ME). γHV68

infected, and hygromycin selected A20.γHV68 (HE2.1) B cells [17] were obtained from

Dr. Sam Speck (Emory University) and cultured in RPMI 1640 with 10% FBS, 1%

Penicillin/Streptomycin + L-glutamine, 50 μM βME and 300 μg/mL Hygromycin B. A20

and A20.γHV68 B cells were treated with vehicle (untreated) or stimulated with 12-O-

tetradecanolphrobol-13-actate (TPA) 20 ng/ml (Sigma) (in DMSO) harvested 24 hr later.

BCBL-1 B cells, a body-cavity based B cell lymphoma cell line that is latently infected

with KSHV (HHV-8), were obtained from the NIH AIDS reagent program (catalog #

3233). BCBL-1 cells were cultured in RPMI containing 20% FBS, 1%

Penicillin/Streptomycin with L-glutamine, 1% HEPES and 50 µM βME. BCBL-1 B cells

were treated with vehicle (untreated) or stimulated with 20 ng/ml TPA (in DMSO) and

Sodium Butyrate (NaB) 0.3 mM (Calbiochem) (in water) and then harvested 72 hr later.

BL41 B cells (negative for KSHV and EBV), were cultured in RPMI with 10% FBS, 1%

Penicillin/Streptomycin with L-glutamine, and 50 µM βME. Mutu I cells, an EBV-infected,

type I latency Burkitt's lymphoma cell line [23] were obtained from Dr. Shannon Kenney

(University of Wisconsin), and cultured in RPMI with 10% FBS, 1%

Penicillin/Streptomycin and L-glutamine. Mutu I or BL41 B cells were either treated with

vehicle (DMSO) or stimulated with 20 ng/ml TPA (in DMSO) and then harvested 48 hr

later. EBV-immortalized B-lymphoblastoid cell lines (LCLs) were cultured in RPMI, 10%

FBS, 1% Penicillin/Streptomycin with L-glutamine. LCL3, LCL9 and LCL209 BM were

generated from Kenyan samples as previously described [35, 36].

<u>EBV infection of primary human B cells.</u> Peripheral blood was obtained from consenting

healthy adult donors and layered over Ficoll-Paque to isolate peripheral blood

mononuclear cells (PBMCs). B cells were isolated from PBMCs through negative

enrichment using EasySep Human B cell isolation kit (Stem Cell Technologies)

following manufacturer's protocol. B cells were plated at 1×10^6 cells per mL in RPMI,

10% FBS, 1% Penicillin/Streptomycin with L-glutamine, then infected with EBV at a MOI

of 10 genome copies per cell or mock infected. 5 days post infection cells were

harvested for flow cytometry analysis. EBV virus stocks were generated from the EBV+

cell line B95.8 which was reactivated with TPA (50:50 EtOH:Acetone) and Sodium

Butyrate (NaB) for 5 days. Supernatant was collected and centrifuged at 4,000xg for 10

min then passed over a 0.7 micron filter. The supernatant was then ultra-centrifuged at

16,000x g for 90 mins and resuspended in 1/200th the initial volume using RPMI, 10%

FBS, 1% Penicillin/Streptomycin with L-glutamine. Viral stocks were quantified following

DNase treatment with qPCR analysis of the EBV BALF 5 gene done as previously

described [37].

<u>Flow cytometric analysis.</u> Cells were harvested at the indicated time points and

processed for flow cytometry using the PrimeFlow^{TM} RNA Assay (Thermo Fisher).

Mouse cells were incubated with an Fc receptor blocking antibody (2.4G2) for 10 min

and then fixed with 2% PFA (Fisher), washed with PBS (Life Technology). Cells were

stained with a rabbit antibody against the γHV68 ORF4 protein, regulator of complement

activation (RCA) [19], labeled with Zenon R-phycoerythrin rabbit IgG label reagent (Life

Technologies) following manufacturer's protocol. Human B cells were incubated in

human Fc receptor blocking antibody then stained with Zombie Aqua fixable viability

dye (1:500 dilution) (BioLegend), according to manufacturer's protocol. Primary human

B cells were subsequently stained with anti-CD19-FITC (Clone HIB19, dilution 1:25),

and anti-CD69-PE antibody (clone FN50, dilution 1:20). Samples were subjected to the

PrimeFlow^{TM} RNA Assay following manufacturer's protocols, using viral and host target

probes conjugated to fluorescent molecules (Table S1). DAPI (BioLegend) was used on

a subset of samples following manufacturer's protocol, prior to PrimeFlow^{TM} probe

hybridization. Flow cytometric analysis was done on LSR II (BD Biosciences), Fortessa

(BD Biosciences), and ZE5 (Bio-Rad) flow cytometers, with compensation values based

on antibody-stained beads (BD Biosciences) and cross-validated using cell samples

stained with individual antibody conjugates, with compensation modified as needed

post-collection using FlowJo.

Flow cytometric analysis utilizing barcoding. Cells were harvested at the indicated time

points and processed for flow cytometry using the PrimeFlow[TM] RNA Assay (Thermo

Fisher). Cells were incubated with an Fc receptor blocking antibody (2.4G2), then fixed

and permeabilized using reagents from the PrimeFlow[TM] RNA Assay (Fixation Buffer 1

and Permeabilization Buffer). Samples from different experimental conditions were

fluorescently barcoded, with cells treated with either: no fluorescent dye, Ghost Dye[TM]

Violet 450 or Ghost Dye[TM] Violet 510 (Tonbo Biosciences, 1:200 dilution) according to

manufacturer's protocol. After washing, the three labeled samples were pooled together

into a single sample and stained with Zenon-labeled polyclonal rabbit antisera against

viral RCA as desribed above. Cells were fixed to cross-link antibody stain with Fixation

Buffer 2 from PrimeFlow[TM] RNA Assay, then subjected to the PrimeFlow[TM] RNA Assay

following manufacturer's protocols. For the analysis of barcoded samples, singlet cells

from barcoded samples were analyzed for fluorescence of either Ghost Dye[TM] Violet

450 or Ghost Dye[TM] Violet 510, to identify the three input populations: cells negative for

Ghost 450 and Ghost 510, cells singly positive for Ghost 450, and cell singly positive for

Ghost 510. To ensure that no artifacts were introduced due to assignment of one

barcode to a specific experimental condition, barcodes used for each experimental

condition were shuffled across independent samples.

Imaging flow cytometry. Cells were treated as described above then harvested, and split

into two aliquots: one for conventional flow cytometry, and one for imaging flow

cytometry, acquired on an Amnis ImageStream®^{X} Mark II imaging flow cytometer

(MilliporeSigma) with a 60X objective and low flow rate/high sensitivity using INSPIRE®

software. Brightfield (BF) and side scatter (SSC) images were illuminated by LED light

and a 785nm laser respectively. Fluorescent probes were excited off 405nm, 488nm,

and 642nm lasers with the power adjusted properly to avoid intensity saturation of the

camera. Single color controls for compensation were acquired by keeping the same

acquisition setting for samples, with the difference of turning the BF LED light and

785nm (SSC) laser off.

The acquired data were analyzed using IDEAS® software (MilliporeSigma).

Single cells that were in focus were defined as a population with a high "gradient RMS"

value, an intermediate "Area" value, and a medium to high "Aspect ratio" value for

subsequent analysis. Positive and negative events for each fluorescent marker were

determined using the "Intensity" feature. TMER nuclear localization was quantified using

"Similarity" feature, the log-transformed Pearson's correlation coefficient by analyzing

the pixel values of two image pairs [27]. The degree of nuclear localization of TMER

was measured by correlating the pixel intensity of two images with the same spatial

registry. The paired TMER and DAPI images were quantified by measuring the

"Similarity Score" which cells with high similarity scores display high TMER nuclear

localization with similar image pairs. By contrast cells with low similarity scores show

low TMER nuclear localization with dissimilar image pairs. Cell cycle stage analysis was

performed on data obtained from the imaging flow cytometer, stratifying cells based on

DAPI content to identify cells in the G0/G1 phase, S phase or G2/M phase of the cell

cycle.

Cell purification and DNA quantitation based on viral gene expression. Mouse

fibroblasts (3T12) were infected with WT γHV68 (MOI=5), harvested and processed for

PrimeFlow^{TM} at 16 hpi, followed by cell sorting using a BD FACSAria to purify cells

based on relative TMER and ORF73 expression. Post-sort purity checks on sorted

populations indicated that cell purities were 100% for TMER-ORF73- cells, 94.1% for

TMER+ ORF73- and 98% for TMER+ORF73+ cells. DNA was isolated from samples

using the DNeasy Blood and Tissue Kit (Qiagen), with an overnight proteinase K

incubation followed by heat inactivation (95°C for 10 min). DNA was precipitated using

ammonium acetate alcohol. 40 ng of DNA per sample was subjected to qPCR analysis

using LightCycler 480 Probe Master-Mix kit (Roche) and primer sets for γHV68 gB and

host NFAT5 (Table S3). gB standard curve was generated using a gB plasmid dilution

series ranging from 10^{10} to 10^{1} copies diluted in background DNA, with a limit of

detection (LOD) of 100 copies [38]. Host NFAT5 standard curve was generated using

HEK293 cells, with $2x10^{5}$ cell equivalents subjected to serial 10-fold dilutions in

background, salmon sperm DNA, from 10^{5} to 10^{1} copies (LOD=10 copies). Quantitation

of viral gB copy number was standardized relative to input material using the formula

(gB copy number / NFAT5 copy number) / 2, based on the assumption that cells have 2

copies of NFAT5 gene.

<u>KSHV genome quantification.</u> BCBL-1 or BL41 cells were plated at 7.5e5 cells/well in a

6 well plate with 20 ng/ml TPA and 0.3 mM NaB or vehicle only (DMSO and H_2O). Cells

and supernatant were harvested at 72 hrs post-treatment, hard-spun for 30 min at 4° C

and DNA was isolated using the DNeasy Blood and Tissue kit following manufacturer's

protocol, except for sample digestion for 1 hour instead of 10 min. 100 ng of DNA per

sample was used for qPCR analysis via SYBR green detection using KSHV ORF50

primers (5' -TCC GGC GGA TAT ACC GTC AC- 3' and 5'- GGT GCA GCT GGT ACA

GTG TG-3') [39]. qPCR was analyzed using relative quantification normalized against

unit mass calculation, ratio = $E^{deltaCt}$ (Real-Time PCR Application Guide, Bio-Rad

Laboratories Inc. 2006).

<u>RNA and qRT-PCR.</u> RNA was isolated using Trizol (Life Technologies) per

manufacturer's protocol and re-suspended in DEPC treated water. 3 μg of RNA was

treated with DNase 1 (Promega) for 2 hours at 37°C, heat inactivated for 10 min at

65°C. 500 ng of RNA was then subjected to reverse transcription using SuperScript II

(Life Technologies) following manufacturer's protocol for gene specific, oligo(dT), or

random primers (Life Technologies). Quantitative PCR (qPCR) was performed using iQ

SYBR Green super mix (Bio-Rad) follow manufacturer's protocol using host and viral

primer sets (Table S2) or using QuantiTech Primer Assay (Qiagen) for 18s (Hs-

RRN18S_1_SG). qPCR conditions: 3 min at 95°C, amplification cycles for 40 cycles of

15 sec at 95°C, annealing/ extension at temperature for specific primer set for 1 min

ending with a melt curve which started at 50°C or 55°C to 95°C increasing 0.5°C for

0:05 sec. A standard curve for each primer set was generated by pooling a portion of

each sample together and doing a 1:3 serial dilution. 75 ng of cDNA of the unknown

samples was loaded per qPCR reaction/primer set, with reactions run on a Bio-Rad 384

CFX LightCycler and data analyzed using Bio-Rad CFX manager software. Data

analysis was done using the 1:3 standard curve as the control Ct value to calculate the

delta ct, and the Pfaffl equation was used to define the fold difference between the gene

of interest and 18s (reference gene) [40]. qPCR products were analyzed by melt curve

analysis, with all reactions having a prominent, uniform product. In the case of primers

with an aberrant melt curve product (e.g. that arose at late cycles), products were

clearly a different product as defined by melt curve analysis.

<u>Software and Statistical analysis.</u> All flow cytometry data were analyzed in FlowJo

(version 8.8.7, 10.5.0, and 10.5.3), with flow cytometry data shown either as histogram

overlays or pseudo-color dot plots (with or without smoothing), showing outliers (low or

high resolution) on $\log_{10}$ scales. Statistical analysis and graphing were done in

GraphPad Prism (Version 6.0d and 7.0d). Statistical significance was tested by

unpaired t test (when comparing two conditions) or by one-way ANOVA (when

comparing three or more samples) subjected to multiple corrections tests using

recommended settings in Prism. *X-shift analysis:* For automated mapping of flow

cytometry data using X-shift, data were obtained from compensated flow cytometry files,

exported from FlowJo, using singlets that were live (defined by sequential gating on

single cells by FSC-H vs. FSC-W and SSC-H vs. SSC- W, that were DAPI bright vs.

SSC-A). These events were imported into the Java based program VorteX

(http://web.stanford.edu/~samusik/vortex/) [26]. Four parameters [TMER (AlexaFluor

(AF) 488), RCA (PE), ORF73 (AF647), and Actin (AF750)] were selected for clustering

analysis using the X-shift algorithm. The following settings were used when importing

the data set into VorteX: i) Numerical transformation: arcsinh(x/f), f=150, ii) noise

threshold: apply noise threshold of 1.0 (automatic and recommended setting), iii) feature

rescaling: none, and iv) normalization: none, v) a Euclidean noise filter was used with a

Minimal Euclidean length of the profile of 1.0, and vi) an import max of 1,000 rows from

each file after filtering was selected. The following settings were used when preparing

the data set for clustering analysis: i) distance measure: angular distance, ii) clustering

algorithm: X-shift (gradient assignment), iii) density estimate: N nearest neighbors (fast),

iv) number of neighbors for density estimate (K): from 150 to 5, with 30 steps, and v)

number of neighbors for mode finding (N): determine automatically. After the cluster

analysis was completed, all results were selected and the K value that corresponded

with optimal clustering (the elbow point) was calculated, in this case K= 50. All clusters

(seven clusters total) for the optimal K value were selected and a Force-Directed Layout

was created. The maximum number of events sampled from each cluster was 20, and

the number of nearest neighbors was 10. All settings used for this analysis were

automated or explicitly recommended (https://github.com/nolanlab/vortex/wiki). Force-

Directed layouts in Fig. 6A were saved as graphml files from VorteX, opened in the

application Gephi v 0.9.1, and colored by different variables (Cluster ID, experimental

group, Actin mRNA, RCA, ORF73, and TMERs respectively) in Adobe Illustrator CC

2017. Full details on use of the X-shfit algorithm and analysis pipeline can be found in

[41]. *tSNE analysis:* Gated events for each of the six identified populations were

exported from FlowJo, and then imported into Cytobank (www.cytobank.org) for

analysis using the viSNE algorithm. Each file was used for a separate viSNE analysis

(six total runs), where all available events were selected for clustering (202,669,

128,028, 29,096, 5,610, 8,956, 35,850 respectively) and four parameters were selected

for clustering (Actin mRNA, RCA, ORF73, and TMERs). The resulting tSNE plots were

colored according to expression using the "rainbow" color option, with individual events

shown using the stacked dot option. The channel range was user-defined for each

marker according to the range in expression established in Fig. 6E.

ACKNOWLEDGMENTS

The authors would like to acknowledge insightful comments made by members of the

Clambey and van Dyk laboratories, technical assistance by Eva Medina, pilot studies

establishing the feasibility of fluorescent barcoding by Shannon Rudolph, support of flow

cytometry services through ClinImmune, the Dept. of Immunology & Microbiology, and

the University of Colorado Cancer Center, and expert technical guidance from Matt

Cato and Dr. Nori Ueno (Thermo Fisher). T.C. and B.A. are employees of EMD Millipore

and Luminex respectively and have a potential conflict of interest. E.T.C. was a

recipient of the 2016 North America Affymetrix Single-cell Grant Recipient, for studies

unrelated to this manuscript.

FUNDING

This research was funded by National Institutes of Health grants R01CA103632 and

R01CA168558 to L.F.V.D., R21AI134084 to E.T.C. and L.F.V.D., and by an American

Heart Association National Scientist Development grant (#13SDG14510023), a

Colorado CTSI Novel methods development grant, and funding from the University of

Colorado Dept. of Anesthesiology to E.T.C.. The Colorado CTSI is supported by

NIH/NCATS Colorado CTSA Grant Number UL1 TR002535. Contents are the authors'

sole responsibility and do not necessarily represent official NIH views. The funders had

no role in study design, data collection and analysis, decision to publish, or preparation

of the manuscript.

AUTHOR CONTRIBUTIONS

<u>Conceptualization:</u> Eric T. Clambey, Linda F. van Dyk

<u>Data curation:</u> Lauren M. Oko, Eric T. Clambey

<u>Formal analysis:</u> Lauren M. Oko, Abigail K. Kimball, Rachael E. Kaspar, Benjamin

Alderete, Tim Chang, Eric T. Clambey

<u>Funding acquisition:</u> Eric T. Clambey, Linda F. van Dyk

<u>Investigation:</u> Lauren M. Oko, Ashley N. Knox, Carrie B. Coleman

<u>Methodology:</u> Lauren M. Oko, Carrie B. Coleman, Rosemary Rochford, Benjamin

Alderete, Tim Chang, Linda F. van Dyk, Eric T. Clambey

<u>Project administration:</u> Eric T. Clambey, Linda F. van Dyk

<u>Resources:</u> Lauren M. Oko, Rosemary Rochford, Benjamin Alderete, Tim Chang

<u>Software:</u> Abigail K. Kimball

<u>Supervision:</u> Eric T. Clambey, Linda F. van Dyk

<u>Validation:</u> Lauren M. Oko, Eric T. Clambey

<u>Visualization:</u> Lauren M. Oko, Abigail K. Kimball, Rachael E. Kaspar, Benjamin

Alderete, Tim Chang, Eric T. Clambey

<u>Writing – original draft:</u> Eric T. Clambey, Linda F. van Dyk

<u>Writing – review & editing:</u> Lauren M. Oko, Linda F. van Dyk, Eric T. Clambey

REFERENCES

1. Pellett PJ, Roizman B. *Herpesviridae.* Fields Virology. 6th ed2013.

2. Longnecker RM, Kieff, E., Cohen, J.I. Epstein-Barr Virus. Fields Virology. 6th ed2013.

3. Ganem D. KSHV infection and the pathogenesis of Kaposi's sarcoma. Annu Rev Pathol. 2006;1:273-96. doi: 10.1146/annurev.pathol.1.110304.100133. PubMed PMID: 18039116.

4. Barton E, Mandal P, Speck SH. Pathogenesis and host control of gammaherpesviruses: lessons from the mouse. Annual review of immunology. 2011;29:351-97. doi: 10.1146/annurev-immunol-072710-081639. PubMed PMID: 21219186.

5. Cesarman E. Gammaherpesviruses and lymphoproliferative disorders. Annu Rev Pathol. 2014;9:349-72. doi: 10.1146/annurev-pathol-012513-104656. PubMed PMID: 24111911.

6. Shalek AK, Satija R, Adiconis X, Gertner RS, Gaublomme JT, Raychowdhury R, et al. Single-cell transcriptomics reveals bimodality in expression and splicing in immune cells. Nature. 2013;498(7453):236-40. doi: 10.1038/nature12172. PubMed PMID: 23685454; PubMed Central PMCID: PMCPMC3683364.

7. Battich N, Stoeger T, Pelkmans L. Control of Transcript Variability in Single Mammalian Cells. Cell. 2015;163(7):1596-610. doi: 10.1016/j.cell.2015.11.018. PubMed PMID: 26687353.

8. Chen KH, Boettiger AN, Moffitt JR, Wang S, Zhuang X. RNA imaging. Spatially resolved, highly multiplexed RNA profiling in single cells. Science.

2015;348(6233):aaa6090. doi: 10.1126/science.aaa6090. PubMed PMID: 25858977;

PubMed Central PMCID: PMCPMC4662681.

9.   Adang LA, Parsons CH, Kedes DH. Asynchronous progression through the lytic

cascade and variations in intracellular viral loads revealed by high-throughput single-cell

analysis of Kaposi's sarcoma-associated herpesvirus infection. J Virol.

2006;80(20):10073-82. doi: 10.1128/JVI.01156-06. PubMed PMID: 17005685; PubMed

Central PMCID: PMCPMC1617294.

10.   Borah S, Nichols LA, Hassman LM, Kedes DH, Steitz JA. Tracking expression

and subcellular localization of RNA and protein species using high-throughput single

cell imaging flow cytometry. RNA. 2012;18(8):1573-9. doi: 10.1261/rna.033126.112.

PubMed PMID: 22745225; PubMed Central PMCID: PMCPMC3404377.

11.   Ma JZ, Russell TA, Spelman T, Carbone FR, Tscharke DC. Lytic gene

expression is frequent in HSV-1 latent infection and correlates with the engagement of a

cell-intrinsic transcriptional response. PLoS Pathog. 2014;10(7):e1004237. doi:

10.1371/journal.ppat.1004237. PubMed PMID: 25058429; PubMed Central PMCID:

PMCPMC4110040.

12.   Shnayder M, Nachshon A, Krishna B, Poole E, Boshkov A, Binyamin A, et al.

Defining the Transcriptional Landscape during Cytomegalovirus Latency with Single-

Cell RNA Sequencing. MBio. 2018;9(2). Epub 2018/03/15. doi: 10.1128/mBio.00013-18.

PubMed PMID: 29535194; PubMed Central PMCID: PMCPMC5850328.

13.   Porichis F, Hart MG, Griesbeck M, Everett HL, Hassan M, Baxter AE, et al. High-

throughput detection of miRNAs and gene-specific mRNA at the single-cell level by flow

cytometry. Nat Commun. 2014;5:5641. doi: 10.1038/ncomms6641. PubMed PMID:

25472703; PubMed Central PMCID: PMCPMC4256720.

14. Baxter AE, Niessl J, Fromentin R, Richard J, Porichis F, Charlebois R, et al.

Single-Cell Characterization of Viral Translation-Competent Reservoirs in HIV-Infected

Individuals. Cell Host Microbe. 2016;20(3):368-80. doi: 10.1016/j.chom.2016.07.015.

PubMed PMID: 27545045; PubMed Central PMCID: PMCPMC5025389.

15. Diebel KW, Claypool DJ, van Dyk LF. A conserved RNA polymerase III promoter

required for gammaherpesvirus TMER transcription and microRNA processing. Gene.

2014;544(1):8-18. doi: 10.1016/j.gene.2014.04.026. PubMed PMID: 24747015.

16. Diebel KW, Oko LM, Medina EM, Niemeyer BF, Warren CJ, Claypool DJ, et al.

Gammaherpesvirus small noncoding RNAs are bifunctional elements that regulate

infection and contribute to virulence in vivo. MBio. 2015;6(1):e01670-14. doi:

10.1128/mBio.01670-14. PubMed PMID: 25691585; PubMed Central PMCID:

PMCPMC4337559.

17. Forrest JC, Speck SH. Establishment of B-cell lines latently infected with

reactivation-competent murine gammaherpesvirus 68 provides evidence for viral

alteration of a DNA damage-signaling cascade. J Virol. 2008;82(15):7688-99. doi:

10.1128/JVI.02689-07. PubMed PMID: 18495760; PubMed Central PMCID:

PMCPMC2493333.

18. Cheng BY, Zhi J, Santana A, Khan S, Salinas E, Forrest JC, et al. Tiled

microarray identification of novel viral transcript structures and distinct transcriptional

profiles during two modes of productive murine gammaherpesvirus 68 infection. J Virol.

2012;86(8):4340-57. doi: 10.1128/JVI.05892-11. PubMed PMID: 22318145; PubMed

Central PMCID: PMCPMC3318610.

19.   Kapadia SB, Molina H, van Berkel V, Speck SH, Virgin HWt. Murine

gammaherpesvirus 68 encodes a functional regulator of complement activation. J Virol.

1999;73(9):7658-70. PubMed PMID: 10438856; PubMed Central PMCID: PMC104293.

20.   Sun R, Lin SF, Gradoville L, Miller G. Polyadenylylated nuclear RNA encoded by

Kaposi sarcoma-associated herpesvirus. Proceedings of the National Academy of

Sciences of the United States of America. 1996;93(21):11883-8. Epub 1996/10/15.

PubMed PMID: 8876232; PubMed Central PMCID: PMCPMC38153.

21.   Marchini A, Longnecker R, Kieff E. Epstein-Barr virus (EBV)-negative B-

lymphoma cell lines for clonal isolation and replication of EBV recombinants. J Virol.

1992;66(8):4972-81. Epub 1992/08/01. PubMed PMID: 1321281; PubMed Central

PMCID: PMCPMC241347.

22.   Cao S, Strong MJ, Wang X, Moss WN, Concha M, Lin Z, et al. High-throughput

RNA sequencing-based virome analysis of 50 lymphoma cell lines from the Cancer Cell

Line Encyclopedia project. J Virol. 2015;89(1):713-29. doi: 10.1128/JVI.02570-14.

PubMed PMID: 25355872; PubMed Central PMCID: PMC4301145.

23.   Gregory CD, Rowe M, Rickinson AB. Different Epstein-Barr virus-B cell

interactions in phenotypically distinct clones of a Burkitt's lymphoma cell line. The

Journal of general virology. 1990;71 (Pt 7):1481-95. Epub 1990/07/01. doi:

10.1099/0022-1317-71-7-1481. PubMed PMID: 2165133.

24.   Covarrubias S, Richner JM, Clyde K, Lee YJ, Glaunsinger BA. Host shutoff is a

conserved phenotype of gammaherpesvirus infection and is orchestrated exclusively

from the cytoplasm. J Virol. 2009;83(18):9554-66. doi: 10.1128/JVI.01051-09. PubMed

PMID: 19587049; PubMed Central PMCID: PMC2738246.

25.   Rowe M, Glaunsinger B, van Leeuwen D, Zuo J, Sweetman D, Ganem D, et al.

Host shutoff during productive Epstein-Barr virus infection is mediated by BGLF5 and

may contribute to immune evasion. Proceedings of the National Academy of Sciences

of the United States of America. 2007;104(9):3366-71. doi: 10.1073/pnas.0611128104.

PubMed PMID: 17360652; PubMed Central PMCID: PMC1805610.

26.   Samusik N, Good Z, Spitzer MH, Davis KL, Nolan GP. Automated mapping of

phenotype space with single-cell data. Nature methods. 2016;13(6):493-6. doi:

10.1038/nmeth.3863. PubMed PMID: 27183440; PubMed Central PMCID:

PMCPMC4896314.

27.   Maguire O, Collins C, O'Loughlin K, Miecznikowski J, Minderman H. Quantifying

nuclear p65 as a parameter for NF-kappaB activation: Correlation between

ImageStream cytometry, microscopy, and Western blot. Cytometry A. 2011;79(6):461-9.

doi: 10.1002/cyto.a.21068. PubMed PMID: 21520400; PubMed Central PMCID:

PMCPMC3140714.

28.   Amir el AD, Davis KL, Tadmor MD, Simonds EF, Levine JH, Bendall SC, et al.

viSNE enables visualization of high dimensional single-cell data and reveals phenotypic

heterogeneity of leukemia. Nat Biotechnol. 2013;31(6):545-52. Epub 2013/05/21. doi:

10.1038/nbt.2594. PubMed PMID: 23685480; PubMed Central PMCID:

PMCPMC4076922.

29.   Weck KE, Barkon ML, Yoo LI, Speck SH, Virgin HI. Mature B cells are required

for acute splenic infection, but not for establishment of latency, by murine

gammaherpesvirus 68. J Virol. 1996;70(10):6775-80. PubMed PMID: 8794315; PubMed

Central PMCID: PMCPMC190721.

30.   Rickabaugh TM, Brown HJ, Martinez-Guzman D, Wu TT, Tong L, Yu F, et al.

Generation of a latency-deficient gammaherpesvirus that is protective against

secondary infection. J Virol. 2004;78(17):9215-23. Epub 2004/08/17. doi:

10.1128/JVI.78.17.9215-9223.2004. PubMed PMID: 15308716; PubMed Central

PMCID: PMCPMC506911.

31.   Rosa MD, Gottlieb E, Lerner MR, Steitz JA. Striking similarities are exhibited by

two small Epstein-Barr virus-encoded ribonucleic acids and the adenovirus-associated

ribonucleic acids VAI and VAII. Mol Cell Biol. 1981;1(9):785-96. Epub 1981/09/01.

PubMed PMID: 9279391; PubMed Central PMCID: PMCPMC369362.

32.   Frei AP, Bava FA, Zunder ER, Hsieh EW, Chen SY, Nolan GP, et al. Highly

multiplexed simultaneous detection of RNAs and proteins in single cells. Nature

methods. 2016;13(3):269-75. Epub 2016/01/26. doi: 10.1038/nmeth.3742. PubMed

PMID: 26808670; PubMed Central PMCID: PMCPMC4767631.

33.   Duckworth AD, Gherardini PF, Sykorova M, Yasin F, Nolan GP, Slupsky JR, et

al. Multiplexed profiling of RNA and protein expression signatures in individual cells

using flow or mass cytometry. Nat Protoc. 2019;14(3):901-20. Epub 2019/02/08. doi:

10.1038/s41596-018-0120-8. PubMed PMID: 30728478.

34.   Virgin HWt, Latreille P, Wamsley P, Hallsworth K, Weck KE, Dal Canto AJ, et al.

Complete sequence and genomic analysis of murine gammaherpesvirus 68. J Virol.

1997;71(8):5894-904. PubMed PMID: 9223479; PubMed Central PMCID: PMC191845.

35.   Simbiri KO, Smith NA, Otieno R, Wohlford EE, Daud, II, Odada SP, et al.

Epstein-Barr virus genetic variation in lymphoblastoid cell lines derived from Kenyan

pediatric population. PLoS One. 2015;10(5):e0125420. Epub 2015/05/02. doi:

10.1371/journal.pone.0125420. PubMed PMID: 25933165; PubMed Central PMCID:

PMCPMC4416826.

36.   Daud, II, Coleman CB, Smith NA, Ogolla S, Simbiri K, Bukusi EA, et al. Breast

Milk as a Potential Source of Epstein-Barr Virus Transmission Among Infants Living in a

Malaria-Endemic Region of Kenya. J Infect Dis. 2015;212(11):1735-42. Epub

2015/05/20. doi: 10.1093/infdis/jiv290. PubMed PMID: 25985902; PubMed Central

PMCID: PMCPMC4633760.

37.   Coleman CB, Wohlford EM, Smith NA, King CA, Ritchie JA, Baresel PC, et al.

Epstein-Barr virus type 2 latently infects T cells, inducing an atypical activation

characterized by expression of lymphotactic cytokines. J Virol. 2015;89(4):2301-12.

Epub 2014/12/17. doi: 10.1128/JVI.03001-14. PubMed PMID: 25505080; PubMed

Central PMCID: PMCPMC4338898.

38.   Niemeyer BF, Oko LM, Medina EM, Oldenburg DG, White DW, Cool CD, et al.

Host Tumor Suppressor p18(INK4c) Functions as a Potent Cell-Intrinsic Inhibitor of

Murine Gammaherpesvirus 68 Reactivation and Pathogenesis. J Virol. 2018;92(6).

Epub 2018/01/05. doi: 10.1128/JVI.01604-17. PubMed PMID: 29298882; PubMed

Central PMCID: PMCPMC5827403.

39.   Li Q, He M, Zhou F, Ye F, Gao SJ. Activation of Kaposi's sarcoma-associated

herpesvirus (KSHV) by inhibitors of class III histone deacetylases: identification of

sirtuin 1 as a regulator of the KSHV life cycle. J Virol. 2014;88(11):6355-67. Epub

2014/03/29. doi: 10.1128/JVI.00219-14. PubMed PMID: 24672028; PubMed Central

PMCID: PMCPMC4093851.

40.   Pfaffl MW. A new mathematical model for relative quantification in real-time RT-

PCR. Nucleic Acids Res. 2001;29(9):e45. PubMed PMID: 11328886; PubMed Central

PMCID: PMCPMC55695.

41.   Kimball AK, Oko LM, Bullock BL, Nemenoff RA, van Dyk LF, Clambey ET. A

Beginner's Guide to Analyzing and Visualizing Mass Cytometry Data. J Immunol.

2018;200(1):3-22. Epub 2017/12/20. doi: 10.4049/jimmunol.1701494. PubMed PMID:

29255085; PubMed Central PMCID: PMCPMC5765874.

FIGURE LEGENDS

Figure 1. RNA-Flow cytometry using the PrimeFlow*^{TM} *method affords robust and

sensitive analysis of endogenous γHV genes at the single-cell level. Viral RNA

analysis in γHV68-infected fibroblasts by qRT-PCR (A, C) or by flow cytometric analysis

using PrimeFlow^{TM} at 16 hpi (B, D, E, F). Samples were either mock, TMER-TKO, or

WT γHV68-infected, with infections done using 5 plaque forming units/cell. qRT-PCR

standardized to 18s RNA, at the indicated times. All flow cytometric events gated on a

generous FSC x SSC gate, followed by singlet discrimination. PrimeFlow^{TM} analysis

quantified probe fluorescence for (B) TMERs, (D) ORF73, or (E) ORF18, relative to

side-scatter area (SSC-A). Probe fluorescence is indicated, with all probes detected

using either AlexaFluor (AF) 647 or 488 conjugates. Data representative of 2

independent experiments, each done with biological replicates.

Figure 2. Heterogeneous gene expression in a γHV68 latently infected B cell line.

Viral, host RNA analysis during γHV68 latency and reactivation in A20.γHV68 (HE2.1)

cells by qRT-PCR (A) and flow cytometric analysis using PrimeFlow^{TM} (B-G), comparing

untreated or TPA-stimulated samples at 24 hrs post-treatment. Analysis includes A20,

virus-negative cells and A20.γHV68 cells. (A) qRT-PCR analysis of TMER-5 expression

relative to 18s RNA in A20 and A20.γHV68 (HE2.1) cells, untreated or stimulated with

TPA for 24 hrs. (B) PrimeFlow^{TM} detection of TMER expression in A20.γHV68 (HE2.1)

cells, comparing either samples that were unstained (solid gray) or stained for the

TMERs (open black line). (C) Analysis of TMER expression in multiple conditions,

comparing untreated and stimulated A20 and A20.γHV68 cells, with gates defining the

frequency of events that expressed either intermediate (mid) or high levels of TMERs.

Data depict lymphocytes that were singlets, defined by sequential removal of doublets.

(D) Quantification of the frequency of TMER^{high} cells in stimulated A20.γHV68 cells. (E)

Histogram overlays of HygroGFP and RCA protein expression in A20.γHV68 cells

comparing TMER^{mid} cells from untreated cultures (top, gray), TMER^{mid} cells from TPA-

stimulated cultures (middle, blue), with TMER^{high} cells from stimulated cultures (bottom,

red). (F) Quantification of the frequencies of HygroGFP+ (left) and RCA protein+ (right)

cells as a function of TMER expression and treatment condition. (G) Flow cytometric

analysis on an imaging flow cytometer, with each row showing an individual cell and

representative images of brightfield (BF), RCA protein (RCA), DAPI, and TMER

localization in TMER^{mid} cells from untreated cultures (left) or TMER^{high} RCA+ cells from

stimulated (right) A20.γHV68 cells. Data are from two independent experiments, with

biological replicates within each experiment for all A20.γHV68 cultures. Graphs depict

the mean ± SEM, with each symbol identifying data from a single replicate. Statistical

analysis was done using an unpaired t test (D) or one-way ANOVA, subjected to

Tukey's multiple comparison test (A, F), with statistically significant differences as

indicated, * $p<0.05$, **** $p<0.0001$.

Figure 3. Single-cell analysis of KSHV PAN RNA expression in the BCBL-1 B cell

lymphoma cell line. (A) Flow cytometric analysis of PAN RNA expression in multiple

conditions, from cells incubated with no probe (left), or cells subjected to hybridization

using a probe for PAN RNA, comparing virus-negative BL41 cells (second from left) with

KSHV+ BCBL-1 cells that were either untreated or stimulated (with TPA and sodium

butyrate (NaB)) for 72 hours. Data depict lymphocytes that were singlets, defined by

sequential removal of doublets. Representative images were defined as samples that

were closest to the median frequency. Quantification of (B) the frequency of PAN RNA+

cells and (C) PAN RNA median fluorescence within PAN RNA+ cells, comparing

untreated or stimulated BCBL-1 cells. D, E) Flow cytometric analysis of untreated

BCBL-1 cells, using histogram overlays, to compare cell size (forward scatter, FSC),

granularity (side scatter, SSC) and ORF73 expression in cells that were either PAN

RNA negative [-] (blue line) or PAN RNA positive [+] (red line). Data show (D) histogram

overlays of these populations with fluorescence quantification provided in panel E.

ORF73 analysis includes samples in which there was no ORF73 probe (i.e. "No Probe",

in solid gray), to define background fluorescence. (F, G) Flow cytometric analysis of

stimulated BCBL-1 cells, using histogram overlays, to compare cell size (forward

scatter, FSC), granularity (side scatter, SSC) and ORF73 expression in cells that are

either PAN RNA negative [-] (blue) or PAN RNA positive [+] (red). Data show (F)

histogram overlays of these populations with fluorescence quantification provided in

panel G, with ORF73 analysis including a "No Probe" sample (gray) to define

background fluorescence. Due to variable baseline fluorescence values for SSC and

ORF73 between experiments, values were internally standardized to fluorescent

intensities within the PAN RNA negative population for each experiment, with data

depicting mean ± SEM. Symbols in panels B and C indicate values from individual

samples. Data are from two independent experiments, with biological replicates within

each experiment, with total number of biological replicates as follows: No Probe (n = 2),

BL41 control (n = 3), BCBL-1 untreated (n = 6), BCBL-1 stimulated (n = 6). Statistical

analysis was done using an unpaired t test with statistically significant differences as

indicated, * p<0.05, ** p<0.01, **** p<0.0001.

1017

Figure 4. Single-cell analysis of EBV EBER expression in the Mutu I B cell

lymphoma cell line. (A) Flow cytometric analysis using the PrimeFlow[TM] method, to

quantify EBER expression in multiple conditions, from cells incubated with no probe

(left), or cells subjected to hybridization using a probe for the EBERs, comparing virus-

negative BL41 cells (second from left) with Mutu I EBV+ cells that were either untreated

(with DMSO) or stimulated (with TPA in DMSO) for 48 hours. Data depict lymphocytes

that were singlets, defined by sequential removal of doublets. Representative images

were defined as samples that were closest to the median frequency, with data depicting

mean ± SEM. (B) Quantitation of median EBER fluorescence within EBER+ Mutu I cells

in untreated and stimulated cultures. Horizontal dashed line indicates the background

fluorescent signal from BL41 controls. Data are from two independent experiments, with

biological replicates within each experiment, with total biological replicates as follows:

No Probe (n = 2), BL41 Control untreated (n = 6), Mutu I untreated (n = 6), Mutu I

stimulated (n = 6). Statistical analysis was done using an unpaired t test with statistically

significant differences as indicated, * p<0.05.

1033

Figure 5. Actin mRNA degradation identifies virally-infected cells experiencing

virus-induced host shutoff. Actin mRNA analysis by qRT-PCR (A,D) or by flow

cytometric analysis using PrimeFlow[TM] (B,C,E-G), comparing cells with variable

infection status. (A) qRT-PCR analysis of beta-actin (Actb) mRNA expression relative to

18s RNA in mock, WT or TMER-TKO infected 3T12 fibroblasts at 18 hpi. (B)

PrimeFlow[TM] analysis of actin mRNA in 3T12 fibroblasts, either unstained ("No probe"),

mock-infected or infected with WT γHV68 or TMER-TKO at 18 hpi. (C) PrimeFlow[TM]

analysis of actin mRNA and TMER expression in WT γHV68 infected fibroblasts at 18

hpi. (D) qRT-PCR analysis of beta-actin (Actb) mRNA expression relative to 18s RNA

in A20, virus-negative cells and A20.γHV68 (HE2.1) cells, untreated or stimulated with

TPA for 24 hrs. (E) PrimeFlow[TM] analysis of actin mRNA and TMER expression in

untreated and stimulated A20.γHV68 cells, with the frequency of TMER[high] actin

mRNA[low] cells indicated, based on the gated events. (F,G) Actin mRNA analysis by

PrimeFlow[TM] using either (F) histogram overlays or (G) quantifying frequencies,

comparing A20.γHV68 cells that were either untreated or stimulated, further stratified by

whether the cells were TMER[mid] or TMER[high] (using the gating strategy defined in Fig.

2C). All flow cytometry data depict single cells, defined by sequential removal of

doublets. Data are from two-three independent experiments, with biological replicates

within each experiment. Graphs depict the mean ± SEM, with each symbol identifying

data from a single replicate. Statistical analysis was done using one-way ANOVA,

subjected to Tukey's multiple comparison test (A, D, G), with statistically significant

differences as indicated, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

1056

Figure 6. Heterogeneous viral gene expression at the single-cell level during lytic

replication. Viral, host RNA flow cytometric analysis in γHV68-infected fibroblasts at 16

hpi defined by the PrimeFlow[TM] method, comparing (A) X-shift clustering analysis and

(B-E) biaxial gating analysis for the indicated features. (A) Automated, clustering

analysis using the X-Shift algorithm on 10,000 events total, compiled from mock- and

γHV68-infected fibroblasts at 16 hrs pi (1,000 events randomly imported per sample,

mock infected n=4, γHV68-infected n=6) identifies multiple clusters of cells with

differential gene expression (7 clusters, colored distinctly, "Cluster ID"), with these

clusters then depicted for expression of TMERs, ORF73, Actin mRNA, and RCA. Range

of expression is identified for each parameter. (B) Analysis of TMER and ORF73 co-

expression in mock (left), WT γHV68-infected (middle), and TMER-TKO-infected (right)

samples, with gates depicting populations with different gene expression profiles,

defined relative to mock and TMER-TKO infected samples. (C) Color-coded populations

from WT-infected sample in panel B, with each color indicating a different gene

expression profile. (D) Histogram overlays of the five populations identified in panel C

for the indicated parameters. (E) Quantitation of gene expression among the five

populations identified in panel C, using the same color-coding strategy. Data are from

three independent experiments, with each experiment containing biological replicates.

Flow cytometry data shows single cells that are DNA+ (DAPI+). Statistical significance

tested by one-way ANOVA, comparing the mean of TMER^{high} ORF73^{high} cells to all

other means, followed by Dunnett's multiple testing correction. Significance identified as

*** p<0.001, **** p<0.0001.

1079

Figure 7. Lytic replication is characterized by heterogeneous TMER localization

and variable penetrance of actin RNA degradation. Viral, host RNA flow cytometric

analysis in γHV68-infected fibroblasts at 16 hpi defined by the PrimeFlow^{TM} method. (A)

The frequency of γHV68-infected fibroblasts with TMERs primarily in the nucleus was

quantified by ImageStream, with data showing the frequency of cells in which

TMER:DAPI colocalization (i.e. similarity score) was >1. (B) Images showing brightfield

(BF), TMER, RCA protein (RCA), DAPI, ORF73 and actin mRNA localization,

comparing cells with nuclear TMER localization (left) versus cytoplasmic TMER

localization (right). (C) Analysis of cell subpopulations stratified by TMER and ORF73

expression (defined in Fig. 6C), subjected to the tSNE dimensionality reduction

algorithm. Data and cell populations are derived from the dataset presented in Fig. 6,

showing all DNA+ (DAPI+) single cells (FSC-A, SSC-A) subjected to the tSNE

algorithm. The tSNE algorithm provides each cell with a unique coordinate according to

its expression of Actin mRNA, RCA, ORF73, and TMERs, displayed on a two-

dimensional plot (tSNE1 versus tSNE2). Visualization grid of tSNE plots, with plots

arranged according to marker expression (rows) relative to phenotype of the cellular

population examined (columns). (D) Biaxial analysis of actin RNA versus RCA protein,

among the five populations identified in Fig. 6. Flow cytometry data shows single cells

that are DNA+ (DAPI+). Data in panels A-B from two independent experiments, in panel

C-D from three independent experiments.

1100

Figure 8. Lytic viral gene expression is influenced by target cell type, changes

over time, is enhanced by conditions which promote lytic replication. Fibroblasts

(3T12 cells or mouse embryonic fibroblasts, MEFs) were infected with WT or C-RTA

γHV68 at 5 or 100 PFU/cell, and harvested at either 6 or 16 hpi as indicated. (A-C)

Analysis of viral gene expression by PrimeFlow™ in either 3T12 or MEFs following WT

γHV68 infection (MOI=5) and harvested at 16 hpi. (A) Analysis of viral gene expression

by PrimeFlow^{TM}, comparing gene expression by histogram overlays between 3T12, WT

MEFs and IFNAR KO MEFs. (B) The frequency of events stratified based on expression

of ORF73 and TMER, or (C) based on all 16 possible combinations of gene expression

with the frequency of events in each population plotted as a heatmap. Data are from

three independent experiments harvested at the indicated time, subjected to fluorescent

barcoding, pooled together and subjected to PrimeFlow^{TM} analysis in a single analysis.

(D-F) 3T12 fibroblasts were infected with WT or C-RTA γHV68 at 5 or 100 PFU/cell, and

harvested at either 6 or 16 hpi as indicated. (D) Analysis of viral gene expression by

PrimeFlow^{TM}, comparing gene expression by histogram overlays between different virus

infection conditions and times post-infection, as indicated. (E) The frequency of events

stratified based on expression of ORF73 and TMER, or (F) based on all 16 possible

combinations of gene expression with the frequency of events in each population

plotted as a heatmap. Data are from three independent experiments harvested at the

indicated time, subjected to fluorescent barcoding, pooled together and subjected to

PrimeFlow^{TM} analysis in a single analysis. Data depict mean values, with SEM plotted in

panels B, D, and heatmap values colored based on the categorical key provided in the

figure. Flow cytometry data depict single cells, defined by sequential removal of

doublets according to SSC-A x SSC-H and FSC-A x FSC-H, after which files were

subjected to debarcoding to identify each of the independent samples in the analysis.

1126

1127

 SUPPORTING INFORMATION LEGENDS

 Supplemental Table 1. Viral and host probes for PrimeFlow^{TM} analysis.

 Supplemental Table 2. qPCR primer sets.

 Supplemental Table 3. TaqMan PCR probe sets used in this study.

 Supplemental Figure 1. Analysis of KSHV genomic DNA in BCBL-1 cells either

 untreated or stimulated. Data depict qPCR analysis of KSHV genomic DNA,

 quantified using PCR primers for the KSHV gene 50. Fold change in KSHV DNA was

 calculated relative to untreated BCBL-1 cells, using ΔCt normalized to mass unit,

 defined using 100 ng DNA. Data compiled from 3 independent experiments for BCBL-1

 cultures, with 3 biological replicates per experiment. Data from BL-41 cells derived from

 a single experiment, with 3 biological replicates. Data depict mean +/- SEM, with

 individual biological replicates depicted using individual symbols. Statistical analysis of

 the difference between untreated and stimulated samples was done using an unpaired t

 test, with Welch's correction, with statistical significance defined by ***, $p<0.001$.

 Supplemental Figure 2. Analysis of EBER expression by PrimeFlow^{TM} in a panel

 of human B cell lines. (A) Comparison of fluorescence in BL41 (EBV-) and Mutu I

 (EBV+) cells, where cell lines were either incubated with no probe or an EBER probe.

 The three populations identified on these plots, indicated by 3 polygons with black lines,

correspond to: left, the true negative population; middle, background fluorescence

observed in BL41 cells stained with EBER probes; right, EBER+ events defined by

expression above these two different thresholds. (B) Quantitation of the frequency of

EBER+ events among viable cells across all the cell lines analyzed, with symbols

depicting individual replicate data, bars showing mean ± SEM. (C) Analysis of EBER

expression in three different LCL cultures, as indicated, comparing background

fluorescence (No probe) with fluorescence following EBER probe hybridization. (D)

Gating hierarchy to analyze EBER fluorescence in viable cells. All flow cytometry plots

show events defined as lymphocytes by forward and side scatter, doublet

discrimination, and viable cells defined by exclusion of a viability dye. Data are from a

single experiment with one to three biological replicates (n=1 Mutu I, n=3 for BL41 and

LCLs).

1163

Supplemental Figure 3. Analysis of EBER expression by PrimeFlow^{TM} in human

primary B cells subjected to in vitro EBV infection. (A) Comparison of EBER

expression in human primary B cells subjected to either mock or EBV infection (10

genome copies/cell) for 5 days. Cells were either incubated with a cocktail of antibodies

and the EBER probe, or with specific exclusion of the EBER probe (i.e. a full minus one

control, No probe). Data depict the frequency of EBER+ events within lymphocytes that

were singlets and CD19+ B cells. Data are from a single experiment. (B) Comparison of

EBER expression in human primary B cells subjected to either mock or EBV infection

(10 genome copies/cell) for 5 days. Data depict the frequency of EBER+ events within

lymphocytes that were viable, singlets, and CD19+ B cells. (C) Quantitation of the

frequency of EBER+ cells among viable B cells in mock or EBV infected cultures, with

symbols depicting individual replicate data, bars showing mean ± SEM. (D) Comparison

of cell characteristics in EBV infected cells, between EBER- (in black) and EBER+ (in

red) events using histogram overlays, with populations defined in panel B. Each

histogram overlay depicts three biological replicates, comparing expression of the

defined parameter between EBER- and EBER+ populations.

1180

Supplemental Figure 4. Measurement of viral gene expression heterogeneity by

multiple measures. 3T12 cells were infected with WT γHV68 at an MOI=5 and

analyzed at 16 hpi. **(A)** Histogram overlays comparing expression of the indicated

parameters between 5 different populations of cells stratified by TMER and ORF73

expression, as defined on the leftmost panel, with data representative of three biological

replicates. **(B)** Histogram overlays comparing expression of the indicated parameter in

mock (black) or WT γHV68 -infected (red) samples, comparing three biological

replicates with events defined as DAPI+ singlets. **(C)** Viral DNA as a function of viral

gene expression defined by TMER and 73 expression. NIH 3T12 cells were either

mock or virus infected (MOI=5), harvested at 16 hpi, collected in bulk (mock and

infected, no sort) or subject to FACS purification based on expression of TMER and 73

expression to isolate TMER- 73-, TMER+ 73- and TMER high 73 high populations. DNA

was harvested from all samples, subjected to qPCR with primers to the gB gene of

gHV68. DNA content was normalized relative to cellular DNA, defined by parallel

amplification of the NFAT5 gene. Data depict mean ± SEM, with symbols indicating

individual samples. Statistical analysis was performed using one-way ANOVA subjected

to Tukey's multiple comparisons test, comparing viral DNA content between sorted cell

populations, with statistical significance as identified, *** p<0.001. **(D)** Viral gene

expression during primary lytic replication with gHV68 is relatively comparable between

cells in different stages of the cell cycle stages. Data are from imaging flow cytometry-

based studies, depicting the frequency of cells stratified by expression of TMERs and

ORF73 (as defined in Fig. 6D) present among total DAPI+ cells (top row) compared to

cells in either the G0/G1, S or G2/M phase of the cell cycle. Data depict results from

three independent experiments with 8 total replicates, with columns plotting mean ±

SEM. Flow cytometry data depict single cells, defined by sequential removal of

doublets according to SSC-A x SSC-H and FSC-A x FSC-H.

1207

Supplemental Figure 5. tSNE analysis of cell size and granularity as a function of

infection and gene expression status. Data show flow cytometry data using

populations defined in Fig. 6. Data show all DNA+ (DAPI+) single cells (FSC-A, SSC-A)

subjected to the tSNE dimensionality reduction algorithm, depicting relative expression

values for cell size (FSC) and granularity (SSC) in rows relative to the defined cell

populations (columns). The tSNE algorithm provides each cell with a unique coordinate,

displayed on a two-dimensional plot (tSNE1 versus tSNE2), such that FSC and SSC

values within cellular islands can be directly compared to the corresponding cell islands

presented in Fig. 7C. The channel range was locally-defined for each individual and

channel via Cytobank. Flow cytometry data shows single cells that are DNA+ (DAPI+).

Data are from three independent experiments.

1219

Supplemental Figure 6. Phosphonoacetic acid treatment alters viral gene

expression during lytic replication. 3T12 fibroblasts were infected with WT gHV68

(MOI=5), either in the absence of phosphonoacetic acid (no PAA) or incubated with

PAA (200 mg/mL, +PAA), harvested at 18 hpi and subjected to PrimeFlow analysis.

(A,B) Analysis of TMER and ORF73 expression profiles on a biaxial plot (A), with total

data shown in (B). (C,D) Analysis of TMER and Actin mRNA expression profiles on a

biaxial plot (C), with total data shown in (D). Events were gated on cells subjected to

doublet discrimination. Data depict mean ± SEM from two independent experiments,

with 4 biological replicates per group denoted by individual symbols. Statistical analysis

done using either a two-way ANOVA with Sidak's multiple correction test (panel B) or an

unpaired t test (panel D), with statistical significance as identified ****, $p < 0.0001$.

Figure 1
Oko et al

Figure 2
Oko et al

Figure 3
Oko et al

A. No probe | BL41 Control (KSHV- EBV-) Untreated | Mutu I (EBV+) Untreated | Mutu I (EBV+) Stimulated

SSC | AF647 | EBERs – AF647

0.047 | 0.05 ± 0.02% | 42.1 ± 8.47% | 45.1 ± 10.2%

B. EBER Median Fluorescence among EBER+ cells

Untreated | Stimulated

Figure 4
Oko et al

Figure 5
Oko et al

Figure 6
Oko et al

Figure 7
Oko et al

Figure 8
Oko et al