Neural sensitization improves encoding fidelity in the primate retina

Todd R. Appleby[1,2,3] and Michael B. Manookin[2,3,*]

[1]Graduate Program in Neuroscience, University of Washington, Seattle, WA 98195, USA

[2]Department of Ophthalmology, University of Washington, Seattle, WA 98195, USA

[3]Vision Science Center, University of Washington, Seattle, WA 98195, USA

[*]Correspondence: manookin@uw.edu

ABSTRACT

Sensory neural circuits must create internal representations of the natural environment and update those representations when the environment changes. Many neurons perform these adjustments by adapting to the statistics of incoming stimuli, allowing these cells to then report when those statistics change. Other neural populations exhibit sensitization. Instead of aligning their input-output properties with stimulus statistics, the outputs of sensitizing cells closely track the variance of incoming stimuli. These forms of short-term plasticity are commonly thought to be opposing and complementary with adapting and sensitizing cells encoding strong and weak stimuli, respectively. We report that the numerically dominant output pathway in the macaque monkey retina, the midget (parvocellular-projecting) pathway, constitutes an exception to this rule. Midget cells strongly sensitize and lack a functional counterpart that displays adaptation. Instead of counterbalancing adapting cells, sensitization in midget cells supports accurate sensory encoding and prevents a loss of responsiveness during dynamic vision.

INTRODUCTION

The fundamental constraints on sensory coding require that neural circuits adjust their outputs based on the statistical properties of their recent inputs [1–3]. Neurons respond to dynamic inputs using two distinct strategies—adaptation and sensitization. Adapting cells respond to strong stimulation by decreasing their sensitivity and this decrease in responsiveness can persist for several seconds after the stimulus intensity decreases [3–9]. Thus, adapting cells are relatively insensitive to weak stimuli occurring during these transition periods. Sensitizing cells show the opposite pattern—increasing their responsiveness at these transitions [10–12]. For this reason, adaptation and sensitization are thought to constitute opposing and complementary forms of short-term neural plasticity [10,11].

This hypothesis requires that a sensitizing cell type have an adapting counterpart that encodes common information [10]. However, this constraint could potentially decrease the amount of information that can be encoded in an neural ensemble and increase the metabolic demands on a sensory tissue [3,13]. Alternatively, adaptation and sensitization could be signatures of fundamentally distinct neural coding strategies [14]. Further, these alternative hypotheses are not mutually exclusive— adapting and sensitizing cells could mirror each other in some species and neural pathways and not in others, depending on the particular coding and metabolic constraints in those systems [2,3,13,15]. However, given that neural sensitization was only recently discovered, relatively little is known about its roles in neural information processing.

To address this issue, we recorded from five types of output neurons in the macaque monkey retina—broad thorny (koniocellular-projecting), On and Off parasol (magnocellular-projecting), and On and Off midget (parvocellular-projecting) ganglion cells. These cells have well described roles in visual processing and no known functional counterparts. We studied how these cells responded to global fluctuations in contrast and other stimulus statistics. We report that whereas broad thorny and parasol cells strongly adapted, midget cells sensitized—increasing their responsiveness to certain types of visual stimulation, including high contrast and simulated eye movements. Synaptic current recordings revealed that this increased sensitivity was present in the excitatory input from midget bipolar cells and was mediated by presynaptic disinhibition. A computational model based on synaptic input recordings further indicated that this increase in sensitivity greatly enhanced the fidelity of encoding natural scenes. Moreover, the lack of an adapting counterpart to midget cells indicates that sensitizing circuits perform a distinct role in primate retina relative to that observed in other vertebrate neural systems [10–12,16].

RESULTS

The midget pathway of the primate retina is commonly believed to lack short-term plasticity mechanisms such as contrast gain control. This belief is based on reports that midget cells did not

59 exhibit noticeable changes in responsiveness following transitions from high to low contrast regimes
60 [9,17]. The assay used to measure adaptation was a sinusoidally modulated drifting grating with bar
61 widths tuned to the size of the midget cell receptive field center, which is narrower than many other
62 retinal cell types. Thus, if plasticity in the midget pathway depended on mechanisms with broader
63 spatial tuning, this assay would not engage such mechanisms.

Figure 1. Parasol and midget cells exhibit opposing forms of plasticity.

(A) Spike rate in an On parasol ganglion cell to a low spatial frequency drifting grating presented for five seconds (temporal frequency, 6 Hz; spatial frequency, 0.35 cycles degree^{-1}). After the offset of high contrast, the spike rate declined below the level prior to grating onset (red dashed line). *Right*, zoom of transition period.

(B) Same as (A) in a broad thorny (On-Off type) ganglion cell.

(C) Same as (A) in an On midget ganglion cell to a high spatial frequency grating (3.5 cycles degree^{-1})

(D) Spike responses from the same cell as in (C) to a low spatial frequency grating (0.35 cycles degree^{-1}).

(E) Change in spike rate for the period directly after grating offset relative to period prior to grating onset in parasol (left) and midget ganglion cells (right).

To determine whether short-term plasticity in the midget pathway depended on the spatial properties of the stimulus, we repeated this assay while varying the spatial tuning of the gratings. At the offset of high contrast, midget cells did not exhibit a notable change in firing relative to the period that preceded high-contrast stimulation [9,17] (Figure 1C; spatial frequency, 3.5 cycles degree^{-1}). To determine whether this lack of either

adaptation or sensitization persisted across a range of stimulus conditions, we varied the spatial
frequency content of the drifting gratings. Following the offset of low spatial frequency gratings,
most midget cells showed an increase in spiking relative to the period preceding grating onset
(Figure 1D; spatial frequency, 0.35 cycles degree^{-1}). This increase in spiking following high contrast
is characteristic of the contrast sensitization observed in other vertebrate retinas [10–12]. The presence

of sensitization at low spatial frequencies suggested that sensitization depended on the ability to
engage elements in the midget cell receptive field with broad spatial tuning relative to the midget
bipolar cell.
99
Parasol and broad thorny cells responded very differently than midgets. At the transition from high
to low contrast, these cells showed a pronounced decrease in spiking relative to the period before
the grating turned on and several seconds were required for the spike rate to recover (Figure 1A, B;
high contrast, 1.0; low contrast, 0.0; spatial frequency, 0.18-3.5 cycles degree^{-1}; grating size, 730 μm
× 730 μm). This behavior is characteristic of contrast adaptation—during periods of high contrast,
circuit mechanisms reduce the gain to avoid saturation and the gain remains low for several seconds
following the transition to a low-contrast regime [9,18,19].
107
Wide-field stimulation evokes contrast sensitization in midget ganglion cells
Our next goal was to determine how this putative wide-field component of the midget cell receptive
field contributed to contrast coding. To accomplish this goal, we sought a more spatially and
temporally precise assay of sensitivity following wide-field adaptation. Contrast tuning of parasol and
midget cells was determined with spots centered on the receptive field (duration, 0.1 s; parasol
diameter, 80-200 μm; midget diameter, 40-80 μm). Contrast responses were measured in isolation
(unadapted condition) or 50-100 msec following the offset of an adapting stimulus (adapted
condition). The adapting stimulus was a large, high-contrast spot modulated at 20-30 Hz (diameter,
730 μm; contrast, 0.5-1.0, duration, 1.25 s). Presentations of the adapted and unadapted stimuli were
interleaved to account for any potential variability in cellular responses over time.
118
Example spike responses to this stimulus paradigm are shown in Figure 2. Parasol cells increased
their spike rate at the onset of the adapting stimulus and the spike rate quickly decreased to a steady-
state rate by ~0.25 s. Test flashes presented after the offset of the adapting stimulus evoked fewer
spikes relative to the unadapted control (Figure 2A). Both of these patterns—a transient increase in
spike rate following the transition to high contrast and a decrease in spiking after the transition to
low contrast—are characteristic of cells undergoing contrast adaptation [4,6,20].
125
We modeled the variation in the contrast-response function following the adapting stimulus as a
change in the slope (gain) and a horizontal shift relative to the control condition (see Methods).
Following the adapting stimulus, parasol cells showed a large decrease in gain ($-30.2 \pm 4.5\%$; n = 5
cells; $p = 1.3 \times 10^{-3}$; Wilcoxon signed rank test, here and below) and a small rightward horizontal
shift ($3.6 \pm 1.5\%$ contrast; $p = 3.0 \times 10^{-2}$) relative to the unadapted control (Figure 2C). This result
confirms previous reports that parasol cells readily adapt to contrast by continuously adjusting their
sensitivity to match the statistics of incoming visual inputs [9,17,18].

Figure 2. Midget ganglion cells display contrast sensitization.

(A) Spike responses from an Off parasol ganglion cell to a series of spots centered over the receptive-field. Spots were either presented alone (left) or 50 ms following the offset of an adapting stimulus (right). Shaded regions indicate sampling windows. *Right*, Average spike rate across the shaded regions. The wide-field adaptation evoked a decrease in the slope (gain) of the contrast-response curve (black) relative to the unadapted control condition (red).

(B) Same as (A) for an Off midget ganglion cell. *Right*, Average spike rate across the shaded regions. The wide-field adaptation evoked a leftward shift in the contrast-response curve (black) relative to the unadapted control condition (red).

(C) *Left*, Population data showing the change in slope (gain) for the adapted condition relative to the unadapted condition in On (open circles) and Off (closed circles) parasol ganglion cells (n = 5). *Right*, Population data showing the x-axis shift for adapted relative to unadapted conditions for small-diameter test flashes in On (open circles) and Off (closed circles) midget cells (n = 14). Gray circle and bars indicate mean ± SEM.

(D) Average spike rate evoked by wide-field test flashes for the Off midget cell in (B).

150　(E) Population data showing the *x*-axis shift for adapted relative to unadapted conditions for wide-
151　field test flashes versus small-diameter test flashes in On (open circles) and Off (closed circles)
152　midget cells. Gray circle and bars indicate mean ± SEM.
153
Midget cells showed several striking differences relative to the pattern observed in parasol cells.
First, the decrease in gain was much smaller in midget cells (–5.4 ± 4.3%; n = 14 cells; p = 0.12).
Second, an increase in spike rate was observed at the offset of the adapting stimulus relative to the
unadapted control (Figure 2B). This increase in spiking was evident at all contrasts tested including
the zero-contrast condition in which the spot intensity was equal to the average background
intensity. The elevation in spiking following the adapting stimulus produced a leftward shift in the
contrast-response curve relative to control (–16.2 ± 2.3% contrast; p = 5.2×10^{-6}). A negative
horizontal shift value occurred when the adapted curve was shifted to the left of the control curve
and this indicated that a weaker stimulus was required to elicit the same spike response from a
midget cell following the adapting stimulus. This observation was consistent with previous reports
demonstrating that a decreased spike threshold, increased baseline response, and slight decrease in
gain are characteristic of contrast sensitization [10,11].
166
Contrast sensitization is reduced for wide-field stimulation
Midget cells show narrow receptive-field centers with strong input from the receptive-field surround
[21–23]. Thus, the effect of sensitization may be diminished following the adapting stimulus depending
on the relative influences of the direct midget bipolar cell input and wide-field mechanisms in
contrast sensitization. To determine whether contrast sensitization varied with the size of the test
flash, we repeated the adaptation experiment but used wide-field test flashes to measure the contrast
tuning of midget cells (diameter, 730 μm). The wide-field test flash evoked a slight leftward shift for
the adapted condition relative to control, but this shift was much smaller than was observed for the
small-diameter test flash in the same cell (compare Figure 2B, D). This trend held true across midget
cells—horizontal shifts were more negative for the small-diameter test flash than for the wide-field
test flash in the same cell and these shifts were not statistically significant for the wide-field test
flashes (x-shift, –2.4 ± 4.6% contrast; p = 0.30; gain change, –5.8 ± 5.8%; n = 9 cells; p = 0.17;
Figure 2E). These data indicated that the relative activations of narrow-field and wide-field
mechanisms during and following the adapting stimulus were critical to contrast sensitization in
midget ganglion cells. Moreover, this result agrees well with previous findings that did not report
contrast sensitization to wide-field noise [18].
183
Time course for the onset and persistence of sensitization
We next sought to determine the amount of stimulation needed to evoke sensitization and also how
long sensitization persisted after its onset. To determine the stimulation period needed to initiate

sensitization, we varied the presentation time of the adapting stimulus and measured the change in
spike rate relative to the unadapted control (adaptation duration, 0.25-1.25 sec; contrast, ±0.5;
duration, 0.1 sec). For each period of adaptation, midget cells showed an elevation in spiking relative
to unadapted controls (Figure 3A). Thus, sensitization could be elicited even with fairly brief
stimulus presentations.
192

193
Figure 3. Time course of contrast sensitization and adaptation.
(A) Change in spike rate for the adapted condition relative to unadapted control for adaptation
periods (contrast, ±0.25-0.5; delay 0.05 s). Adaptation period was varied between 0.25-1.25 s (*x*-
axis).
(B) Duration of contrast sensitization in midget ganglion cells. Test flashes (contrast, ±0.25-0.5)
were presented at different delays (*x*-axis) following the offset of an adapting stimulus. Percent
change in spike rate for the adapted condition relative to the unadapted condition is shown on the *y*-
axis.
(C) Same as (B) for parasol ganglion cells.
Error bars indicate mean ± SEM.
204
To determine the time course of sensitization in midget cells, we measured spot responses at
different times following the offset of the adapting stimulus (delay, 0.025-1.6 sec; contrast, ±0.5;
duration, 0.1 sec). Relative to the unadapted control, the adapting stimulus elicited higher spike rates
to the test flash in midget cells at delays of 0.025-0.4 seconds (Figure 3B). This elevation in spiking,
characteristic of sensitization, was greatest 0.05-0.1 seconds after the offset of the adapting stimulus.
Parasol cells, on the other hand, showed a reduction in spiking to the same stimulus that persisted
for approximately one second (Figure 3C). Together, these data indicated that sensitization in midget
cells could be elicited even with fairly brief stimulus presentations and that it persisted for several
hundred milliseconds.
214

Sensitization enhances chromatic processing in midget cells
Midget ganglion cells in the central retina exhibit strong chromatic opponency which is formed from
differential input from long-wavelength cones (L cones) and middle-wavelength cones (M cones) to
the receptive-field center and surround [21-23]. To determine whether sensitization affected chromatic

processing, we measured contrast responses in midget cells with purely chromatic (isoluminant) test
flashes following the adapting stimulus.

Isoluminant (equiluminant) stimuli are commonly employed to study color mechanisms in isolation.
These stimuli are created by modulating L and M cones in opposing phases to silence achromatic
mechanisms that sum inputs from these cone types (i.e., L+M). We measured contrast-responses to
purely chromatic (isoluminant) flashes (duration, 0.1 sec) in the presence or absence of an
achromatic adapting stimulus, as above. As with the achromatic stimuli, the adapting stimulus
elicited a leftward shift to chromatic test contrasts (Figure 4). This shift was reminiscent of that
observed for achromatic stimulation ($-11.3 \pm 4.1\%$ contrast; n = 8 cells; p = 1.5×10^{-2}). These data
indicated that contrast sensitization enhanced both achromatic and chromatic processing in midget
cells.

While chromatic processing was affected by sensitization, the observation that an achromatic
adapting stimulus was sufficient to evoke sensitization indicated that chromatic circuits were not
necessary to elicit the phenomenon. These data did not, however, rule out contributions from purely
chromatic mechanisms to contrast sensitization.

Sensitization does not arise from a chromatic mechanism
To determine whether such a chromatic mechanism contributed to the observed contrast
sensitization, we presented a chromatic adapting stimulus. This stimulus was specifically designed to
modulate chromatic mechanisms that differentiate L- and M-cone inputs (L–M; isoluminant) while
silencing achromatic mechanisms that sum inputs from the L- and M-cone pathways (L+M;
isochromatic). Following the adapting stimulus, an isoluminant contrast series was used to measure
the input-output relationship. In the same cell, we compared chromatic contrast-responses following
a chromatic (L–M) or achromatic (L+M) adapting stimulus.

The achromatic adapting stimulus produced a leftward shift in the chromatic contrast-response
relation. The chromatic adapting stimulus, however, produced no such shift (x-shift, $-1.1 \pm 4.7\%$
contrast; n = 8 cells; p = 0.41). We interpreted this result as evidence that contrast sensitization
arose from an achromatic mechanism in the midget cell receptive-field. Moreover, given the role of
horizontal cells in forming the L-versus-M opponent receptive-field surround, these data excluded
horizontal cells as the source of sensitization in the midget pathway [21].

Figure 4. Sensitization arises from an achromatic mechanism.

(A) Spike responses from an Off midget ganglion cell to a chromatic (isoluminant) contrast series. Spots were either presented alone (left) or 50 msec following the offset of an achromatic adapting stimulus (right). Shaded regions indicate sampling windows.

(B) Average spike rate across the shaded regions indicated in (A). Achromatic adaptation evoked a leftward shift in the contrast-response curve (black) relative to the unadapted control condition (red) for the chromatic test flash.

(C) Same as (B) for a chromatic adapting stimulus. The chromatic adapting stimulus did not evoke change in the contrast-response curve relative to control.

(D) Population data showing the *x*-axis shift for adapted relative to unadapted conditions for a chromatic adapting stimulus (*x*-axis) relative to an achromatic adapting stimulus (*y*-axis) in On (open circles) and Off (closed circles) midget cells. Gray circle and bars indicate mean ± SEM.

Sensitization is present in excitatory synaptic input from midget bipolar cells

The experiments above found contrast sensitization in the spike output of midget ganglion cells. Our next goal was to understand the circuit mechanisms mediating sensitization. To accomplish this goal, we measured the direct excitatory and inhibitory synaptic inputs to midget ganglion cells with whole-cell, voltage-clamp recordings (see Methods). Excitatory currents were isolated by holding a cell's membrane voltage at the reversal potential for inhibition (–70 mV), and likewise, inhibitory currents were recorded at the excitatory reversal potential (0 mV). An increase in excitatory input to a cell was indicated by a more negative (inward) current relative to the leak current. Indeed, the adapting stimulus evoked larger inward excitatory currents relative to the unadapted control at all

contrasts tested (Figure 5A). Plotting excitatory charge as a function of contrast revealed a similar
pattern to that observed in the spike recordings—the adapting stimulus evoked a leftward shift in
the contrast-response curve relative to the unadapted control (Figure 5B). On average, the adapting
stimulus elicited a horizontal shift of –11% contrast (–11.3 ± 4.2% contrast; n = 8 cells; p = 1.95 ×
10^{-2}). These results indicated that contrast sensitization was present in the excitatory synaptic input
from midget bipolar cells to midget ganglion cells.
281

Figure 5. Sensitization present in excitatory synaptic input from midget bipolar cells.
(A) Excitatory currents from an Off midget ganglion cell to a series of spots (diameter, 40-80 µm)
centered over the receptive field. Spots were either presented alone (*left*) or 50 msec following the
offset of an adapting stimulus (*right*; diameter, 730 µm). Shaded regions indicate sampling windows.
(B) Average spike rate across the shaded regions indicated in (A). The wide-field adaptation evoked
a leftward shift in the contrast-response curve (black) relative to the unadapted control condition
(red).
(C) Population data showing the x-axis shift for adapted relative to unadapted conditions for
excitatory versus inhibitory synaptic currents in On (open circles) and Off (closed circles) midget
cells. Mean values are shown in gray. Error bars indicate mean ± SEM.
(D) Excitatory current recordings from the Off midget cell in (A) under the condition in which the
stimulus intensity returned to the mean luminance after the offset of the adapting stimulus and an
additional test flash was not presented (zero-contrast condition). A sustained increase in excitatory
current was observed at the offset of that stimulus.
(E) Proposed model for contrast sensitization in midget bipolar cells.
298

We also tested for the presence of sensitization in the inhibitory synaptic inputs to midget cells.
Unlike the pattern observed in spiking and excitatory currents, the adapting stimulus did not
consistently elicit leftward shifts in the inhibitory contrast-response functions relative to control
(+5.8 ± 4.3% contrast; n = 8 cells; p = 0.25; Figure 5C). These data indicated that contrast
sensitization arose at or prior to the level of glutamate release from midget bipolar cells. This finding
was consistent with the circuit model for contrast sensitization in bipolar cells in the retinas of fish,
salamander, mice, and rabbits [10–12]. This model posited a mechanism in which a strongly adapting
amacrine cell drove sensitization by a mechanism of presynaptic inhibition at the bipolar cell
terminal [11]. During the adapting stimulus, the amacrine cell adapted such that it decreased release of
inhibitory neurotransmitter to the bipolar cell synaptic terminal relative to the tonic level following
stimulus offset. This presynaptic disinhibition, in turn, depolarized the bipolar cell synaptic terminal,
allowing the cell to utilize its full dynamic range in signaling via glutamate release to postsynaptic
ganglion cells.
312
Cleanly measuring the effects of presynaptic inhibition on circuit function has proven exceedingly
difficult as use of inhibitory receptor antagonists typically cause many off-target effects that make
data interpretation highly tenuous [24]. Indeed, adding inhibitory antagonists in primate retina evoked
significant increases in tonic glutamate release from bipolar cells and changed the contrast polarity of
On parasol cells [25]. Nonetheless, our spike and whole-cell recordings strongly supported the
proposed model in which contrast sensitization arose from disinhibition at the presynaptic bipolar
cell terminal [11]. First, the lack of sensitization to a purely chromatic (isoluminant) adapting stimulus
indicated that sensitization did not arise in the outer retina at the level of horizontal cell feedback
(Figure 4). Second, the effect of presynaptic disinhibition was seen in our excitatory current
recordings (Figure 5D). In one of our stimulus conditions the test flash contrast was zero such that
the stimulus intensity returned to the average background intensity at the offset of the adapting
stimulus. Although this stimulus lacked a change in contrast following the adapting stimulus, we
observed an increase in excitatory synaptic input (Figure 5D). This response pattern was consistent
with a decrease in presynaptic inhibition following the offset of the adapting stimulus, resulting in an
increase in glutamate release from midget bipolar cells. Thus, our recordings in midget pathway of
primate retina were consistent with the circuit motif proposed in other vertebrate species (Figure
5E; [11]).
330
A contrast sensitization model reproduces midget cell responses
Having established the presence of contrast sensitization in midget bipolar cells, we next sought to
understand the relevance of this neural computation to visual processing in primates. To accomplish
this goal, we developed a computational model of the proposed circuit in which bipolar cell
glutamate release was modulated through presynaptic amacrine cell inhibition [10–12]. Model

parameters were determined by recording excitatory and inhibitory synaptic current responses from
midget ganglion cells to a Gaussian white noise stimulus (see Methods).
338
We modeled the midget bipolar and presynaptic amacrine cell pathways using the classical linear-
nonlinear model with two modifications: 1) adaptation occurred at the amacrine cell output and 2)
the amacrine cell output was applied to the bipolar cell model prior to the bipolar cell output
nonlinearity (Figure 6A). The model parameters controlling presynaptic sensitization were fit from
direct excitatory current recordings. In the same cell from which these parameters were determined,
we measured excitatory current responses to the wide-field adapting stimulus (see Figure 5), and the
model qualitatively reproduced the increase in excitatory currents following the offset of this
adapting stimulus (Figure 6C).
347

348
Figure 6. Sensitization model reproduces experimental results.
(A) Sensitization model structure. Visual inputs were convolved with a spatiotemporal linear filter
comprised of a Gaussian in space and a biphasic filter in time. Signals in the amacrine cell pathway
were then passed through an output nonlinearity before passing to the adaptation stage of the
model. The output of the amacrine cell model provided inhibitory input to the midget bipolar cell
pathway upstream of the bipolar cell output nonlinearity.
(B) Inhibitory temporal filter (*left*) and input-output nonlinearity (*right*) determined from noise
recordings. These filters were then used as components of the computational model (A).
(C) Excitatory current recording from an Off midget ganglion cell to the wide-field adapting
stimulus (see Figure 5). Model prediction (orange) was generated from excitatory synaptic current
recordings to the noise stimulus in the same cell.
(D) Model output for drifting grating stimuli at high and low spatial frequencies.
361
We further tested the model using the drifting grating stimuli presented in Figure 1. The model
produced distinct outputs for the high and low spatial frequency gratings. The high frequency
grating produced a relatively small response and, as a result, little adaptation in the presynaptic

amacrine cell (Figure 6D, middle row). This was due to the broad receptive field center size of the
amacrine cell relative to the bars of the grating. Low frequency gratings, however, strongly
modulated the amacrine cell and produced significant adaptation; this adaptation, in turn, caused a
removal of inhibition at the level of the bipolar cell following the offset of the grating, resulting in
sensitization (Figure 6D, bottom row). The model predictions were qualitatively similar to our direct
recordings from midget cells, indicating that contrast sensitization in primate retina can be well
explained via presynaptic disinhibition as in other species [10–12].

Sensitizing circuits more accurately reconstruct natural stimuli than adapting circuits
We next sought to understand how these differing strategies of adaptation and sensitization
impacted encoding during naturalistic vision. This was done by testing the ability of adapting and
sensitizing models to accurately encode natural scenes. We specifically wanted to determine how
accurately downstream visual circuits could reconstruct naturalistic input stimuli based on the
outputs of populations of model On and Off midget ganglion cells. The naturalistic stimuli used in
the model were taken from the DOVES database—a dataset of eye movements in humans recorded
while observing natural images [26]. Reconstruction accuracy was determined by calculating the
correlation between the stimulus and response of each model (see Methods). Periods of fixation
between ballistic eye movements are critically important to visual coding in primates; thus, model
performance was separately calculated for the complete movie or for periods of fixation only.

We considered two different decoding models for estimating the stimulus contrast based on the
outputs of On and Off midget ganglion cells. The first model utilized a linear decoding scheme in
which stimulus contrast was estimated by taking the scaled difference between the On and Off cell
outputs. We also tested a quadratic decoding model that squared the On and Off outputs prior to
differencing (see Methods). Using these decoders, we compared the performance of the sensitization
model with a model in which the midget bipolar underwent contrast adaptation. Regardless of the
decoding scheme used, the sensitizing model showed higher accuracy for reconstructing the entire
stimulus trajectory than the adapting model (linear r^2: sensitization, 0.81 ± 0.05; adaptation, $0.23 \pm$
0.07; $p = 2.7 \times 10^{-54}$; quadratic r^2: sensitization, 0.84 ± 0.05; adaptation, 0.45 ± 0.09; $p = 2.9 \times 10^{-54}$;
$n = 161$ movies; mean $\pm$ SD; Figure 7C). The sensitizing model also outperformed the adapting
model when the analysis was restricted to periods of fixation (Figure 7D).

The sensitizing model showed increased encoding accuracy for periods of fixation relative to periods
of ballistic eye movements (movement r^2, 0.63 ± 0.12; $p = 2.9 \times 10^{-35}$; Figure 7E). This finding
suggested that sensitization could play a particularly important role in vision during periods of
fixation following the offset of global motion. We, thus, sought to determine whether background
motion could evoke contrast sensitization with direct recordings from midget ganglion cells.

Figure 7. Sensitization increases the fidelity of encoding natural movies.

(A) Example image from the DOVES database. The observer's eye trajectory is shown in red.

(B) *Top*, temporal contrast sequence from the eye movement data in (A). *Bottom*, responses of the adaptation and sensitization models to the example contrast sequence.

(C) Performance of the sensitization (*x*-axis) and adaptation (*y*-axis) models at reconstructing 161 natural movies in the database. Performance was measured as the Pearson correlation between the stimulus and model predictions after adjusting for temporal lag. Performance for each movie is indicated by a black dot. Gray dot and bars indicate mean ± SD. The sensitization model outperformed the adaptation model for each of the movies.

(D) Model performances as in (C), but restricted to periods of fixation. The sensitization model outperformed the adaptation model in each case.

(E) Sensitization model performance for periods of fixation versus periods of eye motion. Predictive performance of the model was typically higher during periods of fixation.

Background motion evokes contrast sensitization in midget cells

To determine whether background motion elicited sensitization, we measured contrast responses in midget cells following the offset of a full-field moving texture (speed, 5-11 degrees s^{-1}; duration, 1 s).

The goal was to simulate, as closely as possible, the brief periods of fixation following eye

movements and to test sensitivity during these fixation periods. We interleaved these recordings with

measurements when the texture was stationary throughout the trial. The moving textures elicited an

increase in spiking and a leftward shift in the contrast-response functions relative to the control

condition in which the texture was stationary (Figure 8). On average, the shift was –25% contrast for

spike recordings (–25.4 ± 4.4% contrast; n = 10 cells; p = 2.4 × 10⁻²) and –12% contrast for

excitatory current recordings (–12.5 ± 5.1% contrast; n = 4 cells; p = 2.4 × 10⁻²).

Figure 8. Background motion evokes contrast sensitization in midget cells.

(A) Spike responses from an Off midget ganglion cell to a series of spots centered over the receptive

field. Spots were either presented alone (left) or 50 msec following the offset of background motion

(speed, 11 degrees s⁻¹). Shaded regions indicate sampling windows.

(B) Average spike rate across the shaded regions indicated in (A). The wide-field adaptation evoked

a leftward shift in the contrast-response curve (black) relative to the unadapted control condition

(red).

(C) Horizontal shift (*x*-shift) in contrast-response function following background motion relative to

control condition in which the background was stationary. Data are shown for On and Off midget

ganglion cells (n = 10). Gray circle and bars indicate mean ± SEM.

(D) Same as (A) for an Off parasol ganglion cell.

(E) Same as (B) for the Off parasol cell in (D). The cell showed a decrease in spike output following

the offset of background motion—the opposite pattern to that observed in the Off midget cell.

(F) Change in gain in the contrast-response function following background motion relative to the

control condition. On average, background motion elicited a decrease in gain of ~30% relative to

the control condition in which the background was stationary (n = 6 cells). Gray circle and bars

indicate mean ± SEM.

These data were consistent with our circuit model of contrast sensitization. The amacrine cell

providing presynaptic inhibition to the midget bipolar cell adapted during background motion; at the

offset of motion, the cell hyperpolarized and reduced presynaptic inhibition to the bipolar terminal.

Thus, similar to circuits described in other vertebrates, the midget pathway could utilize presynaptic

inhibition to account for self-motion [11,27,28].

DISCUSSION

Our results support a novel role for neural sensitization in primates relative to the function

proposed in other species. Sensitizing cells are commonly thought to counteract the loss of

responsiveness experienced by adapting cells during transitions from high to low variance

environments [10]. This hypothesis requires that sensitizing cells have an adapting counterpart that

encodes similar information about the environment. Midget (parvocellular-projecting) ganglion cells

are well known for their roles in both chromatic and achromatic vision [21–23]. Functional parallelism

in the midget pathway is achieved by splitting signals between different classes of cone

photoreceptor (L versus M) or bipolar cell (On versus Off) inputs to the midget cell receptive-field.

Further, we found that both On- and Off-type midget cells exhibited sensitization (Figure 1-4, 8),

and the primate retina lacks an adapting functional counterpart to midget cells with similar

chromatic opponency or spatial acuity [29]; thus, sensitization does not counterbalance adaptation in

another functionally parallel pathway.

Instead, our findings indicate that sensitization maintains the responsiveness of the midget pathway

during dynamic visual processes, such as head or eye movements, that cause rapid fluctuations in

light intensity on the retina. We base this conclusion on several key observations. First, sensitization

was strongest following wide-field stimulation (Figure 1-4) or background motion (Figure 8).

Second, sensitization persisted for >0.2 s (Figure 3), a period that roughly corresponds to the

durations of fixations following eye movements in primates (reviewed in [30]). Finally, sensitization

greatly improved the fidelity of encoding natural movies, particularly during periods of fixation

following ballistic eye motion (Figure 7). Thus, sensitization appears to play a unique and crucial role

in neural coding in primates.

Distinct functions of adaptation and sensitization in primate retina

Our findings also speak to the roles of neural adaptation in the parasol and broad thorny ganglion

cell pathways. Previous work proposed that adapting cells could produce a nearly optimal faithful

encoding of sensory inputs [31]. Our computational model, however, indicates that sensitizing circuits

outperform adapting circuits in encoding natural movies (Figure 7). The improved reconstruction

accuracy of the sensitizing model was consistent with a recent theoretical report indicating that

sensitizing cells are better for encoding faithful representations of sensory input than adapting cells

[14]. According to this paradigm, sensitizing cells such as midget ganglion cells would be useful for

directly encoding information about the properties of the input (e.g., contrast, color). Adapting cells,

on the other hand, are optimized for performing inference tasks [14,32].

Adapting cells dynamically adjust their input-output properties to align with the recent stimulus

distribution [4,8]. These adjustments make the cells exquisitely sensitive to changes in stimulus

statistics, allowing them to infer when salient properties of the environment change. For example,

quickly detecting object motion is an ethologically relevant and phylogenetically ancient neural

computation [33,34]; by decreasing their responsiveness during periods in which the background is

either stationary or coherently moving, adapting neural circuits would be poised to report when an

object moves relative to the background [27,35]. Interestingly, both adapting parasol and broad thorny

ganglion cells have been implicated in motion processing [25,35] and project to retinorecipient brain

regions in the lateral geniculate body, superior colliculus, and inferior pulvinar that contribute

significantly to motion vision [36–38].

Relationship to psychophysical measurements in humans

It has long been recognized that eye movements play important computational roles in visual

processing (reviewed in [39,40]). Periods in which an image is stabilized on the retina cause that image

to fade from perception [41] and small fixational eye movements appear to counteract this fading [42,43].

These eye movements can, however, produce large temporal fluctuations in contrast, particularly

when viewing high-contrast objects. This would, in turn, produce fading phenomena in cells that

strongly adapt, such as parasol ganglion cells—a prediction that was confirmed with our

computational model (Figure 7).

Neural mechanisms such as sensitization may serve to counteract adaptation by maintaining the

sensitivity of certain visual pathways during eye movements. Indeed, our computational model and

direct measurements indicated that contrast sensitization in the midget ganglion cell pathway was

engaged well by background motion such as that observed during eye movements (Figure 7, 8).

Thus, contrast sensitization might act to maintain sensitivity of image-forming visual pathways

following eye movements that are commonplace in primate vision. Indeed, psychophysical studies in

humans indicated that contrast sensitivity increases following both ballistic (saccade) and fixational

eye movements [42,43]. Moreover, this increase in sensitivity was limited to chromatic stimuli and high-

spatial-frequency achromatic stimuli, mirroring our results in midget ganglion cells.

METHODS

Experiments were performed in an *in vitro*, pigment-epithelium attached preparation of the macaque monkey retina [44]. Eyes were dissected from terminally anesthetized macaque monkeys of either sex (Macaca *fascicularis*, *mulatta*, and *nemestrina*) obtained through the Tissue Distribution Program of the National Primate Research Center at the University of Washington. All procedures were approved by the University of Washington Institutional Animal Care and Use Committee.

Tissue Preparation and Electrophysiology

The retina was continuously superfused with warmed (32-35°C) Ames' medium (Sigma) at ~6-8 mL min^{-1}. Recordings were performed from macular, mid-peripheral, or peripheral retina (2-8 mm, 10-30° foveal eccentricity), but special emphasis was placed on recording from more centrally located cells. Physiological data were acquired at 10 kHz using a Multiclamp 700B amplifier (Molecular Devices), Bessel filtered at 3 kHz (900 CT, Frequency Devices), digitized using an ITC-18 analog-digital board (HEKA Instruments), and acquired using the Symphony acquisition software package developed in Fred Rieke's laboratory (http://symphony-das.github.io).

Recordings were performed using borosilicate glass pipettes containing Ames medium for extracellular spike recording or, for whole-cell recording, a cesium-based internal solution containing (in mM): 105 CsCH$_3$SO$_3$, 10 TEA-Cl, 20 HEPES, 10 EGTA, 2 QX-314, 5 Mg-ATP, and 0.5 Tris-GTP, pH ~7.3 with CsOH, ~280 mOsm. Series resistance (~3-9 MΩ) was compensated online by 50%. The membrane potential was corrected offline for the approximately –11 mV liquid junction potential between the intracellular solution and the extracellular medium. Excitatory and inhibitory synaptic currents were isolated by holding midget ganglion cells at the reversal potentials for inhibition/chloride (~–70 mV) and excitation (0 mV), respectively.

Visual Stimuli and Data Analysis

Visual stimuli were generated using the Stage software package developed in the Rieke lab (http://stage-vss.github.io) and displayed on a digital light projector (Lightcrafter 4500; Texas Instruments) modified with custom LEDs with peak wavelengths of 405, 505 (or 475), and 640 nm. Stimuli were focused on the photoreceptor outer segments through a 10X microscope objective. Mean light levels were in the low to medium photopic regimes (~$3 \times 10^3 - 3.4 \times 10^4$ photoisomerizations [R*] cone^{-1} sec^{-1}). Contrast values for contrast-response flashes are given in Weber contrast and for periodic stimuli in Michaelson contrast. All responses were analyzed in MATLAB (R2018a+, Mathworks).

For extracellular recordings, currents were wavelet filtered to remove slow drift and amplify spikes relative to the noise [45] and spikes were detected using either a custom k-means clustering algorithm

555 or by choosing a manual threshold. Whole-cell recordings were leak subtracted and responses were

556 measured relative to the median membrane currents immediately preceding stimulus onset (0.25-0.5

557 s window). Summary data are presented in terms of conductance (g), which is the ratio of the current

558 response (I) to the driving force:

$$g = \frac{I}{V_m - E}$$

where V_m is the holding potential (in mV) and E is the reversal potential (in mV). Reversal potentials

of 0 mV and –70 mV were used for excitatory and inhibitory inputs, respectively.

Sensitization and adaptation models

We modeled spatiotemporal integration in bipolar cells and amacrine cells as the product of a

Gaussian spatial filter and a biphasic temporal filter which was then passed through an input-output

nonlinearity. The output of this nonlinear stage of the amacrine cell model was then passed through

an adaptation stage; adaptation in the amacrine cell provided inhibitory input to the bipolar cell

model prior to the output nonlinearity (Figure 6A). Following the subunit output, model midget

ganglion cells and amacrine cells pooled (summed) inputs from bipolar cell subunits and the weights

of these inputs were normalized by the subunit location relative to the receptive field center using a

Gaussian weighting.

To estimate the excitatory and inhibitory circuit components for the computational model, we

recorded excitatory and inhibitory synaptic currents from midget ganglion cells in response to a full-

field Gaussian flicker stimulus. The contrast of each frame was drawn randomly from a Gaussian

distribution and that value was multiplied by the average contrast. Average contrast was updated

every 0.5 s and drawn from a uniform distribution (0.05-0.35 RMS contrast). The linear temporal

filters (F) were calculated by cross-correlating the stimulus sequence (S) and the leak-subtracted

response (R) [4].

$$F(t) = \int R(\tau)S(t + \tau)d\tau$$

581 where τ is the temporal lag. These filters were then modeled as a damped oscillator with an S-shaped

582 onset [46,47]:

$$F(t) = A\frac{(t/\tau_{rise})^n}{1 + (t/\tau_{rise})^n}e^{-\left(\frac{t}{\tau_{decay}}\right)}cos\left(\frac{2\pi t}{\tau_{period}} + \varphi\right)$$

584 where A is a scaling factor, τ_{rise} is the rising-phase time constant, τ_{decay} is the damping time constant,

585 τ_{period} is the oscillator period, and φ is the phase (in degrees).

587 The input-output nonlinearity was calculated by convolving the temporal filter (F) and stimulus (S)

588 to generate the linear prediction (P).

$$P(t) = \int F(\tau)S(t - \tau)d\tau$$

590 The prediction (*x*-axis) and response (*y*-axis) were modeled as a cumulative Gaussian distribution [48].

$$N(x) = \varepsilon + \frac{\alpha}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{\frac{-(\beta t + \gamma)^2}{2}} dt$$

where *a* indicates the maximal output value, ε is the vertical offset, β is the sensitivity of the output

to the generator signal (input), and γ is the maintained input to the cell.

The spatial component of the bipolar and amacrine cell receptive fields was modeled as a Gaussian

function with a 2-SD width of 18 µm and 90 µm, respectively. Each midget ganglion cell was

modeled as receiving input from a single bipolar cell, as is typically the case in the central retina.

Sensitization parameters were determined by fitting linear-nonlinear model predictions relative to

the excitatory currents recorded to the Gaussian flicker stimulus.

The amacrine cell providing direct inhibition to the midget ganglion cells is likely distinct from the

cell providing presynaptic inhibition at the level of the midget bipolar cell (see Figure 5). Thus, our

inhibitory synaptic recordings likely did not grant us direct access to the properties of the amacrine

cell responsible for contrast sensitization. These recordings do, however, provide an estimate of the

time-course of signals passing through the presynaptic amacrine cell to midget bipolar cells. Signals

passing through this amacrine cell proceed from cone photoreceptors to bipolar cells and then to

the amacrine cell in question before providing input to the midget bipolar cell. In the same way, the

amacrine cell providing direct inhibition to midget ganglion cells must pass through an extra

synapse. Thus, our recordings of direct synaptic inhibition were useful in approximating the time

course of presynaptic inhibition at the midget bipolar terminal.

Evaluating model performance to naturalistic movies

We evaluated the performance of the adaptation and sensitization models in reconstructing the

naturalistic movie sequences using linear and quadratic decoding paradigms. To estimate stimulus

contrast, the linear decoder (*f*$_{LINEAR}$) summed the scaled outputs of the model On and Off midget

ganglion cells:

$$f_{\text{LINEAR}}(t) = a_{\text{ON}}r_{\text{ON}}(t) + a_{\text{OFF}}r_{\text{OFF}}(t) + k$$

618 where *a*$_{ON}$ and *a*$_{OFF}$ are scaling constants and *k* is an offset constant. The quadratic model was similar

619 in structure except that the response from each pathways was squared prior to summation:

$$f_{\text{QUADRATIC}}(t) = a_{\text{ON1}}r_{\text{ON}}(t) + a_{\text{ON2}}r_{\text{ON}}^2(t) + a_{\text{OFF1}}r_{\text{OFF}}(t) + a_{\text{OFF2}}r_{\text{OFF}}^2(t) + k$$

621 For each of the 161 movies in the database, the input stimulus was shifted to the peak of the midget

622 temporal filter (~35 ms) and then scaling and offset coefficients were determined using least-squares

623 curve fitting. The Pearson correlation was then calculated between the temporal trajectories of the

624 model and the movie.

Quantification and Statistical Analysis

All statistical analyses were performed in MATLAB (R2018a+, Mathworks). Reported p values in this study were paired and were calculated using the Wilcoxon signed-rank test. Final figures were created in MATLAB, Igor Pro, and Adobe Illustrator.

ACKNOWLEDGEMENTS

We thank Shellee Cunnington, Mark Cafaro, and Jim Kuchenbecker for technical assistance. Tissue was provided by the Tissue Distribution Program at the Washington National Primate Research Center (WaNPRC; supported through NIH grant P51 OD-010425), and we thank the WaNPRC staff, particularly Chris English, for making these experiments possible. Fred Rieke, Raunak Sinha, Max Turner, and Will Grimes assisted in tissue preparation. We thank Jay Neitz and Fred Rieke for helpful discussions. We also thank Alison Weber and Jon Demb for feedback on a previous version of this manuscript. This work was supported in part by grants from the NIH (NEI R01-EY027323 to M.B.M.; NEI P30-EY001730 to the Vision Core), Research to Prevent Blindness Unrestricted Grant (to the University of Washington Department of Ophthalmology), Latham Vision Research Innovation Award (to M.B.M.), and the Alcon Young Investigator Award (to M.B.M.).

AUTHOR CONTRIBUTIONS

Conceptualization, M.B.M.; Methodology, M.B.M.; Software, M.B.M.; Formal Analysis, M.B.M.; Investigation, M.B.M., T.R.A; Resources, M.B.M.; Data Curation, M.B.M., T.R.A.; Writing – Original Draft, M.B.M.; Writing – Review & Editing, M.B.M., T.R.A.; Visualization, M.B.M.; Supervision, M.B.M.; Project Administration, M.B.M.; Funding Acquisition, M.B.M. The ORCID number for M.B.M. is 0000-0001-8116-7619.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Srinivasan, M. V., Laughlin, S. B. & Dubs, A. Predictive coding: a fresh view of inhibition in the retina. *Proc. R. Soc. Lond. B Biol. Sci.* **216,** 427–459 (1982).

2. Barlow, H. B. The coding of sensory messages. in *Current problems in animal behaviour* (eds. Thorpe, W. H. & Zangwill, O. L.) 331–360 (Cambridge University Press, 1961).

3. Laughlin, S. A simple coding procedure enhances a neuron's information capacity. *Z. Naturforsch. C* **36,** 910–912 (1981).

4. Baccus, S. A. & Meister, M. Fast and slow contrast adaptation in retinal circuitry. *Neuron* **36,** 909–919 (2002).

5. Carandini, M. & Ferster, D. A tonic hyperpolarization underlying contrast adaptation in cat visual cortex. *Science* **276,** 949–952 (1997).

6. Kim, K. J. & Rieke, F. Temporal contrast adaptation in the input and output signals of salamander retinal ganglion cells. *J. Neurosci.* **21,** 287–299 (2001).

7. Manookin, M. B. & Demb, J. B. Presynaptic mechanism for slow contrast adaptation in mammalian retinal ganglion cells. *Neuron* **50,** 453–464 (2006).

8. Smirnakis, S. M., Berry, M. J., Warland, D. K., Bialek, W. & Meister, M. Adaptation of retinal processing to image contrast and spatial scale. *Nature* **386,** 69–73 (1997).

9. Solomon, S. G., Peirce, J. W., Dhruv, N. T. & Lennie, P. Profound contrast adaptation early in the visual pathway. *Neuron* **42,** 155–162 (2004).

10. Kastner, D. B. & Baccus, S. A. Coordinated dynamic encoding in the retina using opposing forms of plasticity. *Nat. Neurosci.* **14,** 1317–1322 (2011).

11. Kastner, D. B. & Baccus, S. A. Spatial segregation of adaptation and predictive sensitization in retinal ganglion cells. *Neuron* **79,** 541–554 (2013).

12. Nikolaev, A., Leung, K. M., Odermatt, B. & Lagnado, L. Synaptic mechanisms of adaptation and sensitization in the retina. *Nat. Neurosci.* **16,** 934–941 (2013).

13. Balasubramanian, V., Kimber, D. & Berry, M. J., 2nd. Metabolically efficient information processing. *Neural Comput.* **13,** 799–815 (2001).

14. Młynarski, W. F. & Hermundstad, A. M. Adaptive coding for dynamic sensory inference. *Elife* **7,** (2018).

15. Levy, W. B. & Baxter, R. A. Energy efficient neural codes. *Neural Comput.* **8,** 531–543 (1996).

16. Cohen-Kashi Malina, K., Jubran, M., Katz, Y. & Lampl, I. Imbalance between excitation and inhibition

in the somatosensory cortex produces postadaptation facilitation. *J. Neurosci.* **33,** 8463–8471 (2013).

17. Benardete, E. A., Kaplan, E. & Knight, B. W. Contrast gain control in the primate retina: P cells are not

X-like, some M cells are. *Vis. Neurosci.* **8,** 483–486 (1992).

18. Chander, D. & Chichilnisky, E. J. Adaptation to temporal contrast in primate and salamander retina. *J.*

Neurosci. **21,** 9904–9916 (2001).

19. Benardete, E. A. & Kaplan, E. The dynamics of primate M retinal ganglion cells. *Vis. Neurosci.* **16,** 355–

368 (1999).

20. Brown, S. P. & Masland, R. H. Spatial scale and cellular substrate of contrast adaptation by retinal

ganglion cells. *Nat. Neurosci.* **4,** 44–51 (2001).

21. Crook, J. D., Manookin, M. B., Packer, O. S. & Dacey, D. M. Horizontal cell feedback without cone

type-selective inhibition mediates 'red-green' color opponency in midget ganglion cells of the primate

retina. *J. Neurosci.* **31,** 1762–1772 (2011).

22. De Monasterio, F. M. & Gouras, P. Functional properties of ganglion cells of the rhesus monkey retina.

J. Physiol. **251,** 167–195 (1975).

23. Derrington, A. M., Krauskopf, J. & Lennie, P. Chromatic mechanisms in lateral geniculate nucleus of

macaque. *J. Physiol.* **357,** 241–265 (1984).

24. Cook, P. B., Lukasiewicz, P. D. & McReynolds, J. S. Action potentials are required for the lateral

transmission of glycinergic transient inhibition in the amphibian retina. *J. Neurosci.* **18,** 2301–2308 (1998).

25. Manookin, M. B., Patterson, S. S. & Linehan, C. M. Neural Mechanisms Mediating Motion Sensitivity in

Parasol Ganglion Cells of the Primate Retina. *Neuron* **97,** 1327–1340.e4 (2018).

26. Van Der Linde, I., Rajashekar, U., Bovik, A. C. & Cormack, L. K. DOVES: a database of visual eye

movements. *Spat. Vis.* **22,** 161–177 (2009).

27. Olveczky, B. P., Baccus, S. A. & Meister, M. Segregation of object and background motion in the retina.

Nature **423,** 401–408 (2003).

28. Baccus, S. A., Ölveczky, B. P., Manu, M. & Meister, M. A retinal circuit that computes object motion. *J.*

Neurosci. **28,** 6807–6817 (2008).

29. Wässle, H. Parallel processing in the mammalian retina. *Nat. Rev. Neurosci.* **5,** 747–757 (2004).

30. Rayner, K. Eye movements in reading and information processing: 20 years of research. *Psychol. Bull.*

124, 372–422 (1998).

31. Fairhall, A. L., Lewen, G. D., Bialek, W. & de Ruyter Van Steveninck, R. R. Efficiency and ambiguity in

an adaptive neural code. *Nature* **412,** 787–792 (2001).

32. Wark, B., Fairhall, A. & Rieke, F. Timescales of inference in visual adaptation. *Neuron* **61,** 750–761

(2009).

33. Frost, B. J., Wylie, D. R. & Wang, Y. C. The processing of object and self-motion in the tectofugal and

accessory optic pathways of birds. *Vision Res.* **30,** 1677–1688 (1990).

34. Lettvin, J. Y., Maturana, H. R., McCulloch, W. S. & Pitts, W. H. What the Frog's Eye Tells the Frog's

Brain. *Proceedings of the IRE* **47,** 1940–1951 (1959).

35. Puller, C., Manookin, M. B., Neitz, J., Rieke, F. & Neitz, M. Broad thorny ganglion cells: a candidate for

visual pursuit error signaling in the primate retina. *J. Neurosci.* **35,** 5397–5408 (2015).

36. Rodieck, R. W. & Watanabe, M. Survey of the morphology of macaque retinal ganglion cells that project

to the pretectum, superior colliculus, and parvicellular laminae of the lateral geniculate nucleus. *J. Comp.*

Neurol. **338,** 289–303 (1993).

37. Crook, J. D. *et al.* Y-cell receptive field and collicular projection of parasol ganglion cells in macaque

monkey retina. *J. Neurosci.* **28,** 11277–11291 (2008).

38. Kwan, W. C. *et al.* Unravelling the subcortical and retinal circuitry of the primate inferior pulvinar. *J.*

Comp. Neurol. (2018). doi:10.1002/cne.24387

39. Martinez-Conde, S., Macknik, S. L. & Hubel, D. H. The role of fixational eye movements in visual

perception. *Nat. Rev. Neurosci.* **5,** 229–240 (2004).

40. Rucci, M. & Victor, J. D. The unsteady eye: an information-processing stage, not a bug. *Trends Neurosci.*

38, 195–206 (2015).

41. Troxler, D. Verschwinden gegebener Gegenstände innerhalb unseres Gesichtskreises. *Ophthalmologische*

Bibliothek **2,** 1–53 (1804).

42. Rucci, M., Iovin, R., Poletti, M. & Santini, F. Miniature eye movements enhance fine spatial detail.

Nature **447,** 851–854 (2007).

43. Schütz, A. C., Braun, D. I., Kerzel, D. & Gegenfurtner, K. R. Improved visual sensitivity during smooth

pursuit eye movements. *Nat. Neurosci.* **11,** 1211–1216 (2008).

44. Manookin, M. B., Puller, C., Rieke, F., Neitz, J. & Neitz, M. Distinctive receptive field and physiological

properties of a wide-field amacrine cell in the macaque monkey retina. *J. Neurophysiol.* **114,** 1606–1616

(2015).

45. Wiltschko, A. B., Gage, G. J. & Berke, J. D. Wavelet filtering before spike detection preserves waveform

shape and enhances single-unit discrimination. *J. Neurosci. Methods* **173,** 34–40 (2008).

46. Schnapf, J. L., Nunn, B. J., Meister, M. & Baylor, D. A. Visual transduction in cones of the monkey

Macaca fascicularis. *J. Physiol.* **427,** 681–713 (1990).

47. Angueyra, J. M. & Rieke, F. Origin and effect of phototransduction noise in primate cone

photoreceptors. *Nat. Neurosci.* **16,** 1692–1700 (2013).

48. Chichilnisky, E. J. A simple white noise analysis of neuronal light responses. *Network* **12,** 199–213 (2001).

FIGURE LEGENDS

Figure 1. Parasol and midget cells exhibit opposing forms of plasticity.

(A) Spike rate in an On parasol ganglion cell to a low spatial frequency drifting grating presented for

five seconds (temporal frequency, 6 Hz; spatial frequency, 0.35 cycles degree^{-1}). After the offset of

high contrast, the spike rate declined below the level prior to grating onset (red dashed line). *Right*,

zoom of transition period.

(B) Same as (A) in a broad thorny (On-Off type) ganglion cell.

(C) Same as (A) in an On midget ganglion cell to a high spatial frequency grating (3.5 cycles degree^{-1})

(D) Spike responses from the same cell as in (C) to a low spatial frequency grating (0.35 cycles

degree^{-1}).

(E) Change in spike rate for the period directly after grating offset relative to period prior to grating

onset in parasol (left) and midget ganglion cells (right).

Figure 2. Midget ganglion cells display contrast sensitization.

(A) Spike responses from an Off parasol ganglion cell to a series of spots centered over the

receptive-field. Spots were either presented alone (left) or 50 ms following the offset of an adapting

stimulus (right). Shaded regions indicate sampling windows. *Right*, Average spike rate across the

shaded regions. The wide-field adaptation evoked a decrease in the slope (gain) of the contrast-

response curve (black) relative to the unadapted control condition (red).

(B) Same as (A) for an Off midget ganglion cell. *Right*, Average spike rate across the shaded regions.

The wide-field adaptation evoked a leftward shift in the contrast-response curve (black) relative to

the unadapted control condition (red).

(C) *Left*, Population data showing the change in slope (gain) for the adapted condition relative to the

unadapted condition in On (open circles) and Off (closed circles) parasol ganglion cells (n = 5).

Right, Population data showing the *x*-axis shift for adapted relative to unadapted conditions for

small-diameter test flashes in On (open circles) and Off (closed circles) midget cells (n = 14). Gray

circle and bars indicate mean ± SEM.

(D) Average spike rate evoked by wide-field test flashes for the Off midget cell in (B).

(E) Population data showing the *x*-axis shift for adapted relative to unadapted conditions for wide-

field test flashes versus small-diameter test flashes in On (open circles) and Off (closed circles)

midget cells. Gray circle and bars indicate mean ± SEM.

Figure 3. Time course of contrast sensitization and adaptation.

(A) Change in spike rate for the adapted condition relative to unadapted control for adaptation

periods (contrast, ±0.25-0.5; delay 0.05 s). Adaptation period was varied between 0.25-1.25 s (*x*-

axis).

(B) Duration of contrast sensitization in midget ganglion cells. Test flashes (contrast, ±0.25-0.5)

were presented at different delays (*x*-axis) following the offset of an adapting stimulus. Percent

change in spike rate for the adapted condition relative to the unadapted condition is shown on the *y*-

axis.

(C) Same as (B) for parasol ganglion cells.

Error bars indicate mean ± SEM.

Figure 4. Sensitization arises from an achromatic mechanism.

(A) Spike responses from an Off midget ganglion cell to a chromatic (isoluminant) contrast series.

Spots were either presented alone (left) or 50 msec following the offset of an achromatic adapting

stimulus (right). Shaded regions indicate sampling windows.

(B) Average spike rate across the shaded regions indicated in (A). Achromatic adaptation evoked a

leftward shift in the contrast-response curve (black) relative to the unadapted control condition (red)

for the chromatic test flash.

(C) Same as (B) for a chromatic adapting stimulus. The chromatic adapting stimulus did not evoke

change in the contrast-response curve relative to control.

(D) Population data showing the *x*-axis shift for adapted relative to unadapted conditions for a

chromatic adapting stimulus (*x*-axis) relative to an achromatic adapting stimulus (*y*-axis) in On (open

circles) and Off (closed circles) midget cells. Gray circle and bars indicate mean ± SEM.

Figure 5. Sensitization present in excitatory synaptic input from midget bipolar cells.

(A) Excitatory currents from an Off midget ganglion cell to a series of spots (diameter, 40-80 μm)

centered over the receptive field. Spots were either presented alone (*left*) or 50 msec following the

offset of an adapting stimulus (*right*; diameter, 730 μm). Shaded regions indicate sampling windows.

(B) Average spike rate across the shaded regions indicated in (A). The wide-field adaptation evoked

a leftward shift in the contrast-response curve (black) relative to the unadapted control condition

(red).

(C) Population data showing the x-axis shift for adapted relative to unadapted conditions for

excitatory versus inhibitory synaptic currents in On (open circles) and Off (closed circles) midget

cells. Mean values are shown in gray. Error bars indicate mean ± SEM.

(D) Excitatory current recordings from the Off midget cell in (A) under the condition in which the

stimulus intensity returned to the mean luminance after the offset of the adapting stimulus and an

additional test flash was not presented (zero-contrast condition). A sustained increase in excitatory

current was observed at the offset of that stimulus.

(E) Proposed model for contrast sensitization in midget bipolar cells.

Figure 6. Sensitization model reproduces experimental results.

(A) Sensitization model structure. Visual inputs were convolved with a spatiotemporal linear filter

comprised of a Gaussian in space and a biphasic filter in time. Signals in the amacrine cell pathway

were then passed through an output nonlinearity before passing to the adaptation stage of the

model. The output of the amacrine cell model provided inhibitory input to the midget bipolar cell

pathway upstream of the bipolar cell output nonlinearity.

(B) Inhibitory temporal filter (*left*) and input-output nonlinearity (*right*) determined from noise

recordings. These filters were then used as components of the computational model (A).

(C) Excitatory current recording from an Off midget ganglion cell to the wide-field adapting

stimulus (see Figure 5). Model prediction (orange) was generated from excitatory synaptic current

recordings to the noise stimulus in the same cell.

Figure 7. Sensitization increases fidelity of encoding natural movies.

(A) Example image from the DOVES database. The observer's eye trajectory is shown in red.

(B) *Top*, temporal contrast sequence from the eye movement data in (A). *Bottom*, responses of the adaptation and sensitization models to the example contrast sequence.

(C) Performance of the sensitization (*x*-axis) and adaptation (*y*-axis) models at reconstructing 161 natural movies in the database. Performance was measured as the Pearson correlation between the stimulus and model predictions after adjusting for temporal lag. Performance for each movie is indicated by a black dot. Gray dot and bars indicate mean ± SD. The sensitization model outperformed the adaptation model for each of the movies.

(D) Model performances as in (C), but restricted to periods of fixation. The sensitization model outperformed the adaptation model in each case.

(E) Sensitization model performance for periods of fixation versus periods of eye motion. Predictive performance of the model was typically higher during periods of fixation.

Figure 8. Background motion evokes contrast sensitization in midget cells.

(A) Spike responses from an Off midget ganglion cell to a series of spots centered over the receptive field. Spots were either presented alone (left) or 50 msec following the offset of background motion (speed, 11 degrees s^{-1}). Shaded regions indicate sampling windows.

(B) Average spike rate across the shaded regions indicated in (A). The wide-field adaptation evoked a leftward shift in the contrast-response curve (black) relative to the unadapted control condition (red).

(C) Horizontal shift (*x*-shift) in contrast-response function following background motion relative to control condition in which the background was stationary. Data are shown for On and Off midget ganglion cells (n = 10). Gray circle and bars indicate mean ± SEM.

(D) Same as (A) for an Off parasol ganglion cell.

(E) Same as (B) for the Off parasol cell in (D). The cell showed a decrease in spike output following the offset of background motion—the opposite pattern to that observed in the Off midget cell.

(F) Change in gain in the contrast-response function following background motion relative to the control condition. On average, background motion elicited a decrease in gain of ~30% relative to the control condition in which the background was stationary (n = 6 cells). Gray circle and bars indicate mean ± SEM.