Genetic differentiation and intrinsic genomic features explain variation in recombination hotspots among cocoa tree populations

Enrique J. Schwarzkopf[1], Juan C. Motamayor[2], Omar E. Cornejo[1,a]
Affiliations:
[1] School of Biological Sciences, Washington State University, Pullman, WA, USA
[2] Universal Genetic Solutions, LLC
[a] Corresponding author: ocornejo@gmail.com

Correspondence:

Running title: Recombination hotspots in ten *Theobroma cacao* populations

Keywords: recombination; recombination hotspots; domestication

Abstract:

Our study investigates the possible drivers of recombination hotspots in *Theobroma cacao* using ten genetically differentiated populations. By comparing recombination patterns between multiple populations, we obtain a novel view of recombination at the population-divergence timescale. For each population, a fine-scale recombination map was generated using the coalescent with a standard method based on linkage disequilibrium (LD). These maps revealed higher recombination rates in a domesticated population and a population that has undergone a recent bottleneck. We inferred hotspots of recombination for each population and find that the genomic locations of hotspots correlate with genetic differentiation between populations (F_{ST}). We used randomization approaches to generate appropriate null models to understand the association between hotspots of recombination and both DNA sequence motifs and genomic features. We found that hotspot regions contained fewer known retroelement sequences than expected and were overrepresented near transcription start and termination sites. Our findings indicate that recombination hotspots are evolving in a way that is consistent with genetic differentiation but are also preferentially driven to near coding regions. We illustrate that, consistent with predictions in plant domestication, the recombination rate of the domesticated population is orders of magnitude higher than that of other populations. More importantly, we find two fixed mutations in the domesticated population's FIGL1 protein. FIGL1 has been shown to increase recombination rates in *Arabidopsis* by several orders of magnitude, suggesting a possible mechanism for the observed increased recombination rate in the domesticated population.

Introduction:

Genetic recombination is an important source of genome-wide genetic variation fundamental for evolutionary forces like selection and genetic drift to act. Selection and drift contribute to a loss of variation, which means that in the absence of forces that maintain variation along the genome, populations would be incapable of evolving over prolonged periods of time. Recombination rearranges genetic material onto different backgrounds generating a larger set of haplotype combinations on which selection can act, reducing the magnitude of Hill-Robertson interference (Felsenstein 1974). Different regimes of recombination can strongly influence how efficient selection is at purging deleterious mutations and increasing the frequency of beneficial mutations in the population (Felsenstein 1974).

Studies in a wide range of species have shown that recombination rates are not uniform along
the genome and general patterns of variation have been described (Begun and Aquadro 1992,
Akhunov et al. 2003, Wu et al. 2003, Anderson et al. 2004, McVean et al. 2004, Mézard 2006,
Kim et al. 2007, Gore et al. 2009, Schnable et al. 2009, Branca et al. 2011,Paape et al. 2012).
One pattern that has been observed in multiple species is the reduced recombination rate in
centromeric regions of the chromosomes and the progressive increase of recombination rates
as the physical distance from the telomeres decreases (Begun and Aquadro 1992, Akhunov et
al. 2003, Wu et al. 2003, Anderson et al. 2004, Gore et al. 2009, Schnable et al. 2009). This
pattern has also been shown to arise in simulation studies (e.g. Mackiewicz et al. 2010).
Another interesting pattern that has been observed is that of regions with unusually high rates of
recombination spread throughout chromosomes: recombination hotspots (McVean et al. 2004,
Brunschwig et al. 2012, Paape et al. 2012, Hellsten et al. 2013, Stevison et al. 2016, Shanfelter
et al. 2018). The importance of recombination hotspots lies in their ability to shuffle genetic
variation at higher rates than the rest of the genome, profoundly impacting the dynamics of
selection for or against specific mutations (Felsenstein 1974). In this study, we focus on locally
defined recombination hotspots, requiring that their recombination rate be unusually high when
compared to neighboring regions.
68
A variety of genomic features have been identified as being associated with regions of high
recombination. Recombination hotspots have been linked to transcriptional start sites (TSSs)
and transcriptional termination sites (TTSs) in *Arabidopsis thaliana*, *Taeniopygia guttata*,
Poephila acuticauda, and humans (Myers et al. 2005, Choi et al. 2013, Singhal et al. 2015). In
Mimulus guttatus hotspots were found to be associated with cpg islands (short segments of
cytosine and guanine rich DNA, associated with promoter regions) (Hellsten et al. 2013). CpG
islands were also associated with increased recombination rates in humans and chimpanzees
(Auton et al. 2012). These patterns point to recombination occurring frequently near, but not
within, coding regions. The formation of chiasmata is important for the proper disjunction of
chromosomes during meiosis (Martinez-Perez et al. 2008), but repeated double-strand breaks
can lead to an increased mutation rate (Rodgers and McVey 2015). In coding regions in
particular, this excess mutation rate can have a high evolutionary cost, due to the likelihood of
novel deleterious mutations being higher than that of beneficial ones (Haldane 1937, Crow and
Kimura 1970, Wloch et al. 2001, Sanjuán et al. 2004, Eyre-Walker and Keightley 2007).
Recombination hotspots have also been found to be correlated with particular DNA sequence
motifs. In some mammals, including *Mus musculus* (Brunschwig et al. 2012) and apes (Auton et
al. 2012, Stevison et al. 2016) binding sites for PRDM9, a histone trimethylase with a DNA zinc-
finger binding domain, have been found to correlate with recombination hotspots. In *A. Thaliana*,
proteins that limit overall recombination rate have been identified, leading to a genome-wide
increase in recombination rate in knockout mutants (Fernandes et al. 2018b). However, these
Arabidopsis proteins have not been shown to direct recombination to particular regions and are
therefore not expected to affect the location of recombination hotspots.
91
The dynamics of recombination hotspots shared between related species or populations of the
same species have been investigated in apes, yielding varying results. Hinch et al. (2011) found
that, at finer scales, the genetic maps of European and African human populations were
significantly different. They also found that, when looking at hotspots in the major
histocompatibility complex, the African populations showed a hotspot that was not present in
Europeans, but all European hotspots were found in African populations (Hinch et al. 2011).
Recent work on recombination in apes found little correlation of recombination rates in
orthologous hotspot regions when looking between species, but a strong correlation when
comparing between two populations of the same species (Stevison et al. 2016). Other studies
have also found very little sharing of hotspots between humans and chimpanzees (Ptak et al.

2005, Winckler et al. 2005). Additionally, the dynamic of changing hotspot locations observed in
humans and other apes has been observed in simulations (Mackiewicz et al. 2013). The
disparity of empirical results regarding hotspots shared between related populations suggest
that further work is required to disentangle the relationship between demographics and shared
hotspots.
107
The identification of ten genetically differentiated populations of the cocoa tree, *Theobroma*
cacao, (Motamayor et al. 2008, Cornejo et al. 2017) can be leveraged to study population-level
dynamics of recombination patterns. The ten *T. Cacao* populations originate from different
regions of South and Central America, and include one fully domesticated population (Criollo),
used in the production of fine chocolate, and nine wilder, more resilient populations which
generate higher cocoa yield than the Criollo variety (Fig. S1) (Motamayor et al. 2008,
Henderson et al. 2007, Cornejo et al 2017). These ten populations have been shown to have
strong signatures of differentiation between them (F_{ST} values ranging from 0.16 to 0.65) and
they separate into clear clusters of ancestry (Cornejo et al. 2017). During domestication,
recombination plays an important role in the segregation of traits, and for this reason it has been
hypothesized that recombination rates will increase during the process of domestication
(Moyers et al. 2018). Domestication can be a rapid process and there is theoretical evidence for
the increase of recombination rates during periods of rapid evolutionary change (Otto and
Barton 1997). Empirical evidence for this prediction has been shown in a limited number of
herbaceous plant species with short generation times (Ross-Ibarra 2004). It is not clear if plant
species with longer generation times are also expected to experience increased recombination
rates, and it is also unclear what mechanisms could explain these differences. One possible
explanation for differences in recombination rates between wild and domesticated populations is
polymorphism in genes like those previously demonstrated to suppress recombination in
Arabidopsis thaliana (Girard et al. 2015, Fernandes et al. 2018a). The differences in
recombination rates between wild and domesticated populations is just one of the possible
questions that can be touched on with this system.
130
The ten populations of *T. cacao* also allow us to compare the locations of hotspots between
them, potentially contributing to the understanding of hotspot turnover at the population-
divergence timescale. These comparisons can also contribute to our understanding of how
demographics impact the turnover of recombination hotspot locations. *T. cacao* is unique in this
case for being a long-lived organism with no known driver of recombination hotspots (e.g.
PRDM9). What contributes to the location of recombination hotspots in such a species is, of
course, contingent on our being able to detect recombination hotspots in the different
populations of *T. cacao*.
139
In order to locate recombination hotspots for *T. Cacao* populations, we must first obtain fine-
scale recombination maps for each population, which we did using an LD-based method. Fine-
scale, LD-based recombination maps have been constructed for a number of plant models
(Paape et al. 2012, Choi et al. 2013, Hellsten et al. 2013), identifying a variety of features
correlated to recombination rate. Unlike these model plants with short generation times, *T.*
Cacao is a perennial woody plant with a five-year generation time (Henderson et al. 2007). The
size and long generation time of *T. Cacao* makes direct measurements of recombination
impractical. However, historical recombination can be estimated for *T. Cacao* using coalescent
based methods (Auton and McVean 2007). Theoretical studies have shown that population
structure can generate artificially inflated measures of LD (Li and Nei 1974, Ohta 1982), which
would be detrimental to our estimates of recombination. For this reason, recombination maps
were constructed independently for each population. For each population we aim to describe

the relationship between recombination hotspots and a variety of evolutionary and genomic
factors.
154
We used an LD-based method to estimate recombination rates for ten populations of *T. Cacao*,
which we then analyzed with a maximum likelihood statistical framework to infer the location of
recombination hotspots. The locations of hotspots were compared across populations and a
novel resampling scheme tailored to the genomic architecture of *T. Cacao* was used to generate
null assumptions for the distribution of hotspots along the genome. These null distributions were
used to identify differential representation of known DNA sequence motifs in ubiquitous
recombination hotspots, and of overlap between recombination hotspots and genomic traits for
each population. The re-sampling schemes used to identify these associations are novel in the
context of this work and were designed to take into account the size and distribution of elements
in the genome. In this work we aimed to answer the following questions: (i) How are
recombination rates distributed within 10 highly differentiated populations of *T. Cacao*, and how
do they compare to each other? (ii) How are hotspots distributed along the genome of each of
the ten populations of *T. Cacao*, and can these distributions be explained by patterns of
population genetic differentiation? (iii) Are there identifiable DNA sequence motifs that are
associated with the location of recombination hotspots along the *T. Cacao* genome? (iv) Are
there genomic features (e.g. TSSs, TTSs, exons, introns) consistently associated with
recombination hotspot locations across *T. Cacao* populations? Our findings suggest that
recombination hotspot locations generally follow patterns of diversification between populations,
while also having a strong tendency to occur close to TSSs and TTSs. Moreover, we find a
strong negative association between the occurrence of recombination hotspots and the
presence of retroelements.
176
177 **Results:**
178
179 *Comparing recombination rates between populations*
180
Populations show a mean recombination rate (r) between 2.1 and 525 cm/Mb (Table 1), with a
variety of distributions (Fig. S2). We observe a higher mean than median r indicating that
extreme high values are present for all populations. The extreme recombination rate values
affect the mean, driving it to values consistently higher than the median. The pattern of
recombination rates along the genome varied between populations, as can be seen in the
comparison of the Nanay and Purus third chromosome (Fig. 1). Purus appears to have a higher
average recombination rate than Nanay for chromosome three. More specifically, particular
regions of the chromosome present peaks in one population that are absent in the other. A
similar patter can also be observed for the density of recombination hotspots, e.g. Purus
presenting a high density of hotspots in certain regions that is not observed in Nanay. The
median 95% probability interval for recombination rate across the genome for each population
was found to be several orders of magnitude larger than the uncertainty per site, estimated as
the median 95% Credibility Interval of the trace for each position in the genome for that
population (Table S1).
195
Overall, the mean recombination rate for most of the populations is similar to that found for
Arabidopsis thaliana using LDhat, when using θ=0.1 (Choi et al. 2013) (Table 1). The LDhat
estimates using θ=0.001 were slightly higher than the estimates using θ=0.1 for each
population. We chose to proceed with analyses using the results from the θ=0.1 since more of
the mean population recombination rates fell within the range of values identified in plants
(Stapley et al. 2016) (Table S2).
202

Table 1. Recombination rates in $\rho = 4Ner$ (in Morgans per base) and r (in cM/Mb) for all ten *T. cacao* populations. The *Ne* (from Cornejo et al., 2018) used to transform ρ to r for each population is also reported, as are the lower and upper bounds of a 95% confidence interval for mean r.

Population	Mean $4N_er$	Mean N_e	Mean r (cM/Mb)	Median r (cM/Mb)	L bound (Mean r)	U bound (Mean r)
Amelonado	1.58e-03	15,744	2.51	2.40e-04	2.48	2.54
Contamana	8.53e-03	61,102	3.49	4.92e-01	3.48	3.50
Criollo	1.46e-04	695	525	427	523	527
Curaray	1.04e-04	58,213	4.45	1.78	4.44	4.46
Guianna	8.66e-03	4,651	46.5	7.74e-01	46.3	46.7
Iquitos	4.23e-03	49,984	2.11	5.88e-04	2.10	2.12
Maranon	4.09e-03	34,037	3.01	1.64e-03	2.99	3.02
Nacional	4.66e-03	26,060	4.47	9.76e-03	4.44	4.49
Nanay	6.82e-03	42,429	4.02	1.51e-02	4.00	4.04
Purus	5.95e-03	17,357	8.57	7.74e-01	8.54	8.60

Figure 1. The third chromosomes of the Nanay (A) and Purus (B) populations were selected to exemplify the differences between populations in recombination rates (*r*) and recombination hotspot locations (bars above rates).

In order to compare the average recombination rates of the different populations, a Wilcoxon signed-rank test was performed for every pair of populations. The only pair that did not show a

218 significant difference in mean recombination rate was that of Nacional and Nanay (*p=0.3*). All
219 other pairwise comparisons were highly significant (*p<2e-16*).
220
Two populations, Guianna and Criollo, have a higher average recombination rate than the other
populations by one and two orders of magnitude respectively (Table 1). Guianna and Criollo
also have been estimated to have a lower effective population size (N_e) (Cornejo et al. 2018) by
one and two orders of magnitude respectively. However, there was no significant association
between mean N_e and r (*p=0.1119*), indicating that, for a high enough N_e, the ability to detect
recombination events is not dictated by the effective population size. When Criollo and Guianna
were excluded, the relationship was also not present (*p=0.3886*). When all populations were
included, the inbreeding coefficient (*F*, from Cornejo et al. 2018) showed no significant linear
association with mean r (*p=0.3361*). We also found no linear trend between sample size and
mean r (*p=0.2333*).
231
The FIGL1 and FLIP proteins characterized by Fernandes et al. (2018a) were found to be
responsible for recombination suppression in *Arabidopsis*. Plants with a FIGL1 knockout were
found to increase recombination rates significantly and FLIP knockouts show increases of
recombination at a much lesser extent (Fernandes et al. 2018). Therefore, we explored the
possibility that missense FIGL1 and FLIP orthologs in *T. cacao* explain the between-population
differences in recombination rate. We used a reciprocal BLAST search to identify the orthologs
for both genes and used annotation data from Cornejo et al. (2018) to identify 15 missense
mutations in FIGL1 and 18 missense mutations in FLIP (Fig. 2, Fig. S3, Table S3). We then
used a generalized linear model framework to infer the impact of the 8 uncorrelated missense
mutations found in the *T. cacao* FIGL1 ortholog under the assumption of a full recessive model.
We find that mutations 215KK (Coeff.=426.54, p=2.41e-13), 155II (Coeff.=8.97, p=0.000231),
and 291TT (Coeff.=0.47, p=0.047) significantly explain changes in the recombination rate, but
all other mutations made no significant impact. The same model was run for FLIP but returned
no significant coefficients (after eliminating perfectly correlated mutations with those found in
FIGL1).
247

248
249 **Figure 2.** The left panel shows the frequency of individuals that are homozygous for the alternative allele
250 of amino acid mutations in a *T. cacao* FIGL1 ortholog. Alternative allele is defined in terms of the
251 Amelonado reference genome. The right panel shows the log_e transformed recombination rates (*r*). The
252 populations are in the same order in both panels.
253

  Comparing recombination hotspot locations between populations

256 The majority (55.5%) of hotspots identified were not shared between populations. The 25 most
257 numerous sets of hotspots are represented in Fig. 3. The nine largest of these are sets of
258 hotspots unique to single populations. The hotspots unique to the remaining population (Criollo)
259 formed the eleventh largest set. Effective population size (N_e) is not a good linear predictor of
260 the number of detected hotspots (*p=0.1489*), nor is sample size (*p=0.351*).

262
Figure 3. Upset plot showing number of hotspots in different subsets. Horizontal bars
represent total hotspots detected in a population, each dot on the matrix indicate that
the vertical bar above it is the count of hotspots unique to that population, connected dots
indicate that the vertical bar above them represents hotspots shared between the populations
represented by the connected dots. The 25 largest subsets are shown.

The recombination rate in hotspot regions for nine of the populations was on average between
22 and 237% higher than the average recombination rate of the genome. The exception was
Guianna, which only showed an approximately 1% increase in average recombination rate in

hotspots regions when compared to that of the non-hotspot regions. For Guianna, we also
compared the recombination rate inside hotspots to their surrounding regions (+/- 5kb). We
found that hotspot regions had a rate ~42% higher rate than their neighboring regions. This
result leads us to believe that the 1% higher average recombination rate in the Guianna
hotspots when compared to the entire genome may be due to an increased ability to detect
hotspots in regions of low recombination for this population. Additionally, Guianna presents
unusually large hotspots (average 8.9 kb, Table S4), which points to an especially low resolution
in hotspot detection for this population.
280
Despite the majority of hotspots not being shared between populations, we conducted pairwise
Fisher's exact tests to verify whether there was significantly more hotspot overlap than expected
(if hotspots were randomly distributed along the genome) between populations. For most pairs
of populations, we found significantly more hotspot overlap than expected (Table 2). There were
three comparisons that did not show significantly more overlap than expected: Amelonado-
Nacional, Amelonado-Purus, and Criollo-Nacional. A Mantel test comparing distances between
populations based on shared hotspots and F_{ST} values between populations resulted in a
significant correlation between them (*r=0.66*, *p=0.002*).
289
290 **Table 2.** Fisher's exact test p-values for pairwise comparisons of recombination hotspot locations
291 between populations of *T. cacao*. We conducted 45 comparisons, corresponding to a Bonferroni
292 correction cutoff value of *α=0.0011*.

Population	Ame	Con	Cri	Cur	Gui	Iqu	Mar	Nac	Nan
Amelonado	-	-	-	-	-	-	-	-	-
Contamana	<2e-07	-	-	-	-	-	-	-	-
Criollo	<9e-05	<5e-13	-	-	-	-	-	-	-
Curaray	<3e-05	<3e-37	<5e-08	-	-	-	-	-	-
Guianna	<3e-06	<1e-37	<7e-07	<4e-20	-	-	-	-	-
Iquitos	<4e-08	<6e-87	<2e-11	<3e-16	<2e-29	-	-	-	-
Maranon	<6e-13	<7e-77	<2e-11	<2e-20	<5e-33	<4e-64	-	-	-
Nacional	0.0015	<2e-43	0.0212	<7e-14	<3e-06	<6e-14	<3e-13	-	-
Nanay	0.0004	<2e-44	<9e-11	<4e-16	<2e-21	<3e-39	<2e-38	<9e-06	-
Purus	0.1782	<4e-117	<2e-05	<2e-29	<1e-33	<2e-39	<8e-43	<6e-27	<2e-21

293
294
To study the effects of demographic history more closely, shared hotspots were converted to
dimensions of a multiple correspondence analysis and modeled along a previously constructed
drift tree (Cornejo et al. 2018). Modeling the dimension as a Brownian motion was a better fit
(AIC=79.4) than modeling it as an Ornstein-Uhlenbeck (OU) process (AIC=81.4), which is
consistent with the small number of hotspots shared between populations. The model assuming
Brownian motion is consistent with pure drift driving differentiation of a trait along a genealogy,
while an OU process is consistent with a higher trait maintenance (stabilizing selection).
302
303 *Identifying DNA sequence motifs associated with the locations of recombination hotspots*
304
We used RepeatMasker to analyze the set of recombination hotspots that were present in at
least eight *T. Cacao* populations (17 total hotspots; referred to as ubiquitous hotspots), as well
as the consensus set of recombination hotspots and the reference genome. In order to
determine whether a particular set of DNA sequence repeats was overrepresented in ubiquitous
hotspots, the percentage of DNA sequence that was identified as potentially being from
retroelements or DNA transposon was compared to an empirical distribution. The percentage of
observations from the distribution which were greater than the observed are reported in Table 3.

While retroelements were found to be underrepresented in the ubiquitous hotspots, DNA
transposons were marginally overrepresented.
314
Table 3. Percentage of DNA sequences identified as either retroelements or DNA transposons, and total
interspersed repeats. Observed values for the entire *T. cacao* genome, for all recombination hotspots
(HS), and ubiquitous hotspots (hotspots in the same location in at least eight different populations). Also
presented are mean percentage of these sequences for 1000 simulations of hotspots equivalent in size
and count as the ubiquitous set and the percentile at which the observed value for the ubiquitous set is
found in the distribution of the simulated set (Sim).
321

Measures	Observed % ubiquitous HS	Observed % all HS	Observed % whole genome	Mean % Sim	% Sim > ubiquitous
Retroelements	2.34	9.45	11.12	11.11	99.9
DNA transposons	1.94	1.64	1.10	1.10	5.4
Total	4.28	11.09	12.21	12.22	99.7

322
323
324
325 *Identifying genomic features associated with the location of recombination hotspots*
326
We found an overrepresentation of recombination hotspots at transcriptional start sites (TSSs)
and transcriptional termination sites (TTSs) in all ten of the *T. Cacao* populations (Table 4). The
level of overrepresentation of hotstpots in particular regions was compared to a null expectation
based on simulations of hotspots of the same size as the ones detected, distributed randomly
along the chromosomes. For all populations, all 1000 simulations showed a lower proportion of
overlap with TSSs and TTSs than the observed. In the case of exons and introns, seven
populations (Contamana, Criollo, Iquitos, Maranon, Nacional, Nanay, Purus) had an observed
value that was lower than all, or almost all (Purus for exons), simulations. Three of the
remaining four populations (Amelonado, Curaray, and Nanay) had no clear trend in either
direction (Table 4). The final population (Guianna) showed an overrepresentation of hotspots in
both exons and introns.
338
339 **Table 4.** Proportion of simulated chromosomes that presented a lower number of hotspots intersecting
340 with TSSs, TTSs, exons, and introns than the observed chromosomes. TSSs and TTSs are considered to
341 be the 500bp upstream and downstream of transcribed regions, respectively.
342

Population	TSSs (500bp)	TTSs (500bp)	Exons	Introns
Amelonado	1	1	0.602	0.527
Contamana	1	1	0	0
Criollo	1	1	0	0
Curaray	1	1	0.346	0.058
Guianna	1	1	1	1
Iquitos	1	1	0	0
Maranon	1	1	0	0
Nacional	1	1	0	0
Nanay	1	1	0.027	0.237
Purus	1	1	0.004	0

343
344
345
346

Discussion:

The set of ten *T. cacao* populations, which includes wild, long-established and recently established populations as well as a domesticated population, has provided us a unique opportunity to study differences in recombination in populations of the same species with varying evolutionary histories (Cornejo et al. 2018, Bartley et al. 2005). We also explored differences in the location of recombination hotspots between the populations and found that the conservation of hotspots between them generally mirrors their patterns of pairwise genetic differentiation. Additionally, we found that TSSs and TTSs are strongly associated with recombination hotspots in all populations, which is consistent with previous findings in plants (Paape et al. 2012, Choi et al. 2013, Hellsten et al. 2013). This factor seems to play an important role in determining the location of novel hotspots. Finally, hotspots that are shared by at least eight populations appear to be associated with DNA transposons, pointing to a potential mechanism for the maintenance of recombination hotspots at the population-divergence timescale. Understanding how recombination rates vary between genetically differentiated populations of the same species is an important step toward disentangling the role of recombination in genetic differentiation.

Comparing recombination rates between populations

We found that the eight long-established, wild *T. cacao* populations show an average recombination rate (r) within the range of recombination rates measured for other suporrosids (Table 1; Stapley et al. 2016). The average recombination rate for these *T. cacao*, populations, was 4.07 cM/Mb, which is very similar to that of *Juglans regia* (4.05 cM/Mb; Zhu et al. 2015), and comparable to other woody rosids (e.g. *Populus deltoides*, 8.13 cM/Mb (Mousavi et al. 2016); *Mangifera indica*, 7.15 cM/Mb (Luo et al. 2016); *Citrus clementina*, 3.6 cM/Mb (Ollitrault et al 2012)) (Stapley et al. 2016). This places *T. cacao* on the high end of known recombination rates for its order but comfortably in the range of other long-lived, woody plants. For all ten populations, the mean recombination rate was found to be greater than the median. This is consistent with high rate outlier values; an expected result in the presence of recombination hotspots.

The other two *T. cacao* populations (Criollo, the domesticated variety and Guianna, the recently established wild population) show unusually high average recombination rates when compared to the eight long-established wild populations. Despite a small sample for some populations, including Criollo, we found no linear trend between sample size and recombination rate (Table S5). Additionally, the rates calculated for the two wild, small-sample populations (Curaray and Nacional) were consistent with those of other wild populations. This makes us confident in our estimates for the Criollo and Guianna populations. We also used the effective population size for *Medicago truncatula* from Siol et al. 2007 and the estimate of ρ from Paape et al. 2012, to calculate r for *M. truncatula* (=433 cM/Mb) and found that it was comparable with the mean rate found for the Criollo population (Table 1). One possible explanation for the higher recombination rate observed for the Criollo population is domestication; which has been observed to increase recombination rates, particularly in plants (Ross-Ibarra 2004). Previous work has shown that Criollo is the only population showing a strong signature of domestication, as revealed by its much higher drift parameter compared to other populations (Cornejo et al. 2018). The high recombination rate observed in Guianna (46.5 cM/Mb) can be explained in a similar way; while Guianna does not show a strong signature of domestication, it is the most recently established population (Bartley et al. 2005) and has undergone a recent bottleneck (Cornejo et al. 2018). We hypothesize from this result that the Guianna population is undergoing the initial stages of domestication, and its increased recombination is an early indicator of this. Another possibility is

398   that the high recombination rates estimated for Criollo and Guianna can be explained by biases
399   in estimation caused by errors associated to small samples or low genetic variation. However,
400   the recombination rates for Amelonado (another population with low variation) and Purus (a
401   population with small sample size) did not present this problem.
402
Analyses exploring mutations of putative recombination suppression genes (Fernandes et al.
2018b) could help disentangle the nature of this extreme variation in recombination rate in the
Criollo and Guianna populations. It has been widely discussed that although actual
recombination rates (chiasmata per bivalent) might increase in domestic plant populations,
effective recombination is likely to be reduced in domestic populations as a result of increased
LD and long runs of homozygosity. The interpretation for this observation is that it is likely that
chromosomes might physically recombine at higher rates, but if homologous chromosomes
contain the same sequence then no appreciable exchange of variants among different
chromatid backgrounds can occur (Moyers et al. 2018). We find significantly higher estimates of
median recombination rates for domesticated Criollo populations across the genome. Given
previous research suggesting the key function of FIGL-1 on suppressing recombination rates in
Arabidopsis, we investigated if mutations in this protein could explain the differences in
recombination rate. Our results suggest that missense mutations in FIGL-1 could impair the
activity of this protein in domesticated populations, reducing its efficacy as a recombination
suppressor. The mutation showing the largest effect (KK_215) is close to the domain that has
been previously shown to be involved in the interaction of FIGL-1 with RAD51 and DMC1
(Fernandes et al. 2018a) and thus lead us to propose that FIGL-1 might be responsible for the
increase recombination rate in domesticated cacao populations. Further work using yeast two-
hybrid systems similar to the work performed by Fernandes et al. (2018a) will be necessary in
the future to show if the mutations do indeed interfere with the normal function of FIGL-1. These
mutations were not found in increased frequency in Guianna, suggesting that the underlying
mechanisms leading to increased recombination rates in this population are different to those in
the Criollo population.
426
427
Despite recombination rates for eight of the ten populations being of the same order of
magnitude, pairwise comparisons of average rates indicated that most populations have a
significantly different rate of recombination from the others. The only exception were Nacional
and Nanay whose average rates were not significantly different from each other. These two
populations, however, are not more closely related to each other than they are to other
populations, based on genetic differentiation (Cornejo et al. 2017). We interpret this result as
suggestive that their similarity is not due to genetic similarity, but some other factors, e.g.
epigenetics.
436
The likelihood of detecting hotspots of recombination in the genome will likely be affected by the
amount of uncertainty in the estimates of recombination across sites or regions. Yet, we have
been unable to identify any study where the magnitude of the uncertainty in the estimates of
recombination are assessed to address this issue. We have performed careful comparisons and
assessed the magnitude of the uncertainty in the estimation of recombination rates to show that
this uncertainty is several orders of magnitude smaller than the variation in recombination rates
across the genome (Table S1).
444
445   *Comparing recombination hotspot locations between populations*
446
Understanding the pattern and rate of change of recombination hotspots at the population level
can elucidate their role in shaping genome architecture, impacting how effectively selection

operates (Felsenstein 1974). We found that a large proportion (55.5%) of hotspots detected are
unique to a single population. The observed variability of hotspot location between populations
points to demographic history not being the main driver of recombination hotspot location.
However, the hotspots tend to appear in similar regions, as demonstrated by the Fisher's exact
tests (Table 2). This dichotomy can be explained by considering that the proportion of the
genome occupied by recombination hotspots is very low, so even a small proportion of hotspots
from two different populations being in the same region is enough for the Fisher's exact test to
recognize them as significantly similar. This small but significant similarity can occur by
recombination being limited in its possible positioning along the genome, but not to the point of
forcing hotspots to occur consistently in the same locations, and thus maintaining some level of
stochasticity.
460
Given the significant proportion of overlapping hotspots between populations, it was still
important to explore whether the similarities can be explained by shared genetic history. If
demographic history explains the evolution of hotspot location, we would expect that more
closely related populations would have a higher percent of overlapped hotspots. A significant
relationship was found between population differentiation (F_{ST}) and the distance between
populations based on shared hotspots (Mantel test, *r=0.66, p=0.002*). This result indicates that,
to some extent, the genetic differentiation and the location of hotspots are mirroring each other,
which could be due to recombination hotspots being a product of the shared history between the
populations. However, since recombination rates were estimated using a coalescent-based
method, we expect historical relationships to be represented in our findings to some extent. We
transformed the information of hotspot overlap to model hotspots as quantitative traits changing
along a population tree (Cornejo et al. 2018). Our results show that a Brownian motion model
(AIC=79.4) better fits the shared hotspot data than a model with stabilizing selection Ornstein-
Uhlenbeck model (AIC=81.4), suggesting that, in principle, drift alone could explain the
evolution of the location of recombination hotspots. However, the absolute number of hotspots
that are shared among populations indicates that demographic history is insufficient to explain
the evolution of recombination hotspots in this species.
478
While we do not detect all the hotspots in these populations and not all the hotspots detected
are necessarily true positives, this proportion of unique hotspots can be seen as an indicator
that the turnover rate for hotspots is faster than the time it took the 10 populations to
differentiate. The detection rate for LDhot is approximately 55% under constant population
conditions, and greater when a recent bottleneck has occurred (Auton et al. 2014, Dapper and
Payseur 2017). Only two of the populations in this study (Criollo and Guianna) have a known
recent bottleneck (Cornejo et al. 2018). However, Criollo is the only one of these two with an
unusually low hotspot count (Table S4). Criollo's low number of detected hotspots can be a
product of its increased genome-wide recombination rate, making the signal of hotspots less
pronounced. In a similar fashion, the increased overall recombination rate of Guianna may be
affecting the detectability of hotspot regions, limiting our ability to resolve the limits of hotspots.
It is important to note that our hotspots are unusually large for all populations (Table S4), which
is likely a product of our low sample size leading to low resolution when resolving hotspot
regions. Further work, increasing the sample size per population will contribute to increasing the
resolution of these estimates.
494
While shared recombination hotspots can, to some extent, be explained by patterns of genetic
differentiation, some of the sharing can simply be due to a tendency for hotspots to arise near
TSSs and TTSs. It has been observed in other organisms that hotspots of recombination are
frequently associated to specific genomic features (including TSSs and TTSs) (Auton et al.
2013, Choi et al. 2013, Hellsten et al. 2013, Myers et al. 2005, Singhal et al. 2015) or DNA

sequence motifs (Auton et al. 2012, Brunschwig et al. 2012, Stevison et al. 2016). These factors
can affect the landscape of recombination, contributing to the patterns of shared hotspot
locations between populations that we observe in *T. cacao*.
503
Our ability to measure recombination in ten distinct populations allows us to analyze the
relationship between population genetic processes and recombination. Our results suggest that
the pattern of gains and losses of recombination hotspots is very dynamic and the landscape of
recombination changes rapidly during the process of diversification within a species. This
dynamism can have a tremendous impact on the adaptive dynamics of a species, and it should
be taken into account, considering that theoretical studies tend to assume that recombination
rates are constant during the evolution of populations (Hudson and Kaplan 1988, Donnelly and
Kurtz 1999).
512
513   *Identifying DNA sequence motifs associated with the locations of recombination hotspots*
514
The analysis of 17 hotspots shared between at least eight populations of *T. cacao* found an
underrepresentation of retroelements and a marginal overrepresentation of DNA transposons
when compared to the entire genome (Table 3). These results are not entirely surprising as it
has already been suggested that transposable elements (TEs) tend to be enriched in areas of
low recombination in *Drosophila* as a consequence of selection against TE activity that could
lead to chromosome instability (Rizzon et al. 2002). However, the marginal over-representation
of DNA transposons in the most conserved recombination hostspot is unexpected, given that all
previous observations have shown a reduced representation of mobile elements in areas with
high recombination rate (Rizzon et al. 2002). It is possible that DNA transposons are at least
partly responsible for the maintenance of recombination hotspots as populations diverge, from
which we expect that site-directed recombination is more frequent in these locations of the
genome. However, the low percentage of these sequences observed in the set of all hotspots
(Table 3) indicates that these sequences only have a small effect on the maintenance of
hotspots. It has been observed in humans that short DNA motifs enriched for repeat sequences
determine the location of 40 per cent of hotspots enriched for recurrent non-allelic homologous
recombination (McVean 2010). One potential explanation for why natural selection does not
eliminate hotspots in these regions is the possibility that these regions do not produce a large
enough mutational load for natural selection to remove them from the population (McVean2010
et al.).
534
535   *Identifying genomic features associated with the location of recombination hotspots*
536
For all ten populations, an overrepresentation of hotspots was found in the areas immediately
preceding and following transcribed regions of the chromosome. This matches the findings of
previous studies in *Arabidopsis thaliana* (Choi et al. 2013), *Taenipygia guttata* and *Poephila*
acuticauda (Singhal et al. 2015), and humans (Myers et al. 2005). The most likely explanation is
that recombination events within genes are selected against. The rationale being that a
recombinant chromosome that undergoes a double-strand break in the middle of a coding
region will have a higher risk of being inviable, and therefore not represented in the current set
of chromosomes for its population. Recombination occurring in transcription start and stop sites,
on the other hand, does a much better job at breaking up haplotypes or shuffling alleles in
different genomic backgrounds, while preserving the functionality of coding regions. This
rationale is supported by previous findings of increased recombination rates in these regions
(Choi et al. 2013). It is also supported by results from PRDM9 knock-out *Mus musculus*, which
has shown a reversion to hotspots located near TSSs (Brick et al. 2011). The enrichment of *T.*

cacao hotspots in TSSs and TTSs is thus a reasonable result given that zinc-finger binding
motifs and potential modifiers like PRDM9 have not been identified in this species.
552
553   *Implications for the evolutionary history of* T. cacao
554
Overall, our results show a large, consistent pattern where recombination rates in the ten
populations of *T. cacao* are of a similar magnitude as mutation rates but show a high diversity in
location and number of hotspots of recombination that cannot be explained solely by the
process of diversification of the populations. A potential hypothesis that could explain the rapid
turnover of hotspots of recombination and the relative differences in recombination among
populations is that epigenetic changes are involved in controlling the turnover of recombination
in plants. This hypothesis is not unreasonable given the recent observation of epigenetic control
of recombination in plants (Yelina et al. 2015). Further theoretical and simulation work should be
done in order to better understand the implications of the rapidly changing recombination
hotspots in adaptive dynamics. We also show that there is an overall underrepresentation of
hotspots in exons and introns for most populations, which is consistent with purifying selection
acting against changes that could result in disruptions of gene function. On the other hand, we
observed an overrepresentation of hotspots in TTSs and TSSs for all ten populations. This
could impact the maintenance and spread of beneficial traits in the population by shuffling allelic
variants of genes without causing disruption of their function. We hypothesize that the
enrichment of hotspots of recombination in TTSs and TSSs can have an important impact in the
spread of beneficial mutations across different genomic backgrounds; increasing the rate of
adaptation to selective pressures (e.g. selection for improved pathogen response).
573
574   **Materials and Methods:**
575
576   *Comparing recombination rates between populations*
577
Sequence data were downloaded from the Cacao Genome Database and NCBI (Accession
PRJNA486011), including the reference sequence for each chromosome and the full genome
annotation (*Theobroma cacao* cv. Matina 1-6 v1.1) (Motamayor et al. 2013). Processing was
done using the pipeline from (Cornejo et al 2018) available at the github repository
oeco28/Cacao_Genomics. Full genome data was used from a total of 73 individuals (146
chromosomes) across 10 populations (Cornejo et al. 2018). We filtered single nucleotide
polymorphism data and excluded rare variants (minor allele frequency <= 0.05) per population.
Separate variant files per population per chromosome were then phased using default
conditions with SHAPEIT2 (Delaneau et al. 2011) under default parameters. Haplotype files
were converted back to phased variant calling format (vcf) for its downstream analysis. We have
also phased the data with Beagle (Browning and Browning. 2007), using a burn-in of 10000
iterations, and estimations done over 10000 iterations. No appreciable differences were
observed between the two methods and Beagle phasing was maintained for the analyses. The
reason for performing the phasing separately for each population is that linkage disequilibrium
patterns are expected to be affected by population structure. The ten populations have been
shown to be unique clusters with very little admixture between them (Cornejo et al. 2018), and
the individuals used in this study were those whose ancestry was clearly from a single
population. VCFTools (Danecek et al. 2011) was used to remove all singletons and doubletons
(i.e. SNPs where the minor allele count is less than three). Only bi-allelic single nucleotide
polymorphisms (SNPs) were retained and were exported in LDhat format. The sample size and
post-filtering SNP count for each population can be found in Table S5.
599

In order to estimate recombination rates, we used the *interval* routine from LDhat (Auton and
McVean 2007), a program that implements coalescent resampling methods to estimate
historical recombination rates from SNP data. To reduce computation time, each chromosome
was split into windows, each containing 2000 SNPs. To counteract the overestimation of
recombination rate produced at the ends of the windows, an overlap of 500 SNPs was left
between consecutive windows. The final window for each chromosome did not always match
the general scheme, so the final 2000 SNPs were taken (making the overlap with the second to
last window variable, but never less than 500 SNPs) (Fig. S4). Once these windows were
generated, LDhat was run over each window using 100 million iterations, sampling every 10000
iterations (10000 total points sampled), with a block penalty of 5. Lookup tables with a grid of
100 points, population mutation rate parameters (θ) of 0.1 and 0.001, and a number of
sequences (n) of 50 were used for all populations. Downstream analyses were conducted using
the results from the θ=0.1 runs of LDhat. We used the same θ for all populations since
estimates from (Cornejo et al. 2018) ranged from π=0.27% to π=0.37%, all comfortably within an
order of magnitude of each other. The first 50 million iterations were discarded as burn-in. Once
recombination rates were calculated, 250 positions were cut off from both windows involved in
each overlap, so that the estimates for the first half of the overlap was taken from the end of the
preceding window and the estimates for the second half of the overlap were taken from the
beginning of the following window. The final overlap in each chromosome was split in order to
remove 250 SNPs from the second to last window, regardless of the remaining size of the last
window. The remaining rate estimates were then merged in order to obtain recombination rates
for the entire chromosome. This was done for each chromosome of each population.
622
The estimation of recombination rates with LDhat is approximated using a sampling scheme
with a Markov Chain Monte Carlo (MCMC) algorithm as implemented in the *interval* routine. The
inference of recombination rates is the result of the integration of estimated parameter values
across iterations with the routine *stats*. In the majority of recent studies where LDhat or
LDhelmet are used (Myers et al. 2005, Auton et al. 2012, Brunschwig et al. 2012, Paape et al.
2012, Auton et al. 2013, Choi et al. 2013, Singhal et al. 2015, Stevison et al. 2016), whether
there is convergence of the Markov chains has not been explicitly investigated. One study that
we are aware of has used simulations to assess whether their small sample size affected their
ability to obtain reliable estimates of recombination using LDhelmet (Booker et al. 2017) but did
not assess the uncertainty of the estimates from the MCMC process itself. We argue that
evaluation of convergence is important to assess the confidence in the estimated reported
values, especially if there is interest in analyzing the differences in recombination rate along the
genome. Visual inspection of pilot runs of the analysis demonstrated that convergence was not
achieved after running 40M iterations, which is why the length of the chains was increased to
100M iterations. Additionally, we explored the uncertainty in the estimates of recombination site-
wise by integrating over the trace of the estimates for recombination rate to infer the 95%
Credibility Interval. We then estimated the 95% interval of recombination estimates range
across all sites in the genome to have an overall measure of uncertainty that we compared to
the median 95% Credibility Interval for the trace of each position.
642
643   In order to compare recombination rates, the effective population size (N_e) calculated for each
644   population (Cornejo et al. 2018) was used to convert rates in $N_e r$ to r. We tested the difference
645   between the whole-genome mean rate of recombination (r) between populations using the
646   Wilcoxon signed-rank test `wilcox.test` function from the `stats` package in `R`) (R core team
647   2018). There were 45 comparisons, making the Bonferroni correction cutoff value: α=0.0011.
648
Suppressors of recombination identified in Arabidopsis and other systems, FIGL1 and FLIP,
have not yet been identified in *T. cacao* (Motamayor et al. 2013). We performed a reciprocal

BLAST search using a newly generated databased containing sequences of FIGL1 and FLIP
obtained from ncbi (Altschul et al. 1990, Altschul et al. 1994b, Shiryev et al. 2007). After
identification of orthologous copies of FIGL1 and FLIP, we extracted the annotated variants
responsible for missense mutations in the proteins from data generated in Cornejo et al. 2018.
We then used these mutations to estimate the frequency of homozygous genotypes with
alternative alleles (different to the reference). Because the reference genome belongs to the
Amelonado populations, the population that presents the lowest estimated median
recombination rate in our work, we used the frequency of homozygous alternatives to infer the
impact of missense mutations under the assumption of a recessive model on the recombination
rate. For this, we estimated the homozygosity for missense mutations and eliminated those
found in complete correlation ($r^2 = 1$) and fitted a generalized linear model of the form:
662
663    $$y = \beta_0 + \beta_j + \varepsilon_j.$$
664
665   Where y is a vector of the median recombination rate accross populations, b_0 is the intersect (in
666   this case the Amelonado median recombination rate) and b_j is the effect of the homozygosity in
667   position j. We assumed a Gaussian link function for the model.
668
669
670   *Comparing recombination hotspot locations between populations*
671
Recombination hotspots were estimated with LDhot (Auton and McVean 2007), a likelihood-
based program that tests whether a single distribution model or a two-distribution model better
explains the observed recombination rates in 1 kb sliding windows (default), for each
chromosome. Each chromosome was run in its entirety, with the number of simulations (nsims)
set to 1000. The resulting potential hotspots were refined by an α of 0.001, and overlapping
hotspots were merged. This method therefore detects hotspots by comparing rates in 1 kb
windows to the rates in the surrounding regions.
679
To determine the set of consensus hotspots, the hotspots from all populations were merged.
Two hotspots from different populations were considered to be shared if they both overlapped
with the same hotspot in the consensus set. To summarize all shared hotspots, a Boolean
matrix was constructed, in which a population having a hotspot that overlaps with a hotspot in
the consensus list leads to an indication of presence of the consensus hotspot in that
population. This matrix was used to determine hotspots shared by two or more populations.
686
687   A Fisher's exact test was run for each pair of populations in order to determine whether hotspots
688   for the pair of populations overlap significantly more than expected. The BED files containing
689   the location of the recombination hotspots for each pair of populations were compared using
690   Bedtools:fisher (Quinlan and Hall. 2010). The number of comparisons was 45, making the
691   Bonferroni correction cutoff value: α=0.0011.
692
In order to compare the relationships between populations based on shared hotspots we
calculated Jaccard distances (`distance` function, `philentropy` package, `R`) (Drost 2018) and
compared them to a published F_{ST} matrix (Cornejo et al. 2018) using a Mantel test
(`mantel.rtest` function, `ade4` package, `R`) (Chessel et al. 2004, Dray et al. 2007a, Dray et al.
2007b, Bougeard and Dray 2018). The F_{ST} estimates from (Cornejo et al. 2018) were generated
using Weir and Cockerham's estimator (Weir and Cockerham 1984).
699

In order to model the presence or absence of hotspots along a drift tree, a multiple
correspondence analysis was used on the Boolean matrix of shared hotspots using the MCA
function from the FactoMineR package in R (Le et al. 2008). Nine dimensions were retained
and used as traits along a previously generated drift tree (Cornejo et al. 2018). Using the
Rphylopars package in R (Goolsby et al. 2016), the dimensions were modeled as Brownian
motion and as an Ornstein-Uhlenbeck process. The fit of the two models were compared using
the AIC values for the best fitting models of each type.
707
708 *Identifying DNA sequence motifs associated with the locations of recombination hotspots*
709
Motifs associated with hotspots were found using RepeatMasker (Smith et al. 2016). The entire
genome, the set of consensus hotspots, and a set of ubiquitous hotspots (hotspots shared by at
least eight of the populations) were examined with RepeatMasker, using normal speed and
"theobroma cacao" in the species option. In order to determine whether ubiquitous hotspots
were enriched for particular DNA sequences, a set of the same number and size of sequences
was randomly selected from the genome using Bedtools:shuffle (Quinlan and Hall. 2010) and
examined with RepeatMasker. This simulation was repeated one thousand times and a null
distribution against which observed values were compared was constructed from the results.
718
719 *Identifying genomic features associated with the location of recombination hotspots*
720
Testing whether recombination hotspots were overrepresented near particular genomic features
was done by using a resampling scheme to establish null expectations and then comparing the
observed value to the empirical distribution. For each feature, locations were retrieved and the
number of observed hotspots that overlap with this feature were counted. To determine whether
this number of overlapping hotspots was unusually high or low, a set of hotspots that matched
the number of hotspots and the size of each hotspot was simulated. These simulated hotspots
were placed randomly along the chromosome, using a uniform distribution. The simulation was
run 1000 times and the number of simulated hotspots that overlap with the true genomic
features was measured for each simulation. The simulations generate an expected distribution
of overlap with the genomic feature, and the true value was then compared to the distribution.
When simulated hotspots overlapped, the location of one of them was sampled again. Features
tested were: Transcriptional start sites (TSSs), transcriptional termination sites (TTSs), exons,
and introns. TSSs and TTSs are considered to be the 500bp upstream and downstream of
coding regions respectively.
735
736 The reason for the proposed novel resampling scheme is that, if the size and distribution of
737 genomic features and hotspots were not taken into account, it would set unrealistic expectations
738 for the overlap between features under a null model of no association. In this sense, the null
739 model would be inappropriate and potentially inflate the false positive rate.
740
741 **Data Access:**
742
743 Rate and summary files from LDhat runs as well as hotspots for each population will be placed
744 in a Dryad repository. Scripts for LDhat and LDhot runs as well as the resampling schemes
745 used and additional analysis is available in the following github repository:
746 *ejschwarzkopf/recombination-map*.
747
748
749
750

Acknowledgements:

The authors would like to thank the Noe Higinbotham endowment and the WSU College of Arts and Science for travel funds to EJS to present earlier versions of this work. We would like to thank the Kamiak High Performance Computing Cluster at WSU for the infrastructure support to run the analyses, and the Cornejo, Kelley, and Busch labs at WSU for feedback and edits on the manuscript.

References:

Akhunov E, Goodyear A, Geng S, Qi L, Echalier B, Gill B, Gustafson J, Lazo G, Chao S, Anderson O, et al. 2003. The Organization and Rate of Evolution of Wheat Genomes Are Correlated With Recombination Rates Along Chromosome Arms. *Genome Res*. **13**:753–763. Doi:10.1101/gr.808603.

Altschul S, Gish W, Miller W, Myers E, Lipman D. 1990. Basic local alignment search tool. *J Mol Biol*. **215**:403-410.

Altschul S, Boguski M, Gish W, Wootton J. 1994. Issues in searching molecular sequence databases. *Nature Genet*. **6**:119-129

Anderson L, Salameh N, Bass H, Harper L, Cande W, Weber G, Stack S. 2004. Integrating Genetic Linkage Maps With Pachytene Chromosome Structure in Maize. *Genetics*. **166**:1923–1933.

Auton A, McVean G. 2007. Recombination rate estimation in the presence of hotspots. *Genome res*. **17**:1219–1227.

Auton A, Myers S, McVean G. 2014. Identifying recombination hotspots using population genetic data. bioRxiv doi: 1403.4264v1.

Auton A, Fledel-Alon A, Pfeifer S, Venn O, Ségurel L, Street T, Leffler EM, Bowden R, Aneas I, Broxholme J, et al. 2012. A fine-scale chimpanzee genetic map from population sequencing. *Science*. **336**:193–198. doi:10.1126/science.1216872.

Auton A, Rui Li Y, Kidd J, Oliveira K, Nadel J, Holloway JK, Hayward JJ, Cohen PE, Greally JM, Wang J, et al. 2013. Genetic Recombination Is Targeted towards Gene Promoter Regions in Dogs. *PLoS Genet*. **9**. doi:10.1371/journal.pgen.1003984.

Bartley BGD. 2005. *The genetic diversity of cacao and its utilization*. CABI. Wallingford, United Kingdom.

Begun DJ, Aquadro CF. 1992. Levels of naturally occurring DNA polymorphism correlate with recombination rates in *D. melanogaster*. *Nature*. **356**. doi:10.1038/356519a0.

Booker TR, Ness RW, Keightley PD. 2017. The Recombination Landscape in Wild House Mice Inferred Using Population Genomic Data. *Genetics*. **207**:297–309. doi:10.1534/genetics.117.300063.

Bougeard S, Dray S. 2018. Supervised Multiblock Analysis in R with the ade4 Package. *J Stat Softw*. **86**:1–17. doi:10.18637/jss.v086.i01.

Branca A, Paape T, Zhou P, Briskine R, Farmer A, Mudge J, Bharti A, Woodward J, May G, Gentzbittel L, et al. 2011. Whole-genome nucleotide diversity, recombination, and linkage disequilibrium in the model legume Medicago truncatula. *P Natl Acad Sci USA*. **108**:E864–E870.

Brick K, Smagulova F, Khil P, Camerini-Otero RD, Petukhova GV. 2012. Genetic recombination is directed away from functional genomic elements in mice. *Nature*. **485**:642–645. doi:10.1038/nature11089.

Browning SR, Browning BL. 2007. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering.

Am J Hum Genet. **81**.

Brunschwig H, Levi L, Ben-David E, Williams R, Yakir B, Shifman S. 2012. Fine-Scale Maps of
Recombination Rates and Hotspots in the Mouse Genome. *Genetics*. **191**:757–64.

Chessel D, Dufour A, Thioulouse J. 2004. The ade4 Package – I: One-Table Methods. *R News*,
4, 5–10.

Choi K, Zhao X, Kelly K, Venn O, Higgins J, Yelina N, Hardcastle T, Ziolkowski P, Copenhaver
G, Franklin F, et al. 2013. *Arabidopsis* meiotic crossover hot spots overlap with H2A.Z
nucleosomes at gene promoters. *Nature Genet*. **45**:1327–36.

Cornejo OE, Yee M-C, Dominguez V, Andrews M, Sockell A, Strandberg E, Livingstone D,
Stack C, Romero A, Umaharan P, et al. 2018. Population genomic analyses of the
chocolate tree, *Theobroma cacao* L., provide insights into its domestication process.
Commun biol. **1**:167–167. doi:10.1038/s42003-018-0168-6.

Crow JF, Kimura, Motoo. 1970. An introduction to population genetics theory. Harper & Row,
New York.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics*.
27:2156–2158. doi:10.1093/bioinformatics/btr330.

Dapper AL, Payseur BA. 2018. Effects of Demographic History on the Detection of
Recombination Hotspots from Linkage Disequilibrium. *Mol Biol Evol*. **35**:335–353.
doi:10.1093/molbev/msx272.

Delaneau O, Marchini J, Zagury J. 2012. A linear complexity phasing method for thousands of
genomes. *Nat Methods*. **9**:179–81. doi:10.1038/nmeth.1785.

Donnelly P, Kurtz TG. 1999. Genealogical Processes for Fleming-Viot Models with Selection
and Recombination. *Ann Appl Probab*. **9**:1091–1148.

Dray S, Dufour A. 2007. The ade4 Package: Implementing the Duality Diagram for Ecologists. *J
Stat Softw*, **22**:1–20. doi: 10.18637/jss.v022.i04.

Dray S, Dufour A, Chessel D. 2007. The ade4 Package – II: Two-Table and K-Table Methods. *R
News*, **7**:47–52.

Drost HG. 2018. Philentropy: Information Theory and Distance Quantification with R. *J Open
Source Softw*. **3**:765.

Eyre-Walker A, Keightley P. 2007. The distribution of fitness effects of new mutations. *Nat Rev
Genet*. **8**:610–618. doi:10.1038/nrg2146.

Felsenstein J. 1974. The evolutionary advantage of recombination. *Genetics*. **78**:737–56.

Fernandes J, Duhamel M, Seguéla-Arnaud M, Froger N, Girard C, Choinard S, et al. 2018a.
FIGL1 and its novel partner FLIP form a conserved complex that regulates homologous
recombination. *PLoS Genet.* **14**:e1007317. doi:10.1371/journal.pgen.1007317

Fernandes JB, Séguéla-Arnaud M, Larchevêque C, Lloyd AH, Mercier R. 2018b. Unleashing
meiotic crossovers in hybrid plants. *P Natl Acad Sci USA*. **115**:2431–2436.
doi:10.1073/pnas.1713078114.

Girard C, Chelysheva L, Choinard S, Froger N, Macaisne N, Lehmemdi A, et al. 2015. AAA-
ATPase FIDGETIN-LIKE 1 and Helicase FANCM Antagonize Meiotic Crossovers by
Distinct Mechanisms. *PLoS Genet.* **11**: e1005369. doi:10.1371/journal.pgen.1005369

Goolsby EW, Bruggeman J, Ané C. 2017. Rphylopars: fast multivariate phylogenetic
comparative methods for missing data and within-species variation. *Methods in Ecol
Evol*. **8**:22–27. doi:10.1111/2041-210X.12612.

Gore MA, Chia J-M, Elshire RJ, Sun Q, Ersoz ES, Hurwitz BL, Peiffer JA, Mcmullen MD, Grills

GS, Ross-Ibarra J, et al. 2009. A First-Generation Haplotype Map of Maize. *Science*.
326:1115–1117. doi:10.1126/science.1177837.
Haldane J. 1937. The Effect of Variation on Fitness. *Am Nat*. **71**.
Hellsten U, Wright K, Jenkins J, Shu S, Yuan Y, Wessler S, Schmutz J, Willis J, Rokhsar D.
2013. Fine-scale variation in meiotic recombination in *Mimulus* inferred from population
shotgun sequencing. *P Natl Acad Sci USA*. **110**.
Henderson J, Joyce R, Hall G, Hurst W, McGovern P. 2007. Chemical and archaeological
evidence for the earliest cacao beverages. *P Natl Acad Sci USA*. **104**.
Hinch A, Tandon A, Patterson N, Song Y, Rohland N, Palmer C, Chen G, Wang K, Buxbaum S,
Akylbekova E, et al. 2011. The landscape of recombination in African Americans.
Nature. **476**:170–5. doi:10.1038/nature10336.
Hudson RR, Kaplan NL. 1988. The coalescent process in models with selection and
recombination. *Genetics*. **120**:831–840.
Kim S, Plagnol V, Hu T, Toomajian C, Clark R, Ossowski S, Ecker J, Weigel D, Nordborg M.
2007. Recombination and linkage disequilibrium in *Arabidopsis thaliana*. *Nat Genet*.
39:1151–1155. doi:10.1038/ng2115.
Lê S, Josse J, Husson F. 2008. FactoMineR : An R Package for Multivariate Analysis. *J Stat*
Softw. **25**. doi:10.18637/jss.v025.i01.
Li W-H, Nei M. 1974. Stable linkage disequilibrium without epistasis in subdivided populations.
Theor Popul Biol. **6**:173–183. doi:10.1016/0040-5809(74)90022-7.
Luo C, Shu B, Yao Q, Wu H, Xu W, Wang S. 2016. Construction of a High-Density Genetic Map
Based on Large-Scale Marker Development in Mango Using Specific-Locus Amplified
Fragment Sequencing (SLAF-seq). *Front Plant Sci*. **7**. doi:10.3389/fpls.2016.01310.
Mackiewicz D, de Oliveira PMC, Moss de Oliveira S, Cebrat S, Lustig AJ. 2013. Distribution of
Recombination Hotspots in the Human Genome – A Comparison of Computer
Simulations with Real Data. *PLoS ONE*. **8**. doi:10.1371/journal.pone.0065272.
Mackiewicz D, Zawierta M, Waga W, Cebrat S. 2010. Genome analyses and modelling the
relationships between coding density, recombination rate and chromosome length. *J*
Theor Biol. **267**:186–192. doi:10.1016/j.jtbi.2010.08.022.
Martinez-Perez E, Schvarzstein M, Barroso C, Lightfoot J, Dernburg AF, Villeneuve AM. 2008.
Crossovers trigger a remodeling of meiotic chromosome axis composition that is linked
to two-step loss of sister chromatid cohesion. *Gene Dev*. **22**:2886–901.
doi:10.1101/gad.1694108.
McVean G. 2010. What drives recombination hotspots to repeat DNA in humans? *Philos T Roy*
Soc B. **365**:1213–1218. doi:10.1098/rstb.2009.0299.
Mcvean G, Myers S, Hunt S, Deloukas P, Bentley D, Donnelly P. 2004. The fine-scale structure
of recombination rate variation in the human genome. *Science*. **304**:581–4.
Mézard C. 2006. Meiotic recombination hotspots in plants. *Biochem Soc T*. **34**:531–4.
Motamayor J, Lachenaud P, Loor R, Kuhn D, Brown J, Schnell R. 2008. Geographic and
Genetic Population Differentiation of the Amazonian Chocolate Tree (*Theobroma cacao*
L). *PLoS One*. **3**. doi:10.1371/journal.pone.0003311.
Motamayor J, Mockaitis K, Schmutz J, Haiminen N, Livingstone D, Cornejo O, Findley S, Zheng
P, Utro F, Royaert S, et al. 2013. The genome sequence of the most widely cultivated
cacao type and its use to identify candidate genes regulating pod color. *Genome Biol*.
14:r53–r53. doi:10.1186/gb-2013-14-6-r53.
Mousavi M, Tong C, Liu F, Tao S, Wu J, Li H, Shi J. 2016. De novo SNP discovery and genetic
linkage mapping in poplar using restriction site associated DNA and whole-genome
sequencing technologies. *BMC Genomics*. **17**. doi:10.1186/s12864-016-3003-9.
Moyers B, Morrell P, McKay J. 2018. Genetic Costs of Domestication and Improvement. *J*
Hered. **109**: 103–116.
Myers S, Bottolo L, Freeman C, McVean G, Donnelly P. 2005. A fine-scale map of

recombination rates and hotspots across the human genome. *Science*. **310**:321–324.

Ohta T. 1982. Linkage disequilibrium due to random genetic drift in finite subdivided
populations. *P Natl Acad Sci USA*. **79**:1940–1944. doi:10.1073/pnas.79.6.1940.

Ollitrault P, Terol J, Chen C, Federici C, Lotfy S, Hippolyte I, Ollitrault F, Bérard A, Chauveau A,
Cuenca J, et al. 2012. A reference genetic map of *C. clementina* hort. ex Tan.; citrus
evolution inferences from comparative mapping. *BMC Genomics*. **13**. doi:10.1186/1471-
2164-13-593.

Otto S, Barton N. 1997. The evolution of recombination: removing the limits to natural selection.
Genetics. **147**:879–906.

Paape T, Zhou P, Branca A, Briskine R, Young N, Tiffin P. 2012. Fine-scale population
recombination rates, hotspots, and correlates of recombination in the *Medicago*
truncatula genome. *Genome Biol Evol*. **4**:726–737. doi:10.1093/gbe/evs046.

Pickrell J, Pritchard J. 2012. Inference of Population Splits and Mixtures from Genome-Wide
Allele Frequency Data. *PLoS Genetics*. **8**. doi: 10.1371/journal.pgen.1002967.

Ptak S, Hinds D, Koehler K, Nickel B, Patil N, Ballinger D, Przeworski M, Frazer K, Pääbo S.
2005. Fine-scale recombination patterns differ between chimpanzees and humans.
Nature Genet. **37**:429–434.

Quinlan A, Hall I. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. **26**:841–842. doi:10.1093/bioinformatics/btq033.

R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL http://www.R-project.org/.

Revelle W. 2019. psych: Procedures for Psychological, Psychometric, and Personality
Research. Northwestern University, Evanston, Illinois. R package version 1.9.12,
https://CRAN.R-project.org/package=psych.

Rizzon C, Marais G, Gouy M, Biémont C. 2002. Recombination rate and the distribution of
transposable elements in the Drosophila melanogaster genome. *Genome Res*. **12**:400–
407.

Rodgers K, McVey M. 2016. Error-Prone Repair of DNA Double-Strand Breaks. *J Cell Physiol*.
231:15–24. doi:10.1002/jcp.25053.

Ross-Ibarra J. 2004. The Evolution of Recombination under Domestication: A Test of Two
Hypotheses. *Am Nat*. **163**:105–112. doi:10.1086/380606.

Sanjuán R, Moya A, Elena SF, Ohta T. 2004. The Distribution of Fitness Effects Caused by
Single-Nucleotide Substitutions in an RNA Virus. *P Natl Acad Sci USA*. **101**:8396–8401.

Schnable P, Ware D, Fulton R, Stein J, Wei F, Pasternak S, Liang C, Zhang J, Graves L, Minx
T, et al. 2009. The B73 maize genome: complexity, diversity, and dynamics. *Science*.
326:1112–1115. doi:10.1126/science.1178534.

Shanfelter A, Archambeault S, White M, Hurst L. 2019. Divergent Fine-Scale Recombination
Landscapes between a Freshwater and Marine Population of Threespine Stickleback
Fish. *Genome Biol Evol*. **11**:1573–1585. doi:10.1093/gbe/evz090.

Shiryev S, Papadopoulos J, Schäffer A, Agarwala R. 2007. Improved BLAST searches using
longer words for protein seeding. *Bioinformatics*. **23**:2949-51.

Singhal S, Leffler E, Sannareddy K, Turner I, Venn O, Hooper D, Strand A, Li Q, Raney B,
Balakrishnan C, et al. 2015. Stable recombination hotspots in birds. *Science*. **350**:928–
932. doi:10.1126/science.aad0843.

Siol M, Bonnin I, Olivieri I, Prosperi J, Ronfort J. 2007. Effective population size associated with
self-fertilization: lessons from temporal changes in allele frequencies in the selfing
annual Medicago truncatula. *J Evolution Biol*. **20**:2349–2360. doi:10.1111/j.1420-
9101.2007.01409.x.

Smith A, Hubley R, Green P. 2016. *RepeatMasker Open-4.0* (2013-2015).
<http://www.repeatmasker.org>.

Stapley J, Feulner P, Johnston S, Santure A, Smadja C. 2017. Variation in recombination

frequency and distribution across eukaryotes: patterns and processes. Philos T Roy Soc
B. **372**. doi:10.1098/rstb.2016.0455.

Stevison L, Woerner A, Kidd J, Kelley J, Veeramah K, McManus K, Bustamante C, Hammer M,
Wall J. 2016. The Time Scale of Recombination Rate Evolution in Great Apes. *Mol Biol*
Evol. **33**:928–945. doi:10.1093/molbev/msv331.

Weir B, Cockerham C. 1984. Estimating f-statistics for the analysis of population structure.
Evolution. **38**:1358–1370. doi:10.1111/j.1558-5646.1984.tb05657.x.

Winckler W, Myers S, Richter D, Onofrio R. 2005. Comparison of Fine-Scale Recombination
Rates in Humans and Chimpanzees. *Science*. **308**:107–11.

Wloch D, Szafraniec K, Borts R, Korona R. 2001. Direct estimate of the mutation rate and the
distribution of fitness effects in the yeast Saccharomyces cerevisiae. *Genetics*. **159**:441–
452.

Wu J, Mizuno H, Hayashi-Tsugane M, Ito Y, Chiden Y, Fujisawa M, Katagiri S, Saji S, Yoshiki S,
Karasawa W, et al. 2003. Physical maps and recombination frequency of six rice
chromosomes. *Plant J*. **36**:720–730. doi:10.1046/j.1365-313X.2003.01903.x.

Yelina N, Diaz P, Lambing C, Henderson IR. 2015. Epigenetic control of meiotic recombination
in plants. *Sci China Life Sci*. **58**:223–231. doi:10.1007/s11427-015-4811-x.

Zhu Y, Yin Y, Yang K, Li J, Sang Y, Huang L, Fan S. 2015. Construction of a high-density
genetic map using specific length amplified fragment markers and identification of a
quantitative trait locus for anthracnose resistance in walnut (*Juglans regia* L.). *BMC*
Genomics. **16**:614.

 Supplementary Materials:

998 Genetic differentiation and intrinsic genomic features explain variation in recombination hotspots
999 among cocoa tree populations

1000

1001 Enrique J. Schwarzkopf, Juan C. Motamayor, Omar E. Cornejo (ocornejo@gmail.com)

1002

1003 **Tables:**

1004

Table S1. The median of the upper and lower bounds of the 95% Credibility Interval for the trace of
estimates of r (in cM/Mb) from all positions in the genome are presented for each population (i.e. Position
L95 and Position U95). The upper and lower bounds of the 95% probability interval for the median
estimate of r for each population is also presented (i.e. Genome L95 and Genome U95). The quotients of
the upper and lower bounds for each of the two intervals point to a much larger genome-wide variation in
r than per-position variation in the trace for the estimate of r.

1011

Population	Position L95	Position U95	Genome L95	Genome U95	Position Range Quotient	Genome Range Quotient
Amelonado	6.35e-05	7.67e-03	2.33e-05	31.3	120.75	1.34e+06
Contamana	1.63e-01	1.34	1.40e-04	26.4	8.22	1.88e+05
Criollo	87.5	43.1	5.35e-02	1,660	4.92	3.11e+04
Curaray	5.15e-01	2.63	2.72e-04	20.2	5.11	7.40e+04
Guianna	9.76e-01	12.6	1.81e-04	296	12.90	1.63e+06
Iquitos	5.50e-05	6.52e-03	1.58e-05	24.5	186.29	1.55e+06
Maranon	1.98e-04	7.40e-02	2.31e-05	35.2	373.35	1.52e+06
Nacional	4.80e-05	6.50e-03	2.76e-05	36.6	135.60	1.33e+06
Nanay	1.65e-04	3.06e-02	2.06e-05	35.2	185.32	1.71e+06
Purus	3.98e-03	5.24e-01	2.00e-04	63.5	131.87	3.18e+05

1012

1013 **Table S2.** Mean and median genome-wide recombination rates (r) in cM/Mb for all ten *T. cacao*
1014 populations obtained using LDhat with θ= 0.001.

1015

Population	Mean r (cM/Mb)	Median r (cM/Mb)
Amelonado	4.04	4.78e-03
Contamana	4.55	9.75e-01
Criollo	715.35	668.90
Curaray	5.85	3.12
Guianna	63.24	4.34
Iquitos	2.68	7.95e-03
Maranon	4.16	3.38e-02
Nacional	6.00	1.32e-01
Nanay	5.43	1.61e-01
Purus	11.27	1.96

1022 **Table S3.** Name of *T. cacao* gene coding for FIGL1 and FLIP and amino acid mutations for FIGL1 and
1023 FLIP orthologs.
1024

Protein	FIGL1	FLIP
Gene	*Thecc1EG017182t2*	*Thecc1EG020410t1*
Mutations	K9Q, F66L, D114Y, T155I, D189V, K195M, E215K, S235N, R270Q, A285S, S291T, V548I, G566S, A567V, M580T	L102F, S153T, K174T, Y140N, V125A, H203R, Q237K, V406L, S430Y, R446Q, M497I, L521M, D696V, L729I, M739I, A746G, K875Q, P906L

1025
1026 **Table S4.** Average hotspot size (in kb) and count for hotspots detected in each population
1027 and average for all populations.
1028

Population	Mean hotspot size (kb)	Hotspot count
Amelonado	6.9	1325
Contamana	6.1	5184
Criollo	6.1	887
Curaray	5.8	2303
Guianna	8.6	3655
Iquitos	7.0	3258
Maranon	6.8	3296
Nacional	6.9	2202
Nanay	7.6	3818
Purus	6.3	3972
Average	6.9	2989.9

1029
1030 **Table S5.** Sample size and post-filtering SNP count for all ten populations of *Theobroma cacao* for which
1031 recombination maps were generated. The proportion of the genome that is callable is also reported.

Population	Sample size (chromosome)	SNP count	Proportion callable
Amelonado	11 (22)	373,789	0.68
Contamana	9 (18)	2,097,618	0.87
Criollo	4 (8)	309,818	0.69
Curaray	5 (10)	1,106,871	0.70
Guianna	9 (18)	770,729	0.915
Iquitos	7 (14)	1,575,711	0.922
Maranon	14 (28)	1,783,226	0.955
Nacional	4 (8)	718,099	0.89
Nanay	10 (20)	830,885	0.94
Purus	6 (12)	1,184,181	0.9

1052
1053
1054 **Figures:**
1055
1056

1057
1058 **Figure S1.** Drift tree constructed using treemix (Pickrell and Pritchard, 2012) for the 10 *T. Cacao*
1059 populations. Distances between populations are based on the drift parameter.
1060 Modified from Cornejo et al. (2018)
1061

Figure 2. Distribution of log_{10} recombination rates ($log_{10}(r)$) along the genomes of the ten *T. Cacao* populations. The sample size (N) is reported for each population.

1066
1067 **Figure S3**. The left panel shows the frequency of individuals that are homozygous for the alternative
1068 allele of amino acid mutations in a *T. cacao* FLIP ortholog. Alternative allele is defined in terms of the
1069 Amelonado reference genome. The right panel shows the log_e transformed recombination rates (r). The
1070 populations are in the same order in both panels.

1071
1072

1073
1074 **Figure S4.** Example of the window layout for a 10,750 SNP chromosome. The 2,000 SNP
1075 long windows are represented by alternating horizontal and vertical lines and the overlaps
1076 between them are represented by square crosshatches. Braces above the chromosome indicate
1077 the regions from which recombination rates are extracted to generate the chromosome-wide
1078 recombination rates.
1079