1   **Beyond a single patch: local and regional processes explain biodiversity patterns in a**
2   **seagrass epifaunal metacommunity**
3

4   Keila A Stark[1]*, Patrick L Thompson[1], Jennifer Yakimishyn[2], Lynn Lee[3], Mary I O'Connor[1]
5
6
7   1. Department of Zoology
8   Biodiversity Research Centre
9   University of British Columbia
10 V6T 1Z4
11 Vancouver, BC, Canada
12
13 2. Pacific Rim National Park Reserve
14 P.O. Box 280
15 Ucluelet, BC, Canada
16
17 3. Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve, and
18 Haida Heritage Site
19 60 Second Beach Road
20 Skidegate, BC, Canada
21
22 ***Correponding author:** keilastark@zoology.ubc.ca
23

24
25 **Abstract:** 260 words
26 **Main text:** 5167 words
27 **Number of references:** 57
28 **Number of figures:** 5
29 **Number of tables:** 1
30
31

32 **Data accessibility:** Data will be made available on Dryad should the manuscript be accepted. All
33 code for analysis can be found at https://github.com/keilast/HMS-Seagrass.
34
35 **Competing interests:** We have no competing interests.

Abstract

Ecological communities are structured by dispersal, environmental niche filtering, and

biotic interactions. Metacommunity ecology provides a framework for understanding how these

processes combine to determine community composition in sites that are connected through the

dispersal of multiple potentially interacting populations. Previous approaches for describing

metacommunity dynamics from observational data have had limited success due to statistical and

theoretical limitations. We used the Hierarchical Modelling of Species Communities (HMSC)

statistical framework to assess the contributions of these metacommunity processes to faunal

community structure in an important coastal habitat, seagrass meadows, across approximately

1000 km along the coast of British Columbia. Following the HMSC framework, we parametrized

a hierarchical joint species distribution model describing the relationship between environmental

variables, spatial distance between sites, species co-occurrence patterns, and random effects on

epifaunal species distributions across thirteen eelgrass (*Zostera marina* (L.)) meadows. Species

distribution patterns suggested that sites are well-connected by dispersal, and that dispersal rates

are not so high that they homogenize regional diversity. We found evidence of niche filtering in

several species along an abiotic gradient; however, environmental conditions only explained a

small fraction of the overall variation in species distributions (22%), suggesting variable niche

width among species. Strong structuring in pairwise species co-occurrences after controlling for

environmental and spatial covariates led us to hypothesize the importance of priority effects and

trophic interactions. This work has yielded refined hypotheses about the specific local and

regional dynamics influencing seagrass epifaunal community structure; these hypotheses in turn

can inform targeted experiments, monitoring efforts, and management decisions for preserving

diversity in this important coastal ecosystem.

Key words: Hierarchical Modelling of Species Communities, priority effects, dispersal, niche

filtering, British Columbia, *Zostera marina* (L.)

Introduction

Understanding how local environmental conditions, biotic interactions, and regional

connectivity jointly structure the composition of communities is a central challenge in ecology

(Ricklefs and Schluter 2003, Vellend 2010, 2016, Leibold and Chase 2017). Determining the

relative importance of these assembly processes in particular ecosystems is a critical step in

understanding the spatial and temporal scales of stability and resilience in spatially structured

communities (Economo et al. 2011). Metacommunity theory offers a promising framework for

understanding drivers of community assembly across spatial scales; however, previous attempts

to quantitatively apply metacommunity concepts to real-world systems have had limited success

(Logue et al. 2011, Brown et al. 2017). Over the past decade and a half, the primary focus of

metacommunity studies has been seeking in natural systems signatures of one of four idealized

paradigms: the neutral model, patch-dynamics, species sorting, and mass effects (Leibold et al.

2004, Cottenie 2005). An emerging perspective holds that natural metacommunities often do not

fit neatly under these paradigms because they do not reflect all possible combinations of the

various assembly processes (Logue et al. 2011, Brown et al. 2017, Leibold and Chase 2017).

Thus, the revised goal is to determine how dispersal, environmental conditions, and biotic

interactions combine to structure species composition and relative abundance within a

metacommunity.

Many marine ecosystems that host high biodiversity occupy spatially structured habitats (e.g.

rocky reefs, coral reefs, seagrass meadows), and marine species have dispersing life histories that

connect populations in distinct habitat patches. Marine systems are characterized by strong local

species interactions (Berlow 1999, Sala and Graham 2002), and the importance of 'supply-side'

or dispersal-driven population dynamics has long been recognized (Levin and Paine 1974,

Gaines and Roughgarden 1985, Lee and Bruno 2009). These combined roles of dispersal, the

local environment, and biotic interactions suggest that marine biodiversity patterns may reflect

processes characteristic of metacommunities (Guichard et al. 2003). Despite this,

metacommunity dynamics have only been explicitly investigated in a few marine systems

(Guichard et al. 2003, Yamada et al. 2014, Valanko et al. 2015, Whippo et al. 2018).

Seagrasses, like kelps, corals, and salt marsh plants, are foundation species that support high

productivity and faunal diversity (Orth et al. 1984, Duffy et al. 2015). Seagrasses form meadows

separated from each other by deeper water, un-vegetated seafloor, or other vegetated habitats

such as kelp forests, and associated fauna disperse among meadows (Boström et al. 2006). This

scenario is consistent with the type of landscape that is governed by metacommunity dynamics

(Mouquet and Loreau 2003, Leibold et al. 2004). Yet, our understanding of the causes and

conservation implications of seagrass-associated faunal diversity patterns has not accounted for

processes operating across numerous meadows and adjacent habitats in a seascape. Local

characteristics such as habitat structure (Orth et al. 1984, Heck and Valentine 2006), primary

biomass (Cébrian and Duarte 1998, Gullström et al. 2012), or trophic interactions (Amunudrud et

al. 2015, Huang et al. 2015) affect faunal species diversity, but the importance of ecological

processes operating on a regional or inter-site scale remains unclear. Our limited understanding

of the importance of these regional processes could lead to underestimates of the amount of

habitat required to sustain biodiversity and ecosystem function (Boström et al 2006).

To date, evidence for the importance of metacommunity dyamics and regional-scale

processes in seagrass associated communities is based on two different quantitative

metacommunity approaches (Yamada et al. 2014, Whippo et al. 2018). Inferences were based on

statistical methods available at the time that assessed the contribution of environment and space,

but were not able to identify possible biotic interactions and were therefore did not account for

this critical community assembly process. Using variation partitioning on constrained ordinations

(Borcard et al. 1992, Cottenie 2005), Yamada et al. (2014) found that the relative contribution of

space and environment varied among invertebrate functional groups. However, the variation

partitioning analysis was not able to distinguish between the possible metacommunity dynamics

at play owing to the ambiguous statistical significance of the results (Yamada et al. 2014).

Simulation studies have shown that this method can over- or under-estimate spatial and

environmental components of variation, and produce inflated R^2 values depending on the

researcher's statistical choices (Gilbert and Bennett 2010, Tuomisto et al. 2012). The other

approach applied to seagrass epifaunal systems, known as elements of metacommunity structure

(EMS), seeks idealized metacommunity structures in species occurrence patterns (Leibold and

Mikkelson 2002, Presley et al. 2010, Whippo et al. 2018). While useful for visualizing regional

diversity patterns, EMS does not robustly elucidate the processes contributing to these patterns

(Brown et al. 2017).

These limitations, along with the revised goal of individually assessing the contribution of

dispersal, niche filtering, and biotic interactions to diversity patterns, invite novel methodological

approaches. A new statistical framework, Hierarchical Modelling of Species Communities

(HMSC) (Ovaskainen et al. 2017), shows promise for achieving this goal. HMSC uses Bayesian

inference to parametrize a hierarchical joint species distribution model summarizing the

influence of space, environment, phylogeny, and traits on species distributions. The framework

allows the partitioning of variation in species distributions due to the environment, spatial

distance, and random effects for all species distributions, allowing the observer to discern

patterns at the whole-community level as well as in individual species. Additionally, it specifies

a way to visualize patterns of positive or negative pairwise species co-occurences. For these

reasons, HMSC achieves what researchers aimed to achieve with the previous methods, and

much more.

Here we test the hypothesis that seagrass-associated epifaunal invertebrate communities are

jointly structured by the spatial distances between meadows (a proxy for inferring dispersal

dynamics), local environmental conditions, and biotic interactions. We follow the HMSC

framework to answer the following questions: (1) Is the metacommunity characterized by

dispersal limitation, intermediate dispersal, or high dispersal? (2) Do species distribution patterns

suggest environmental niche filtering, and if so along which environmental axes? (3) Do species

co-variance patterns suggest possible biotic interactions influencing community assembly? The

order in which these questions are posed is significant. For example, if evidence for dispersal

limitation is found, this knowledge affects how we may interpret observed species co-occurrence

patterns. We address these three questions with a biodiversity survey of invertebrates in 13

Zostera marina (L.) meadows in British Columbia, Canada. We find evidence of intermediate

dispersal rates among meadows across 1000 km, but no sign of complete regional

homogenization resulting from high dispersal. We also find that local environmental conditions

have some influence on species distributions across the 13 sites, but that individual species'

responses to environmental gradients vary dramatically among taxa. Strong species' covariance

patterns led us to hypothesize possible biotic interactions influencing local community

composition.

Methods

Study sites and field sampling of eelgrass and associated epifauna

We sampled epifaunal biodiversity in 13 eelgrass *Zostera marina* (L.) meadows across a

spatial scale of 1000 km from late June through early August in 2017. The sites were located

within four coastal areas: Haida Gwaii within Gwaii Haanas National Park Reserve, National

Marine Conservation Area Reserve and Haida Heritage Site; southern Clayoquot Sound (one site

was within Pacific Rim National Park Reserve); Barkley Sound on the West coast of Vancouver

Island; and the Southern Gulf Islands on the south-eastern tip of Vancouver Island within Gulf

Islands National Park Reserve (Fig. 1).

We sampled three sites in Barkley Sound (SA, DC, RB) with SCUBA because they

required boat access, and the deepest parts of the meadow reached a depth of 5 m. All other sites

were sampled by wading or snorkeling at a low tide of 0.6 m or less. We used a standardized,

nested sampling regime with 0.25 m x 0.25 m quadrats to collect aboveground eelgrass,

epifaunal invertebrates, detritus, and macroalgae (Whippo et al. 2018). Six quadrats were set in a

15 m x 30 m array, such that every quadrat was situated 15 metres from its nearest neighbour

(Appendix S1: Fig. 1). The array was placed in the middle of the eelgrass meadow to avoid edge

effects where possible. We removed all above-ground biomass (eelgrass, detritus, macroalgae)

within each quadrat by hand and immediately placed the contents into a mesh or ziploc bag.

Acquiring biotic and abiotic environmental variables

We acquired water quality and biotic data associated with hypothesized niche differences

among epifauna. The water quality variables included in-situ measurements of pH, salinity,

dissolved oxygen, nitrates and chlorophyll-*a* at each meadow, and averages acquired from Bio-

ORACLE (Assis et al. 2018), including surface chlorophyll-*a*, dissolved oxygen, surface nitrates,

surface silicates, surface phosphates, pH, salinity, and sea surface temperature. Bio-ORACLE is

a database of marine environmental variables gathered from several satellite and in-situ sources

at a spatial resolution of 0.08° (Assis et al. 2018). All sites (meadows) were situated in different

spatial cells, therefore all sites had distinct values for the aforementioned parameters. We also

quantified fetch, which is a proxy for site exposure.

For each quadrat sampled, we measured five biotic attributes summarizing habitat

structure and food availability: eelgrass leaf area index (LAI) and shoot density, eelgrass dry

mass, total algal dry mass, and detritus dry mass. To quantify LAI, we measured leaf length,

width, and number of blades in five haphazardly chosen shoots within each quadrat. Leaf area

index was calculated by multiplying the average blade area (m^2) per shoot by the shoot density

(number of shoots per m^2). We separated and dried eelgrass, detritus, and macroalgae in a

desiccator oven (60° C for 48 hours) to measure dry mass.

Quantifying epifaunal diversity

We rinsed the eelgrass blades from each quadrat to remove epifaunal invertebrates, and

passed samples through a 500 μm sieve. Invertebrates were transferred into a centrifuge tube and

preserved in 95% EtOH for identification with light microscopy (10 times magnification). We

identified invertebrates to the lowest taxonomic level possible using keys in the Light and Smith

Manual (Carlton 2007) and Kozloff (1996).

Analysis

We followed the methodology of Ovaskainen et al. (2017) to construct a hierarchical

joint species distribution model describing the relationship between measured environmental

variables, spatial distance, species co-occurrences and species distributions using the HMSC

package (Guillaume Blanchet et al. 2018) in R (version 3.4.2; R Development Core Team 2014).

The model is composed of two linear predictor terms: one describing fixed effects of

environment (nine water quality and five biotic variables) and space, and one describing random

effects that cannot be attributed to environment or space (Equations 2 & 3 in Ovaskainen et al.

2017). The random-effects linear predictor term describes random variation in individual species

distributions as well as covariation for every species pair in the analysis.

We used Markov chain Monte Carlo to estimate the model parameters. Species'

responses to environmental variables were assumed to follow a multivariate Gaussian

distribution. We assumed a Poisson distribution for species count data and used the same priors

as in the examples presented in Ovaskainen et al. (2017). The Markov Chain included 100 000

iterations, a burn-in length of 1000 iterations, and thinning of 100. We confirmed that parameters

were well mixed by visually inspecting trace plots. We tested the model's performance by

calculating explained deviance (D^2_{adj}) as described in Appendix S1: Eq. S1. D^2_{adj} is comparable

to R^2_{adj} in least square regression analysis (Guisan and Zimmermann 2000); in this context, it

describes the model fit where a D^2_{adj} value of 1 means the model perfectly predicts the observed

data. Our calculations yielded a D^2_{adj} value for each individual species distribution model, and

the mean D^2_{adj} across all species gave a single metacommunity-level value of the model's

predictive power.

To answer our three main questions about the roles of metacommunity processes, we

visualized the parametrized model in several ways (Table 1). We investigated three possible

dispersal conditions: dispersal limitation, intermediate dispersal, and dispersal surplus. To

investigate dispersal limitation at large spatial scales, we plotted distance-decay of community

similarity. To do so, we randomly drew 200 pairwise distances between sampling plots, and

plotted predicted community similarity against distance. Distance-decay of community similarity

could reflect dispersal limitation due to isolation by distance, or it could reflect differences in

environmental conditions with increasing spatial scale. To distinguish these possibilities, we

compared predicted species compositional similarity as a function of both distance and

environmental covariates (the full HSMC model), and predicted compositional similarity using

an alternative model in which the effect of the environment was removed (all environmental

covariates were set to their mean value across all sites; Equation 5 in Ovaskainen et al. 2017). To

distinguish between dispersal sufficiency (intermediate) and dispersal surplus (high), we visually

inspected our site-by-species matrix for signatures of regional homogenization by high dispersal.

We used the degree of environmental niche filtering (as estimated by D^2_{adj}) as another line of

evidence to distinguish intermediate and high dispersal: theory predicts that intermediate

dispersal allows species to colonize sites where local environmental conditions are optimal for

growth and reproduction, whereas high dispersal overrides this (Mouquet and Loreau 2003).

To estimate the strength of niche filtering, we conducted variation partitioning analysis of

the fixed and random effects as defined in Ovaskainen et al. (2017). This summarized the

relative contribution of all environmental variables, spatial distance, and random effects to the

distribution of each species in the model; we visualized this in a stacked bar plot. We then

estimated the pure and shared fractions of variation explained by environment and space, as well

as the unexplained fraction using type III sum of squares (Borcard et al. 1992), but using the

HMSC model as opposed to constrained ordination. To do so, we followed the steps and

equations in Appendix S1: Eq. 2.

To identify species co-occurrence patterns resulting from possible biotic interactions, we

represented modelled pairwise associations in a correlation matrix describing the extent to which

species pairs co-occurred more negatively or positively than by random chance, controlling for

the effect of the environment. We then conducted hierarchical cluster analysis on sites according

to predicted species abundances from the HMSC model using Ward's criterion (Ward 1963). We

mapped the cluster dendrogram onto an abundance heatmap of the 20 species in our study with

the highest mean abundance, and colour coded the cells according to the species co-occurrence

groupings (groups of species that tended to positively co-occur amongst themselves). Full

versions of these figures with all species are provided in Appendix S1: Figs. S4 and S5.

Results

Species abundance and distribution patterns

We counted and identified 39 644 individuals representing at least 58 species across the

region (Appendix 1: Table S1). The represented taxa span multiple phyla, diet types (herbivores,

detritivores, suspension feeders, and carnivores), and life history strategies (brooding, broadcast

spawning). The mean plot or quadrat-level species richness was approximately 13 species, and

mean meadow or site-level richness was approximately 16 species. A full summary of traditional

diversity metrics can be found in Appendix S1: Table S2.

Several taxa were present at all sites (Appendix S1: Fig. S2). These include harpacticoid

copepod *Porcellidium* sp., bivalves *Saxidomus gigantea* and *Clinocardium nutallii*, snails

Alvania compacta, *Lacuna* spp., and the tanaid *Leptochelia* sp. Genera *Mytilus* (a mussel) and

Nereis (a polychaete worm) were also represented at all sites, however we were unable to

confidently identify these organisms to species. Ten species were present in at least one site in

every region: pycnogonid *Anoplodactylus viridintestinalis,* isopod *Idotea resecata,* gammarid

amphipods *Monocorophium insidiosum, Ampithoe valida, Pontogeneia rostrata, Ampithoe dalli,*

Aoroides spp*.,* snails *Amphissa columbiana* and *Alia carinata,* and limpet *Lottia pelta.*

Dispersal: Distance-decay of community similarity due to spatial distance and environment

280   In the full HMSC model summarizing species' responses to environment and distance

between sites, predicted community similarity decreased with increasing distance between

meadows (Fig. 2, red points and line). However, community similarity showed a slight increase

with distance when the effect of environmental variables was removed from the model (all

environmental variables set to their mean value), suggesting that spatial distance alone did not

correspond with differences in invertebrate compositional similarity (Fig. 2, blue points and

line).

Environmental niche filtering: variation partitioning

289   Patterns in abundance and distribution were consistent with weak environmental niche

filtering along an abiotic axis; however, the importance of environmental conditions varied

across species. The relative contributions of all 14 environmental variables, spatial distance and

random effects in the HMSC model can be seen in Fig. 3. Out of all the measured environmental

variables, the water quality variables (all blue bars in Fig. 3) explained the largest proportion of

variation across nearly all species distributions. Across all taxa, herbivore food availability

(seagrass, algae and detritus biomass) and habitat structure (eelgrass shoot density and LAI) had

smaller influences on species distributions than abiotic predictors (Fig. 3). The relative

contribution of abiotic parameters to species abundance and distribution patterns also varied

among species and major taxonomic groups. While the mean deviance explained (D^2_{adj}) across

all species distribution models was 0.29, individual values varied dramatically (0 to 0.82,

Appendix 1: Table S1), implying high variation in the match between our model and the

observed species distributions. Accounting for this, the true fraction of variation explained by

environment was 22.4%, space explained -3%, the shared fraction between environment and

space was 4%, and the residual fraction of variation was 0.76 (Appendix S1: Fig. S3).

Species interactions: Patterns in pairwise species co-occurrences

We identified, post hoc, three main site-level species co-occurrence groupings after

controlling for effects of environmental covariates. The first group of species indicated by purple

cells (*Nereis* sp. through *Caprella laeviuscula* along the vertical axis in Fig. 4a), henceforth

known as the *Nereis* group, tended to positively co-occur with each other, and negatively co-

occur with the second group. This second group of species indicated by blue cells, henceforth

known as the *C. californica* group (*Caprella californica* through *Ampithoe valida,* and

Harpacticoid 2 through *Lacuna* sp. along the vertical axis of Fig. 4a), positively co-occurred with

members of the same group and negatively co-occurred with members of the *Nereis* group.

Alvania compacta and *Lacuna* sp. negatively co-occurred with members of the *C. californica*

group and positively co-occurred with members of the *Nereis* group, however their co-

occurrence values tended to be weaker. The third co-occurrence grouping was comprised of

species that did not show strong positive or negative co-occurrence with other species. *Spirorbis*

sp. was the only species in this category among the 20 species with highest mean abundances.

Species representing each of the three major co-occurrence groupings were present at all

sites (Fig. 4b), indicating that members from these groups can be simultaneously present.

However, the abundances of members from the different co-occurrence groups tended to

negatively co-vary; when species from one group were abundant, species from the others were

less abundant (Fig. 4b). At sites JB, HL, CB, RA, EB, DC, and DK, members of the *Nereis*

group were predicted to be most abundant, whereas *C. californica* group members were

predicted to be most abundant at SS, SA, LH, GB and RB. The site IN was predicted to be

strongly dominated by *Spirorbis* sp. in. The emergent species assemblages identified in Fig. 4a

from the HMSC modeled distributions do not clearly correspond with the spatial distance

because the cluster analysis did not group sites by spatial proximity or regional membership

(black symbols in Fig. 4b).

Discussion

By applying a new statistical framework, we have gained new insight into the possible

metacommunity processes structuring seagrass-associated biodiversity in coastal British

Columbia. We report evidence that both biotic interactions and environmental conditions

influence seagrass meadow invertebrate composition. Spatial patterns of species abundance and

distribution suggest regional-scale dispersal patterns are possible, such that even distant

meadows may host the same species. However, these dispersal rates are clearly not so high as to

overwhelm the local meadow scale effects of biotic interactions and environmental conditions.

Below, we review these conclusions and the evidence to support them.

Epifaunal communities may be well-connected and experience intermediate dispersal rates

Seagrass epifaunal communities appear to be well-connected by dispersal at the present

spatial scale and distribution of seagrass habitat along the British Columbia coast. This is true for

the seven species that were present at all sites, and the additional ten species that were not

present at every site but present in all regions (Appendix S1: Fig. S2). It is noteworthy that these

widely distributed groups had representation from across phyla and life history traits, suggesting

that no single taxonomic group or dispersal life history strategy had a consistently greater spatial

distribution among the sites sampled. Many of the meadows sampled were small (< 1 ha), and

epifaunal abundances can vary substantially within meadows from year to year such that local

extinctions followed by rescue from nearby populations are likely important for diversity and

composition (Thom et al. 1995). However, the specific pathway the organisms travelled to arrive

at these sites remains unknown. We used Euclidean distances in our analysis, which preserved

the rank order of distances across sites, but actual distances are likely greater due to

oceanographic circulation patterns (Kinlan and Gaines 2003, Mitarai et al. 2008, Treml et al.

2015).

It is unclear whether distant sites share the same species because they are (1) linked by

direct dispersal; (2) indirectly linked by dispersal via unsampled "stepping-stone" sites; or (3)

were colonized by populations of the same species in a historical dispersal event, but have not

seen the exchange of individuals since. Seagrass-associated epifauna generally disperse passively

either by larval transport in currents or by "rafting" on floating pieces of seagrass or macroalgae.

Kelp rafts have been observed hosting benthic and epifaunal invertebrates across phyla

(echinoderms; peracarids; molluscs; annelids), and are important in maintaining connectivity

between coastal ecosystems across 100s of kilometres (Wichmann et al. 2012). Thus, while

incidences of direct dispersal between the farthest sites may be rare, they are not impossible.

Species in our analysis that were only found at one site or region (e.g. *Jassa marmorata,*

Neoamphitrite robusta, Appendix S1: Fig. S2) may be dispersal limited for reasons other than

prohibitively long travel distances. Those with low abundances (fewer than five individuals per

meadow) may not disperse in appreciable numbers, and therefore cannot establish populations at

all sites as readily as more abundant species. This is supported by our observation of a generally

positive relationship between range size (number of sites a species was represented at) and

average abundance (Appendix 1: Fig. S6).

We can also conclude that dispersal rates are high enough for several species to be

present at most sites, but not so high as to completely overwhelm the effects of the environment

and biotic interactions. Thus, dispersal rates are likely intermediate on average. A subset of

species was present at all sites, but often at varying abundances (Fig 4b); this may be indicative

of weak mass effects, wherein immigration allows populations to persist even if local conditions

are not optimal (Schmida and Wilson 1985, Mouquet and Loreau 2003). Metacommunity theory

predicts that such weak mass effects are likely to be found whenever dispersal rates are not

limiting (Thompson et al. 2017), particularly in organisms that cannot control their own dispersal

(Leibold and Chase 2017) such as seagrass-associated epifauna. This may be supported by our

result on the contribution of environmental conditions, particularly in ubiquitous species for

which the model fit was poor (e.g. *Aoroides* spp., *Spirorbis* sp., $D^2_{adj} = 0$); the poor model fit

may indicate that these persist at sites even if environmental conditions are not optimal, due to

sufficient immigration. An alternative explanation for the model's poor predictive power in these

species is that these taxa have broader environmental niches, and thus the regional environmental

gradient is not sufficient to drive clear presence-absence patterns.

Seagrass epifauna show varying degrees of environmental niche filtering

The most important environmental axis informing species distributions is the abiotic one,

with phosphates, salinity, and dissolved oxygen explaining the largest fractions of variation (Fig.

3). However, after accounting for the model fit, the total variation explained by the environment

was small (22%). In some species for which the model had high predictive power (e.g. *Ampithoe*

dalli and *Caprella natalensis,* Appendix 1: Table S1*)*, niche filtering may be important, but

overall there remains a large amount of variation unexplained by the environment.

Environmental heterogeneity may influence species distributions in one of three ways.

First, environmental conditions may influence food availability through primary productivity.

This is possible despite the fact that primary biomass explained less variance than the abiotic

parameters in our HMSC model (Figure 3); the primary biomass measurements were taken at a

single time point. Conversely, the abiotic parameters represent a four-year average (Assis et al.

2018) and may therefore better represent the growing conditions these organisms experience

over their lifetimes. Second, sea surface temperature, dissolved oxygen, and salinity may affect

species' distributions by via their environmental tolerance ranges. For example, estuarine isopod

Idotea baltica (a relative of *I. resecata* and *I. montereyensis* in our study) experiences

significantly lower survival in prolonged exposure to higher water temperatures and lower

salinities than those experienced in their home habitat (Rugiu et al. 2018). Third, temperature

may influence the metabolic demands of the epifauna (many of which are herbivores) thereby

influencing grazing rates and altering the available primary biomass (O'Connor 2009). Decreased

food resources may increase competition, thereby influencing the number and identity of species

a seagrass patch can host. Determining whether individual species' responses to competitive

stress influence whole-community structure may provide key information on how environment

and species interactions interact to influence community structure.

Environmental conditions in temperate eelgrass meadows fluctuate seasonally, driving

changes in the relative abundance of epifaunal taxa throughout the year (Wlodarska-Kowalczuk

et al. 2014, Markel et al. 2017). As a result, the epifaunal metacommunity may not show these

distributions year-round; sampling several times in one year to capture these temporal changes

would give a clearer picture of species' responses to all metacommunity processes.

Species co-variance patterns may suggest priority effects and trophic interactions

Three main co-occurrence groupings emerged showing antagonistic abundances across

sites (Fig. 4b). Stochasticity alone is unlikely to have produced such a strong pattern, especially

given deterministic processes tend to be more important in systems that are not dispersal-limited

(Lee and Bruno 2009, Ron et al. 2018). While there are multiple possible explanations for non-

random species occurrence patterns (Connor and Simberloff 1979), environmental heterogeneity

was ruled out as the co-occurrence values represented pairwise covariation after controlling for

spatial and environmental covariates (Ovaskainen et al. 2017). This, along with the high

unexplained fraction in our variation partitioning result (Appendix S1: Fig. 3), leads us to

suggest the potential importance of biotic processes. Several types of interactions may be

occurring in the metacommunity; many species in the study are herbivores or detritivores, and

thus may compete for primary biomass. Laboratory experiments have demonstrated that

behaviours such as grazing rate and habitat selection in amphipods are altered in the presence of

morphologically and functionally similar intrrerspecific competitors (Beermann et al. 2018).

Other species in our study are filter feeders, and thus may not be competing for food resources

but possibly predator-free space or substrate on the eelgrass blades.

The community structure of a given site with certain local environmental conditions and

access to a certain regional pool may have multiple possible compositional states depending on

the arrival order of species; this phenomenon is known as priority effects (Fukami et al. 2016, Ke

and Letten 2018) and has been documented in marine fouling communities (Vieira et al. 2018).

The antagonistic co-occurrence groupings in our analysis may suggest priority effects. We

hypothesize that following the establishment of a new meadow or a disturbance event, the first

few species to colonize the meadow determine the success of subsequent colonizers, regardless

of local environmental conditions. As an illustrative example, *Lacuna* spp. might have been

among the first to colonize sites JB, HL, GB, and CB (Fig. 4b), thereby gaining a numerical

advantage over species in the alternative cohort. Distibution patterns in the sessile calcifying

polychaete *Spirorbis* sp. may also indicate priority effects. While it did not strongly negatively or

positively co-occur with other species, it dominated the HL site; 16 300 individuals were found

across the six quadrats, several orders of magnitude higher than its abundance at other sites (Data

S1). Site HL also had the lowest species richness of all 13 sites (9 species). Eelgrass shoots there

were completely covered by these *Spirorbis* sp., possibly preventing grazers in the community

from accessing epiphytic algae or the seagrass itself. The age, colonization history, and

disturbance regime of the meadows likely matter if priority effects are important.

The negatively correlated abundance patterns could also reflect the presence of a predator

such as *Nereis* at some sites and not others (Fig. 4b). Field experiments have demonstrated that

changes in predation pressure by fish, shorebirds, and predaceous invertebrates can shift

seagrass-associated epifaunal assemblages in a matter of weeks (Amundrud et al. 2015; Huang et

al. 2015; Lewis & Anderson 2016). Differential predation pressure could also lead to the pattern

of diverging community composition that we have witnessed (Guzman et al. 2018). A more

comprehensive sampling effort of fish and bird predators would be required to confirm the

degree to which trophic interactions structure these communities.

The positive co-occurrence patterns between certain species may additionally be a result

of positive biotic interactions. An example is seen in *Orchomenella recondita* - a gammarid

amphipod that lives in the gastrovascular cavity of the anemone *Anthopleura elegans* (Carlton

2007). This species was only recorded at the SS site, and specifically only found in quadrats

where *A. elegans* was collected with the seagrass shoots.

Conclusion

Using a new statistical approach, we have arrived at refined hypotheses about the local

and regional metacommunity processes driving temperate seagrass epifaunal diversity. Our study

also gives a baseline spatial scale (1000 km) for comparing the contribution of these processes

across spatial scales and other ecological systems (Viana and Chase 2018). Manipulative

experiments and monitoring are now needed to test our refined hypotheses. One could test the

hypothesis that species sorting is relatively weak by measuring fitness differences along the most

important environmental axes. Given our hypothesis on the importance of priority effects, one

might experimentally manipulate the colonization order of species from the different co-

occurrence groupings to determine if resulting compositional states differ (Fukami et al. 2016).

Testing our hypothesis that populations are not dispersal limited is more challenging, but could

be done using particle tracking models similar to those used to estimate connectivity between

coral reefs via larval dispersal across thousands of kilometres (Treml et al. 2008). A population

genetic analysis could also ground-truth the extent and time scale of dispersal between allopatric

populations. All together, we suggest that a modern metacommunity perspective combined with

new advances in our statistical toolbox provide a powerful and relevant framework for studying

natural ecosystems.

Our findings are particularly relevant to conservation initiatives that aim to maintain

diversity in seagrass meadows; identifying the contribution of local and regional processes

allows the conservation practitioner to better predict how habitat fragmentation and climate

change might influence diversity. Seagrass-associated epifaunal communities may be quite

resilient, given our findings that environmental conditions make a minimal contribution to

species distributions, and several species across phyla and functional groups are consistently

present across a fairly large spatial scale.

Acknowledgements

KAS conceived of the study with guidance from MIO. JY provided samples from Clayoquot

Sound. LL provided samples from Haida Gwaii. KAS collected and processed samples from the

Gulf Islands and Barkley Sound, analyzed data, and wrote the first manuscript draft. PLT

provided recommendations for the analysis. KAS, PLT, and MIO contributed to writing the

manuscript. This research was financially supported by a UBC Zoology SURE Grant, and

NSERC Discovery Grants, Canadian Foundation for Innovation (CFI), and is sponsored by the

NSERC Canadian Healthy Oceans Network and its Partners: Department of Fisheries and

Oceans Canada and INREST (representing the Port of Sept-Îles and City of Sept-Îles). PLT is

supported by NSERC and Killam postdoctoral fellowships. We are grateful to Duarte Viana for

providing critical recommendations and code for the D^2_{adj} and variation partitioning calculations.

Matt Whalen pulled the Bio-ORACLE dataset, and John Cristiani made the map in Fig. 1.

Special thanks to Michelle Paleczny and Gulf Islands National Park Reserve for boat and

laboratory space during their eelgrass field season. We acknowledge staff, students, and

volunteers from Pacific Rim National Park Reserve for their assistance collecting eelgrass

samples in Clayoquot Sound. Thanks goes to Rob Underhill, Abbie Sherwood, Emily

Adamczyk, Minako Ito, Ariane Comeau, Giordano Bua for assistance with sample collection,

processing, and ID. We thank Coreen Forbes, Matt Whalen and Jeff Groh for their comments on

the manuscript drafts.

510   Literature Cited

Amundrud, S.L., D. S. Srivastava, and M. I. O'Connor. 2015. Indirect effects of predators

control herbivore richness and abundance in a benthic eelgrass (*Zostera marina*)

mesograzer community. Journal of Animal Ecology 84:1092–1102.

Assis J., L. Tyberghein L. S. Bosch, H. Verbruggen, E. A. Serrão, and O. De Clerck. 2018. Bio-

ORACLE v2.0: Extending marine data layers for bioclimatic modelling. Global Ecology

and Biogeography 27:277–284.

Beermann, J., K. Boos, L. Gutow, M. Boersma, and A. C. Peralta. 2018. Combined effects of

predator cues and competition define habitat choice and food consumption of amphipod

mesograzers. Oecologia 186:645–654.

Berlow, E. L. 1999. Strong effects of weak interactions in ecological communities. Nature 398:

330–334.

Boström, C., E. L. Jackson, and C. A. Simenstad. 2006. Seagrass landscapes and their effects on

associated fauna: A review. Estuarine, Coastal and Shelf Science 68:383-403.

Borcard, D., P. Legendre, and P. Drapeau. 1992. Partialling out the Spatial Component of

Ecological Variation. Ecology. 73:1045–1055.

Brown, B.L., E. R. Sokol, J. Skelton, and B. Tornwall. 2017. Making sense of metacommunities:

dispelling the mythology of a metacommunity typology. Oecologia 183:643–652.

Carlton, J. 2007. The Light and Smith Manual: Intertidal invertebrates from Southern California

to Oregon. Fourth edition. University of California Press, Oakland, California, USA.

Cébrian, J. and C. M. Duarte. 1998. Patterns in leaf herbivory on seagrasses. Aquatic Botany

60:67–82.

Connor, E. F., and D. Simberloff. 1979. The assembly of species communities: Chance or

competition? Ecology 60:1132–1140.

Cottenie, K. 2005. Integrating environmental and spatial processes in ecological community

dynamics. Ecology Letters 8:1175–1182.

Duffy J.E, et al. 2015. Biodiversity mediates top-down control in eelgrass ecosystems: A global

comparative-experimental approach. Ecology Letters 18:696–705.

Economo, E. P. 2011. Biodiversity Conservation in Metacommunity Networks: Linking Pattern

and Persistence. The American Naturalist 177:167-180.

Fukami, T., E. A. Mordecai, and A. Ostling. 2016. A framework for priority effects. Journal of

Vegetation Science 27:655–657.

Gaines, S., and J. Roughgarden. 1985. Larval settlement rate: A leading determinant of structure

in an ecological community of the marine intertidal zone. Proceedings of the National

Academy of Sciences. 82:3707–3711.

Gilbert, B, and J. R. Bennett. 2010. Partitioning variation in ecological communities: do the

numbers add up? Journal of Applied Ecology 47: 1071–1082.

Guichard, F. 2003. Mussel disturbance dynamics: Signatures of oceanographic forcing from

local interactions. American Naturalist 161:889–904.

Guillaume Blanchet, F., with contributions from G.Tikhonov and A. Norberg. 2018. HMSC:

Hierarchical Modelling of Species Community. R package version. 2.1-2.

Guisan, A., and N. E. Zimmermann. 2000. Predictive habitat distribution models in ecology.

Ecological Modelling 145:147–186.

Gullström, M., S. Baden, and M. Lindegarth. 2012. Spatial patterns and environmental correlates

in leaf-associated epifaunal assemblages of temperate seagrass (*Zostera marina*) meadows.

Marine Biology 159:413–425.

Guzman, L. M., R. M. Germain, C. Forbes, S. Straus, M. I. O'Connor, D. Gravel, D. S. Srivastava,

and P. L. Thompson. 2018. Towards a multi-trophic extension of metacommunity ecology.

In press. Ecology Letters doi: 10.1111/ele.13162.

Heck K.L., and J. F. Valentine. 2006. Plant-herbivore interactions in seagrass meadows. Journal

of Experimental Marine Biology 330:420–436.

Huang A.C., M. Essak, and M. I. O'Connor. 2015. Top-down control by great blue herons *Ardea*

herodias regulates seagrass-associated epifauna. Oikos 124:1492–1501.

Ke, P. J., and A. D. Letten. 2018. Coexistence theory and the frequency-dependence of priority

effects. Nature Ecology and Evolution 2:1691–1695.

Kinlan, B.P., and S. D. Gaines. 2003. Propagule Dispersal in Marine and Terrestrial

Environments: A Community Perspective. Ecology 84:2007–2020.

Kozloff, E. 1996. Marine Invertebrates of the Pacific Northwest. University of Washington

Press, Seattle, Washington, USA.

Leibold, M.A., et al. 2004. The metacommunity concept: A framework for multi-scale

community ecology. Ecology Letters 7:601–613.

Leibold, M. A. and J. M. Chase. 2017. Metacommunity Ecology, Volume 59. Princeton

University Press, Princeton, New Jersey, USA.

Leibold, M. A., and G. M. Mikkelson. 2002. Species Turnover , and Boundary Clumping:

Elements of Meta-Community Structure. Oikos 97:237–250.

Lee, S. C, and J. F. Bruno. 2009. Propagule supply controls grazer community structure and

primary production in a benthic marine ecosystem. Proceedings of the National Academy

of Sciences 106:7052–7057.

Levin, S. A., and R. T. Paine. 1974. Disturbance, patch formation, and community

structure. Proceedings of the National Academy of Sciences. 71:2744–2747.

Lewis, L. S., and T. W. Anderson. 2016. Top-down control of epifauna by fishes enhances

seagrass production. Ecology 93:2746–2757.

Markel, R. W., K. E. Lotterhos, and C. K. Robinson. 2017. Temporal variability in the

environmental and geographic predictors of spatial-recruitment in nearshore rockfishes.

Marine Ecology Progress Series 574:97–111.

Mitarai, S., D. A. Siegel, and K. B. Winters. 2008. A numerical study of stochastic larval

settlement in the California Current system. Journal of Marine Systems 69:295–309.

Mouquet, N., and M. Loreau. 2003. Community patterns in source-sink metacommunities.

American Naturalist 162:554–557.

O'Connor, M.I. 2009. Warming strengthens an herbivore-plant interaction. Ecology 90:388–398.

Orth, R. J., K. L. Heck, and J. van Montfrans. 1984. Faunal Communities in Seagrass Beds: A

Review of the Influence of Plant Structure and Prey Characteristics on Predator-Prey

Relationships. Estuaries 7:339-350.

Ovaskainen, O., G. Tikhonov, A. Norberg, F. Guillaume Blanchet, L. Duan, D. Dunson, T.

Roslin, and N. Abrego. 2017. How to make more out of community data? A conceptual

framework and its implementation as models and software. Ecology Letters 20:561–576.

Presley, S.J., C. L. Higgins, and M. R. Willig. 2010. A comprehensive framework for the

evaluation of metacommunity structure. Global Ecology and Evolution 19:908–917.

R Development Core Team. 2014. R: a language and environment for statistical computing. R

Foundation for Statistical Computing. Vienna, Austria. http://www.R-project.org.

Ricklefs, R. E., and D. Schluter. 1993. Species Diversity in Ecological Communities: Historical

and Geographical Perspectives. University of Chicago Press, Chicago, Illinois, USA.

Ron, R., O. Fragman-Sapir, and R. Kadmon. 2018. Dispersal increases ecological selection by

increasing effective community size. Proceedings of the National Academy of Sciences

115:11280–11285.

Rugiu, L., I. Manninen, J. Sjöroos, and V. Jormalainen. 2018. Variations in tolerance to climate

change in a key littoral herbivore. Marine Biology 165:1–11.

Sala, E., and M. H. Graham. 2002. Community-wide distribution of predator-prey interaction

strength in kelp forests. Proceedings of the National Academy of Sciences 99:3678–3683.

Schmida, A., and M.V. Wilson. 1985. Biological determinants of species diversity. Journal of

Biogeography 12:1–20.

Thom, R., B. Miller, and M. Kennedy. 1995. Temporal patterns of grazers and vegetation in a

temperate seagrass system. Aquatic Botany 50:201–205.

Thompson, P. L., B. Rayfield, and A. Gonzalez. 2017. Loss of habitat and connectivity erodes

species diversity, ecosystem functioning, and stability in metacommunity networks.

Ecography 40:98–108.

Treml, E.A., P. N. Halpin, D. L. Urban, and L.F. Pratson. 2008. Modeling population

connectivity by ocean currents: a graph-theoretic approach for marine conservation.

Landscape Ecology 23:19–36.

Treml, E. A., J. R. Ford, K. P. Black, and S. E. Swearer. 2015. Identifying the key biophysical

drivers, connectivity outcomes, and metapopulation consequences of larval dispersal in the

sea. Movement Ecology 3:17.

Tuomisto, H., L. Ruokolainen, and K. Ruokolainen. 2012. Modelling niche and neutral

dynamics: on the ecological interpretation of variation partitioning. Ecography 35:961-971.

Valanko, S., J. Heino, J. M. Westerbom, M. Viitasalo, and A. Norkko. 2015. Complex

metacommunity structure for benthic invertebrates in a low-diversity coastal system.

Ecology and Evolution 5:5203–5215.

Viana, D. S., and J. M. Chase. 2018. Spatial scale modulates the inference of metacommunity

assembly processes. In press. Ecology doi: 10.1002/ecy.2576

Vieira, E. A., A. A. V. Flores, and G. M. Dias. 2018. Persistence and space preemption explain

species-specifc founder effects on the organization of marine sessile communities. Ecology

and Evolution 8:3430-3442.

Vellend, M. 2010. Conceptual Synthesis in Community Ecology. The Quarterly Review of

Biology, 85:183–206.

Vellend, M. 2016. The Theory of Ecological Communities: Princeton University Press.

Ward, J. H. 1963. Hierarchical Grouping to Optimize an Objective Function. Journal of the

American Statistical Association 58:236-244.

Wichmann, C. S., I. A. Hinojosa, and M. Thiel. 2012. Floating kelps in Patagonian Fjords: An

important vehicle for rafting invertebrates and its relevance for biogeography. Marine

Biology 159:2035–2049.

Whippo, R., N. S. Knight, C. Prentice, J. Cristiani, M. R. Siegle, and M. I. O'Connor. 2018.

Epifaunal diversity patterns within and among seagrass meadows suggest landscape-scale

biodiversity processes. Ecosphere 9:02490.10.1002/ecs2.2490.

Wlodarska-Kowalczuk, M., E. Jankowska, L. Kotwicki, and P. Balazy. 2014. Evidence of

Season-Dependency in Vegetation Effects on Macrofauna in Temperate Seagrass

Meadows (Baltic Sea). PLoS One 9:e100788.

Yamada, K., Y. Tanaka, T. Era, and M. Nakaoka. 2014. Environmental and spatial controls of

macroinvertebrate functional assemblages in seagrass ecosystems along the Pacific coast of

northern Japan. Global Ecology and Conservation 2:47–61.

666   **Table 1.** Summary of evidence used to answer questions about metacommunity processes

Question	Analyses	Figure number
(1) Is the metacommunity characterized by dispersal limitation, intermediate dispersal, or high dispersal?	Dispersal limitation: Distance-decay of community similarity from HMSC joint species distribution models with and without accounting for environmental similarity	Fig. 2
	Intermediate versus high dispersal: Biodiversity patterns across sites for signatures of high dispersal (compositional homogenization); HMSC model results indicating structured co-occurrence patterns and variation in composition explained by environmental	Fig. 4a, 4b
(2) Do species distribution patterns suggest environmental niche filtering, and if so along which environmental axis?	Variation partitioning of modelled site and region-level fixed and random effects of environmental covariates and calculating predictive power (proportion of deviance) of the fixed effects model	Fig. 3
(3) Do species co-variance patterns suggest possible biotic interactions influencing community assembly?	Correlation matrix of pairwise species co-occurences, after controlling for the effect of space and environment	Fig. 4a & 4b

670   **Figure 1.** Map of coastal British Columbia showing eelgrass field sites in Haida Gwaii (HL,

671   RA), Clayoquot Sound (DK, EB, IN), Barkley Sound (SA, DC, RB), and the Southern Gulf

672   Islands (GB, JB, CB, LH, SS).

685 **Figure 2**. HMSC model predictions of community similarity relative to the spatial distance

686 (Euclidean) between communities. The red circles show similarity predictions based on the full

687 model. The blue triangles show predictions wherein the effect of all environmental parameters (9

688 water quality and 5 biotic) has been removed (standardized to their mean values), thus showing

689 predicted community similarity as a result of spatial distance only.

Figure 3. Variance partitioning of fixed (environmental variables, spatial distance) and random

effects across all species (numbers correspond to species names in Appendix 1: Table S1) in the

HMSC model. Purple cells represent site and region-level random effects. Blue cells represent

abiotic water quality parameters. Yellow cells represent food availability (sum of variances

explained by eelgrass, algae and detritus biomass). Green cells represent habitat structure, which

is the sum of variances explained by eelgrass leaf area index and shoot density. Bars are

organized according to broad taxonomic groupings as indicated in Appendix 1: Table S1. The

percentages shown next to the legend labels indicate mean variance explained by that

environmental variable across all species. Adjusted deviance explained (D^2_{adj}) for the individual

species distribution models varied (Appendix 1: Table S1). The mean D^2_{adj} across all species was

0.29.

Figure 4a. A correlation matrix showing modelled site-level co-occurrence of species pairs in

the 20 most abundant species. Purple cells represent positively co-occurring species pairs, and

turquoise cells represent negatively co-occurring species pairs. Species names along both axes

are ordered according to the output of hierarchical clustering with Ward's criterion on pairwise

co-occurrence values. A version of this figure with all 58 species is shown in Appendix S1: Fig.

S4.

Figure 4b. Heatmap and cluster dendrogram depicting modelled (HMSC) species abundance and

compositional similarity across sites. The heat map shows the twenty species with the highest

predicted mean proportional abundance at the site level, and is ordered from highest to lowest

predicted mean proportional abundance. Cell colours correspond to the HMSC co-occurrence

groups shown in Fig. 4a: *Nereis* (turquoise group), *C. californica* (purple) group, and *Spirorbis*

(grey) group. Cell shade strength represents proportional abundance at a given site (darker means

higher abundance). The symbols to the left of the site abbreviations indicate region membership;

circles show Haida Gwaii sites, squares show Barkley Sound sites, diamonds show Clayoquot

Sound sites, and triangles show Southern Gulf Islands sites.

 Appendix S1: Supporting Information

729 **Table S1.** Species names, taxonomic groupings, and proportion of deviance (predictive power).

730 Numbers correspond to the axes in Figure 3. The top twenty species with highest predicted

731 abundances according to the HMSC model are bolded.

Number	Species name	Broad taxonomic group	Adjusted deviance explained for the HMSC model (D^2_{adj})
1	*Alia carinata*	Gastropod (snail)	0.15
2	***Alvania compacta***	**Gastropod (snail)**	**0.4**
3	*Amphissa columbiana*	Gastropod (snail)	0
4	*Crepidula* sp.	Gastropod (snail)	0.64
5	*Euspira lewisii*	Gastropod (snail)	0.46
6	*Harmothoe imbricata*	Gastropod (snail)	0
7	***Lacuna*** **spp. (*L. variegata* and *L. vincta*)**	**Gastropod (snail)**	**0.45**
8	*Lirularia parcipincta*	Gastropod (snail)	0.46
9	*Lirularia* sp.	Gastropod (snail)	0
10	*Littorina* sp.	Gastropod (snail)	0.5
11	*Margarites pupillus*	Gastropod (snail)	0.45
12	*Lottia pelta*	Gastropod (limpet)	0.53
13	***Clinocardium nuttallii***	**Bivalve**	**0**
14	***Mytilus*** **sp.**	**Bivalve**	**0.28**
15	*Saxidomus gigantea*	Bivalve	0
16	*Unknown clam 1*	Bivalve	0.68
17	*Unknown clam 2*	Bivalve	0.35
18	*Dorvillea longicornis*	Polychaete	0.52
19	*Exogone sp.*	Polychaete	0.25
20	*Glycinde armigera*	Polychaete	0.05
21	*Lepidonotus squamatus*	Polychaete	0.55
22	*Neoamphitrite robusta*	Polychaete	0.22
23	*Unknown polychaete (Family Oenoidae)*	Polychaete	0.15
24	*Unknown polychaete (Family Opheliidae)*	Polychaete	0.36
25	***Spirorbis*** **sp.**	**Polychaete**	**0**

26	*Nereis* **sp.**	**Polychaete**	**0.55**
27	***Ampithoe dalli***	**Gammarid amphipod**	**0.47**
28	*Ampithoe lacertosa*	Gammarid amphipod	0.34
29	***Ampithoe valida***	**Gammarid amphipod**	**0**
30	***Aoroides* spp.**	**Gammarid amphipod**	**0**
31	*Ceradocus spinicauda*	Gammarid amphipod	0
32	*Grandidierella japonica*	Gammarid amphipod	0
33	*Ischyocerus anguipes*	Gammarid amphipod	0.33
34	*Jassa marmorata*	Gammarid amphipod	0
35	***Monocorophium insidiosum***	**Gammarid amphipod**	**0.23**
36	*Orchomenella recondita*	Gammarid amphipod	0.73
37	***Photis brevipes***	**Gammarid amphipod**	**0**
38	***Pontogeneia rostrata***	**Gammarid amphipod**	**0.65**
39	*Unknown gammarid (Family Hyalidae)*	Gammarid amphipod	0.12
40	*Unknown gammarid (Family Isaeidae)*	Gammarid amphipod	0.10
41	*Unknown gammarid (Family Ischyoceridae)*	Gammarid amphipod	0.24
42	***Caprella californica***	**Caprellid amphipod**	**0.36**
43	***Caprella laeviuscula***	**Caprellid amphipod**	**0.06**
44	***Caprella natalensis***	**Caprellid amphipod**	**0.82**
45	*Cumella vulgaris*	Cumacean	0.36
46	***Leptochelia* sp.**	**Tanaid**	**0.49**
47	*Nebalia gerkinae*	Leptostracan	0.39
48	***Porcellidium* sp.**	**Copepod**	**0.33**
49	***Harpacticoid 1***	**Copepod**	**0**
50	***Harpacticoid 2***	**Copepod**	**0**
51	*Harpacticoid 3*	Copepod	0.50
52	*Anoplodactylus viridintestinalis*	Pycnogonid	0.01
53	*Idotea montereyensis*	Isopod	0.73
54	***Idotea resecata***	**Isopod**	**0.36**
55	*Munna sp.*	Isopod	0.20
56	*Pugettia producta*	Brachyuran crab	0.36
57	*Pugettia richii*	Brachyuran crab	0.34
58	*Hippolyte californiensis*	Caridean shrimp	0.45

Table S2. Summary of traditional biodiversity metrics for each site

Site abbrev	Region	Species richness	Raw abundance	Effective diversity (Hill number)
DC	Barkley Sound	21	1168	11.8
RP	Barkley Sound	17	1590	8.8
SA	Barkley Sound	17	924	6.8
DK	Clayoquot Sound	21	1168	11.8
EB	Clayoquot Sound	21	4812	8.5
IN	Clayoquot Sound	16	1803	7.5
CB	Gulf Islands	18	2112	5.0
GB	Gulf Islands	11	2239	4.3
JB	Gulf Islands	22	3522	7.1
LH	Gulf Islands	15	2259	6.6
SS	Gulf Islands	10	740	6.6
HL	Haida Gwaii	9	16743	1.3
RA	Haida Gwaii	14	934	8.6

Equation S1. Equations for D^2_{adj} calculations

Explained deviance or D^2 (Guisan and Zimmerman 2000) was calculated by the equation:

$$D^2 = 1 - Residual\ deviance\ /\ Null\ deviance$$

Where the residual deviance refers to the deviance in predicted species occurrences, and the null
deviance refers to deviance in observed species occurrences.

Adjusted D^2 (Guisan and Zimmerman 2000) was calculated by the equation:

$$D^2_{adj} = 1 - (((n-1)/(n-p-1)*(D^2-1))$$

Where n is the number of sites (13) in our study, and p is the number of parameters the model
estimated (4, Ovaskainen et al. 2017, Supporting Information). Negative D^2_{adj} values (those which
had greater residual deviance than null deviance) were set to 0.

Equation S2. Calculations for type III variation partitioning on HMSC model (Borcard et al. 1992)

We estimated three versions of the HMSC model: 1) environmental variables only with the random
effects (*m2*), and 2) spatial distances only with random effects, and 3) the original global model
which included environmental and spatial variables, site- and region-level random effects (*m1*).
From each model, we extracted the total variation explained by environment and space, the
variation explained by environment, and the variation explained by space. To determine the shared
fraction of variation between environment and space, we multiplied each whole fraction by the
calculated community-level D^2_{adj} to reflect the model deviance in our variance partitioning
estimates, and followed the equations

m1 = the fraction of variation explained by all environmental variables in the HMSC model
estimated with only environmental variables;

m2 = the fraction of variation by spatial distances in the version of the HMSC model estimated
with only spatial distances;

m3 = the global model including both;

Fraction *ab* (pure environment + shared fraction) = D^2_{adj} *m1*

Fraction *bc* (pure space + shared fraction) = D^2_{adj} *m2*

Fraction *abc* (pure environment + shared fraction + pure space) = D^2_{adj} *m3*

Fraction *a* (pure environment) = *abc - bc*

Fraction *c* (pure space) = *abc - ab*

Fraction *b* (shared fraction) = *ab + bc - abc*

Fraction *d* (residuals) = *1 - abc*

Fractions *a* ("Environmental conditions"), *c* ("Spatial distance"), *b* (shared fraction), and *d*
("Residuals") are shown in the Venn diagram in Fig. S3.

Figure S2. Site-by-species presence-absence matrix of all 58 invertebrate species in the study

listed in alphabetical order. Black cells indicate species presences.

809   **Figure S3.** Venn diagram summarizing the fractions of variation explained by environmental

810   covariates only (nine water quality and five biotic variables), spatial distance only, and the shared

811   fraction explained by environment and space.

 Figure S4. A correlation plot showing modelled site-level co-occurrence of species pairs across

 all species. Purple cells represent positively co-occurring species pairs, and turquoise cells

 represent negatively co-occurring species pairs. Species names along both axes are ordered

 according to the output of hierarchical clustering with Ward's criterion (Ward 1963) on pairwise

 co-occurrence values. This figure is a supplement to Fig. 4a in the main text.

Figure S5. Heat map depicting abundance patterns in all 58 species in the study. Species names

are ordered from highest to lowest predicted mean proportional abundance. Cell colours

correspond to the co-occurrence groups in Fig. 4b, with turquoise cells representing members of

the *Nereis* group, purple cells representing members of the *C. californica* assemblage, and grey

cells indicating species that did not significantly co-occur negatively or positively with other

species. Cell shade strength represents proportional abundance at a given site (darker means higher

abundance). Most species outside the top twenty most abundant had extremely low predicted

proportional abundances owing to their low raw abundances. This figure is a supplement to Fig.

4b in the main text.

Figure S6. Plot showing relationship between maximum site-level abundance and site occupancy for all species. Black points represent broadcast spawners and white points represent brooders. Spirorbid polychaetes were excluded from this figure due to their unusually high maximum abundance (approximately 15000 at one site). A two-tailed t-test showed that there was no significant difference in the number of sites colonized between brooders and broadcast spawners (t = 0.052, df = 42.5, p = 0.96).