Beyond a single patch: local and regional processes explain diversity patterns in a seagrass epifaunal metacommunity

Keila A Stark[1]*, Patrick L Thompson[1], Jennifer Yakimishyn[2], Lynn Lee[3], Emily M Adamczyk[1], Margot Hessing-Lewis[4], Mary I O'Connor[1]

1. Biodiversity Research Centre
University of British Columbia
Vancouver, BC, Canada

2. Pacific Rim National Park Reserve
P.O. Box 280
Ucluelet, BC, Canada

3. Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve, and
Haida Heritage Site
60 Second Beach Road
Skidegate, BC, Canada

4. Hakai Institute
PO Box 309
Heriot Bay, BC, Canada

*Correponding author: keilastark@zoology.ubc.ca

Abstract: 286
Main text: 5389
Number of references: 65
Number of figures: 3

Data accessibility: Data will be made permanently available on Dryad should the manuscript be accepted. All data and code for the analysis can be currently found at https://github.com/keilast/HMS-Seagrass.

Competing interests: We have no competing interests.

Abstract

Ecological communities are jointly structured by dispersal, density-independent responses to

environmental conditions, and density-dependent biotic interactions. Metacommunity ecology

provides a framework for understanding how these processes combine to determine community

composition among local sites that are regionally connected through dispersal. In 17 temperate

seagrass meadows in British Columbia, Canada, we tested the hypothesis that eelgrass (*Zostera*

marina L.) faunal assemblages are dispersal limited, such that meadows separated by greater

spatial distances host distinct epifaunal invertebrate fauna. We used hierarchical joint species

distribution modelling to understand the relative contribution of environmental conditions,

spatial distance between meadows, and unexplained variation on species distributions. We found

that modelled species compositional similarity did not decay as a function of spatial distance,

and sites in the same region were no more similar to each other than sites farther away. These

results suggest that dispersal limitation of these invertebrates does not strongly influence

diversity over 1000 km of coastline. Abiotic environmental conditions such as temperature,

nitrate levels, and salinity partially explained regional patterns of abundance, but their relative

importance varied across taxa. We found co-occurrence patterns between taxa that could not be

explained by shared responses to the environment, suggesting the possibility of interspecific

interactions influencing community structure. Our results support findings from previous studies

that seagrass faunal communities tend to be somewhat homogenized across regional spatial

scales even up to 1000 km. Minor variations in community structure across sites could be

attributed to the abiotic environment, or intraspecific interactions at the site or meadow-level

spatial scale. Our results give refined hypotheses about the specific ecological mechanisms

driving community structure at various spatial scales, which in turn can inform targeted

experiments, monitoring, and management decisions for maintaining diversity in this important

ecosystem.

Key words: Hierarchical Modelling of Species Communities, priority effects, dispersal, niche

filtering, British Columbia, *Zostera marina*

Introduction

Understanding how local environmental conditions, regional connectivity by dispersal,

biotic interactions, and ecological drift jointly structure the composition of communities is a

central challenge in ecology (Ricklefs and Schluter 2003, Vellend 2010, Leibold and Chase

2017). Metacommunity theory (Leibold et al. 2004, Holyoak 2005, Leibold and Chase 2018)

offers a promising framework for understanding drivers of community assembly across spatial

scales. Applications of the framework to date have often focused on categorizing

metacommunities into one of four discrete archetypes (e.g. Cottenie 2005, Logue 2011) that each

reflect key hypothesized differences in the mechanisms that structure dynamics (Leibold et al.

2003). However, recent papers have called for metacommunity theory to move beyond the four

archtypes and instead focus on the underlying processes that govern the dynamics of

communities (Brown et al. 2017, Leibold and Chase 2018, Thompson et al. 2019). Thompson et

al. (2019) reframed the metacommunity concept based on three fundamental processes that

together govern the dynamics of populations and communities. These are: 1) Densitiy-

independent responses to environmental heterogeneity, 2) Density-dependent biotic interactions

(ie. inter and intra-specific competition influencing population growth and co-existence), and 3)

Dispersal influencing connectivity across a landscape or seascape. We apply this framework to

understand the relative contribution of core assembly processes to a seagrass-associated

invertebrate metacommunity.

Many coastal environments that host high biodiversity occupy spatially structured habitats

(e.g. coral reefs, kelp forests, seagrass meadows), and many marine species have dispersing life

histories that connect populations in distinct habitat patches. The importance of interspecific

interactions (Berlow 1999, Stachowicz 2001, Sala and Graham 2002), and dispersal-driven

population dynamics (Levin and Paine 1974, Gaines and Roughgarden 1985) in coastal and

marine communities has long been recognized. These combined roles of dispersal, the local

environment, and biotic interactions suggest patterns of marine biodiversity likely reflect more

than simple abiotic variation, and instead likely reflect metacommunity processes (Boström et al.

2006). To date, marine metacommunity dynamics have been investigated in sandy-bottom

benthic invertebrates (Valanko et al. 2015, Rodil et al. 2018), coral reef fishes (MacNeil et al.

2009), kelp-associated invertebrates and fishes (Lamy et al. 2018), rocky intertidal communities

(Salois et al. 2018), and temperate seagrass epifauna (Yamada et al. 2014, Whippo et al. 2018).

Seagrasses, like corals, salt marsh plants, and mangroves, are foundation species that support

high productivity and faunal diversity (Orth et al. 1984, Duffy et al. 2015). Seagrasses are often

prioritized in coastal protection plans because of the ecosystem services they provide, and many

of these services stem from their capacity to support high biodiversity and secondary

productivity. Seagrasses form meadows separated from each other by deeper water, un-vegetated

seafloor, or other vegetated habitats, and associated fauna disperse among meadows (Boström et

al. 2006, 2011). Highly diverse and productive assemblages of invertebrates live in and among

seagrass leaves, providing rich food sources for fish and birds. Invertebrates in this system such

as snails, gammarid amphipods, isopods, peracarid shrimp and polychaete worms are not only

important to seagrass-based food webs, but also process energy and organic matter in these

productive coastal ecosystems. The grazers within this group consume detritus, macroalgae, the

seagrass itself (Valentine and Heck 1999), and many of them exert top-down control on

epiphytic microalgae that compete with the seagrass for light and nutrients in the water column

(Duffy et al. 2015). The animals that live in seagrass meadows exhibit a range of dispersal modes

(fast swimming by adult isopods; slow crawling as in snails; and permanent attachment as in

some bivalves) and reproductive strategies (brooding versus broadcast spawning) that likely

influence dispersal rates and distances. Seagrass meadows can occur in a wide range of

temperature, salinity and hydrodynamic environments, such that meadows may differ in their

abiotic or biotic suitability for some invertebrate species. Finally, invertebrates on eelgrass

interact through facilitation, competition and predation.

Decades of research have identified the influence of local characteristics such as habitat

complexity (Orth et al. 1984, Hovel and Lipcius 2001), primary biomass (Cébrian and Duarte

1998, Gullström et al. 2012), and trophic interactions (Amunudrud et al. 2015, Huang et al.

2015) to faunal richness and abundance. However, more recent evidence suggests that when one

expands the spatial scale of observation to include many meadows within a region, these local

effects are overridden by regional effects such as habitat fragmentation and landscape

heterogeneity (Bostrom et al. 2010, Lefcheck et al. 2016, Yeager et al. 2019).

The conservation value of seagrass may reflect more than the simple presence of this

foundation species. The capacity of seagrasses to support biodiversity may also depend on how

meadows can support metacommunities of invertebrates in spatially structured landscapes

(Bostrom et al 2006). Metacommunity processes may play important roles maintaining the high

diversity of animals within and among seagrass meadows, and if this is true, then management of

seagrass for its ecosystem services may be more effective if biodiversity connectivity and

metacommunity processes are considered explicitly in the spatial scales of protection plans.

The main aim of this study is to better understand the ecological processes that form faunal

diversity across a seascape. To address this aim we ask three main questions: (1) What is the

contribution of spatial distance (a rough proxy for dispersal limitation) to community similarity?

(2) Does variation in local environmental factors drive cariation in the presence and abundance

of taxa across sites, and if so, which factors are most important? (3) Do species co-occurrence

patterns reveal any biotic interactions that might influence community structure?

We hypothesize that: (1) Spatial distance does not confer community dissimilarity because

sites are well-connected by dispersal, at even at larger spatial scales (Kinlan and Gaines 2003);

(2) Differences in local-scale environmental factors such seagrass shoot density and biomass,

temperature, salinity, and water column nutrient levels will drive differences in diversity, but not

necessarily taxonomic identity. (3) Species will not show any strong co-occurrence patterns due

to high dispersal and environmental responses overriding abiotic interactions.

We address these questions by fitting a joint species distribution model with Bayesian

estimation to an invertebrate biodiversity survey in 17 *Zostera marina* (L.) meadows spanning

the entire coast of British Columbia, Canada (approximately 1000 km). We found evidence of

connectivity between sites, even among the most distant sites within our dataset. We also found

that species responded differently to different environmental variables, with abiotic water quality

variables explaining the most variation in taxonomic abundance. Strong co-occurrence patterns

that could not be attributed to shared responses to the environment led us to hypothesize possible

biotic interactions influencing community structure.

Methods

Study sites and field sampling of eelgrass and associated epifauna

We sampled epifaunal biodiversity in 17 eelgrass *Zostera marina* (L.) meadows along

1000 km of coastline from late June through early August in 2017. The sites were located within

five major regions: Haida Gwaii within Gwaii Haanas National Park Reserve, National Marine

Conservation Area Reserve and Haida Heritage Site; the central coast of BC near Calvert Island

and surrounding areas; southern Clayoquot Sound (one site was within Pacific Rim National

Park Reserve); Barkley Sound on the West coast of Vancouver Island; the Southern Gulf Islands

within Gulf Islands National Park Reserve (Fig. 1a). We used a spatially hierarchical sampling

method to estimate biodiversity at three nested spatial scales: the quadrat (or plot) -level) scale,

the meadow (or site) -level scale, and the regional scale (Appendix 1: Fig. A1). These sites

varied in environmental conditions, such as seagrass shoot density and meadow size, site

exposure and hydrological regimes, freshwater outflow influencing seawater salinity, and others.

We sampled three sites in Barkley Sound (SA, DC, RB) and the central coast (TB, TB,

CI, CS) with SCUBA. All other sites were sampled by wading or snorkeling at low tide. We

used 0.25 m x 0.25 m quadrats to collect all above ground eelgrass, epifaunal invertebrates,

eelgrass detritus, and macroalgae in each sampled area. Six quadrats were set in a 15 m x 30 m

array (Appendix 1: Fig. A1); the array was placed in the middle of the meadow to avoid edge

effects where possible. We uprooted eelgrass shoots at the first node to avoid infaunal effects

where possible, removed all other above-ground biomass (detritus, macroalgae, associated

epifauna) within each quadrat by hand, and immediately placed all the contents into a 300 μm

mesh or plastic Ziploc bag.

Abiotic environmental variables

We acquired water quality and biotic data. The water quality covariates were annual

means pulled from Bio-ORACLE (Assis et al. 2018), including surface chlorophyll-*a*, dissolved

oxygen, nitrates, salinity, maximum current velocity and temperature (mean and range). Bio-

ORACLE is a database of marine environmental layers gathered from several satellite and in-situ

sources at a spatial resolution of 0.08°, and has been been demonstrated in accurate species

distribution modelling of shallow-water organisms (Tyberghein et al. 2012, Assis et al. 2018).

Sea surface temperature and chlorophyll-a layers were taken from the Aqua-MODIS satellite,

and values for nitrates, dissolved oxygen, pH, and salinity were interpolated from in-situ

measurements reported in the World Ocean Database (Tyberghein et al. 2012). All sites

(meadows) were situated in different spatial cells, and thus had distinct values for the

aforementioned variables.

Seagrass, algae, and detritus biometrics

For each quadrat sampled, we measured five biotic attributes summarizing habitat

structure and food availability: eelgrass leaf area index (LAI) and shoot density, and eelgrass dry

mass, eelgrass detritus dry mass, and algal dry mass. To quantify LAI, we measured leaf length,

width, and blade number in five haphazardly-chosen shoots from each quadrat, and multiplied

the average blade area (m^2) per shoot by the shoot density (number of shoots per m^2). We

separated and dried eelgrass, detritus, and macroalgae in a desiccator oven (60° C for 48 hours).

Invertebrate sample processing and identification

Immediately after collection from the field, we rinsed eelgrass blades with fresh water to

remove epifaunal invertebrates; the collected water containing invertebrates was passed through

a 500 μm sieve. Invertebrates were preserved in 95% EtOH for identification with light

microscopy (10x magnification). We identified epifauna to the lowest taxonomic level possible

using keys in the Light and Smith manual (Carlton 2007) and Kozloff (1996). A complete

species list including rare species can be found in the Appendix 1.

Analysis

All statistical analyses were conducted in R (version 3.4.2; R Development Core Team 2014). We

followed the Hierarchical Modelling of Species Communities (HMSC) framework (Ovaskainen et

al. 2017) using the 'Hmsc-R' package (Tikhonov et al. 2019) to fit a hierarchical joint species

distribution model (JDSM) with Bayesian inference. The framework uses traditional single-

species distribution modelling by estimating the species responses to environmental covariates

across space, but does so for all species simultaneously. Residual variation from environmental

covariates can be used to explain species co-occurrences that are not resulting from shared

response to the environment (Ovaskainen et al. 2017). It is also capable of accounting for spatially

hierarchical sampling structures. In these ways, the HMSC framework overcomes statistical

limitations of previous methods used in metacommunity ecology (Gilbert and Bennett 2010,

Tuomisto et al. 2012, Brown et al. 2017).

Each species distribution model is given by a generalized linear model that describes the

abundance of species j (where $j = 1… n$), where y_{ij} *is* the count of species j at site i, D is the

statistical distribution of the data (Poisson distribution for count data in this study), is the linear

predictor, and σ^2_j is a variance term for count of species j.

$$y_{ij}^{D} \sim D(L_{ij}, \sigma_j^{2}).$$

230   The linear predictor L_{ij} is described by fixed (F) and random (R) effect terms:

232
$$L_{ij} = L^F_{ij} + L^R_{ij}$$

233
234   The fixed effect term L^F_{ij} is modelled as a regression:

235
$$L^F_{ij} = \beta_{jk} x_{ik}$$

237   where x_{ik} represents the measured value for environmental covariates k for a given sample i (ie.,

238   the measured eelgrass biomass in quadrat i), and parameter β represents the relationship between

239   that environmental covariate k, and the abundance of a particular species j.

240   The random component L^R_{ij}, captures the variation in species occurrences and co-occurrences that

241   cannot be explained by the measured covariates. It is further denoted as

242
$$L^R_{ij,} = \varepsilon_{ij}^Q + \varepsilon_{ij}^S + \varepsilon_{ij}^R + \varepsilon_{ij}^D$$

where the terms describe three hierarchical random effects associated with our sampling scheme

(ε_{ij}^Q, ε_{ij}^S, ε_{ij}^R for Quadrat, Site, and Region), as well as a fourth spatially explicit random effect.

The spatially explicit random effect ε_{ij}^D was calculated with latent factor analysis that takes into

account spatial distance between all pairwise combinations of sampling sites (see Ovaskainen et

al. 2016a for details).

The three sampling random effect terms are assumed to have multivariate normal distributions ε_{ij}^Q

$\sim N(0, \Omega^Q)$, $\varepsilon_{ij}^S \sim N(0, \Omega^S)$, $\varepsilon_{ij}^R \sim N(0, \Omega^R)$. Variance terms Ω^Q, Ω^S, Ω^R, are variance-covariance

matrices (square matrices containing all taxa in the model), where the diagonal elements give

species-specific residual variance in occurrences among sites, and the off-diagonal elements give

residual co-variances between species pairs. The term "residual" refers to the fact that it is the

amount that is not explained by environmental covariates in the fixed effects predictor described

above. These variance-covariance matrices (Ω^Q, Ω^S, Ω^R, and Ω^D) are parameters estimated using

a latent variable approach described in Ovaskainen et al. (2016b). They were used to represent co-

occurrence matrix R, where j_1 and j_2 refer to two species within the model, and

$R_{j1j2} = \Omega_{j1j1}/\sqrt{\Omega_{j1j1}\Omega_{j2j2}}$. R describes the extent to which a given species pair co-occurs more

positively or negatively than by chance, after controlling for possible shared responses to the same

environmental covariates.

We estimated the parameters of the joint species distribution model with Markov chain

Monte Carlo. The Markov chains were run to 200 000 iterations, a burn-in length of 1000

iterations, and was thinned to retain every 100[th] sample of the posterior distributions. We

confirmed that Markov chains were well-mixed by visually inspecting trace plots. Parameter

estimates of β_j (environmental responses across taxa) and Ω (variance-covariance matrices used

to visualize species co-occurrence patterns independent of the environment) were extracted as

95% credible intervals. We evaluated the model fit using 2-fold cross-validation.

To test our three hypotheses about the contribution of metacommunity processes, we

visualized different aspects of the parametrized JDSM (See Appendix 1: Table A1 for summary).

We investigated three possible dispersal conditions: dispersal limitation, intermediate dispersal,

and high dispersal. To investigate dispersal limitation at large spatial scales, we plotted distance-

decay of similarity by randomly drawing 3000 pairwise distances between sampling units, and

plotting predicted community similarity against spatial distance. Distance-decay of community

similarity could reflect dispersal limitation due to prohibitively far travel distances, *or* due to

differences in environmental conditions with increasing spatial scale. To distinguish between

these possibilities, we compared the predicted community similarity using the full model

(including environmental covariates) with the predicted community similarity using an

alternative model where the effect of the environment was removed (all environmental covariates

were set to their mean values).

To test the hypothesis that differences in environmental conditions among meadows

allowed different species to colonize and persist in each meadow ('environmental niche

filtering'), we first partitioned variation among environmental covariates, spatial autocorrelation,

and random effects associated with the sampling design (quadrat, site, and region), and

visualized the contribution of these in a stacked bar plot.

To test the hypothesis that some species co-occurred more often than would be expected

by chance or by their shared environmental niches, we extracted correlations between species

from the residual variation in our model, with latent factor loading (Ovaskainen et al. 2017) and

represented modelled pairwise associations with correlation plots describing the extent to which

species pairs co-occurred more negatively or positively than predicted by their environmental

niches alone (Ovaskainen et al. 2017, Aivelo and Norberg 2018). We show species co-

occurrences at the quadrat, site, and region random effect levels. To visualize patterns in site-

level co-occurrences across sites, we conducted hierarchical cluster analysis on sites according to

observed species abundances using Ward's criterion (Ward 1963). We mapped the cluster

dendrogram onto an abundance heatmap of the 10 most abundant taxa in our analysis, and

colour-coded the cells according to the species co-occurrence groupings from the correlation

plots (groups of species that tended to positively co-occur amongst themselves). Full versions of

these figures with all species are provided in Appendix 1: Fig. A4 and Fig. A5.

Results

Species abundance and distribution patterns

We counted and identified 52 282 individuals representing at least 50 taxa across the

region (Appendix 1: Table A2). We excluded rare species (those found in fewer than 5% of all

samples, or fewer than 5 out of 102 quadrats) from our JDSM to avoid statistically over-inflating

the importance of model covariates; this left 33 taxa in our analysis. The represented taxa span

multiple phyla, diet types (herbivores, detritivores, suspension feeders, carnivores), and life

history strategies (brooding, broadcast spawning). Mean quadrat-level species richness was 13

species, and mean site-level richness was 16 species. A full summary of diversity metrics can be

found in Appendix 1: Table A3.

Several taxa were present at all 17 sites (Appendix 1: Fig. A2). These include

harpacticoid copepod *Porcellidium* sp., bivalves *Saxidomus gigantea* and *Clinocardium nutallii*,

snails *Alvania compacta*, *Lacuna* spp., and the tanaid *Leptochelia* sp. Genera *Mytilus* (a mussel)

and *Nereis* sp. (a polychaete worm) were represented at all sites, however we were not confident

the same species comprised these groups. Ten species were present in at least one site in every

region: isopod *Penidotea resecata,* gammarid amphipods *Monocorophium insidiosum, Ampithoe*

valida, Pontogeneia rostrata, Ampithoe dalli, Aoroides spp*.,* snails *Amphissa columbiana* and

Alia carinata, and limpet *Lottia pelta.*

Dispersal: Distance-decay of community similarity due to spatial distance and environment

In the full joint species distribution model summarizing species' responses to

environment and spatial distance, predicted community similarity decreased with increasing

distance between sites (Fig. 1b, red points and line). However, community similarity showed a

slight increase with distance when the effect of environmental variables was removed from the

model (all environmental variables set to their mean value). We infer that spatial distance alone

did not explain differences in invertebrate compositional similarity (Fig. 1b, blue points and

line), but that differences in habitat conditions among meadows better explained divergence in

community composition.

Environmental niche filtering: variation partitioning

Consistent with the previous analysis, we found that patterns in abundance and

distribution were consistent with weak environmental niche filtering along an abiotic axis. Water

quality variables (yellow bars in Fig. 2) explained the largest proportion of variation across

nearly all species distributions. Herbivore food availability (seagrass, algae and detritus biomass)

and habitat structure (eelgrass shoot density and LAI) had minimal influences on species

distributions (pink bars in Fig. 2). The importance of environmental covariates varied across

species; for example, Fig 2). While the mean R^2 across all species distribution models was 0.27,

individual R^2 values were highly variable (0 to 0.68, Appendix 1: Table A2).

Species interactions: Patterns in pairwise species co-occurrences

At the scale of interactions among individuals (site-level samples), positive co-

occurrences tended to be more common than negative co-occurences, suggesting facilitation as a

possible important species interaction. Co-occurrence patterns were not strong at the quadrat and

region–levels (Fig 3a and 3b).

At the site level (between-sites), we identified, *post hoc*, two main species co-occurrence

groupings. Members of the first group (hereafter we call this the *"Nereis"* group), *Leptochelia*

sp. (tanaid), *Nereis* polychaetes, spirorbid polychaetes, and *Caprella laeviuscula* (caprellid

amphipod) (Fig. 3b), tended to positively co-occur more than expected by chance or by

environmental filtering, and negatively co-occur members of the second group. The second

group included a Harpacticoid copepod species, the isopod *Porcellidium* sp., and an gammarid

amphipod within the family Ischyroceridae (Fig. 3b). Other species did not show strong positive

or negative co-occurrence values with others.

The two co-occurrence groups were present at all sites, indicating that members from

these groups can be simultaneously present (Fig. 3d). However, their abundances tended to

negatively co-vary; at sites JB, HL, CB, RA, EB, DC, and DU, members of the *Nereis* group

were predicted to be most abundant, whereas other group members were predicted to be most

abundant at SS, SA, LH, GB and RB. The site HL was strongly dominated by *Spirorbis* sp. The

emergent species groupings identified in Fig. 3b do not clearly correspond with spatial distance

because the cluster analysis did not group sites by spatial proximity (black symbols in Fig. 3d);

this is another line of evidence suggesting that dispersal is not limited at this spatial scale.

Discussion

This work is the largest, spatially explicit seagrass metacommunity study in geographical

extent to our knowledge. We tested the hypotheses that dispersal, environmental heterogeneity

and species interactions influenced spatial patterns in this system, consistent with

metacommunity dynamics. We found support for the importance of environmental niche filtering

and biotic interactions as drivers of biodiversity patterns, but surprisingly, we found little support

for dispersal limitation even across over 1000 km of coastline. Overall, these results suggest that

seagrass-associated biodiversity patterns reflect ecological processes spanning local (meadow-

scale) to regional scales.

Is there evidence of dispersal limitation among eelgrass meadows?

Our finding that the effect of spatial distance alone does not confer community

dissimilarity, along with the low spatial autocorrelation with environmental covariates, suggests

that spatial distance between sites likely does not result in dispersal limitation. This is true for the

six taxa that were present at all sites, and the additional ten taxa that were not present at every

site but present in all regions (Appendix 1: Fig. A2). These widely distributed groups had

representation from across phyla and life history traits, suggesting that no single taxonomic

group or dispersal life history strategy had a consistently greater spatial distribution than any

other. Many of the meadows were small (< 1 ha), and epifaunal abundances can vary

substantially within meadows from year to year (Adamczyk et al. in prep) such that local

extinctions followed by rescue from nearby populations are likely important (Thom et al. 1995).

However, the specific pathway the organisms travelled to arrive at these sites remains unknown.

We used Euclidean distances in our analysis, which preserved the rank order of distances

between sites, but actual distances are likely greater due to oceanographic circulation patterns

(Kinlan and Gaines 2003, Mitarai et al. 2008).

It is unclear whether distant sites share the same species because they are 1) linked by

direct dispersal via oceanographic processes such as currents; 2) indirectly linked by dispersal

via unsampled "stepping-stone" sites; or 3) were colonized by populations of the same species in

a historical dispersal event, but have not seen the exchange of individuals since. Seagrass-

associated epifauna generally disperse passively between meadows, either by larval transport in

currents or by "rafting" on floating pieces of seagrass or macroalgae. Rafts have been observed

hosting benthic and epifaunal invertebrates across phyla (echinoderms; peracarids; molluscs;

annelids), and contribute to connectivity between coastal ecosystems across 100s of kilometres

(Wichmann et al. 2012).

Species that were only found at one site or region (e.g. *Jassa marmorata,* Appendix 1:

Fig. A2) may be dispersal limited for reasons other than prohibitively long travel distances.

Populations with low abundances may not disperse in appreciable numbers, and therefore cannot

establish populations at all sites as readily as abundant species. This is supported by our

observation of a generally positive relationship between range size (the number of sites a species

was represented at) and abundance (Appendix 1: Fig. A6).

Theory predicts that intermediate dispersal (dispersal sufficiency) allows species to

colonize sites where local environmental conditions are optimal for growth and reproduction,

whereas high dispersal (dispersal surplus) overrides this (Mouquet and Loreau 2003). Therefore,

it is likely that dispersal rates are high enough for several species to be present at most sites, but

not so high as to completely overwhelm the effects of environmental conditions. A subset of

species was present at all sites, but often at varying abundances (Fig 3d). This may indicate weak

mass effects, where immigration allows populations to persist even if local conditions are not

optimal (Schmida and Wilson 1985, Mouquet and Loreau 2003). Metacommunity theory

predicts that such weak mass effects are likely to be found whenever dispersal rates are not

limiting (Thompson et al. 2017), particularly in organisms that cannot control their own

dispersal, (Leibold and Chase 2017) such as seagrass-associated invertebrates. This may be

supported by our result on the contribution of the environment, particularly in ubiquitous species

for which the model fit was poor (e.g. *Aoroides* spp., *Spirorbis* sp., $D^2_{adj} = 0$); the poor model fit

may indicate that these persist at sites even if local environmental conditions are not optimal, due

to sufficient immigration. An alternative explanation for the model's poor predictive power for

these species is that these taxa have broader environmental niches, and thus the regional

environmental gradient is not sufficient to drive clear presence-absence patterns.

Seagrass epifauna show varying degrees of environmental niche filtering

The abiotic environmental axis had the largest influence on species distributions, with

phosphates, salinity, and sea surface temperature explaining the largest fractions of variation on

average in the joint species distribution model (Fig. 2). In some species distribution models that

had a higher fit (e.g. *Ampithoe dalli* and *Caprella natalensis*, Appendix 1: Table A2),

environmental niche filtering may be important, but overall there remains a large amount of

variation unexplained by the environment.

Environmental heterogeneity may be influencing species distributions and abundances

across the region in one of three ways. First, the most important abiotic environmental covariates

(mean and range in ocean temperatures, nitrates, dissolved oxygen) may influence food

availability through primary productivity. This is possible despite the fact that primary biomass

explained less variation than abiotic parameters (Figure 2). While the primary biomass

measurements were taken at a single time point, the abiotic parameters represent a four-year

average (Assis et al. 2018) and may therefore better represent the conditions these organisms

experience over their lifetimes or across generations. Second, sea surface temperature, dissolved

oxygen, and salinity may affect species' distributions via environmental tolerance ranges. For

example, the isopod *Idotea baltica* (a relative of *P. resecata* and *P. montereyensis* in our study)

experiences significantly lower survival in prolonged exposure to higher temperatures and lower

salinities than those experienced in their home habitat (Rugiu et al. 2018). Third, temperature

may influence the metabolic demands of invertebrates (many of which are herbivores), thereby

influencing grazing rates and altering the available primary biomass (O'Connor 2009). Decreased

food resources may increase competition, thus influencing the number and identity of species a

patch of seagrass can host. Determining whether population-level responses to competitive stress

influence whole-community structure would improve our understanding of how the environment

and species interactions combine to influence community structure.

Environmental conditions in temperate eelgrass meadows fluctuate seasonally, driving

changes in epifaunal community structure throughout the year (Wlodarska-Kowalczuk et al.

2014). Because these species may not show the same distributions year-round, sampling several

times in a year to capture these temporal changes would give a clearer picture of species'

responses to all metacommunity processes.

Species co-occurrence patterns may suggest priority effects and trophic interactions

We did not observe strong quadrat-level co-occurrence patterns; this is consistent with

previous findings that epifaunal diversity varies randomly within meadows (between quadrats)

(Whippo et al. 2018). We also did not observe strong region-level co-occurrence patterns,

suggesting that there do not appear to be clear regional assemblages organized according to the

bioregions we have defined (Fig. 1a). At the site-level, two main co-occurrence groupings

emerged showing generally antagonistic abundances across sites (Fig. 3d). Stochasticity alone is

unlikely to have produced such a strong pattern, especially given deterministic processes tend to

be more important in systems that are not dispersal limited (Lee and Bruno 2009, Ron et al.

2018). While there are multiple possible explanations for non-random species co-occurrence

patterns (Connor and Simberloff 1979), space and the measured environmental covariates were

ruled out since co-occurrence values were extracted from residual variation unexplained by

environment or spatial autocorrelation (Ovaskainen et al. 2017). This, along with the high

unexplained fraction in the variation partitioning leads us to hypothesize the importance of biotic

processes. Several types of interactions may be occurring in the metacommunity; the majority of

species are herbivores or detritivores, and thus may compete for primary biomass. Laboratory

experiments have shown that behaviours such as grazing rate and habitat selection in amphipods

are altered in the presence of morphologically and functionally similar interspecific competitors

(Beermann et al. 2018). Other species in our study are filter feeders, and thus may not be

competing for food resources but possibly predator-free space or substrate on the eelgrass blades.

The community structure of a given site with certain local environmental conditions and

access to a certain regional pool may have multiple possible compositional states depending on

the arrival order of species; this phenomenon is known as priority effects (Fukami et al. 2016, Ke

and Letten 2018) and has been documented in marine fouling communities (Vieira et al. 2018).

The antagonistic co-occurrence groupings in our analysis not arising from shared responses to

the environment may suggest priority effects at the site-level or meadow spatial scale (Fig. 3b). It

is possible that, following the establishment of a new meadow or a disturbance event, the first

few species to colonize a seagrass meadow determine the success of subsequent colonizers,

regardless of local environmental conditions. As an illustrative example, *Leptochelia* sp. might

have been among the first to colonize sites CS, DC, IN, and HL (Fig. 3b and 3d), thereby gaining

a numerical advantage over species in the alternative cohort. Distibution patterns in the sessile

calcifying polychaete *Spirorbis* sp. may also indicate priority effects. It dominated the HL site;

16 300 individuals were found, several orders of magnitude higher than its abundance elsewhere

(Data S1). Site HL also had the lowest species richness of all 13 sites (9 species). Eelgrass shoots

there were completely covered by these *Spirorbis* sp., possibly preventing grazers in the

community from accessing epiphytic algae or the seagrass itself. The age, colonization history,

and disturbance regime of the meadows likely matter if priority effects are occurring; further

work must be done to better understand these temporal dynamics in temperate seagrasses.

Negative co-occurrence patterns could also reflect the higher abundance of a predator at

some sites. Field experiments have shown that changes in predation pressure by fish, shorebirds,

and predatory invertebrates can shift seagrass-associated epifaunal assemblages in a matter of

weeks (Amundrud et al. 2015; Huang et al. 2015). Differential predation pressure could also lead

to the pattern of diverging community composition that we have witnessed (Guzman et al. 2018).

A more comprehensive sampling effort of fish and bird predators would be required to determine

the degree to which trophic interactions structure these communities.

Positive co-occurrence patterns between certain species may additionally be a result of

positive biotic interactions. An example is seen in *Orchomenella recondita* - a gammarid

amphipod that lives in the gastrovascular cavity of the anemone *Anthopleura elegans* (Carlton

2007). This species was only recorded at the SS site, and specifically only found in quadrats

where *A. elegans* was collected with the seagrass shoots.

Conclusion

Using a new statistical approach to characterizing dispersal, environmental niche

filtering, and biotic interactions, we have developed refined hypotheses about the

metacommunity processes driving temperate seagrass epifaunal diversity on a regional scale. Our

study also gives a baseline spatial scale (1000 km) for comparing the contribution of these

processes across spatial scales and other ecological systems (Viana and Chase 2018).

Manipulative experiments and monitoring are now needed to test our refined hypotheses. One

could test the hypothesis that species sorting is relatively weak by measuring fitness differences

along the most important environmental axes. Given our hypothesis on the importance of priority

effects, one might experimentally manipulate the colonization order of key species in our study

to determine if resulting compositional states differ (Fukami et al. 2016). Testing our hypothesis

that populations within the metacommunity are not dispersal limited is more challenging, but

could be done using particle tracking models similar to those used to estimate larval dispersal

between coral reefs across thousands of kilometres (Treml et al. 2008). A population genetic

analysis could also ground-truth the extent of dispersal between allopatric populations.

Our findings are particularly relevant to conservation initiatives that aim to maintain

diversity in seagrass meadows; identifying the importance of local and regional processes allows

the conservation practitioner to better predict how habitat fragmentation and climate change

might influence diversity. Seagrass epifaunal communities may be quite resilient, given our

findings that environmental conditions minimally contribute to species distributions, and several

species across functional groups are consistently present across a fairly large spatial scale.

Acknowledgements

KAS conceived of the study with guidance from MIO. JY provided samples from IN, DK, and

EB. LL provided samples from RA and HL. EMA and MHL provided samples from TN, TB, CI,

and CS. KAS collected and processed samples, analyzed data, and wrote the first manuscript

draft. PLT provided recommendations for the analysis. KAS, PLT, and MIO contributed to

writing the manuscript. This research was financially supported by a UBC Zoology SURE Grant,

and NSERC Discovery Grants, Canadian Foundation for Innovation (CFI), and is sponsored by

the NSERC Canadian Healthy Oceans Network and its Partners: Department of Fisheries and

Oceans Canada and INREST (representing the Port of Sept-Îles and City of Sept-Îles). PLT is

supported by NSERC and Killam postdoctoral fellowships. We are grateful to the Tula

Foundation for providing logistical support and facilities during Calvert Island field collections.

Special thanks go to Michelle Paleczny and Gulf Islands National Park Reserve for boat and

laboratory space, and staff, students, and volunteers from Pacific Rim National Park Reserve for

their assistance collecting eelgrass. We thank Minako Ito, Ariane Comeau, Rob Underhill, Abbie

Sherwood, Giordano Bua for assistance with samples and Coreen Forbes, Matt Whalen, and

anonymous reviewers for feedback on the manuscript.

Literature Cited

Aivelo, T. and Norberg, A. 2018. Parasite-microbiota interactions potentially affect intestinal

communities in wild mammals. J. Anim. Ecol. 18:438-447.

Amundrud, S. L. et al. 2015. Indirect effects of predators control herbivore richness and

abundance in a benthic eelgrass (*Zostera marina*) mesograzer community. – J. Anim. Biol.

84: 1092–1102.

Assis, J. et al. 2018. Bio-ORACLE v2.0: Extending marine data layers for bioclimatic modelling.

– Global Ecol. Biogeogr. 27:277–284.

Beermann, J. et al. 2018. Combined effects of predator cues and competition define habitat

choice and food consumption of amphipod mesograzers. - Oecologia 186:645–654.

Berlow, E. L. 1999. Strong effects of weak interactions in ecological communities. - Nature 398:

330–334.

Boström, C. et al. 2006. Seagrass landscapes and their effects on associated fauna: A review. -

Estuarine, Coastal Shelf Sci. 68: 383-403.

Boström, C. et al. 2010. Invertebrate dispersal and habitat heterogeneity: Expression of

biological traits in a seagrass landscape. J. Exp. Mar. Biol. Ecol. 390:106-117.

Borcard, D et al. 1992. Partialling out the spatial component of ecological variation. - Ecology

73:1045–1055.

Brown, B. L. et al. 2017. Making sense of metacommunities: dispelling the mythology of a

metacommunity typology. - Oecologia 183:643–652.

Carlton, J. 2007. The Light and Smith Manual: Intertidal invertebrates from Southern California

to Oregon. Fourth edition. University of California Press, Oakland, California, USA.

Cébrian, J. and Duarte, C. M. 1998. Patterns in leaf herbivory on seagrasses. – Aquat. Bot.

60:67–82.

Connor, E. F. and Simberloff, D. 1979. The assembly of species communities: Chance or

competition? - Ecology 60:1132–1140.

Cottenie, K. 2005. Integrating environmental and spatial processes in ecological community

dynamics. – Ecol. Lett. 8:1175–1182.

Duffy J. E. et al. 2015. Biodiversity mediates top-down control in eelgrass ecosystems: A global

comparative-experimental approach. – Ecol. Lett. 18:696–705.

Economo, E. P. 2011. Biodiversity conservation in metacommunity networks: Linking pattern

and persistence. – Amer. Nat. 177:167-180.

Fukami, T. et al. 2016. A framework for priority effects. – J. Veg. Sci. 27:655–657.

Gaines, S. and Roughgarden, J. 1985. Larval settlement rate: A leading determinant of structure

in an ecological community of the marine intertidal zone. – Proc. Natl. Acad. Sci.

82:3707–3711.

Gilbert, B. and Bennett, J. R. 2010. Partitioning variation in ecological communities: do the

numbers add up? – J. Appl. Ecol. 47:1071–1082.

Guichard, F. 2003. Mussel disturbance dynamics: Signatures of oceanographic forcing from

local interactions. – Amer. Nat. 161:889–904.

573 Guillaume Blanchet, F. et al. 2018. HMSC: Hierarchical Modelling of Species Community. R

574   package version. 2.1-2.

575 Guisan, A. and Zimmermann, N. E. 2000. Predictive habitat distribution models in ecology. -

576   Ecol. Modell. 145:147–186.

577 Gullström, M. et al. 2012. Spatial patterns and environmental correlates in leaf-associated

578   epifaunal assemblages of temperate seagrass (*Zostera marina*) meadows. – Mar. Biol.

579   159:413–425.

580 Guzman, L. et al. 2018. Towards a multi-trophic extension of metacommunity ecology. – Ecol.

581   Lett. 22:19-33.

582 Huang A. C. et al. 2015. Top-down control by great blue herons *Ardea herodias* regulates

583   seagrass-associated epifauna. - Oikos 124:1492–1501.

584 Ke, P. J. and Letten, A. D. 2018. Coexistence theory and the frequency-dependence of priority

585   effects. – Nat. Ecol. Evol. 2:1691–1695.1–7.

586 Kinlan, B. P. and Gaines, S. D. 2003. Propagule dispersal in marine and terrestrial environments:

587   A community perspective. - Ecology 84:2007–2020.

588 Kozloff, E. 1996. Marine Invertebrates of the Pacific Northwest. University of Washington

589   Press, Seattle, Washington, USA.

590 Leibold, M. A. et al. 2004. The metacommunity concept: A framework for multi-scale

591   community ecology. – Ecol. Lett. 7:601–613.

592 Leibold, M. A. and Chase, J. M. 2017. Metacommunity Ecology, Volume 59. Princeton

593   University Press, Princeton, New Jersey, USA.

594 Leibold, M.A. and Mikkelson, G. M. 2002. Species turnover, and boundary clumping: Elements

595   of meta-community structure. - Oikos 97:237–250.

Lee, S. C. and Bruno, J. F. 2009. Propagule supply controls grazer community structure and

primary production in a benthic marine ecosystem. – Proc. Natl. Acad. Sci. 106:7052–

7057.

Lefcheck J. S. et al. 2016. Faunal Communities Are Invariant to Fragmentation in Experimental

Seagrass Landscapes. - PLoS ONE 11(5): e0156550.

Levin, S. A. and Paine, R. T. 1974. Disturbance, patch formation, and community structure. –

Proc. Natl. Acad. Sci. 71:2744–2747.

Logue, J. et al. 2011. Empirical approaches to metacommunities: a review and comparison with

theory. – Trends Ecol. Evol. 26:482–491.

MacNeil, M. A. et al. 2009. Hierarchical drivers of reef-fish metacommunity structure. –

Ecology 90:252-264.

Mitarai, S. D. et al. 2008. A numerical study of stochastic larval settlement in the California

Current system. – J. Marine Syst. 69:295–309.

Mouquet, N. and Loreau, M. 2003. Community patterns in source-sink metacommunities. –

Amer. Nat. 162:554–557.

O'Connor, M. I. 2009. Warming strengthens an herbivore-plant interaction. - Ecology 90:388–

398.

Orth, R. J. et al. 1984. Faunal communities in seagrass beds: A review of the influence of plant

structure and prey characteristics on predator-prey relationships. - Estuaries 7:339–350.

Ovaskainen, O. et al. 2016a. Uncovering hidden spatial structure in species communities with

spatially explicit joint species distribution models. - Methods Ecol. Evol. 7:428–436.

Ovaskainen, O. et al. 2016b. Using latent variable models to identify large networks of species-

to-species associations at different spatial scales. - Methods Ecol. Evol. 7:549–555.

Ovaskainen, O. et al. 2017. How to make more out of community data? A conceptual framework

and its implementation as models and software. – Ecol. Lett. 20:561–576.

Presley, S.J. et al. 2010. A comprehensive framework for the evaluation of metacommunity

structure. – Glob. Ecol. Biogeogr. 19:908–917.

R Development Core Team. 2014. R: a language and environment for statistical computing. R

Foundation for Statistical Computing. Vienna, Austria. http://www.R-project.org.

Ricklefs, R. E. and Schluter, D. 1993. Species Diversity in Ecological Communities: Historical

and Geographical Perspectives. University of Chicago Press, Chicago, Illinois, USA.

Rodil, I. et al. 2018. The importance of environmental and spatial factors in the metacommunity

of exposed sandy beach benthic invertebrates. – Estuar. Coast. 41:206-217.

Ron, R. et al. 2018. Dispersal increases ecological selection by increasing effective community

size. – Proc. Natl. Acad. Sci. 115:11280–11285.

Rugiu, L. et al. 2018. Variations in tolerance to climate change in a key littoral herbivore. – Mar.

Biol. 165:1–11.

Sala, E. and Graham, M. H. 2002. Community-wide distribution of predator-prey interaction

strength in kelp forests. – Proc. Natl. Acad. Sci. 99:3678–3683.

Schmida, A. and Wilson, M. V. 1985. Biological determinants of species diversity. – J.

Biogeogr. 12:1–20.

Thom, R. et al. 1995. Temporal patterns of grazers and vegetation in a temperate seagrass

system. – Aquat. Bot. 50:201–205.

Thompson, P. L. et al. 2017. Loss of habitat and connectivity erodes species diversity, ecosystem

functioning, and stability in metacommunity networks. - Ecography 40:98–108.

Treml, E. A. et al. 2008. Modeling population connectivity by ocean currents: a graph-theoretic

approach for marine conservation. – Landsc. Ecol. 23:19–36.

Tuomisto, H. et al. 2012. Modelling niche and neutral dynamics: on the ecological interpretation

of variation partitioning. - Ecography 35:961–971.

Tyberghein, L. et al. 2011. Bio-ORACLE: a global environmental dataset for marine species

distribution modelling. – Glob. Ecol. Biogeogr. 21: 272-281.

Valanko, S. et al. 2015. Complex metacommunity structure for benthic invertebrates in a low-

diversity coastal system. – Ecol. Evol. 5:5203–5215.

Viana, D. S. and Chase, J. M. 2018. Spatial scale modulates the inference of metacommunity

assembly processes. - Ecology doi:10.1002/ecy.2576.

Vieira, E. et al. 2018. Persistence and space preemption explain species-specifc founder effects

on the organization of marine sessile communities. - Ecology and Evolution 8:3430–3442.

Vellend, M. 2010. Conceptual synthesis in community ecology. – Q. Rev. Biol. 85:183–206.

Ward, J. H. 1963. Hierarchical grouping to optimize an objective ounction. - Journal of the

American Statistical Association 58:236–244.

Wichmann, C. S. et al. 2012. Floating kelps in Patagonian Fjords: An important vehicle for

rafting invertebrates and its relevance for biogeography. – Mar. Biol. 159:2035–2049.

Whippo, R. et al. 2018. Epifaunal diversity patterns within and among seagrass meadows suggest

landscape-scale biodiversity processes. - Ecosphere 9:02490.10.1002/ecs2.2490.

Wlodarska-Kowalczuk, M. et al. 2014. Evidence of season-dependency in vegetation effects on

macrofauna in temperate seagrass meadows (Baltic Sea). - PLoS One 9:e100788.

Yamada, K. et al. 2014. Environmental and spatial controls of macroinvertebrate functional

assemblages in seagrass ecosystems along the Pacific coast of northern Japan. - Glob. Ecol.

Cons. 2:47–61.

Yeager, L. A. et al. 2019. Trait sensitivities to seagrass fragmentation across spatial scales shape

benthic community structure. – J. Anim. Ecol. 88:1743-1754.

Figure legends

Figure 1. a) Map of coastal British Columbia showing eelgrass field sites in Haida Gwaii (HL,

RA), Clayoquot Sound (DK, EB, IN), Barkley Sound (SA, DC, RB), near Calvert Island in the

central coast (TB, TN, CS, CI), and the Southern Gulf Islands (GB, JB, CB, LH, SS). **b**) HMSC

model predictions of community similarity relative to the spatial distance (Euclidean) between

communities. The red circles show similarity predictions based on the full model. The blue

triangles show predictions wherein the effect of all environmental variables has been removed

(standardized to their mean values).

708   **Figure 1a)**

716   **Figure 1b)**

Pairwise site distance (km)

○ Space + environmental covariates
△ Space only

Figure 2.

Figure 2. Variation partitioning of fixed and random effects within the joint species distribution

model. Grey cells represent the contribution of random effects associated with the sampling design.

Yellow cells represent the contribution of abiotic water quality covariates. Pink cells represent

biotic covariates (food availability, habitat structure). Bars are organized according to species

abundances in descending order from left to right (most abundant to least abundant). The

percentages next to the legend labels indicate mean variation explained by that covariate/ random

effect across all species distribution models. The black circles below each stacked bar describe the

model fit to the data (R^2_{adj}). The mean R^2_{adj} of all species distribution models was 0.27.

Figure 3. a-c) Correlation plots showing modelled a) quadrat-level, b) site-level, and c) region-level pairwise co-occurrences after removing the effect of shared responses to the environmental covariates in our model. Only the 10 most abundant species are represented. Purple cells represent positively co-occurring species pairs, and turquoise cells represent negatively co-occurring species pairs. Species names are ordered according to the output of hierarchical clustering with Ward's criterion on pairwise co-occurrence values. A version of this figure with all 33 species is shown in Appendix S1: Fig. S4. **d)** Heatmap and cluster dendrogram depicting species relative abundance and compositional similarity across sites. Species are ordered by

decreasing abundance from left to right. Cell colours correspond to the site-level co-occurrence

groupings shown in Fig. 3b: turquoise or *Nereis* group, purple or *C. californica* group, and grey

Spirorbis group. Cell shade strength indicates proportional abundance at a given site (darker

means higher abundance). The black symbols to the left of the site abbreviations indicate region

membership; stars indicate Haida Gwaii sites, diamonds indicate Central Coast sites, triangles

indicate Barkley Sound sites, squares indicate Clayoquot Sound sites, and triangles indicate

Southern Gulf Islands sites. A version of this figure with all 33 species is shown in Appendix S1:

Fig. S5.

755