Regional connectivity, local environmental conditions, and biotic interactions jointly structure a temperate seagrass epifaunal metacommunity

Keila A Stark[1]*, Patrick L Thompson[1], Jennifer Yakimishyn[2], Lynn Lee[3], Mary I O'Connor[1]

1. Department of Zoology
Biodiversity Research Centre
University of British Columbia
V6T 1Z4
Vancouver, BC, Canada

2. Pacific Rim National Park Reserve
P.O. Box 280
Ucluelet, BC, Canada

3. Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve, and
Haida Heritage Site
60 Second Beach Road
Skidegate, BC, Canada

*Correponding author: keila.stark@zoology.ubc.ca

Abstract: 317 words
Main text: 6101 words
Number of references: 57
Number of figures: 5
Number of tables: 2

Competing interests: We have no competing interests.

Abstract

Dispersal, environmental niche filtering, and biotic interactions all structure patterns of abundance and distribution in ecological communities. Metacommunity ecology provides a framework for understanding how these processes combine to determine community composition in sites across a region that are connected through the dispersal of multiple potentially interacting populations. Previous metacommunity approaches have had limited success due to statistical limitations and incompatibilities between theory and methods. Here, we use the Hierarchical Modelling of Species Communities statistical framework to individually assess the contribution of the aforementioned assembly processes to biodiversity patterns in an important coastal habitat, seagrass meadows, across approximately 1000 km along the coast of British Columbia, Canada. Following the HMSC framework, we parametrized a hierarchical joint species distribution model describing the relationship between multiple measured environmental covariates, spatial distance between sites, and unexplained random effects on epifaunal species distributions across thirteen eelgrass (*Zostera marina* (L.) meadows. Observed species distribution patterns suggest that seagrass epifauna form well-connected metacommunities at the present spatial scale, but that dispersal rates are not so high that they homogenize the region through mass effects. We found evidence of niche filtering in several species along an abiotic gradient, with surface phosphates, temperature and dissolved oxygen being the most important environmental variables. However, environmental conditions only explained a small fraction of the variation in observed species occurrences at the community level, and the predictive power (D^2_{adj}) of the model was poor for some species, suggesting that either some species have broader environmental niches, or niche filtering is not important at this spatial scale. Strong structuring in pairwise species co-occurrences led us to hypothesize that

local community composition may be influenced by priority effects resulting from biotic

interactions. Our approach has yielded refined hypotheses about the specific local and regional

dynamics influencing seagrass epifaunal community structure; these hypotheses in turn can

inform targeted experiments, monitoring efforts, and management decisions for preserving

biodiversity in this important coastal ecosystem.

Key words: Hierarchical Modelling of Species Communities, priority effects, dispersal, niche

filtering, British Columbia, *Zostera marina* (L.)

Introduction

Understanding how local environmental conditions, biotic interactions, and regional

connectivity combine to structure the composition of communities is a central challenge in

ecology (Vellend 2016, Leibold and Chase 2017). Determining the relative importance of these

assembly processes in particular ecosystems is a critical step in understanding the spatial and

temporal scales of stability and resilience in spatially structured communities (Leibold and Chase

2017). Metacommunity theory offers an integrative framework for understanding drivers of

diversity across spatial scales, however previous attempts to quantitatively apply

metacommunity concepts to real-world ecological systems have had limited success (Logue et al.

2011, Brown et al. 2017). Over the past decade and a half, the primary focus of metacommunity

studies has been seeking signatures of one of four idealized metacommunity paradigms in natural

systems: the neutral model, patch-dynamics, species sorting and mass effects (Leibold et al.

2004, Cottenie 2005). An emerging perspective holds that natural metacommunities do not fit

neatly under one of the four paradigms because they do not reflect all possible combinations of

the various assembly processes (Logue et al. 2011, Brown et al. 2017, Leibold and Chase 2017).

Thus, the revised goal is to determine how dispersal, environmental conditions, and biotic

interactions combine to structure species composition and relative abundance within a

metacommunity.

Many marine ecosystems that host high biodiversity occupy spatially structured habitats such

as rocky reefs, coral reefs, and seagrass meadows, and marine species have various dispersal

strategies that connect populations in distinct habitat patches. Marine systems are characterized

by strong local species interactions (Berlow 1999, Sala and Graham 2002) and the importance of

'supply-side' or dispersal-driven population dynamics has long been recognized (Levin and

Paine 1974, Gaines and Roughgarden 1985). These combined roles of dispersal, local

environment, and species interactions suggest that marine biodiversity patterns may in fact

reflect local and regional processes characteristic of metacommunities (Guichard et al. 2003).

Despite the historically strong focus on dispersal and species interactions in marine ecology,

metacommunity dynamics have only been explicitly investigated in a few marine systems

(Guichard et al. 2003, Yamada et al. 2014, Valanko et al. 2015, Whippo et al. 2018).

Consequently, the potential benefits to conservation prioritization that could be gleaned by

applying metacommunity theory to marine systems have not been fully realized (Economo

2011).

Seagrasses, like kelps, corals and salt marsh plants, are foundation species that support high

productivity and faunal diversity (Heck and Thoman 1984, Orth et al. 1984, Duffy et al. 2015).

Seagrasses form meadows separated from each other by deeper water, un-vegetated seafloor, or

other vegetated habitats such as kelp forests. Faunal species disperse among meadows and interact

through predation, competition, and facilitation (Boström et al. 2006). This scenario is consistent

with the type of landscape that supports communities governed by metacommunity dynamics

(Mouquet and Loreau 2003, Leibold et al. 2004). Yet, our understanding of seagrass-associated

faunal diversity and its consequences for conservation and ecosystem function does not yet reflect

the possibility that seagrass biodiversity is maintained by processes operating across numerous

seagrass meadows and adjacent habitats in a seascape. Numerous seagrass community ecology

studies have focused on the influence of local, site-level characteristics such as habitat structure

(Orth et al. 1984), primary biomass (Cébrian and Duarte 1998, Gullström et al. 2012), or trophic

interactions (Amunudrud et al. 2015, Huang et al. 2015) on faunal diversity. Few studies have

elucidated ecological processes occurring operating on a regional or inter-site scale. Failure to

identify the contribution of metacommunity processes could lead to underestimates of the amount

of habitat required to sustain biodiversity and ecosystem functions (Boström et al 2006).

Two studies that applied different metacommunity approaches to seagrass invertebrate

communities found evidence for the importance of regional-scale processes (Yamada et al. 2014,

Whippo et al. 2018), but their ability to characterize the individual contribution of all assembly

processes driving metacommunity structure was limited by the statistical methods available at

the time. Furthermore, neither approach was able to identify possible biotic interactions, thereby

entirely missing a critical aspect of community assembly. Using variation partitioning on

constrained ordinations (Borcard et al. 1992, Cottenie 2005), Yamada et al. (2014) found that the

relative contribution of space and environment varied among different invertebrate functional

groups, suggesting that the relative importance dispersal levels versus niche filtering could be

mediated by functional traits. However, the variation partitioning analysis was not able to

distinguish between multiple possible metacommunity dynamics at play owing to the ambiguous

statistical significance of the results (Yamada et al. 2014). Simulation studies have shown that

this variation partitioning method can over- or under-estimate spatial and environmental

components of variation, and produce inflated R^2 values depending on the researcher's statistical

choices (Gilbert and Bennett 2010, Tuomisto et al. 2012). Whippo et al. (2018) used another

method, elements of metacommunity structure (EMS) (Leibold and Mikkelson 2002, Presley et

al. 2010). While useful for visualizing regional diversity patterns, EMS does not elucidate the

potential processes contributing to these patterns (Brown et al. 2017).

These limitations, along with the revised goal of individually assessing the contribution of

dispersal, niche filtering, and biotic interactions to diversity in a metacommunity, invite novel

methodological approaches. A new statistical framework, Hierarchical Modelling of Species

Communities (HMSC) (Ovaskainen et al. 2017), shows promise for achieving this goal. HMSC

uses Bayesian inference to parametrize a hierarchical joint species distribution model

summarizing the influence of space, environment, patterns of co-occurrence, phylogeny and

traits on species distributions. The framework allows the partitioning of explained variation due

to environmental covariates, spatial distance, and random effects for all species distributions,

allowing the observer to discern patterns at the whole community level as well as in individuals.

Additionally, the framework models the residual variance in species distributions unexplained by

environment and space as a variance-covariance matrix, which can be used to visualize patterns

of positive or negative pairwise species co-occurences. For these reasons, HMSC achieves what

researchers aimed to achieve with the previous methods, and much more.

Here we test the hypothesis that seagrass-associated epifaunal invertebrate communities are

jointly structured by the spatial distances (a proxy for inferring dispersal dynamics) between

meadows, local environmental conditions, and biotic interactions. Our approach moves away

from the four original metacommunity paradigms, instead focusing on individually

characterizing the effects of dispersal, environmental niche filtering, and biotic interactions on

the community structure. We follow the HMSC framework (Ovaskainen et al. 2017) to answer

the following questions: (1) Is the metacommunity characterized by dispersal limitation,

intermediate dispersal, or high dispersal? (2) Do species distribution patterns suggest

environmental niche filtering, and if so along which environmental axes? (3) Do species co-

variance patterns suggest possible biotic interactions influencing community assembly? The

order in which these questions are posed matters. For example, if dispersal limitation is found to

be significant, this knowledge affects how we may interpret observed species co-occurrence

patterns. In answering these questions with a biodiversity survey of invertebrates in 13 meadows

in British Columbia, we also provide a comprehensive assessment of species abundance and

diversity from local to regional scales. We find evidence of intermediate dispersal rates among

meadows across 1000 km, but no sign of complete biodiversity homogenization resulting from

high dispersal. We also find that local environmental conditions have some influence on species

distributions across the 13 sites, but that individual species' responses to environmental gradients

vary dramatically. Strong species covariance patterns led us to hypothesize possible biotic

interactions influencing local community composition.

Methods

Study sites

We sampled epifaunal biodiversity in 13 eelgrass *Zostera marina* (L.) meadows across a

spatial scale of 1000 km from late June through early August in 2017. The sites were located

within four coastal areas: Haida Gwaii within Gwaii Haanas National Park Reserve, National

Marine Marine Conservation Area Reserve, and Haida Heritage Site; southern Clayoquot Sound

(one site was within Pacific Rim National Park Reserve), Barkley Sound on the West coast of

Vancouver Island, and the Southern Gulf Islands on the south-eastern tip of Vancouver Island

within Gulf Islands National Park Reserve (Fig. 1).

Zostera marina (L.) is the most common subtidal seagrass species in the Northeast Pacific

(Alaska to southern California). In British Columbia, Canada, *Z. marina* meadows are found in

discrete patches along the coast. They vary in several biotic and abiotic conditions, such as

seagrass shoot surface area and density, macroalgal species composition, fish community

structure, temperature, salinity, and water column nutrient levels.

Field sampling of eelgrass and associated epifauna

We sampled three sites in Barkley Sound (SA, DC, RB) with SCUBA because they

required boat access, and the deepest parts of the meadow reached a depth of 5m during

sampling. All other sites were sampled by wading or snorkeling at a low tide of 0.6 m or less.

We used a standardized, nested sampling regime with 0.25 m x 0.25 m quadrats to collect

aboveground eelgrass, epifaunal invertebrates, detritus, and macroalgae (Whippo et al. 2018).

Six quadrats were set in a 15 m x 30 m array, such that every quadrat was situated 15 metres

from its nearest neighbour (Appendix S1: Fig. 1). The array was placed in the middle of the

eelgrass meadow to avoid edge effects where possible. We removed all above-ground biomass

(eelgrass, detritus, macroalgae) within each quadrat by hand and immediately placed the contents

into a 250 μm mesh bag at the Barkley Sound and Southern Gulf Islands (GB, LH, SS, CB, JB)

sites, and into a Ziploc bag at the Clayoquot Sound (IN, EB, DK) and Haida Gwaii (HL, RA)

sites.

Acquiring biotic and abiotic environmental variables

To test our hypothesis on environmental niche filtering, we quantified 14 environmental

attributes of each meadow and quadrat sampled. We used in-situ field measurements and

satellite-derived averages from the Bio-ORACLE dataset (Assis et al. 2018) as estimates of

water quality conditions associated with hypothesized niche differences among epifauna. We

gathered in-situ measurements of water quality parameters (pH, salinity, dissolved oxygen,

nitrates and chlorophyll-*a*) at each meadow. We also acquired data on mean surface chlorophyll-

a ($\mu g\ L^{-1}$), dissolved oxygen, surface nitrates, surface silicates, surface phosphates, pH, salinity,

and mean sea surface temperature (°C) from Bio-ORACLE: a global database of marine

ecological variables gathered from several satellite and in-situ sources at a spatial resolution of

0.08° (Assis et al. 2018). All sites (meadows) were situated in different spatial cells, therefore all

sites had distinct values for the aforementions parameters. However, all quadrats within a given

meadow (spaced 5-15m from its adjacent neighbor) were situated in the same spatial cell and

thus were given the same site-level value for all the Bio-ORACLE variables; we accounted for

this spatially nested random effect while constructing our model. We also quantified fetch, which

is a proxy for site exposure. We chose these abiotic variables because we expected them to be the

most important variables influencing species distributions, either through variation in

physiological tolerances, or through influencing primary productivity and food quality.

For each sample, we measured five biotic attributes: eelgrass leaf area index (LAI), shoot

density, eelgrass dry mass, total algal dry mass, and detritus dry mass. We measured leaf length,

width, and number of blades in five representative shoots within each quadrat. Leaf Area Index

(m^2 eelgrass blade area per m^2 of ground) was calculated by multiplying the average blade area

(m^2) per shoot by the shoot density (number of shoots per m^2). We standardized blade length

measurements from the end of the blade to just above the first rhizome node because some

shoots had intact rhizomes while others were missing after uprooting in the field. We separated

and dried eelgrass, detritus, and macroalgae in a desiccator oven (60° C for 48 hours) to measure

dry mass.

Response variable: Quantifying epifaunal diversity

We rinsed the eelgrass blades from each quadrat to remove epifaunal invertebrates, and

passed samples through a 500 μm sieve. Invertebrates were transferred into a centrifuge tube and

preserved in 95% EtOH for identification with light microscopy (10 times magnification). We

identified invertebrates to the lowest taxonomic level possible using keys in the Light and Smith

Manual (Carlton 2007) and Kozloff (1996).

Analysis

We followed Ovaskainen et al. (2017) to construct a hierarchical joint species distribution

modelling the relationship between measured environmental variables, spatial distance, species

co-occurrences and species distributions using the HMSC package (Guillaume Blanchet et al.

2018) in R (version 3.4.2; R Development Core Team 2014). The model comprises two linear

predictor terms: one describing fixed effects of environment (9 water quality and 5 biotic

variables) and space, and one describing random effects that cannot be attributed to environment

or space (Equations 2 & 3 in Ovaskainen et al. 2017). The random-effects linear predictor term

describes random variation in individual species distributions as well as covariation for every

species pair in the analysis.

We used Markov chain Monte Carlo to estimate the model parameters. Species'

responses to environmental variables were assumed to follow a multivariate Gaussian

distribution. We assumed a Poisson likelihood distribution for species count data and used the

same priors as in the examples presented in Ovaskainen et al. (2017). The Markov Chain

included 100 000 iterations, a burn-in length of 1000 iterations, and thinning of 100. We

confirmed that parameters were well mixed by visually inspecting trace plots. We tested the

model's performance by calculating explained deviance (D^2_{adj}) as described in Appendix S1: Eq.

S1. D^2_{adj} is comparable to R^2_{adj} in least square regression analysis (Guisan and Zimmermann

2000); in this context, it describes the model fit where a D^2_{adj} value of 1 means the model

perfectly predicts the observed data. This yielded a D^2_{adj} value for each individual species

distribution model, and the mean explained deviance across all species gave a single

metacommunity-level value of the model's predictive power.

To answer our three main questions about the relative roles of dispersal, environment,

and species interactions, we visualized the parametrized model in several ways. We investigated

three possible dispersal conditions: dispersal limitation, intermediate dispersal, and dispersal

surplus. To investigate dispersal limitation at large spatial scales, we plotted distance-decay of

community similarity. To do so we randomly drew 200 pairwise distances between sampling

plots, and plotted predicted community similarity against distance. Distance-decay of community

similarity could reflect dispersal limitation, or it could reflect differences in environmental

conditions with increasing spatial scale. To distinguish these two possibilities, we compared

predicted species compositional similarity as a function of both distance and environmental

covariates (the full HSMC model), and predicted compositional similarity using an alternative

model in which the effect of the environment was removed (all environmental covariates were

set to their mean value across all sites; Equation 5 in Ovaskainen et al. 2017). To distinguish

between dispersal sufficiency (intermediate) and dispersal surplus (high), we looked for evidence

of biodiversity homogenization across sites- a signature of swamping by high dispersal- through

visual inspection of our site-by-species matrix. We used estimates of environmental niche

filtering strength (as measured by the proportion of deviance explained) as another line of

evidence to distinguish intermediate and high dispersal: theory predicts that intermediate

dispersal allows species to settle at sites where local environmental conditions are optimal for

growth and reproduction, whereas high dispersal overrides this (Mouquet and Loreau 2003).

To estimate the strength of environmental niche filtering, we conducted variation

partitioning analysis of the fixed and random effects as defined in Ovaskainen et al. (2017) in the

HMSC R package (Guillaume Blanchet et al. 2018). This summarized the relative contribution

of every abiotic and biotic environmental variable, spatial distance, and random effects to the

distribution of each species in the HMSC model; we visualized this in a stacked bar plot. We

then estimated the fractions of variation explained by environment only, space only, the shared

variation between space and environment, and the residual or unexplained variation in a way

similar to Borcard et al. (1992), but using the HMSC model as opposed to constrained

ordination. To do so, we estimated three versions of the HMSC model: 1) the original global

model which included environmental and spatial variables, site- and region-level random effects,

2) environmental variables only with the random effects, and 3) spatial distances only with

random effects. From each model, we extracted the total variation explained by environment and

space, the variation explained by environment, and the variation explained by space. To

determine the shared fraction of variation between environment and space, we multiplied each

whole fraction by the calculated community-level D^2_{adj} to reflect the model deviance in our

variance partitioning estimates, and followed the equations in Appendix S1: Eq. 2. A Venn

diagram illustrating all fractions is shown in the Appendix S1: Fig. 3.

To identify species co-occurrence patterns possibly resulting from biotic interactions, we

represented modelled pairwise associations in a correlation matrix describing the extent to which

species pairs co-occurred more negatively or positively than by random chance, controlling for

the effect of environmental covariates. We included only the twenty most abundant species in the

main figure, but provide a full species co-occurrence matrix in Appendix S1: Fig. S4. We then

conducted hierarchical cluster analysis on sites according to predicted species proportional

abundances from the HMSC model using Ward's criterion (Ward 1963). We mapped this

dendrogram onto an abundance heatmap of the 20 most abundant species, and colour coded the

cells according to the species co-occurrence groupings (groups of species that tended to

positively co-occur amongst themselves). A version of this figure with all species is provided in

Appendix S1: Fig. S5.

Results

Species abundance and distribution patterns

The aim of this section is to describe basic biodiversity patterns across the seagrass

meadows. Specific results aimed at answering our three questions about metacommunity

processes follow in the subsequent sections. Seagrass-associated biodiversity varied spatially,

both within meadows and among meadows. We counted and identified 39 644 individuals

representing at least 58 species across all sites (Table 2). The metacommunity represented

multiple phyla, diet types (herbivores, detritivores, suspension feeders, and carnivores), and life

history strategies (brooding, broadcast spawning). A full summary of traditional diversity metrics

can be found in Appendix S1: Table S1.

Several taxa were present at all sites (Appendix S1: Fig. S2). These include the

harpacticoid copepod *Porcellidium* sp., bivalves *Saxidomus gigantea* and *Clinocardium nutallii*,

snails *Alvania compacta*, *Lacuna variegata* and *Lacuna vincta*, and the tanaid *Leptochelia* sp.

Genera *Mytilus* (a mussel) and *Nereis* (a polychaete worm) were also present at all sites, however

the number of species represented remains unknown because we were unable to confidently

identify these organisms to species. Ten species were present in at least one site in every region:

the pycnogonid *Anoplodactylus viridintestinalis,* isopod *Idotea resecata,* gammarid amphipods

Monocorophium insidiosum, Ampithoe valida, Pontogeneia rostrata, Ampithoe dalli, Aoroides

spp.*,* snails *Amphissa columbiana* and *Alia carinata,* and limpet *Lottia pelta*.

Dispersal: Distance-decay of community similarity due to spatial distance and environment

In the full HMSC model with latent variables summarizing species' responses to

environment and distance between sites, predicted community similarity decreased with

increasing distance between meadows (Fig. 2, red points and line). However, this decrease in

community similarity with spatial distance disappeared, and community similarity showed a

slight increase with distance when the effect of environmental variables was removed from the

model (all environmental variables set to their mean value), suggesting that spatial distance alone

did not correspond with differences in invertebrate compositional similarity (Fig. 2, blue points

and line). The range of predicted similarities was greater at small distances than larger distances,

indicating that spatial proximity between sites does not necessarily confer compositional

similarity resulting from the exchange of individuals.

Environmental niche filtering: variation partitioning

Patterns in abundance and distribution are consistent with weak environmental niche

filtering along an abiotic axis, however the importance of environmental conditions varies across

species. The relative contribution of all 14 environmental variables, spatial distance and random

effects in the HMSC model can be seen in Fig. 3. Out of all the measured environmental

variables, the water quality variables (all blue bars in Fig. 3) explained the largest proportion of

variation across nearly all species distributions Across all taxa, herbivore food availability

(seagrass, algae and detritus biomass) and habitat structure (eelgrass shoot density and Leaf Area

Index) had smaller influences on species distributions than abiotic predictors (Fig. 3). The

relative contribution of abiotic parameters to species abundance and distribution patterns also

varied among species and major taxonomic groups. For instance, site and region membership,

food availability and habitat structure generally explained less variation in gastropod

distributions than gammarid amphipod distributions. While the mean deviance explained (D^2_{adj})

across all species distribution models was 0.29, individual values varied dramatically (0 to 0.82,

Table 2), implying high variation in the match between our model and the observed data.

Accounting for this, the true fraction of variation explained by environment was 22.4%, space

explained -3%, the shared fraction between environment and space was 4%, and the residual

fraction of variation was 0.76 (Appendix S1: Fig. S3).

Species interactions: Patterns in pairwise species co-occurrences

We identified, post hoc, three main site-level species co-occurrence groupings across all

78 samples from 13 meadows and 4 regions after controlling for effects of environmental

covariates. The six species in the first group (*Nereis* sp. through *Caprella laeviuscula* in Fig. 4a)

henceforth known as the purple group, tended to positively co-occur with each other, and

negatively co-occur with non-members of this group. The second group- henceforth known as

the turquoise group- included *Caprella californica* through *Ampithoe valida,* and Harpacticoid 2

through *Lacuna* sp. *Spirorbis* sp. – which had an unusually high abundance of approximately 16

000 individuals at the HL site- did not fall into either category. *Alvania compacta* and *Lacuna* sp.

negatively co-occurred with members of the purple group and positively co-occurred with

members of the turquoise group, however their co-occurrence values tended to be weaker.

Species interactions: Mapping co-occurrence groupings onto a site by species matrix

Species associated with all three major co-occurrence groupings were present at all sites

(Figure 4b), indicating that presences of these species are not mutually exclusive. It is their

abundances that tended to negatively co-vary; when species of one assemblage were abundant,

species of the other assemblages were less abundant (Figure 4b). Fig. 4b shows clustering of

sites according to the identity of the most abundant species by co-occurrence colour grouping. At

JB, HL, CB, RA, EB, DC, and DK, members of the blue group tended to be most abundant,

whereas purple group members were most abundant at SS, SA, LH, GB and RB. IN was an

outlier site, as *Spirorbis* sp. was predicted to vastly outnumber other species. The emergent

species assemblages identified in Fig. 4a from the HMSC modeled distributions do not clearly

correspond with the spatial distance because the cluster analysis did not group sites by spatial

proximity or regional membership (black symbols in Fig. 4b).

Discussion

By applying a new statistical framework, we have gained new insight into the

metacommunity processes structuring seagrass-associated biodiversity in coastal British

Columbia. We report evidence suggesting that both biotic interactions and environmental

conditions have an influence on seagrass meadow invertebrate composition. Spatial patterns of

species abundance and distribution suggest regional-scale dispersal patterns such that even

distant meadows may host the same species. However, these dispersal rates are clearly not so

high as to overwhelm the local meadow scale effects of biotic interactions and environmental

conditions. Below, we review these conclusions and the evidence to support them.

Seagrass animal communities are well-connected and may experience intermediate dispersal

rates

Seagrass epifaunal communities appear to be well-connected by dispersal at the present

spatial scale and distribution of seagrass habitat along the British Columbia coast. This is true for

the seven species that were present at all sites, and the additional ten species that were not

present at every site but present in all regions (Appendix S1: Fig. S2). It is noteworthy that these

groups had representation from across phyla and life history traits (polychaete worms, gammarid

amphipods, snails, bivalves, tanaids), suggesting that no single taxonomic group or dispersal life

history strategy had a consistently greater spatial distribution among the sites sampled. However,

the specific pathway the organisms travelled to arrive at these sites remains unknown. We used

Euclidan distances in our analysis, which preserved the rank order of distances across sites, but

actual distances are likely greater due to oceanographic circulation patterns (Kinlan and Gaines

2003, Mitarai et al. 2008, Treml et al. 2015). It is unclear whether distant sites share the same

species because they are (1) linked by direct dispersal; (2) indirectly linked by dispersal via

unsampled "stepping-stone" sites; or (3) were colonized by populations of the same species in a

historical dispersal event, but have not seen the exchange of individuals since. Incidences of

direct dispersal between the farthest sites (ie. between Haida Gwaii and the Southern Gulf

Islands) may be rare, but are not impossible. Seagrass-associated epifauna are generally passive

dispersers, with individuals colonizing neighbouring sites either by larval transport in currents or

by "rafting" on floating pieces of seagrass, detritus, or macroalgae. Kelp rafts have been

observed hosting benthic and epifaunal invertebrates across phyla (echinoderms; peracarids;

molluscs; annelids), and are important in maintaining connectivity between coastal ecosystems

across 100's of kilometres (Wichmann et al. 2012). Thus, our results suggest that this passive

dispersal is sufficient to facilitate connectivity of these sites across the region so that species

distributions are not limited by dispersal through isolation by distance.

Species in our analysis that were only found at one site or region may be dispersal limited

for reasons other than isolation by distance. Those with low site-level abundances (fewer than

five individuals detected across all plots from a site) may not disperse in appreciable numbers,

and thus are not able to establish populations at all sites as easily as more abundant species. This

group included gammarid amphipods *Jassa marmorata*, *Orchomenella recondita,* polychaete

worms *Glycinde armigera, Neoamphitrite robusta, Lepidonotus squamatus,* and others

(Appendix S1: Fig. S2).

We can also conclude that dispersal rates are high enough for several species to be

present at most sites, but not so high as to completely overwhelm the effects of environmental

differences at the local scale. Thus, dispersal rates are likely intermediate. Co-occurrence

patterns indicate that a subset of species was present at all sites, but often at varying abundances.

This may be indicative of weak mass effects, which allow these populations to persist even if

local conditions are not sufficient for them to persist without immigration (Mouquet and Loreau

2003). Metacommunity theory predicts that such weak mass effects are likely to be found

whenever dispersal rates are not limiting (Thompson et al. 2017), particularly in organisms that

cannot control their own dispersal (Leibold and Chase 2017), such as seagrass-associated

epifauna. This may be supported by our result on the contribution of environmental conditions,

particularly in ubiquitous species for whom the model fit was poor (e.g.'s *Aoroides* spp.,

Spirorbis sp., D^2_{adj} = 0); the poor model fit may indicate that they may persist sites even if

environmental conditions are not optimal, due to sufficient immigration. An alternative

explanation is that species distribution models with low D^2_{adj} values indicate species with broader

environmental niches, or their abundances along environmental gradients do not follow a

Gaussian distribution as assumed in the model. As a result, the HMSC model could not

accurately predict their occurrences given local environmental conditions. It is difficult to

discern between these possibilities without empirical tests of species niches.

Seagrass epifauna show varying degrees of environmental niche filtering

The fact that environmental covariates across space- and not space alone- are associated

with differences in community composition may suggest some influence of environmental niche

filtering (Fig. 2). The most important environmental axis informing species distributions is the

abiotic one, with phosphates, salinity, and dissolved oxygen being the top three parameters by

percent variation explained (Fig. 4a). However, the actual amount of variation explained by the

environment was small (Appendix S1: Fig. 3), leaving a great deal of unexplained variation in

the observed data at the whole-community level. However, in some species for whom our model

had high predictive power (including *Ampithoe dalli, Pontogeneia rostrata, Caprella natalensis)*,

environmental niche filtering may be important.

Species may respond to environmental heterogeneity across the metacommunity in one of

four ways. First, the most important abiotic environmental parameters (nitrates, phosphates, sea

surface temperature, and dissolved oxygen) may indirectly influence epifaunal presence and

abundance via primary productivity levels. This is possible despite the fact that primary biomass

explained less variance than the abiotic parameters in our HMSC model (Figure 3). Our measure

of primary biomass was based on a measurement at a single time point. But, our abiotic

parameters represent a four-year average (Assis et al. 2018) and so may provide a better

representation of what the organisms experience over their lifetimes. Second, sea surface

temperature, dissolved oxygen, and salinity may also directly influence optimal tolerance ranges.

For example, estuarine isopod *Idotea baltica* (a relative of *I. resecata* and *I. montereyensis* in our

study) experiences significantly lower survival in prolonged exposure to higher water

temperatures and lower salinities than those experienced in their home habitat (Rugiu et al.

2018). Third, temperature may influence the metabolic demands of the epifauna (most of which

are grazers such as some gammarid amiphopids, *Idotea* spp., *Lacuna* spp.), thereby influencing

grazing rates and altering the available primary biomass (O'Connor 2009). Decreased food

resources may result in competition, thereby influencing the number of individuals, number of

species, or identity of species a seagrass patch can host. It is not easy to determine whether

individual species' responses to competitive stress can influence whole-community structure,

however doing so may provide key information on how environmental conditions and

interspecific interactions interact to influence biodiversity.

Environmental characteristics of temperate eelgrass meadows fluctuate with the seasons,

likely driving changes in the relative abundance of epifaunal taxa throughout the year

(Wlodarska-Kowalczuk et al. 2014, Markel et al. 2017). As a result, the epifaunal

metacommunity may not evince these abundances or distributions year-round; sampling several

times in one year to capture temporal changes in community structure would give a clearer

picture of species' responses to environmental conditions, as well as biotic interactions and

dispersal events.

Do species co-variance patterns reveal possible biotic processes influencing community

assembly?

After controlling for effects of space and environment, two main co-occurrence

groupings emerged showing antagonistic abundances across sites rather than mutually exclusive

presences (Fig. 4b). Stochasticity alone is unlikely to have produced such a strong pattern. While

there may be multiple possible explanations for non-random observations in species occurrences,

we were able to rule out environmental heterogeneity (Connor and Simberloff 1979) as the co-

occurrence values represented pairwise species covariation unexplained by the spatial and

environmental covariates. This along with the high residual variation in our variation partitioning

result (Appendix S1: Fig. 3) leads us to suggest the potential importance of biotic processes in

structuring the epifaunal metacommunity. Several types of interactions may be occurring in the

metacommunity; the majority of species in the present study are known to be herbivores or

detritivores, and thus may compete for primary biomass. Laboratory experiments have

demonstrated that behaviours such as grazing rate and habitat selection in the marine amphipods

are altered in the presence of morphologically and functionally similar intrerspecific competitors

(Beermann et al. 2018)*,* however it is difficult to empirically test limited food resources result in

competition in natural systems. Other species in our study are filter feeders, and thus may not be

competing for food resources but possibly space.

The community structure of a given site with a certain regional pool and local

environmental conditions may have multiple stable states depending on the arrival order of

species and the structure of competitive interactions; this phenomenon is known as priority

effects (Waters et al. 2013, Fukami et al. 2016, Ke and Letten 2018) and has been documented in

marine fouling communities (Vieira et al. 2018). The fact that we found antagonistic co-

occurrence groupings that did not result from space or environment may suggest priority effects

(Fig. 4a and 4b). We hypothesize that following the establishment of a new meadow or a

disturbance event, the first few species to colonize a seagrass meadow greatly determine the

success of subsequent colonizers, regardless of the environmental conditions in the meadow. The

high abundance and negative co-occurrence between *Nereis* sp. and *Lacuna* sp. suggests that

these species may be key determinants of community structure (Fig. 4b). As an illustrative

example, *Nereis* sp. might have been among the first to colonize sites SS, SA, LH, and RB (Fig.

4b), thereby gaining a numerical advantage over *Lacuna* sp. and others. Distibution patterns in

the sessile calcifying polychaete *Spirorbis* sp. may also be indicative of priority effects. While it

did not appear to strongly negatively or positively co-occur with other species approximately, it

dominated other species in abundance at the HL sites. 16 300 individuals were found across the

six quadrats, several orders of magnitude higher than its abundance at other sites (Data S1). HL

also had the lowest species richness of all 13 sites. Eelgrass shoots there were completely

covered by these *Spirorbis* sp., possibly preventing grazers in the community from accessing

epiphytic algae or the seagrass itself. The age, colonization history, and disturbance regime of

the meadows themselves likely matter if priority effects are an important influence on

community assembly. A key piece of information that would confirm the nature of biotic

processes is whether species from different co-occurrence groupings compete more strongly with

each other than with themselves (Ke and Letten 2018), but this would require an experimental

test.

The positive co-occurrence patterns between certain species may be a result of positive

interspecific interactions. An example is *Orchomenella recondita*: a gammarid amphipod found

living in the gastrovascular cavity of the anemone *Anthopleura elegans* (Carlton 2007). This

species was only recorded at the SS site, and specifically only found in quadrats where *A.*

elegans was collected with the seagrass shoots.

A potentially important biotic interaction missed in our analysis is the trophic effect of

predators. Field experiments have demonstrated that changes in predation pressure by fish,

shorebirds, and predaceous invertebrates can shift seagrass-associated epifaunal assemblages in a

matter of weeks (Amundrud et al. 2015; Huang et al. 2015; Lewis & Anderson 2016).

Differential predation pressure or composition of predators could also lead to the pattern of

diverging community composition that we have witnessed (Guzman et al. 2018). Further study is

required to determine the degree to which these seagrasses associated communities are structured

by trophic interactions.

Conclusion

Using a new statistical approach to characterizing dispersal, environmental niche

filtering, and biotic interactions, we have gained greater insight into the local and regional

metacommunity processes driving temperate seagrass epifaunal diversity. Our study also gives a

baseline spatial scale (1000 km) for comparing the contribution of these assembly processes

across spatial scales and marine systems (Viana and Chase, 2018). We have generated new

hypotheses about the relative importance of dispersal, environmental heterogeneity, and biotic

interactions in shaping these communities- manipulative experiments and ongoing monitoring

are now needed test and refine these hypotheses. We could test our hypothesis that species

sorting is relatively weak by measuring fitness differences along the most important

environmental axes in key species identified in this study, or constructing an experiment that

allows individuals to disperse to sites where we expect them to be optimally adapted to local

abiotic conditions. Given our hypothesis that priority effects play a potential role in structuring

these communities, experiments are now required to test whether the composition of seagrass

meadows depends on the order of species colonization (Fukami et al. 2016). Testing our

hypothesis that these communities are not dispersal limited is more challenging, but could be

studied using particle tracking models similar to those used to estimate connectivity between

coral reefs via larval dispersal across thousands of kilometres (Treml et al. 2008). A population

genetic analysis could also ground-truth the extent and time scale of dispersal by determining

whether populations share alleles, or have been isolated for some time. All together, we suggest

that a modern metacommunity perspective combined with new advances in our statistical

toolbox provide a powerful and relevant framework for studying natural ecosystems.

Our findings are particularly relevant to conservation initiatives that aim to maintain

diversity or prevent species loss in seagrass meadows. Specifically, identifying the contribution

of dispersal, environmental conditions, and interspecific interactions allows the conservation

practitioner to better predict how habitat fragmentation and climate change might influence

diversity. Epifaunal invertebrate communities in British Columbia seagrass meadows may be

quite resilient, given our findings that environmental conditions make a minimal contribution to

species distributions, and several species across phyla and functional traits are consistently

present and abundant across a fairly large spatial scale.

Acknowledgements

KAS conceived the study with guidance from MIO. JY provided samples from IN, DK, and EB sites. LL provided samples from RA and HL sites. KAS collected samples from GB, JB, CB, LH, SS, RB, SA, & DC sites, processed samples, analyzed data, and wrote the first draft of the manuscript. PLT provided substantial recommendations for the analysis. KAS, PLT, and MIO contributed to writing the manuscript. This research was financially supported by a UBC Zoology SURE Grant to KAS, and funding from NSERC Discovery Grants, Canadian Foundation for Innovation (CFI). PLT is supported by NSERC and Killam postdoctoral fellowships. This research is sponsored by the NSERC Canadian Healthy Oceans Network and its Partners: Department of Fisheries and Oceans Canada and INREST (representing the Port of Sept-Îles and City of Sept-Îles). Matt Whalen pulled the Bio-ORACLE dataset. Special thanks to Michelle Paleczny and Gulf Islands National Park Reserve for boat and laboratory space during their 2017 eelgrass sampling campaign. We acknowledge staff, students, and volunteers from Pacific Rim National Park Reserve for their assistance collecting eelgrass samples in Clayoquot Sound. We also sincerely thank Rob Underhill, Abbie Sherwood, Emily Adamczyk, Minako Ito, Ariane Comeau, Giordano Bua for support and assistance with collection, processing, and identification of epifaunal samples. We are grateful to Duarte Viana for providing critical recommendations and code for the D^2_{adj} and variation partitioning calculations. We also thank Coreen Forbes and Matt Whalen for their comments on the manuscript drafts.

Literature Cited

Amundrud, S.L., D. S. Srivastava, and M. I. O'Connor. 2015. Indirect effects of predators control herbivore richness and abundance in a benthic eelgrass (*Zostera marina*)

mesograzer community. Journal of Animal Ecology 84:1092–1102.

Assis J., L. Tyberghein L. S. Bosch, H. Verbruggen, E. A. Serrão, and O. De Clerck. 2018. Bio-

ORACLE v2.0: Extending marine data layers for bioclimatic modelling. Global Ecology

and Biogeography 27:277–284.

Beermann, J., K. Boos, L. Gutow, M. Boersma, and A. C. Peralta. 2018. Combined effects of

predator cues and competition define habitat choice and food consumption of amphipod

mesograzers. Oecologia 186:645–654.

Berlow, E. L. 1999. Strong effects of weak interactions in ecological communities. Nature 398:

330–334.

Boström, C., E. L. Jackson, and C. A. Simenstad. 2006. Seagrass landscapes and their effects on

associated fauna: A review. Estuarine, Coastal and Shelf Science 68:383-403.

Borcard, D., P. Legendre, and P. Drapeau. 1992. Partialling out the Spatial Component of

Ecological Variation. Ecology. 73:1045–1055.

Brown, B.L., E. R. Sokol, J. Skelton, and B. Tornwall. 2017. Making sense of metacommunities:

dispelling the mythology of a metacommunity typology. Oecologia 183:643–652.

Carlton, J. 2007. The Light and Smith Manual: Intertidal invertebrates from Southern California

to Oregon. Fourth edition. University of California Press, Oakland, California, USA.

Cébrian, J. and C. M. Duarte. 1998. Pattems in leaf herbivory on seagrasses. Aquatic Botany

60:67–82.

Connor, E. F., and D. Simberloff. 1979. The assembly of species communities: Chance or

competition? Ecology 60:1132–1140.

Cottenie, K. 2005. Integrating environmental and spatial processes in ecological community

dynamics. Ecology Letters 8:1175–1182.

Duffy, J.E. 2006. Biodiversity and the functioning of seagrass ecosystems. Marine Ecology

Progress Series 311:233–250.

Duffy J.E, et al. 2015. Biodiversity mediates top-down control in eelgrass ecosystems: A global

comparative-experimental approach. Ecology Letters 18:696–705.

Economo, E. P. 2011. Biodiversity Conservation in Metacommunity Networks: Linking Pattern

and Persistence. The American Naturalist 177:167-180.

Fukami, T., E. A. Mordecai, and A. Ostling. 2016. A framework for priority effects. Journal of

Vegetation Science 27:655–657.

Gaines, S., and J. Roughgarden. 1985. Larval settlement rate: A leading determinant of structure

in an ecological community of the marine intertidal zone. Proceedings of the National

Academy of Sciences. 82:3707–3711.

Gilbert, B, and J. R. Bennett. 2010. Partitioning variation in ecological communities: do the

numbers add up? Journal of Applied Ecology 47: 1071-1082.

Guichard, F. 2003. Mussel disturbance dynamics: Signatures of oceanographic forcing from

local interactions. American Naturalist 161:889–904.

Guillaume Blanchet, F., with contributions from G.Tikhonov and A. Norberg. 2018. HMSC:

Hierarchical Modelling of Species Community. R package version. 2.1-2.

Guisan, A., and N. E. Zimmermann. 2000. Predictive habitat distribution models in ecology.

Ecological Modelling 145:147–186.

Gullström, M., S. Baden, and M. Lindegarth. 2012. Spatial patterns and environmental correlates

in leaf-associated epifaunal assemblages of temperate seagrass (*Zostera marina*) meadows.

Marine Biology 159:413–425.

Guzman, L. M., R. M. Germain, C. Forbes, S. Straus, M. I. O'Connor, D. Gravel, D. S. Srivastava,

and P. L. Thompson. 2018. Towards a multi-trophic extension of metacommunity ecology.

In press. Ecology Letters doi: 10.1111/ele.13162.

Heck K.L., and J. F. Valentine. 2006. Plant-herbivore interactions in seagrass meadows. Journal

of Experimental Marine Biology 330:420–436.

Huang A.C., M. Essak, and M. I. O'Connor. 2015. Top-down control by great blue herons *Ardea*

herodias regulates seagrass-associated epifauna. Oikos 124:1492–1501.

Jaschinski, S., and U. Sommer. 2008. Functional diversity of mesograzers in an eelgrass-

epiphyte system. Marine Biology 154:475–482.

Ke, P. J., and A. D. Letten. 2018. Coexistence theory and the frequency-dependence of priority

effects. Nature Ecology and Evolution 2:1691–1695.

Kinlan, B.P., and S. D. Gaines. 2003. Propagule Dispersal in Marine and Terrestrial

Environments: A Community Perspective. Ecology 84:2007–2020.

Kozloff, E. 1996. Marine Invertebrates of the Pacific Northwest. University of Washington

Press, Seattle, Washington, USA.

Leibold, M.A., et al. 2004. The metacommunity concept: A framework for multi-scale

community ecology. Ecology Letters 7:601–613.

Leibold, M.A., and G. M. Mikkelson. 2002. Species Turnover , and Boundary Clumping:

Elements of Meta-Community Structure. Oikos 97:237–250.

Levin, S.A., and R. T. Paine. 1974. Disturbance, patch formation, and community

structure. Proceedings of the National Academy of Sciences. 71:2744–2747.

Lewis, L.S., and T. W. Anderson. 2016. Top-down control of epifauna by fishes enhances

seagrass production. Ecology 93:2746–2757.

Loreau, M., N. Mouquet, and A. Gonzalez. 2003. Biodiversity as spatial insurance in

heterogeneous landscapes. Proceedings of the National Academy of Sciences 28:12765–

12770.

Markel, R. W., K. E. Lotterhos, and C. K. Robinson. 2017. Temporal variability in the

environmental and geographic predictors of spatial-recruitment in nearshore rockfishes.

Marine Ecology Progress Series 574:97–111.

Mitarai, S., D. A. Siegel, and K. B. Winters. 2008. A numerical study of stochastic larval

settlement in the California Current system. Journal of Marine Systems 69:295–309.

Mouquet, N., and M. Loreau. 2003. Community patterns in source-sink metacommunities.

American Naturalist 162:554–557.

O'Connor, M.I. 2009. Warming Strengthens an Herbivore- Plant Interaction. Ecology 90:388–

398.

Orth, R. J., K. L. Heck, and J. van Montfrans. 1984. Faunal Communities in Seagrass Beds: A

Review of the Influence of Plant Structure and Prey Characteristics on Predator-Prey

Relationships. Estuaries 7:339-350.

Ovaskainen, O., G. Tikhonov, A. Norberg, F. Guillaume Blanchet, L. Duan, D. Dunson, T.

Roslin, and N. Abrego. 2017. How to make more out of community data? A conceptual

framework and its implementation as models and software. Ecology Letters 20:561–576.

Presley, S.J., C. L. Higgins, and M. R. Willig. 2010. A comprehensive framework for the

evaluation of metacommunity structure. Global Ecology and Evolution 19:908–917.

R Development Core Team. 2014. R: a language and environment for statistical computing. R

Foundation for Statistical Computing. Vienna, Austria. http://www.R-project.org.

Rugiu, L., I. Manninen, J. Sjöroos, and V. Jormalainen. 2018. Variations in tolerance to climate

change in a key littoral herbivore. Marine Biology 165:1–11.

Sala, E., and M. H. Graham. 2002. Community-wide distribution of predator-prey interaction

strength in kelp forests. Proceedings of the National Academy of Sciences 99:3678–3683.

Thompson, P. L., B. Rayfield, and A. Gonzalez. 2017. Loss of habitat and connectivity erodes

species diversity, ecosystem functioning, and stability in metacommunity networks.

Ecography 40:98–108.

Treml, E.A., P. N. Halpin, D. L. Urban, and L.F. Pratson. 2008. Modeling population

connectivity by ocean currents , a graph-theoretic approach for marine conservation.

Landscape Ecology 23:19–36.

Treml, E. A., J. R. Ford, K. P. Black, and S. E. Swearer. 2015. Identifying the key biophysical

drivers, connectivity outcomes, and metapopulation consequences of larval dispersal in the

sea. Movement Ecology 3:17.

Tuomisto, H., L. Ruokolainen, and K. Ruokolainen. 2012. Modelling niche and neutral

dynamics: on the ecological interpretation of variation partitioning. Ecography 35:961-971.

Valanko, S., J. Heino, J. M. Westerbom, M. Viitasalo, and A. Norkko. 2015. Complex

metacommunity structure for benthic invertebrates in a low-diversity coastal system.

Ecology and Evolution 5:5203–5215.

Viana, D. S., and J. M. Chase. 2018. Spatial scale modulates the inference of metacommunity

assembly processes. In press. Ecology doi: 10.1002/ecy.2576

Vieira, E. A., A. A. V. Flores, and G. M. Dias. 2018. Persistence and space preemption explain

species-specifc founder effects on the organization of marine sessile communities. Ecology

and Evolution 8:3430-3442.

Vellend, M. 2016. The Theory of Ecological Communities: Princeton University Press.

Ward, J. H. 1963. Hierarchical Grouping to Optimize an Objective Function. Journal of the

American Statistical Association 58:236-244.

Waters, J. M., et al. 2013. Founder takes all: density-dependent processes structure biodiversity.

Trends in Ecology and Evolution 28:78-85.

Wichmann, C. S., I. A. Hinojosa, and M. Thiel. 2012. Floating kelps in Patagonian Fjords: An

important vehicle for rafting invertebrates and its relevance for biogeography. Marine

Biology 159:2035–2049.

Whippo, R., N. S. Knight, C. Prentice, J. Cristiani, M. R. Siegle, and M. I. O'Connor. 2018.

Epifaunal diversity patterns within and among seagrass meadows suggest landscape-scale

biodiversity processes. Ecosphere 9:02490.10.1002/ecs2.2490.

Wlodarska-Kowalczuk, M., E. Jankowska, L. Kotwicki, and P. Balazy. 2014. Evidence of

Season-Dependency in Vegetation Effects on Macrofauna in Temperate Seagrass

Meadows (Baltic Sea). PLoS One 9:e100788.

Yamada, K., Y. Tanaka, T. Era, and M. Nakaoka. 2014. Environmental and spatial controls of

macroinvertebrate functional assemblages in seagrass ecosystems along the Pacific coast of

northern Japan. Global Ecology and Conservation 2:47–61.

735   **Table 1.** Summary of evidence used to answer questions about metacommunity processes

Question	Analyses	Figure number
(1) Is the metacommunity characterized by dispersal limitation, intermediate dispersal, or high dispersal?	*Dispersal limitation:* Distance-decay of community similarity from HMSC joint species distribution models with and without accounting for environmental similarity	Fig. 2
	Intermediate versus high dispersal: Biodiversity patterns across sites for signatures of high dispersal (compositional homogenization); HMSC model results indicating structured co-occurrence patterns and variation in composition explained by environmental	Fig. 4a, 4b
(2) Do species distribution patterns suggest environmental niche filtering, and if so along which environmental axis?	HMSC model: Variation partitioning of modelled site and region-level fixed and random effects of environmental covariates and calculating predictive power (proportion of deviance) of the fixed effects model	Fig. 3
(3) Do species co-variance patterns suggest possible biotic interactions influencing community assembly?	HMSC model: Correlation matrix of pairwise species co-occurences, after controlling for the effect of space and environment	Fig. 4a & 4b

739   **Table 2.** Species names, taxonomic groupings, and proportion of deviance (predictive power).

740   Numbers correspond to the axes in Figure 3. The top twenty species with highest predicted

741   abundances according to the HMSC model are bolded.

Number	Species name	Broad taxonomic group	Adjusted deviance explained for the HMSC model (D^2_{adj})
1	*Alia carinata*	Gastropod (snail)	0.15
2	***Alvania compacta***	**Gastropod (snail)**	**0.4**
3	*Amphissa columbiana*	Gastropod (snail)	0
4	*Crepidula* sp.	Gastropod (snail)	0.64
5	*Euspira lewisii*	Gastropod (snail)	0.46
6	*Harmothoe imbricata*	Gastropod (snail)	0
7	***Lacuna* spp. (*L. variegata* and *L. vincta*)**	**Gastropod (snail)**	**0.45**
8	*Lirularia parcipincta*	Gastropod (snail)	0.46
9	*Lirularia* sp.	Gastropod (snail)	0
10	*Littorina* sp.	Gastropod (snail)	0.5
11	*Margarites pupillus*	Gastropod (snail)	0.45
12	*Lottia pelta*	Gastropod (limpet)	0.53
13	***Clinocardium nuttallii***	**Bivalve**	**0**
14	***Mytilus* sp.**	**Bivalve**	**0.28**
15	*Saxidomus gigantea*	Bivalve	0
16	*Unknown clam 1*	Bivalve	0.68
17	*Unknown clam 2*	Bivalve	0.35
18	*Dorvillea longicornis*	Polychaete	0.52
19	*Exogone sp.*	Polychaete	0.25
20	*Glycinde armigera*	Polychaete	0.05
21	*Lepidonotus squamatus*	Polychaete	0.55
22	*Neoamphitrite robusta*	Polychaete	0.22
23	*Unknown polychaete (Family Oenoidae)*	Polychaete	0.15
24	*Unknown polychaete (Family Opheliidae)*	Polychaete	0.36
25	***Spirorbis* sp.**	**Polychaete**	**0**
26	***Nereis* sp.**	**Polychaete**	**0.55**
27	***Ampithoe dalli***	**Gammarid amphipod**	**0.47**

28	*Ampithoe lacertosa*	Gammarid amphipod	0.34
29	***Ampithoe valida***	**Gammarid amphipod**	**0**
30	***Aoroides* spp.**	**Gammarid amphipod**	**0**
31	*Ceradocus spinicauda*	Gammarid amphipod	0
32	*Grandidierella japonica*	Gammarid amphipod	0
33	*Ischyocerus anguipes*	Gammarid amphipod	0.33
34	*Jassa marmorata*	Gammarid amphipod	0
35	***Monocorophium insidiosum***	**Gammarid amphipod**	**0.23**
36	*Orchomenella recondita*	Gammarid amphipod	0.73
37	***Photis brevipes***	**Gammarid amphipod**	**0**
38	***Pontogeneia rostrata***	**Gammarid amphipod**	**0.65**
39	*Unknown gammarid (Family Hyalidae)*	Gammarid amphipod	0.12
40	*Unknown gammarid (Family Isaeidae)*	Gammarid amphipod	0.10
41	*Unknown gammarid (Family Ischyoceridae)*	Gammarid amphipod	0.24
42	***Caprella californica***	**Caprellid amphipod**	**0.36**
43	***Caprella laeviuscula***	**Caprellid amphipod**	**0.06**
44	***Caprella natalensis***	**Caprellid amphipod**	**0.82**
45	*Cumella vulgaris*	Cumacean	0.36
46	***Leptochelia sp.***	**Tanaid**	**0.49**
47	*Nebalia gerkinae*	Leptostracan	0.39
48	***Porcellidium sp.***	**Copepod**	**0.33**
49	***Harpacticoid 1***	**Copepod**	**0**
50	***Harpacticoid 2***	**Copepod**	**0**
51	*Harpacticoid 3*	Copepod	0.50
52	*Anoplodactylus viridintestinalis*	Pycnogonid	0.01
53	*Idotea montereyensis*	Isopod	0.73
54	***Idotea resecata***	**Isopod**	**0.36**
55	*Munna sp.*	Isopod	0.20
56	*Pugettia producta*	Brachyuran crab	0.36
57	*Pugettia richii*	Brachyuran crab	0.34
58	*Hippolyte californiensis*	Caridean shrimp	0.45

747 **Figure 1.** Map of coastal British Columbia showing eelgrass field sites in Haida Gwaii (HL,

748 RA), Clayoquot Sound (DK, EB, IN), Barkley Sound (SA, DC, RB), and the Southern Gulf

749 Islands (GB, JB, CB, LH, SS).

762   **Figure 2**. HMSC model predictions of community similarity relative to the spatial distance

763   (Euclidean) between communities. The red circles show similarity predictions based on the full

764   model. The blue triangles show predictions wherein the effect of all environmental parameters (9

765   water quality and 5 biotic) has been removed (standardized to their mean values), thus showing

766   predicted community similarity as a result of spatial distance only.

Figure 3. Variance partitioning of fixed (environmental variables, spatial distance) and random

effects across all species (numbers correspond to Table 2) in the HMSC model. Purple cells

represent site and region-level random effects. Blue cells represent abiotic water quality

parameters. Yellow cells represent food availability (sum of variances explained by eelgrass,

algae and detritus biomass). Green cells represent habitat structure, which is the sum of variances

explained by eelgrass leaf area index and shoot density. Bars are organized according to broad

taxonomic groupings as indicated in Table 2. The percentages shown next to the legend labels

indicate mean variance explained by that environmental variable across all species. Adjusted

deviance explained (D^2_{adj}) for the individual species distribution model fits varied (see Table 2).

The mean D^2_{adj} across all species was 0.29.

Figure 4a. A correlation plot showing modelled site-level co-occurrence of species pairs in the

20 most abundant species. Purple cells represent positively co-occurring species pairs, and

turquoise cells represent negatively co-occurring species pairs. Species names along both axes

are ordered according to the output of hierarchical clustering with Ward's criterion on pairwise

co-occurrence values. A version of this figure with all 58 species is shown in Appendix S1: Fig.

S4.

Figure 4b. Heatmap and cluster dendrogram depicting modelled (HMSC) species abundance and compositional similarity across sites. The dendrogram was produced using hierarchical cluster analysis with Ward's criterion on predicted species proportional abundances. The heat map shows the twenty species with the highest predicted mean proportional abundance at the site level, and is ordered from highest to lowest predicted mean proportional abundance. Cell colours correspond to the HMSC co-occurrence groups shown in Fig. 4a: turquoise group, purple group, and grey group. Cell shade strength represents proportional abundance at a given site (darker means higher abundance). The symbols to the left of the site abbreviations indicate region membership; circles show Haida Gwaii sites, squares show Barkley Sound sites, diamonds show Clayoquot Sound sites, and triangles show Southern Gulf Islands sites. A version

803   of this figure with all 58 species is shown in Appendix S1: Fig. S5.

Appendix S1: Supporting Information

Equation S1. Equations for D^2_{adj} calculations

Explained deviance or D^2 (Guisan and Zimmerman 2000) was calculated by the equation:

$$D^2 = 1 - Residual\ deviance\ /\ Null\ deviance$$

Where the residual deviance refers to the deviance in predicted species occurrences, and the null
deviance refers to deviance in observed species occurrences.

Adjusted D^2 (Guisan and Zimmerman 2000) was calculated by the equation:

$$D^2_{adj} = 1-(((n-1)/(n-p-1)*(D^2-1))$$

Where n is the number of sites (13) in our study, and p is the number of parameters the model
estimated (4, Ovaskainen et al. 2017, Supporting Information). Negative D^2_{adj} values (those which
had greater residual deviance than null deviance) were set to 0.

Equation S2. Calculation for type III variation partitioning result

m1 = the fraction of variation explained by all environmental variables in the HMSC model
estimated with only environmental variables;

m2 = the fraction of variation by spatial distances in the version of the HMSC model estimated
with only spatial distances;

m3 = the global model including both;

Fraction *ab* (pure environment + shared fraction) = D^2_{adj} *m1*

Fraction *bc* (pure space + shared fraction) = D^2_{adj} *m2*

Fraction *abc* (pure environment + shared fraction + pure space) = D^2_{adj} *m3*

Fraction *a* (pure environment) = *abc* - *bc*

Fraction *c* (pure space) = *abc* - *ab*

Fraction *b* (shared fraction) = *ab* + *bc* - *abc*

Fraction *d* (residuals) = 1 - *abc*

Fractions *a* ("Environmental conditions"), *c* ("Spatial distance"), *b* (shared fraction), and *d*
("Residuals") are shown in the Venn diagram in Fig. S3.

882   **Figure S2.** Site-by-species presence-absence matrix of all 58 invertebrate species in the study

883   listed in alphabetical order. Black cells indicate species presences.

892   **Figure S3.** Venn diagram summarizing the fractions of variation explained by environmental

893   covariates only (9 water quality and 5 biotic variables), spatial distance only, and the shared

894   fraction explained by environment and space.

Figure S4. A correlation plot showing modelled site-level co-occurrence of species pairs across all species. Purple cells represent positively co-occurring species pairs, and turquoise cells represent negatively co-occurring species pairs. Species names along both axes are ordered according to the output of hierarchical clustering with Ward's criterion (Ward 1963) on pairwise co-occurrence values. This figure is a supplement to Fig. 4a in the main text.

Figure S5. Heat map depicting abundance patterns in all 58 species in the study. Species names

are ordered from highest to lowest predicted mean proportional abundance. Cell colours

correspond to the co-occurrence groups in Fig. 4b, with turquoise cells representing members of

the turquoise assemblage, purple cells indicating members of the purple assemblage, and grey cells

indicating species that did not significantly co-occur negatively or positively with other species.

Cell shade strength represents proportional abundance at a given site (darker means higher

abundance). Most species outside the top twenty most abundant had extremely low predicted

proportional abundances owing to their low raw abundances. This figure is a supplement to Fig.

4b in the main text.