Beyond a single patch: local and regional processes explain diversity patterns in a seagrass epifaunal metacommunity

Keila A Stark[1]*, Patrick L Thompson[1], Jennifer Yakimishyn[2], Lynn Lee[3], Emily M Adamczyk[1], Margot Hessing-Lewis[4], Mary I O'Connor[1]

1. Biodiversity Research Centre and Department of Zoology
University of British Columbia
Vancouver, BC, Canada

2. Pacific Rim National Park Reserve
P.O. Box 280
Ucluelet, BC, Canada

3. Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve, and Haida Heritage Site
60 Second Beach Road
Skidegate, BC, Canada

4. Hakai Institute
PO Box 309
Heriot Bay, BC, Canada

*Corresponding author: keilastark@zoology.ubc.ca

Abstract: 251
Main text: 5000
Number of references: 68
Number of figures: 3

Data accessibility: Data will be made permanently available on Dryad should the manuscript be accepted. All data and code for the analysis can be currently found at https://github.com/keilast/HMS-Seagrass.

Competing interests: We have no competing interests.

Acknowledgements

KAS and MIO conceived of the study. JY provided samples from IN, DK, and EB. LL provided

samples from RA and HL. EMA and MHL provided samples from TN, TB, CI, and CS. KAS

collected and processed samples, analyzed data, and wrote the first manuscript draft. PLT

provided recommendations for the analysis. KAS, PLT, and MIO contributed to writing the

manuscript. This research was financially supported by a UBC Zoology SURE Grant, and

NSERC Discovery Grants, Canadian Foundation for Innovation (CFI), and is sponsored by the

NSERC Canadian Healthy Oceans Network and its Partners: Department of Fisheries and

Oceans Canada and INREST (representing the Port of Sept-Îles and City of Sept-Îles). PLT is

supported by NSERC and Killam postdoctoral fellowships. We are grateful to the Tula

Foundation for providing logistical support and facilities during Calvert Island field collections.

Special thanks go to Michelle Paleczny and Gulf Islands National Park Reserve for boat and

laboratory space, and staff, students, and volunteers from Pacific Rim National Park Reserve for

their assistance collecting eelgrass. We thank Minako Ito, Ariane Comeau, Rob Underhill, Abbie

Sherwood, Giordano Bua for assistance with samples and Coreen Forbes, Matt Whalen, and

anonymous reviewers for feedback on the manuscript.

Abstract

Ecological communities are jointly structured by dispersal, density-independent responses to

environmental conditions and density-dependent biotic interactions. Metacommunity ecology

provides a framework for understanding how these processes combine to determine community

composition among local sites that are regionally connected through dispersal. In 17 temperate

seagrass meadows along the British Columbia coast, we tested the hypothesis that eelgrass

(*Zostera marina* L.) epifaunal invertebrate assemblages are influenced by local environmental

conditions, but that high dispersal rates at larger spatial scales dampen effects of environmental

differences. We used hierarchical joint species distribution modelling to understand the

contribution of environmental conditions, spatial distance between meadows, and species co-

occurrences to epifaunal invertebrate abundance and distribution across the region. We found

that patterns of taxonomic compositional similarity among meadows were inconsistent with

dispersal limitation and meadows in the same region were often no more similar to each other

than meadows over 1000 km away. Abiotic environmental conditions (temperature, dissolved

oxygen) explained a small fraction of variation in taxonomic abundances patterns across the

region. We found novel co-occurrence patterns among taxa that could not be explained by shared

responses to environmental gradients, suggesting the possibility that interspecific interactions

influence seagrass invertebrate abundance and distribution. Our results add to mounting evidence

that suggests that the biodiversity and ecosystem functions provided by seagrass meadows reflect

ecological processes occurring both within meadows and across seascapes, and suggest that

management of eelgrass habitat for biodiversity may be most effective when both local and

regional processes are considered.

Key words: Hierarchical Modelling of Species Communities, priority effects, dispersal, niche

filtering, British Columbia, *Zostera marina*

Introduction

Understanding how local environmental conditions, regional connectivity by dispersal and

biotic interactions jointly structure the composition of communities is a central challenge in

ecology (Ricklefs and Schluter 2003, Vellend 2010, Leibold and Chase 2017). Metacommunity

ecology (Leibold et al. 2004, Leibold and Chase 2017, Thompson et al. 2020) offers a

framework for understanding community assembly processes across spatial scales. In recent

years, applications of the metacommunity framework have emphasized the underlying processes

that give rise to abundance and diversity patterns (Brown et al. 2017, Leibold and Chase 2018,

Thompson et al 2020). Thompson et al. (2020) framed the metacommunity concept based on

three fundamental processes that together govern the dynamics of populations and communities:

1) Density-independent responses to environmental conditions, 2) Density-dependent biotic

interactions (ie. inter and intra-specific competition, predation or facilitation influencing

population growth and co-existence), and 3) Dispersal influencing connectivity across a

landscape or seascape. We apply this framework to understand the contribution of these

processes to a seagrass-associated invertebrate metacommunity.

Many coastal environments that host high biodiversity occupy spatially structured habitats

(e.g. coral reefs, kelp forests, seagrass meadows), and many marine species have dispersing life

histories that link populations in distinct habitat patches. The importance of interspecific

interactions (Berlow 1999, Sala and Graham 2002), and dispersal-driven population dynamics

(Levin and Paine 1974, Gaines and Roughgarden 1985) in coastal and marine communities has

long been recognized. These combined roles of dispersal, the local environment, and biotic

interactions suggest patterns of marine biodiversity reflect metacommunity processes (Boström

et al. 2006).

Seagrasses are foundation species that support high productivity and faunal diversity (Orth et

al. 1984, Duffy et al. 2015). They form meadows separated by deeper water, un-vegetated

seafloor, or other vegetated habitats, and associated fauna disperse among meadows (Boström et

al. 2006, 2011). Diverse invertebrate assemblages including snails, amphipods, isopods, and

polychaete worms live among seagrass leaves, providing food sources for larger invertebrates,

fish, and birds (Best and Stachowicz 2014, Huang et al. 2015). The grazers within this group

consume detritus, macroalgae, or the seagrass itself (Valentine and Heck 1999), and many taxa

exert top-down control on epiphytic microalgae that compete with the seagrass for light and

nutrients (Sand-Jensen 1977, Duffy and Stachowicz 2006). Seagrass-associated epifauna exhibit

a range of dispersal modes (fast swimming by isopods; slow crawling by snails; permanent

attachment in bivalves) and reproductive strategies (brooding versus broadcast spawning) which

influence dispersal rates and the distances over which meadows are demographically connected.

Seagrass meadows occur in a wide range of temperature, salinity and hydrodynamic

environments, such that meadows may differ in their suitability for various invertebrate taxa

(Yamada et al. 2007, 2014). Evidence for the importance of environmental filtering -- or the

tendency for taxa to exist where local environmental conditions are ideal -- has been reported in

several seagrass-associated fish and invertebrate systems (Baden et al. 2010, Robinson et al.

2011, Iacarella et al. 2018).

Though interspecific interactions among seagrass-associated invertebrates are not extensively

documented, there is some evidence of competition and predation influencing community

structure (Nelson 1979, Best and Stachowicz 2014). Predation can reduce abundances of

vulnerable invertebrate taxa, allowing others to increase in abundance (Nelson 1979, Baden et al.

2010, Best and Stachowicz 2014, Amundrud et al. 2015, Huang et al. 2015). Despite some

evidence for competition among epiphytic grazers for shared food sources (Edgar 1990, Bruno

and O'Connor 2005) there is no evidence of competitive dominance to the point of exclusion in

the field (Nelson 1979, Best and Stachowicz 2014).

While research in past decades has found evidence for direct and indirect influences of local

characteristics such as habitat complexity (Orth et al. 1984), primary biomass (Cébrian and

Duarte 1998, Gullström et al. 2012), and nutrient availability (Virnstein and Howard 1987) on

faunal diversity and abundance, recent studies have focused more on regional scale or multi-

meadow processes such as dispersal (Whippo et al. 2018, Lefcheck et al. 2016, Stier et al. 2019,

Yeager et al. 2019), and have found that the influence of local-scale factors are often overriden

by regional patterns of inter-meadow connectivity (Lefcheck et al. 2016, Stier et al. 2019, Yeager

et al. 2019). This pattern has been attributed to relatively rapid life histories and high inter-

meadow dispersal in seagrass-associated organisms (Lefcheck et al. 2016).

We designed this study to better understand the contributions of dispersal, local

environmental conditions, and interspecific interactions with the largest spatially explicit (in

geographic extent) dataset of seagrass metacommunity diversity. Drawing upon metacommunity

theory, we tested the hypotheses that: (H1) Spatial distance *per se* within a region does not

confer community dissimilarity because meadows are well-connected by dispersal, even at larger

spatial scales (*dispersal limitation hypothesis*); (H2) Differences in local-scale environmental

factors such habitat structure, temperature, and salinity drive some differences in invertebrate

presence and abundance owing to differences in environmental niches among taxa, however the

contribution of environmental factors is smaller than the contribution of spatial distance/

dispersal (*environmental filtering hypothesis*); (H3) No clear patterns of spatial co-occurrence

among epifaunal species emerge because high dispersal and environmental responses override

signatures of biotic interactions (predation, competition) that would produce such patterns (*biotic*

interactions hypothesis).

Methods

Study sites

We sampled epifaunal invertebrate diversity in 17 *Zostera marina* (L.) meadows

spanning the entire coast of British Columbia (approximately 1000 km distance) from late June

through August of 2017. Within five regions that each contain numerous eelgrass meadows, we

sampled a few meadows for our study (Fig. 1a): Haida Gwaii (two meadows); the central coast

of BC near Calvert Island (four meadows); southern Clayoquot Sound (three meadows); Barkley

Sound on the West coast of Vancouver Island (three meadows); the Southern Gulf Islands (five

meadows). Meadows varied in environmental conditions, such as seagrass shoot size and

density, hydrological regimes, and freshwater outflow influencing salinity.

Field sampling

We sampled three subtidal meadows with SCUBA in the Barkley Sound (SA, DC, RB)

and Central Coast (TB, TB, CI, CS) regions. All others were sampled by wading or snorkeling

at low tide. We conducted collections within a six-week period of peak seagrass growth. We

used 0.25 m x 0.25 m quadrats to collect all above-ground eelgrass, epifaunal invertebrates,

eelgrass detritus, and macroalgae in each sample, following Whalen et al. (2013). Six quadrats

were arranged in a 15 m x 30 m array (Appendix 1: Fig. S1) in the middle of the meadow to

avoid edge effects. We uprooted eelgrass shoots at the first node, leaving rhizomes to avoid

sampling infauna that live in the sediments, removed all other above-ground biomass (detritus,

macroalgae, associated epifauna) within each quadrat by hand, and immediately placed all the

contents into a 300 μm mesh or plastic Ziploc bag for transport to the lab.

Environmental data

We acquired water quality data from regional data sources. The abiotic water quality data

were annual means pulled from Bio-ORACLE (Assis et al. 2018), including oxygen, nitrates,

phosphates, silicates, salinity, maximum current velocity and sea surface temperature. Bio-

ORACLE is a database of marine environmental layers gathered from several satellite and in-situ

sources at a spatial resolution of 0.08°, and has been shown to accurately model distributions of

shallow-water invertebrate species (Tyberghein et al. 2012, Assis et al. 2018). Sea surface

temperature layers were taken from the Aqua-MODIS satellite, and values for nitrates, dissolved

oxygen, and salinity were interpolated from in-situ measurements reported in the World Ocean

Database (Tyberghein et al. 2012). All meadows were situated in distinct spatial cells, and thus

had distinct values for the aforementioned variables. We excluded phosphate and silicate

concentrations from our final analysis, as they strongly covaried with nitrates.

In each quadrat sample, we measured four biotic attributes summarizing habitat structure

and food availability: eelgrass leaf area index (LAI), and eelgrass, eelgrass detritus, and algal dry

mass. To quantify LAI, we measured leaf length, width, and blade number in five haphazardly-

chosen shoots from each quadrat, and multiplied the average blade area per shoot by the quadrat-

level shoot density. We dried eelgrass, detritus, and macroalgae in a desiccator oven (60° C for

48 hours) to measure ash-free dry mass.

Invertebrate identification

Immediately after collection, we rinsed eelgrass shoots with fresh water, passing the

water through a 500 μm sieve to remove epifaunal invertebrates. Invertebrates were preserved in

95% EtOH for identification with light microscopy (10x magnification). We identified

invertebrates to the lowest taxonomic level possible using keys in the Light and Smith manual

(Carlton 2007) and Kozloff (1996). In many cases, the lowest taxonomic resolution possible was

family or genus, therefore our biodiversity survey likely underestimates full species diversity.

Analysis

All statistical analyses were conducted in R (version 3.6.0; R Development Core Team

2019. We followed the Hierarchical Modelling of Species Communities (HMSC) framework

(Ovaskainen et al. 2017) using the 'Hmsc-R' package (Tikhonov et al. 2019) to fit a hierarchical

joint species distribution model (JDSM) with Bayesian inference. The framework uses traditional

single-species distribution modelling by estimating species responses (presence or abundance) to

environmental covariates across samples, but does so for all species simultaneously. It can use

residual variation in occurrences to infer species co-occurrence patterns that do not result from

shared responses to environmental covariates in the model (Ovaskainen et al. 2017). HMSC can

also account for spatially hierarchical sampling structures. In these ways, the HMSC framework

overcomes statistical limitations of previous methods used in metacommunity studies (Gilbert and

Bennett 2010, Tuomisto et al. 2012, Brown et al. 2017).

Each individual species distribution model within the JDSM is a generalized linear model

that describes the abundance of species j (where $j = 1 \dots n$), where y_{ij} is the abundance of species j

in sample i, D is the statistical distribution of the abundance data (Poisson distribution in this study),

L_{ij} is the linear predictor to link species' presence with environmental covariates, and σ^2_j is a

variance term for the abundance of species j:

$$y_{ij}^D \sim D(L_{ij}, \sigma_j^2).$$

224
The linear predictor L_{ij} is described by fixed (F) and random (R) effect terms:

$$L_{ij} = L^F_{ij} + L^R_{ij},$$

The fixed effect term L^F_{ij} is modelled as a regression:

$$L^F_{ij} = c_{jk}x_{ik}$$

where x_{ik} represents the measured value for environmental covariate k for a given sample i (e.g.,

biomass in quadrat i), and parameter β represents the relationship between environmental

covariate k, and the abundance of species j.

The random effect term L^R_{ij}, captures the variation in species abundances that cannot be explained

by the measured covariates. It is further denoted as:

$$L^R_{ij,} = \varepsilon_{ij}^Q + \varepsilon_{ij}^S + \varepsilon_{ij}^R + \varepsilon_{ij}^D$$

where the terms describe three random effects associated with our spatially nested sampling

scheme (ε_{ij}^Q, ε_{ij}^S, ε_{ij}^R for Quadrat (sample), Site (meadow), and Region), as well as a fourth spatially

explicit random effect (spatial autocorrelation in species abundances). The spatially explicit

random effect ε_{ij}^D was calculated with latent factor analysis that takes into account spatial distance

between all pairwise combinations of samples (see Ovaskainen et al. 2016a for details).

The three non-spatial random effect terms are assumed to have multivariate normal

distributions $\varepsilon_{ij}^Q \sim N(0, \Omega^Q)$, $\varepsilon_{ij}^S \sim N(0, \Omega^S)$, $\varepsilon_{ij}^R \sim N(0, \Omega^R)$. Variance terms Ω^Q, Ω^S, Ω^R, are

variance-covariance matrices (square matrices containing all taxa in the model), where the

diagonal elements give species-specific residual variance in occurrences among samples, and the

off-diagonal elements give residual co-variances between species pairs. The term "residual" refers

to the fact that it is the variance unexplained by environmental covariates in the fixed effects

predictor described above. These variance-covariance matrices (Ω^Q, Ω^S, Ω^R, and Ω^D) are

parameters estimated using a latent variable approach described in Ovaskainen et al. (2016b). They

were used to represent co-occurrence matrix R, where j_1 and j_2 refer to two species within the

model, and $R_{j1j2} = \Omega_{j_1j_1}/\sqrt{\Omega_{j_1j_1}\Omega_{j_2j_2}}$. R describes the extent to which a given species pair co-

occurs more positively or negatively than by chance, after controlling for possible shared responses

to the same environmental covariates.

We estimated parameters with Markov chain Monte Carlo. Markov chains were run to

200 000 iterations, a burn-in length of 1000 iterations, and was thinned to retain every 10^{th}

sample of the posterior distributions. We confirmed that Markov chains were well-mixed by

visually inspecting trace plots. Estimates of parameter β (environmental responses across taxa)

and Ω (variance-covariance matrices used to visualize species co-occurrence patterns

independent of the environment) were extracted as 95% credible intervals. We evaluated the

model fit (R^2 and Root-Mean-Square Error or RMSE) using 4-fold cross-validation.

To test our three hypotheses, we visualized different aspects of the parametrized JDSM.

To test for dispersal limitation (H1), we looked for increased pairwise community dissimilarity

among samples with increased pairwise spatial distance. We predicted species abundances in

every sample using our trained model, then calculated pairwise Bray-Curtis dissimilarity index

(Bray and Curtis 1957) on these predicted abundances. We then conducted log-linear regression

of predicted pairwise dissimilarity against pairwise spatial distance (km) on a logarithmic scale.

Increased community dissimilarity could reflect dispersal limitation due to prohibitively far

dispersal distances, *or* due to increasing differences in environmental conditions with increasing

spatial scale. To distinguish between these possibilities, we compared a regression of predicted

dissimilarity based on the full JDSM (including environmental covariates) with the predicted

dissimilarity calculated from the same trained model, but with no effect of the environment (all

environmental covariates were set to their mean values).

To test the environmental filtering hypothesis (H2), we analyzed β for each combination

of species and environmental covariate to determine whether they had a positive or negative

relationship. We also partitioned the variance in abundance explained by environmental

covariates, spatial autocorrelation, and sampling design random effects.

To test the biotic interactions hypothesis (H3), we represented co-occurrence matrix R as

a correlation plot for each random effect level, demonstrating the extent to which species pairs

co-occurred more negatively or positively than predicted by their modelled correlation with

environmental covariates at each spatial scale (Ovaskainen et al. 2017, Aivelo and Norberg

2018).

Results

Taxonomic abundance and distribution patterns

We identified 52 282 individuals representing at least 50 taxa across the region

(Appendix 2: Table 2). Of these, 3% of individuals were bivalves, 12% were gastropods (snails

and limpets), 14% were copepods, 41% were polychaetes (most of which were calcifying

polychaetes *Spirorbis* sp.), 13% were gammarid amphipods, 6% were caprellid amphipods, and

the remaining 11% included other crustaceans (isopods, tanaids, cumaceans, crabs, shrimp).

These taxa span several phyla, diet types (herbivores, detritivores, suspension feeders), and

dispersal strategies (brooding, broadcast spawning). Mean meadow-level taxonomic richness

was 20 taxa per 0.0625 m^2 of seagrass area.

Several taxa were present at all 17 meadows: harpacticoid copepods *Porcellidium* sp.

(4% of all individuals across the entire region), snails *Lacuna* spp. (7%), and tanaids *Leptochelia*

sp. (10%). We found genera *Mytilus* sp. (mussel) and *Nereis* sp. (polychaete worm) in all

meadows, however we were not confident the same species were present at each meadow. Ten

taxa were present in every region, but not necessarily every meadow: isopod *Pentidotea*

resecata, gammarid amphipods *Monocorophium insidiosum, Ampithoe valida, Pontogeneia*

rostrata, Ampithoe dalli, Aoroides spp.*,* snails *Amphissa columbiana* and *Alia carinata,* and

limpet *Lottia pelta*.

We excluded rare species (those found in fewer than 5% of samples, or fewer than 5 out

of 102 quadrats) from our JDSM, leaving 33 taxa in our modelling analysis (Appendix 1: Table

S2). This was necessary to avoid statistically over-inflating the importance of model covariates.

Fig. 1A Map of coastal British Columbia showing eelgrass meadow sites in Haida Gwaii (HL, RA), Clayoquot Sound (DK, EB, IN), Barkley
sound (SA, DC, RB), near Calvert island in the central coast (TB, TN, CS, CI), and the southern gulf islands (GB, JB, CB, LH, SS).

Spatial connectivity via dispersal driving community similarity

The highest Bray-Curtis dissimilarity value was 0.99 for a pair of sites spaced 787 km apart (HL in Haida Gwaii and SA in Clayoquot Sound). The two communities at the highest pairwise distance (997 km, RA in Haida Gwaii and SS in the Gulf Islands) had a dissimilarity index of 0.8. The lowest dissimilarity index value was 0.59 at a pairwise spatial distance of 0.6 km (LH and SS in the Gulf Islands). A log-linear regression of Bray-Curtis dissimilarity calculated from predicted abundances using the JDSM showed that community dissimilarity increased with increased spatial distance (Fig. 1b, $y = 0.75 + 0.023\log(x)$, n = 272, $R^2 = 0.16$). The maximum and minimim pairwise dissimilarity values were reduced when the effect of the environment was removed ($y = 0.74 + 0.021\log(x)$, n = 272, $R^2 = 0.07$), suggesting environmental variables drive some dissimilarity in species composition with increased spatial distance.

Figure 1b. Bray-Curtis pairwise community dissimilarity index as a function of spatial distance (Euclidean) between every pairwise combination
of meadows in our study. The purple points and line show pairwise dissimilarity predicted by the full joint species distribution model using the
original environmental and spatial data. The yellow points and line show pairwise dissimilarity predicted by the same trained model, but the
effect of all environmental variables has been removed (set to their mean values). The yellow points and line show calculated pairwise
dissimilarity based on the raw abundance data.

Environmental conditions

The assessment of model fit and variance partitioning analysis both showed that the importance

of environmental covariates differed among taxa, and that overall the environment explained

relatively a low proportion of variation in species abundances across the region (Fig 2a). While

the mean R^2 across individual species distribution models was 0.27, individual R^2 values varied

(0 to 0.68, Appendix 1: Table A2) indicating variability in model fit.

Figure 2a. Variation partitioning of fixed and random effects within the joint species distribution model. Variances explained
have been adjusted to reflect the model fit (Multiplied by Pseudo R^2). An unadjusted version of the variance partitioning analysis
can be found in Appendix 1, Fig. S3 of the Supplementary Information. Dashed lines indicate broad invertebrate taxonomic
groupings. Grey cells represent the contribution of random effects associated with the sampling design. Yellow cells represent
abiotic water quality covariates. Pink cells represent biotic covariates (food availability, habitat structure). The percentages next
to the legend labels indicate mean variation explained by that covariate/ random effect across all species distribution models.

Overall, abiotic water quality variables explained more variation than biotic variables
(Fig. 2a). Mean and range of sea surface temperatures and dissolved oxygen explaining the
largest proportions of variation on average. Herbivore food availability (seagrass, algae biomass)
and habitat structure (seagrass shoot density and LAI) had lower influence on abundance (pink
bars, Fig. 2a). The importance of some environmental covariates differed markedly among taxa;
for example, nitrate levels explained approximately 27% of variance in the abundance of
gastropod *Alvania compacta* across meadows, but only 5% of variance in gastropod *Alia*
carinata (Fig 2a). The mean variance in taxon abundance explained by spatial autocorrelation

was 2.2% (pale gray bars, Fig. 2a), and other hierarchical random effects associated with the

sampling design explained 2.92% (quadrat-level), 2.59 (meadow or site-level), and 1.42%

(region-level) of variation.

The strength and directionality of responses to environmental variables also differed

among taxa, based on inspection of the β parameters that estimate the relationships between

environmental covariates and species abundances (Fig. 2b). None of the taxa in our study

showed statistically supported responses (β within 95% credible interval) to all environmental

variables. The few that showed a significant response to eelgrass LAI responded positively

(higher LAI meant higher abundances). Thirteen taxa responded negatively to higher temperature

ranges, and two responded positively (Fig. 2b). The majority of taxa (22 out of 33) had

significant responses to dissolved oxygen levels. Of these, 8 responded negatively (high

dissolved O_2 yields lower abundances), and 14 responded positively (high dissolved O_2 yields

lower abundances).

Figure 2b. Heat plot summarizing beta (β) parameters, or responses to all environmental covariates in the joint species distribution model. Coloured cells indicate that the response is statistically supported (β falls within 95% credible interval). Red cells indicate positive responses to a given environmental covariate (higher abundance correspond with higher values of the environmental covariate), whereas blue cells indicate negative responses (lower abundance with higher values of the environmental covariate).

Co-occurrence patterns

Co-occurrence patterns were less strong at the between-quadrat and between-region spatial scales (Fig 3a, 3c). In comparisons among quadrats, we observed more positive species co-occurrences than negative co-occurences (Fig. 3a). In comparisons among meadows, we identified, *post hoc*, two main species co-occurrence groupings (Fig. 3b). Members of the first

group (hereafter referred to as *Leptochelia* group) included *Leptochelia* sp. (tanaid), *Photis*

brevipes (gammaridean amphipod), *Amphissa columbiana* (snail), *Spirorbis* sp. (calcifying

polychaete), *Nereis* sp. (polychaete worm), *Caprella laeviuscula* and *Caprella natalensis*

(caprellid amphipods) (Fig. 3b). These taxa positively co-occurred more often than expected by

chance or by environmental filtering. They negatively co-occurred with members of the second

group, which included a harpacticoid copepod species, *Mytilus* sp. (mussel), *Porcellidium* sp.

(copepod), *Ischyrocerus* sp. (gammarid amphipod), and *Penidotea resecata* (isopod) (hereafter

referred to as harpacticoid group). *Alvania compacta* and *Lirularia* sp. (snails) positively co-

occurred with each other but did not strongly co-occur with members of the two main groups.

Remaining taxon pairs did not exhibit strong positive or negative co-occurrence patterns.

The two co-occurrence groups were represented at all meadows, suggesting that members

from these groups can be simultaneously present (Fig. 3d). However, their abundances tended to

negatively co-vary; at meadows JB, HL, CB, RA, EB, DC, and DU, members of the *Leptochelia*

group were most abundant, whereas Harpacticoid group members were most abundant at SS, SA,

LH, GB and RB. Meadow HL was an outlier, as it was strongly dominated by *Spirorbis* sp. The

emergent species groupings identified in Fig. 3b do not clearly correspond with geographical

structure because the cluster analysis did not group meadows by region membership (black

symbols, Fig. 3d).

Figure 3. a-c) Correlation plots showing modelled a) quadrat-level, b) site-level, and c) region-level pairwise co-occurrences after removing the effect of shared responses to the environmental covariates in our model. Only the 20 most abundant species (according to total abundance across the metacommunity or study region) are represented. Purple cells represent positively co-occurring species pairs, and turquoise cells represent negatively co-occurring species pairs. Species names are ordered according to the output of hierarchical clustering with Ward's criterion on pairwise co-occurrence values. **d)** Heatmap and cluster dendrogram depicting species relative abundance and compositional similarity across sites. Species are ordered by decreasing abundance from left to right. Cell colours correspond to the two main site-level co-occurrence groupings shown in Fig. 3b: purple or "*Leptochelia*" group, and blue or "Harpacticoid" group. Cell shade strength indicates proportional abundance at a given site (darker means higher relative abundance). The black symbols to the left of the site abbreviations indicate region membership; stars indicate Haida Gwaii sites, diamonds indicate Central Coast sites, triangles indicate Barkley Sound sites, squares indicate Clayoquot Sound sites, and triangles indicate Southern Gulf Islands sites.

Discussion

With a biodiversity survey of 17 seagrass meadows across an approximately 1000km

spatial extent, we fitted a joint species distribution model to test hypotheses about the

contributions of dispersal, environmental filtering, and species interactions to epifaunal

invertebrate abundance and distribution. We also documented at least two previously

unrecognized species' associations in the region. This study is the largest (in geographical

extent) spatially explicit seagrass-based metacommunity study to our knowledge. We found little

support for complete dispersal limitation even across over 1000 km of coastline, some support

for the importance of environmental niche filtering, and co-occurrence patterns that might have

arisen from interspecific interactions. Overall, these results suggest that seagrass-associated

biodiversity patterns reflect ecological processes spanning local (meadow-scale) to regional

scales, and provide additional support for recent findings that eelgrass-associated diversity

reflects regional-scale ecological processes in addition to local scale processes (Lefcheck et al.

2016, Whippo et al. 2018, Stier et al. 2019, Yeager et al. 2019).

Dispersal limitation is unlikely at this regional spatial scale

Our findings that: 1) Many species were present in all regions or all meadows; 2) The

effect of spatial distance alone did not explain decay in community similarity (Fig. 1b); 3) There

was a low contribution of spatial autocorrelation to variance in species abundance (Fig. 2a); and

4) Hierarchical cluster analysis did not group meadows according to spatial regions (black

symbols in Fig. 3d), all suggest that distance between meadows likely does not confer dispersal

limitation preventing populations at distant sites from exchanging individuals. It is however

likely that meadows that are physically near each other likely experience higher rates of

exchange in individuals than distant ones, given our observation that dissimilarity increases

logarithmically with increased spatial distance. The six taxa that were present at all meadows and

an additional ten taxa that were present in all regions had representation from across phyla and

life history traits (including rare taxa), suggesting that no single taxonomic group or dispersal

strategy had a consistently larger spatial distribution than others. Many meadows were small (< 1

ha), and epifaunal abundances can vary substantially within meadows from year to year (Nelson

1997, Douglass et al. 2010) such that local extinctions followed by rescue from nearby

populations are likely (Thom et al. 1995). However, the specific pathway that dispersing

organisms travelled to arrive at these meadows remains unknown. We used Euclidean distances

in our analysis, which preserved the rank order of distances between meadows, but actual

distances are likely greater due to oceanographic circulation patterns (Kinlan and Gaines 2003,

Mitarai et al. 2008).

It is unclear whether distant meadows share taxa because they: 1) Are linked by direct

dispersal via oceanographic processes (currents); 2) Are indirectly linked by dispersal via

unsampled "stepping-stone" meadows or other habitat types; or 3) Were colonized by

populations of the same species in a historical dispersal event, but have not seen the exchange of

individuals since (Lefcheck et al. 2016). Seagrass-associated epifauna generally disperse

passively between meadows, either through larval transport in currents or by "rafting" on

floating pieces of seagrass or macroalgae. Rafts have been observed dispersing benthic and

epifaunal invertebrates across phyla (echinoderms, peracarids, molluscs, annelids), and facilitate

connectivity between coastal ecosystems across 100s of kilometres (Wichmann et al. 2012).

Our inferences are based on an analysis that necessarily emphasized common taxa –

either numerically dominant or present in most meadows, or both. We observed seventeen rare

taxa that were found in fewer than 5% of our samples, often at extremely low abundances (1-2

individuals per meadow, or per 6 quadrats) in our survey of over 50 000 individuals. The spatial

extent of their range could be dispersal-limited for reasons other than prohibitively long travel

distances. Populations with low abundances may not disperse in appreciable numbers, and

therefore cannot establish populations as readily as abundant species; this is supported by our

observation of a positive relationship between range size (the number of meadows a species was

observed at) and abundance (Appendix 1: Fig. S4). This highlights a potential bias in studies like

ours against detecting dispersal limitation if it is most severe for rare taxa. Increasing the sample

size might reduce the risk of this type of bias, but would cost a substantial amount of time

associated with invertebrate identification. More efficient biodiversity sampling methods (e.g.

eDNA) could overcome this time cost.

Theory predicts that intermediate dispersal ('dispersal sufficiency', Leibold and Chase

2017) allows species to colonize sites where local environmental conditions are optimal for

growth and reproduction, whereas high dispersal ('dispersal surplus') overrides this, allowing

species to persist in habitats that cannot sustain positive population growth without substantial

immigration ('mass effects', Mouquet and Loreau 2013). A subset of species was present at all

meadows, but their abundances varied (Fig 3d); this pattern may suggest weak mass effects,

where dispersal rates are high enough for several taxa to occur at most meadows even if local

conditions are suboptimal, but not so high as to completely overwhelm the signature of

environmental conditions (Schmida and Wilson 1985, Mouquet and Loreau 2003).

Metacommunity theory predicts that such weak mass effects are likely whenever dispersal rates

are not limiting (Thompson et al. 2017), particularly in organisms that cannot control their own

dispersal (Leibold and Chase 2017) such as seagrass-associated invertebrates. This may explain

the poor model fit in some ubiquitous taxa such as *Amphissa columbiana* ($R^2 = 0.05$, Appendix

1: Table S2). The poor model fit may suggest that these taxa either 1) persist at several meadows

even if local environmental conditions are not optimal, due to sufficient immigration, or 2) have

broader environmental niches, and thus the environmental gradient observed across the study

region is not sufficient to observe clear responses (lower or higher abundances) in these taxa.

Regional species abundance patterns suggest weak environmental filtering

We found that differences in local-scale environmental factors account for some

differences in invertebrate presence and abundance (H2), however there remains a great degree

of unexplained variation given the imperfect model fit (mean $R^2 = 0.27$). Abiotic environmental

variables had a greater influence on invertebrate distribution and abundance than biotic variables

(Fig. 2a). The estimated β parameters suggest that, generally speaking, environments that are

saltier, have a higher seagrass surface area, smaller annual temperature range, and higher

dissolved oxygen levels have higher abundances of many taxa in this study (Fig. 2b). However,

there were several exceptions to this trend, highlighting the importance of understanding

individual species responses to the environment as opposed to the community as a whole.

Overall, we conclude that, at least in taxa for whom the model that had a higher fit (e.g.

Leptochelia sp., $R^2 = 0.52$), environmental filtering influences regional patterns in abundance

and distribution.

Differences in environmental conditions may influence distribution and abundance across

the region in one of three ways. First, abiotic variables such as temperature and nutrient

availability may influence food availability through primary productivity. This is possible despite

the fact that biometric variables (seagrass LAI, biomass) explained a relatively low proportion of

variance in abundance; while these measurements were taken at a single time point during field

sampling, the abiotic variables represented annual averages (Assis et al. 2018) and therefore

better represent long-term productivity. Second, temperature and salinity may affect species'

abundances via environmental tolerance ranges; for example, the isopod *Idotea baltica* (a

relative of *P. resecata* and *P. montereyensis* in our study) experiences significantly lower

survival in prolonged exposure to higher temperatures and lower salinities than those in their

home habitat (Rugiu et al. 2018). Third, temperature may influence invertebrate metabolic

demands, thereby influencing consumption rates and available primary biomass (O'Connor

2009). Reduced food availability may increase competition, thus altering the number of

individuals a seagrass patch can host.

Environmental conditions in temperate seagrass meadows fluctuate seasonally, driving

changes in epifaunal community structure throughout the year (Nelson et al. 1979, Wlodarska-

Kowalczuk et al. 2014, Whippo et al. 2018). Sampling meadows several times within a year to

capture these temporal changes would clarify the importance of the environment if invertebrate

assemblages shift with changes in environmental variables following the linear relationships

uncovered in our analysis.

Species co-occurrence patterns may suggest the importance of interspecific interactions

We did not observe strong between-region co-occurrence patterns, suggesting that there

do not appear to be clear regional assemblages corresponding with the five subregions in our

analysis (Figs. 3c, black symbols in 3d). We also did not observe strong co-occurrence patterns

between quadrats, however we did observe strong co-occurrence patterns at the meadow-level

spatial scale. The explanation is that meadows had distinct assemblages, and these assemblages

tended to be homogenized within meadows; if two taxa co-occurred in a given meadow (Fig.

3b), then they likely co-occurred in all quadrats within the meadow (Fig. 3a). This is consistent

with previous findings that epifaunal diversity patterns do not differ from random patterns within

meadows (between quadrats) (Whippo et al. 2018).

At the meadow-level, two main co-occurrence groupings showed antagonistic abundance

patterns across meadows that could not be explained by spatial structure or shared responses to

the environment (Fig. 3d). These co-occurrence groups have not, to our knowledge, been

explicitly documented before in this region. While there are multiple possible explanations for

non-random species co-occurrence patterns (Connor and Simberloff 1979), we ruled out distance

between meadows and measured environmental variables, because co-occurrence values were

extracted from residual variation unexplained by the environment or space (Ovaskainen et al.

2017). This residual variation therefore likely can be explained by a combination of unmeasured

environmental variables, stochasticity, and biotic interactions.

Several interaction types may influence abundance and diversity. The majority of species

in our survey are herbivores or detritivores, and thus may compete for primary (mostly epiphytic

algae) biomass. Laboratory experiments have shown that grazing rate and habitat selection in

amphipods are altered in the presence of morphologically and functionally similar interspecific

competitors (Howard 1985, Brooks and Bell 2001, Beermann et al. 2018). Other species in our

study are suspension feeders, and thus may not compete for food but possibly predator-free space

or substrate on eelgrass blades.

The invertebrate assemblage at a given meadow may have multiple possible

compositional states depending on the arrival order of species; this phenomenon is known as

priority effects (Fukami et al. 2016, Ke and Letten 2018) and has been documented in marine

fouling communities (Vieira et al. 2018). The antagonistic co-occurrence groupings in our

analysis could suggest priority effects at the meadow-level. It is possible that, following seasonal

declines in abundance or a disturbance event, the first few populations to increase in abundance

or colonize a meadow determine the success of others. Abundance patterns in the sessile

calcifying polychaete *Spirorbis* sp. might demonstrate an example of priority effects. *Spirorbis*

sp. dominated the HL meadow on Haida Gwaii; there were approximately 16 300 individuals,

several orders of magnitude higher than its abundance elsewhere. Meadow HL also had the

lowest taxonomic richness of all 17 meadows (9 species), and we observed fewer micro and

macro-epiphytes on eelgrass blades. Instead, the eelgrass was completely covered with *Spirorbis*

sp. This phenomenon has also been observed in *Z. marina* meadows in Akkeshi-Ko estuary,

Japan (Clark et al. 2018) and in *Thalassia testudinum* meadows in the Northwestern Gulf of

Mexico (Dirnberger 1990). Experimental evidence suggests that *Spirorbis* spp. larvae tend to

settle lower in the water column on newer seagrass growth, away from epiphytic algae and

previously settled conspecifics (Dirnberger 1990). Settlement rates were determined by

planktonic larval density rather than space availability on seagrass blades (Dirnberger 1990).

Given this, it is possible that *Spirorbis* sp. dominates HL from a combination of density-

dependent processes (high larval recruitment) and environmental conditions (high salinity, low

nitrates). The low epiphyte load on seagrass at HL, whether mostly driven by high *Spirorbis* sp.

densities or by the environment, may explain the low abundance and diversity of all other

invertebrate taxa. Overall, the importance of priority effects likely depends on the age,

colonization history, and disturbance regime of seagrass meadows.

Negative co-occurrence patterns in taxonomic distributions could also reflect the higher

abundance of predators at some meadows. Field experiments have shown that changes in

predation pressure by fish, shorebirds, and predatory invertebrates can shift seagrass-associated

epifaunal assemblages in a matter of weeks (Amundrud et al. 2015; Huang et al. 2015). Previous

studies involving some of the meadows in this analysis (Haida Gwaii, Clayoquot Sound, and

Barkley Sound) found high variation (beta diversity) in fish assemblages among meadows

(Robinson et al. 2011, Iacarella et al. 2018). A comprehensive survey of fish and bird predators

is required to determine the extent to which top-down trophic interactions structure these

invertebrate communities. Positive co-occurrence patterns may additionally be a result of

positive biotic interactions. An example is *Orchomenella recondita* - a gammarid amphipod that

lives in the gastrovascular cavity of the anemone *Anthopleura elegans* (Carlton 2007). This

species was only recorded at the SS meadow, and specifically only found in quadrats where *A.*

elegans was collected with the seagrass shoots.

While inferences from data taken at a single time point have limitations, our JDSM

approach suggests that there is more to learn about species interactions and the interplay between

interactions and dispersal in structuring these communities. The hypothesized processes driving

seagrass epifaunal diversity patterns that can be further tested with experiments to test for

priority effects or trophic interactions, with population genetics to test for demographic

connectivity, or with particle tracking models (e.g. Treml et al. 2008) to estimate dispersal

velocity.

Conclusion

Seagrasses are important coastal foundation species, valued throughout the world for

supporting diverse and productive food webs (Constanza et al. 1997, Williams and Heck 2001).

The extent of these ecosystems is declining at rates believed to exceed those of coral reef and

rainforest habitat loss, suggesting major losses of associated biodiversity (Waycott et al. 2009).

Our finding that several seagrass-associated faunal taxa across functional groups are consistently

present across a fairly large spatial scale is consistent with notion that "the mobile, fast-

reproducing, and generally omnipresent animal community is keenly responsive to the presence

of habitat" (Lefcheck et al. 2016). Our study suggests that the management of seagrass for its

ecosystem services may be more effective if both local and regional processes are considered

explicitly in habitat protection plans (Economo 2011).

Literature Cited

Aivelo, T. and Norberg, A. 2018. Parasite-microbiota interactions potentially affect intestinal

communities in wild mammals. J. Anim. Ecol. 18:438-447.

Amundrud, S. L. et al. 2015. Indirect effects of predators control herbivore richness and

abundance in a benthic eelgrass (*Zostera marina*) mesograzer community. – J. Anim. Biol.

84: 1092–1102.

Assis, J. et al. 2018. Bio-ORACLE v2.0: Extending marine data layers for bioclimatic modelling.

– Global Ecol. Biogeogr. 27:277–284.

Beermann, J. et al. 2018. Combined effects of predator cues and competition define habitat

choice and food consumption of amphipod mesograzers. - Oecologia 186:645–654.

Best, R., and Stachowicz, J. J. 2014. Phenotypic and phylogenetic evidence for the role of food

and habitat in the assembly of communities of marine amphipods. - Ecolog. 95: 775-786.

Berlow, E. L. 1999. Strong effects of weak interactions in ecological communities. - Nature 398:

330–334.

Boström, C. et al. 2006. Seagrass landscapes and their effects on associated fauna: A review. -

Estuarine, Coastal Shelf Sci. 68: 383-403.

Boström, C. et al. 2010. Invertebrate dispersal and habitat heterogeneity: Expression of

biological traits in a seagrass landscape. J. Exp. Mar. Biol. Ecol. 390:106-117.

Bray, J. R. and J. T. Curtis. 1957. An ordination of upland forest communities of southern

Wisconsin. – Ecol. Monogr. 27:325-349.

Brooks R.A., and Bell S.S. 2001. Mobile corridors in marine landscapes: Enhancement of faunal

exchange at seagrass/sand ecotones. 2001. – J. Exp. Mar. Bio. Ecol. 264: 67–84.

Brown, B. L. et al. 2017. Making sense of metacommunities: dispelling the mythology of a

metacommunity typology. - Oecologia 183:643–652.

Carlton, J. 2007. The Light and Smith Manual: Intertidal invertebrates from Southern California

to Oregon. Fourth edition. University of California Press, Oakland, California, USA.

Cébrian, J. and Duarte, C. M. 1998. Patterns in leaf herbivory on seagrasses. – Aquat. Bot.

60:67–82.

Connor, E. F. and Simberloff, D. 1979. The assembly of species communities: Chance or

competition? - Ecology 60:1132–1140.

Costanza, R., et al. 1997. The value of the world's ecosystem services and natural capital. -

Nature 387:253–260.

Cottenie, K. 2005. Integrating environmental and spatial processes in ecological community

dynamics. – Ecol. Lett. 8:1175–1182.

Duffy J. E. et al. 2015. Biodiversity mediates top-down control in eelgrass ecosystems: A global

comparative-experimental approach. – Ecol. Lett. 18:696–705.

Economo, E. P. 2011. Biodiversity conservation in metacommunity networks: Linking pattern

and persistence. – Amer. Nat. 177:167-180.

Fukami, T. et al. 2016. A framework for priority effects. – J. Veg. Sci. 27:655–657.

Gaines, S. and Roughgarden, J. 1985. Larval settlement rate: A leading determinant of structure

in an ecological community of the marine intertidal zone. – Proc. Natl. Acad. Sci.

82:3707–-3711.

Gilbert, B. and Bennett, J. R. 2010. Partitioning variation in ecological communities: do the

numbers add up? – J. Appl. Ecol. 47:1071–1082.

Guillaume Blanchet, F. et al. 2018. HMSC: Hierarchical Modelling of Species Community. R

package version. 2.1-2.

Gullström, M. et al. 2012. Spatial patterns and environmental correlates in leaf-associated

epifaunal assemblages of temperate seagrass (*Zostera marina*) meadows. – Mar. Biol.

159:413–425.

Huang, A. C. et al. 2015. Top-down control by great blue herons *Ardea herodias* regulates

seagrass-associated epifauna. - Oikos 124:1492–1501.

Howard, R. K. 1985. Measurements of short-term turnover of epifauna within seagrass beds

using an in situ staining method. – Mar. Ecol. Prog. Ser. 22: 163–168.

Ke, P. J. and Letten, A. D. 2018. Coexistence theory and the frequency-dependence of priority

effects. – Nat. Ecol. Evol. 2:1691–1695.1–7.

Kinlan, B. P. and Gaines, S. D. 2003. Propagule dispersal in marine and terrestrial environments:

A community perspective. - Ecology 84:2007–2020.

Kozloff, E. 1996. Marine Invertebrates of the Pacific Northwest. University of Washington

Press, Seattle, Washington, USA.

Leibold, M. A. et al. 2004. The metacommunity concept: A framework for multi-scale

community ecology. – Ecol. Lett. 7:601–613.

Leibold, M. A. and Chase, J. M. 2017. Metacommunity Ecology, Volume 59. Princeton

University Press, Princeton, New Jersey, USA.

Lefcheck, J. S. et al. 2016. Faunal Communities Are Invariant to Fragmentation in Experimental

Seagrass Landscapes. - PLoS ONE 11(5): e0156550.

Levin, S. A. and Paine, R. T. 1974. Disturbance, patch formation, and community structure. –

Proc. Natl. Acad. Sci. 71:2744–2747.

Logue, J. et al. 2011. Empirical approaches to metacommunities: a review and comparison with

theory. – Trends Ecol. Evol. 26:482–491.

Mitarai, S. D. et al. 2008. A numerical study of stochastic larval settlement in the California

Current system. – J. Marine Syst. 69:295–309.

Morlon, H. et al. 2008. A general framework for the distance-decay of similarity in ecological

communities. - Ecol. Lett. 11:904-917.

Mouquet, N. and Loreau, M. 2003. Community patterns in source-sink metacommunities. –

Amer. Nat. 162:554–557.

Nelson, W. G. 1979. An analysis of structural pattern in an eelgrass (*Zostera marina* L.)

amphipod community. - J. exp. Mar. Biol. Ecol. 39:231-264.

O'Connor, M. I. 2009. Warming strengthens an herbivore-plant interaction. - Ecology 90:388–

398.

Orth, R. J. et al. 1984. Faunal communities in seagrass beds: A review of the influence of plant

structure and prey characteristics on predator-prey relationships. - Estuaries 7:339–350.

Ovaskainen, O. et al. 2016a. Uncovering hidden spatial structure in species communities with

spatially explicit joint species distribution models. - Methods Ecol. Evol. 7:428–436.

Ovaskainen, O. et al. 2016b. Using latent variable models to identify large networks of species-

to-species associations at different spatial scales. - Methods Ecol. Evol. 7:549–555.

Ovaskainen, O. et al. 2017. How to make more out of community data? A conceptual framework

and its implementation as models and software. – Ecol. Lett. 20:561–576.

R Development Core Team. 2014. R: a language and environment for statistical computing. R

Foundation for Statistical Computing. Vienna, Austria. http://www.R-project.org.

Ricklefs, R. E. and Schluter, D. 1993. Species Diversity in Ecological Communities: Historical

and Geographical Perspectives. University of Chicago Press, Chicago, Illinois, USA.

Rugiu, L. et al. 2018. Variations in tolerance to climate change in a key littoral herbivore. – Mar.

Biol. 165:1–11.

Sala, E. and Graham, M. H. 2002. Community-wide distribution of predator-prey interaction

strength in kelp forests. – Proc. Natl. Acad. Sci. 99:3678–3683.

Sand-Jensen. 1977. Effect of epiphytes on eelgrass photosynthesis. – Aquat. Bot. 3:55-63.

Schmida, A. and Wilson, M. V. 1985. Biological determinants of species diversity. – J.

Biogeogr. 12:1–20.

Stier, A. C. et al. 2019. Temporal variation in dispersal modifies dispersal-diversity relationships

in an experimental seagrass metacommunity. – Mar. Ecol. Prog. Ser. 613:67-76.

Thom, R. et al. 1995. Temporal patterns of grazers and vegetation in a temperate seagrass

system. – Aquat. Bot. 50:201–205.

Thompson, P. L. et al. 2017. Loss of habitat and connectivity erodes species diversity, ecosystem

functioning, and stability in metacommunity networks. - Ecography 40:98–108.

Thompson, P. L. et al. 2020. A process-based metacommunity framework linking local and

regional scale community ecology. –Ecol. Lett. (In press)

Treml, E. A. et al. 2008. Modeling population connectivity by ocean currents: a graph-theoretic

approach for marine conservation. – Landsc. Ecol. 23:19–36.

Tuomisto, H. et al. 2012. Modelling niche and neutral dynamics: on the ecological interpretation

of variation partitioning. - Ecography 35:961–971.

Tyberghein, L. et al. 2011. Bio-ORACLE: a global environmental dataset for marine species

distribution modelling. – Glob. Ecol. Biogeogr. 21: 272-281.

Vieira, E. et al. 2018. Persistence and space preemption explain species-specific founder effects

on the organization of marine sessile communities. - Ecology and Evolution 8:3430–3442.

Virnstein, R. W., and R. K. Howard. 1987. Motile epifauna of marine macrophytes in the Indian

River Lagoon, Florida. II. Comparisons between drift algae and three species of seagrasses.

Bulletin of Marine Science 41:13–26.

Vellend, M. 2010. Conceptual synthesis in community ecology. – Q. Rev. Biol. 85:183–206.

Ward, J. H. 1963. Hierarchical grouping to optimize an objective ounction. - Journal of the

American Statistical Association 58:236–244.

Waycott, M. et al. 2009. Accelerating loss of seagrasses across the globe threatens coastal

ecosystems. – Proc. Natl. Acad. Sci. 106: 12377-12381.

Wichmann, C. S. et al. 2012. Floating kelps in Patagonian Fjords: An important vehicle for

rafting invertebrates and its relevance for biogeography. – Mar. Biol. 159:2035–2049.

Williams, S. L., and K. L. Heck, Jr. 2001. Seagrass community ecology. Pages 317–337 in M. D.

Bertness, M. E. Hay, and S. D. Gaines, editors. Marine community ecology. Sinauer

Associates, Sunderland, Massachusetts, USA.

Whalen, M. A., et al. 2013. Temporal shifts in top-down vs. bottom-up control of epiphytic algae

in a seagrass ecosystem. - Ecology 94:510-520.

Whippo, R. et al. 2018. Epifaunal diversity patterns within and among seagrass meadows suggest

landscape-scale biodiversity processes. - Ecosphere 9:02490.10.1002/ecs2.2490.

Wlodarska-Kowalczuk, M. et al. 2014. Evidence of season-dependency in vegetation effects on

macrofauna in temperate seagrass meadows (Baltic Sea). - PLoS One 9:e100788.

Yamada et al. 2007. Temporal and spatial macrofaunal community changes along a salinity

gradient in seagrass meadows of Akkeshi-ko estuary and Akkeshi Bay, northern Japan. –

Hydrobiologia 592: 345-358.

Yamada, K. et al. 2014. Environmental and spatial controls of macroinvertebrate functional

assemblages in seagrass ecosystems along the Pacific coast of northern Japan. - Glob. Ecol.

Cons. 2:47–61.

Yeager, L. A. et al. 2019. Trait sensitivities to seagrass fragmentation across spatial scales shape

benthic community structure. – J. Anim. Ecol. 88:1743-1754.