1 **Beyond a single patch: local and regional processes explain diversity patterns in a seagrass**
2 **epifaunal metacommunity**
3

4 Keila A Stark[1]*, Patrick L Thompson[1], Jennifer Yakimishyn[2], Lynn Lee[3], Mary I O'Connor[1]
5
6
7 1. Department of Zoology
8 Biodiversity Research Centre
9 University of British Columbia
10 V6T 1Z4
11 Vancouver, BC, Canada
12
13 2. Pacific Rim National Park Reserve
14 P.O. Box 280
15 Ucluelet, BC, Canada
16
17 3. Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve, and
18 Haida Heritage Site
19 60 Second Beach Road
20 Skidegate, BC, Canada
21
22 ***Correponding author:** keilastark@zoology.ubc.ca

23
24
25 **Abstract:** 241 words
26 **Main text:** 4756 words
27 **Number of references:** 56
28 **Number of figures:** 3
29
30

31 **Data accessibility:** Data will be made available on Dryad should the manuscript be accepted. All
32 code for analysis can be found at https://github.com/keilast/HMS-Seagrass.
33
34 **Competing interests:** We have no competing interests.

Abstract

Ecological communities are jointly structured by dispersal, environmental niche filtering, and biotic interactions. Metacommunity ecology provides a framework for understanding how these processes combine to determine community composition in sites that are connected through dispersal. Here, we use the Hierarchical Modelling of Species Communities (HMSC) statistical framework to assess the contributions of metacommunity processes to community structure in an important coastal habitat, seagrass meadows, across approximately 1000 km along the coast of British Columbia. We parametrized a hierarchical joint species distribution model describing the relationship between environmental conditions, spatial distance between sites, and random effects on epifaunal invertebrate species distributions across thirteen eelgrass (*Zostera marina* (L.)) meadows. Compositional similarity did not decrease at large spatial distances suggesting sites are well-connected by dispersal, but observed species distribution patterns suggested dispersal rates are not so high that they homogenize regional diversity. We found evidence of niche filtering in several species along an abiotic gradient; however, environmental conditions only explained a small fraction of the overall variation in species distributions (22%), suggesting variable niche width among species and the contribution of processes other than niche fitering. Strong structuring in pairwise species co-occurrences led us to hypothesize the importance of priority effects and trophic interactions. This work has yielded refined hypotheses about the specific local and regional dynamics influencing seagrass epifaunal community structure, which in turn can inform targeted experiments, monitoring, and management decisions for preserving diversity in this important ecosystem.

Key words: Hierarchical Modelling of Species Communities, priority effects, dispersal, niche filtering, British Columbia, *Zostera marina*

Introduction

Understanding how local environmental conditions, biotic interactions, and regional connectivity jointly structure the composition of communities is a central challenge in ecology (Ricklefs and Schluter 2003, Vellend 2010, Leibold and Chase 2017). Determining the relative importance of these assembly processes is critical to understanding the spatial and temporal scales of stability and resilience in spatially structured communities (Economo et al. 2011). Metacommunity theory offers a promising framework for understanding drivers of community assembly across spatial scales; however, previous attempts to quantitatively apply metacommunity concepts to real-world systems have had limited success (Logue et al. 2011, Brown et al. 2017). Over the past decade and a half, the primary focus of metacommunity studies has been seeking in natural systems signatures of one of four idealized paradigms: the neutral model, patch-dynamics, species sorting, and mass effects (Leibold et al. 2004, Cottenie 2005). An emerging perspective holds that natural metacommunities often do not fit neatly under these paradigms because they do not reflect all possible combinations of the various assembly processes (Logue et al. 2011, Brown et al. 2017, Leibold and Chase 2017). Thus, the revised goal is to determine how dispersal, environmental conditions, and biotic interactions combine to structure species composition and abundance within metacommunities.

Many marine ecosystems that host high biodiversity occupy spatially structured habitats (e.g. coral reefs, kelp forests, seagrass meadows), and marine species have dispersing life histories that connect populations in distinct habitat patches. Marine systems are characterized by strong species interactions (Berlow 1999, Sala and Graham 2002), and the importance of dispersal-driven population dynamics has long been recognized (Levin and Paine 1974, Gaines and Roughgarden 1985). These combined roles of dispersal, the local environment, and biotic

interactions suggest that marine diversity patterns may reflect processes characteristic of

metacommunities (Guichard et al. 2003). Despite this, metacommunity dynamics have only been

explicitly investigated in a few marine systems (Guichard et al. 2003, Yamada et al. 2014,

Valanko et al. 2015).

Seagrasses, like kelps, corals and salt marsh plants, are foundation species that support high

productivity and faunal diversity (Orth et al. 1984, Duffy et al. 2015). Seagrasses form meadows

separated from each other by deeper water, un-vegetated seafloor, or other vegetated habitats,

and associated fauna disperse among meadows (Boström et al. 2006). This scenario is consistent

with the type of landscape that is governed by metacommunity dynamics (Leibold et al. 2004).

Yet, our understanding of the causes and conservation implications of seagrass-associated faunal

diversity patterns has not accounted for processes operating across numerous meadows and

adjacent habitats in a seascape. Local characteristics such as habitat structure (Orth et al. 1984),

primary biomass (Cébrian and Duarte 1998, Gullström et al. 2012), and trophic interactions

(Amunudrud et al. 2015, Huang et al. 2015) all affect faunal diversity, but the importance of

processes operating on a regional or inter-site scale remains unclear. Our limited understanding

of the importance of these regional processes could lead to underestimates of the amount of

habitat required to sustain biodiversity and ecosystem function (Boström et al. 2006).

To date, evidence for the importance of metacommunity dyamics in seagrass-associated

communities is based on two different quantitative approaches (Yamada et al. 2014, Whippo et

al. 2018). Using variation partitioning on constrained ordinations (Borcard et al. 1992, Cottenie

2005), Yamada et al. (2014) found that the relative contribution of space and environment varied

among invertebrate functional groups. However, the variation partitioning could not distinguish

between the possible metacommunity dynamics at play owing to the ambiguous statistical

significance of the results (Yamada et al. 2014). Simulation studies have shown that this method

can over- or under-estimate spatial and environmental components of variation, and produce

inflated R^2 values depending on the user's statistical choices (Gilbert and Bennett 2010,

Tuomisto et al. 2012). The other approach, known as elements of metacommunity structure

(EMS), seeks idealized structures in species occurrence patterns (Leibold and Mikkelson 2002,

Presley et al. 2010). While useful for visualizing species distribution patterns across a region,

EMS does not robustly elucidate the processes contributing to these patterns (Brown et al. 2017).

These limitations, along with the revised goal of individually assessing the contribution of

dispersal, niche filtering, and biotic interactions to diversity patterns, invite novel methodological

approaches. The Hierarchical Modelling of Species Communities statistical framework (HMSC,

Ovaskainen et al. 2017), shows promise for achieving this goal. HMSC uses Bayesian inference

to parametrize a hierarchical joint species distribution model, and allows the partitioning of

variation in species distributions due to the environment, spatial distance, and random effects,

allowing inference of patterns at the whole-community level as well as in individual species.

Additionally, it specifies a way to visualize positive and negative pairwise species co-

occurences, which can inform hypotheses on possible species interactions. For these reasons,

HMSC achieves what researchers aimed to achieve with the previous methods, and much more.

Here we test the hypothesis that seagrass-associated epifaunal invertebrate communities are

jointly structured by the spatial distances between meadows (a proxy for inferring dispersal

dynamics), local environmental conditions, and biotic interactions. We follow the HMSC

framework to answer the following questions: (1) Is the metacommunity characterized by

dispersal limitation, intermediate dispersal, or high dispersal? (2) Do species distribution patterns

suggest environmental niche filtering, and if so along which environmental axes? (3) Do species

co-variance patterns suggest possible biotic interactions influencing community assembly? The

order in which these questions are posed is significant. For example, evidence for dispersal

limitation would affect how we interpret observed species co-occurrence patterns.

We address these three questions with a biodiversity survey of invertebrates in 13 *Zostera*

marina (L.) meadows in British Columbia, Canada. We find evidence of intermediate dispersal

rates. We also find variable species responses to environmental conditions, and large overall

variation unexplained by environment and space in our model. Strong species' covariance

patterns led us to hypothesize possible biotic interactions influencing community structure.

Methods

Study sites and field sampling of eelgrass and associated epifauna

We sampled epifaunal biodiversity in 13 eelgrass *Zostera marina* (L.) meadows across a

1000 km spatial scale from late June through early August in 2017. The sites were located within

four coastal areas: Haida Gwaii within Gwaii Haanas National Park Reserve, National Marine

Conservation Area Reserve and Haida Heritage Site; southern Clayoquot Sound (one site was

within Pacific Rim National Park Reserve); Barkley Sound on the West coast of Vancouver

Island; and the Southern Gulf Islands within Gulf Islands National Park Reserve (Fig. 1a).

We sampled three sites in Barkley Sound (SA, DC, RB) with SCUBA because the

meadows reached a depth of 5 m. All other sites were sampled by wading or snorkeling at low

tide. We used a nested sampling regime with 0.25 m x 0.25 m quadrats to collect eelgrass,

epifaunal invertebrates, detritus, and macroalgae. Six quadrats were set in a 15 m x 30 m array

(Appendix S1: Fig. 1); the array was placed in the middle of the meadow to avoid edge effects

where possible. We removed all above-ground biomass (eelgrass, detritus, macroalgae) within

each quadrat by hand and immediately placed the contents into a mesh or ziploc bag.

Acquiring biotic and abiotic environmental variables

162   We acquired water quality and biotic data associated with hypothesized niche differences

among epifauna. The water quality variables included in-situ measurements of pH, salinity,

dissolved oxygen, nitrates and chlorophyll-*a* at each meadow, and averages acquired from Bio-

ORACLE, including surface chlorophyll-*a*, dissolved oxygen, nitrates, silicates, phosphates, pH,

salinity, and temperature. Bio-ORACLE is a database of marine environmental variables

gathered from several satellite and in-situ sources at a spatial resolution of 0.08° (Assis et al.

2018). All sites (meadows) were situated in different spatial cells, and thus had distinct values

for the aforementioned variables. We also calculated fetch, which quantifies site exposure.

170   For each quadrat sampled, we measured five biotic attributes summarizing habitat

structure and food availability: eelgrass leaf area index (LAI) and shoot density, and eelgrass dry

mass, algal dry mass, and detritus dry mass. To quantify LAI, we measured leaf length, width,

and blade number in five haphazardly-chosen shoots from each quadrat, and multiplied the

average blade area (m^2) per shoot by the shoot density (number of shoots per m^2). We separated

and dried eelgrass, detritus, and macroalgae in a desiccator oven (60° C for 48 hours).

Quantifying epifaunal diversity

177   We rinsed the eelgrass blades to remove epifaunal invertebrates, and passed samples

178   through a 500 μm sieve. Invertebrates were preserved in 95% EtOH for identification with light

179   microscopy (10x magnification). We identified epifauna to the lowest taxonomic level possible

180   using keys in the Light and Smith manual (Carlton 2007) and Kozloff (1996).

181   *Analysis*

182   We followed the methodology of Ovaskainen et al. (2017) to construct a hierarchical

joint species distribution model describing the relationship between environmental variables,

spatial distance, species co-occurrences and species distributions using the HMSC package

(Guillaume Blanchet et al. 2018) in R (version 3.4.2; R Development Core Team 2014). The

model is composed of two linear predictor terms: one describing fixed effects of environment

and space, and one describing random effects that cannot be attributed to environment or space

(Equations 2 & 3 in Ovaskainen et al. 2017). The random-effects linear predictor term describes

random variation in individual species' distributions as well as covariation for every species pair.

We used Markov chain Monte Carlo to parametrize the model. Species' responses to

environmental variables were assumed to follow a multivariate Gaussian distribution. We

assumed a Poisson distribution for species count data and used the same priors as in the

examples in Ovaskainen et al. (2017). The Markov Chain included 100 000 iterations, a burn-in

length of 1000 iterations, and thinned to retain every 100^{th} sample of the posterior distributions.

We confirmed that parameters were well mixed by visually inspecting trace plots. We tested the

model's performance by calculating explained deviance (D^2_{adj}) as described in Appendix S1: Eq.

S1. D^2_{adj} is comparable to R^2_{adj} in least square regression analysis (Guisan and Zimmermann

2000); in this context, it describes the model fit where a value of 1 means the model perfectly

predicts the observed data. Our calculations yielded a D^2_{adj} value for each individual species

distribution model. The mean D^2_{adj} across all species gave a metacommunity-level value of

predictive power.

To answer our three main questions about the various metacommunity processes, we

visualized the parametrized model in several ways (Appendix S1: Table S1). We investigated

three possible dispersal conditions: dispersal limitation, intermediate dispersal, and high

dispersal. To investigate dispersal limitation at large spatial scales, we plotted distance-decay of

similarity by randomly drawing 200 pairwise distances between sampling plots, and plotting

predicted community similarity against distance. Distance-decay of similarity could reflect

dispersal limitation due to prohibitively far travel distances, or differences in environmental

conditions with increasing spatial scale. To distinguish these possibilities, we compared

predicted community similarity as a function of distance and environmental covariates (the full

HSMC model), and predicted community similarity using an alternative model in which the

effect of the environment was removed (all environmental covariates were set to their mean

values). To distinguish between intermediate and high dispersal, we visually inspected the

occurrence data for signatures of regional homogenization by high dispersal. We used niche

filtering strength (estimated by D^2_{adj}) as another line of evidence to distinguish intermediate and

high dispersal: theory predicts that intermediate dispersal (dispersal sufficiency) allows species

to colonize sites where local environmental conditions are optimal for growth and reproduction,

whereas high dispersal (dispersal surplus) overrides this (Mouquet and Loreau 2003).

To estimate niche filtering strength, we conducted variation partitioning of the fixed and

random effects (Ovaskainen et al. 2017). This summarized the relative contribution of

environmental variables, spatial distance, and random effects to each species distribution model;

we visualized this in a stacked bar plot. We then estimated the pure and shared fractions of

variation explained by environment and space, as well as the unexplained fraction using type III

sum of squares (Borcard et al. 1992), but using the HMSC model instead of constrained

ordination. To do so, we followed the equations in Appendix S1: Eq. S2.

To identify species co-occurrence patterns resulting from possible biotic interactions, we

represented modelled pairwise associations in a correlation matrix describing the extent to which

species pairs co-occurred more negatively or positively than by random chance, controlling for

the effect of the environment and space. We then conducted hierarchical cluster analysis on sites

according to predicted species abundances from the model using Ward's criterion (Ward 1963).

We mapped the cluster dendrogram onto an abundance heatmap of the 20 species in our study

with the highest mean abundance, and colour coded the cells according to the species co-

occurrence groupings (groups of species that tended to positively co-occur amongst themselves).

Full versions of these figures with all species are provided in Appendix S1: Fig. S4 and Fig. S5.

Results

Species abundance and distribution patterns

We counted and identified 39 644 individuals representing at least 58 species across the

region (Appendix 1: Table S2). The represented taxa span multiple phyla, diet types (herbivores,

detritivores, suspension feeders, carnivores), and life history strategies (brooding, broadcast

spawning). Mean quadrat-level species richness was 13 species, and mean site-level richness was

16 species. A full summary of diversity metrics can be found in Appendix S1: Table S3.

Several taxa were present at all sites (Appendix S1: Fig. S2). These include harpacticoid

copepod *Porcellidium* sp., bivalves *Saxidomus gigantea* and *Clinocardium nutallii*, snails

Alvania compacta, *Lacuna* spp., and the tanaid *Leptochelia* sp. Genera *Mytilus* (a mussel) and

Nereis (a polychaete worm) were also represented at all sites, however we were unable to

confidently identify these organisms to species. Ten species were present in at least one site in

every region: pycnogonid *Anoplodactylus viridintestinalis,* isopod *Idotea resecata,* gammarid

amphipods *Monocorophium insidiosum, Ampithoe valida, Pontogeneia rostrata, Ampithoe dalli,*

Aoroides spp.*,* snails *Amphissa columbiana* and *Alia carinata,* and limpet *Lottia pelta*.

Dispersal: Distance-decay of community similarity due to spatial distance and environment

In the full HMSC model summarizing species' responses to environment and spatial

distance, predicted community similarity decreased with increasing distance between sites (Fig.

1b, red points and line). However, community similarity showed a slight increase with distance

when the effect of environmental variables was removed from the model (all environmental

variables set to their mean value), suggesting that spatial distance alone did not correspond with

differences in invertebrate compositional similarity (Fig. 1b, blue points and line).

Environmental niche filtering: variation partitioning

Patterns in abundance and distribution were consistent with weak environmental niche

filtering along an abiotic axis; however, the importance of environmental conditions varied

across species. The relative contributions of all 14 environmental variables, spatial distance and

random effects in the HMSC model can be seen in Fig. 2. Out of all environmental variables, the

water quality variables (blue bars in Fig. 2) explained the largest proportion of variation across

nearly all species distributions. Herbivore food availability (seagrass, algae and detritus biomass)

and habitat structure (eelgrass shoot density and LAI) had smaller influences on species

distributions (Fig. 2). While the mean deviance explained (D^2_{adj}) across all species distribution

models was 0.29, individual values varied (0 to 0.82, Appendix 1: Table S2), implying high

variation in model fit. Accounting for this, the true fraction of variation explained by

environment was 22%, space explained -3%, the shared fraction between environment and space

was 4%, and the residual fraction of variation was 76% (Appendix S1: Fig. S3).

Species interactions: Patterns in pairwise species co-occurrences

We identified, post hoc, three main site-level species co-occurrence groupings after

controlling for fixed effects. Members of the first group, including *Nereis* sp. through *Caprella*

laeviuscula along the vertical axis in Fig. 3a, tended to positively co-occur with each other, and

negatively co-occur members of the second group. This first group is henceforth known as the

Nereis group. The second group, henceforth known as the *C. californica* group, included

Caprella californica through *Ampithoe valida,* and Harpacticoid 2 through *Lacuna* spp. in Fig.

3a. *Alvania compacta* and *Lacuna* spp. showed co-occurrence patterns consistent with the *C.*

californica group, but their co-occurrence values were generally weaker. The third group was

comprised of species that did not show strong positive or negative co-occurrence values with

others; *Spirorbis* sp. was the only representative in this category among the 20 most abundant

species.

282   Species from each of the three co-occurrence groupings were present at all sites,

indicating that members from these groups can be simultaneously present (Fig. 3b). However,

the abundances of members from different co-occurrence groups tended to negatively co-vary; at

sites JB, HL, CB, RA, EB, DC, and DK, members of the *Nereis* group were predicted to be most

abundant, whereas *C. californica* group members were predicted to be most abundant at SS, SA,

LH, GB and RB. The site IN was predicted to be strongly dominated by *Spirorbis* sp. The

emergent species groupings identified in Fig. 3a do not clearly correspond with spatial distance

because the cluster analysis did not group sites by spatial proximity (black symbols in Fig. 3b).

Discussion

291   By applying a new statistical framework, we have gained new insight into the possible

metacommunity processes structuring seagrass-associated epifaunal diversity in coastal British

Columbia. We report evidence that both biotic interactions and environmental conditions

influence seagrass meadow invertebrate composition. Spatial patterns of species abundance and

distribution suggest regional-scale dispersal patterns are possible, such that even distant

meadows may host the same species. However, these dispersal rates are clearly not so high as to

overwhelm the local meadow scale effects of biotic interactions and environmental conditions.

Below, we review these conclusions and the evidence to support them.

Epifaunal communities may be well-connected and experience intermediate dispersal rates

Epifaunal communities appear to be well-connected by dispersal at the present spatial

scale and seagrass habitat distribution in coastal British Columbia. This is true for the seven

species that were present at all sites, and the additional ten species that were not present at every

site but present in all regions (Appendix S1: Fig. S2). These widely distributed groups had

representation from across phyla and life history traits, suggesting that no single taxonomic

group or dispersal life history strategy had a consistently greater spatial distribution. Many of the

meadows were small (< 1 ha), and epifaunal abundances can vary substantially within meadows

from year to year such that local extinctions followed by rescue from nearby populations are

likely important (Thom et al. 1995). However, the specific pathway the organisms travelled to

arrive at these sites remains unknown. We used Euclidean distances in our analysis, which

preserved the rank order of distances between sites, but actual distances are likely greater due to

oceanographic circulation patterns (Kinlan and Gaines 2003, Mitarai et al. 2008).

It is unclear whether distant sites share the same species because they are 1) linked by

direct dispersal; 2) indirectly linked by dispersal via unsampled sites; or 3) were colonized by

populations of the same species in a historical dispersal event, but have not seen the exchange of

individuals since. Seagrass-associated epifauna generally disperse passively between meadows,

either by larval transport in currents or by "rafting" on floating pieces of seagrass or macroalgae.

Rafts have been observed hosting benthic and epifaunal invertebrates across phyla (echinoderms;

peracarids; molluscs; annelids), and contribute to connectivity between coastal ecosystems

across 100s of kilometres (Wichmann et al. 2012). Thus, while incidences of direct dispersal

between the farthest sites may be rare, they are not impossible.

Species that were only found at one site or region (e.g. *Jassa marmorata,* Appendix S1:

Fig. S2) may be dispersal limited for reasons other than prohibitively long travel distances.

Those with low abundances (less than five individuals per site) may not disperse in appreciable

numbers, and therefore cannot establish populations at all sites as readily as abundant species.

This is supported by our observation of a generally positive relationship between range size (the

number of sites a species was represented at) and abundance (Appendix 1: Fig. S6).

327   We can also conclude that dispersal rates are high enough for several species to be

present at most sites, but not so high as to completely overwhelm the effects of environmental

conditions. Thus, dispersal rates are likely intermediate on average. A subset of species was

present at all sites, but often at varying abundances (Fig 3b). This may be indicative of weak

mass effects, wherein immigration allows populations to persist even if local conditions are not

optimal (Schmida and Wilson 1985, Mouquet and Loreau 2003). Metacommunity theory

predicts that such weak mass effects are likely to be found whenever dispersal rates are not

limiting (Thompson et al. 2017), particularly in organisms that cannot control their own dispersal

(Leibold and Chase 2017) such as seagrass-associated epifauna. This may be supported by our

result on the contribution of the environment, particularly in ubiquitous species for which the

model fit was poor (e.g. *Aoroides* spp., *Spirorbis* sp., $D^2_{adj} = 0$); the poor model fit may indicate

that these persist at sites even if local conditions are not optimal, due to sufficient immigration.

An alternative explanation for the model's poor predictive power for these species is that these

taxa have broader environmental niches, and thus the regional environmental gradient is not

sufficient to drive clear presence-absence patterns.

Seagrass epifauna show varying degrees of environmental niche filtering

343   The abiotic environmental axis had the largest influence on species distributions, with

phosphates, salinity, and sea surface temperature explaining the largest fractions of variation on

average in the joint species distribution model (Fig. 2). However, after accounting for the model

fit, the total variation explained by the environment was small (22%). In some species for which

the model had high predictive power (e.g. *Ampithoe dalli* and *Caprella natalensis*, Appendix 1:

Table S2*)*, niche filtering may be important, but overall there remains a large amount of variation

unexplained by the environment.

Environmental heterogeneity may influence species distributions and abundances across

the metacommunity in one of three ways. First, the most important abiotic environmental

parameters may influence food availability through primary productivity. This is possible despite

the fact that primary biomass explained less variation than abiotic parameters (Figure 2). While

the primary biomass measurements were taken at a single time point, the abiotic parameters

represent a four-year average (Assis et al. 2018) and may therefore better represent the

conditions these organisms experience over their lifetimes. Second, sea surface temperature,

dissolved oxygen, and salinity may affect species' distributions by via environmental tolerance

ranges. For example, isopod *Idotea baltica* (a relative of *I. resecata* and *I. montereyensis* in our

study) experiences significantly lower survival in prolonged exposure to higher temperatures and

lower salinities than those experienced in their home habitat (Rugiu et al. 2018). Third,

temperature may influence epifaunal metabolic demands (many of which are herbivores),

thereby influencing grazing rates and altering the available primary biomass (O'Connor 2009).

Decreased food resources may increase competition, thus influencing the number and identity of

species a seagrass patch can host. Determining whether individual species' responses to

competitive stress influence whole-community structure may provide key information on how

the environment and species interactions combine to influence community assembly.

Environmental conditions in temperate eelgrass meadows fluctuate seasonally, driving

changes in epifaunal community structure throughout the year (Wlodarska-Kowalczuk et al.

2014). Because these species may not show the same distributions year-round, sampling several

times in a year to capture these temporal changes would give a clearer picture of species'

responses to all metacommunity processes.

Species co-variance patterns may suggest priority effects and trophic interactions

Three main co-occurrence groupings emerged showing antagonistic abundances across

sites (Fig. 3b). Stochasticity alone is unlikely to have produced such a strong pattern, especially

given deterministic processes tend to be more important in systems that are not dispersal limited

(Lee and Bruno 2009, Ron et al. 2018). While there are multiple possible explanations for non-

random species co-occurrence patterns (Connor and Simberloff 1979), space and environmental

heterogeneity were ruled out as co-occurrence values represented pairwise covariation

unexplained by environment and space (Ovaskainen et al. 2017). This, along with the high

unexplained fraction in the variation partitioning leads us to suggest the importance of biotic

processes. Several types of interactions may be occurring in the metacommunity; the majority of

species are herbivores or detritivores, and thus may compete for primary biomass. Laboratory

experiments have shown that behaviours such as grazing rate and habitat selection in amphipods

are altered in the presence of morphologically and functionally similar intrerspecific competitors

(Beermann et al. 2018). Other species in our study are filter feeders, and thus may not be

competing for food resources but possibly predator-free space or substrate on the eelgrass blades.

The community structure of a given site with certain local environmental conditions and

access to a certain regional pool may have multiple possible compositional states depending on

the arrival order of species; this phenomenon is known as priority effects (Fukami et al. 2016, Ke

and Letten 2018) and has been documented in marine fouling communities (Vieira et al. 2018).

The antagonistic co-occurrence groupings in our analysis may suggest priority effects (Fig. 3a

and 3b). We hypothesize that following the establishment of a new meadow or a disturbance

event, the first few species to colonize a seagrass meadow determine the success of subsequent

colonizers, regardless of local environmental conditions. As an illustrative example, *Lacuna* spp.

might have been among the first to colonize sites JB, HL, GB, and CB (Fig. 3b), thereby gaining

a numerical advantage over species in the alternative cohort. Distibution patterns in the sessile

calcifying polychaete *Spirorbis* sp. may also indicate priority effects. While it did not strongly

negatively or positively co-occur with other species, it dominated the HL site; 16 300 individuals

were found, several orders of magnitude higher than its abundance elsewhere (Data S1). Site HL

also had the lowest species richness of all 13 sites (9 species). Eelgrass shoots there were

completely covered by these *Spirorbis* sp., possibly preventing grazers in the community from

accessing epiphytic algae or the seagrass itself. The age, colonization history, and disturbance

regime of the meadows likely matter if priority effects are occurring.

The negatively correlated abundance patterns could also reflect the presence of a predator

such as *Nereis* at some sites and not others (Fig. 3b). Field experiments have shown that changes

in predation pressure by fish, shorebirds, and predatory invertebrates can shift seagrass-

associated epifaunal assemblages in a matter of weeks (Amundrud et al. 2015; Huang et al.

2015). Differential predation pressure could also lead to the pattern of diverging community

composition that we have witnessed (Guzman et al. 2018). A more comprehensive sampling

effort of fish and bird predators would be required to determine the degree to which trophic

interactions structure these communities.

Positive co-occurrence patterns between certain species may additionally be a result of

positive biotic interactions. An example is seen in *Orchomenella recondita* - a gammarid

amphipod that lives in the gastrovascular cavity of the anemone *Anthopleura elegans* (Carlton

2007). This species was only recorded at the SS site, and specifically only found in quadrats

where *A. elegans* was collected with the seagrass shoots.

Conclusion

Using a new statistical approach to characterizing dispersal, environmental niche

filtering, and biotic interactions, we have developed refined hypotheses about the

metacommunity processes driving temperate seagrass epifaunal diversity on a regional scale. Our

study also gives a baseline spatial scale (1000 km) for comparing the contribution of these

processes across spatial scales and other ecological systems (Viana and Chase 2018).

Manipulative experiments and monitoring are now needed to test our refined hypotheses. One

could test the hypothesis that species sorting is relatively weak by measuring fitness differences

along the most important environmental axes. Given our hypothesis on the importance of priority

effects, one might experimentally manipulate the colonization order of key species in our study

to determine if resulting compositional states differ (Fukami et al. 2016). Testing our hypothesis

that populations within the metacommunity are not dispersal limited is more challenging, but

could be done using particle tracking models similar to those used to estimate larval dispersal

between coral reefs across thousands of kilometres (Treml et al. 2008). A population genetic

analysis could also ground-truth the extent of dispersal between allopatric populations. All

together, we suggest that a modern metacommunity perspective combined with new advances in

statistical methods provide a powerful and relevant framework for studying natural ecosystems.

Our findings are particularly relevant to conservation initiatives that aim to maintain

diversity in seagrass meadows; identifying the importance of local and regional processes allows

the conservation practitioner to better predict how habitat fragmentation and climate change

might influence diversity. Seagrass epifaunal communities may be quite resilient, given our

findings that environmental conditions minimally contribute to species distributions, and several

species across functional groups are consistently present across a fairly large spatial scale.

Acknowledgements

KAS conceived of the study with guidance from MIO. JY provided samples from IN, DK, and

EB. LL provided samples from RA and HL. KAS collected and processed samples, analyzed

data, and wrote the first manuscript draft. PLT provided recommendations for the analysis. KAS,

PLT, and MIO contributed to writing the manuscript. This research was financially supported by

a UBC Zoology SURE Grant, and NSERC Discovery Grants, Canadian Foundation for

Innovation (CFI), and is sponsored by the NSERC Canadian Healthy Oceans Network and its

Partners: Department of Fisheries and Oceans Canada and INREST (representing the Port of

Sept-Îles and City of Sept-Îles). PLT is supported by NSERC and Killam postdoctoral

fellowships. We are grateful to Duarte Viana for providing critical recommendations and code

for the D^2_{adj} and variation partitioning analyses. Matt Whalen pulled the Bio-ORACLE dataset,

and John Cristiani made the map in Fig. 1a. Special thanks goes to Michelle Paleczny and Gulf

Islands National Park Reserve for boat and laboratory space. We acknowledge staff, students,

and volunteers from Pacific Rim National Park Reserve for their assistance collecting eelgrass in

Clayoquot Sound. We thank Rob Underhill, Abbie Sherwood, Emily Adamczyk, Minako Ito,

Ariane Comeau, Giordano Bua for assistance with sample collection, processing, and ID, and

Coreen Forbes, Matt Whalen, and Jeff Groh for feedback on the manuscript.

Literature Cited

Amundrud, S. L., D. S. Srivastava, and M. I. O'Connor. 2015. Indirect effects of predators

control herbivore richness and abundance in a benthic eelgrass (*Zostera marina*)

mesograzer community. Journal of Animal Ecology 84: 1092–1102.

Assis J., L. Tyberghein, S. Bosch, H. Verbruggen, E. A. Serrão, and O. De Clerck. 2018. Bio-

ORACLE v2.0: Extending marine data layers for bioclimatic modelling. Global Ecology

and Biogeography 27:277–284.

Beermann, J., K. Boos, L. Gutow, M. Boersma, and A. C. Peralta. 2018. Combined effects of

predator cues and competition define habitat choice and food consumption of amphipod

mesograzers. Oecologia 186:645–654.

Berlow, E. L. 1999. Strong effects of weak interactions in ecological communities. Nature 398:

330–334.

Boström, C., E. L. Jackson, E. L., and C. A. Simenstad. 2006. Seagrass landscapes and their

effects on associated fauna: A review. Estuarine, Coastal and Shelf Science 68: 383-403.

Borcard, D., P. Legendre, and P. Drapeau. 1992. Partialling out the spatial component of

ecological variation. Ecology 73:1045–1055.

Brown, B. L., E. R. Sokol, J. Skelton, and B. Tornwall. 2017. Making sense of

metacommunities: dispelling the mythology of a metacommunity typology. Oecologia

183:643–652.

Carlton, J. 2007. The Light and Smith Manual: Intertidal invertebrates from Southern California

to Oregon. Fourth edition. University of California Press, Oakland, California, USA.

Cébrian, J. and C. M. Duarte. 1998. Patterns in leaf herbivory on seagrasses. Aquatic Botany

60:67–82.

Connor, E. F., and D. Simberloff. 1979. The assembly of species communities: Chance or

competition? Ecology 60:1132–1140.

Cottenie, K. 2005. Integrating environmental and spatial processes in ecological community

dynamics. Ecology Letters 8:1175–1182.

Duffy J. E, et al. 2015. Biodiversity mediates top-down control in eelgrass ecosystems: A global

comparative-experimental approach. Ecology Letters 18:696–705.

Economo, E. P. 2011. Biodiversity conservation in metacommunity networks: Linking pattern

and persistence. The American Naturalist 177:167-180.

Fukami, T., E. A. Mordecai, and A. Ostling. 2016. A framework for priority effects. Journal of

Vegetation Science 27:655–657.

Gaines, S., and J. Roughgarden. 1985. Larval settlement rate: A leading determinant of structure

in an ecological community of the marine intertidal zone. Proceedings of the National

Academy of Sciences 82:3707–-3711.

Gilbert, B, and J. R. Bennett. 2010. Partitioning variation in ecological communities: do the

numbers add up? Journal of Applied Ecology 47:1071–1082.

Guichard, F. 2003. Mussel disturbance dynamics: Signatures of oceanographic forcing from

local interactions. The American Naturalist 161:889–-904.

Guillaume Blanchet, F., with contributions from G. G. Tikhonov and A. A. Norberg. 2018.

HMSC: Hierarchical Modelling of Species Community. R package version. 2.1-2.

Guisan, A., and N. E. Zimmermann. 2000. Predictive habitat distribution models in ecology.

Ecological Modelling 145:147–186.

Gullström, M., S. Baden, and M. Lindegarth. 2012. Spatial patterns and environmental correlates

in leaf-associated epifaunal assemblages of temperate seagrass (*Zostera marina*) meadows.

Marine Biology 159:413–425.

Guzman, L. M., R. M. Germain, C. Forbes, S. Straus, M. I. O'Connor, D. Gravel, D. S.

Srivastava, and P. L. Thompson. 2018. Towards a multi-trophic extension of

metacommunity ecology. In press. Ecology Letters doi:10.1111/ele.13162.

Huang A. C., M. Essak, and M. I. O'Connor. 2015. Top-down control by great blue herons *Ardea*

herodias regulates seagrass-associated epifauna. Oikos 124:1492–1501.

Ke, P. J., and A. D. Letten. 2018. Coexistence theory and the frequency-dependence of priority

effects. Nature Ecology and Evolution 2:1691–1695.1–7.

Kinlan, B. P., and S. D. Gaines. 2003. Propagule dispersal in marine and terrestrial

environments: A community perspective. Ecology 84:2007–2020.

Kozloff, E. 1996. Marine Invertebrates of the Pacific Northwest. University of Washington

Press, Seattle, Washington, USA.

Leibold, M. A., et al. 2004. The metacommunity concept: A framework for multi-scale

community ecology. Ecology Letters 7:601–613.

Leibold, M. A., and J. M. Chase. 2017. Metacommunity Ecology, Volume 59. Princeton

University Press, Princeton, New Jersey, USA.

Leibold, M.A., and G. M. Mikkelson. 2002. Species turnover, and boundary clumping: Elements

of meta-community structure. Oikos 97:237–250.

Lee, S. C, and J. F. Bruno. 2009. Propagule supply controls grazer community structure and

primary production in a benthic marine ecosystem. Proceedings of the National Academy

of Sciences 106:7052–7057.

Levin, S. A., and R. T. Paine. 1974. Disturbance, patch formation, and community

structure. Proceedings of the National Academy of Sciences 71:2744–2747.

Logue, J., et al. 2011. Empirical approaches to metacommunities: a review and comparison with

theory. Trends in Ecology and Evolution 26:482–491.

Mitarai, S., D. A. Siegel, and K. B. Winters. 2008. A numerical study of stochastic larval

settlement in the California Current system. Journal of Marine Systems 69:295–309.

Mouquet, N., and M. Loreau. 2003. Community patterns in source-sink metacommunities. The

American Naturalist 162:554–557.

O'Connor, M. I. 2009. Warming strengthens an herbivore-plant interaction. Ecology 90:388–

398.

Orth, R. J., K. L. Heck, and J. van Montfrans. 1984. Faunal communities in seagrass beds: A

review of the influence of plant structure and prey characteristics on predator-prey

relationships. Estuaries 7:339–350.

Ovaskainen, O., G. Tikhono, A. Norberg, F. Guillaume Blanchet, L. Duan, D. Dunson, T.

Roslin, and N. Abrego. 2017. How to make more out of community data? A conceptual

framework and its implementation as models and software. Ecology Letters 20:561–576.

Presley, S.J., C. L. Higgins, and M. R. Willig. 2010. A comprehensive framework for the

evaluation of metacommunity structure. Global Ecology and Evolution 19:908–917.

R Development Core Team. 2014. R: a language and environment for statistical computing. R

Foundation for Statistical Computing. Vienna, Austria. http://www.R-project.org.

Ricklefs, R. E., and D. Schluter. 1993. Species Diversity in Ecological Communities: Historical

and Geographical Perspectives. University of Chicago Press, Chicago, Illinois, USA.

Ron, R., O. Fragman-Sapir, and R. Kadmon. 2018. Dispersal increases ecological selection by

increasing effective community size. Proceedings of the National Academy of Sciences

115:11280–11285.

Rugiu, L., I. Manninen, J., Sjöroos, and V. Jormalainen. 2018. Variations in tolerance to climate

change in a key littoral herbivore. Marine Biology 165:1–11.

Sala, E., and M. H. Graham. 2002. Community-wide distribution of predator-prey interaction

strength in kelp forests. Proceedings of the National Academy of Sciences 99:3678–3683.

Schmida, A., and M. V. Wilson. 1985. Biological determinants of species diversity. Journal of

Biogeography 12:1–20.

Thom, R., B. Miller, and M. Kennedy. 1995. Temporal patterns of grazers and vegetation in a

temperate seagrass system. Aquatic Botany 50:201–205.

Thompson, P. L., B. Rayfield, and A. Gonzalez. 2017. Loss of habitat and connectivity erodes

species diversity, ecosystem functioning, and stability in metacommunity networks.

Ecography 40:98–108.

Treml, E. A., P. N. Halpin, D. L. Urban, and L. F. Pratson. 2008. Modeling population

connectivity by ocean currents: a graph-theoretic approach for marine conservation.

Landscape Ecology 23:19–36.

Tuomisto, H., L. Ruokolainen, and K. Ruokolainen. 2012. Modelling niche and neutral

dynamics: on the ecological interpretation of variation partitioning. Ecography 35:961–

971.

Valanko, S., J. Heino, J. M. Westerbom, M. Viitasalo, and A. Norkko. 2015. Complex

metacommunity structure for benthic invertebrates in a low-diversity coastal system.

Ecology and Evolution 5:5203–5215.

Viana, D. S., and J. M. Chase. 2018. Spatial scale modulates the inference of metacommunity

assembly processes. In press. Ecology doi:10.1002/ecy.2576.

Vieira, E. A., A. A. V. Flores, and G. M. Dias. 2018. Persistence and space preemption explain

species-specifc founder effects on the organization of marine sessile communities. Ecology

and Evolution 8:3430–3442.

Vellend, M. 2010. Conceptual synthesis in community ecology. The Quarterly Review of

Biology, 85:183–206.

Ward, J. H. 1963. Hierarchical grouping to optimize an objective ounction. Journal of the

American Statistical Association 58:236–244.

Wichmann, C. S., I. A. Hinojosa, and M. Thiel. 2012. Floating kelps in Patagonian Fjords: An

important vehicle for rafting invertebrates and its relevance for biogeography. Marine

Biology 159:2035–2049.

Whippo, R., N. S. Knight, C. Prentice, J. Cristiani, M. R. Siegle, and M. I. O'Connor. 2018.

Epifaunal diversity patterns within and among seagrass meadows suggest landscape-scale

biodiversity processes. Ecosphere 9:02490.10.1002/ecs2.2490.

Wlodarska-Kowalczuk, M., E. Jankowska, E. L. Kotwicki, and P. Balazy. 2014. Evidence of

season-dependency in vegetation effects on macrofauna in temperate seagrass meadows

(Baltic Sea). PLoS One 9:e100788.

Yamada, K., Y. Tanaka, T. Era, and M. Nakaoka. 2014. Environmental and spatial controls of

macroinvertebrate functional assemblages in seagrass ecosystems along the Pacific coast of

northern Japan. Global Ecology and Conservation 2:47–61.

 Figure legends

Figure 1. a) Map of coastal British Columbia showing eelgrass field sites in Haida Gwaii (HL,

RA), Clayoquot Sound (DK, EB, IN), Barkley Sound (SA, DC, RB), and the Southern Gulf

Islands (GB, JB, CB, LH, SS). **b)** HMSC model predictions of community similarity relative to

the spatial distance (Euclidean) between communities. The red circles show similarity

predictions based on the full model. The blue triangles show predictions wherein the effect of all

environmental variables has been removed (standardized to their mean values).

Figure 2. Variation partitioning of fixed and random effects across all HMSC species

distribution models (numbers correspond to species names in Appendix 1: Table S2). Black cells

represent the contribution of spatial distance. Purple cells represent random effects. Blue cells

represent the contribution of water quality parameters. Yellow cells represent food availability

(eelgrass, algae and detritus biomass). Green cells represent habitat structure, which is the sum of

variation explained by eelgrass LAI and shoot density. Bars are organized according to broad

taxonomic groupings as shown in Appendix 1: Table S2. The percentages next to the legend

labels indicate mean variation explained by that variable across all species. Adjusted deviance

explained (D^2_{adj}) varied (Appendix 1: Table S2). The mean D^2_{adj} across all species was 0.29.

Figure 3. a.) A correlation plot showing modelled site-level pairwise co-occurrences in the 20

most abundant species. Purple cells represent positively co-occurring species pairs, and turquoise

cells represent negatively co-occurring species pairs. Species names are ordered according to the

output of hierarchical clustering with Ward's criterion on pairwise co-occurrence values. A

version of this figure with all 58 species is shown in Appendix S1: Fig. S4. **b)** Heatmap and

cluster dendrogram depicting modelled species abundance and compositional similarity across

sites. The heat map shows the twenty species with the highest predicted mean proportional

abundances ordered from highest to lowest. Cell colours correspond to the HMSC co-occurrence

groups shown in Fig. 3a: turquoise or *Nereis* group, purple or *C. californica* group, and grey

Spirorbis group. Cell shade strength indicates proportional abundance at a given site (darker

means higher abundance). The symbols to the left of the site abbreviations indicate region

membership; circles show Haida Gwaii sites, squares show Barkley Sound sites, diamonds show

Clayoquot Sound sites, and triangles show Southern Gulf Islands sites. A version of this figure

with all 58 species is shown in Appendix S1: Fig. S5.

Figure 1

Figure 2.

 Figure 3

 Appendix S1: Supporting Information

630 **Table S1.** Summary of evidence used to answer questions about metacommunity processes

Question	Analyses	Figure number
(1) Is the metacommunity characterized by dispersal limitation, intermediate dispersal, or high dispersal?	Dispersal limitation: Distance-decay of community similarity from HMSC joint species distribution models with and without accounting for environmental similarity	Fig. 1b
	Intermediate versus high dispersal: Biodiversity patterns across sites for signatures of high dispersal (compositional homogenization); HMSC model results indicating structured co-occurrence patterns and variation in composition explained by environmental	Fig. 3a, 3b
(2) Do species distribution patterns suggest environmental niche filtering, and if so along which environmental axis?	Variation partitioning of modelled site and region-level fixed and random effects of environmental covariates and calculating predictive power (proportion of deviance) of the fixed effects model	Fig. 2
(3) Do species co-variance patterns suggest possible biotic interactions influencing community assembly?	Correlation matrix of pairwise species co-occurences, after controlling for the effect of space and environment	Fig. 3a, 3b

633 **Table S2.** Species names, taxonomic groupings, and proportion of deviance (predictive power).

634 Numbers correspond to those in Figure 2. The top twenty species with highest predicted

635 abundances according to the HMSC model are bolded.

Number	Species name	Broad taxonomic group	Adjusted deviance explained for the HMSC model (D^2_{adj})
1	*Alia carinata*	Gastropod (snail)	0.15
2	***Alvania compacta***	**Gastropod (snail)**	**0.4**
3	*Amphissa columbiana*	Gastropod (snail)	0
4	*Crepidula* sp.	Gastropod (snail)	0.64
5	*Euspira lewisii*	Gastropod (snail)	0.46
6	*Harmothoe imbricata*	Gastropod (snail)	0
7	***Lacuna* spp. (*L. variegata* and *L. vincta*)**	**Gastropod (snail)**	**0.45**
8	*Lirularia parcipincta*	Gastropod (snail)	0.46
9	*Lirularia* sp.	Gastropod (snail)	0
10	*Littorina* sp.	Gastropod (snail)	0.5
11	*Margarites pupillus*	Gastropod (snail)	0.45
12	*Lottia pelta*	Gastropod (limpet)	0.53
13	***Clinocardium nuttallii***	**Bivalve**	**0**
14	***Mytilus* sp.**	**Bivalve**	**0.28**
15	*Saxidomus gigantea*	Bivalve	0
16	*Unknown clam 1*	Bivalve	0.68
17	*Unknown clam 2*	Bivalve	0.35
18	*Dorvillea longicornis*	Polychaete	0.52
19	*Exogone sp.*	Polychaete	0.25
20	*Glycinde armigera*	Polychaete	0.05
21	*Lepidonotus squamatus*	Polychaete	0.55
22	*Neoamphitrite robusta*	Polychaete	0.22
23	*Unknown polychaete (Family Oenoidae)*	Polychaete	0.15
24	*Unknown polychaete (Family Opheliidae)*	Polychaete	0.36
25	***Spirorbis* sp.**	**Polychaete**	**0**
26	***Nereis* sp.**	**Polychaete**	**0.55**
27	***Ampithoe dalli***	**Gammarid amphipod**	**0.47**

28	*Ampithoe lacertosa*	Gammarid amphipod	0.34
29	***Ampithoe valida***	**Gammarid amphipod**	**0**
30	***Aoroides* spp.**	**Gammarid amphipod**	**0**
31	*Ceradocus spinicauda*	Gammarid amphipod	0
32	*Grandidierella japonica*	Gammarid amphipod	0
33	*Ischyocerus anguipes*	Gammarid amphipod	0.33
34	*Jassa marmorata*	Gammarid amphipod	0
35	***Monocorophium insidiosum***	**Gammarid amphipod**	**0.23**
36	*Orchomenella recondita*	Gammarid amphipod	0.73
37	***Photis brevipes***	**Gammarid amphipod**	**0**
38	***Pontogeneia rostrata***	**Gammarid amphipod**	**0.65**
39	*Unknown gammarid (Family Hyalidae)*	Gammarid amphipod	0.12
40	*Unknown gammarid (Family Isaeidae)*	Gammarid amphipod	0.10
41	*Unknown gammarid (Family Ischyoceridae)*	Gammarid amphipod	0.24
42	***Caprella californica***	**Caprellid amphipod**	**0.36**
43	***Caprella laeviuscula***	**Caprellid amphipod**	**0.06**
44	***Caprella natalensis***	**Caprellid amphipod**	**0.82**
45	*Cumella vulgaris*	Cumacean	0.36
46	***Leptochelia sp.***	**Tanaid**	**0.49**
47	*Nebalia gerkinae*	Leptostracan	0.39
48	***Porcellidium sp.***	**Copepod**	**0.33**
49	***Harpacticoid 1***	**Copepod**	**0**
50	***Harpacticoid 2***	**Copepod**	**0**
51	*Harpacticoid 3*	Copepod	0.50
52	*Anoplodactylus viridintestinalis*	Pycnogonid	0.01
53	*Idotea montereyensis*	Isopod	0.73
54	***Idotea resecata***	**Isopod**	**0.36**
55	*Munna sp.*	Isopod	0.20
56	*Pugettia producta*	Brachyuran crab	0.36
57	*Pugettia richii*	Brachyuran crab	0.34
58	*Hippolyte californiensis*	Caridean shrimp	0.45

640   **Table S3.** Summary of traditional biodiversity metrics for each site

Site abbrev	Region	Species richness	Raw abundance	Effective diversity (Hill number)
DC	Barkley Sound	21	1168	11.8
RP	Barkley Sound	17	1590	8.8
SA	Barkley Sound	17	924	6.8
DK	Clayoquot Sound	21	1168	11.8
EB	Clayoquot Sound	21	4812	8.5
IN	Clayoquot Sound	16	1803	7.5
CB	Gulf Islands	18	2112	5.0
GB	Gulf Islands	11	2239	4.3
JB	Gulf Islands	22	3522	7.1
LH	Gulf Islands	15	2259	6.6
SS	Gulf Islands	10	740	6.6
HL	Haida Gwaii	9	16743	1.3
RA	Haida Gwaii	14	934	8.6

Equation S1. Equations for D^2_{adj} calculations

Explained deviance or D^2 (Guisan and Zimmerman 2000) was calculated by the equation:

$$D^2 = 1 - Residual\ deviance\ /\ Null\ deviance$$

Where the residual deviance refers to the deviance in predicted species occurrences, and the null

deviance refers to deviance in observed species occurrences.

Adjusted D^2 (Guisan and Zimmerman 2000) was calculated by the equation:

$$D^2_{adj} = 1-(((n-1)/(n-p-1)*(D^2-1))$$

Where n is the number of sites (13) in our study, and p is the number of parameters the model

estimated (4, Ovaskainen et al. 2017, Supporting Information). Negative D^2_{adj} values (those which

had greater residual deviance than null deviance) were set to 0.

669 **Equation S2.** Calculations for type III variation partitioning on the HMSC model (Borcard et al.
670 1992)
671

672 We estimated three versions of the HMSC model: 1) environmental variables only with the random

673 effects (*m2*), and 2) spatial distances only with random effects, and 3) the original global model

674 which included environmental and spatial variables, site- and region-level random effects (*m1*).

676 *m1* = the fraction of variation explained by all environmental variables in the HMSC model
677 estimated with only environmental variables;
678 *m2* = the fraction of variation by spatial distances in the version of the HMSC model estimated
679 with only spatial distances;
680 *m3* = the global model including both;
681

682 From each model, we extracted the total variation explained by environment and space, the

683 variation explained by environment, and the variation explained by space. To determine the shared

684 fraction of variation between environment and space, we multiplied each whole fraction by the

685 calculated community-level D^2_{adj} to reflect the model deviance in our variance partitioning

686 estimates, and followed the equations to calculate fractions *a, b, c,* and *d*.

687
688 Fraction *ab* (pure environment + shared fraction) = D^2_{adj} *m1*
689 Fraction *bc* (pure space + shared fraction) = D^2_{adj} *m2*
690 Fraction *abc* (pure environment + shared fraction + pure space) = D^2_{adj} *m3*
691
692 Fraction *a* (pure environment) = *abc - bc*
693 Fraction *c* (pure space) = *abc - ab*
694 Fraction *b* (shared fraction) = *ab + bc - abc*
695 Fraction *d* (residuals) = 1 - *abc*
696
697 Fractions *a* ("Environmental conditions"), *c* ("Spatial distance"), *b* (shared fraction), and *d*

698 ("Residuals") are shown in the Venn diagram in Fig. S3.

709 **Figure S1.** Schematic diagram field sampling regime. Black lines represent transect lines, and red

710 boxes represent 0.25 x 0.25 m quadrats.

Figure S2. Site-by-species presence-absence matrix of all 58 invertebrate species in the study

listed in alphabetical order. Black cells indicate species presences.

Environmental conditions   Spatial distance

0.22   0.04   -0.03

Residual variation 0.76

729 **Figure S3.** Venn diagram summarizing the fractions of variation explained by environmental

730 covariates only (nine water quality and five biotic variables), spatial distance only, and the shared

731 fraction explained by environment and space.

Figure S4. A correlation plot showing modelled site-level co-occurrence of species pairs across all species. Purple cells represent positively co-occurring species pairs, and turquoise cells represent negatively co-occurring species pairs. Species names along both axes are ordered according to the output of hierarchical clustering with Ward's criterion (Ward 1963) on pairwise co-occurrence values. This figure is a supplement to Fig. 4a in the main text.

Figure S5. Heat map depicting abundance patterns in all 58 species in the study. Species names are ordered from highest to lowest predicted mean proportional abundance. Cell colours correspond to the co-occurrence groups in Fig. 3a, with purple cells representing members of the *Nereis* assemblage, and turquoise cells indicating members of the *C. californica* assemblage, and grey cells indicating species that did not significantly co-occur negatively or positively with other species. Cell shade strength represents proportional abundance at a given site (darker means higher abundance). Most species outside the top twenty most abundant had extremely low predicted proportional abundances owing to their low raw abundances. This figure is a supplement to Fig. 3b in the main text.

Figure S6. Plot showing relationship between maximum site-level abundance and site occupancy

for all species. Black points represent broadcast spawners and white points represent brooders.

Spirorbid polychaetes were excluded from this figure due to their unusually high maximum

abundance (approximately 16 300 at one site). A two-tailed t-test showed that there was no

significant difference in the number of sites colonized between brooders and broadcast spawners

(t = 0.052, df = 42.5, p = 0.96).

Literature Cited

Borcard, D., P. Legendre, and P. Drapeau. 1992. Partialling out the spatial component of

ecological variation. Ecology 73:1045–1055.

Guisan, A., and N. E. Zimmermann. 2000. Predictive habitat distribution models in ecology.

Ecological Modelling 145:147–186.

Ovaskainen, O., G. Tikhonov, A. Norberg, F. Guillaume Blanchet, L. Duan, D. Dunson, T. Roslin,

and N. Abrego. 2017. How to make more out of community data? A conceptual framework

and its implementation as models and software. Ecology Letters 20:561–576.

Ward, J. H. 1963. Hierarchical Grouping to Optimize an Objective Function. Journal of the

American Statistical Association 58:236-244.