Immunological Diversity with Similarity

Rohit Arora, PhD[1], Harry M. Burke[2], and Ramy Arnaout, MD, DPhil[1,3]*

[1]Division of Clinical Pathology, Department of Pathology and [2]Division of Clinical Informatics,

Department of Medicine, Beth Israel Deaconess Medical Center, Boston, MA 02215. [3]Harvard

Medical School, 25 Shattuck St, Boston, MA 02115

*To whom correspondence should be addressed at rarnaout@gmail.com

Summary

A diverse immune repertoire is considered a hallmark of good health. However, other things

being equal, current methods for measuring repertoire diversity do not distinguish between a

repertoire that is composed of similar sequences, clonotypes, or clones and a repertoire that is

composed of different ones, even though the latter is intuitively more diverse. Here we describe

a framework for incorporating similarity into diversity measures, and illustrate using … define

diversity with binding similarity as functional diversity, and measure functional diversity on 391

large-scale antibody and T-cell receptor (TCR) repertoires. We find that while repertoires often

contain millions of unique sequences, functional diversity reveals a landscape defined by at

most a few thousand unrelated CDR3 binding targets. Naïve/IgM repertoires have more unique

sequences than memory/IgG, but memory/IgG repertoires are more functionally diverse.

Functional diversity is sensitive to vaccination, infection, and aging, and unlike raw diversity is

robust to sampling error. Finally, according to functional diversity, repertoires from different

people overlap significantly, suggesting a definable ceiling for the functional diversity of

humanity. Similarity redefines diversity in complex systems.

Introduction

Immune repertoires are famously diverse. Collectively, a person's ~10^{12} B and T cells express

many millions of unique recombined antibody and TCR genes as part of millions of clonal

lineages, more unique sequence than in the entire germline genome (Jiang et al., 2013). At the

sequence level, the repertoires of any two people overlap by only a fraction of a percent,

indicating still higher diversity in the population (Arnaout et al., 2011; Robins et al., 2010). Yet

repertoires are formed from V, D, and J gene segments that almost all people share and that

are expressed at similar frequencies across individuals targets (de Bourcy et al., 2017; DeWitt

et al., 2016), and repertoires are shaped by similar antigenic exposures and a consequent need

to recognize and bind similar. Squaring the diversity that is seen with the similarity that must

exist is a major goal in immunology.

This goal has relevance for disease stratification and clinical management across a range of

conditions. B- and T-cell diversity fall with age, as specific exposures expand a few select

lineages at the expense of others (Messaoudi et al., 2004). Chronic infection appears to have a

similar effect, impairing vaccination (Jiang et al., 2013). Low B-cell diversity is associated with

physiological frailty, a syndrome seen alongside conditions that are traditionally considered to

be unrelated to adaptive immunity (e.g., atherosclerotic cardiovascular disease), independent of

chronological age (Gibson et al., 2009). In cancer, a rise in sequence-level T-cell diversity is

thought to predict a successful response to immune-checkpoint inhibitors, drugs that make

tumors more visible to the immune system (Hopkins et al., 2018).

Traditionally, diversity has been measured as a simple count of the number of different

sequences, lineages, or clones in a sample, a measure known formally as species richness.

However, species richness ignores a key feature of repertoire diversity, species frequency: the

fact that some sequences are common and others rare. In an intuitive sense, a repertoire with a

single dominant (e.g., leukemic) clone is less diverse than a repertoire that has the same

number of clones but no dominant clone. To incorporate frequency into measurements of

diversity, there exist a family of measures that includes Shannon entropy, the Simpson index

(and related Gini coefficient), and the Berger-Parker index (Hill, 1973). These differ from each

other in how much weight they place on frequency: i.e., how much more a large clone adds to

the total diversity than a small one. Mathematically, weight can be represented as a parameter,

q, in the so-called Hill framework, a master equation for diversity in which species richness,

Shannon entropy, the Simpson index, and the Berger-Parker index, among others, have been

shown to correspond to different values of q (q=0, 1, 2, and ∞, respectively). It is understood

that no single diversity measure is best: the different measures provide complementary

information about a given complex system (Morris et al., 2014). Robust methods exist for

correcting sampling error for species richness and the frequency-weighted measures, and these

methods are becoming standard for measuring immunological diversity (Greiff et al., 2015;

Kaplinsky and Arnaout, 2016).

However, there is a second key feature of repertoire diversity that the frequency-weighted

measures fail to capture: species similarity. A repertoire made up of all-different sequences is

intuitively more diverse than a repertoire that has the same number of sequences, present at

the same frequencies as in the first repertoire, but all drawn from the same lineage or clone. In

the literature, this fact is sometimes addressed indirectly by grouping sequences together before

measuring diversity, for example by clustering reads, collapsing clones, or binning by V(D)J

segment usage (DeWitt et al., 2016; Jiang et al., 2013; Ju et al., 2018; Kaplinsky et al., 2014;

Vollmers et al., 2013). However, grouping usually imposes a binary threshold—in or out—on

what is by nature a continuous and overlapping relationship among sequences and their

encoded proteins. Grouping also usually zeros out or ignores any diversity that might exist

within groups. It is unclear what is lost by ignoring similarity, or what might be gained from a

more complete synthesis of diversity with similarity. This is true not only for the immunome, but

for other complex systems such as microbiomes, images, and tissues. Here we sought to

develop and explore a continuous framework for measuring diversity-with-similarity on B- and T-

cell repertoires.

Results

Framework. We measured diversity-with-similarity on high-throughput B- and T-cell repertoires

using a robust mathematical framework initially proposed for studying diversity in ecology and

environmental settings (Leinster and Cobbold, 2012). This framework provides "with-similarity"

counterparts for species richness and the frequency-weighted diversity measures: species

richness with similarity (^0D_s, which places a very small weight on frequency, and $^{\emptyset}D_s$, which like

^0D ignores frequency altogether), the exponential form of entropy with similarity (^1D_s, henceforth

simply "entropy with similarity," and likewise for other named indices), and so on. In ^qD_s

notation, q is the frequency-weighting parameter, D_s denotes diversity-with-similarity, and D

without the subscript means diversity without similarity, which we refer to as "raw diversity."

Mathematically, the key innovation of diversity-with-similarity relative to raw diversity is inclusion

of a similarity matrix whose entries quantify how similar each pair of species (sequences,

clonotypes, etc.) is. Constructing the similarity matrix necessitates a choice of similarity

measure. (Note the difference between similarity measures and diversity measures: a similarity

measure is used to build the similarity matrix, which then is used to calculate diversity

measures.) The choice of similarity measure depends on the biological feature(s) of interest. For

example to study somatic hypermutation, one might use the Hamming distance. Just as

different weightings provide complementary information about rare vs. frequent species—for

example, the number of new thymic emigrants (species richness; ^0D or $^{\varnothing}D_s$) vs. large leukemic

clones (Berger-Parker index; $^{\infty}D$)—different similarity measures are expected to reveal different

systems-level features of repertoires' sequence-level configuration. Also as with raw diversity

measures, expressing results for the new diversity-with-similarity measures as effective

numbers, also known as number equivalents, (Macarthur, 1965; Hill, 1973; Jost, 2007; Marion

et al., 2015), as opposed to as bits or nats (for ^1D_s) or as various fractions (for $^{>1}D_s$), makes it

possible to compare them to each other, regardless of weighting or similarity measure, on a

single intuitive scale (Box 1).

Similarity measure. We were interested in the single most fundamental mechanistic feature of

antibodies and TCRs: binding to specific targets (Fig. 1). Therefore for our similarity measure,

we used a proxy for binding affinity that follows from the empirically observed changes in

dissociation constant (K_d) associated with amino-acid substitution in antibody and TCR CDRs

(Jankauskaite et al., 2018). We found that on average, a single amino-acid substitution at an

antibody-antigen or TCR-peptide binding surface lowers affinity by 4-5 fold (geometric mean),

with a long tail corresponding to rare orders-of-magnitude effects (Fig. 2a). We focused on

CDR3, the third complementarity determining region, of IgH and TCRβ, since this is the single

most important contributor to binding specificity (Xu and Davis, 2000); however, our approach

can be applied to other regions. Because the relationship between sequence and specificity

remains non-predictive and therefore complex, for any given sequence pair the similarity

imputed from the observed distribution will be approximate; however, averaged over the many

millions of pairs in each repertoire, it was expected to be a reasonably accurate first-pass

repertoire-level view of immunological diversity with binding similarity.

Using this similarity measure, diversity-with-similarity is interpreted as the effective number of

sequences in a repertoire if the sequences were equally common and had no binding overlap

with each other (Box 1), or equivalently, the number of equally common non-overlapping binding

targets that a repertoire can recognize. We therefore refer to this version of diversity-with-

similarity as "functional diversity" (Fig. 1). Functional diversity can be interpreted in the context

of a "shape space" (Perelson and Oster, 1979) that contains all possible CDR3 binding targets,

with nearby targets having similar three-dimensional shapes and conformations (Fig. 1a). Each

CDR3 binds a (possibly overlapping) subset of targets; together, a repertoire's CDR3s cover

some part of shape space (Fig. 1b). Functional diversity measures the size of this region,

controlling for similarity and overlap in binding among different CDR3s (Fig. 1c).

Validity. We first established that our similarity measure behaved sensibly, with closely related

sequences scoring high and unrelated sequences scoring low (Fig. 2b). We next established

that it resulted in intuitive values for functional similarity by testing against expectations on

simple *in silico* repertoires. In a representative test, we constructed four repertoires with 34

unique sequences each and 752 sequences total (Fig. 2c-d). In each repertoire, a few

sequences were common (larger circles) while most were rare (smaller circles), representing

the long-tailed frequency distribution seen in real repertoires (Arnaout et al., 2011; Weinstein et

al., 2009). Importantly, the species-frequency distribution for all four repertoires was identical,

meaning that raw diversity was also identical across the repertoires, for all frequency

weightings. The only difference between the repertoires was in the pairwise similarity among

sequences.

For the first repertoire (Fig. 2d, top row), we chose closely related sequences from a single real-

world CDR3 clone. We expected that species richness with similarity—functional species

richness—would be close to 1. (We used $^{0}D_{s}$ here; $^{\varnothing}D_{s}$ performed similarly.) We observed a

value of 1.5; the extra 0.5 reflected sequence diversity within the clone. For the second

repertoire (Fig. 2d, second row), we swapped out half the unique CDR3s with CDR3s from a

different, unrelated real-world clone. As expected, we observed a rough doubling of functional

diversity, to 2.4. For the third repertoire (Fig. 2d, third row), we replaced all the sequences with

34 randomly chosen real-world CDR3s. We expected a functional diversity that was much

higher than in the first two repertoires but less than 34 because of the inherent sequence

similarities that make a CDR3 a CDR3, and, consistent with this expectation, observed a value

of 22. For the final repertoire (Fig 2d, bottom row), we replaced the CDR3s with random amino-

acid sequences (controlling for length), expecting a functional similarity of nearly 34, and this

again was observed ($^{0}D_{s}$=32). In contrast to these differences in functional diversity, raw

diversity was indistinguishably 34 for all four repertoires. In every example, functional diversity fit

an intuitive sense of what diversity should mean (Fig. 2c), while raw diversity failed to detect a

difference. These results support the validity of our functional-diversity framework for immune
repertoires.

Robustness. Sampling error—the "missing-species" problem (Bunge and Fitzpatrick,
1993(Bunge and Fitzpatrick, 1993))—is known to be a major potential confounder when
measuring raw diversity, necessitating large (e.g.) blood volumes and/or post-hoc statistical
correction for measurements on the sample to reflect repertoire diversity in the individual as a
whole (Kaplinsky and Arnaout, 2016). A practical feature of functional similarity is that the
values are smaller than those of raw diversity (reflecting clustering of similar sequences; Fig. 1-
2). The effective coverage is therefore greater, meaning that less information about the
functional diversity of an individual's overall repertoire is lost upon sampling than is the case for
raw diversity. This observation suggested that functional diversity is more robust to sampling
error, possibly even making it accurate enough to use without statistical correction, and thus
useful for the sample sizes typically available for sequencing (10,000-1 million cells).

To test this possibility, we systematically downsampled from a representative TCRβ repertoire
and two representative IgH (IgG) repertoires, one prepared from mRNA and one from genomic
DNA, each with ~10^6 unique sequences, and compared raw vs. functional diversity on the
subsamples to those of the full sample (Fig. 3). (We wished to consider possibility of lower
diversity from mRNA than DNA, since transcriptionally less active cells may be less likely to be
sampled.) For TCRβ, we found that functional species richness saturated at a sample size of
~30,000 sequences and functional entropy at ~10,000 sequences (Fig. 3b, first and third
columns). Functional diversity for higher frequency weightings (q) saturated with even fewer
sequences. For IgH from mRNA, functional species richness did not saturate but did plateau,
with a final increase of ≤2 percent per order of magnitude. Assuming that each unique sequence
corresponds to a cell and 10^{10} B cells in the body, this final measured rate of increase means
that the individual's total functional species richness is no more than 50 percent higher than the
value measured on the sample (Fig. 3, middle row). This is the maximum expected sampling
error. For IgH from DNA, functional species richness had begun to plateau at the full sample
size, resulting in the value for the individual being no more than three times as much as in the
sample (maximum three-fold error). Meanwhile, functional entropy saturated at 30,000 cells for
IgH from mRNA and 300,000 cells for IgH from DNA. This behavior was in marked contrast to
that of raw diversity, which did not saturate or plateau for species richness (Fig. 3, white
symbols), consistent with previous reports and illustrating the need for statistical correction
(Kaplinsky and Arnaout, 2016).

We then asked whether the robustness of functional diversity can be expected to generalize for

any IgH or TCR repertoire. We reasoned that a "meta-repertoire" comprising sequences drawn

uniformly (i.e., without regard to frequency) from many individuals will be more diverse, by any

measure, than any single repertoire (which will have fewer sequences, and in which the same

sequence may appear multiple times). Downsampling from a meta-repertoire therefore provides

an upper bound or worst-case scenario for the sampling requirements for any single repertoire.

To build meta-repertoires, we pooled and then uniqued CDR3s from 114 different IgH

repertoires from 79 individuals including Americans of African, European, and Hispanic descent

(Bild et al., 2002; DeWitt et al., 2016; Vollmers et al., 2013) to build an IgH meta-repertoire of

roughly 36 million unique sequences—as many as or more than ever observed or currently

estimated to be in a typical individual—and similarly for CDR3s from TCRβ repertoires from 69

healthy individuals (of mostly European but some Asian descent) (Emerson et al., 2017) to build

a TCRβ meta-repertoire of 10 million unique sequences, and downsampled from each of these

meta-repertoires as above (Fig. 3, large circles). We found that functional diversity plateaued for

all q, saturated for $q{\geq}1$ and reflected overall diversity to within a few percent from sample sizes

of 50,000 for TCRβ and IgH RNA and 100,000 for IgH DNA for q=0, and 30,000 for TCRβ and

IgH RNA and 300,000 for IgH DNA for $q{\geq}1$ (Fig. 3, large colored circles). Together, these results

confirmed that functional diversity measured on samples is an accurate measure of overall

functional diversity in the individual, at conventional sample sizes.

Raw and functional diversity. We measured raw and functional diversity on 141 healthy

human subjects (Fig. 4). For IgH, we found a (geometric) mean functional species richness

(D_s) of 677 (range, 487-916) from mRNA and 2,205 (range, 2,042-2,485) from DNA,

suggesting that on average, the human antibody repertoire is capable of recognizing the

equivalent of no more than a few thousand unique non-overlapping heavy-chain CDR3 binding

targets. (As above, lower diversity from mRNA was not unexpected, since inactive cells, which

produce less IgH mRNA than active cells, may be underrepresented.) For TCRβ the mean

functional diversity was 140 targets (range, 115-167). Functional diversity can be thought of as

clustering similar sequences together, although functional clusters can overlap and sequences

can belong to multiple clusters. An indication of the average size of these clusters can be

obtained by taking the ratio of raw to functional diversity measures. For species richness, we

found that IgH typically had hundreds of sequences per cluster, while TCRβ had thousands.

Thus by both functional diversity and average functional-cluster size, IgH CDR3 repertoires are

roughly 5-10 times as diverse as TCRβ (for small q). Repertoires with higher raw diversity might

be expected to be more functionally diverse, but we found no consistent trend across all

repertoire types. Thus, functional diversity generally complements raw diversity, adding

information not captured by raw diversity alone.

Naïve vs. memory B cells. We next sought to investigate what this information might add to our

understanding of adaptive immunity. We began with two widely studied B-cell subsets, naïve

(IgM) and memory cells (predominantly IgG). Previous studies have shown that naïve

repertoires have higher raw diversity than memory repertoires (DeWitt et al., 2016). This is at

least superficially consistent with the fact that only a subset of naïve cells are selected to enter

into the memory compartment. However, in a functional sense there is a case to be made that

memory/IgG repertoires should be more diverse, since somatic hypermutation differentiates

memory cells from naïve cells, and indeed from each other. Using well-characterized publicly

available repertoires from DNA from three healthy human subjects, we confirmed that by raw

species richness, naïve (CD27$^-$IgM$^+$) B-cell repertoires are ~10 times as diverse as memory

(CD27$^+$IgM$^-$) repertoires (DeWitt et al., 2016) (Fig. 5a-b). Yet by functional species richness, we

found that memory repertoires were at least as diverse as naïve (Figs. 5a-b). Comparing raw

and functional diversity for 34 IgM and 32 IgG repertoires from mRNA (repertoires with less than

100,000 total sequences were discarded) from 28 additional healthy individuals from a separate

dataset showed a similar pattern as for the three DNA repertoires: in all but a few outliers, IgM

had higher raw diversity but IgG had higher functional diversity (Fig. 5c-d). For raw diversity, the

IgM:IgG ratio rose from ~3:1 at q=0 to peak at 10:1 around q=1, due to a large fraction of rare

IgG sequences (Fig. 5d). This effect was more pronounced for naïve:memory (Fig. 5b). For

functional diversity, the absence of a peak in the IgM:IgG ratio suggests that these many rare

sequences must nonetheless be similar to others in the repertoire, possibly because they are

members of clones (Fig. 5b,d).

Cytomegalovirus (CMV) exposure. CMV is a herpesvirus to which half of the adult population

has been exposed and results in life-threatening opportunistic infections in newborns, transplant

recipients, and immunocompromised individuals (Emery, 2001). In most healthy individuals, it

causes a chronic infection marked by clonal expansion of both B cells and T cells and a

consequent fall in raw diversity, an effect also seen during aging (see below) (de Bourcy et al.,

2017; Qi et al., 2014). We measured raw and functional TCRβ CDR3 diversity for 120

individuals: 69 CMV-seronegative and 51 CMV-seropositive subjects aged 19-35 (Emerson et

al., 2017), the narrow age range helping control for any age-related effects. There was a clear

trend toward lower diversity in the CMV-seropositive group relative to the CMV-seronegative

group by both raw and functional diversity, for all weighting parameters (Fig. 6a). Combining raw

with functional diversity facilitated identification of two subgroups among the subjects with

known CMV status: subjects with a high raw Berger-Parker Index ($^{\infty}D$) were almost always

CMV-seronegative (Fig. 6b), whereas subjects with low functional Berger-Parker Index ($^{\infty}D_s$)

were almost always CMV-seropositive (Fig. 6c). The reverse—low $^{\infty}D$ or high $^{\infty}D_s$—did not

distinguish between the groups. Using both measures gave a better indication of CMV status

than did either one alone (Fig. 6d). The conclusion is that CMV is unlikely in the absence of

large clones/expanded lineages, as has been reported, but is likely only if the large

clones/expanded lineages that are present exhibit high similarity to other clones/lineages in the

repertoire, or else are indeed very large (Fig. 6e). Again, the addition of functional diversity

offers insight that raw diversity alone does not.

Flu vaccination. Vaccination with a seasonal trivalent influenza vaccine (TIV) triggers clonal

expansion in B cells. Previous work on five vaccinees showed likely flu-specific memory IgG

lineages emerging by day 7 post-vaccination (Vollmers et al., 2013). We found that combining

raw and functional diversity reveals a signature of clonal expansion and selection without the

need for lineage tracking (Fig. 7). We measured raw and functional diversity for IgM and IgG at

day 0 (pre-administration) and day 7 from all 14 vaccinees in Vollmers' dataset. We found that

for most subjects, for IgG, raw species richness rose from day 0 to day 7 while functional

species richness fell (Fig. 7a-b). This means that even as the number of sequences increased,

many of the new sequences were similar to each other (or to existing sequences), and they

tended to replace different-looking sequences. Meanwhile, there was no obvious pattern in IgM

(Fig. 7c). Together, these results are what we would expect from clonal expansion and selection

in a memory response, and thus represent a repertoire-level signature of these phenomena.

Interestingly, in most cases, raw and functional entropy both fell (Fig. 7a-b, right panels). This

suggests that most of the new sequences at day 7 were rare, while at the same time a subset of

sequences and functional clusters grew. Thus overall, the addition of functional diversity reveals

a key feature of clonal dynamics, which is not evident from raw diversity alone.

Aging. To explore the effect of age, we measured raw and functional diversity for TCRβ CDR3

repertoires from 41 healthy individuals aged 6-90 years old (Britanova et al., 2014) (Fig. 8). We

found that raw diversity falls with age regardless of weighting parameter; a fall in raw species

richness had been reported previously (Britanova et al., 2014). Functional diversity also fell,

regardless of weighting parameter. However, for species richness, four septuagenarians bucked

the trend (Fig. 8, arrows): even as their raw species richness was unremarkable relative to that

of other individuals of similar age, their functional species richness was similar to that of

children. Only one of these four had a high likelihood of being CMV-negative; the probability that

all four were CMV-negative was low. We therefore consider CMV unlikely as an explanation for

their high functional species richness. Unlike their peers, these four appear to have retained

functional diversity among their rarest sequences. (The alternative hypothesis is that these four

saw a rise in functional species richness from a lower level earlier in life, but we think this

unlikely given the overall downward trend across individuals.) We considered but excluded

PCR/sequencing artifacts as the cause, as we expected such artifacts would have led to larger

raw species richness, which was not observed. Thus, functional diversity identified for further

study individuals who were unremarkable by raw diversity alone.

Discussion

Diversity both affects function, and reflects it. In the adaptive immune system, the defining

tradeoff is breadth vs. depth: a repertoire must be sufficiently diverse to contain sequences that

can recognize a given target and lead to useful clones, but not so diverse that cells that express

such sequences are too rare to find the target on biologically relevant timescales (Schober et

al., 2018; Zarnitsyna et al., 2013). To monitor immunological diversity, either diagnostically or

therapeutically, we must be able to measure it, and to measure it, we must define it. It is

increasingly recognized that a reasonable definition of immunological diversity must account for

differences in species frequency. Here we argue such a definition must also account for

species' pairwise similarity, and show that binding similarity, which leads to what we call

functional diversity, provides useful insight into repertoire function.

Pairwise similarity can be seen as governed by a tunable parameter that helps define the

similarity matrix, analogous to how q is a tunable parameter that governs the effect of

differences in frequency (Chao et al.). In our study, the similarity matrix is defined by the

average single amino-acid change in K_d, an average based on over 1,300 independent

measurements, and the assumption of multiplicative independence. This source data is not

systematic, but to our knowledge is the best available. While our study is to our knowledge the

first of its kind, it follows a long tradition of attempts to estimate the number of binding targets

that can be recognized by the adaptive immune system immunization (Bachmann et al., 1994;

Obar et al., 2008). In these past studies, typically a sample of B or T cells was diluted until

binding to/protection from a given target was abolished, using whatever thresholds the

investigators deemed appropriate. If the limiting frequency for binding/protection was found to

be, for example, 1:3,000, the conclusion was that the repertoire could recognize 3,000 different

targets. This conclusion was based on the assumption that on average, all targets behave the

same as the one under study. Such studies gave functional diversities of 100 to 100,000 targets

for various T-cell populations and for B cells after antigen exposure, a wide range (Bachmann et

al., 1994; Obar et al., 2008). The width of this range may reflect real differences in the

frequencies of cells that are specific for different targets, or variability in stringency or

experimental setup. Interestingly, this "how-many-can-fit" logic seems not to have been used

when testing so-called natural antibodies, which bind many targets at low affinity (Frank, 2002;

Holodick et al., 2017; Notkins, 2004). For example, when one in five natural antibodies were

found to bind insulin (Chen et al., 1998), this was not taken to mean that the repertoire could

recognize only five targets, because of presumed overlapping specificity of these antibodies for

other targets. Meanwhile, theoretical studies have suggested a need for ≤10,000 binding

targets, and fewer for T cells than B cells, because of major histocompatibility complex (MHC)

restriction (Langman and Cohn, 1987; Zarnitsyna et al., 2013)

For raw diversity, we found that a typical repertoire contains on the order of 10 million unique

CDR3s, well above the upper end of previous estimate for the number of binding targets. These

findings are in line with other recent estimates that were likewise based on a combination of

deep sequencing and statistical correction (Britanova et al., 2014; DeWitt et al., 2016; Kaplinsky

and Arnaout, 2016). On average, these raw species richnesses mean that each of 100-100,000

putative binding targets can be bound by 100-100,000 unique CDR3s ($10^7/10^5$ to $10^7/10^2$). From

a medical perspective, such redundancy is good for treatment, because it supports the

prevailing view that there are many ways to design an antibody- or TCR-based drug that will

recognize a given target, but potentially a complicating factor for attempts to diagnose specific

diseases based on repertoire sequence, because it suggests that signatures of exposure to a

given target may be quite variable.

One of our key findings is that functional diversity is much lower than raw diversity: repertoires

contain only a few hundred functional clusters for TCRβ CDR3s and at most a few thousand for

IgH. The fact that functional diversity is based on K_ds suggests that functional diversity should

correlate with the number of structurally unique, non-overlapping target clusters that CDR3s can

recognize (Fig. 1). Yet our measurements of functional species richness lie at the low end of the

range of prior estimates. We propose two explanations. First, our measurements are limited to

CDR3s; variability in the rest of the antibody or TCR protein must add to the total number of

potential binding targets. This possibility is testable by extending our method to more or indeed

all of the antibody or TCR sequence. Second, functional diversity may be providing a less

detailed description of shape space than limiting-dilution studies: i.e., functional diversity may be

a coarse graining of the target-binding landscape (Fairlie-Clarke et al., 2009; Smith et al., 1997)

(Fig. 9). A pair of sequences may be similar enough to lie near each other in shape space, but

only one may bind a given target above a threshold level of specificity in a binding study. In

short, binding studies may be counting peaks while functional diversity is counting mountains. If

true, our results suggest that the landscape of TCRβ CDR3 binding is more clustered than that

of IgH, such that there are on average several times as many functional IgH clusters as TCRβ.

This prediction is testable, at least in principle, through large-scale systematic binding assays to

measure K_d, or by measuring binding as a binary outcome at multiple stringency thresholds.

Both explanations may contribute. To our knowledge ours is the first attempt a quantitative

summary of this landscape using data from large-scale binding studies and high-throughput

repertoire sequencing.

Why is CDR3 functional diversity higher for IgH than for TCRβ? We hypothesize that it is for the

same three reasons that there is more sequence diversity for IgH than TCRβ. First, humans

have 23 D_H gene segments vs. only 2 D_β segments, and D is the largest germline contributor to

CDR3. V and J segments tend to directly contribute little more than the canonical starts and

ends of CDR3s, and besides there are similar numbers of V-J combinations in IgH as TCRβ

(49×6=294 and 48×13=624, respectively). Second is somatic hypermutation, which diversifies

IgH but not TCRβ. And third, IgH CDR3s are longer than TCRβ CDR3s, allowing for a larger

number of possible sequences. Further analysis will be needed to test these hypotheses.

We have shown how functional diversity complements raw diversity to offer insight into the

difference between naïve and memory repertoires, to aid in identification of disease states, and

to illustrate clonal selection and other repertoire dynamics. We hope these examples will

encourage others to use and/or expand our framework to investigate repertoire dynamics in

other conditions, in other subsets, in the other chains (TCRα and IgL), and in other model

systems such as zebrafish (Weinstein et al., 2009) and mouse (Arnaout et al., 2011; Kaplinsky

et al., 2014). We draw attention to the fascinating difference between the number of unique

sequences, which ran into the millions in most of the repertoires we investigated, and the much

smaller numbers of what we call functional clusters (the effective numbers of functional

diversity). The result is a "functional degeneracy" among sequences that are organized into

functional clusters. Characterizing these clusters is an interesting topic for future work.

Functional similarity also offers a new perspective on similarities and differences between

people. To show that functional diversity is robust to sampling, we generated "meta-repertoires"

by pooling sequences from scores and in some cases over a hundred people, including people

from different ethnic backgrounds. Surprisingly, and somewhat unexpectedly given the low

sequence overlap between pairs of individuals, the functional diversity of these meta-repertoires

never exceeded the functional diversity of any given repertoire by more than a few fold;

moreover, the functional diversity trended toward saturating in samples of just a million

sequences (Fig. 3). Together, these findings predict that any two individuals share a majority of

their functional clusters, in stark contrast to the vanishingly small fraction of sequences they

share. Further, these findings suggest that the functional diversity of the entire population is only

a few hundred clusters for TCRβ CDR3s and a few thousand for IgH, and imply that these

clusters can be sampled exhaustively by sequencing fewer than 20 individuals. It will be

fascinating to test this finding with additional ethnically and geographically diverse populations,

to further examine our prediction that, contrary to conventional wisdom, the functional limits of

the adaptive immune system are in a practical sense both finite and within reach.

The focus of this study was binding similarity, but we expect that the utility of the diversity-with-

similarity framework will extend to other facets of immunology (e.g., somatic hypermutation) and

to other fields, most readily metagenomics, sociology, oncology, and cellular cartography

(Almendro et al., 2014; Heindl et al., 2016; Koopmans and Schaeffer, 2013; Li et al., 2012;

Taraska, 2015). We hope this study will serve as a template for incorporating similarity into the

study of other complex systems.

Acknowledgements

This research was supported by grants from the National Institutes of Health (NIAID

K08AI11495801) and the American Heart Association (15GPSPG23830004), contracts

HHSN268201500003I, N01-HC-95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-

HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168

and N01-HC-95169 from the National Heart, Lung, and Blood Institute, and by grants UL1-TR-

000040, UL1-TR-001079, and UL1-TR-001420 from NCATS, as well as support from the

Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by

National Science Foundation grant number ACI-1548562. The authors thank the investigators,

staff, and participants of the MESA study for their valuable contributions (a full list of

participating MESA investigators and institutions can be found at http://www.mesa-nhlbi.org), as

well as the Research Computing Group at the High Performance Computing Cluster at Harvard
Medical School.

Online methods

High-throughput repertoires. We obtained 391 quantitative high-throughput IgH and TCRβ
repertoires from 202 human subjects. These included IgH from naïve and memory B cells from
DNA (*n*=3 individuals) (DeWitt et al., 2016); TCRβ chains from DNA from healthy subjects
known to be serologically negative for cytomegalovirus (CMV) (*n*=69 individuals) (Emerson et
al., 2017) and from healthy subjects whose CMV serostatus was unknown (*n*=41 individuals)
(Britanova et al., 2014); pooled barcoded IgG and IgM heavy chains from mRNA from healthy
subjects before and seven days after administration of one of two influenza vaccines (*n*=28
individuals) (Vollmers et al., 2013); quantitative pooled TCRβ chains from DNA for subjects who
were otherwise healthy but serologically CMV positive (*n*=51 individuals) (Emerson et al., 2017);
and IgH chains from DNA for subjects enrolled in the Multi-Ethnic study of Atherosclerosis
(MESA; *n*=41 individuals) (Bild et al., 2002). CDR3 annotation was performed using our in-
house pipeline as previously reported (Kaplinsky et al., 2014) and standard tools (e.g. IMGT).
Details for obtaining these datasets are available from the primary publications referenced
above.

Similarity measures. A functional measure of similarity between polypeptides is how well they
bind the same target. We were interested in similarity as a function of the number of amino acid
substitutions (i.e., as a function of edit distance). The effect of substitutions on binding is
complex and depends on the position and identity of the specific amino acids involved; many
substitutions may have little or no effect, while a few may abolish binding entirely (Lunzer et al.,
2010). When comprehensive data are available, detailed statistical models can offer reasonable
predictions of the effect of specific amino-acid substitutions (Hopf et al., 2017; Salinas and
Ranganathan, 2018; Lee et al., 2008). However, this type of data does not yet exist across
entire antibody and TCR repertoires, and so simpler models are required. These models are not
expected to precisely predict the effects of specific substitutions, but should accurately reflect
the effects of substitutions when averaged over many pairs of proteins, such as the millions of
pairs in megacell-scale repertoires.

To develop a model for our similarity measure, *s*, we downloaded SKEMPI 2.0, which is to our
knowledge the largest and best-curated database of experimentally measured effects of amino-

acid substitution on protein-protein binding (Jankauskaite et al., 2018). Each entry includes a

Protein Data Bank (PDB) identifier (Berman et al., 2000), the type of structural region (Levy,

2010) that contains the substitution(s), one or more PDB coordinates, and (in nearly all cases)

the dissociation constant (K_d) of each member of the pair (referred to in the database and Fig.

2a as "wild type" and "mutant"). We extracted entries for all single amino-acid substitutions for

which K_d for both wild type and mutant were recorded, and considered only entries that involved

binding between antibody and antigen (*n*=797) or TCR and peptide/MHC (*n*=531; total

n=1,328). Although amino-acid substitutions anywhere in a protein may affect binding,

substitutions at the core of the binding interface are more likely to affect binding than

substitutions elsewhere (Levy, 2010). Therefore we split the data into core (*n*=584) and non-

core (*n*=744) groups and analyzed the effect of substitution binding, measured as

$|\log_{10}(K_{d\text{mut}}/K_{d\text{wt}})|$, separately for each group.

As expected, the probability distributions for the two groups differed substantially from each

other (Mann-Whitney U *p*-value $2.0\text{x}10^{-33}$). Substitution of a core residue had a 13-fold

(geometric) mean effect on binding, consistent with prior reports (Whittaker et al., 2001), while

substitution of a non-core residue had a 4-fold effect. Both probability distributions were long-

tailed, and were reasonably well described by exponential probability-density functions (i.e., of

the form ke^{-kx}). We confirmed that the distributions for antibody-antigen core residues (*n*=352)

and TCR-peptide/MHC core residues (*n*=232) were similar to each other, that the distributions

for antibody-antigen non-core residues (*n*=445) and TCR-peptide/MHC non-core residues

(*n*=299) were also similar to each other, that within each of the antibody-antigen and TCR-

peptide/MHC subgroups the distributions for core and non-core residues were different, and that

these results held separately for human and non-human (nearly all of which were mouse)

sequences (using the Structural Antibody Database (Dunbar et al., 2014) and the Structural

TCR Database (Leem et al., 2018) to assign species), all using Mann-Whitney U and visualized

as histograms. Using PyMol v2.2.0 (The PyMOL Molecular Graphics System, Version 2.0

Schrödinger, LLC), we next manually reviewed nine structures containing substitutions in

human IgH or TCRβ CDR3s (1BD2, 1OGA, 3BN9, 3QDJ, 3SE8, 3SE9, 4I77, 5C6T, 5E9D) and

found that to a good approximation, a constant fraction 0.15±0.05 of CDR3 amino acids consist

of core residues, with no obvious difference between chain types. To estimate the effect of a

single amino-acid substitution in a CDR3 in our datasets, we therefore combined core and non-

core distributions with a weighting of 0.15:0.85.

The resulting distribution was again long-tailed, with most substitutions having small effects and

a few having effects of many orders of magnitude (Fig. 2a). There were small spikes in the tail

for substitutions with $\gtrsim$60-fold effects, i.e. $|\log_{10}(K_{dmut}/K_{dwt})| \gtrsim 1.8$. A review of sources cited by

SKEMPI suggested that these spikes likely reflect ascertainment bias: selective experimentation

on amino acids with unusually strong effects (e.g. Pons et al., 1999; Taylor et al., 1998). To

counteract such bias, we therefore built a high-confidence dataset using 1.8 as the cutoff.

Ascertainment bias in two- and three-amino-acid substitutions is expected to follow the square

and cube of the bias in single-amino-acid substitutions, respectively, precluding rigorous

conclusions from being drawn from independence testing. However, comparison with those

groups was broadly consistent with either multiplicative ($s=c^m$, where s=similarity; c=the cost of

binding, i.e., 1/(fold effect); m=edit distance) or additive ($s=c/m$ for $m \geq 1$) independence.

Because additive effects result in higher pairwise similarities and therefore smaller repertoire

diversities than multiplicative effects, the multiplicative-independence model is more

conservative for studying the effects of similarity on diversity. We therefore chose the

multiplicative model for further analysis.

To determine the similarity between two CDR3s with edit distance m, we sampled

independently from the high-confidence dataset m times, and multiplied the costs together. We

confirmed that on average, the results of this stochastic sampling were the same as

deterministic calculation of $s=c^m$ with $c \approx 0.55$. We performed sensitivity analysis based on lower-

confidence cutoffs (down to c=0.48) and alternative assumptions (up to c=0.60). This resulted in

somewhat higher or lower diversity values, but qualitative patterns were robust to these

perturbations.

Diversity measures. We calculated ^qD as previously described (Hill, 1973; Kaplinsky and

Arnaout, 2016) and ^qD_s according to Leinster and Cobbold (Leinster and Cobbold, 2012). We

corrected ^qD for sampling error using Recon (default settings) as previously described

(Kaplinsky and Arnaout, 2016). We note Hill's framework (Hill, 1973) has inspired several

methods for incorporating similarity into diversity measurements, each of which retains useful

features of Hill's framework (Chao et al., 2018; Chiu and Chao, 2014; Leinster and Cobbold,

2012; Scheiner, 2012). Two of the new frameworks were introduced with explicit discussion of

how to decompose population-level diversity into within- and between-group components

(Leinster and Cobbold, 2012; Chiu and Chao, 2014). Each of these has advantages and

disadvantages over the other (discussed in Botta-Dukát, 2018) . We chose Leinster and

Cobbold's framework here because we found it easier to apply and interpret. For readability, we

made a minor change to the notation, from $^qD^Z$ to ^qD_s.

Use of this framework raised two issues that we addressed. First, its q=0 measure, ^0D_s,

depends on frequency albeit to a very small extent, unlike the Hill framework's q=0 measure, ^0D,

which is species richness (and is independent of frequency). Therefore, as a more direct

comparison to species richness, we calculated ^0D_s both with frequency information and without

it (i.e., setting the frequencies of each of the n species to 1/n). We refer to the latter as $^{\varnothing}D_s$ ("D-

null"). Second, it has been shown that this framework can result in unreasonably low diversity

values when most of the off-diagonal entries of the similarity matrix are far from zero, resulting

in an insensitivity to q (Botta-Dukát, 2018; Chiu and Chao, 2014). We expected most of our off

diagonals to be close to zero, since our similarity measures directly or indirectly involve

exponential decay, which generates small values, but confirmed that most of the off diagonals in

our similarity matrices were indeed close to zero by plotting histograms. Consequently, our

measures were sensitive to q, as desired and expected.

Robustness analyses. For robustness analyses, IgH and TCRβ were analyzed separately. The

upper-bound/worst-case scenario for IgH was evaluated by constructing a "meta-repertoire" by

combining IgG sequences of subjects before vaccination (n=28 individuals; Vollmers et al.

2013), sequences from memory cells from healthy subjects from public database (n=3; DeWitt

et al. 2016), and sequences from subjects enrolled in MESA study (n=41; Bild et al. 2002), and

sampling from this meta-repertoire without regard to the frequency of sequences. We chose

IgG/memory sequences where possible because those sets exhibited higher functional diversity

than naïve sets, and we were interested in maximizing diversity. We ignored the frequency of

sequences for the same reason: uniform frequency maximizes diversity, other things equal. For

TCRβ, we constructed a meta-repertoire by combining sequences from CMV seronegative

individuals (n=69; Emerson et al. 2017) and again sampling at uniform frequency. We chose

CMV seronegative individuals for the same reason as we chose memory/IgG sequences above:

seronegative individuals exhibited higher diversity. For both IgH and TCRβ, including all

sequences lowered diversities slightly. The representative samples were from subject D3 for

IgH (from DNA), subject SRR960344 for IgH (from mRNA), and subject Keck0070 for TCRβ

(CMV seronegative). CDR3 sequences were sampled proportional to their frequency in the

repertoire.

Figure titles and legends

Figure 1: Functional diversity. (a) Each dot represents a binding target (e.g. an epitope) with a different shape. Nearby targets have similar shapes (inset). Targets form clusters of similarity. **(b)** Each colored region represents the targets that can be bound by one of six unique CDR3 sequences in a representative repertoire; this repertoire has a raw species richness of 6. Together, the colored regions cover the part of shape space that can be bound by the repertoire. (The unbound region might include, e.g., self-antigens.) Note the substantial overlap in binding targets for the orange, yellow, green, and blue CDR3s. This overlap reflects binding similarity among these CDR3s. **(c)** Because of this similarity, the repertoire covers only the region denoted by the four identical non-overlapping squares. The functional species richness of this repertoire is therefore 4: this repertoire has the same species richness as a repertoire comprising four CDR3s that have zero overlap in binding specificity.

Figure 2: Validity. (a) Single amino-acid mutations in antibody and TCR molecules have a range of effects on affinity, as measured by change in dissociation constant, K_d (gray). This was well fit by a simple exponential (black line), providing parameterization for the similarity metric. **(b)** CDR3s with high sequence identity have high similarity, while different CDR3s have low similarity. Shown are two clones, represented by red and white subnetworks, each composed of 17 unique CDR3 sequences drawn from clonotypes of two real IgH repertoires. Node size corresponds to the frequency of each sequence; edges connect pairs of sequences that differ at a single amino-acid position. **(c)-(d)** Functional similarity agrees with an intuitive sense of repertoire diversity. Each of the four repertoires in (c) has the same number of unique sequences, present at the same frequencies; as a result, they all have identical raw diversity (for every value of q) despite their obvious quantitative and qualitative differences. In contrast, functional diversity increases with the number of, and increasing difference between, repertoires' constituent sequences. Node size denotes sequence frequency. Shades denote different clonotypes. Comparing the third and fourth rows, note that even when two repertoires have the same number and frequencies of unique sequences, the repertoire whose sequences are more different from each other (random peptides) has the higher functional diversity.

Figure 3: Robustness. Results for raw and functional species richness ($q=0$; ^0D and ^0D_s) and raw and functional entropy ($q=1$; ^1D and ^1D_s). Raw, white shapes; functional, colored shapes. Large symbols give an upper bound/worst-case scenario based on sampling meta-repertoires; small symbols give results for a representative sample from DNA (circles) and mRNA

(triangles). First row: sample diversity is plotted as the effective number of sequences. For $q=0$,

functional diversity plateaus for TCRβ and IgH RNA and trends toward a plateau for IgH DNA at

the tested sample sizes; all three plateau for $q=1$; raw diversity does not plateau for either $q=0$

or $q=1$. Second row: discovery rate is the probability that the next sampled sequence will add to

the diversity. For example, for the IgH DNA representative sample, for $q=0$ raw diversity, at a

sample size of 1 million sequences there is still a probability of ~0.5 (a 50 percent chance) that

the next sequence to be sampled will be one that has not yet been seen and will therefore add

to the diversity. Third row: maximum error is the maximum fraction by which the diversity in the

sample can underestimate the diversity in the individual from whom the sample was taken.

Horizontal dashed lines indicate the threshold for two-fold error. For example, for the worst-case

scenario for TCRβ, $q=0$ functional diversity measured on a sample of 10,000 sequences will be

no more than a two-fold underestimate of diversity in the individual as a whole; in other words,

the sample value will be at least 50 percent of the overall value.

Figure 4: Diversity in individuals. Raw (black lines; left vertical axis) and functional (colored

bars; right vertical axis) species richness (q=∅) for 179 CDR3 repertoires from healthy

individuals representing **(a)** IgH from mRNA (all isotypes: IgA, IgG, IgM, IgD, and IgE), **(b)** IgM

and **(c)** IgG from mRNA from the subjects in (a), **(d)** IgH from DNA (all isotypes), **(e)** naïve IgH

from DNA, **(f)** memory IgH from DNA, and **(g)** TCRβ from DNA. See Methods for references.

Matched pairs of symbols below the horizontal axis denote replicates. Note the difference in the

scale for functional diversity between IgH and TCRβ. Note also a general lack of correlation

between raw and functional species richness, except in (c).

Figure 5: Naïve vs. memory. (a) Diversity profiles for naïve (red) and memory (black)

compartments from three deeply sequenced individuals. A diversity profile is a way to show

diversity across a range frequency-weighting parameter values at once. By raw diversity (left),

the naïve compartment is more diverse across the range of weightings. By functional diversity

(right), this distinction disappears. In **(b)**, this disappearance is highlighted by plotting the ratio of

naïve:memory diversity for raw diversity (red) and functional diversity (black). According to

functional diversity, the naïve compartment is no more diverse, and indeed sometimes

somewhat less diverse, than the memory compartment. This reversal is even more prominent in

comparisons of repertoires from an additional 28 healthy subjects **(c,d)**.

Figure 6: Infection. (a) Diversity profiles showing effective number of species as a function of

weighting parameter q for diversity without similarity (left) and diversity with similarity (right)

showing a trend toward lower diversity in CMV-seropositive individuals (red) relative to CMV-
seronegative individuals (black), especially for large q. **(b)** Raw Berger-Parker Index ($q=\infty$),
which measures the largest clones, showing that high diversity—an absence of large clones—is
rare in CMV-seropositive individuals. **(c)** Functional Berger-Parker Index, showing that low
diversity—the presence of large clones with similarity to other clones in the repertoire—is rare in
CMV-seronegative individuals. **(d)** Combining raw and functional Berger-Parker Indices (first
principal component of PCA, which explains 72 percent of variance) illustrates both of the trends
in (c): for the third of subjects beyond the cutoffs indicated by the horizontal dashed lines CMV
serological status is assigned with an accuracy of 95 percent. **(e)** Schematic representation of
the three classes revealed by combining diversity with and without similarity. Each circle is a
clone; each collection of clones is a representative repertoire. Top: subjects without large clones
are almost always CMV seronegative. Bottom: subjects with large clones that are similar to
other clones in the sample (shown in red) are almost always CMV seropositive. Middle:
repertoires with large clones that are not similar to other clones in the repertoire may be either
CMV seropositive or CMV seronegative. Receiver-operator characteristic (ROC) analysis gave
an area under the curve (AUC) of 0.79.

Figure 7: Vaccination. Raw and functional diversity together reveal clonal expansion and
selection without needing lineage analysis. **(a)** In the IgG compartment, raw species richness
rises while functional species richness falls in most vaccinees (left). Meanwhile raw and
functional entropy both fall (right). The difference vs. species richness suggests most new
sequences at day 7 are rare. (b) Meanwhile, the IgM compartment changes less by these
measures.

Figure 8: Aging. Raw and functional species richness ($q=\varnothing$) for TCRβ CDR3 repertoires from
41 healthy individuals. Arrows denote four septuagenarians who bucked the trend of lower
functional species richness with age. Note that for each individual, the raw species richness is
~10-fold higher than previously reported (Britanova 2014), likely because the method we used
to correct for missing species (Recon) is more sensitive than the method used in the previous
report (Fisher).

Figure 9: Binding landscape. (a) Schematic of the target-binding landscape. The gray
distribution represents CDR3 sequences that bind a given target. Sequences are ordered by
their similarity to each other. (In reality, similarity is a multidimensional property that makes it
impossible to order sequences in a single dimension as shown here; this is done for illustrative

purposes only.) The height at each sequence denotes the affinity with which it binds the target

(vertical axis; measured e.g. by K_d). Many more sequences bind the target at low affinity than at

high affinity, resulting in a "mountain-and-peak" appearance. This schematic is useful for

interpreting functional diversity as described in this study, and the raw diversity estimates based

on previous binding studies, as described in the Discussion. (Note that in this schematic, the

raw diversity as measured simply corresponds to the total number of sequences along, i.e. the

width of, the horizontal axis.) At high affinity, very few sequences bind a given target. At medium

affinity, more sequences bind, and can be binned into two small clusters, represented by the

small circles. At low affinity, many sequences bind, and can be binned into a single large

cluster, represented by the large circle. In **(b)**-**(d)**, many targets are shown. Each color

corresponds to a different target; nearby targets are structurally similar. As in (a), each colored

area denotes the sequences that bind a given target, as a function of binding affinity (vertical

axis). Experiments usually detect the highest-affinity sequences: the peaks of the landscape

(above the horizontal dotted line). The narrower the peak when it crosses the experimental

threshold, the rarer specific sequences are, and the larger the number of targets that the

repertoire will be estimated to bind. (For example, if 100,000 sequences are shown across the

horizontal axis in each plot, and only one crosses the experimental threshold for a given target,

the frequency of sequences specific for that target is 1:100,000, and the conclusion will be that

there must be 100,000 such targets that the repertoire can bind. If 100 cross the experimental

threshold, the conclusion will be that the repertoire can bind only 1,000 targets.) Functional

diversity measures the overall contours of the landscape. Conceptually, this can be thought of

as measuring the number and size of the "mountains" at a lower affinity threshold (horizontal

solid lines). The differences in functional diversity between (b) memory IgH, (c) TCRβ, and (d)

naïve IgH correspond to different landscapes. The raw species richness of memory IgH and

TCRβ are comparable, represented here by the same width of all the plots. In addition, a similar

number of sequences per target cross the experimental threshold, so estimates of the total

number of targets that the repertoires can bind will also be comparable. However, less low-

affinity overlap between the targets of the IgH sequences in (b) gives it higher functional species

richness than the TCRβ repertoire in (c): here, six functional clusters (white circles) vs. three.

(The sizes of the clusters are related to frequency-weighted functional diversity measures, i.e.

larger q.) The sequences in the naïve IgH repertoire in (d) have only low affinity for the six

colored targets, and many recognize more than one target (overlapping colored areas). Note the

lower experimental threshold (horizontal dotted line), consistent with the ~10% or more of

antibodies that recognize a target and the high degree of cross-reactivity in studies of natural

antibodies (Frank, 2002; Holodick et al., 2017). The functional diversity threshold controlled by

the average effect on K_d of a single-amino-acid change in CDR3. If the effect were larger—or if

it were amplified by e.g. raising it to a power when building the similarity matrix—the threshold

would be higher, and vice versa.

Box 1: Interpreting effective numbers

Consider two repertoires of 100 clones each. In the first repertoire, one clone is large and accounts for 91 percent of all cells (e.g. a leukemic clone); the other 99 clones are small and account for the remaining 9 percent. In the second repertoire, all 100 clones are equally common, each accounting for 1 percent of cells. The Shannon entropies of the two repertoires are 1.0 bit and 6.6 bits. Entropy is converted to an effective number—^1D—by exponentiation: the effective number of clones in the first repertoire is $2^{1.0}=2$, while in the second repertoire it is $2^{6.6}=100$. Thus per entropy, the first repertoire can be thought of as "effectively" consisting of just two clones: the 99 rare clones collectively count the same as the one large clone. In other words, the first repertoire has the same effective diversity as a repertoire that consists of just two clones that are equally common. The second repertoire already consists of clones that are equally common, so the effective number of clones in this repertoire, $2^{6.6}=100$, is the same as its species richness. Diversity with similarity is interpreted in a similar fashion: a repertoire with a ^qD_s of n species has the same effective diversity as a repertoire with n species that are equally common (as above), with the additional constraint that these species are now also completely unrelated to/dissimilar from each other.

Figure 1

Figure 2

Figure 3

679 Figure 4

Figure 5

 Figure 6

Figure 7

Figure 8

Figure 9

a **Target-binding landscape** | Low-affinity mountains vs. high-affinity peaks

b **Memory IgH** | Many mountains/peaks → high functional/predicted diversity

c **TCRβ** | Few mountains/many peaks → low functional/high predicted diversity

d **Naïve IgH** | Some mountains/no peaks → medium functional diversity/no prediction

References

Almendro, V., Kim, H.J., Cheng, Y.-K., Gönen, M., Itzkovitz, S., Argani, P., van Oudenaarden,
A., Sukumar, S., Michor, F., and Polyak, K. (2014). Genetic and phenotypic diversity in breast
tumor metastases. Cancer Res *74*, 1338–1348.

Arnaout, R., Lee, W., Cahill, P., Honan, T., Sparrow, T., Weiand, M., Nusbaum, C., Rajewsky,
K., and Koralov, S.B. (2011). High-Resolution Description of Antibody Heavy-Chain Repertoires
in Humans. PLOS ONE *6*, e22365.

Bachmann, M.F., Kündig, T.M., Kalberer, C.P., Hengartner, H., and Zinkernagel, R.M. (1994).
How many specific B cells are needed to protect against a virus? J. Immunol. *152*, 4235–4241.

Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N.,
and Bourne, P.E. (2000). The Protein Data Bank. Nucleic Acids Res. *28*, 235–242.

Bild, D.E., Bluemke, D.A., Burke, G.L., Detrano, R., Diez Roux, A.V., Folsom, A.R., Greenland,
P., Jacob, D.R., Kronmal, R., Liu, K., et al. (2002). Multi-Ethnic Study of Atherosclerosis:
objectives and design. Am. J. Epidemiol. *156*, 871–881.

Botta-Dukát, Z. (2018). The generalized replication principle and the partitioning of functional
diversity into independent alpha and beta components. Ecography *41*, 40–50.

de Bourcy, C.F.A., Angel, C.J.L., Vollmers, C., Dekker, C.L., Davis, M.M., and Quake, S.R.
(2017). Phylogenetic analysis of the human antibody repertoire reveals quantitative signatures
of immune senescence and aging. Proc Natl Acad Sci U S A *114*, 1105–1110.

Britanova, O.V., Putintseva, E.V., Shugay, M., Merzlyak, E.M., Turchaninova, M.A., Staroverov,
D.B., Bolotin, D.A., Lukyanov, S., Bogdanova, E.A., Mamedov, I.Z., et al. (2014). Age-related
decrease in TCR repertoire diversity measured with deep and normalized sequence profiling. J.
Immunol. *192*, 2689–2698.

Bunge, J., and Fitzpatrick, M. (1993). Estimating the Number of Species: A Review. Journal of
the American Statistical Association *88*, 364–373.

Chao, A., Chiu, C.-H., Villéger, S., Sun, I.F., Thorn, S., Lin, Y., Chiang, J.-M., and B. Sherwin,
W. (2018). An attribute-diversity approach to functional diversity, functional beta diversity, and
related (dis)similarity measures. Ecological Monographs.

Chao, A., Chiu, C.-H., Villéger, S., Sun, I.-F., Thorn, S., Lin, Y.-C., Chiang, J.-M., and Sherwin,
W.B. An attribute-diversity approach to functional diversity, functional beta diversity, and related
(dis)similarity measures. Ecological Monographs *0*.

Chen, Z.J., Wheeler, C.J., Shi, W., Wu, A.J., Yarboro, C.H., Gallagher, M., and Notkins, A.L.
(1998). Polyreactive antigen-binding B cells are the predominant cell type in the newborn B cell
repertoire. Eur. J. Immunol. *28*, 989–994.

Chiu, C.-H., and Chao, A. (2014). Distance-Based Functional Diversity Measures and Their
Decomposition: A Framework Based on Hill Numbers. PLOS ONE *9*, e100014.

DeWitt, W.S., Lindau, P., Snyder, T.M., Sherwood, A.M., Vignali, M., Carlson, C.S., Greenberg,
P.D., Duerkopp, N., Emerson, R.O., and Robins, H.S. (2016). A Public Database of Memory
and Naive B-Cell Receptor Sequences. PLoS ONE *11*, e0160853.

Dunbar, J., Krawczyk, K., Leem, J., Baker, T., Fuchs, A., Georges, G., Shi, J., and Deane, C.M.
(2014). SAbDab: the structural antibody database. Nucleic Acids Res. *42*, D1140-1146.

Emerson, R.O., DeWitt, W.S., Vignali, M., Gravley, J., Hu, J.K., Osborne, E.J., Desmarais, C.,
Klinger, M., Carlson, C.S., Hansen, J.A., et al. (2017). Immunosequencing identifies signatures
of cytomegalovirus exposure history and HLA-mediated effects on the T cell repertoire. Nat
Genet *49*, 659–665.

Emery, V. (2001). Investigation of CMV disease in immunocompromised patients. J Clin Pathol
54, 84–88.

Fairlie-Clarke, K.J., Shuker, D.M., and Graham, A.L. (2009). Why do adaptive immune
responses cross-react? Evol Appl *2*, 122–131.

Frank, S.A. (2002). Immunology and Evolution of Infectious Disease (Princeton (NJ): Princeton
University Press).

Gibson, K.L., Wu, Y.-C., Barnett, Y., Duggan, O., Vaughan, R., Kondeatis, E., Nilsson, B.-O.,
Wikby, A., Kipling, D., and Dunn-Walters, D.K. (2009). B-cell diversity decreases in old age and
is correlated with poor health status. Aging Cell *8*, 18–25.

Greiff, V., Bhat, P., Cook, S.C., Menzel, U., Kang, W., and Reddy, S.T. (2015). A bioinformatic
framework for immune repertoire diversity profiling enables detection of immunological status.
Genome Med *7*.

Heindl, A., Lan, C., Rodrigues, D.N., Koelble, K., and Yuan, Y. (2016). Similarity and diversity of
the tumor microenvironment in multiple metastases: critical implications for overall and
progression-free survival of high-grade serous ovarian cancer. Oncotarget *7*, 71123–71135.

Hill, M.O. (1973). Diversity and Evenness: A Unifying Notation and Its Consequences. Ecology
54, 427–432.

Holodick, N.E., Rodríguez-Zhurbenko, N., and Hernández, A.M. (2017). Defining Natural
Antibodies. Front Immunol *8*.

Hopf, T.A., Ingraham, J.B., Poelwijk, F.J., Schärfe, C.P.I., Springer, M., Sander, C., and Marks,
D.S. (2017). Mutation effects predicted from sequence co-variation. Nat. Biotechnol. *35*, 128–
135.

Hopkins, A.C., Yarchoan, M., Durham, J.N., Yusko, E.C., Rytlewski, J.A., Robins, H.S., Laheru,
D.A., Le, D.T., Lutz, E.R., and Jaffee, E.M. (2018). T cell receptor repertoire features associated
with survival in immunotherapy-treated pancreatic ductal adenocarcinoma. JCI Insight *3*.

Jankauskaite, J., Jiménez-García, B., Dapkunas, J., Fernández-Recio, J., and Moal, I.H. (2018).
SKEMPI 2.0: An updated benchmark of changes in protein-protein binding energy, kinetics and
thermodynamics upon mutation. Bioinformatics.

Jiang, N., He, J., Weinstein, J.A., Penland, L., Sasaki, S., He, X.-S., Dekker, C.L., Zheng, N.-Y.,
Huang, M., Sullivan, M., et al. (2013). Lineage structure of the human antibody repertoire in
response to influenza vaccination. Sci Transl Med *5*, 171ra19.

Jost, L. (2007). Partitioning diversity into independent alpha and beta components. Ecology *88*,
2427–2439.

Ju, C.-H., Blum, L.K., Kongpachith, S., Lingampalli, N., Mao, R., Brodin, P., Dekker, C.L., Davis,
M.M., and Robinson, W.H. (2018). Plasmablast antibody repertoires in elderly influenza vaccine
responders exhibit restricted diversity but increased breadth of binding across influenza strains.
Clin. Immunol. *193*, 70–79.

Kaplinsky, J., and Arnaout, R. (2016). Robust estimates of overall immune-repertoire diversity

from high-throughput measurements on samples. Nat Commun *7*, 11881.

Kaplinsky, J., Li, A., Sun, A., Coffre, M., Koralov, S.B., and Arnaout, R. (2014). Antibody

repertoire deep sequencing reveals antigen-independent selection in maturing B cells. Proc.

Natl. Acad. Sci. U.S.A. *111*, E2622-2629.

Koopmans, R., and Schaeffer, M. (2013). De-composing diversity: In-group size and out-group

entropy and their relationship to neighbourhood cohesion (WZB Berlin Social Science Center).

Langman, R.E., and Cohn, M. (1987). The E-T (elephant-tadpole) paradox necessitates the

concept of a unit of B-cell function: the protection. Mol. Immunol. *24*, 675–697.

Lee, J., Natarajan, M., Nashine, V.C., Socolich, M., Vo, T., Russ, W.P., Benkovic, S.J., and

Ranganathan, R. (2008). Surface Sites for Engineering Allosteric Control in Proteins. Science

322, 438–442.

Leem, J., de Oliveira, S.H.P., Krawczyk, K., and Deane, C.M. (2018). STCRDab: the structural

T-cell receptor database. Nucleic Acids Res. *46*, D406–D412.

Leinster, T., and Cobbold, C.A. (2012). Measuring diversity: the importance of species similarity.

Ecology *93*, 477–489.

Levy, E.D. (2010). A simple definition of structural regions in proteins and its use in analyzing

interface evolution. J. Mol. Biol. *403*, 660–670.

Li, K., Bihan, M., Yooseph, S., and Methé, B.A. (2012). Analyses of the Microbial Diversity

across the Human Microbiome. PLOS ONE *7*, e32118.

Lunzer, M., Golding, G.B., and Dean, A.M. (2010). Pervasive cryptic epistasis in molecular

evolution. PLoS Genet. *6*, e1001162.

Macarthur, R.H. (1965). Patterns of Species Diversity. Biological Reviews *40,* 510–533.

Marion, Z.H., Fordyce, J.A., and Fitzpatrick, B.M. (2015). Extending the Concept of Diversity

Partitioning to Characterize Phenotypic Complexity. Am. Nat. *186*, 348–361.

Messaoudi, I., Lemaoult, J., Guevara-Patino, J.A., Metzner, B.M., and Nikolich-Zugich, J.
(2004). Age-related CD8 T cell clonal expansions constrict CD8 T cell repertoire and have the
potential to impair immune defense. J. Exp. Med. *200*, 1347–1358.

Morris, E.K., Caruso, T., Buscot, F., Fischer, M., Hancock, C., Maier, T.S., Meiners, T., Müller,
C., Obermaier, E., Prati, D., et al. (2014). Choosing and using diversity indices: insights for
ecological applications from the German Biodiversity Exploratories. Ecol Evol *4*, 3514–3524.

Notkins, A.L. (2004). Polyreactivity of antibody molecules. Trends Immunol. *25*, 174–179.

Obar, J.J., Khanna, K.M., and Lefrançois, L. (2008). Endogenous naive CD8+ T cell precursor
frequency regulates primary and memory responses to infection. Immunity *28*, 859–869.

Perelson, A.S., and Oster, G.F. (1979). Theoretical studies of clonal selection: minimal antibody
repertoire size and reliability of self-non-self discrimination. J. Theor. Biol. *81*, 645–670.

Pons, J., Rajpal, A., and Kirsch, J.F. (1999). Energetic analysis of an antigen/antibody interface:
alanine scanning mutagenesis and double mutant cycles on the HyHEL-10/lysozyme
interaction. Protein Sci. *8*, 958–968.

Qi, Q., Liu, Y., Cheng, Y., Glanville, J., Zhang, D., Lee, J.-Y., Olshen, R.A., Weyand, C.M.,
Boyd, S.D., and Goronzy, J.J. (2014). Diversity and clonal selection in the human T-cell
repertoire. Proc Natl Acad Sci U S A *111*, 13139–13144.

Robins, H.S., Srivastava, S.K., Campregher, P.V., Turtle, C.J., Andriesen, J., Riddell, S.R.,
Carlson, C.S., and Warren, E.H. (2010). Overlap and effective size of the human CD8+ T cell
receptor repertoire. Sci Transl Med *2*, 47ra64.

Salinas, V.H., and Ranganathan, R. (2018). Coevolution-based inference of amino acid
interactions underlying protein function. Elife *7*.

Scheiner, S.M. (2012). A metric of biodiversity that integrates abundance, phylogeny, and
function. Oikos *121*, 1191–1202.

Schober, K., Buchholz, V.R., and Busch, D.H. (2018). TCR repertoire evolution during
maintenance of CMV-specific T-cell populations. Immunol. Rev. *283*, 113–128.

Smith, D.J., Forrest, S., Hightower, R.R., and Perelson, A.S. (1997). Deriving Shape Space
Parameters from Immunological Data. Journal of Theoretical Biology *189*, 141–150.

Taraska, J.W. (2015). Cell biology of the future: Nanometer-scale cellular cartography. J Cell
Biol *211*, 211–214.

Taylor, M.G., Rajpal, A., and Kirsch, J.F. (1998). Kinetic epitope mapping of the chicken
lysozyme.HyHEL-10 Fab complex: delineation of docking trajectories. Protein Sci. *7*, 1857–
1867.

Vollmers, C., Sit, R.V., Weinstein, J.A., Dekker, C.L., and Quake, S.R. (2013). Genetic
measurement of memory B-cell recall using antibody repertoire sequencing. Proc. Natl. Acad.
Sci. U.S.A. *110*, 13463–13468.

Weinstein, J.A., Jiang, N., White, R.A., Fisher, D.S., and Quake, S.R. (2009). High-throughput
sequencing of the zebrafish antibody repertoire. Science *324*, 807–810.

Whittaker, J., Groth, A.V., Mynarcik, D.C., Pluzek, L., Gadsbøll, V.L., and Whittaker, L.J. (2001).
Alanine scanning mutagenesis of a type 1 insulin-like growth factor receptor ligand binding site.
J. Biol. Chem. *276*, 43980–43986.

Xu, J.L., and Davis, M.M. (2000). Diversity in the CDR3 region of V(H) is sufficient for most
antibody specificities. Immunity *13*, 37–45.

Zarnitsyna, V.I., Evavold, B.D., Schoettle, L.N., Blattman, J.N., and Antia, R. (2013). Estimating
the diversity, completeness, and cross-reactivity of the T cell repertoire. Front Immunol *4*, 485.

PyMOLThe PyMOL Molecular Graphics System, Version 1.8, Schrödinger, LLC.