Evolution of the HIV-1 RRE during natural infection reveals key nucleotide changes that

correlate with altered structure and increased activity over time.

Chringma Sherpa[a]*, Patrick E. H. Jackson [b,c]*, Laurie R. Gray[c,d], Kathryn Anastos[e],

Stuart F. J. Le Grice[a]#, Marie-Louise Hammarskjold[c,d], David Rekosh[c,d]#

[a]Basic Research Laboratory, Center for Cancer Research, National Cancer Institute,

National Institute of Health, Frederick, Maryland, USA

[b]Division of Infectious Diseases and International Health, Department of Medicine,

University of Virginia, Charlottesville, Virginia, USA

[c]Myles H. Thaler Center for HIV and Human Retrovirus Research, University of Virginia,

Charlottesville, Virginia, USA

[d]Department of Microbiology, Immunology and Cancer Biology, University of Virginia,

Charlottesville, Virginia, USA

[e]Albert Einstein College of Medicine and Montefiore Medical Center, Department of

Medicine, Bronx, New York, USA

Running Head: HIV-1 RRE structural and functional evolution.

#Address correspondence to: David Rekosh, dr4u@virginia.edu, or Stuart Le Grice,

legrices@mail.nih.gov.

*C.S. and P.E.H.J. contributed equally to this work.

ABSTRACT

Abstract.

The HIV-1 Rev Response Element (RRE) is a *cis*-acting RNA element characterized by multiple stem-loops. Binding and multimerization of the HIV Rev protein on the RRE promotes nucleocytoplasmic export of incompletely spliced mRNAs, an essential step in HIV replication. Most of our understanding of the Rev-RRE regulatory axis comes from studies of lab-adapted HIV clones. However, in human infection, HIV evolves rapidly and mechanistic studies of naturally occurring Rev and RRE sequences are essential to understanding this system. We previously described the functional activity of two RREs found in circulating viruses in a patient followed during the course of HIV infection. The "early" RRE was less functionally active than the "late" RRE despite differing in sequence by only four nucleotides. In this study, we describe the sequence, function, and structural evolution of circulating RREs in this patient using plasma samples collected over six years of untreated infection. RRE sequence diversity varied over the course of infection with evidence of selection pressure that led to sequence

convergence as disease progressed. An increase in RRE functional activity was

observed over time, and a key mutation was identified that correlates with a major

conformational change in the RRE and increased functional activity. Additional

mutations were found that may have contributed to increased activity as a result of

greater Shannon entropy in RRE stem-loop II, which is key to primary Rev binding.

Importance.

HIV-1 replication requires interaction of the viral Rev protein with a *cis*-acting regulatory

RNA, the Rev Response Element (RRE), whose sequence changes over time during

infection within a single host. In this study, we show that the RRE is subject to selection

pressure and that RREs from later time points in infection tend to have higher functional

activity. Differences in RRE functional activity are attributable to specific changes in

RNA structure. Our results suggest that RRE evolution during infection may be

important for HIV pathogenesis and that efforts to develop therapies acting on this viral

pathway should take this into account.

INTRODUCTION

All retroviruses produce mRNAs that retain introns, and these mRNAs must be

exported to the cytoplasm for packaging into viral particles and translation of essential

viral proteins. Eukaryotic cells have RNA surveillance mechanisms that would normally

restrict nucleocytoplasmic export and translation of these mRNA species (1). Thus,

retroviruses have evolved specific mechanisms to overcome this restriction (2). In HIV,

the Rev protein, in conjunction with its RNA binding partner, the Rev Response Element

(RRE), mediates this important function (3-6).

HIV Rev contains several well-characterized functional domains that facilitate

shuttling between the nucleus/nucleolus and the cytoplasm by accessing cellular

pathways for nuclear import and export. One of these domains is a basic, arginine-rich

motif that functions as a nuclear/nucleolar localization signal (7-10). This domain also

serves as an RNA-binding domain that binds specifically to the RRE (11, 12). It is

flanked on both sides by oligomerization domains that are required for Rev

multimerization on the RRE (13, 14). Towards the carboxy terminus is a leucine-rich

domain which docks on the Crm1-RanGTP complex (15) and functions as the nuclear

export signal (NES) (12, 16, 17).

The RRE is a *cis*-acting RNA element located in a highly conserved and

functionally important region of *env* at the N-terminus of the gp41 fusion protein (3-6). At

this position, it is present in all HIV RNAs that retain introns. The minimal functional

RRE, or "short" RRE, has been mapped in infectious laboratory clones to a 234-nt

region (see Figure 3). It forms a highly branched structure where the 5' and 3' ends pair

to form stem-loop I (SL-I) (18). SL-I opens into a central loop, from which several

additional stem loops branch out. Stem loop II contains stem IIA that branches out of

the central loop and opens into a three-way junction. The junction opens into stem-loops

IIB and IIC (19, 20). SL-III is a small stem-loop that comes off of the central loop, while

SL-IV and SL-V can adopt alternative topologies (21). Maximal functional activity of the

RRE requires a somewhat larger structure, often referred to as the "long" (351-nt) RRE,

characterized by an extended SL-I (18, 22, 23).

Primary RRE binding initiates cooperative assembly of additional Rev molecules

(about 6-8) in a process that requires electrostatic Rev-RRE interactions and

hydrophobic Rev-Rev interactions (13, 18, 24-32). Rev oligomerization on the RRE

increases its binding affinity ~500 fold (30). Oligomerization also arranges the NES

domains for binding to the RAN-GTP bound Crm1 dimer (15) forming an export-

competent ribonucleoprotein complex (33, 34). The complex is targeted to the nuclear

pore, where it interacts with the nucleoporins, resulting in its translocation to the

cytoplasmic side. Once in the cytoplasm, RRE-containing mRNAs are translated into

Gag, Gag-Pol, Env, and accessory viral proteins.

Molecular details of the Rev-RRE pathway have been delineated mostly from

studies of lab-adapted HIV clones. However, there is mounting evidence that subtle

variation in Rev and RRE sequences among primary isolates may contribute to

pathogenesis. Differences in Rev-RRE functional activity up to 24-fold were observed

for naturally occurring viruses in different patients (35). In addition, a study performed

with a Thai cohort demonstrated RRE changes over the course of infection, and higher

RRE activity was associated with a more rapid CD4 count decline (36). Low Rev activity

has also been associated with slower disease progression (37, 38) and reduced

susceptibility of infected cells to T-cell killing (39). In equine infectious anemia virus, a

related lentivirus that also utilizes the Rev/RRE axis, functional evolution of Rev has

been observed during infection, and Rev activity has been found to correlate with

disease state in ponies (40-42). Further studies of the structural and functional evolution

of the Rev-RRE system in natural infection are necessary to understand the role that

this regulatory axis plays in adaptation of HIV to diverse immune environments, and this

may benefit development of Rev-RRE targeted therapeutics.

We previously investigated the activity of Rev-RRE cognate pairs from HIV

isolated from five different patients at two time-points during their course of infection

(43). The sequences were obtained by single genome sequencing of viruses from blood

plasma samples and their functional activity was determined using a sub-genomic

reporter assay. We observed significant activity differences between Rev-RRE cognate

pairs from different patients and from different time points in the same patient. Evolution

of Rev and the RRE observed in patient SC3 was particularly striking. In this patient, the

RRE converged on one predominant sequence at the later time point (the M57A RRE,

designated here as V20-1), suggesting it was subject to strong selective pressures.

Furthermore, this RRE had only four nucleotide changes (mut 1-4) relative to the early

time point RRE (M0-A RRE, designated here as V10-2) but was 2-3 fold more

functionally active. Gel mobility shift assays revealed that the V20-1 RRE promoted Rev

multimerization at a lower concentration of Rev protein compared to the V10-2 RRE. It

was also notable that the predominant Rev sequence present at the early time point

persisted at the late time point, suggesting that the limited nucleotide changes in the

RRE were the major driver of differential Rev-RRE activity.

In the current study, we conducted a detailed examination of RRE evolution in

the blood plasma of patient SC3 using samples collected at six-month intervals, from

the time viral RNA was first detected through year six of infection. DNA deep

sequencing, selective 2′ hydroxyl acylation analyzed by primer extension (SHAPE)

chemical probing, and a Rev-RRE functional assay were used to determine sequence

evolution, RRE secondary structure, and Rev-RRE functional activity over time and to
explore the mechanism underlying the observed activity differences. This study
highlights, for the first time, the structure-function relationship of longitudinal RRE
sequence evolution in a single patient and the underlying molecular mechanisms.

RESULTS:

RRE sequence evolution in an HIV-infected individual followed over many years.

Plasma samples from patient SC3 were collected over a period of 10 years
during standardized visits spaced about 6 months apart as part of the Women's
Interagency HIV Study (WIHS) Consortium in the Bronx, New York. Patient SC3 was
enrolled in the WIHS cohort prior to HIV seroconversion, was followed through visit 20,
and never received antiretroviral therapy (ART). Samples for this patient were available
for all time points with the exception of four missed visits (2, 3, 17 and 18). Plasma
samples from each WIHS visit were tested for p24 antibody and CD4 count (Figure 1).
Viral load was measured once the patient tested positive for p24 antibodies. In this
report, each visit is denominated as VXX with the numeral representing the visit
number. Although p24 antibodies were not detected under the original WIHS protocol
until V10, a significant fall in her CD4 count was noted at V08 and we were able to
readily amplify HIV from plasma taken at this time point, suggesting infection between
V07 and V08. After a short-lived rebound at V09, the CD4 count continued to fall and
reached a nadir of 73 cells/μl at V20. Seroconversion and viremia were noted in the
data obtained from the WIHS protocol at V10, and the viral load continued to rise
through V20 reaching a peak of 1.4 x10^6 copies/ml. As expected without ART treatment,

we observed a general increase in viral load concomitant with a decline in CD4 count as

disease progressed.

We investigated RRE sequence evolution in viruses isolated from V08 and

onwards. A total of 12 plasma samples representing visits 7, 8, 9, 10, 11, 12, 13, 14, 15,

16, 19 and 20 were received from the WIHS consortium. Viral RNA isolation was

attempted from each sample and, if successful, RRE sequences were determined by

next-generation sequencing. As stated above, the first visit to yield HIV sequences that

could be amplified by PCR was V08. The prevalence of each RRE sequence at a given

visit was calculated as a percentage of the total number of sequences present at that

time point (Figure S1). Sequences present with a frequency of <5% were then excluded

from consideration, as it was not possible to distinguish true rare RREs from artifacts

due to PCR errors, and the prevalence of significant variants was recalculated after

exclusion of these minority sequences (Figures 2, S2). RRE sequences were assigned

a code in the form VXX-Y where XX refers to the visit number and Y refers to the

prevalence rank order of that sequence.

As shown in Figure 1, data obtained from the WIHS indicated that patient SC3

seroconverted to HIV-positive status at V10. Thus, in our previous study, V10 was

believed to represent a timepoint within six months of infection and was designed month

0 (M0). An RRE was identified at this visit by single genome sequencing that was

believed to be the major species present at that time and designated an "early" RRE.

Our deep DNA sequencing results showed that this "early" RRE sequence represented

17% of the sequences present at that visit and it is therefore labeled as V10-2 in the

present study. This sequence is very similar to the single founder RRE sequence found

at V08 (V08-1), differing only by a single nucleotide near the 3' end and mapping to the

right-hand side of SL-1 (designated here as mut 5). Our previous study also identified a

single RRE from V20 which we designated a "late" RRE. This RRE is identical with the

V20-1 sequence generated by deep sequencing.

Since the V10-2 RRE previously described as the "early" RRE has been studied

intensively by us, we decided to compare its properties to the founder V08-1 RRE. Both

RREs were transcribed *in vitro* and their secondary structures chemically probed by

1M7 (1-methyl-7-nitroisatoic anhydride) using selective 2′ hydroxyl acylation analyzed by

primer extension and mutational profiling (SHAPE-MaP) (44-50). In this procedure,

RNAs are first modified with chemical reagents that selectively acylate unpaired

ribonucleotides at their 2′-hydroxyl positions, then reverse transcribed under conditions

that introduce mutations in cDNA opposite the sites of modification (47, 49). The

position and frequencies of these mutations are used to create reactivity profiles

indicating which RNA nucleotides are likely to be single- or double-stranded (Figure S3).

This information is in turn used to guide the RNAstructure software (51) to generate

lowest Gibbs free energy secondary structure models.

SHAPE-MaP studies revealed that V08-1 and V10-2 form similar structures with

only minor local differences (Figure 3A and 3B). These structures are similar to the

previously reported ARv2/SF2 RRE structure (22) where a part of the central loop (nt

125-130) between SL-III and SL-IV paired with nucleotides from the upper stem of SL-I

(193-198) forming a stem that bridges the central loop and SL-IV and SL-V.

To determine activity of each of these RREs, as well as V20-1, each RNA was

cloned into a two-color fluorescent-based proviral reporter vector (Jackson et al,

manuscript in preparation) which carried inactivating mutations in *rev*. RRE activity was

measured by co-transfecting a separate plasmid that expressed the previously

described M0-B/M57A-SC3 Rev. This Rev protein has been shown by single genome

sequencing to pair with the V10-2 RRE (43). It was also the unique founder Rev

sequence present at V08. Unexpectedly, this sequence still persisted at V20 and was

found in single genomes with the V20-1 RRE (43). The reporter construct expressed

eGFP from the intron-containing Rev-dependent *gag* mRNA and TagBFP from the

intron-less Rev-independent *nef*-like transcript. Thus, in this assay, the ratio of the

fluorescent signals (eGFP:TagBFP) is a measure of Rev-RRE functional activity. All

three RREs displayed clear Rev responsiveness as the fluorescent ratio in the non-Rev

containing transfections was at least 500-fold less than in Rev-containing transfections

(data not shown). Despite their local structural differences, both V08-1 and V10-2 RRE

showed similar levels of RRE activity (p=0.999) (Figure 4). Therefore, although mutation

5 arose after infection and increased in prevalence through V20, it did not increase RRE

functional activity or significantly change RRE structure. Additionally, activities of V08-1

and V10-2 RREs were substantially less than that of V20-1 (Figure 4) (p=0.022 and

p=0.021, respectively), consistent with our previous finding that V20-1 RRE is more

active (43). Notably, we also replicated the finding that V20-1 is significantly more

active than V10-2 using a lentiviral vector system, where packaging of genomic RNA

and titer is dependent on Rev-RRE function (Figure S4) (35, 52) .

V10-2 and V20-1 RREs form distinctly different secondary structures.

We previously reported that there are only four single nucleotide changes (mut 1-

4) between the V10-2 and V20-1 RREs (43). A fifth mutation (mut 5) is present in both

V10-2 and V20-1 and therefore did not score as a difference between the two RREs

identified as "early" and "late" in our previous study. On a canonical 234-nt five stem-

loop (5SL) RRE structure (21), these changes map near the apical loop of SL-IIB [nt 61:

mutation (mut) 1] and SL-IIC [nt 84: mutation (mut) 2], the central loop region between

SL-V and SL-I [nt 194: mutation (mut) 3], and at the base of stem of SL-I [nt 228:

mutation (mut) 4] (see Figures 3B and 3D). We next explored whether the differences in

activity between the two RREs could be attributed to structural differences caused by

any individual nucleotide change.

In vitro transcribed and folded V10-2 and V20-1 short RREs migrated at a

significantly different rate when analyzed by native polyacrylamide gel electrophoresis

(Figure 3E), suggesting major structural differences. Consistent with the gel migration

data, SHAPE-MaP of the two 234-nt RREs revealed that the V10-2 and V20-1 RREs

adopt distinctly different conformations (Figures 3B and 3D). Unlike V10-2, which had

part of the central loop region base-paired, V20-1 formed the canonical 5SL structure

where five distinct stem-loop structures (SL-I to SL-V) radiate out directly from the single

stranded central loop.

Closer analysis of the location of each of the four nucleotide changes suggested

that the major structural shift in the two RREs might have been caused by mut 3. The

nucleotide at the mut 3 position (G) is base-paired in the bridging stem in V10-2. In V20-

1, it mutates to C disrupting base pairing at this position thereby destabilizing the stem

formed by nt 125-130 and 193-198. Consequently, nt 34-36 were significantly more

reactive while nt 125, 127, and 130 were significantly less reactive in V10-2 compared

to V20-1. Mut 1 and mut 2 produced more limited local structural changes, i.e. opening

and formation of the base-pair near the apical loop of SL-IIB and SL-IIC in V20-1

relative to V10-2. Since mut 4 was located at the primer-hybridization site (see the

PCR1 step of SHAPE-MaP in Methods) for the 234-nt RREs, we were unable to assess

its effect on the secondary structure.

A 3-dimensional structural analysis of a 351-nt RRE by small angle X-ray

scattering has shown that the extended SL-I folds back on regions in and around SL-II

to expose a cryptic Rev-binding site (22). Data obtained in connection with our previous

study (43), showed that the extended SL-I of the 351-nt V10-2 and V20-1 RREs has an

additional four nucleotide changes at nucleotide positions 13, 27, 29, and 57 (using the

351-nt RRE numbering system). We therefore investigated if these nucleotide changes

in the extended SL-I affect RRE structure. Additionally, probing these extended RREs

provided an opportunity to determine the effect of mut 4 on their structure, as this

position no longer fell on the primer hybridization site during SHAPE analysis of the

351-nt RREs. The long RREs were investigated by capillary electrophoresis-based

SHAPE (Figure 5) (CE-SHAPE) using N-methylisatoic anhydride (NMIA) as the

electrophile (53, 54). CE-SHAPE differs from SHAPE-MaP in that the chemically

modified nucleotides are identified as stops during primer extension. CE-SHAPE

analysis of the 351-nt V10-2 and V20-1 revealed that mut 4 did not produce any

structural changes between V10-2 and V20-1 RREs, as it changed a U-A base pair to

U-G. Additionally, the nucleotide differences in the extended SL-I region of these RREs

produced only highly localized structural changes, if any. Changes at nt 13 and nt 57

produced no structural changes at all, whereas nucleotide differences at nt 27 and nt 29

induced opening of base pairs at these positions in V20-1. As these mutations in SL-1

did not alter the structure of the comparable region of the 234-nt RRE, subsequent

studies investigated their effects only in the short RRE forms.

Structural and functional analysis of patient SC3 RREs containing individual

single nucleotide mutations.

We next investigated the contribution of each of the four nucleotides which

differed between the V10-2 and the V20-1 RREs to alterations in structure and

functional activity. To this end, we created four new synthetic 234-nt RRE sequences,

each containing one of the single nucleotide changes in a V10-2 RRE background. The

new RREs containing mut 1, 2, 3, and 4 were respectively termed M1, M2, M3, and M4.

The secondary structure of each RRE was probed by SHAPE-MaP (Figure 6A-D). As

predicted, M1, M2, and M4 RREs formed V10-2 like structures, whereas M3 RRE

formed a V20-1 like structure. M1 RRE resembled V10-2 RRE with the exception that

the G to A mutation eliminated a base pair adjacent to the SL-II apical loop. Similarly,

M2 RRE differed from V10-2 RRE only in the formation of an extra base pair next to the

apical loop of SL-IIC. M4 RRE formed a structure identical to V10-2 RRE, since the mut

4 A to G change preserved non-canonical base pairing at this position by replacing A-U

with a G-U base pair. The major structural shift involving the central loop, SL-IV, SL-V,

and SL-I between the V10-2 and V20-1 RREs was reproduced by the single G to C

mutation in the M3 RRE. Therefore, we reasoned that mut 3 might be the major driver of

the functional difference between the V10-2 and V20-1 RREs.

We next measured the contribution of each mutation to the functional activity

difference between the V10-2 and V20-1 RREs (Figure 6E). Activity of RREs M1-M4

was determined using the fluorescent-based transient proviral reporter with the same

cognate SC3 Rev used in Figure 4. As before, in the presence of SC3 Rev, functional

activity of V20-1 RRE was significantly higher than that of V10-2 RRE (p=0.001). Among

all single SC3 mutants, only M3 displayed activity that was statistically significantly

higher than V10-2 (p=0.072) even though it did not quite reach the activity level of V20-

1. M1 and M2 RREs were only slightly more active than V10-2 but these differences did

not reach statistical significance. Activity of M4 RRE was also not statistically

distinguishable from that of V10-2. However, M4 had a tendency towards lower, rather

than higher, functional activity. Thus, the difference in functional activity between V10-2

and V20-1 could not be explained by any single nucleotide mutation.

Structural and functional analysis of selected RRE sequences derived from

patient SC3 at different time points.

To further understand how specific sequence changes between V10-2 and V20-1

RREs affected functional activity, we studied intermediate RRE sequences from viruses

that arose during disease progression in patient SC3. SC3 RREs that allowed us to test

combinations of these mutations in the order they appeared in the natural isolates were

selected from this evolutionary data set (Figure 2; Figure S2). The secondary structure

and functional activity of these RREs were then determined by SHAPE-MaP and the

fluorescent-based reporter assay, respectively.

The position designated mut 3 starts as a G in V08-1, remains a G in V10-2,

changes to A in many sequences in V09 through V16, and then to C in V19 and V20. As

this is the position that causes the major structural shift between V10-2 and V20-1, we

sought to determine if the A at the mut 3 position would result in the same structure as

its replacement by C. To test this, we determined the secondary structure of V09-2 RRE

by SHAPE-MaP and compared it to the structure of V08-1 and V10-2 (Figure 3A-C).

The V09-2 RRE sequence is identical to V08-1 except at the mut 3 position, where the

G in V08-1 is replaced with A. We hypothesized that if disruption of the G-C base-pair at

this position induced the structural shift observed between V10-2 and V20-1, then such

a shift should also be reflected between V08-1 and V09-2. Indeed, the secondary

structure of V09-2 resembled that of V20-1 RRE, confirming our hypothesis. V09-2 RRE

was also slightly more active than V08-1 RRE (though this trend did not reach statistical

significance) (p=0.116) and V10-2 (p=0.027) (Figure 7). Consistent with our previous

observation with M3 RRE, the structural shift invoked in V20-1 and V09-2 relative to

V08-1 and V10-2 shows a trend towards increased SC3 RRE activity.

Mut 2 was first noted in the sample from V11 and was present in all RRE

sequences from subsequent visits. We tested the V11-1 RRE as it provided the

opportunity of investigating the combination of mut 2 together with A in the position of

mut 3. Although it was predicted to have higher activity than M3 and V09-2 RRE, V11-1

RRE had similar level of RRE activity as the founder RRE, V08-1, suggesting that while

A at the position of mut 3 increases functional activity alone, it does not do so in the

context of the additional mut 2 (p=1.000).

We first detected mut 1 in plasma samples from visit 13. This mutation was

frequently observed on subsequent visits and was present in the majority of RREs from

visit 15 onwards. The V13-1 RRE contained not only mut 1 but also mut 2 and A in the

mut 3 position. This RRE had significantly higher activity than both the V08-1 and V09-2

variants ($p<0.001$ for both comparisons). Therefore, mut 1 not only contributes to RRE

activity in itself but also rescues the activity of the combination of mut 2 and an A in the

mut 3 position.

We also tested the activity of V14-2, which allowed us to test the combined

phenotype of mut 1, mut 2, A in the mut 3 position, and mut 5. Activity of this RRE did

not significantly differ from V13-1, suggesting that acquisition of mut 5 does not

contribute to RRE activity ($p=0.332$). However, activity of the V20-1 RRE, which harbors

C in the position of mut 3 rather than the A of V13-1, as well the other mutations, was

significantly higher than V13-1 ($p<0.001$). This suggests that C in the mut 3 position is

important for imparting higher RRE activity when all other mutations are present. It is

also possible that although mut 4 does not contribute to RRE activity individually, it can

contribute to activity when combined with the other additional changes.

This set of SC3 RREs permitted examination of changes in functional activity

over time. Activities of V08-1, V10-2, and V11-1 were very similar, demonstrating that

the combinations of mutations observed in these sequences are not sufficient in

themselves to confer higher functional activity. V13-1 was the first tested RRE to arise,

other than V09-2, that showed significantly higher activity than V08-1 and it did so with

the combination of mut 1, mut 2, and A in the mut 3 position. The accumulation of

additional changes in V14-2 (addition of mut 5), V19-2 (addition of mut 4, the A to C

change at mut 3, and an additional A to G change at nt 142), and V20-1 (reversion of nt

142 to A) corresponded to a trend towards increasing functional activity at each step.

This steady trend suggests functional evolution of the RRE in the later phase of the

disease course of patient SC3, as new combinations of mutations arose and those

conferring greater functional activity showed preferential selection.

Shannon entropy profile of SC3 RRE variants and single mutants.

Since the energetically most favorable secondary structures of SC3 RRE variants

and single mutants were insufficient to explain the functional activity differences

observed between the V08-1and V20-1 RREs, we next assessed the ability of the RREs

to adopt alternative conformers by evaluating their SHAPE-guided Shannon entropies

and base pairing probability at single nucleotide resolution. Shannon entropies were

calculated based on the probability for each base-pair appearing across all possible

structures predicted for the RNA. Regions with highly stable well-defined RNA

structures are characterized by lower Shannon entropies. Conversely, regions with high

Shannon entropy are likely to form alternative conformers.

We compared Shannon entropy profiles of the V08-1, V09-2, V10-2, V20-1, M1,

M2, M3, and M4 SC3 RREs (Figure 8). Most regions of these RREs exhibited low

Shannon entropy, suggesting their overall secondary structures are highly stable. This

result is consistent with the observation that these RRE RNAs migrate as a single

discrete band on a native agarose gel, suggesting a high degree of structural

homogeneity (Figure S5).

However, we observed differences in Shannon entropy values at two different

regions. Specifically, values of the loop regions between SL-IIA and SL-IIB (Region I)

and between SL-IIB and SL-IIC (Region II) varied between RREs. The values for the

V20-1 RRE were, respectively, 5-6 fold and 4 fold higher than for the V08-1 and V10-2

RREs, suggesting that the nucleotides in these two regions of the V20-1 RRE are not

always single-stranded as depicted in the secondary structures generated by SHAPE-

MaP. This finding is further corroborated by SHAPE-MaP-guided base pairing

probability calculated for each of these RREs (Figures S6-S9), which shows that the two

higher entropy regions of V20-1 RRE can base-pair with each other with a probability of

10-80%. Both the Shannon entropy and pairing probability data suggest that these two

regions of V20-1 RRE are structurally dynamic, a feature that confers accessibility to

protein interaction with the RNA by serving as landing pads for protein cofactors (44).

This accessibility may have implications for Rev-binding, possibly contributing to the

higher activity of V20-1 RRE.

Furthermore, both the V09-2 and M3 RREs have 2-fold higher Shannon entropy

for the loop region between SL-IIA and SL-IIB (Region I) compared to V08-1 and V10-2

RREs. Their entropy values for the nucleotides between SL-IIB and SL-IIC (Region II),

however, did not change relative to V08-1 and V10-2. While both V09-2 and M3

undergo the structural shift characteristic of V20-1 and are modestly more active than

V08-1 and V10-2, the inability of V09-1 and M3 RREs to reach the degree of structural

flexibility of the V20-1 RRE in the internal loop of SL-II might explain why these RREs

are not as functionally active as V20-1. The Shannon entropy profile of M2 RRE is

similar to the M3 and V09-1 RREs, whereas M1 RRE has 2-fold higher Shannon

entropy than V08-1 and V10-2 in both regions of SL-II. The modestly higher entropy in

the M1, M2, and M3 RREs relative to V08-1 and V10-2 might help to explain their trend

towards higher functional activity (Figure 6). The M4 RRE, on the other hand, did not

show any increase in entropy over V08-1 and V10-2 and also displayed similar or lower

functional activity.

Taken together, our analysis suggests that both the major structural shift

observed between the V08-1 and V20-1 RREs, and the difference in structural

dynamism in the internal loop of SL-II contribute to the significant increase in functional

activity over time.

DISCUSSION

In this study, we describe a complex relationship between RRE structure and

functional activity using naturally occurring sequences obtained from a single patient.

This study is the first detailed examination of longitudinal evolution of the HIV-1 RRE

over about six years of naturally occurring infection and demonstrates clear selection

pressures acting on the RRE sequence with a tendency towards increased functional

activity over time. Increased activity can be explained by large-scale conformational

changes within the RRE and a decrease in base-pairing stability at the initial Rev-

binding site in SL-II.

The evolution of the RRE sequence is strongly suggestive of selection pressures

operating on the RRE itself. The single founder RRE sequence found at visit 8

diversifies initially, and by visit 20 converges on only a single circulating RRE sequence

differing from V08-1 by five single nucleotides. The early appearance and persistence of

mutations at only a few locations, as well as the reduced viral diversity by visit 20, argue

that the phenotype resulting from these changes was subject to strong selection

pressures. Functional activity analysis of the SC3 RREs suggests that the RREs with

higher activity were selected as disease progressed, with the early time-point RREs

(V08-1 and V10-2) displaying the lowest and the late time-point RRE (V20-1) displaying

the highest functional activity. The tendency towards increased functional activity

corresponded with gradual accumulation, in different combinations, of five mutations

that characterize V20-1 relative to V08-1.

There is no direct correspondence between specific mutations and functional

activity phenotypes. Changes at the mutation 3 position predicted to disrupt base-

pairing appear to result in an activity increase. However, V09-2 RRE containing this

change alone still has significantly lower activity than V20-1, showing that the other

mutations also contribute to a change in RRE activity in combination. While all

mutations 1-5 are necessary to achieve the highest activity, their contributions are not

merely additive. For example, V11-1 resembles V09-2 with the addition of mut 2, but it

does not have increased activity relative to the founder. Additionally, mutations which

have no impact on activity in isolation can increase activity in combination. For example,

mut 4 does not change activity alone, but is responsible for the increase in activity

between V14-2 and V20-1 in the context of the other four mutations.

Two types of structural changes were uncovered in this study that can explain

differences in activity: major conformational changes and regions of increased base-

pairing entropy. Previously, we have shown that alternative conformations of the

laboratory strain NL4-3 RRE display significantly different activities, with a four stem-

loop conformation supporting lower replicative capacity than a five stem-loop structure (21). In this study, we found that the higher activity RRE (V20-1) forms a five stem-loop conformation resembling that observed for NL4-3, while the lower activity RREs (V08-1 and V10-2) adopt a different conformation with a more collapsed central loop. As expected from functional analysis of the M3 and V09-2 RREs, both the G to A (V09-2) and G to C (M3) changes at the mutation 3 position induce a five stem-loop structure associated with increased activity. The additional mutations were not individually associated with large-scale conformational changes in SHAPE-MaP modeling.

The combination of mutations 1-5, though not each mutation in isolation, also corresponds to increased Shannon entropy in stem-loop II. This is the region where the first and second dimers of Rev bind (55). The disordered base pairing in this region, implied by greater entropy, may facilitate Rev-RNA binding at the primary Rev binding site and subsequent multimerization leading to enhanced activity. This is consistent with our previous *in vitro* gel-shift observations (43) that demonstrated the ability of the V20-1 RRE to promote Rev multimerization more efficiently than the V10-2 RRE. In the absence of major conformational changes attributable to mutations 1, 2, 4, and 5, this mechanism may account for the difference in functional activity between the RREs containing unpaired nucleotides at mutation 3 position (e.g. V09-2 and M3) that lack these additional mutations and the V20-1 RRE. It remains unclear how the presence of these mutations in combination results in increased stem-loop II entropy.

This study also provides further support for the hypothesis that the Rev-RRE regulatory axis plays an important role in HIV pathogenesis. An important consideration in the interpretation of these findings is that all five mutations in V20-1 relative to V08-1

are nonsynonymous in the overlapping gp41 coding sequence, raising the possibility

that changes in Env could contribute to the selective pressure. We searched for known

CTL epitopes in the Env ORF overlapping the V08-1 RRE sequence using the Los

Alamos HIV database Epitope Location Finder tool (www.hiv.lanl.gov). CTL epitopes

have been described in the V08-1 regions containing three of the five mutations

(mutations 3, 4, and 5). Thus, selection of these mutations could plausibly reflect

immune evasion, though it is notable that neither mutation 3 nor 5 were consistently

found in all RRE sequences at all time points after they were first observed at visits 9

and 10.

While the selection evidenced by sequence convergence at visit 20 might be due

to pressure on Env in terms of immune evasion, replication efficiency, or other factors, it

is likely that functional differences in Rev-RRE activity also play a significant role in

overall viral fitness. Nucleotide changes at the mutation 3 position highlight this

likelihood. The nonsynonymous mutations from G to A at visit 9 and A to C at visit 19

are both consistent with CTL evasion. From the standpoint of the RRE, however, the G

to A change shifts the RRE structure to a higher activity conformation while A to C is

structurally and functionally neutral. Thus, the A to C mutation preserves high RRE

activity while also modifying the Env amino acid sequence. Convergence of circulating

RRE sequences by visit 20 is particularly striking in light of the very low CD4 count at

this time point, as viral fitness would likely have been more influenced by Rev-RRE

activity than by immune surveillance.

We were unable to assess functional evolution of the Rev protein over time in

this study, due to the sequencing strategy used, which created reads too short to allow

linkage of both of the Rev coding exons and the RRE. However, previous work using

single genome sequencing with samples from this patient (43) demonstrated that a

single Rev amino acid sequence was most prevalent at both visits V10 and V20. This is

the Rev sequence that was used in this study. Its persistence throughout the entire

infection period suggests that it likely occurred in conjunction with most or all of the

RREs tested here.

The generalizability of the findings described here awaits further longitudinal

studies, analyzing both Rev and RRE sequences, in patients with varying courses of

disease. A clear understanding of RRE evolution during natural infection will help to

further our understanding of the role that this regulatory axis may play in key aspects of

the viral life cycle, including transmission and the establishment of latency. Furthermore,

since the Rev-RRE interaction is an essential step in viral replication, it is also a

promising target for drug development. This study demonstrates that RRE structures

can vary over the course of natural infection, and thus that rational drug design must

account for the spectrum of potential Rev-RRE interaction conformations.

.

MATERIALS AND METHODS

Clinical samples and ethics statement.

All studies involving human subjects were approved by the institutional review

board at Montefiore Hospital, Bronx, NY, as part of the Women's Interagency HIV Study

(WIHS). Written informed consent was provided by all study subjects. Plasma samples

from one patient (SC3), collected over a period of seven years, were obtained from the

WIHS Consortium, Bronx, NY (Kathryn Anastos, principal investigator), as previously

described (43). A total of 11 samples representing different time points were received

for analysis.

Viral RNA extraction and cDNA synthesis.

Viral RNA was extracted from 1 ml of plasma using a guanidinium extraction

protocol (56). Viral cDNA synthesis was performed immediately after viral RNA was

extracted as described (43). Briefly, each viral RNA pellet was resuspended in 40 µl of 5

mM Tris-HCL, pH 8.0, prior to cDNA synthesis with 50µM of oligo(dT) and 100 U of

Superscript III per reaction (Invitrogen, Carlsbad, CA). The reverse transcription

reaction was incubated at 50°C for 1 hour, then 55°C for 1 hour and then 70°C for 15

minutes, after which 4 units of RNase H (Invitrogen) were added, and samples

incubated at 37°C for 20 minutes. Samples were stored at -80°C prior to amplicon

generation.

PCR amplification and purification.

Amplicons (~3 kb) spanning both HIV Rev exons and the RRE, were produced

for each time point. The first round PCR reaction was performed by adding 1 µl of

cDNA template to a 20 µl reaction containing 0.005 U of Platinum Taq Hi-Fidelity

polymerase (Invitrogen) as previously described (57). Using 1 µl of the first round PCR

product as a template, nested PCR resulted in amplicons ~3 kb in length. Primers for

the first round PCR reaction were 2302 (5'-aagccacctttgcctagtg-3') and 2278 (5'-

ttgctacttgtgattgctccatgt-3') while those for the nested PCR were 2277 (5'-

tagagccctggaagcatccaggaag-3') and 2280 (5'-gtctcgagatactgctcccaccc-3') (43).

Amplicon size was verified by agarose gel electrophoresis.

Each 20 µl PCR reaction was purified with 36 µl of AMPure XP beads

(Agencourt). After addition of beads, samples were pipetted up and down 10 times and

incubated at room temperature for 10 minutes prior to being placed on a magnetic

stand. After a 2 minute incubation, supernatant was removed and the beads were

washed twice with 80% ethanol. After beads were air-dried on a magnetic stand for 10

minutes, they were removed from the magnet, resuspended in 52.5 µl of elution buffer,

and incubated at room temperature for 2 minutes. Beads were then placed on the

magnetic stand for 2 minutes, or until the supernatant was cleared. The cleared

supernatant containing the purified PCR amplicons was removed and placed in a new

96-well plate then stored at -20°C.

DNA library preparation and sequencing.

Prior to preparing the DNA library, purified amplicon products were quantified

with the dsDNA HS (high sensitivity) assay kit (Invitrogen) using the Qubit 2.0

fluorometer (Invitrogen). The DNA library was prepared using the standard Nextera XT

DNA library preparation protocol (Illumina). A total of 1 ng of input DNA for each sample

was added to the reaction buffer which fragments the input DNA and adds adapter

sequences to the ends to allow for PCR reactions downstream in the library preparation

process. A brief PCR cycle followed the fragmentation protocol to add indexes used to

identify each individual sample and additional sequences for proper cluster formation.

PCR samples were purified and size-selected for 300-500 bp amplicons using 25 µl of

AMPure XP beads. Amplicons were pooled together in equimolar concentrations then

sequenced with the Illumina MiSeq Reagent Kit v3 (600 cycles).

RRE sequence analysis.

Paired end reads for each time point were generated using an Illumina MiSeq.

The two overlapping paired reads were merged using FLASH (58), a plugin in Geneious

(Biomatters, Auckland, New Zealand) to produce a single read for each pair. These

reads were then aligned to the NL4-3 genome and reads overlapping the RRE were

extracted. Extracted reads were filtered for individual reads that spanned the full length

of the short RRE. These reads were assembled into contigs using the Geneious *de*

novo assembly tool, set for a minimum overlap identity of 100% and a minimum overlap

of the entire 234-nt minimal RRE. The frequency of each contig was calculated based

on the number of reads assigned to each contig. Contigs comprising less than 5% of the

total reads were then removed from each time point (to eliminate minor HIV species

and/or potential PCR errors) and the frequencies of the remaining contigs were

recalculated based on the total number of reads remaining. The sequence of each

contig was then aligned to the sequence from the major V08 contig to generate the

alignment shown in Figure 2 (see also Figures S1 and S2). RREs were named using

the format VXX-Y where XX corresponds to the WIHS cohort visit number at which the

plasma sample generating the sequence was obtained and Y corresponds to the

frequency rank order of the sequence. Thus, V10-2 refers to the second most prevalent

RRE from the sequencing of the plasma sample obtained at WIHS visit 10.

***In vitro* transcription of RRE RNAs.**

RRE RNAs were *in vitro* transcribed by T7 RNA polymerase using the

MEGAshort-script kit (Life Technologies) per manufacturer's guidelines. Templates for

in vitro transcription were generated by PCR amplification of RREs from plasmids

containing synthetic sequences (Integrated DNA technologies) using oligonucleotides

designed to introduce both a T7 promoter sequence at the 5' end of the 234-nt and 351-

nt RREs and a structure cassette at the 3' end of the 351-nt RREs (Table S1). Plasmids

used for generating transcription DNA template for the 234-nt V08-1, V09-2, V10-2,

V20-1, M1, M2, M3, and M4 RREs were designated pHR 5334, pHR 5766, pHR 4784,

pHR 4788, pHR 5326, pHR 5328, pHR 5330, and pHR 5332, respectively. Those used

to generate 351-nt V10-2 and V20-1 RREs were designated pHR 5321 and pHR 5322,

respectively. *In vitro* transcription reactions were treated with Turbo DNase I (Life

Technologies) for 1 hr at 37°C to digest the DNA template, heated to 85°C for 2 min,

and RNA was fractionated on a denaturing gel (5% polyacrylamide (19:1), 1x TBE, 7M

urea) at constant temperature (45°C, 30W max). RRE bands were located by UV

shadowing, excised, electroeluted at 200 V for 2 hours at 4°C, ethanol precipitated, and

stored at 4°C in TE buffer (10 mM Tris, pH 7.6; 0.1 mM EDTA).

Folding and SHAPE-MaP profiling of RRE RNA.

For each RNA, 1M7 (+), 1M7 (-), and denaturation control reactions were

generated. Approximately 5 pmoles of RRE RNA was incubated with renaturation buffer

(1X: 10 mM Tris, pH 8.0, 100 mM KCl, 0.1 mM EDTA) in a total volume of 5 µl, heated

to 85°C, and renatured by slow cooling (0.1°C/sec) to 25°C. Renatured RNA was

incubated with RNA folding buffer (1X: 35 mM Tris pH 8.0, 90 mM KCl, 0.15 mM EDTA,

5 mM MgCl2, 2.5% glycerol) in a total volume of 9 μl at 37°C for 30 min. RNA was

modified by addition of 1 μL of 25 mM 1M7. The negative control reaction contained 1

μL of DMSO instead of 1M7. The denaturation control was produced by incubating 5

pmoles of RNA with denaturation buffer (1X: 50% formamide, 50 mM HEPES, and 4

mM EDTA) in a volume of 9 μL at 95 °C for 1 min. 1 μL of 20 mM 1M7 was added and

the mixture incubated at 95 °C for another 1 min. RNA from 1M7 (+), 1M7 (-), and

denatured control reactions were recovered by ethanol precipitation, resuspended in TE

buffer (10 mM Tris, pH 7.6; 0.1 mM EDTA), and stored at -20°C.

Mutational profiling of modified RNA.

RNA from 1M7 (+), 1M7 (-), and denatured control reactions were reverse

transcribed using corresponding oligos to generate a cDNA library (Table S1). Reverse

transcription was performed by first annealing 2 μM of RT oligo to the RNA in a reaction

volume of 11 μL by incubation at 65 °C for 5 min, followed by cooling on ice. cDNA

synthesis was initiated by incubating the annealing mixture with 8 μL of 2.5X RT

reaction mixture (2.5X: 125 mM Tris (pH 8.0), 187.5 mM KCl, 15 mM MnCl$_2$, 25 mM

DTT and 1.25 mM dNTPs) and 1 μL of Superscript II RT (Thermo Fisher, 200 U/μL) for

42 °C for 3 hours. The RNA template was then hydrolyzed by adding 1 μL 2 N NaOH,

followed by neutralization by addition of 1 μL 2N HCl. The cDNA library was purified

using G50 spin columns (GE Healthcare).

For mutational profiling, cDNA was converted to dsDNA with Illumina adapters

for high throughput sequencing on an Illumina platform. This was achieved in two

consecutive PCR reactions, namely PCR1 and PCR2. The first reaction (PCR1)

appended partial Illumina adapters to the ends of the amplicons. The entire cDNA

library was used as template in a 100 μl PCR1 reaction (1.1 μL each of 50 pmoles of

forward and reverse oligo, 2 μL of 10 mM dNTPs, 20 μL of 5x Q5 reaction buffer, 1 μL

of Hot Start High-Fidelity DNA polymerase (NEB)) using cycling conditions: 98°C for 30

sec, 15 cycles of (98 °C for 10 sec, 50 °C for 30 sec, 72 °C for 30 sec), 72 °C for 2 min.

PCR product was gel purified and 10% was used as template DNA in the PCR2

reaction. The PCR2 completed the Illumina adapters while adding appropriate

barcoding indices. The PCR2 reaction and cycling conditions were as those for PCR1.

The resulting amplicon library was fractionated through a 2% agarose gel and

amplicons purified from the gel slices by electro-elution at room temperature for 2 hr

followed by ethanol precipitation. Each amplicon library was quantified by real time PCR

using the KAPA Universal Library Quantification Kit (Illumina) per manufacturer's

protocol. Sequencing libraries were pooled and mixed with 20% PhiX and sequenced

on an Illumina MiSeq to generate 2 X 150 paired-end reads. Sequence files were fed

into ShapeMapper (v1.2) software (49) to generate SHAPE reactivity profiles by aligning

the reads to 234-nt RRE RNA sequence using the software with default settings.

Reactivity values obtained from ShapeMapper were input to RNAstructure software (51)

to generate the minimum free-energy RNA secondary structure model.

RRE gel migration assay.

RRE structural homogeneity was assessed by comparing the migration rate of the

folded RREs on native agarose and polyacrylamide gels. Approximately 20 pmoles of

RNA was suspended in 5 µl renaturation buffer (1X: 10mM Tris, pH 8.0, 100mM KCl,

0.1mM EDTA), heated to 85°C, and renatured by slow cooling (0.1°C/sec) to 25°C.

Renatured RNA was incubated with 5 µl of 2X RNA folding buffer (1X:70 mM Tris pH

8.0, 180 mM KCl, 0.3 mM EDTA, 10 mM $MgCl_2$, 5% of glycerol) at 37°C for 30 min, and

fractionated through a native 8% polyacrylamide gel (29:1) run for 22 hours at 4°C or on

a 2% native agarose. RRE bands were visualized by UV shadowing and ethidium

bromide staining, respectively.

Plasmid constructs for functional assays of Rev-RRE activity.

Functional activity of Rev-RRE pairs was determined by means of a

fluorescence-based transient transfection assay. To test RRE sequences, a plasmid

containing a near full length NL4-3 HIV-1 sequence (59) was modified to express blue

fluorescent protein (BFP) in a Rev-independent fashion and green fluorescent protein

(GFP) in a Rev-dependent fashion. The native 351-nt RRE within *env* was flanked by

XmaI and XbaI sites to permit exchange of this sequence with other RREs of interest.

Additional modifications were made to ensure the construct was not replication-

competent and could not express Rev. To create additional RRE constructs, the native

RRE was removed by digesting the plasmid with XmaI and XbaI. Commercially

synthesized 234-nt RRE sequences of interest with appropriate flanking sequences

(Integrated DNA Technologies) were then cloned into the opened *env* region using

Gibson assembly.

Rev was provided *in* trans using one of two constructs. The predominant Rev

previously identified at V10 and V20 by single genome sequencing had identical amino

acid sequences. This Rev was designated M0-B/M57-A and used for functional assays

except as otherwise noted (43).

For the analysis described in Figure 4, the M0-B/M57-A *rev* sequence was

cloned into a CMV expression plasmid. For the remaining functional assays, a separate

Rev-expressing construct was created by modifying the murine stem cell virus

construct, pMSCV-IRES-mCherry FP (Addgene plasmid # 52114), to express both the

M-0B/M57-A Rev and the mCherry fluorescent protein by means of a bicistronic

transcript including an internal ribosome entry site (60). The MSCV vector was a gift

from Dario Vignali (unpublished). Constructs utilized in the functional assays are listed

in Table S2.

Selection of RREs for functional analysis.

Both naturally occurring RREs sequenced from patient SC3 and synthetic RREs

containing individual mutations of interest were tested for functional activity. All RREs

tested were 234-nt in length. The naturally occurring RREs assessed were V08-1, V09-

2, V10-2, V11-1, V13-1, V14-2, V19-2, and V20-1. Additionally, synthetic RREs were

created based on single nucleotide differences between the sequences of V10-2 and

V20-1. The mutation 1 RRE (M1) is identical to V10-2 with the exception of a G to A

change at position 61, reflecting the G to A change seen at that position in V20-1. The

mutation 2 RRE (M2) is identical to V10-2 with the exception of a T to A change at

position 84. Of note, this RRE sequence was also observed to occur naturally as

sequence V12-1. The mutation 3 RRE (M3) is identical to V10-2 with the exception of a

G to C change at position 194. The mutation 4 RRE (M4) is identical to V10-2 with the

exception of an A to G change at position 228.

Rev-RRE functional activity assays.

RRE-containing constructs were designed such that GFP expression occurs in a

Rev-dependent fashion while BFP expression occurs in a Rev-independent fashion. In

cells transfected with both the RRE-containing and Rev-containing constructs, the

degree of GFP expression relative to BFP expression is proportional to the functional

activity of the tested Rev-RRE pair (Jackson et al.. manuscript in preparation).

Except as noted, transfections were performed using the polyethylenamine

method and $8x10^5$ 293T/17 cells in each well of a 12-well plate. Cells were maintained

in 1 mL IMDM supplemented with 5% bovine calf serum. Each RRE-containing

construct was tested individually with the SC3 Rev-containing construct. Additionally,

each RRE-containing construct was transfected into cells without the addition of Rev to

ensure that GFP expression in this system was truly Rev-dependent. In transfections

performed without the Rev-containing construct, an empty CMV construct was used to

compensate such that each transfection was performed with a constant mass of

plasmid. For each transfection, 1000 ng of an RRE-containing construct and 100 ng of

either the MSCV-Rev construct or empty vector was used. Simultaneously, different

cultures of 293T/17 cells were transfected with constructs expressing GFP, BFP, or

mCherry alone as single-color controls to permit color compensation during flow

cytometry. Each set of transfections was performed in duplicate with the exception of

V09-2 and V13-1, for which four replicates were performed. After transfection, cells

were incubated for 48 h, then suspended in phosphate buffered saline. Flow cytometry

was performed on the resulting suspension using an Attune NxT flow cytometer with

autosampler attachment (Thermo Fischer Scientific). Data acquisition was performed on

the Attune NxT software package using the following channels:

Laser line	488	405	561
Emission filter	530/30	440/50	620/15
Fluorescent protein	eGFP	TagBFP	mCherry

Post-acquisition color compensation and data analysis was performed using

FlowJo v10 (FlowJo, LLC). For each analysis, gating was performed on single 293T/17

cells. Next, a daughter population of cells that expressed the RRE-containing transcript

was identified by gating on positivity for GFP or BFP. Using this population, the ratio of

arithmetic mean fluorescent intensity (MFI) of eGFP to TagBFP for successfully

transfected cells was calculated in both the presence and the absence of Rev. A low

ratio in the absence of Rev and a high ratio in its presence indicated that GFP

expression was Rev-dependent.

After determining that the RRE-containing constructs expressed GFP in a Rev-

dependent fashion, an additional analysis was performed to determine the functional

activity of specific Rev-RRE pairs. In this further analysis, only those transfections

including both an RRE-containing construct and the Rev-containing construct were

considered. An additional daughter population was created from that described above

by gating on mCherry positivity. This final population consisted only of single 293T/17

cells expressing GFP and/or BFP and also mCherry. This pattern of expression ensured

that selected cells were successfully co-transfected with both plasmid constructs. Once

this population was defined, the ratio of GFP to BFP MFI was re-calculated. This final

GFP:BFP ratio was used to determine the relative functional activity of different RREs

assessed in the presence of SC3 Rev. Differences in functional activity values between

RREs were evaluated using SPSS Statistics v25 (IBM). P-values were calculated using

a two-tailed one-way ANOVA with Tukey's HSD test for Figures 4 and 7 and with a one-

tailed one-way ANOVA with Dunnett's test for Figure 6E.

Analysis of the V08-1, V10-2, and V20-1 RREs described in Figure 4 was

performed with minor modifications. For this analysis, Rev was provided *in trans* by the

CMV-SC3 Rev construct. Each well of a 12-well plate was seeded with 4×10^5 293T/17

cells maintained in 2 mL IMDM supplemented with 10% bovine calf serum. One day

after plating, medium was removed from each well and replaced with 1 mL IMDM

supplemented with 5% bovine calf serum. Cells in each well were then transfected with

1000 ng of an RRE-containing construct and 50 ng of the CMV-Rev construct using the

polyethylenamine method. Four transfection replicates were performed. Simultaneously,

different cultures of 293T/17 cells were transfected with constructs expressing GFP or

BFP alone as single-color controls to permit color compensation during flow cytometry.

Cells were harvested and flow cytometry was performed 24 hours after transfection.

Gating was performed as above identifying cells successfully transduced with the RRE-

containing construct and expressing GFP or BFP, and the ratio of GFP:BFP was

calculated. Additional gating on mCherry positive cells was not performed.

CE-SHAPE.

RNA was folded as in the gel migration assay except that the final volume of the

folded RNA mix was 150 ul and glycerol was excluded from the folding buffer. Folded

RNAs were probed using 3 mM NMIA. For this, RNAs were divided into experimental

(NMIA+) and control (NMIA-) aliquots (72 µl each), to which 8 µl 30 mM NMIA in

anhydrous DMSO or DMSO alone was added, respectively. Modification reactions were

incubated at 37°C for 50 min, ethanol precipitated, and re-suspended in 13 ul nuclease-

free water. Reverse transcription of modified RNAs, cDNA processing/fractionation, and

SHAPE data analysis were conducted as previously described (61).

Creating probability pairing and Shannon entropy profiles.

Pairing probability (Figure S5-S8) and Shannon entropy values were calculated

by feeding SuperFold (49) with the 1M7 reactivities of the RREs. Shannon entropy

measurements were calculated over a centered 11-nt sliding window and plotted

against nucleotide position.

Nucleotide sequence accession numbers.

Nucleotide sequences generated through the procedure above corresponding to

RREs from viral quasispecies found in samples from patient SC3 were deposited in

Genbank under accession numbers MK190736 through MK190867. The SC3 Rev

sequence used in the functional assays was previously deposited under accession

number KF559146. The RRE sequences listed here as V10-2 and V20-1 were

previously deposited under accession numbers KF559160 and KF559162, respectively.

ACKNOWLEDGMENTS

This work was supported by grants GM 110009, AI134208, AI087505 and

AI068501 as well as the Women's Interagency HIV Study (WIHS) grant UO1-AI-35004

from the National Institutes of Health (NIH). The patient samples used in this manuscript

were collected by the Women's Interagency HIV Study (WIHS). The Bronx WHIS

repository provided the initial patient materials and the WIHS central repository provided

the materials from the intermediate time points (Bronx WIHS [Kathryn Anastos and

Anjali Sharma], U01-AI-035004). Jason W Rausch (NCI) is acknowledged for his

contribution in setting up the SHAPE-MaP analysis pipeline. S.L.G. and C.S. were

supported by the Intramural Research Program of the National Cancer Institute,

National Institutes of Health, Department of Health and Human Services. P. E. H. J. was

supported by grant K08AI136671 from the National Institutes of Health. Salary support

for M.-L.H. and D.R. was provided by the Charles H. Ross, Jr., and Myles H. Thaler

Endowments at the University of Virginia. The contents of this publication are solely the

responsibility of the authors and do not represent the official views of NIH or the WIHS.

The funders had no role in study design, data collection and interpretation, or the

decision to submit the work for publication.

REFERENCES

1.   Rekosh D, Hammarskjold ML. 2018. Intron retention in viruses and cellular

genes: Detention, border controls and passports. Wiley Interdiscip Rev RNA

9:e1470.

2.   Leblanc J, Weil J, Beemon K. 2013. Posttranscriptional regulation of retroviral

gene expression: primary RNA transcripts play three roles as pre-mRNA, mRNA,

and genomic RNA. Wiley Interdiscip Rev RNA 4:567-80.

3.   Fernandes JD, Booth DS, Frankel AD. 2016. A structurally plastic

ribonuceloprotein complex mediates post-transcriptional gene regulation in HIV-

1. Wiley Interdiscip Rev RNA doi:10.1002/wrna.1342.

4.   Hammarskjold ML, Heimer J, Hammarskjold B, Sangwan I, Albert L, Rekosh D.

1989. Regulation of human immunodeficiency virus env expression by the rev

gene product. J Virol 63:1959-66.

5.   Hadzopoulou-Cladaras M, Felber BK, Cladaras C, Athanassopoulos A, Tse A,

Pavlakis GN. 1989. The rev (trs/art) protein of human immunodeficiency virus

type 1 affects viral mRNA and protein expression via a cis-acting sequence in the

env region. J Virol 63:1265-74.

6.   Malim MH, Hauber J, Le SY, Maizel JV, Cullen BR. 1989. The HIV-1 rev trans-

activator acts through a structured target sequence to activate nuclear export of

unspliced viral mRNA. Nature 338:254-7.

7.   Berger J, Aepinus C, Dobrovnik M, Fleckenstein B, Hauber J, Bohnlein E. 1991.

Mutational analysis of functional domains in the HIV-1 Rev trans-regulatory

protein. Virology 183:630-5.

8.   Cochrane AW, Perkins A, Rosen CA. 1990. Identification of sequences important

in the nucleolar localization of human immunodeficiency virus Rev: relevance of

nucleolar localization to function. J Virol 64:881-5.

9.   Kubota S, Nosaka T, Cullen BR, Maki M, Hatanaka M. 1991. Effects of chimeric

mutants of human immunodeficiency virus type 1 Rev and human T-cell

leukemia virus type I Rex on nucleolar targeting signals. J Virol 65:2452-6.

10. Dundr M, Leno GH, Hammarskjold ML, Rekosh D, Helga-Maria C, Olson MO.

1995. The roles of nucleolar structure and function in the subcellular location of

the HIV-1 Rev protein. J Cell Sci 108 (Pt 8):2811-23.

11. Kjems J, Frankel AD, Sharp PA. 1991. Specific regulation of mRNA splicing in

vitro by a peptide from HIV-1 Rev. Cell 67:169-78.

12. Battiste JL, Mao H, Rao NS, Tan R, Muhandiram DR, Kay LE, Frankel AD,

Williamson JR. 1996. Alpha helix-RNA major groove recognition in an HIV-1 rev

peptide-RRE RNA complex. Science 273:1547-51.

13. Daugherty MD, Liu B, Frankel AD. 2010. Structural basis for cooperative RNA

binding and export complex assembly by HIV Rev. Nat Struct Mol Biol 17:1337-

42.

14. Thomas SL, Oft M, Jaksche H, Casari G, Heger P, Dobrovnik M, Bevec D,

Hauber J. 1998. Functional analysis of the human immunodeficiency virus type 1

Rev protein oligomerization interface. J Virol 72:2935-44.

15. Booth DS, Cheng Y, Frankel AD. 2014. The export receptor Crm1 forms a dimer

to promote nuclear export of HIV RNA. Elife 3.

16. Wen W, Meinkoth JL, Tsien RY, Taylor SS. 1995. Identification of a signal for

rapid export of proteins from the nucleus. Cell 82:463-473.

17.   Fischer U, Huber J, Boelens WC, Mattaj IW, Luhrmann R. 1995. The HIV-1 Rev

863   activation domain is a nuclear export signal that accesses an export pathway

864   used by specific cellular RNAs. Cell 82:475-483.

865   18.   Mann DA, Mikaelian I, Zemmel RW, Green SM, Lowe AD, Kimura T, Singh M,

866   Butler PJ, Gait MJ, Karn J. 1994. A molecular rheostat. Co-operative rev binding

867   to stem I of the rev-response element modulates human immunodeficiency virus

868   type-1 late gene expression. J Mol Biol 241:193-207.

869   19.   Dayton ET, Powell DM, Dayton AI. 1989. Functional analysis of CAR, the target

870   sequence for the Rev protein of HIV-1. Science 246:1625-9.

871   20.   Kjems J, Brown M, Chang DD, Sharp PA. 1991. Structural analysis of the

872   interaction between the human immunodeficiency virus Rev protein and the Rev

873   response element. Proc Natl Acad Sci U S A 88:683-7.

874   21.   Sherpa C, Rausch JW, Le Grice SF, Hammarskjold ML, Rekosh D. 2015. The

875   HIV-1 Rev response element (RRE) adopts alternative conformations that

876   promote different rates of virus replication. Nucleic Acids Res 43:4676-86.

877   22.   Bai Y, Tambe A, Zhou K, Doudna JA. 2014. RNA-guided assembly of Rev-RRE

878   nuclear export complexes. Elife 3:e03656.

879   23.   Watts JM, Dang KK, Gorelick RJ, Leonard CW, Bess JW, Jr., Swanstrom R,

880   Burch CL, Weeks KM. 2009. Architecture and secondary structure of an entire

881   HIV-1 RNA genome. Nature 460:711-6.

882   24.   Cook KS, Fisk GJ, Hauber J, Usman N, Daly TJ, Rusche JR. 1991.

883   Characterization of HIV-1 REV protein: binding stoichiometry and minimal RNA

884   substrate. Nucleic Acids Res 19:1577-83.

25.   Daly TJ, Rennert P, Lynch P, Barry JK, Dundas M, Rusche JR, Doten RC, Auer

M, Farrington GK. 1993. Pertubation of the carboxy terminus of HIV-1 rev affects

multimerization on the rev responsive element. Biochemistry 32:8945-8954.

26.   Malim MH, McCarn DF, Tiley LS, Cullen BR. 1991. Mutational definition of the

human immunodeficiency virus type 1 Rev activation domain. J Virol 65:4248-54.

27.   Heaphy S, Finch JT, Gait MJ, Karn J, Singh M. 1991. Human immunodeficiency

virus type 1 regulator of virion expression, rev, forms nucleoprotein filaments

after binding to a purine-rich "bubble" located within the rev-responsive region of

viral mRNAs. Proc Natl Acad Sci U S A 88:7366-70.

28.   Zapp ML, Hope TJ, Parslow TG, Green MR. 1991. Oligomerization and RNA

binding domains of the type 1 human immunodeficiency virus Rev protein: a dual

function for an arginine- rich binding motif. Proc Natl Acad Sci U S A 88:7734-

7748.

29.   Zemmel RW, Kelley AC, Karn J, Butler PJ. 1996. Flexible regions of RNA

structure facilitate co-operative Rev assembly on the Rev-response element. J

Mol Biol 258:763-77.

30.   Daugherty MD, D'Orso I, Frankel AD. 2008. A solution to limited genomic

capacity: using adaptable binding surfaces to assemble the functional HIV Rev

oligomer on RNA. Mol Cell 31:824-34.

31.   DiMattia MA, Watts NR, Stahl SJ, Rader C, Wingfield PT, Stuart DI, Steven AC,

Grimes JM. 2010. Implications of the HIV-1 Rev dimer structure at 3.2 A

resolution for multimeric binding to the Rev response element. Proc Natl Acad

Sci U S A 107:5810-4.

32.   Jain C, Belasco JG. 2001. Structural model for the cooperative assembly of HIV-
1 Rev multimers on the RRE as deduced from analysis of assembly-defective
mutants. Mol Cell 7:603-14.

33.   Fornerod M, Ohno M, Yoshida M, Mattaj IW. 1997. CRM1 is an export receptor
for leucine-rich nuclear export signals. Cell 90:1051-60.

34.   Otero GC, Harris ME, Donello JE, Hope TJ. 1998. Leptomycin B inhibits equine
infectious anemia virus Rev and feline immunodeficiency virus rev function but
not the function of the hepatitis B virus posttranscriptional regulatory element. J
Virol 72:7593-7.

35.   Jackson PE, Tebit DM, Rekosh D, Hammarskjold ML. 2016. Rev-RRE Functional
Activity Differs Substantially Among Primary HIV-1 Isolates. AIDS Res Hum
Retroviruses 32:923-34.

36.   Phuphuakrat A, Paris RM, Nittayaphan S, Louisirirotchanakul S, Auewarakul P.
2005. Functional variation of HIV-1 Rev Response Element in a longitudinally
studied cohort. J Med Virol 75:367-73.

37.   Iversen AK, Shpaer EG, Rodrigo AG, Hirsch MS, Walker BD, Sheppard HW,
Merigan TC, Mullins JI. 1995. Persistence of attenuated rev genes in a human
immunodeficiency virus type 1-infected asymptomatic individual. J Virol 69:5743-
53.

38.   Churchill MJ, Chiavaroli L, Wesselingh SL, Gorry PR. 2007. Persistence of
attenuated HIV-1 rev alleles in an epidemiologically linked cohort of long-term
survivors infected with nef-deleted virus. Retrovirology 4:43.

39. Bobbitt KR, Addo MM, Altfeld M, Filzen T, Onafuwa AA, Walker BD, Collins KL.

2003. Rev activity determines sensitivity of HIV-1-infected primary T cells to CTL

killing. Immunity 18:289-99.

40. Baccam P, Thompson RJ, Li Y, Sparks WO, Belshan M, Dorman KS,

Wannemuehler Y, Oaks JL, Cornette JL, Carpenter S. 2003. Subpopulations of

equine infectious anemia virus Rev coexist in vivo and differ in phenotype. J Virol

77:12122-12131.

41. Belshan M, Baccam P, Oaks JL, Sponseller BA, Murphy SC, Cornette J,

Carpenter S. 2001. Genetic and biological variation in equine infectious anemia

virus Rev correlates with variable stages of clinical disease in an experimentally

infected pony. Virology 279:185-200.

42. Belshan M, Harris ME, Shoemaker AE, Hope TJ, Carpenter S. 1998. Biological

characterization of Rev variation in equine infectious anemia virus. Journal of

virology 72:4421-6.

43. Sloan EA, Kearney MF, Gray LR, Anastos K, Daar ES, Margolick J, Maldarelli F,

Hammarskjold ML, Rekosh D. 2013. Limited nucleotide changes in the Rev

response element (RRE) during HIV-1 infection alter overall Rev-RRE activity

and Rev multimerization. J Virol 87:11173-86.

44. Smola MJ, Christy TW, Inoue K, Nicholson CO, Friedersdorf M, Keene JD, Lee

DM, Calabrese JM, Weeks KM. 2016. SHAPE reveals transcript-wide

interactions, complex structural domains, and protein interactions across the Xist

lncRNA in living cells. Proc Natl Acad Sci U S A 113:10322-7.

45. Sztuba-Solinska J, Rausch JW, Smith R, Miller JT, Whitby D, Le Grice SFJ.

2017. Kaposi's sarcoma-associated herpesvirus polyadenylated nuclear RNA: a

structural scaffold for nuclear, cytoplasmic and viral proteins. Nucleic Acids Res

45:6805-6821.

46. Lu YF, Mauger DM, Goldstein DB, Urban TJ, Weeks KM, Bradrick SS. 2015.

IFNL3 mRNA structure is remodeled by a functional non-coding polymorphism

associated with hepatitis C virus clearance. Sci Rep 5:16037.

47. Siegfried NA, Busan S, Rice GM, Nelson JA, Weeks KM. 2014. RNA motif

discovery by SHAPE and mutational profiling (SHAPE-MaP). Nat Methods

11:959-65.

48. Smola MJ, Calabrese JM, Weeks KM. 2015. Detection of RNA-Protein

Interactions in Living Cells with SHAPE. Biochemistry 54:6867-75.

49. Smola MJ, Rice GM, Busan S, Siegfried NA, Weeks KM. 2015. Selective 2'-

hydroxyl acylation analyzed by primer extension and mutational profiling

(SHAPE-MaP) for direct, versatile and accurate RNA structure analysis. Nat

Protoc 10:1643-69.

50. Sherpa C, Rausch JW, Le Grice SFJ. 2018. Structural characterization of

maternally expressed gene 3 RNA reveals conserved motifs and potential sites

of interaction with polycomb repressive complex 2. Nucleic Acids Res

doi:10.1093/nar/gky722.

51. Bellaousov S, Reuter JS, Seetin MG, Mathews DH. 2013. RNAstructure: Web

servers for RNA secondary structure prediction and analysis. Nucleic Acids Res

41:W471-4.

52.   Srinivasakumar N, Chazal N, Helga-Maria C, Prasad S, Hammarskjold ML,

976   Rekosh D. 1997. The effect of viral regulatory protein expression on gene

977   delivery by human immunodeficiency virus type 1 vectors produced in stable

978   packaging cell lines. J Virol 71:5841-8.

979   53.   Deigan KE, Li TW, Mathews DH, Weeks KM. 2009. Accurate SHAPE-directed

980   RNA structure determination. Proc Natl Acad Sci U S A 106:97-102.

981   54.   McGinnis JL, Dunkle JA, Cate JH, Weeks KM. 2012. The mechanisms of RNA

982   SHAPE chemistry. J Am Chem Soc 134:6617-24.

983   55.   Tiley LS, Malim MH, Tewary HK, Stockley PG, Cullen BRJPotNAoS. 1992.

984   Identification of a high-affinity RNA-binding site for the human immunodeficiency

985   virus type 1 Rev protein. 89:758-762.

986   56.   Palmer S, Wiegand AP, Maldarelli F, Bazmi H, Mican JM, Polis M, Dewar RL,

987   Planta A, Liu S, Metcalf JA, Mellors JW, Coffin JM. 2003. New real-time reverse

988   transcriptase-initiated PCR assay with single-copy sensitivity for human

989   immunodeficiency virus type 1 RNA in plasma. J Clin Microbiol 41:4531-6.

990   57.   Kearney M, Maldarelli F, Shao W, Margolick JB, Daar ES, Mellors JW, Rao V,

991   Coffin JM, Palmer S. 2009. Human immunodeficiency virus type 1 population

992   genetics and adaptation in newly infected individuals. J Virol 83:2715-27.

993   58.   Magoc T, Salzberg SL. 2011. FLASH: fast length adjustment of short reads to

994   improve genome assemblies. Bioinformatics 27:2957-63.

995   59.   Adachi A, Gendelman HE, Koenig S, Folks T, Willey R, Rabson A, Martin MA.

996   1986. Production of acquired immunodeficiency syndrome-associated retrovirus

in human and nonhuman cells transfected with an infectious molecular clone. J

Virol 59:284-91.

60.   Shaner NC, Campbell RE, Steinbach PA, Giepmans BN, Palmer AE, Tsien RY.

2004. Improved monomeric red, orange and yellow fluorescent proteins derived

from Discosoma sp. red fluorescent protein. Nat Biotechnol 22:1567-72.

61.   Lusvarghi S, Sztuba-Solinska J, Purzycka KJ, Pauly GT, Rausch JW, Grice SF.

2013. The HIV-2 Rev-response element: determining secondary structure and

defining folding intermediates. Nucleic Acids Res 41:6637-49.

1005

FIGURE LEGENDS:

Figure 1: Patient SC3 HIV viral load and CD4 count at each visit during participation in

the WIHS cohort. Sequential study visits occurred about six months apart and blood

samples were obtained at each visit. V01 refers to the first visit, V04 to the fourth, etc.

Data is missing for unattended visits. HIV sequences could first be amplified from

plasma samples obtained at V08 and onwards. However, HIV seroconversion was first

appreciated under the WIHS protocol at V10 and viral loads were only performed from

this visit on. The patient died after V20.

1014

1015 Figure 2: Evolutionary alignment of SC3 RREs. Deep sequencing of viral RNA was

1016 performed from plasma collected at each study visit. This alignment includes RRE

1017 contigs that were present in at least 5% of the total sequences as explained in the

1018 methods. Sequence labels are in the form of VXX-Y where XX refers to visit number

1019 and Y refers to rank order of the contig within that visit plasma sample. Nucleotide

changes relative to the presumptive founder sequence, V08-1, are highlighted. Mutations (Mut) 1-5 refer to single nucleotide changes that occur in V20-1 relative to V08-1. Mut 5 is highlighted in red, as this change is also found in V10-2 and was not recognized in our previous study of the "early" and "late" RREs from patient SC3 (43). (See also Figures S1 for the alignment including minority variants and S2 for the complete nucleotide sequences included here.)

Figure 3: Structures of selected SC3 RREs. Secondary structures of [A] V08-1, [B] V10-2, [C] V09-2, and [D] V20-1 234-nt RREs were determined by SHAPE-MaP. 1M7 reactivities of RRE nucleotides are color coded and superimposed on the structures. The positions of the mut 1, mut 2, mut 3, and mut 4 single nucleotide changes are represented by black arrows. Green labelled arrows show where the nucleotide at this position varies from V08-1, the presumptive founder sequence, and V10-2. In V09-2, the mut 3 position shows a G to A change, while the mut 3 position in V20-1 shows a G to C change. The mut 5 position is shown with a red arrow and is additionally labelled with "mut 5" in structures where the nucleotide varies from V08-1. [E] Migration of V10-2 and V20-1 RRE on a native PAGE gel was visualized after 22 hours by UV shadowing.

Figure 4: Functional activity of SC3 founder (V08-1), "early" (V10-2), and "late" (V20-1) RREs. Functional activity of the three RREs was determined by transfecting reporter constructs containing the different RREs into 293T/17 cells along with 50 ng of SC3 Rev and measuring the ratio of the mean fluorescent intensity of GFP to BFP. N=4, SEM is

1042   represented by error bars. P-values of <0.05 are represented by *, NS=non-significant

1043   difference.

1044

Figure 5: Secondary structures of full-length (351-nt) V10-2 and V20-1 SC3 RREs.

Secondary structures of [A] V10-2 and [B] V20-1 SC3 RREs, determined by CE-

SHAPE. NMIA nucleotide reactivities are color coded and superimposed on the

structures. The positions of the eight nucleotides differing between the two RREs are

represented by arrows. Mutations within the short 234-nt RRE are designated mut 1,

mut 2, mut 3, and mut 4 and shown in green on the V20-1 RRE. Four additional

mutations within stem SL-I are shown as green arrows without labels on the V20-1

RRE. The position of each of these changes is shown by black arrows on the V10-2

RRE.

1054

Figure 6: Secondary structures and function of V10-2 RRE single mutants. Secondary

structures of the [A] M1, [B] M2, [C] M3, and [D] M4 SC3 RREs were determined by

SHAPE-MaP. 1M7 reactivities are color coded and superimposed on the structures. The

positions of mut 1-5 are represented by black and green arrows. Mut 1, mut 2, mut 3,

and mut 4 are indicated in green on the RREs where the nucleotide at that position

differs from V10-2, while mut 5 is represented in red and is present on every structure.

[E] Rev-RRE functional activity of M1-M4 RREs compared with V10-2 and V20-1 in the

presence of 100 ng SC3 Rev. N=2, SEM are represented by error bars. Selected p-

values of <0.05 and <0.001 are represented by * and ***, respectively.

1064

1065 Figure 7: Rev-RRE activity of SC3 RRE haplotypes. Activity of selected RREs was

1066 determined in the presence of 100 ng Rev using the fluorescent assay system. N=2

1067 except for V9-2 and V13-1 where N=4. SEM are represented by the error bars. Selected

1068 p-values of <0.05 and <0.001 are represented by * and ***, respectively.

1069

Figure 8: Shannon entropy profiles of SC3 RRE variants and single mutants. Shannon

entropy values of the SHAPE-MaP generated RRE structures, smoothed over centered

11-nt sliding windows are plotted as a function of nucleotide position. Higher Shannon

entropy suggests regions that are structurally dynamic. The boxed regions correspond

to the loop between SL-IIA and SL-IIB (Region I) and the loop between SL-IIB and SL-

IIC (Region II). These regions display high Shannon entropy in the high-activity V20-1

structure and low entropy in the lower activity V08-1 structure.

1077

A

SL-III

SL-IIC

SL-IIA

SL-IIB

SL-I

SL-IV

SL-V

High (>0.7)
Medium (0.5-0.7)
Low (0.3-0.5)
None (<0.3)
Unquantifiable

351 nt V10-2 RRE

B

SL-III

SL-IIC

SL-IIA

SL-IIB

SL-I

SL-IV

SL-V

mut 1
mut 2
mut 3
mut 4

351 nt V20-1 RRE