Genomic changes underlying host specialization in the bee gut

symbiont *Lactobacillus* Firm5

Ellegaard KM[1], Brochet S[1], Bonilla-Rosso G[1], Emery O[1], Glover N[2], Hadadi N[1], Jaron

KS[2,4], van der Meer JR[1], Robinson-Rechavi M[2,4], Sentchilo V[1], Tagini F[3], SAGE class

2016-17, Engel P[1]*

[1]Department of Fundamental Microbiology, University of Lausanne, 1015 Lausanne,

Switzerland

[2]Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne,

Switzerland

[3]Institute of Microbiology, Department of Laboratory Medicine, University of

Lausanne & Lausanne University Hospital, Lausanne, Switzerland

[4]Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland

*Correspondence:

Prof. Philipp Engel

Department of Fundamental Microbiology

University of Lausanne, CH-1015 Lausanne, Switzerland

Tel.: +41 (0)21 692 56 12

e-mail: philipp.engel@unil.ch

Abstract

Bacteria that engage in longstanding associations with particular hosts are expected to evolve host-specific adaptations that limit their capacity to strive in other environments. Consistent with this, many gut symbionts seem to have a limited host range, based on community profiling and phylogenomics. However, few studies have experimentally investigated host specialization of gut symbionts and underlying mechanisms have largely remained elusive. Here, we studied host specialization of a dominant gut symbiont of social bees, *Lactobacillus* Firm5. We show that Firm5 strains isolated from honey bees and bumble bees separate into deep-branching phylogenetic lineages. Despite their divergent evolution, colonization experiments show that bumble bee strains are capable of colonizing the honey bee gut. However, they were less successful than honey bee strains, and competition with honey bee strains completely abolished their colonization, whereas honey bee strains were able to coexist. This suggests that both host selection and interbacterial competition play important roles for host specialization. Using comparative genomics of 27 Firm5 isolates, we identified candidate genomic changes underlying host specialization. We found that honey bee strains harbored a larger and more diverse gene pool of carbohydrate-related functions than bumble bee strains. As dietary-derived carbohydrates are the main energy source for strains of the Firm5 phylotype, the metabolic flexibility of honey bee strains may give these bacteria a competitive advantage over bumble bee strains in colonizing the gut niche and hence contribute to host specialization.

Introduction

Symbiotic relationships between bacteria and eukaryotes are pervasive and range from loose associations to obligate interdependencies (McFall-Ngai *et al.* 2013; Kostic *et al.* 2013). The evolution of a host-associated lifestyle is typically accompanied by the loss of generalist characteristics, limiting a symbionts' capacity to compete and survive in other environments. This in turn results in host specialization (Bobay & Ochman 2017; Sriswasdi *et al.* 2017). In particular, bacteria with longstanding associations are often host-specific and undergo marked genomic changes (Toft & Andersson 2010). Among the most extreme examples are primary endosymbionts of plant-sap feeding insects. These obligate mutualists reside within host cells, are vertically inherited through the germ-line, and have experienced extreme genome reduction due to population bottlenecks and genetic drift (McCutcheon & Moran 2012).

Based on phylogenetic analyses, host specialization has also been inferred for many gut symbionts, as bacterial lineages are frequently found to be exclusively associated with particular hosts (Ley *et al.* 2008; Oh *et al.* 2010; Ochman *et al.* 2010; Eren *et al.* 2015; Moeller *et al.* 2016; Kwong *et al.* 2017). This is remarkable considering that gut symbionts are horizontally transmitted, and are exposed at least at some point to the environment outside the host, which in principle provides opportunities for host switching. This leads to the question whether the realized niche of gut symbionts (the conditions where the bacteria actually live) is different from their fundamental niche (the conditions where they can live). If there is a difference, the next question is which factors limit niche realization, such as

competition with other bacteria, dispersion, or host selection (Macarthur & Levins

2015; Hutchinson 1957).

In the case of the gut symbiont *Lactobacillus reuteri*, strains isolated from mice are

capable of colonizing the mouse gut, whereas those from humans, pigs, or chickens

are not, suggesting that host association in this case has resulted in the restriction of

the fundamental niche (Oh *et al.* 2010; Frese *et al.* 2011). In contrast, in another

study it was shown that bacterial communities from diverse habitats can colonize

and persist in the mouse gut (despite the fact that these species naturally do not

occur in the mouse gut), suggesting that a species' realized niche is frequently more

restricted than its fundamental niche (Seedorf *et al.* 2014). A notable difference

between the two studies is that host specificity of *L. reuteri* was tested in mice that

were free of *Lactobacilli*, but otherwise harbored a conventionalized microbiota,

whereas in the second study, most experiments were carried out in microbiota-free

mice. Hence, interbacterial competition may be an important factor defining the

realized niche (Seedorf *et al.* 2014).

Given that both experimental and phylogenetic evidence is required to determine

host specialization, our understanding of host specialization is limited for most gut

symbionts. Moreover, little is known about the underlying mechanisms and the

genomic changes accompanying host-specific evolution of gut symbionts. Selective

forces acting on gut symbionts may differ between hosts due to varying degrees of

population bottlenecks during transmission, or due to differences in dietary

preferences, gut structure or host physiology, resulting in distinct evolutionary

patterns.

A good model to study host specialization of bacterial inhabitants is the gut

microbiota of corbiculate bees (Kwong & Moran 2015). Most species of honey bees,

bumble bees, and stingless bees share a specialized core gut microbiota that is

composed of five phylotypes (strains sharing ≥97% 16S rRNA sequence identity as

estimated from amplicon sequencing studies): the gammaproteobacterium

Gilliamella, the betaproteobacterium *Snodgrassella alvi*, two Lactobacilliales (Firm5

and Firm4), and a Bifidobacterium (Cox-Foster *et al.* 2007; Moran *et al.* 2012; Corby-

Harris *et al.* 2014). These phylotypes are likely to have been acquired in a last

common ancestor of the corbiculate bees, as they are widely distributed among

contemporary species of honey bees, bumble bees, and stingless bees (Kwong *et al.*

2017). Moreover, there is evidence for host specialization and coevolution, because

strains isolated from the three groups of corbiculate bees separate into divergent

sublineages for most phylotypes (Koch *et al.* 2013; Kwong *et al.* 2014; Ellegaard *et*

al. 2015; Zheng *et al.* 2016; Kwong *et al.* 2017; Steele *et al.* 2017).

The best-studied member of the bee gut microbiota with respect to host

specialization is *S. alvi* (Kwong & Moran 2015). Reciprocal mono-colonization

experiments of microbiota-depleted bees showed that *S. alvi* isolates from honey

bees (*Apis mellifera*) colonize poorly the gut of bumble bees (*Bombus impatiens*),

and vice versa, suggesting that the host-specific evolution of these isolates has led to

specialization (Kwong *et al.* 2014). Based on the comparison of three *S. alvi*

genomes, it was suggested that bumble bee isolates tend to have smaller genomes

and contain larger amounts of mobile elements than honey bee isolates. Genomic

differences were also identified among isolates from different host groups (honey

bees and bumble bees) for the phylotype *Gilliamella*: honey bee isolates encoded

more carbohydrate-related functions than bumble bee isolates (Kwong *et al.* 2014).

However, recent genome sequencing of a larger number of *Gilliamella* strains

revealed that some isolates from honey bees have genomes as small as those from

bumble bees, suggesting that a large metabolic repertoire is not necessarily needed

for colonization of the honey bee gut (Zheng *et al.* 2016; Ludvigsen *et al.* 2017;

Steele *et al.* 2017).

For the other phylotypes of the bee gut microbiota, little is known about the link

between phylogeny, host range, and genome features (Ellegaard *et al.* 2015; Kwong

et al. 2017). One of the most widely distributed and abundant phylotypes of the bee

gut microbiota is *Lactobacillus* Firm5. In the gut of honey bees (*Apis mellifera*)*,* four

deep-branching monophyletic sublineages have been identified for this phylotype,

and different species names have been proposed (Olofsson *et al.* 2014). Strains

belonging to these four sublineages typically co-occur in individual bees *(Ellegaard*

and Engel, in revision), and can vary by up to 40% in gene content (Ellegaard *et al.*

2015), suggesting that these sublineages may have adapted to distinct metabolic or

spatial niches within the honey bee gut. Interestingly, Firm5 strains isolated from

other corbiculate bees seem to belong to different sublineages than the honey bee

isolates, as indicated by single gene phylogenies (Kwong *et al.* 2017). Moreover, a

divergent Firm5 strains from bumble bees has been isolated and described as a new

species, *Lactobacillus bombicola* (Praet *et al.* 2015). Given that Firm5 strains can be

cultured, and that the honey bee is amenable to experimental colonization, this

phylotype represents an excellent opportunity to study evolutionary trajectories of

host adaptation and the consequences for the fundamental and realized niche of this

gut symbiont.

Here, we used gnotobiotic bee experiments, genome sequencing, and comparative

genomics to address host specialization in the Firm5 phylotype. First, we show that

isolates from honey bees and bumble bees belong to distinct sublineages suggesting

longstanding host-specific associations. Second, we provide experimental evidence

that both host selection and interbacterial competition contribute to host

specialization. Third, our comparative genome analysis reveals marked differences

in carbohydrate utilization capacities between honey bee and bumble bee isolates

suggesting that metabolic flexibility gives honey bee isolates a competitive

advantage over bumble bee isolates to establish in the honey bee gut.

Results

Bumble bee and honey bee isolates belong to separate sublineages of the Firm5 phylotype.

We sequenced the genomes of 15 new isolates of the Firm5 phylotype. Five isolates were obtained from the honey bee *Apis mellifera* and ten isolates from three different bumble bee species (five from *Bombus pascuorum*, four from *Bombus bohemicus*, and one from *Bombus terrestris*). All bees were collected in Western Switzerland (**Table S1**). We also included 12 previously sequenced isolates (one from a bumble bee, the others from honey bees) to be more comprehensive in our analyses. All 27 isolates shared >95% sequence identity across the full-length 16S rRNA gene (**Table S2**), suggesting that they all belong to the Firm5 phylotype. Isolates from conspecific hosts tended to have higher 16S rRNA sequence identities. The draft genomes of the 15 newly sequenced isolates consisted of 11-24 contigs with total lengths of 1.63-2.11 Mb, which is in the range of the previously sequenced Firm5 strains (**Table S1**). While the genomes of the bumble bee isolates tended to be smaller (1.63-1.70 Mb) than those of the honey bee isolates (1.68-2.15 Mb), genome synteny was largely conserved across the entire Firm5 phylotype (**Figure S1 and S2**).

To assess the evolutionary relationship between the 27 sequenced Firm5 strains, we inferred a genome-wide phylogeny (including 15 close and three more distant outgroup strains, see methods) (**Figure 1**), and calculated pairwise average nucleotide identities (ANI) (**Figure S3, Table S3**). This analysis showed that the

Firm5 strains fall into six monophyletic sublineages with >91% ANI for within-

lineage divergence and <86% ANI for between-lineage divergence. Four of these six

sublineages consisted of only honey bee isolates and corresponded to the previously

identified Firm5 sublineages (Ellegaard *et al.* 2015). The two other sublineages

consisted of only bumble bee isolates and formed a monophyletic clade within

Firm5 (**Figure 1**). One sublineage comprised isolates from three different bumble

bee species (*B. lapidarus, B. terrestris, B. pascuorum*) including the previous isolate

described as species *L. bombicola* (Praet *et al.* 2015)*. The other sublineage

comprised exclusively isolates from *B. bohemicus*. Based on its deep divergence

from the other sublineages (ANI <80%, **Figure S3**), this second sublineage of

bumble bee isolates is likely to represent a novel species, for which we propose the

Candidatus name '*Lactobacillus bohemicus'*.

Out of the 27 Firm5 isolates included in the current study, five isolates (ESL0262,

ESL0234, ESL0236, ESL0245, ESL0247) from three different sublineages were

identical or almost identical to other isolates (ANI >99.99%, **Table S3**). In all cases,

the nearly identical isolates were obtained from the same individual. Hence, they

were excluded from all subsequent analyses to avoid biases due to repeated

sampling of the same genotype.

In summary, our phylogenetic analysis of the Firm5 phylotype revealed a pattern

suggesting host specialization, because strains of each of the six deep-branching

sublineages were exclusively associated with either honey bees or bumble bees.

Bumble bee strains can colonize microbiota-depleted honey bees, but are outcompeted by honey bee strains.

To test if the differences in the realized niche of Firm5 isolates are due to host specialization, we experimentally tested the ability of bumble bee strains to colonize the honey bee gut. In the absence of competitors (i.e. when microbiota-depleted bees were mono-colonized), we found that all bumble bee strains were able to colonize the honey bee hindgut (**Figure 2A**). The number of recovered bacterial cells at day 5 post colonization (10^6 – 10^8 CFUs per gut) was substantially higher than in the inoculum (**Figure S5A**) indicating active growth of the bumble bee strains in the honey bee gut. Thus, we can conclude that the fundamental niche of the bumble bee strains includes the honey bee gut. However, the colonization levels were slightly lower compared to mono-colonizations with honey bee strains, which reached 10^8 – 10^9 CFUs per gut (**Figure 2A**). Moreover, the percentage of successfully colonized bees varied between strains and was overall also lower for bumble bee (10-80%) than for honey bee strains (80-100%).

To test if the colonization success depends on the number of cells in the inoculum, we colonized microbiota-depleted honey bees with different inocula of the bumble bee Firm5 strain ESL0228 (**Figure S5B**). With the lowest inoculum, no colonization was obtained at day 5 post colonization (n=10), while with the highest inoculum, all bees were colonized, yielding again between 10^6 – 10^8 CFUs per gut (**Figure 2B**). The relatively high number of bacteria that was needed to achieve a robust colonization suggests that stronger selection is at play on bumble bee than on honey bee Firm5 strains for gut colonization.

To test for the effect of competitive exclusion between strains, we co-colonized

microbiota-depleted bees with the bumble bee Firm5 strain ESL0228 and a mix of

four honey bee strains (ESL0183, ESL0184, ESL0185, and ESL0186), each from one

of the four divergent sublineages. We kept the number of bacteria in the inoculum

constant for the honey bee strains (1:1:1:1), but provided the bumble bee strain at

ratios of 1:1, 10:1, or 100:1 relative to the honey bee strains (**Figure S5B**). All bees

in the experiment (n=30, n=10 per treatment) were successfully colonized by the

Firm5 phylotype and the total numbers of CFUs per gut were in the same range as

for the mono-colonizations (10^8 – 10^9 CFUs). We used amplicon sequencing of a

short fragment of a conserved housekeeping gene that allowed us to determine the

relative abundance of the five Firm5 strains tested in the community (see Methods).

This analysis revealed that overall all four honey bee strains successfully colonized

and coexisted in the gut, except for strain ESL0184 which was absent from a few

samples (**Figure 2C**). In contrast, the bumble bee strain ESL0228 was detected in

only a few bees and at very low relative abundance (<0.1%), even when inoculated

with a ratio of 100:1.

Collectively, these experiments show that bumble bee strains of the Firm5

phylotype are capable of colonizing the honey bee gut, but consistent colonization

can only be achieved with a relatively high inoculum and when honey bee strains

are absent. Therefore, we conclude that both host selection and interbacterial

competition contribute to the restriction of the realized niche of bumble bee strains.

Firm5 strains harbor a large gene pool of phylotype-specific functions of which few are conserved.

In order to identify genomic characteristics that may contribute to host specialization among Firm5 strains, we carried out a detailed comparative genome analysis. We first determined the distribution of the entire pan genome across the analyzed Firm5 strains. We included 15 divergent outgroup strains in this analysis (i.e. strains not belonging to the Firm5 phylotype, see **Figure 1** and methods) to identify Firm5-specific gene families that could play a role in adaptation to the bee gut environment. In total, 8,248 pan genome gene families were identified across the 37 genomes (i.e. gene families present in at least one genome), of which 2,131 gene families were specific to the Firm5 strains. Of those, 571 and 1,222 gene families were only found in bumble bee and honey bee strains, respectively, and 338 gene families were shared across the two hosts (**Figure 3A**).

Despite this relatively large gene pool of Firm5-specific functions, few gene families were conserved (5.6% of the shared gene families and 1.4% and 1.6% of the host-specific gene families, **Figure 3A**). Among the gene families conserved across all Firm5 strains, we found an ABC transporter system for branched chain amino acids and two putative adhesin genes (DUF4097). Amino acid transporter genes were also present among the few conserved gene families specific to the honey bee strains, while those specific to the bumble bee strains were almost all annotated as either hypothetical proteins or transcriptional regulators, providing few clues about their functional roles for host adaptation (**Dataset S1**). Altogether, this analysis revealed

very few phylotype- or host-specific gene functions as potential candidates for

general determinants of host adaptation across the analyzed Firm5 strains.

Firm5-specific gene content is restricted to sublineages

As strains from the same host can belong to divergent sublineages, it is possible that

sublineage-specific gene functions are involved in host specialization, e.g. by

adaptation to different metabolic niches within the gut. Such genes could also

explain the ability of the four honey bee sublineages of Firm5 to coexist in bees

(Ellegaard and Engel, in review).

Indeed, we found that a relatively large number of the Firm5-specific pan gene

content is confined to strains of the same sublineage (840 of 1,222 for honey bee

strains, 532 of 571 for bumble bee strains). However, as for the previous analysis,

relatively few of these sublineage-specific gene families were conserved, i.e. present

in all strains of a given sublineage (**Figure 3B**). In the honey bee sublineage Firm5-2

(*L. helsingborgensis*) 53 gene families were conserved (i.e. 34% of the sublineage-

specific gene content), including several sugar transporter genes and a genomic

island for the breakdown of rhamnogalacturonan, a major polysaccharide of pectin

(**Dataset S1**). This genomic island was also found in a recent metagenomic study to

correlate in abundance with core genes of this sublineage *(Ellegard and Engel, in*

review), suggesting that rhamnogalacturonan utilization is a conserved function of

strains belonging to Firm5-2 (*L. helsingborgensis*). For the other three honey bee

sublineages, only 9-20 gene families (0.3%-1.1%) were conserved (**Figure 3B**). In

sublineage Firm5-3 (*L. melliventris*), functions for rhamnose utilization were found

to be conserved, whereas the annotations of the gene families in the other two

sublineages provided little functional insights (**Dataset S1**). The same was the case

for the conserved gene families found in the two bumble bee sublineages (9 and 59

gene families, **Figure 3B**), of which most were annotated as hypothetical proteins

(**Dataset S1**). Overall, these results indicate a high degree of gene content variability

among strains from the same host and from the same sub-lineage.

Honey bee strains harbor a larger diversity of carbohydrate-related functions

than bumble bee strains.

The high degree of gene content plasticity within the Firm5 phylotype prompted us

to look at the functional composition of the entire Firm5-specific gene pool. This

analysis revealed marked differences between honey bee and bumble bee strains

with respect to carbohydrate-related functions. While 'Carbohydrate transport and

metabolism' was by far the most dominant COG category (COG 'G') among the gene

families specific to the honey bee strains (184 gene families, 50% of those with COG

annotation), this category was nearly absent among the gene families specific to the

bumble bee strains (4 gene families, 10% of those with COG annotation) (**Figure**

3C). In fact, most gene families specific to bumble bee strains had no COG annotation

at all. Analysis of the sublineage-specific gene content revealed a similar pattern. For

three of the four honey bee sublineages, 'Carbohydrate transport and metabolism'

was by far the most abundant COG category among the sublineage-specific gene

content, while for the two bumble bee sublineages this category was much less

prominent (**Figure 3D**).

This trend was confirmed by analyzing the overall abundance of carbohydrate-

related functions per Firm5 strain. Both relative and total number of genes assigned

to COG category 'G' was higher for most honey bee strains compared to bumble bee

strains or outgroup strains (**Figure 4A, Figure S6)**. However, the Firm5-1

sublineage represented an exception to this pattern. All three strains of this

sublineage encoded fewer COG category 'G' genes than other honey bee strains. A

large proportion of the genes assigned to COG category 'G' encoded

phosphotransferase systems (PTSs), i.e. transporters involved in sugar.

Correspondingly, these gene families showed a similar distribution as the COG

category 'G' genes across the Firm5 strains, with most honey bee strains harboring a

much larger number of PTS genes than bumble bee strains (**Figure 4B)**.

PTS transport systems are often genetically linked to glycoside hydrolases (GHs),

which mediate the cleavage of sugar residues from polysaccharides or other

glycosylated compounds. To assess if bee gut bacteria harbor a specific arsenal of

these sugar-cleaving enzymes, we identified all GH genes in the analyzed genomes.

As for COG category 'G' and PTS transporters, we found a larger number of GH genes

for honey bee strains compared to bumble bee strains (**Figure 4C, Dataset S3)**.

Some honey bee strains harbored twice as many GH genes than bumble bee strains.

However, there was remarkable variation in the number of GH genes among the

honey bee strains, both within and across sublineages. Specifically, all strains of

sublineages Firm5-3 and Firm5-4 harbored a relatively high number of GH genes,

while strains of sublineage Firm5-1 varied substantially in the number of GH genes,

and those of sublineage Firm5-2 were consistently low.

The identified GH genes belonged to 79 different gene families (**Figure 4D, Dataset S3**), of which 43 were specific to the Firm5 phylotype. Most of these (67%) were only detected among honey bee strains, 19% were shared, and only 14% were specific to bumble bee strains. Moreover, honey bee strains also shared more GH gene families with the outgroup strains than bumble bee strains (18 vs 1 gene families).

While the substrate specificity of GH gene families cannot be unambiguously inferred from sequence data, many of the Firm5-specific GH gene families included glucosidases, fucosidases, mannosidases, xylosidases, and arabinofuranosidases (e.g. GH29, GH38, GH39, GH43, and GH51), as based on the CAZY (Carbohydrate-Active enZYmes) database classification (**Figure 4E).** A similarity search against the publicly available non-redundant database NCBI nr (NCBI Resource Coordinators 2018) revealed that many of the Firm5-specific GH families, especially those exclusively present among the honey bee strains, have best hits to other taxonomic groups than lactobacilliales (**Figure S7**). While these gene families may have been acquired by horizontal gene transfer or secondarily lost in other lactobacillus, their limited distribution among lactobacilliales suggests specific functions in the bee gut environment.

Overall, the analysis of the carbohydrate-related gene content shows that Firm-5 strains from honey bees harbor a larger diversity of PTS transporters and glycoside hydrolases than bumble bee strains. However, differences in the type and abundance of these functions between strains and sublineages suggest that honey bee strains have diversified in their ability to utilize different sugar resources.

Firm5 strains from bumble bees, but not from honey bees harbor class II bacteriocins

Most gene families specific to the bumble bee strains were annotated as hypothetical proteins (**Figure 3C and D**), providing no insights about the possible genetic basis of adaptation to the bumble bee gut environment. However, we found several short open-reading frames encoding putative class II bacteriocins. Bacteriocins are small peptide toxins that act against closely related bacterial strains (Cotter *et al.* 2003). In the case of class II bacteriocins, an ABC-like transporter usually facilitates toxin secretion, and a dedicated immunity protein provides self-protection. Except for strain ELS0228, all bumble bee strains harbored at least one class II bacteriocin gene with homology to lactococcin 972, described to inhibit septum formation (Martínez *et al.* 2000). Consistent with the genetic organization of lactococcin loci in other species (Letzel *et al.* 2014), putative immunity proteins and ABC transporter genes were encoded downstream of the bacteriocin gene **(Figure 5)**. We identified four distinct genomic regions with this genetic organization. All four regions exhibited a high degree of genomic plasticity, with many non-conserved open reading frames close by (**Figure 5 and Figure S8**). Each bacteriocin locus was specific to one of the two bumble bee sublineages and only present in a subset of the analyzed strains. In sublineage Firm5-5, one region encoded two adjacent bacteriocin loci, and in several instances, one of the immunity protein or toxin genes was pseudogenized **(Figure 5)**. Strikingly, none of these genomic regions were present in the honey bee strains analyzed, suggesting that this genetic feature may be specific to bumble bee strains. However, we found

homologs of genes for helveticin-J in honey bee strains, another protein with known

bactericidal activity against related bacteria. This gene family was conserved in all

strains of Firm5 as well as in some of the outgroup strains **(Figure S9**).

In summary, while the two bumble bee sublineages of Firm5 harbored a large pool

of host-specific gene families, bacteriocins were the only conserved genes with

annotated functions, and thus the only identified candidates to play a role for niche

specialization in the present state of our knowledge.

Discussion

In this study, we combined honey bee colonization experiments with comparative genomics to investigate host specialization of Lactobacillus Firm5, a dominant gut symbiont of social bees. Our results show that strains isolated from honey bees and bumble bees belong to separate, highly divergent sublineages of the Firm5 phylotype, which parallels phylogenetic analysis of other bee gut symbionts (Kwong *et al.* 2014; Zheng *et al.* 2016; Kwong *et al.* 2017; Steele *et al.* 2017).

Interestingly, all tested Firm5 strains from bumble bees were able to colonize the gut of microbiota-depleted honey bees, indicating that the divergent evolution of Firm5 strains from different bee species has not resulted in strict host specialization. However, the percentage of successfully colonized bees as well as the number of bacterial cells per gut were both lower for bumble bee strains compared to honey bee strains. Only by increasing the number of bacterial cells in the inoculum by 100-fold, were we able to achieve reliable colonization, which suggests strong negative selection of bumble bee strains during passage through the honey bee gut, possibly due to the lack of host-specific adaptations.

However, we currently do not know whether, inversely, bumble bee strains would perform better, and honey bee strains worse, in microbiota-depleted bumble bees, which would provide further evidence for host-specific adaptations. Nevertheless, our findings show that the fundamental niche of Firm5 strains is larger than the realized niche and includes host species from other bee genera. This is in agreement with a previous study showing that selected bacteria from diverse environments,

including zebrafish or termite gut, can establish in the gut of germ-free mice

(Seedorf *et al.* 2014). Moreover, the gut symbionts *S. alvi* (social bee gut) and *L.*

reuteri (vertebrate gut) - for which host specialization has been experimentally

demonstrated – are both able to colonize non-native hosts, although at much lower

levels than native hosts (Frese *et al.* 2011; Kwong *et al.* 2014).

If related species have overlapping fundamental niches, such as in the case of the

Firm5 isolates from honey bees and bumble bees, classical ecological theory

predicts that differences in the realized niche are due to interactions leading to

competitive exclusion of one of the species (Macarthur & Levins 2015). This is

exactly what we observed when we colonized microbiota-depleted honey bees with

a community of five Firm5 strains, including one bumble bee strain and four honey

bee strains. The bumble bee strain did not establish in any of the tested bees,

although we inoculated the microbiota-depleted bees with up to 100x more

bacterial cells of the bumble bee strain than the four honey bee strains. This clearly

shows that the tested bumble bee strain has a competitive disadvantage in the

honey bee gut compared to honey bee strains. Similar results were obtained for *S.*

alvi, when a non-native strain was challenged with a native competitor for gut

colonization (Kwong *et al.* 2014).

Notably, the phylogenetic similarity between the bumble bee and the honey bee

strains of the Firm5 phylotype cannot be the underlying reason for the competitive

exclusion of the bumble bee strain, because the four more closely related honey bee

strains were able to coexist. Therefore, it is more likely that coevolution of the

honey bee strains in the honey bee gut has resulted in reciprocal adaptations

facilitating coexistence.

Honey bees live in large colonies and engage in frequent social interactions. This

results in constant exposure to bacteria from nestmates, thereby providing few

opportunities for bacteria from non-native hosts to establish in the gut of young

worker bees during community assembly. However, even when given the ecological

opportunity for gut colonization (as in our colonization experiments), bumble bee

strains cannot reliably colonize the gut. Hence we conclude that the competitive

disadvantage relative to honey bee strains as well as the suboptimal host adaptation

are two important factors contributing to the exclusion of bumble bee Firm5 strains

from the honey bee gut in natural populations.

The bacteria-mediated exclusion of the bumble bee Firm5 strains from the honey

bee gut could arise via direct antagonistic interactions between bacteria (e.g. via

bacterial toxins), or from resource competition. We identified a number of genes

encoding bacteriocins, which are known to mediate interbacterial killing

(Kommineni *et al.* 2015). These genes were either shared by strains isolated from

both hosts, or they were specific to the bumble bee strains. However, we did not

identify any toxin genes specific to the honey bee strains, which could mediate

possible antagonistic effects towards bumble bee strains and hence hinder their

colonization in the gut. Therefore, we conclude that based on the current

knowledge, it is unlikely that direct antagonistic interactions explain the

competitive exclusion of bumble bee strains from the honey bee gut.

Biofilm formation at the host epithelium has been shown to be a crucial factor for

the colonization success and competitiveness of the murine gut symbiont *L. reuteri*

(Frese *et al.* 2011; Duar *et al.* 2017). The honey bee gut symbiont *S. alvi* also

colonizes the epithelial surface and forms biofilm-like structures, making it

conceivable that competition for adherence is also a critical factor for colonization

in the bee gut (Kwong & Moran 2016). However, bacteria of the Firm5 phylotype do

not attach to the host epithelium, as shown by fluorescence in situ hybridization

experiments, but rather colonize the gut lumen in the rectum, where competition for

space seems less likely to be a predominant limiting factor (Martinson *et al.* 2012).

Moreover, our genomic analysis did not identify genes involved in host interaction

or adherence to be specific to strains from one of the two host groups.

Instead our analyses suggest competition for nutrients to be an important

underlying factor for competitiveness. We found several metabolic genes that were

conserved across all analyzed honey bee strains but not present in any of the

bumble bee strains. Among them were several genes encoding amino acid

transporters, which may facilitate the acquisition of organic nitrogen from the host

diet. The predominant energy metabolism of the Firm5 phylotype is predicted to be

fermentation of dietary carbohydrates, which is not surprising given that the diet of

social bees (pollen and nectar) is rich in simple sugars, polysaccharides (pectin,

hemicellulose and cellulose), and other glycosylated compounds (e.g. flavonoids)

(Engel *et al.* 2012; Ellegaard *et al.* 2015; Kešnerová *et al.* 2017). Strikingly, we found

that most honey bee strains harbored a much larger arsenal of gene functions for

carbohydrate metabolism and transport compared to bumble bee and outgroup

strains. This suggests that honey bee strains have a greater capacity to utilize diet-

derived carbohydrates, which may give them a growth advantage over bumble bee

strains in the bee gut. A similar trend has also been observed for strains of the gut

symbiont *G. apicola* (Kwong *et al.* 2014)*.*

Bumble bees and honey bees have a similar dietary regime, as both eat nectar and

pollen. Hence, the reason why bumble bee strains harbor significantly fewer

carbohydrate-related functions is currently unclear. One possibility could be that

these strains colonize a different niche in the gut. However in this case, we would

not expect to see competition for colonization among honey bee and bumble bee

strains. Another possible factor could be differences in the life cycle of bumble bees

and honey bees. Honey bees maintain perennial colonies of large population sizes,

while bumble bees build smaller colonies from a single overwintering queen every

year. This presents a population bottleneck for the bacterial community in the gut of

bumble bees, which reduces genetic variation among gut symbionts by genetic drift,

which in turn might lead to gene loss and decrease the selective pressure imposed

by related bacteria. Consequently, it would slow the acquisition of genes that would

allow bacteria to use diverse carbohydrates. The genomes of bumble bee strains

tended to be slightly smaller than those of honey bees, which seems to be consistent

with this hypothesis. Strikingly, also in the host-specialized vertebrate gut symbiont

L. reuteri, it was speculated that genomic differences in genome size and pan

genome diversity may be due to differences in population bottlenecks across hosts

(Frese *et al.* 2011).

Interestingly, almost none of the carbohydrate-related gene families specific to

honey bee strains of the Firm5 phylotype were conserved across all analyzed

genomes, suggesting that the genetic basis of host adaptation in regard of

carbohydrate metabolism differs between strains. A large proportion of the

carbohydrate-related gene content was specifically associated with one of the four

sublineages of honey bee Firm5 strains. However, only a few of these functions were

conserved (e.g rhamnogalacturonan and rhamnose utilization in Firm5-2 and

Firm5-3, respectively) and the number of carbohydrate-related functions varied

markedly among strains of some sublineages. Together these findings suggest that

metabolic functions with possible roles for adaptation are frequently gained and/or

lost, raising the possibility that metabolic flexibility in itself represents an important

adaptation to honey bees compared to bumble bees. This may also explain why the

four tested honey bee strains were able to coexist in individual bees. In this context,

it is important to highlight that the total number of CFUs per bee was similar

between colonizations with the individual honey bee strains and the community,

suggesting that the four honey bee strains in the community have overlapping

niches, but segregate (spatially or metabolically) when present together. Moreover,

these results provide additional evidence that interbacterial competition within

phylotypes plays an important role in the bee gut environment.

In conclusion, our study advances the understanding of host specialization of gut

symbionts. While previous studies on *L. reuteri* have shown that host interaction,

and specifically colonization of the gut surface, determine host specificity, we

provide evidence for metabolic flexibility that may facilitate adaptation to the host

diet and hence the competitive exclusion of non-adapted strains. As specific dietary

preferences are common among animals, similar processes may also be a

determining factor of host specialization among other gut symbionts.

Materials and methods

544

545 **Bee sampling, bacterial culturing and DNA isolation.**

546 Bumble bees were collected from flowers in different locations in Western

547 Switzerland as indicated in **Table S1**. Honey bees were sampled from two healthy

548 looking colonies in the same region located at the University of Lausanne. Within 6h

549 after sampling, bees were immobilized on ice and the entire gut was dissected with

550 sterile scissors and forceps. Each gut tissue was individually placed into a screw cap

551 tube containing 1ml 1x PBS and glass beads (0.75-1mm, SIGMA) and homogenized

552 with a bead-beater (FastPrep-24 5G, MP Biomedicals) for 30s at speed 6.0. Serial

553 dilutions of the gut homogenates were plated on MRS agar and incubated in an

554 incubator at 34°C in an anaerobic chamber (Coy laboratories, MI, USA) containing a

555 gas mix of 8% H_2, 20% CO_2 and 72% N_2. After 3-5 days of incubation, single colonies

556 were picked, restreaked on fresh MRS agar and incubated for another 2-3 days.

557 Then, a small fraction of each restreaked bacterial colony was resuspended in lysis

558 buffer (10 mM Tris-HCl, 1 mM EDTA, 0.1% Triton, pH 8, 2 mg/ml lysozyme and

559 1mg/ml proteinase K) and incubated in a thermocycler (10 min 37°C, 20 min 55°C,

560 10 min 95°C). Subsequently, a standard PCR with universal bacterial primers (5'-

561 AGR GTT YGA TYM TGG CTC AG-3', 5'-CCG TCA ATT CMT TTR AGT TT-3') was

562 performed on 1 μl of the bacterial lysate and the resulting PCR products were sent

563 for Sanger sequencing. Sequencing reads were inspected with Geneious v6

564 (Biomatters Limited) and compared to the NCBI nr database (NCBI Resource

565 Coordinators 2018) using BLASTN. Isolates identified to have high similarity (i.e.

566 >95% sequence identity) to honey bee strains of the Firm-5 phylotype were stocked

in MRS broth containing 25% glycerol at -80°C. Genomic DNA was isolated from

fresh bacterial cultures of the strains of interest using the GenElute Bacterial

Genomic DNA Kit (SIGMA) according to manufacturers instructions. Bumble bees

were genotyped based on the COI gene by performing a PCR on DNA extracted from

the carcass, sending the PCR product for Sanger sequencing, and searching the

resulting sequence read by BLASTN against the NCBI nr database.

Genome sequencing, assembly and annotation.

Genome sequencing libraries were prepared with the TruSeq DNA kit and

sequenced on the MiSeq platform (Illumina) using the paired-end 2x250-bp

protocol at the Genomic Technology facility (GTF) of the University of Lausanne. The

preliminary genome sequence analysis was carried out in the framework of the

student course 'Sequence-a-genome (SAGE)' at the University of Lausanne in 2016-

2017. In short, the resulting sequence reads were quality-trimmed with

trimmomatic v0.33 (Bolger *et al.* 2014) to remove adapter sequences and low

quality reads using the following parameters: ILLUMINACLIP:TruSeq3-PE.fa:3:25:6

LEADING:9 TRAILING:9 SLIDINGWINDOW:4:15 MINLEN:60. The quality-trimmed

reads were assembled with SPAdes v.3.7.1 (Bankevich *et al.* 2012), using the "--

careful" flag and multiple k-mer sizes (-k 21,33,55,77,99,127). Small contigs (less

than 500 bp) and contigs with low kmer coverage (less than 5) were removed from

the assemblies, resulting in 11-22 contigs per assembly. The contigs of each

assembly were re-ordered according to the complete genome of the honey bee

strain ESL0183 using MAUVE v2.4 (Rissman *et al.* 2009). The origin of replication

was set to the first base of the *dnaA* gene, which coincided with the sign change of

the GC skew. The ordered assemblies were checked by re-mapping the quality-

trimmed reads (**Figure S2**). Except for a few prophage regions that showed

increased read coverage, no inconsistencies in terms of read coverage or GC skew

were revealed suggesting that the overall order of the contigs was correct. The

median read coverage of the sequenced genomes ranged between 135x-223x

(**Figure S2**). The genomes were annotated using the 'Integrated Microbial Genomes

and Microbiomes' (IMG/mer) system (Markowitz *et al.* 2014).

Inference of a genome-wide phylogeny.

Gene families, i.e. groups of homologous genes, were determined using OrthoMCL

(Li *et al.* 2003) between all publicly available and newly sequenced genomes of the

Firm5 phylotype as well as a set of outgroup genomes of other lactobacilli strains.

The outgroup strains were selected based on their phylogenetic relatedness with

the Firm5 phylotype using a previously published phylogeny of the entire genus

Lactobacillus (Zheng *et al.* 2015). Based on this analysis, we included the genomes of

15 closely related outgroup strains that belong to the same *Lactobacillus* clade as

Firm5 ('delbrueckii group') and three more distantly related strains for rooting the

phylogeny. All-against-all BLASTP searches were conducted with the proteomes of

the selected genomes, and hits with an e-value of $\leq 10^{-5}$ and a relative alignment

length of >50% of the query and the hit protein lengths were kept for OrthoMCL

analysis. All steps of the OrthoMCL pipeline were executed as recommended in the

manual and the mcl program was run with the parameters '--abc -I 1.5'.

The core genome phylogeny was inferred from 408 single copy orthologs extracted

from the OrthoMCL output (i.e. gene families having exactly one representative in

every genome in the analysis). The protein sequences of each of these core gene

families were aligned with mafft (Katoh *et al.* 2017). Alignment columns

represented by less than 50% of all sequences were removed and then the

alignments were concatenated. Core genome phylogenies were inferred on the

concatenated trimmed alignments using RAxML (Stamatakis 2014) with the

PROTCATWAG model and 100 bootstrap replicates.

Comparison of genome structure, genome divergence, and gene content.

To compare and visualize whole genomes we used the R-package genoPlotR (Guy *et*

al. 2010). BLASTN comparison files were generated with DoubleACT (www. hpa-

bioinfotools.org.uk) using a bit score cutoff of 100. To estimate sequence divergence

between genomes, we calculated pairwise average nucleotide identity (ANI) with

OrthoANI (Lee *et al.* 2016) using the exectutable 'OAT_cmd.jar' with the parameter '-

method ani'.

For analyzing the distribution of gene families across Firm5 sublineages and closely

related outgroup strains, we carried out a second OrthoMCL analysis, in which we

excluded the three distantly related outgroup strains. To remove redundancy in our

database, we also excluded the genomes of five Firm5 isolates that were identical, or

almost identical, to other Firm5 strains in the analysis, based on ANI values of

>99.9%. This resulted in a total 37 genomes (22 Firm5 genomes and 15 outgroup

genomes) that were included in the analysis. BLASTP and OrthoMCL were run with

the same parameters as before. Gene family subsets of interest (e.g. families specific

to honey bee, bumble bee or outgroup strains) were extracted from the OrthoMCL

output file using custom-made Perl scripts. COGs (Cluster of Orthologous Groups)

were retrieved from IMG/mer genome annotations (Markowitz *et al.* 2014).

For the detection and visualization of the genomic regions encoding bacteriocin

genes, Bagel3 (van Heel *et al.*) and MultiGeneBlast (Medema) were used. For the

MultiGeneBlast analysis, bacteriocins-encoding genomic regions of strains ESL0233

and ESL0247 served as query sequences for searching a custom-made database

composed of all non-redundant Firm5 genomes.

Identification of glycoside hydrolase gene families.

Glycoside hydrolase gene families were identified in all analyzed genomes

(excluding the redundant Firm5 strains and the three distant outgroup strains)

using the command-line version of dbCAN (**D**ata**b**ase for automated **C**arbohydrate-

active enzyme **An**notation) (Yin *et al.*). In short, we searched each genome against

dbCAN using hmmscan implemented in HMMER v3 (Eddy 2009). The output was

processed with the parser script 'hmmscan-parser.sh', and genes with hits to Hidden

Markov Models of glycoside hydrolase families were extracted.

For determining the taxonomic distribution of related genes, we searched one

homolog of each glycoside hydrolase gene family against the NCBI *nr* database

(NCBI Resource Coordinators 2018) using BLASTP. The taxonomy of the first 50

BLASTP hits (e-value <10-5) was extracted at the family level using the Perl script

'Tax_trace.pl' and the database files nodes.dmp and names.dmp. The latter two files

contain the NCBI taxonomy nodes and names.

Bee colonization experiments.

Newly emerged, microbiota-depleted bees were generated as described in (Emery

et al. 2017) and colonized within 24-36h after pupal eclosion. To this end, bacterial

strains were grown on MRS agar containing 2% fructose and 0.2% L-cysteine-HCl

from glycerol stocks for two days in an anaerobic chamber at 34°C. Then, 1-10

colonies were inoculated into 5ml of carbohydrate-free MRS supplemented with 4%

fructose, 4% glucose and 1% L-cysteine-HCl and incubated for another 16-18h

without shaking. Bacteria were spun down and resuspended in 1xPBS/sugar water

(1:1). The optical density (600nm) was adjusted according to the experimental

condition (OD=0.0001, 0.001, 0.01, or 0.1) and 5 µl of the final bacterial suspension

was fed to each newly emerged bee. Before feeding, the bees were starved for 2-3h.

After colonization, bees were given 1 ml of sterilized polyfloral pollen and sugar

water ad libitum. Bees were co-housed in groups of 20-40 bees. For the competition

experiment, each of the four honey bee strains was adjusted to an optical density

(600nm) of 0.001. The bumble bee strain ESL0228 was adjusted to an optical

density (600nm) of either 0.001, 0.01, or 0.1. Then equal volumes of the five strains

were mixed together and fed to newly emerged bees as described before. As

negative control, bees were fed with 5 µl of 1xPBS/sugar water. Dilutions of the

bacterial inocula were plated on MRS agar containing 2% fructose and 0.2% L-

cysteine-HCl and incubated as described before to determine how many CFUs

correspond to a given optical density (see Figure S5).

Ten bees per condition were dissected on day 5 after colonization. The hindgut was

separated from the midgut with a sterile scalpel and tweezers, and added to 1 ml or

500 ul of 1x PBS (depending on the experiment). The tissues were homogenized by

bead-beating as described before, dilutions plated on MRS agar containing 2%

fructose and 0.2% L-cysteine-HCl and the number of CFUs counted two to three

days after incubation. For the negative control, bacterial colonies were detected for

only one out of 30 bees with a relatively low abundance (10^3 CFUs per gut).

Moreover, the colonies looked different from the colonies of the Firm5 strains and

were identified as being *E. coli* and *Staphylococcus aureus* by 16S rRNA gene

sequencing.

The relative abundance of the five strains in the competition experiment was

analyzed using amplicon sequencing of a 199-bp fragment of a conserved

housekeeping gene (COG0266). To this end, a two-step PCR protocol was

established. In the first PCR, the 199-bp fragment of COG0266 was amplified from

crude cell lysates of gut homogenates with primers 1133 (5' -

CGTACGTAGACGGCCAGTATGCCNGAAATGCCRGARGTTGA – 3') and 1134 (5' -

GACTGACTGCCTATGACGACTAARCGATAYTTRCCYTCCATRCG) (3' – 95°C; 25x: 30'' -

95°C, 30'' – 64°C, 30'' – 72°C; 5' – 72°C). After removing primers with exonuclease

and shrimp alkaline phosphatase, barcoded Illumina adapters were added in the

second PCR. The resulting PCR products were pooled at equal volumes, gel purified

(MinElute Gel Extraction Kit, Qiagen) and loaded on an Illumina MiniSeq instrument

in mid-output mode. Reads were demultiplexed and filtered on quality using

trimmomatic (LEADING:28 TRAILING:28 SLIDINGWINDOW:4:15 MINLEN:90)

(Bolger *et al.* 2014). Then, each forward and reverse read pair was assembled using

PEAR (-m 290 -n 284 -j 4 -q 26 -v 10 -b 33) (Zhang *et al.* 2014). The resulting contigs

were assigned to the five strains based on base positions with discriminatory SNP

variants with the help of a custom-made Perl script.

Acknowledgments

We thank the School of Biology of the University of Lausanne for financial support of this project. We would also like to thank Ambrin Farizah Babu, Melvin Berard, Sarah Bergerm, Laurent Casini, Joaquim Claivaz, Yassine El Chazli, Jonas Garessus, Nastassia Gobet, Charlotte Griessen, Olivier Gustarini, Karim Hamidi, Dominique Jacques-Vuarambon, Titouan Laessle, Mirijam Mattei, Cyril Mattheiy-Doret., Jennifer Mayor, Sandrine Pinheiro, Claire Pralong, Virginie Ricci, Shalonie Sheppard, Tatiana Sokoloff, Anthony Sonrel, and Gaëlle Spack who participated as students in the SAGE class 2016/2017 and were involved in a preliminary analysis of the genomic data.

Data accessibility

Genome sequences and short read datasets are available under NCBI Bioproject accession PRJNA392822. Annotations of the Firm5 strains used for this study can be found in IMG/mer. Data analyses including custom scripts and intermediate output files are available under the following link: https://drive.switch.ch/index.php/s/kJiY2tjndjrBwJ5.

Figures

0.05

Figure 1. Core genome phylogeny. The tree was inferred using maximum likelihood on the concatenated protein alignments of 408 single-copy core genes (i.e. present in all Firm5 strains and in the outgroup strains). The collapsed outgroup consisted of 18 strains that were used to root the tree (see **Figure S4** for the complete tree). The two lineages of bumble bee strains and the four lineages of honey bee isolates are shown in green and blue color shades, respectively. Black and grey circles indicate bootstrap support values of 100 and ≥80, respectively, out of 100 replicates. The strain designation of each isolate is given and the species names of the typing strains indicated. The candidatus species name *Lactobacillus bohemicus* is depicted by hyphens. The length of the bar indicates 0.05 amino acid substitutions/site.

Figure 2. Colonization of microbiota-depleted honey bees with Firm5 strains from bumble bees and honey bees. (A) Mono-colonizations of microbiota-depleted honey bees (n=10 per treatment) with six bumble bee strains and four

honey bee strains. Each bee was inoculated with 5 µL of an optical density of 0.001.

Different letters indicate statistically significant differences between groups,

according to two-way ANOVA and Tukey HSD post hoc test (adjusted p-value <0.05).

The dashed red line indicates the detection threshold. Data points below the

detection limit show bees that had no detectable colonization levels. The percentage

of successfully colonized bees is shown. Horizontal lines indicate median. **(B)** Mono-

colonizations of microbiota-depleted honey bees with increasing inocula of the

bumble bee strain ESL0228. Colony forming units (CFUs) per gut were determined

at day 5 post colonization. The graph has the same layout as in panel A. (**C**)

Community profiles of microbiota-depleted bees colonized with a community

consisting of the bumble bee strain ESL0228 and four honey bee strains (ESL0183,

ESL0184, ESL0185, and ESL0186). Three different inoculation ratios of bumble bee

strain versus honey bee strains were used. The optical density of the bumble bee

strain in the inoculum is given in brackets. Due to the absence or the very low

abundance of ESL0228, the red fraction of the graph is not visible. Asterisks indicate

samples for which at least a few reads of strain ELS0228 were detected.

Figure 3. Pan genome analysis of Firm5 strains from honey bees and bumble bees and comparison to outgroup strains (i.e. closely related lactobacilli). (A) Venn diagram showing gene family distribution into the three major groups: Firm5 strains from bumble bees, Firm5 strains from honey bees, and outgroup strains. Numbers in bold indicate pan genome gene families (i.e. present in at least one genome of a given group). Numbers in regular font indicate core genome gene families (i.e. present in all genomes of a given group). **(B)** Number of Firm5-specific gene families exclusively present in strains of one sublineage. The fraction of core genome (present in every genome of a given sublineage) and pan genome (present in at least one genome of a given sublineage) gene families is indicated by grey and white color, respectively. Numbers above the graph indicate total number of

lineage-specific gene families for each host group. **(C)** Upper plot shows number of

gene families with COG annotation, and lower plot shows COG category distribution

of the annotated gene families for each subset of the Venn diagram in panel A. B,

specific to bumble bee strains; H, specific to honey bee strains; O, specific to

outgroup strains; BH, shared between honey bee and bumble bee strains; BO,

shared between bumble bee and outgroup strains; HO, shared between honey bee

and outgroup strains; BHO, shared between all three groups. **(D)** Same as in panel C,

but for the sublineage-specific gene families shown in panel B. Complete lists of all

gene families and their annotations can be found in **Datasets S1 and S2**. The

dominant COG category 'G' is shown in dark red and corresponds to 'Carbohydrate

transport and metabolism'. Other COG category abbreviations are given in **Dataset**

S2.

Figure 4. Distribution of carbohydrate-related gene families across strains of the Firm5 phylotype. (A) Total number of COG category 'G' gene families per genome per sublineage. **(B)** Total number of PTS (Phosphotransferase system) gene families per genome per sublineage. **(C)** Total number of glycoside hydrolase gene families per genome per sublineage. In all three panels, the genomes of the outgroup strains were included as a reference. **(D)** Venn diagram of glycoside hydrolase gene family distribution into the three major phylogenetic groups: Firm5 strains isolated

from bumble bees, Firm5 strains isolated from honey bees, and outgroup strains

belonging to related lactobacillus species. **(E)** Heatmap showing the distribution of

the identified glycoside hydrolase (GH) families across the analyzed genomes. The

dendrogram on the left shows a hierarchical clustering based on glycoside

hydrolase distribution. Strains are colored according to the major groups (green,

bumble bee strains; blue, honey bee strains; yellow, outgroup) and sublineage (color

tones). GH families specific to the Firm5 phylotype are indicated by grey boxes.

Figure 5. Genomic region encoding class II bacteriocins in Firm5 strains of bumble bee strains. Genomic regions encoding bacteriocin genes were identified and visualized with MultiGeneBlast v1.1.14 (Medema). Arrows present genes and same color indicates homology. A black line indicates the lactococcin 972 locus (*lcl*) and vertical grey blocks connect the homologous genes in other strains. An enlarged version of the three genes of the *lcl* locus with annotation is shown in the lower right. Grey shading over strain names indicates two sublineages of bumble bee strains; the four honey bees strains are representatives of the four sublineages. Other genomic regions encoding bacteriocins genes are given in Figure S8.

Supplementary Figures

Figure S1. Whole genome alignments of divergent Firm5 strains. ESL0247 and ESL0233 are bumble bee strains. ESL0183-186 are honey bee strains. Vertical grey lines indicate blocks of nucleotide sequence similarity. Different color intensities correspond to different degree of similarity based on BLASTN hits with a bit score of at least 100. Genes in color correspond to Firm5-specific genes relative to the outgroup (blue, genes specific to honey bee strains; red, genes specific to bumble bee strains; green, genes shared between honey bee and bumble bee strains). Other genes are shown in grey.

Figure S2. Read coverage and GC skew of the final genome assemblies.

Assembly positions are shown on the x-axis for each sequenced strain. y-Axis shows

Illumina read coverage for a sliding window of 100 bp. Red dashed lines indicate

contig breaks. Contigs were ordered according to the fully sequenced reference

strains ESL0183. Small contigs were left at the end of the assembly. Inset shows the

circular form of the assembly with the GC skew indicated. We found a higher read

coverage at the origin of replication, which is characteristic for replicating bacteria.

Moreover, our assemblies showed the typical GC skew of bacterial genomes. Both

characteristics indicate that the contigs of the assemblies were correctly assembled

and ordered. Notably, regions of extremely high coverage correspond to prophages

that apparently were amplified during culturing of some of the strains.

Figure S3. Average nucleotide identity between all analyzed Firm5 genomes.
Intensity of heatmap indicates pairwise ANI. White areas correspond to genomes, which were too divergent for ANI calculation. The names of each strain included in the analysis are given next to the plot area (see also **Table S1**). Grey shading indicates the six different sublineages of Firm5. ANI values are given in **Table S3**.

Figure S4. Complete core genome phylogeny. The tree was inferred using maximum likelihood on the concatenated protein alignments of 408 single-copy core gene families (i.e. present in all Firm5 strains and the outgroup strains). The two lineages of bumble bee strains and the four lineages of honey bee strains are shown in green and blue color shades, respectively. As outgroup, 15 representative strains of the *L. delbrueckii* group (to which Firm5 belongs to) were included in the analysis (shown in yellow) based on a previously published phylogeny of the entire genus *Lactobacillus* (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4579461/). In addition, we included three more distantly related strains to root the tree. Noteworthy, these three distantly related lactobacilli were excluded for all subsequent comparative analysis of the Firm5 strains. Filled circles indicate 100 bootstrap support values. The strain designation of each isolate is given. The length of the bar indicates 0.05 amino acid substitutions/sites.

Figure S5. Number of bacterial cell in the inocula used to colonize microbiota-

depleted honey bees with Firm5 strains. (A) CFUs in the inocula used for the

monocolonization experiments with individual strains. CFUs are given per 5μl, as

each bee was inoculated with 5 μl of an OD600 of 0.0001. Despite the adjustment to

the same OD600, the amount of live bacteria in each inoculum varied across strains.

The inoculum of strain ESL0237 could not be assessed due to a handling mistake

during dilution plating. **(B)** CFUs in the incocula used for the colonization

experiment with the bumble bee strain ESL0228 (left part) and for the colonization

experiment with the five-member community consisting of the bumble bee strain

ES0228 (left panel, OD=0.001, 0.01, and 0.1) and the four different honey bee strains

(right panel).

Figure S6. Percentage of gene families annotated as COG category 'G' per genome per sublineage. Same data as in Figure 4A, but expressed in relative numbers (percentage of all gene families per genome).

Figure S7. BlastP hit distribution of glycoside hydrolase (GH) gene families

specific to Firm5. A representative protein sequence of each gene family was

blasted against the NCBI nr database (NCBI Resource Coordinators 2018). The

distribution of the taxonomic classification of the first 50 Blast hits is shown at the

family level. The family of lactobacilliales is shown in red with black outlines. For

each gene family, the gene family identifier and the glycoside hydrolase enzyme

family (GHxx) are given.

Figure S8. Additional genomic regions encoding class II bacteriocins in Firm5

strains of bumble bee strains. Genomic regions encoding bacteriocin genes were

identified and visualized with MultiGeneBlast v1.1.14 (Medema). Arrows represent

genes, and same color indicates homology. A black line indicates the lactococcin 972

locus (*lcl*) and vertical grey blocks connect the homologous genes in other strains.

An enlarged version of the three genes of the *lcl* locus with annotation is shown in

the lower right of panel A. Grey shading over strain names indicates two sublineages

of bumble bee strains; the four honey bees strains are representatives of the four

sublineages.

Figure S9. Genomic regions encoding helviticin-J, a class III bacteriocin.

Genomic regions encoding bacteriocin genes were identified and visualized with

MultiGeneBlast v1.1.14 (Medema). Arrows represent genes, and same color

indicates homology. An arrow points at the helveticin-J gene homolog and vertical

grey blocks connect the homologous genes in other strains. Strains with the two

different types of grey shadings indicate strains from bumble bees and honey bees.

(A) Genomic region encoding helveticin-J in honey bee strains, and (B) genomic

region encoding helveticin-J in bumble bee strains.

Supplementary Tables and Datasets

Table S1. Strain list and genome features.

Table S2. Pairwise 16S rRNA gene sequence identities.

Table S3. ANI values.

Dataset S1. List of all pan genome gene families and their distribution according the three major groups: honey bee strains, bumble bee strains, outgroup strains.

Dataset S2. List of sublineage-specific gene families and COG category abbreviations.

Dataset S3. List of genes per genome with hits to the Carbohydrate-active enzyme (CAZY) database.

References

Bankevich A, Nurk S, Antipov D *et al.* (2012) SPAdes: a new genome assembly

algorithm and its applications to single-cell sequencing. *Journal of computational*

biology : a journal of computational molecular cell biology, **19**, 455–477.

Bobay L-M, Ochman H (2017) The Evolution of Bacterial Genome Architecture.

Frontiers in genetics, **8**, 829.

Bolger AM, Lohse M, Usadel B (2014) *Trimmomatic: a flexible trimmer for*.

NCBI Resource Coordinators (2018) Database resources of the National Center for

Biotechnology Information. *Nucleic Acids Research*, **46**, D8–D13.

Corby-Harris V, Maes P, Anderson KE (2014) The bacterial communities associated

with honey bee (*Apis mellifera*) foragers. *PloS one*, **9**, e95056.

Cotter PD, Ross RP, Hill C (2013) Bacteriocins - a viable alternative to antibiotics?

Nature reviews. Microbiology, **11**, 95–105.

Cox-Foster DL, Conlan S, Holmes EC *et al.* (2007) A metagenomic survey of microbes

in honey bee colony collapse disorder., **318**, 283–287.

Duar RM, Frese SA, Lin XB *et al.* (2017) Experimental Evaluation of Host Adaptation

of Lactobacillus reuteri to Different Vertebrate Species. *Applied and*

environmental microbiology, **83**, e00132–17.

Eddy SR (2009) A new generation of homology search tools based on probabilistic

inference. *Genome informatics. International Conference on Genome Informatics*,

23, 205–211.

Ellegaard KM, Tamarit D, Javelind E *et al.* (2015) Extensive intra-phylotype diversity

in lactobacilli and bifidobacteria from the honeybee gut. *BMC genomics*, **16**, 284.

Emery O, Schmidt K, Engel P (2017) Immune system stimulation by the gut

symbiont *Frischella perrara* in the honey bee (*Apis mellifera*). *Molecular Ecology*,

50, 735.

Engel P, Martinson VG, Moran NA (2012) Functional diversity within the simple gut

microbiota of the honey bee. *Proceedings of the National Academy of Sciences of*

the United States of America, **109**, 11002–11007.

Eren AM, Sogin ML, Morrison HG *et al.* (2015) A single genus in the gut microbiome

reflects host preference and specificity. *The ISME journal*, **9**, 90–100.

Frese SA, Benson AK, Tannock GW *et al.* (2011) The Evolution of Host Specialization

in the Vertebrate Gut Symbiont Lactobacillus reuteri (DS Guttman, Ed,). *PLoS*

genetics, **7**, e1001314.

Guy L, Kultima JR, Andersson SGE (2010) genoPlotR: comparative gene and genome

visualization in R. *Bioinformatics*, **26**, 2334–2335.

Hutchinson GE (1957) Concluding remarks. Cold Spring Harbor Symposia on

Quantitative Biology, *22: 415–427*.

Katoh K, Rozewicki J, Yamada KD (2017) MAFFT online service: multiple sequence

alignment, interactive sequence choice and visualization. *Briefings in*

bioinformatics, **30**, 3059.

Kešnerová L, Mars RAT, Ellegaard KM *et al.* (2017) Disentangling metabolic

functions of bacteria in the honey bee gut. *PLoS biology*, **15**, e2003467.

Koch H, Abrol DP, Li J, Schmid-Hempel P (2013) Diversity and evolutionary patterns

of bacterial gut associates of corbiculate bees. *Molecular Ecology*, **22**, 2028–

2044.

Kommineni S, Bretl DJ, Lam V *et al.* (2015) Bacteriocin production augments niche

competition by enterococci in the mammalian gastrointestinal tract. *Nature*,

526, 719–722.

Kostic AD, Howitt MR, Garrett WS (2013) Exploring host-microbiota interactions in

animal models and humans. *Genes & development*, **27**, 701–718.

Kwong WK, Moran NA (2015) Evolution of host specialization in gut microbes: the

bee gut as a model. *Gut microbes*, **6**, 214–220.

Kwong WK, Moran NA (2016) Gut microbial communities of social bees. *Nature*

reviews. Microbiology, **14**, 374–384.

Kwong WK, Engel P, Koch H, Moran NA (2014) Genomics and host specialization of

honey bee and bumble bee gut symbionts. *Proceedings of the National Academy*

of Sciences of the United States of America, **111**, 11509–11514.

Kwong WK, Medina LA, Koch H *et al.* (2017) Dynamic microbiome evolution in

social bees. *Science Advances*, **3**, e1600513.

Lee I, Kim YO, Park S-C, Chun J (2016) OrthoANI: An improved algorithm and

software for calculating average nucleotide identity. *International journal of*

systematic and evolutionary microbiology, **66**, 1100–1103.

Letzel A-C, Pidot SJ, Hertweck C (2014) Genome mining for ribosomally synthesized

and post-translationally modified peptides (RiPPs) in anaerobic bacteria. *BMC*

genomics, **15**, 983.

Ley RE, Hamady M, Lozupone C *et al.* (2008) Evolution of mammals and their gut

microbes. *Science*, **320**, 1647–1651.

Li L, Stoeckert CJ, Roos DS (2003) OrthoMCL: identification of ortholog groups for

eukaryotic genomes. *Genome research*, **13**,2178-89.

Ludvigsen J, Porcellato D, L'Abée-Lund TM, Amdam GV, Rudi K (2017)

Geographically widespread honeybee-gut symbiont subgroups show locally

distinct antibiotic-resistant patterns. *Molecular Ecology*, **26**, 6590–6607.

Macarthur R, Levins R (2015) The Limiting Similarity, Convergence, and Divergence

of Coexisting Species. *The American Naturalist*, **101**, 377–385.

Markowitz VM, Chen I-MA, Chu K *et al.* (2014) IMG/M 4 version of the integrated

metagenome comparative analysis system. *Nucleic Acids Research*, **42**, D568–73.

Martinson VG, Moy J, Moran NA (2012) Establishment of characteristic gut bacteria

during development of the honeybee worker. *Applied and environmental*

microbiology, **78**, 2830–2840.

Martínez B, Rodríguez A, Suárez JE (2000) Lactococcin 972, a bacteriocin that

inhibits septum formation in lactococci. *Microbiology*, **146**, 949–955.

McCutcheon JP, Moran NA (2012) Extreme genome reduction in symbiotic bacteria.

Nature reviews. Microbiology, **10**, 13–26.

McFall-Ngai M, Hadfield MG, Bosch TCG *et al.* (2013) Animals in a bacterial world, a

new imperative for the life sciences. *Proceedings of the National Academy of*

Sciences, **110**, 3229–3236.

Medema MH, Takano E, Breitling R (2013) Detecting Sequence Homology at the

Gene Cluster Level with MultiGeneBlast. *Molecular Biolology Evolution*, **30**,

1218-23.

Moeller AH, Caro-Quintero A, Mjungu D *et al.* (2016) Cospeciation of gut microbiota

with hominids. *Science*, **353**, 380–382.

Moran NA, Hansen AK, Powell JE, Sabree ZL (2012) Distinctive gut microbiota of

honey bees assessed using deep sampling from individual worker bees. *PloS one*,

7, e36393.

Ochman H, Worobey M, Kuo C-H *et al.* (2010) Evolutionary Relationships of Wild

Hominids Recapitulated by Gut Microbial Communities. *PLoS biology*, **8**,

e1000546.

Oh PL, Benson AK, Peterson DA *et al.* (2010) Diversification of the gut symbiont

Lactobacillus reuteri as a result of host-driven evolution. *The ISME journal*, **4**,

377–387.

Olofsson TC, Alsterfjord M, Nilson B, Butler E, Vásquez A (2014) Lactobacillus

apinorum sp. nov., Lactobacillus mellifer sp. nov., Lactobacillus mellis sp. nov.,

Lactobacillus melliventris sp. nov., Lactobacillus kimbladii sp. nov., Lactobacillus

helsingborgensis sp. nov. and Lactobacillus kullabergensis sp. nov., isolated from

the honey stomach of the honeybee Apis mellifera. *International journal of*

systematic and evolutionary microbiology, **64**, 3109–3119.

Praet J, Meeus I, Cnockaert M *et al.* (2015) Novel lactic acid bacteria isolated from

the bumble bee gut: Convivina intestini gen. nov., sp. nov., Lactobacillus

bombicola sp. nov., and Weissella bombi sp. nov. *Antonie van Leeuwenhoek*, **107**,

1337–1349.

Rissman AI, Mau B, Biehl BS *et al.* (2009) Reordering contigs of draft genomes using

the Mauve aligner. *Bioinformatics*, **25**, 2071–2073.

Seedorf H, Griffin NW, Ridaura VK *et al.* (2014) Bacteria from diverse habitats

colonize and compete in the mouse gut. *Cell*, **159**, 253–266.

Sriswasdi S, Yang C-C, Iwasaki W (2017) Generalist species drive microbial

dispersion and evolution. *Nature communications*, **8**, 1162.

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics*, **30**, 1312–1313.

Steele MI, Kwong WK, Whiteley M, Moran NA (2017) Diversification of Type VI

Secretion System Toxins Reveals Ancient Antagonism among Bee Gut Microbes.

mBio, **8**, e01630–17.

Toft C, Andersson SGE (2010) Evolutionary microbial genomics: insights into

bacterial host adaptation. *Nature Reviews Genetics*, **11**, 465–475.

van Heel AJ, de Jong A, Montalban-Lopez M (2013)*.* BAGEL3: automated

identification of genes encoding bacteriocins and (non-)bactericidal

posttranslationally modified peptides. *Nucleic Acid Research*, **41**, W448-53.

Yin Y, Mao X, Yang J *et al.* dbCAN: a web resource for automated carbohydrate-active

enzyme annotation. *Nucleic Acid Research*, **40**, W445-51.

Zhang J, Kobert K, Flouri T, Stamatakis A (2014) PEAR: a fast and accurate Illumina

Paired-End reAd mergeR. *Bioinformatics*, **30**, 614–620.

Zheng H, Nishida A, Kwong WK *et al.* (2016) Metabolism of Toxic Sugars by Strains

of the Bee Gut Symbiont Gilliamella apicola. *mBio*, **7**, e01326–16.

Zheng J, Ruan L, Sun M, Gänzle M (2015) A Genomic View of Lactobacilli and

Pediococci Demonstrates that Phylogeny Matches Ecology and Physiology.

Applied and environmental microbiology, **81**, 7233–7243.

1047