Computational tools to detect signatures of mutational processes in DNA from tumours: a review and empirical comparison of performance

Omichessan Hanane[1,2], Severi Gianluca[1,2,3] ,Perduca Vittorio[4,1]

[1]CESP (UMR INSERM 1018), Université Paris-Saclay, UPSud, UVSQ, Villejuif Cedex, F-94805, France

[2]Gustave Roussy, Villejuif, F-94805, France

[3]Cancer Epidemiology Centre, Cancer Council Victoria, and Centre for Epidemiology and Biostatistics, Melbourne School for Population and Global Health, The University of Melbourne, Australia

[4]Laboratoire de Mathématiques Appliquées – MAP5 (UMR CNRS 8145), Université Paris Descartes, 75006, Paris, France

* Coressponding author
E-mail : vittorio.perduca@parisdescartes.fr (VP)

Contributorship

Short title: Review and empirical comparison of mutational signatures tools

Abstract

Mutational signatures refer to patterns in the occurrence of somatic mutations that reflect

underlying mutational processes. To date, after the analysis of tens of thousands of exomes and

genomes from about 40 different cancers types, tens of mutational signatures characterized by a

unique probability profile across the 96 trinucleotide-based mutation types have been identified,

validated and catalogued. At the same time, several concurrent methods have been developed for

either the quantification of the contribution of catalogued signatures in a given cancer sequence

or the identification of new signatures from a sample of cancer sequences. A review of existing

computational tools has been recently published to guide researchers and practitioners through

their mutational signature analyses, but other tools have been introduced since its publication and,

a systematic evaluation and comparison of the performance of such tools is still lacking. In order

to fill this gap, we have carried out an empirical evaluation of the main packages available to

date, using both real and simulated data.

Keywords: *sequencing; cancer; mutational signatures; decomposition algorithms; probabilistic*

models; simulation study.

Introduction

It is well-known that cancer originates from a DNA modification in a single normal cell that is then propagated through cell divisions and that accumulates with further DNA modifications finally leading to abnormal, cancerous cells [1]. Such changes that include Single Nucleotide Variants (SNVs), insertions or deletions, Copy Number Variation (CNVs) and chromosomal aberrations, have been termed as "somatic mutations" to distinguish them with those inherited and transmitted from parents (germline mutations). With the development and the improvement of sequencing technologies collectively referred to as *High-Throughput Sequencing (HTS)* and the availability of cancer exome and genome data from most human cancers, much has been learnt about these mutations and the set of genes (oncogenes and tumour suppressor genes) that operate in human cancers. Beyond the relatively small number of "driver" mutations, those that confer a selective advantage for tumour development and progression, the vast majority of mutations, so-called "passenger" mutations, are "neutral" that means they do not affect cancer cells' survival. Together, passenger and driver mutations constitute a record of all cumulative DNA damage and repair activities occurred during the cellular lineage of the cancer cell [2].

Among all types of somatic mutations, a particular focus has been placed on single base substitutions that have been classified in six types according to the mutated pyrimidine base (C or T) in a strand-symmetric model of mutation. Such 6 substitutions (C>A, C>G, C>T, T>A, T>C and T>G) may be further classified in different types when considering the sequence pattern in which they are located (sequence context). For practical reasons, the sequence context is typically defined using the 5' and 3' bases proximal to the mutated base, that results in substitutions being

classified in 96 types (6 ∗ *4* ∗ *4*). In this paper we adopt this convention and use "mutations" to

refer to these 96 types of contextualized substitutions.

It has been hypothesised that specific patterns of mutations are the results of the action of

particular underlying mutational process. To identify such patterns, computational models, such

as matrix decomposition algorithms or probabilistic models, have been developed. The first of

such methods was published in 2013 by Alexandrov and colleagues [2,3], and, as for most of all

the other models that followed, is based on the idea that a mutational signature can be seen as a

probability distribution of the 96 types of mutations. Mutational signatures contribute to the total

mutational burden of a cancer genome, the so called mutational "catalogue" or "spectrum".

Such original methodological framework, applied to tens of thousands of genomes and exomes

from 40 different cancers types from large data repositories such as TCGA (The Cancer Genome

Atlas), has led to the identification of 30 mutational signatures characterized by a unique

probability profile across the 96 mutation types. These validated mutational signatures are listed

in a repertory on the COSMIC (Catalogue Of Somatic Mutations In Cancer) website [3] and have

been widely used as references. More recently, Alexandrov et al. have introduced an updated set

of signatures identified from an even larger collection of both exome and whole-genome

sequences (including the sequences from the PCAWG project) using two different methods (a

new version of the original framework and a Bayesian alternative) [4]. The new repertory

includes 49 new mutational signatures based on single base substitutions as in COSMIC, and also

mutational signatures built in the context of other types of mutations such as double base

substitutions (11 signatures), clustered based substitutions (4 signatures) and small insertions and

deletions (17 signatures).

After the introduction of the original framework, several other mathematical methods and

computational tools have been proposed to detect mutational signatures and estimate their

contribution to a given catalogue. These methods can be grouped in two categories with different

goals. The first class of methods aims to discover novel signatures while the second class aims to

detect the known and validated mutational signatures in the mutational catalogue of a given

sample. The approaches used in the first class are referred to as "*de novo*" (or "signature

extraction") while those in the second class as "refitting" (or "signature fitting"). All methods

have been implemented in open source tools, mainly R packages, but some of them are available

through command line, the Galaxy project or a web interface.

Signatures identified with de novo methods can be compared to reference signatures (for instance

those listed in COSMIC) through measures such as cosine [5] or bootstrapped cosine similarity

[6], which is a distance metric between two non-zero vectors. In this step of the analysis,

extracted signatures are matched to the most similar reference signature, provided that their

similarity is greater than a fixed threshold.

Recently, Baez-Ortega and Gori published a paper that discussed the mathematical models and

computational techniques for mutational signature analysis [7] . However, to our knowledge no

systematic evaluation of the performance of the existing methods has been conducted and the

issue of the choice of an appropriate cosine similarity threshold when matching a newly extracted

signature to the most similar counterpart in a reference set has not been addressed yet.

The present work has been designed to use simulated and real-world cancer sequences in order to

evaluate and compare for the first time the methods available to date, and assess their ability to

accurately detect the underlying mutational signatures in mutational catalogues.

Mathematical definition of mutational catalogues and signatures

In this section we briefly review the mathematical framework and notations based on the original model developed by Alexandrov and colleagues [2]. The data structure that is used as input by most current methods is the *mutational catalogue*, or *spectrum*, of a genome (or exome), with the counts for each of the 96 different mutation *features*, or *types*. In most approaches, the notion of mutation type refers to base substitutions within the trinucleotide context described above.

The mutational catalogue of a genome (or exome) g is defined as the vector $(m_g^1, \dots, m_g^K)^T$, where each m_g^k is the number of mutations of type k found in the genome and K, the number of possible mutation types, is equal to 96. Note that in this setting, information about mutation locations in the sequence is lost. The catalogue is built by comparing the sequence to a reference sequence in order to detect mutations and then by simply counting the occurrences of each type. The reference sequence can either be a standard reference (e.g. the assembly GRCh38) or a sequence from a "normal" tissue from the same individual (e.g. DNA from blood or from normal tissue surrounding tumours when available). For the purposes of the present work we will use the generic term "samples" for both genomes and exomes as the concepts and models used may be applied to both.

The basic idea underlying all computational models proposed is that the mutational catalogue of a sample results from the combination of all the mutational processes operative during lifetime, and therefore it can be seen as the weighted superposition of simpler mutational *signatures*, each corresponding to a specific process. The weight is larger if the process has a larger role in the

final catalogue of mutations: for example, for exposures to mutagens that last longer and/or are

more intense.

Formally, the signature of a mutational process n is a vector $p_n = (p_n^1, \dots, p_n^K)^T$, where each p_n^k

represents the probability that the mutational process will induce a mutation of type k. In other

words, p_n^k is the expected relative frequency of type k mutations in genomes exposed to n. Note

that $\sum_{k=1}^K p_n^k = 1$ and $0 \le p_n^k \le 1$ for all k.

The intensity of the exposure to a mutational process n in a sample g is measured by the number

of mutations e_g^n in g that are due to n. For this reason, e_g^n is referred to as the "exposure" of g to

n. The expected number of mutations of type k due to the process n in sample g is therefore

$p_n^k e_g^n$. If sample g has been exposed to N mutational processes, then the total number of

mutations of type k is

$$m_g^k = \sum_{n=1}^N p_n^k e_g^n + \epsilon_g^k, \qquad (1)$$

where ϵ_g^k is an error term reflecting sampling variability and non-systematic errors in sequencing

or subsequent analysis.

Matrix notation is effectively used when dealing with several samples and signatures. In this

situation, the collection of G samples is represented by the $K \times G$ matrix, with catalogues in

columns:

$$M = \begin{pmatrix} m_1^1 & m_2^1 & \dots & m_G^1 \\ \vdots & \vdots & & \vdots \\ m_1^K & m_2^K & \dots & m_G^K \end{pmatrix},$$

the N signatures are represented by the $K \times N$ matrix

148
$$P = \begin{pmatrix} p_1^1 & p_2^1 & \cdots & p_N^1 \\ \vdots & \vdots & & \vdots \\ p_1^K & p_2^K & \cdots & p_N^K \end{pmatrix},$$

149 and the exposures by the $N \times G$ matrix

150
$$E = \begin{pmatrix} e_1^1 & e_2^1 & \cdots & e_G^1 \\ \vdots & \vdots & & \vdots \\ e_1^N & e_2^N & \cdots & e_G^N \end{pmatrix}.$$

151 Equation (1) then becomes

152
$$M \approx P \times E$$

153 where we omitted the error term.

de novo **extraction vs. refitting**

155 *de novo* signature extraction methods aim at estimating P and E given M. Non-negative matrix

156 factorization (NMF) is an appealing solution to this unsupervised learning problem, because, by

157 definition, all involved matrices are non-negative [8]. NMF identifies two matrices P and E that

158 minimize the distance between M and $P \times E$. In particular, NMF finds an approximated solution

159 to the non-convex optimization problem

160
$$\underset{P \geq 0, E \geq 0}{\arg\min} ||M - P \times E||_F^2, \quad (2)$$

where the Frobenius matrix norm of the error term is considered. NMF requires the number of

signatures N, an unknown parameter, to be predefined. An approach for selecting this parameter

consists in obtaining a factorization of M for several of its values and then choosing the best N

with respect to some performance measure such as the reconstruction error or the overall

reproducibility. NMF is at the core of the Welcome Trust Sanger Institute (WTSI) Mutational

Signature Framework, the first published method for signature extraction [2]. An alternative to

numerical approaches based on NMF is given by statistical modelling and algorithms. With these

latter approaches, the number of mutations of a given type can be modelled by a Poisson

distribution

$$170 \qquad m_g^k \sim \mathcal{P}\left(\sum_{n=1}^N p_n^k e_g^n\right)$$

where mutational processes are assumed to be mutually independent. In order to estimate E and

P, it has been proposed to consider E as latent data and P as a matrix of unknown parameters and

to apply an expectation-maximization algorithm [9] or use Bayesian approaches [10–12]. One

important advantage of statistical approaches is the availability of model selection techniques for

the choice of N. We refer to the recent review [7] for a comprehensive overview of the

mathematical and computational aspects of existing methods, including alternative techniques not

mentioned above.

With respect to de novo extraction of signatures, the refitting approaches consider that the

signatures P are known and the goal is to estimate E given M and P. Refitting can be done for

individual mutational catalogues (i.e. individual samples) and, from a linear algebra perspective,

can be seen as the problem of projecting a catalogue living in the K-dimensional vector space (the

space spanned by all mutation types) onto its subset of all linear combinations of the given

mutational signatures with non-negative coefficients (the cone spanned by the given signatures).

A current practice consists in first performing a de novo extraction of signatures followed by a

comparison of the newly identified signatures with the reference signatures (e.g. COSMIC

signatures) by means of a similarity score, typically cosine similarity ranging from 0 (completely

different) to 1 (identical) [6,13]. Finally, a "novel" signature is considered to reflect a specific

reference signature if the similarity is larger than a fixed cut-off. If similarity is observed with

more than one reference signature, the one with the largest value of similarity is chosen, S1 Fig.

This assignment step crucially depends on the choice of the cut-off h that has been so far

inconsistent in the literature with some studies using a value of 0.75 [14] whereas others 0.80

[15,16]. Another difficulty is that different signatures might have very close cosine similarity, as

it happens also between COSMIC signatures, S2 Fig, so that a unique assignment is not always

possible.

Overview of the available tools

An similar number of *de novo* and refitting methods exist and all of them are available as open

source tools, mainly as R packages, or web interfaces (Table 1). The typical input of the tools is a

file including the mutations counts but some tools derive the mutation counts from ad-hoc input

files that may include for each individual a list of mutated bases, their position within the genome

and the corresponding bases from a reference genome. The typical format of such input files is

MAF, VCF or less common formats such as MPF and MFVF. For biologists or those who are

not familiar with programming, a set of tools were also developed and provided with user-

friendly interfaces (see also Table 1). Some tools include additional features such as the

possibility to search for specific patterns of mutations (e.g. APOBEC-related mutations [17]) and

differentially expressed genes [5].

Software	Available platform/ model	Input files	Additional features
de novo approaches			
WTSI	MATLAB/ NMF		Original framework
EMu [9] https://github.com/andrej-fischer/EMu	Command line/EM algorithm	- Mutation counts file - With respect to other tools, the counts file is transposed (the rows correspond to the samples)	- Opportunity matrix - Selection of the optimal number of signatures
SomaticSignatures [17] https://bioconductor.org/packages/release/bioc/html/SomaticSignatures.html	R/NMF and PCA	Variant Call Format	- Group-wise comparisons - Genomic visualization - Hierarchical clustering
pmsignature [18] https://github.com/friend1ws/pmsignature	R/mixed-membership model	- Mutation Position Format -Mutation Feature Vector Format	- Reduction of complexity - Mutation types defined by one or two flanking bases - Selection of the optimal number of signatures - Transcriptional strand bias - Background signature
bayesNMF [11,19,20] https://github.com/jburos/bayesNMF https://software.broadinstitute.org/cancer/cga/msp	R/Bayesian NMF	Mutation counts file	- Selection of the optimal number of signatures

signeR [10] https://bioconductor.org/packages/release/bioc/html/signeR.html	R/Bayesian NMF	Variant Call Format	- Opportunity matrix - Selection of the optimal number of signatures - Group-wise comparison (differential analysis)
mutSignatures [21] https://cran.r-project.org/web/packages/mutSignatures/index.html	R/NMF	Mutation counts file	- R-based implementation of [2]
Maftools [5] https://bioconductor.org/packages/release/bioc/html/maftools.html	R-Bioconductor /NMF	- Mutation Annotation - Format	- Genomic visualization - Cosine similarity - Selection of the optimal number of signatures - Group-wise comparisons (differential analysis) - APOBEC enrichment analysis
Helmsman [22] https://github.com/carjed/helmsman	Python/ NMF and PCA	- Variant Call Format - Mutation Annotation Format	- Able to run in parallel and designed for large datasets - Connection to external packages (in R) - may generate mutational catalogues from sequence data
SigAnalyzer [23]	R/ Bayesian NMF	Mutation counts file	- Automatic selection of the optimal number of signatures - Sparse signature profiles and contributions
SigProfiler [2,23]	Matlab/ NMF	Mutation counts file	- Further development of the original framework - Two steps: 1) extraction of a minimal set of signatures, 2) estimation of their contributions to individual samples

SparseSignatues [24] https://bioconductor.org/packages/release/bioc/html/SparseSignatures.html	R/ NMF with Lasso-penalized cost function	Mutation counts file	- Integration of DNA replication error signature - Sparse signature matrix - Number of signatures estimated with cross-validations - Scalable to large datasets
Refitting approaches			
deconstructSigs [25] https://github.com/raerose01/deconstructSigs	R/linear regression	Mutation counts file	- Opportunity matrix
QPsig [26] https://f1000researchdata.s3.amazonaws.com/supplementary/8918/0d25c07c-16ba-4b14-91e7-71749dcbbdd5.pdf	R/quadratic programming	Mutation counts file	
SignatureEstimation [27] https://www.ncbi.nlm.nih.gov/CBBresearch/Przytycka/index.cgi\#signatureestimation	R/quadratic programming and simulated alienation	Mutation counts file	
MutationalPatterns [28] http://bioconductor.org/packages/release/bioc/html/MutationalPatterns.html	R/Non-Negative Least Squares	Mutation counts file	- Also de novo identification - Cosine similarity comparison - Strand bias analyses - Enrichment and depletion
YAPSA [29] http://bioconductor.org/packages/release/bioc/html/YAPSA.html	R/Linear Combination Decomposition	Mutation counts file	- Cut-off for normalized exposure - Enrichment and depletion
decompTumor2Sig [30] https://github.com/rmpiro/decompTumor2Sig	R/quadratic programming	- Variant Call Format - Mutation Position Format - Mutation Feature Vector Format	- Converts a set of "Alexandrov's signatures" [13] to "Shiraishi's signatures" [18] - Decomposes a mutational catalogue in "Shiraishi's signatures"

MutationalCone [See Supporting information]	R/cone projection	Mutation counts file	- Fast in comparison to others refitting tools
Sigfit [12] https://github.com/kgori/sigfit	R/ Bayesian NMF	Mutation counts file	- Provides a new model for combining de novo and refitting approaches - Possible application to indel or rearrangement count data - Also implements Emu [9] model and allows conversion to genome-or exome- relative signatures
Pipelines and web-interfaces			
Mutspec [31] https://toolshed.g2.bx.psu.edu/repository/view_repository?id=f5c1f75e9fb33f8e	Galaxy pipeline/NMF	Variant Call Format	- de novo identification - Includes MS analysis in mouse cancer
MutaGene [32] https://www.ncbi.nlm.nih.gov/research/mutagene/	Web-interface	TCGA and ICGC data	- Refitting and de novo identification - Clustering of samples according to mutational profiles - Identification of potential driver's mutations
mSignatureDB [6] http://tardis.cgu.edu.tw/msignaturedb/	Web-interface	- Variant Call Format - Mutation Annotation Format - TSV	- Refitting and de novo identification - Bootstrapped cosine similarity - Comparison with either hg19 or hg38
Mutalisk [33] http://mutalisk.org	Web-interface	Variant Call Format	- Refitting and de novo identification - Transcriptional strand bias - Localization of kaetegis - Histones modifications - Cosine similarity comparison

MuSiCa [34] http://bioinfo.ciberehd.org:3838/MuSiCa/?	Web-interface	- Variant Call Format - Mutation Annotation Format - TSV - Excel	- Refitting and de novo identification - Cosine similarity - Samples classification

de novo approaches

Most tools that have been developed to identify mutational signatures were based on

decomposition algorithms including NMF or a Bayesian version of NMF [10,11]. The original

method developed by Alexandrov et al. was based on NMF and was implemented in MATLAB

[2] and is available also as an R package developed independently [21]. An updated and

elaborated version named SigProfiler, was proposed recently for extracting a minimal set of

signatures and estimating their contribution to individual samples [4]. The latter article also

discusses an alternative method based on Bayesian NMF, called SignatureAnalyzer, that led to

the identification of 49 reference signatures. Another tool that utilises NMF is Maftools that is

one of the few *de novo* tools that allows systematic comparison with the 30 validated signatures

in COSMIC by computing cosine similarity and assigning the identified signatures to the

COSMIC one with the highest cosine similarity [5].

Other tools such as SomaticSignatures [17] or the recent Helmsman [22] allows the identification

of mutational signatures through Principal Component Analysis (PCA) in addition to NMF. For

the sake of our formal comparison of the tools performance, we have tested SomaticSignatures

only with the NMF implementation because with PCA the factors are orthogonal and the values

inside the matrix can potentially be null or negatives, which is biologically unrealistic. Developed

in the Python language, Helmsman allows the rapid and efficient analysis of mutational

signatures directly from large sequencing datasets with thousands of samples and millions of

variants.

A promising recent method called SparseSignatures [24] proposes an improvement of the

traditional NMF algorithm based on two innovations, namely the default incorporation of a

background signature due to replication errors during cellular division and the enforcement of

sparsity in identified signatures through a Lasso penalty. This latter feature allows the

identification of signatures with well-differentiated profiles, thus reducing the risk of overfitting.

In addition to decomposition methods, an approach based on the Expectation Maximization (EM)

algorithm has been proposed to infer the number of mutational processes operative in a

mutational catalogue and their individual signatures. This approach is implemented in the EMu

tool [9], where the underlying probabilistic model is based on the assumption that the input

samples are independent and the number of mutational signatures is estimated using the Bayesian

Information Criterion (BIC).

Another tool that uses a probabilistic model named mixed-membership model is pmsignature

[18]. This tool utilises a flexible approach that at the same time reduces the number of estimated

parameters and allows to modify key contextual parameters such as the number of flanking bases.

The latter feature makes it possible to consider more-fine grained signatures and could potentially

lead to new biological insights [23]. It is important to note that for the purpose of our comparison

with the other tools, we set to one the number of flanking bases, and therefore to 96 mutation

types.

EMu, SigneR and pmsignature (and the refitting tool deconstructSigs) have been designed to take

into account the distribution of triplets in a reference exome or genome for example from a

sequence of normal tissue in the same individual. This is done by "normalizing" the input

mutational catalogues with respect to the distribution of triplets in the reference exome or

genome using an "opportunity matrix".

Refitting with known mutational signatures

In addition to the identification of novel mutational signatures, we are often interested in

evaluating whether a signature observed in an individual tumour is one of an established set of

signatures (e.g. the COSMIC signatures). This task is performed by "refitting tools" that aim to

search for the "best" combination of established signatures that explains the observed mutational

catalogue by projecting the latter (i.e. mapping) into the multidimensional space of all non-

negative linear combinations of the N established signatures.

The deconstructSigs [25] tool searches for the best linear combination of the established

signatures through an iterative process based on multiple linear regression aimed at minimizing

the distance between the linear combination of the signatures and the mutational catalogue. All

the other tools minimise the distance through equivalent approaches based on quadratic

programming [26,27,30], non-negative least square [28] linear combination decomposition [29]

and simulated annealing [27].

A faster implementation

We propose an alternative implementation of the decomposition performed in [28,29] based on a

simple geometric framework. Finding the linear decomposition of the input catalogue M on a set

of given signatures minimizing the distance can be seen as the problem of projecting M on the

geometric cone whose edges are the reference signatures. We propose to solve this problem by

applying the very efficient R package called coneproj [35]. More details about our algorithm,

which we called MutationalCone, together with the R code implementing it, can be found in the

Supporting information.

Combining de novo and refitting procedures

Sigfit [12] is a recently introduced R package for Bayesian inference based on two alternative

probabilistic models. The first of such models is a statistical formulation of classic NMF where

signatures are the parameters of independent multinomial distributions and catalogues are

sampled according to a mixture of such distributions with weights given by the exposures, while

the second model is a Bayesian version of the EMu model. An interesting innovation of Sigfit is

that it allows the fitting of given signatures and the extraction of undefined signatures in the same

Bayesian process. As argued by the authors, this unique feature might be helpful in cases where

the small sample of catalogues makes it difficult to try to identify new signatures or when the aim

is to study the heterogeneity between the primary tumour and metastasis in terms of the

signatures they show.

In this work we empirically evaluate the methods that have been already presented in a peer

review published paper and for which an implementation in R is available. To this aim, we adopt

the COSMIC set as reference for the analysis of simulated and real mutational catalogues because

we evaluate tools that were developed at the time when COSMIC was the only available database

of reference.

Data and experimental settings

Real cancers data from TCGA

In order to evaluate the performance of the available algorithms on real data, we used the exome sequences in The Cancer Genome Atlas (TCGA) repository (https://cancergenome.nih.gov/) for four cancer types: breast cancer, lung adenocarcinoma, B-cell lymphoma, and melanoma.

Mutation Annotation Format (MAF) files with the whole-exome somatic mutation datasets from these cohorts were downloaded from the portal gdc.cancer.gov on 6 March 2018. Data were annotated with MuSE [36] and the latest human reference genome (GRCh38). Mutational catalogues from these cohorts were obtained by counting the number of different mutation types using maftools [5]. The distribution of the number of mutations for each sample and separately for each cancer type is depicted in Fig 1.

Fig 1. Barplot with the number of mutations in each sample in four TCGA cohorts. Each bar represents a sample, with the number of mutations shown in the y-axis.

According to the COSMIC website, 13 and 7 signatures have been found for breast cancer and lung adenocarcinoma respectively, 6 for B-cell lymphomas and 5 for melanoma.

Simulated data

The first key assumption of our model of simulated mutational catalogues is that the number of mutations in a sample g that are induced by process n follows a zero-inflated Poisson (ZIP) distribution. According to this two-component mixture model, e_g^n is either 0 with probability π or is sampled according to a Poisson distribution $\mathcal{P}(\lambda)$ with probability $1 - \pi$. Such a model

depends on two parameters: the expectation λ of the Poisson component, and the probability π of

extra zeros. The ZIP model allows for frequent zeroes and is therefore more suitable for

modelling a heterogeneous situation where some samples are not exposed to a given mutational

process ($e_g^n = 0$) while some others are ($e_g^n > 0$). Realistically, the mutation counts due to

process n in each of the G samples, $e_1^n, \dots, e_G^n$, are assumed to be independent and identically

distributed according to a ZIP model where the expectation of the Poisson component is specific

to n:

$$e_g^n \sim \text{ZIP}(\lambda_n, \pi), \text{for all } n = 1, \dots, N.$$

Note that the expected number of mutations in sample g due to process n is $(1 - \pi)\lambda_n$. This

flexibility given by process-specific average counts is the second important characteristics of our

model and reflects the possibility that the mutagenic actions of different processes are

intrinsically different with respect to their intensity. Obviously, it would have been possible to do

one step further and allow for parameters $\lambda_{n,g}$ specific to both processes and samples, thus

representing the realistic situation in which the exposures of different samples to the same

process have different duration (e.g. smokers/non-smokers). However, this would have resulted

in too many parameters to tune, thus making it difficult to interpret the results of our simulation

study. For the same reason we considered one fixed value of π.

The parameter λ_n depends on both the average total number r of mutations in a sample and the

relative contribution of n. We therefore imposed the parameterization $\lambda_n(1 - \pi) = q_n r$, where

q_n is the average proportion of mutations due to the process n.

When taking a unique value of r, this model produces realistic simulations even though it

underrepresents extreme catalogues with very large or small total numbers of mutations (see S3

Fig (c)). While considering a specific value of r for each sample, or group of samples, would

definitely make it possible to obtain a more realistic distribution of simulated catalogues (see S3

Fig (b)), such multidimensional parameter would complicate unnecessarily our empirical

assessment of mutational signatures detection methods by introducing too many specifications.

We therefore decided to consider a unique r for each set of simulations. This formulation allowed

us to study empirically the performance of a given signature detection method as a function of the

average number of mutations r while fixing the average proportion of mutations due to each

mutational process q_n, according to different profiles that mimic real cancer catalogues.

Interestingly, the ZIP model appeared to be more appropriate to represent mutational catalogues

than the pure Poisson model used in previous publications [9,10] (S3 Fig (d)).

Our simulation protocol was the following:

1.   We chosed N signatures from the COSMIC database, thus obtaining the matrix P.

2.   For each sample g and process n we sampled e_g^n from a ZIP distribution with parameters

$\lambda_n = q_n r / (1 - \pi)$ and π and obtained E. Here q_n, r and π are fixed parameters to set.

3.   We computed the product $P \times E$. In order to obtain the final simulated catalogue M, we

added some noise to the latter matrix by taking $m_g^k \sim \mathcal{P}\big((P \times E)_g^k\big)$.

We generated four alternative sets of simulated catalogues, referred to as Pofiles 1, 2, 3 and 4,

each set mimicking a particular cancer: breast cancer, lung adenocarcinoma, B-cell lymphoma

and melanoma. In order to do so, for each tumour type, we applied MutationalCone to the

corresponding TCGA datasets and we calculated the mean contribution across all samples of each

signature known to contribute to the specific cancer type q_n. Signatures with $q_n = 0$ do not

contibute to the final catalogue and were not in the matrix P S4 Fig depicts the resulting four sets

of configurations $(q_1, \ldots, q_N)$ used for the simulations. We fixed the frequency of structural null

contributions to the catalogues to $\pi = 0.6$ in all simulations. This value was chosen because it

leads to a limited number of hypermutated catalogues, as it is often encountered in practice. At

last, we varied the number of r from as little as 10 to as much as 100,000 mutations. This allowed

us to study the performance of methods on a large spectrum of catalogues: from a limited number

of mutations as in exomes, to a very large number, as in whole cancer genome sequences.

For each of the four tumour types and for each value of r, we simulated a catalogue matrix with

G samples. Fig 2 shows examples with $G = 30$ for three values of r.

Fig 2. Sample of simulated catalogues. Catalogues are simulated according to the model

described in the text. Tuning parameters are the average number of mutations, here $r = 10^2, 10^4$

and 10^5 and the cancer type-specific average contribution of each signature (also see S4 Fig).

Coloured bars indicate the contribution of each signature to the total mutational load of each

sequence.

Methods for measuring the performance of algorithms

369 All methods for identifying signatures find solutions to the minimization problem (2). A

370 straightforward way to measure the accuracy of the reconstructed catalogue is, therefore, to

371 calculate the Frobenius norm of the reconstruction error

372
$$||M - \hat{M}||_F^2 = \sum_{g=1}^{G} \sum_{n=1}^{N} (m_g^k - \hat{m}_g^k)^2,$$

where $\hat{M} = \hat{P} \times \hat{E}$ is the matrix of catalogues reconstructed from the estimated signature and

exposure matrices. Some of the algorithms involve stochastic steps such as resampling and/or

random draws of initial parameters. For these algorithms, one simple way to assess the robustness

of the estimates is to look at the variability of the reconstruction error when the same catalogues

are analyzed several times with the same algorithm.

In order to make decisions about whether an extracted signature is the same as validated

signatures (e.g. COSMIC signatures) a cut-off for cosine similarity needs to be defined. We

decided to apply six different cut-offs and consider as "new" all identified mutational signatures

for which the maximal cosine similarity is lower than the cut-off value.

Specificity and sensitivity for de novo extraction and assignment

In most applications, signature extraction is done in two steps: first, signatures are found using a

de novo extraction tool and then for the extracted signatures a cosine similarity with each of the

COSMIC signatures is calculated. In order to measure the performance of both these steps

combined, we used simulated catalogues and computed false and true positive rates and false and

true negative rates. In a simulated catalogue, the set of *true* signatures $p_1, \ldots, p_N$ that do

contribute to the catalogue are known, thus allowing the comparison of the latter to the estimated

signatures $\hat{p}_{i_1}, \ldots, \hat{p}_{i_N}$. Note that, for the sake of simplicity, we explicitly look for the same

number of signatures that we used to simulate the catalogues, not addressing questions about

model selection performance.

Estimated signatures that belong to the set of "true signatures" are considered as true positives,

while all "true signatures" that are not extracted count as false negatives. False positives are all

estimated signatures that do not have a match in the set of "true signatures". This can happen for

two reasons: the estimated signature is assigned to a COSMIC signature not used to build the

catalogue, or it is not sufficiently similar to any COSMIC signature. This last situation usually

takes place when setting a very high cosine similarity threshold h. In this case, signatures that

have maximal cosine similarity lower than the cutoff, will be termed as "new". At last, true

negatives are all COSMIC signatures not used for the simulation, nor estimated. From these four

measures, we compute specificity (number of true negatives divided by the total number of

negatives) and sensitivity (number of true positives divided by the total number of positives).

In our empirical study, for each simulation setting described in the Simulated data section (that is

for each profile given by a choice of proportions $(q_1, \dots, q_N)$ and for a choice of total number of

mutations r) we built 50 replicates, each made of a matrix of G samples. Then we extracted

signatures from all replicates with a given tool. Importantly, we set the expected number of

signatures to be the same as the number of signatures used in the simulation, thus not addressing

the problem of model selection. Then we compared the extracted signatures to the COSMIC

signatures using a cosine similarity threshold h. Finally, we computed specificity and sensitivity

and obtained Monte-Carlo estimates based on the means over all replicates.

Bias of refitting procedures

Because refitting algorithms assume that the matrix of signatures is known, a simple way to

assess their performance is to look at the bias of the exposure estimates e_g^n. In order to do so, we

simulated 50 replicates each consisting of one lymphoma-mimicking catalogue (Profile 2) with

an average number of mutations set to $r = 10^4$. Then, for each process n, we obtained Monte-

Carlo estimates of the bias $E[\hat{e}_g^n] - e_g^n$ by averaging the differences $\hat{e}_g^n - e_g^n$ over all replicates.

A global measure of performance that considers all exposure estimates is given by the mean

squared error (MSE), that is the expected value of the loss function $\sum_{n=1}^{30}(\hat{e}_g^n - e_g^n)^2$. We

obtained Monte-Carlo estimates of the mean squared error by averaging the loss function values

across all 50 replicates and calculate asymptotic confidence intervals.

Analyses were performed using R 3.5.1 on a Linux Ubuntu (14.04) machine with 32 GB RAM

and Intel Xeon E5-2630 v4 CPUs @ 2.2 GHz x 16.

Comparisons of algorithm performance

Performance of de novo tools

Fig 3 shows the distribution of the reconstruction error when a given computational tool is

applied several times to the same real trinucleotide matrix. Reconstruction errors show limited

variability due to stochastic steps in the algorithms and no variability whatsoever for EMu and

Maftools. All methods under evaluation are roughly equivalent in terms of their ability to

properly reconstruct the initial matrix of mutational catalogues. This is not surprising, given that

all methods are meant to solve the optimization problem in (2). The error value also appears to be

cancer-dependent. This is expected because cancers differ with regards to the total number of

mutations and the number of operating mutational signatures, making the decomposition more or

less difficult.

Fig 3. Reconstruction errors and their variability due to stochastic steps in the algorithms.

Each program under evaluation is applied 50 times on the same matrix of real catalogues shown

in Fig 1; boxplots represent the distribution of the squared Frobenius distance between the

original catalogue and its reconstruction. Boxplots look like flat segments because of the scale of

the y-axis.

We used realistic simulations to evaluate the performance of each method for de novo extraction

followed by a classification step in which the extracted signatures are assigned to the most similar

COSMIC signature.

Figs 4 and 5 show the specificity and sensitivity of such two-stage procedure as functions of the

number of samples G in each catalogue and the cosine similarity cut-off h, while Figs 6 and 7

show the specificity and sensitivity as functions of the number of mutations in each catalogue and

h.

Figs 4 and 5. Performance of extraction methods and mapping on COSMIC signatures as

the number of analysed catalogues and the cosine cut-off h vary. Specificity and sensitivity

are estimated from 50 replicates each made of G genomes. The average number of mutations in

each catalogue is $r = 10,000$. The model used to simulate realistic replicates according to the

four Profiles and the estimation methods are described in the section Data and experimental

settings.

Figs 6 and 7. Performance of extraction methods and mapping on COSMIC signatures as

the average number of mutations and the cosine cut-off h vary. Specificity and sensitivity are

estimated from 50 replicates each made of $G = 30$ catalogues. The model used to simulate

realistic replicates according to the four Profiles and the estimation methods are described in the

section Data and experimental settings.

We do not see very large differences in the tools' performance (specificity and sensitivity)

between profiles with respect to the number of samples. For Profiles 2, 3 and 4 there is almost no

gain in specificity and sensitivity when the sample size increases whereas for Profile 1, that is

characterised by small contributions from several signatures (S4 Fig), sensitivity is poor for small

samples sizes and increases with the sample size.

Sensitivity is poor when the average number of mutations is poor. Sensitivity increases with the

average number of mutations with a large variability according to the cancer profile and method.

In general, sensitivity is higher for profiles characterized by a signature with a stronger effect

than other contributing signatures (e.g. Profile 2 vs. Profile 4). Specificity is less variable and

generally high.

Specificity and sensitivity slightly deteriorate for higher cut-off values. This is expected because

by setting a higher cut-off, the number of found signatures that are not similar enough to

COSMIC signatures increases. Because these estimated signatures are considered as novel, they

are false positives (that is found signatures not used for simulations), leading to a greater number

of false positives and therefore to a lower specificity. Moreover, if the cut-off is too stringent, the

number of false negatives will be high because some signatures used for the simulations are

correctly found but do not score a high enough cosine similarity and therefore count as false

negatives. This will make the resulting sensitivity low.

Fig 8 shows the running time when tools are applied to real lung datasets with a varying number

of samples. While all methods show a fast-growing running time with increasing number of

samples, SomaticSignatures and maftools are much faster than the others for more than 100

samples, making it possible to analyse large number of samples in few seconds. For example, for

two hundred samples, the slowest method (SigneR), the running time is 913.72s while for the

fastest (maftools), the value is 5.97s.

Fig 8. Running times of refitting tools. Methods were applied to subsets of the TCGA Lung

cohort of different sizes.

Performance of refitting tools

Fig 9 shows the distribution of the differences between the estimated and true contribution of all

n signatures $e_g^1, \ldots, e_g^{30}$ for the different refitting methods under evaluation. Sample catalogues

were simulated mimicking Lung cancer profiles (Profile 3), with signatures 1,2,4,5,6, 13 and 17

actually contributing as shown in S4 Fig. All methods give almost identical results.

Fig 9. Bias of the estimates of each signature contribution for several refitting methods. For

each signature, the bias estimates are obtained by averaging the exposure estimates across 50

samples. Mean square errors, together with 95% confidence intervals, are reported on the top of

each plot. Simulations were done according to the model described in the Data and experimental

settings section.

By comparison with the true exposure profile given in S4 Fig, it is clear that all refitting methods

provide good estimates of the contributions of all but signatures 1,5 and, 6 and to a lesser extent

signature 4. Moreover, all methods correctly estimate a zero contribution for signatures 3 and 16

even though these are very similar to signature 5, S2 Fig.

Interestingly, we note that signatures 2 and 13 (both attributed to APOBEC activity) are well

identified by all methods. This finding is in line with previous claims about the stability of these

two signatures.

In terms of running time, deconstructSigs and SignatureEstimation based on simulated annealing

are more than two orders of magnitude slower than the other methods (Fig 10). All other

methods run in a fraction of second. As expected, the running time increases linearly with the

number of samples. MutationalCone, our custom implementation of the solution to the

optimization problem solved by YAPSA and MutationalPatterns outperforms all other methods.

The second fastest method is SignatureEstimation based on Quadratic Programing. As example,

for two hundred samples, the execution time of deconstrucSigs is 86.148s and for

MutationalCone is 0.028s.

Fig 10. Running times of refitting tools. Methods were applied to subsets of the TCGA Lung

cohort of different sizes.

Discussion

In this work we complement and expand a recent review of the available methods to identify mutational signatures [37] by empirically comparing their performance, using both real world TCGA data and simulated data. To our knowledge, this is the first formal comparison of both de novo discovery and refitting of mutational signatures. The research about mutational signatures is very active and in rapid development, with preprints about new methods regularly published on bioRxiv.org; in this work we thoroughly assessed and formally compared methods that have already been published in peer reviewed journals, and we briefly described other more recent methods. The results of this work can lead to a better understanding of the strengths and limitations of each method as well as to the identification of the key parameters influencing their performance, namely the number of mutations and the "complexity" of the contributing signatures.

Since the publication of the first work about mutational signatures in 2013 [2], multiple algorithms have been developed, leading to similar but not identical results, a source of concern for researchers interested in this type of analysis. Conceptually, this is not surprising: mutational signatures are naturally defined in terms of non-negative matrix factorization, a well-known ill-posed problem (a unique solution does not exist). Although this limitation has cast doubts on the biological validity of mutational signatures, this has been somehow validated using experimental and computational approaches by Zho et al. [27].

An original aspect of this paper is the empirical performance study and in particular the probabilistic model of mutational catalogues used to generate simulations. This model is based on

the zero-inflated Poisson distribution that allows for sparse contribution of signatures and thus

makes it possible to build mutation count data that are more realistic than the pure Poisson model

previously considered [12,13]. As argued in the Simulated data section, our model

underrepresents the few samples with extremely large total mutation counts. Because catalogues

of this type are more informative for mutational signatures, it is likely that our results slightly

underestimate the methods performance. However, our main objective was the comparison of the

different methods, and this is not affected by this systematic bias. As discussed, it would have

been possible to consider even more realistic models, however these would have led to results

that depend on too many parameters thus making the interpretation harder. For the opposite

reason, we did not simulate catalogues from real data using the bootstrap: this would have

produced almost real samples but without the possibility to evaluate the performance of methods

according to different parametric scenarios. Alternative models that were recently proposed are

based on the negative binomial distribution [24] and on the Dirichlet distributions for the

exposures and signatures and the multinomial distribution for the catalogues [12]. We suggest

that developers should assess their new methods on simulations based on realistic models such as

ours or the latter.

Another innovative aspect of our work is the assessment of the performance of de novo

approaches. This is in general a slippery problem given the non-supervised learning nature of

NMF and concurrent methods. We addressed it by building confusion matrices after comparing

the "true" signatures used for simulations with the newly identified signatures in terms of their

cosine similarity.

At last, we introduced MutationalCone, a new implementation of existing refitting tools that is

based on a very simple geometric model and proved to be faster than all other methods. The R

code implementing MutationalCone can be found in the Supporting information.

The results of our study can be helpful to guide researchers through the planning of mutational

signature analysis. In particular, we show that the performance of de novo methods depends on

the complexity of the analysed sequences, the number of mutations and to a lesser degree the

number of samples analysed. We showed that cancer profiles for which it is "easier" to detect

signatures are characterised by one predominant signature only (as in our profiles 2 and 3), while

cancers with more complex profiles, characterized by non-negligible contributions by several

signatures (profiles 1 and 4), are more difficult to identify. Indeed, recent evidence shows that the

majority of cancers harbour a large number of mutational signatures [23] and therefore belong to

the latter scenario.

With regards to the mutation number, we observe that with the number of mutations that could be

found in some cancer exomes the performance is generally poor (i.e. low specificity and

sensitivity). A clear example is illustrated in a recent comparison between a smaller set of

genomes (G=2,780) and a much larger sample of exomes (G=19,184), where it is shown that in

the former set a much larger number is signatures is detected than in the latter [23]. This problem

is likely to be mitigated if counts were normalized by the expected number of each type's

trinucleotides in the analysed region under healthy condition, that is if an opportunity matrix was

provided. We do not address this important aspect in our comparison study as only a few methods

can account for opportunity matrices.

We showed that when comparing identified signatures with COSMIC signatures, the choice of a

cosine similarity cut-off has a relatively small impact on the overall performance. If the aim is to

identify novel signatures it would be preferable to choose a lower value (0.75 or less). On the

contrary, if the aim is to access the presence of known signatures in mutational catalogues

(cancer genomes or exomes), we recommend turning to refitting methods.

Not all the de novo methods we evaluated offer the possibility to automatically choose the

number of signatures to be found. For instance, the popular SomaticSignatures only provides a

graphical visualisation of the residual sum of squares for several choices of the number of

signatures; the user can choose the optimal number by identifying the inflexion point. For this

reason, this crucial aspect was not addressed in our empirical assessment. Similarly, we only

considered mutation types defined by the trinucleotide motifs, as currently only pmsignature [18]

can consider more than one flanking basis on each side of the substitution.

For well-studied cancers, refitting approaches are a faster and more powerful alternative to de

novo methods, even with just one input sample. As the COSMIC database has been built and

validated by analysing tens of thousands of sequences of most cancer types, we recommend

borrowing strength from previous studies and using refitting tools when performing standard

analysis not aimed at the discovery of new signatures.

A more recent set of 49 signatures have been estimated from a new large dataset including

thousands of additional sequences. The article introducing this new set has not been published yet

and the 30 COSMIC signatures can still be considered as the reference. In the future it will be

interesting to evaluate the performance of the different tools using the new set of signatures.

It is important to note that our empirical comparison did not allow to identify a method that is

clearly superior to others; for this reason we recommend to systematically perform a sensitivity

analysis based on the application of one or more alternative methods based on different

algorithms, in order to assess the robustness of results.

Acknowledgements

609 We acknowledge the support of a grant from La Ligue contre le Cancer (Appel à projets 2016

610 'Recherche en Épidémiologie'). HO is supported by a PhD fellowship from the French Institut

611 National du Cancer (INCa_11330). VP would like to thank Grégory Nuel for the useful

612 discussions around zero-inflated models. The authors would like to thank an anonymous reviewer

613 for the very useful remarks and constructive suggestions.

References

1.   Stratton MR, Campbell PJ, Futreal PA. The cancer genome. Nature. 2009;458: 719–724.
doi:10.1038/nature07943
2.   Alexandrov LB, Nik-Zainal S, Wedge DC, Campbell PJ, Stratton MR. Deciphering
Signatures of Mutational Processes Operative in Human Cancer. Cell Reports. 2013;3: 246–259.
doi:10.1016/j.celrep.2012.12.008
3.   Forbes SA, Beare D, Boutselakis H, Bamford S, Bindal N, Tate J, et al. COSMIC:
somatic cancer genetics at high-resolution. Nucleic Acids Research. 2017;45: D777–D783.
doi:10.1093/nar/gkw1121
4.   Alexandrov LB, Kim J, Haradhvala NJ, Huang MN, Ng AW, Boot A, et al. The
Repertoire of Mutational Signatures in Human Cancer. bioRxiv. 2018; 322859.
doi:10.1101/322859
5.   Mayakonda A, Koeffler HP. Maftools: Efficient analysis, visualization and

summarization of MAF files from large-scale cohort based cancer studies. bioRxiv. 2016;
052662. doi:10.1101/052662

6.   Huang P-J, Chiu L-Y, Lee C-C, Yeh Y-M, Huang K-Y, Chiu C-H, et al. mSignatureDB: a
database for deciphering mutational signatures in human cancers. Nucleic Acids Research.
2018;46: D964–D970. doi:10.1093/nar/gkx1133

7.   Baez-Ortega A, Gori K. Computational approaches for discovery of mutational signatures
in cancer. Briefings in Bioinformatics. 2017; doi:10.1093/bib/bbx082

8.   Lee DD, Seung HS. Learning the parts of objects by non-negative matrix factorization.
Nature. 1999;401: 788.

9.   Fischer A, Illingworth CJ, Campbell PJ, Mustonen V. EMu: probabilistic inference of
mutational processes and their localization in the cancer genome. Genome Biology. 2013;14:
R39. doi:10.1186/gb-2013-14-4-r39

10.   Rosales RA, Drummond RD, Valieris R, Dias-Neto E, da Silva IT. signeR: an empirical
Bayesian approach to mutational signature discovery. Bioinformatics. 2017;33: 8–16.
doi:10.1093/bioinformatics/btw572

11.   Kim J, Mouw KW, Polak P, Braunstein LZ, Kamburov A, Tiao G, et al. Somatic ERCC2
mutations are associated with a distinct genomic signature in urothelial tumors. Nature Genetics.
2016;48: 600–606. doi:10.1038/ng.3557

12.   Gori K, Baez-Ortega A. sigfit: flexible Bayesian inference of mutational signatures.
bioRxiv. 2018; doi:10.1101/372896

13.   Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SAJR, Behjati S, Biankin AV, et al.
Signatures of mutational processes in human cancer. Nature. 2013;500: 415–421.
doi:10.1038/nature12477

14.   Letouzé E, Shinde J, Renault V, Couchy G, Blanc J-F, Tubacher E, et al. Mutational
signatures reveal the dynamic interplay of risk factors and cellular processes during liver
tumorigenesis. Nature Communications. 2017;8. doi:10.1038/s41467-017-01358-x

15.   Kim SY, Jung S-H, Kim MS, Han M-R, Park H-C, Jung ES, et al. Genomic profiles of a
hepatoblastoma from a patient with Beckwith-Wiedemann syndrome with uniparental disomy on
chromosome 11p15 and germline mutation of APC and PALB2. Oncotarget. 2017;8.
doi:10.18632/oncotarget.20515

16.   Han M-R, Shin S, Park H-C, Kim MS, Lee SH, Jung SH, et al. Mutational signatures and
chromosome alteration profiles of squamous cell carcinomas of the vulva. Experimental &
Molecular Medicine. 2018;50: e442. doi:10.1038/emm.2017.265

17.   Gehring JS, Fischer B, Lawrence M, Huber W. SomaticSignatures: inferring mutational
signatures from single-nucleotide variants: Fig. 1. Bioinformatics. 2015; btv408.
doi:10.1093/bioinformatics/btv408

18.   Shiraishi Y, Tremmel G, Miyano S, Stephens M. A Simple Model-Based Approach to
Inferring and Visualizing Cancer Mutation Signatures. Marchini J, editor. PLOS Genetics.
2015;11: e1005657. doi:10.1371/journal.pgen.1005657

19.   Kasar S, Kim J, Improgo R, Tiao G, Polak P, Haradhvala N, et al. Whole-genome
sequencing reveals activation-induced cytidine deaminase signatures during indolent chronic
lymphocytic leukaemia evolution. Nature Communications. 2015;6. doi:10.1038/ncomms9866

20.   Tan VYF, Fevotte C. Automatic Relevance Determination in Nonnegative Matrix
Factorization with the /spl beta/-Divergence. IEEE Transactions on Pattern Analysis and Machine
Intelligence. 2013;35: 1592–1605. doi:10.1109/TPAMI.2012.240

21.   Fantini D, Huang S, Asara JM, Bagchi S, Raychaudhuri P. Chromatin association of
XRCC5/6 in the absence of DNA damage depends on the XPE gene product DDB2. Tansey WP,

editor. Molecular Biology of the Cell. 2017;28: 192–200. doi:10.1091/mbc.E16-08-0573

22.   Carlson J, Li JZ, Zöllner S. Helmsman: fast and efficient mutation signature analysis for
massive sequencing datasets. BMC Genomics. 2018;19. doi:10.1186/s12864-018-5264-y

23.   Alexandrov L, Kim J, Haradhvala NJ, Huang MN, Ng AWT, Boot A, et al. The
Repertoire of Mutational Signatures in Human Cancer. 2018; doi:10.1101/322859

24.   Ramazzotti D, Lal A, Liu K, Tibshirani R, Sidow A. De Novo Mutational Signature
Discovery in Tumor Genomes using SparseSignatures. bioRxiv. 2018; doi:10.1101/384834

25.   Rosenthal R, McGranahan N, Herrero J, Taylor BS, Swanton C. deconstructSigs:
delineating mutational processes in single tumors distinguishes DNA repair deficiencies and
patterns of carcinoma evolution. Genome Biology. 2016;17. doi:10.1186/s13059-016-0893-4

26.   Lynch AG. Decomposition of mutational context signatures using quadratic programming
methods. F1000Research. 2016;5: 1253. doi:10.12688/f1000research.8918.1

27.   Huang X, Wojtowicz D, Przytycka TM. Detecting presence of mutational signatures in
cancer with confidence. Bioinformatics. 2018;34: 330–337. doi:10.1093/bioinformatics/btx604

28.   Blokzijl F, Janssen R, van Boxtel R, Cuppen E. MutationalPatterns: comprehensive
genome-wide analysis of mutational processes. Genome Medicine. 2018;10. doi:10.1186/s13073-
018-0539-0

29.   Huebschmann D, Gu Z, Schelsner M. YAPSA: Yet Another Package for Signature
Analysis R package version 1.6.0. R package version 1.6.0. 1 Jan 2015.

30.   Krüger S, Piro RM. Identification of mutational signatures active in individual tumors.
doi:10.7287/peerj.preprints.3257v1

31.   Ardin M, Cahais V, Castells X, Bouaoun L, Byrnes G, Herceg Z, et al. MutSpec: a
Galaxy toolbox for streamlined analyses of somatic mutation spectra in human and mouse cancer
genomes. BMC Bioinformatics. 2016;17. doi:10.1186/s12859-016-1011-z

32.   Goncearenco A, Rager SL, Li M, Sang Q-X, Rogozin IB, Panchenko AR. Exploring
background mutational processes to decipher cancer genetic heterogeneity. Nucleic Acids
Research. 2017;45: W514–W522. doi:10.1093/nar/gkx367

33.   Lee J, Lee AJ, Lee J-K, Park J, Kwon Y, Park S, et al. Mutalisk: a web-based somatic
MUTation AnaLyIS toolKit for genomic, transcriptional and epigenomic signatures. Nucleic
Acids Research. 2018; doi:10.1093/nar/gky406

34.   Díaz-Gay M, Vila-Casadesús M, Franch-Expósito S, Hernández-Illán E, Lozano JJ,
Castellví-Bel S. Mutational Signatures in Cancer (MuSiCa): a web application to implement
mutational signatures analysis in cancer samples. BMC Bioinformatics. 2018;19.
doi:10.1186/s12859-018-2234-y

35.   Liao X, Meyer MC. **coneproj** : An *R* Package for the Primal or Dual Cone Projections
with Routines for Constrained Regression. Journal of Statistical Software. 2014;61.
doi:10.18637/jss.v061.i12

36.   Fan Y, Xi L, Hughes DST, Zhang J, Zhang J, Futreal PA, et al. MuSE: accounting for
tumor heterogeneity using a sample-specific error model improves sensitivity and specificity in
mutation calling from sequencing data. Genome Biology. 2016;17. doi:10.1186/s13059-016-
1029-6

37.   Perduca V, Omichessan H, Baglietto L, Severi G. Mutational and epigenetic signatures in
cancer tissue linked to environmental exposures and lifestyle: Current Opinion in Oncology.
2018;30: 61–67. doi:10.1097/CCO.0000000000000418

Supporting information

S1 Fig. Comparison of newly identified signatures with COSMIC signatures. Signatures a-g were identified in a de novo extraction using the maftools R package from the TCGA Lung Adenocarcinoma cohort of 563 cancer genomes. The novel signatures were then compared to the 30 signatures validated in the COSMIC database in terms of cosine similarity. Each signature is then assigned to the most similar COSMIC signature provided that their cosine similarity is above a fixed threshold. For instance, signature f is matched to signature 5 at a cut-off of 0.75 but is considered as a completely new signature if the cut-off is at 0.80. Also note that a unique assignment can be controversial: for instance, signature g is similar both to signatures 12 and 26.

S2 Fig. Cosine similarity plot of COSMIC signatures. Some COSMIC signatures are very similar to others. For instance, signature 8 is similar to signatures 3, 4 and 5.

S3 Fig. Simulations of 563 lung adenocarcinoma catalogues according to different models. **(a)** Real catalogues from the TCGA lung adenocarcinoma cohort. **(b)-(c)** Catalogues sampled from the ZIP model described in the main text. The relative contribution q_n of each signature n is the mean of the relative contributions of n in all samples as estimated by Maftools. In **(b)** simulated and real catalogues are in a 1 to 1 correspondence: for each simulated sample g, the total number of mutations r_g in the corresponding real catalogue is taken. In **(c)** all samples are simulated according to $r = 306$, the average total number of mutations in the real data. The latter example illustrates the parametric model used for the main empirical study. **(d)** Catalogues

sampled according to the Poisson model $e_g^n \sim \mathcal{P}(\lambda_n)$, where λ_n is the mean number of mutations

due do to n in the real samples as estimated by Maftools.

S4 Fig. Choice of parameters q_n in the simulations. Four different configurations $(q_1, \dots, q_{30})$

were considered for simulating realistic data. Each configuration represents the average share of

mutations due to the different COSMIC signatures and was chosen to mimic real exposure

profiles for four cancer types: estimates were obtained from Breast Cancer (Profile 1),

Lymphoma (Profile 2), Lung Adenocarcinoma (Profile 3) and Melanoma (Profile 4) TCGA

cohorts.

S1 File. Mutational Cone. Description of our original method for signature refitting and the

corresponding R script.

Fig 1

Fig 2

Number of mutations

Average number of mutations

Profile 1
Profile 2
Profile 3
Profile 4

100
10000
100000

Signature 1   Signature 3   Signature 5   Signature 7   Signature 9   Signature 11   Signature 17   Signature 20   Signature 30
Signature 2   Signature 4   Signature 6   Signature 8   Signature 10   Signature 13   Signature 18   Signature 26

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7

Fig 8

Fig 9

Fig 10

S1 Fig

	Signature 1	Signature 2	Signature 3	Signature 4	Signature 5	Signature 6	Signature 7	Signature 8	Signature 9	Signature 10	Signature 11	Signature 12	Signature 13	Signature 14	Signature 15	Signature 16	Signature 17	Signature 18	Signature 19	Signature 20	Signature 21	Signature 22	Signature 23	Signature 24	Signature 25	Signature 26	Signature 27	Signature 28	Signature 29	Signature 30
a	0.16	0.03	0.49	0.86	0.38	0.25	0.08	0.6	0.25	0.16	0.11	0.21	0.08	0.37	0.15	0.33	0.05	0.62	0.23	0.43	0.08	0.18	0.12	0.75	0.36	0.13	0.09	0.03	0.7	0.14
b	0.26	0.31	0.51	0.27	0.55	0.24	0.28	0.4	0.67	0.15	0.32	0.41	0.12	0.25	0.21	0.53	0.83	0.17	0.38	0.32	0.22	0.13	0.26	0.23	0.39	0.35	0.09	0.62	0.23	0.4
c	0.95	0.22	0.23	0.18	0.44	0.88	0.33	0.29	0.23	0.31	0.35	0.18	0.08	0.6	0.68	0.21	0.11	0.19	0.51	0.43	0.28	0.05	0.34	0.2	0.33	0.31	0.08	0.05	0.33	0.47
d	0.17	0.08	0.59	0.95	0.46	0.23	0.14	0.77	0.32	0.27	0.16	0.25	0.06	0.43	0.18	0.41	0.07	0.73	0.31	0.41	0.07	0.3	0.16	0.77	0.5	0.14	0.13	0.05	0.78	0.21
e	0.16	0.86	0.36	0.14	0.23	0.09	0.48	0.17	0.13	0.17	0.28	0.12	0.8	0.12	0.09	0.26	0.09	0.14	0.22	0.11	0.06	0.02	0.11	0.21	0.23	0.12	0.07	0.04	0.13	0.46
f	0.34	0.11	0.67	0.48	0.77	0.27	0.52	0.49	0.45	0.18	0.58	0.47	0.12	0.3	0.2	0.63	0.19	0.28	0.66	0.39	0.24	0.25	0.56	0.32	0.55	0.37	0.16	0.14	0.3	0.65
g	0.31	0.08	0.57	0.39	0.76	0.46	0.15	0.5	0.5	0.14	0.27	0.8	0.1	0.42	0.56	0.7	0.17	0.23	0.44	0.55	0.7	0.27	0.31	0.36	0.6	0.79	0.18	0.17	0.38	0.34

S2 Fig

S3 Fig

S4 Fig

S1 File

MutationalCone

We report here the R code implementing our original method for signature refitting.

Let be the linear subspace of spanned by the reference signatures. Our function MutationalCone() projects the input mutational catalogue onto the cone in spanned by the reference signatures with the very fast coneproj R package (https://cran.r-project.org/web/packages/coneproj). Because projections are simply calculated as scalar products, this function requires the user to specify an orthonormal basis of toghether with the components of the reference signatures with respect to it. These two input matrices can be calculated with the function SignatureSubspace() once and for all, before iterating MutationalCone() on all catalogues. SignatureSubspace() finds an orthonormal basis of with the Gram-Schmidt algorithm.

```r
SignatureSubspace <- function(signatures){
# signatures: (K,N)-matrix with reference signatures in columns (e.g.   # COSMIC signatures)
# with K = number of mutation types (e.g. 96), N = number of reference    # signatures

# Orthonormalization of the subspace generated by reference signatures
S <- signatures
S.qr <- qr(S)
Q <- qr.Q(S.qr) # orthonormal basis of the subspace
R <- qr.R(S.qr) # components of the reference signatures in the orthonormal basis

return(list(Q=Q, R=R))
}

MutationalCone <- function(catalogue, Q, R){
# catalogue: vector of length K with the mutational catalogue,
# Q: matrix with the orthonormal basis of the subspace generated
# by the reference signatures in columns
# R: matrix with the components of the reference signatures wrt
# the orthonormal basis in columns. Q and R are found with            # SignatureSubspace()

require(coneproj)

# Projection of the catalogue onto the subspace generated by
# reference signatures
proj.subspace <- t(Q) %*% catalogue

# Projection onto the cone spanned by the signatures
weights <- as.vector(coneB(y=as.vector(proj.subspace),delta=R)$coefs)
return(weights)
}
```