Discovery of Novel Glycerolated Quinazolinones from *Streptomyces* sp. MBT27

Nataliia Machushynets [1], Changsheng Wu [1,2] *, Somayah S. Elsayed [1], Thomas Hankemeier [3] and Gilles

P. van Wezel [1], *

[1] Molecular Biotechnology, Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE, The

Netherlands

[2] State Key Laboratory of Microbial Technology, Institute of Microbial Technology, Shandong

University, Qingdao 266237, P.R. China

[3] Leiden Academic Centre for Drug Research (LACDR), Leiden University, Einsteinweg 55, 2333 CC,

Leiden, The Netherlands

* Author for correspondence. Tel: +31 71 5274310; email: g.wezel@biology.leidenuniv.nl,

or wcs0313@gmail.com

Abstract

Actinobacteria are a major source of novel bioactive natural products. A challenge in the screening of

these microorganisms lies in finding the favorable growth conditions for secondary metabolite

production and dereplication of known molecules. Here, we report that *Streptomyces* sp. MBT27

produces 4-quinazolinone alkaloids in response to elevated levels of glycerol, whereby quinazolinones

A (**1**) and B (**2**) form a new sub-class of this interesting family of natural products. Global Natural

Product Social molecular networking (GNPS) resulted in a quinazolinone-related network that included

anthranilic acid (**3**), anthranilamide (**4**), 4(3H)-quinazolinone (**5**) and 2,2-dimethyl-1,2-

dihydroquinazolin-4(3*H*)-one (**6**). Actinomycins D (**7**) and X2 (**8**) were also identified in the extracts of

Streptomyces sp. MBT27. The induction of quinazolinone production by glycerol combined with

biosynthetic insights provide evidence that glycerol is integrated into the chemical scaffold. The

unprecedented 1,4-dioxepane ring that is spiro-fused into the quinazolinone backbone, is most likely

formed by intermolecular etherification of two units of glycerol. Our work underlines the importance

of varying the growth conditions for the discovery of novel natural products and for understanding

their biosynthesis.

Keywords: 4(3*H*)-quinazolinones, molecular networking*,* metabolomics, carbon source

Introduction

Actinobacteria are a major source of bioactive compounds, producing some two third of all antibiotics as well as molecules with a wide variety of activities such as anticancer, antifungal and immunosuppressant [2,4]. Traditionally, microbial natural product (NP) discovery has been done via high-throughput screening followed by iterative bioassay-guided fractionation and structure elucidation. While such pipelines were extremely successful and delivered a plethora of therapeutic agents, in the modern era the large pharmaceutical companies move out of NP-discovery programs due to high cost and chemical redundancy [1,9,28]. At the same time, the power of genome sequencing brought the recognition that microorganisms harbor a vast and yet untapped biosynthetic potential, and it rapidly became clear that the potential for metabolic diversity of even the best-studied model organisms as producers of natural products had been grossly underestimated [3,11,18]. How could these compounds have been missed by the very extensive HT screening campaigns of the 20th century? The answer is that many of the biosynthetic gene clusters (BGCs) discovered by genome mining are poorly expressed or cryptic under laboratory conditions [21,27]. A drug-discovery pipeline that is rapidly gaining momentum involves combining genome mining with fluctuating the culturing conditions to achieve differential synthesis of NPs, followed by the metabolic profiling-based identification of the bioactivity of interest [13,46]. A major challenge thereby lies in finding the appropriate chemical triggers or ecological cues to elicit the production of cryptic antibiotics (recently reviewed in [33,48,49]). The use of chemical elicitors is thereby a promising approach [10,49].

Manipulation of fermentation conditions for promising producer strains, known as the "one strain many compounds" (OSMAC) approach, is an effective way of enhancing the production of secondary metabolites [32,5]. The regulatory networks that control the production of bacterial natural products respond strongly to changes in carbon, nitrogen or phosphate concentration [40,39]. The re-isolation of known metabolites is a major bottleneck in the discovery of new bioactive natural products. A crucial step in this regard is the early identification of already known substances, in order to concentrate the efforts on the discovery of new ones, a process known as dereplication [15]. Current

dereplication strategies include hyphenated techniques, such as LC-MS, LC-NMR, LC-NMR-MS, and LC-

SPE-NMR. Bioactivity fingerprinting has also been used to dereplicate natural products based on their

biological modes of action [35], while molecular networking is a powerful tool for the visualization and

dereplication of natural products [47,44]. Mass spectrometry-based molecular networking relies on

clustering of molecules based on similarities in their MS/MS fragmentation patterns, which depends

on the structural features of the ionized molecules. The resulting clusters allow scientists to visually

explore the metabolites produced by a given strain under a specific growth condition, allowing rapid

dereplication of known compounds by automated spectral library searches, and to visualize their

unknown structural analogues [43].

In this work we analyzed the potential of *Streptomyces* sp. MBT27 as producer of natural

products in response to changes in the carbon source. The strain had previously been identified as a

promising producer of NPs [50]. Extensive fluctuations in the secondary metabolite profiles were

observed depending on the carbon source used, and statistical methods combined with GNPS

molecular networking identified a family of known as well as novel quinazolinone compounds, in

response to high concentrations of glycerol. Quinazolinones are heterocyclic compounds with a wide

range of medical applications, such as antimicrobial, antiviral, antituberculosis and as enzyme

inhibitors (reviewed in [23,16,19]). Combination of MS and NMR methods identified the novel

quinazolinones A (**1**) and B (**2**), which further expands the chemical space of this rich family of natural

products.

Materials and Methods

Bacterial strain, growth conditions

Streptomyces sp. MBT27 was obtained from the Leiden University strain collection and had previously

been isolated from the Qingling Mountains, Shanxi province, China [50]. Cultures were grown in

triplicate in 100 mL Erlenmeyer flasks with 30 mL of liquid minimal medium (MM; [20]), supplemented

with various carbon sources, and inoculated with 10 µL of 10^9/mL spore suspension. The carbon

sources (percentages in w/v) were: 1% mannitol + 1% glycerol, 1% mannitol, 2% mannitol, 1% glycerol,

2% glycerol, 1% glucose, 2% glucose, 1% fructose, 1% arabinose or 1% *N*-acetylglucosamine (GlcNAc).

The cultures were incubated in a rotary shaker at 30 °C at 220 rpm for seven days.

General Experimental Procedures

NMR spectra were recorded in deuterated methanol (CD_3OD) on a Bruker 600 MHz, and were

referenced using the residual ^1H signal of deuterated solvent at 3.30 ppm. FT-IR was measured on

Perkin-Elmer FT-IR Spectrometer Paragon 1000. UV measurements were performed using a Shimadzu

UV mini-1240. Optical rotations were measured on a JASCO P-1010 polarimeter. HPLC purification was

performed on Waters preparative HPLC system comprised of 1525 pump, 2707 autosampler, 2998 PDA

detector, and Water fraction collector III. The columns used were SunFire C_{18} column (5 µm, 100 Å, 10

× 250 mm) and SunFire C_{18} column (10 µm, 100 Å, 19 × 150 mm). TLC was performed using aluminum

plates coated with silica gel 60 F_{254} (Merck). All organic solvents and chemicals were of analytical

or LCMS grade, depending on the experiment.

Metabolite profiling

Following fermentation, culture supernatants were extracted with ethyl acetate (EtOAc) and

evaporated under reduced pressure. For LC-ESI-QTOFMS analyses, extracts were dissolved in MeOH

to a final concentration of 1 mg/mL, and 1 µL was injected into Waters Acquity UPLC system equipped

with Waters Acquity HSS C_{18} column (1.8 µm, 100 Å, 2.1 × 100 mm), which is coupled to Agilent 6530

QTOF MS equipped with Agilent Jet Stream ESI source (Agilent Technologies, Inc., Palo Alto, CA, USA).

For the LC, solvent A was 95% H_2O, 5% acetonitrile (ACN) and 0.1% formic acid; solvent B was 100%

ACN and 0.1% formic acid. The gradient used was 2% B for 1 min, 2–85% for 9 min, 85–100% for 1 min,

and 100% for 3 min. The flow rate used was 0.5 mL/min. As for the MS, the following ESI source

parameters were used: capillary voltage 3 kV, source temperature 325 °C, drying gas flow rate 10

L/min, and fragmentor 175 V. Full MS spectra were acquired in positive mode in the range of 100-

1700 *m/z*, in the extended dynamic range mode. Internal reference masses of purine and Agilent HP-

921 were continuously delivered to the ESI source through an Agilent 1260 isocratic pump.

Thermo Instruments MS system (LTQ Orbitrap XL, Bremen, Germany) equipped with an

electrospray ionization source (ESI) was used for LC-MS/MS analysis. The Waters Acquity UPLC system

equipped with Waters Acquity PDA was run using a SunFire Waters C_{18} column (3.5 μm, 100 Å, 4.6 ×

150 mm), at a flow rate of 0.9 mL/min. Solvent A was 95% H_2O, 5% acetonitrile (ACN) and 0.1% formic

acid; solvent B was 100% ACN and 0.1% formic acid. The gradient used was 2% B for 1 min, 2-85% for

15 min, 85–100% for 3 min, and 100% for 3 min. As for the MS, the following ESI parameters were

used: capillary voltage 5 V, spray voltage 3.5 kV, capillary temperature 300 °C, auxiliary gas flow rate

10 arbitrary units, and sheath gas flow rate 50 arbitrary units. Full MS spectra were acquired in the

Orbitrap in positive mode at a mass range of 100–2000 *m/z*, and FT resolution of 30000. Data

dependent MS^2 spectra were acquired in the ion trap for the three most intense ions using collision

induced dissociation (CID). The resulting chemical data were compared with those in SciFinder and

Antibase [24].

Up-scale fermentation and isolation of metabolites (1) and (2)

For up-scale fermentation, *Streptomyces* sp. MBT27 was grown in eight 2000 mL Erlenmeyer flasks,

each containing 500 mL liquid MM supplemented with 2% w/v glycerol at 30 °C for seven days. The

metabolites were extracted from the culture broth using EtOAc, then the solvent was evaporated

under reduced pressure at 40 ºC. The crude extract (1.4 g) was adsorbed onto 1.4 g silica gel (pore

size 60 Å, 70-230 mesh, Sigma Aldrich), and loaded on silica column, which was eluted using gradient

mixtures of *n*-hexane, acetone, and MeOH. The fractions eluted with acetone 100% were combined,

reconstituted in MeOH, and injected into the preparative SunFire column (19 x 150 mm), which was

eluted with H_2O:MeOH gradient of 7–100% in 30 min, at a flow rate of 15 mL/min, to yield five

fractions. Quinazolinones A and B were further purified on the semi-preparative SunFire column (10 x

250 mm), run at 3 mL/min. Fraction 2 was eluted using H_2O:MeOH gradient of 20–40% in 20 min, to

yield Quinazolinone A (**1**, 1 mg). On the other hand, Fraction 3 was eluted using H_2O:MeOH gradient

of 30-40% in 20 min, to yield Quinazolinone B (**2**, 1.5 mg).

Quinazolinone A (1): colorless, amorphous powder; $[\alpha]_D^{20}$ 2.3 (c 0.1, MeOH); UV (MeOH) λ_{max} (log ϵ)

224 (4.09), 322 (2.98) nm; IR v_{max} 3334, 2922, 1652, 1475, 1052 cm^{-1}; ^1H and ^{13}C NMR data, see Table

1; HRESIMS (positive mode) *m/z* 281.1130 $[M + H]^+$ (calcd. for $C_{13}H_{17}N_2O_5$, 281.1132).

Quinazolinone B (2): colorless, amorphous powder, UV (MeOH) λ_{max} (log ϵ) 226 (4.18), 347 (3.03) nm;

IR v_{max} 3350, 2950, 1649, 1526, 1049, 751 cm^{-1}; ^1H and ^{13}C NMR data, see Table 1; HRESIMS (positive

mode) *m/z* 209.0915 $[M + H]^+$ (calcd. for $C_{10}H_{13}N_2O_3$, 209.0921).

Computation of mass spectral networks

MS/MS raw data were converted to a 32 bit mzXML file using MSConvert (ProteoWizard) [6] and

spectral networks were assembled using Global Natural Product Social molecular networking (GNPS)

(https://gnps.ucsd.edu) as described [43]. For both parent and MS/MS fragment ions, the mass

tolerance was set to 0.5 Da, while the minimum cosine score was set to 0.7. The data were clustered

using MSCluster with a minimum cluster size of three spectra. The spectra in the network were also

searched against GNPS spectral libraries. A minimum score of 0.5 was set for spectral library search,

with at least two fragment peaks matching. Cytoscape 3.5.1. was used for visualization of the

generated molecular networks [37]. The edge thickness was set to represent the cosine score, with

thicker lines indicating higher similarity between nodes. LC-MS/MS data were deposited in the

MassIVE Public GNPS data set (MSV000082988). The molecular networking job in GNPS can be found

at   https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=8fd1fcfa0ff744a9808e80bc7be12115.   The

annotated MS/MS spectra were deposited in the GNPS spectral library for quinazolinone A

(CCMSLIB00004684355), and B (CCMSLIB00004684354).

Statistical analysis

Prior to statistical analysis, mzXML files were imported into Mzmine 2.31 [29] for data processing. Mass

ion peaks were detected using the exact mass algorithm with a noise level set to 1.0×10^4. Afterwards,

chromatograms were built for the detected masses with a minimum time span of 0.05 min, m/z

tolerance of 0.001 m/z and minimum height of 1.0×10^4. Chromatogram deconvolution was then

performed using local minimum search algorithm (search minimum in RT range 0.1 min,

chromatographic threshold 90%, minimum relative height 1%, minimum absolute height 1.0×10^4,

minimum ratio of peak top/edge 2 and peak duration range 0.05–3 min). In the generated peak lists,

isotopes were identified using isotopic peaks grouper (m/z tolerance 0.001 m/z and retention time

tolerance 0.1 min), and variations in retention time were reduced using retention time normalizer (m/z

tolerance 0.001 m/z and retention time tolerance 1 min). All the peak lists were subsequently aligned

using join aligner (m/z tolerance 0.001 m/z, m/z weight 20, retention time tolerance 0.1 min, and

retention time weight 20), and missing peaks were detected through gap filling using peak finder

(intensity tolerance 1.0%, m/z tolerance 0.001 m/z, and retention time tolerance of 0.2 min). Finally,

the aligned peak list was exported as a comma separated file for statistical analysis.

Statistical analysis was performed using MetaboAnalyst [7], where log transformation and

pareto scaling was initially applied to the data. A heat map of all detected masses, among the different

growth conditions, was generated in MetaboAnalyst, to which additional hierarchical clustering

analysis (HCA) was performed using Euclidean distance measure and Ward clustering algorithm.

Student's t-tests with multiple testing correction (Benjamini Hochberg false discovery rate or FDR)

were used to determine significant differences in the intensities of the metabolites, under two

different growth conditions. The thresholds set for statistically significant differences were a fold

change ≥ 4, together with FDR corrected p-value ≤ 0.05 Based on these criteria, a volcano plot was

generated. To identify the difference in intensity of a single mass feature among multiple growth

conditions, one-way ANOVA was performed, followed by a *post hoc* Tukey's honest significant

difference (HSD) test.

RESULTS AND DISCUSSION

The influence of carbon sources on secondary metabolite production

Previous screening of our in-house actinomycete collection, obtained from remote mountain soils, showed it is a promising source of new bioactive compounds [50]. Under specific growth conditions, these isolates exert potent inhibitory activity against the so-called ESKAPE pathogens [30] *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacter* spp. In the current study, the potential of one of the *Streptomyces* strains in our collection, namely *Streptomyces* sp. MBT27, was analyzed to study the effect of carbon sources on its metabolite profile. Traditional approaches such as changes in fermentation conditions, are known to induce significant changes in the microbial metabolome. Culture medium components, and particularly the carbon source, have major effects on the production of secondary metabolites [41,34,5,25]. The differential production of small molecules is ideal for metabolomics studies, whereby metabolic variations are correlated statistically to bioactivity, thus facilitating the identification of the bioactive molecule of interest [46,49]. To establish the potential of *Streptomyces* sp. MBT27, the strain was grown in liquid minimal media containing different carbon sources, namely (percentages in w/v): 1% of both mannitol and glycerol, 1% mannitol, 2% mannitol, 1% glycerol, 2% glycerol, 1% glucose, 2% glucose, 1% fructose, 1% arabinose, or 1% *N*-acetylglucosamine (GlcNAc). The latter is an elicitor of antibiotic produciton, via metabolic interference with the global nutrient sensory network controlled by DasR [38,31]. Phosphate buffer was omitted from the MM as it repressed the production of secondary metabolites by *Streptomyces* sp. MBT27 (data not shown). To identify the secondary metabolites in the cultures, supernatants were extracted with EtOAc, and the resulting crude extracts subjected to LC-MS analysis. Using the LC-MS data, a heat map with added hierarchical clustering was generated, to visualize the production of different metabolites under different culture conditions (Fig. 1). Hierarchical clustering analysis of the LC-MS data allows effective comparative analysis of metabolomics data, and the heat map revealed major

differences in the metabolic profiles of *Streptomyces* sp. MBT27, whereby different groups of

metabolites were enhanced depending on the carbon source used. Interestingly, not only the type of

carbon source, but especially also the concentration resulted in large changes in the metabolic profiles.

Doubling the concentration of either glycerol or glucose from 1% to 2% had a particularly profound

change on the metabolic profile. Thin layer chromatography (TLC) was conducted to compare

metabolic profiles of the 1% and 2% glycerol-grown cultures. Interestingly, this revealed that several

fluorescent compounds were differentially produced in the extracts of 2% glycerol-grown cultures

relative to those produced in 1% glycerol (Fig. 2a). In order to provide statistical relevance to the data,

the metabolic profiles of 1% and 2% glycerol-grown cultures were compared using a volcano plot (Fig.

2b). The volcano plot was then searched for the mass features which increased in production in 2%

glycerol as compared to 1%. A mass of *m/z* 281.1151 (**1**) stood out as its intensity had increased by

around 7000 fold (*p*-value = 0.002) in cultures fermented in 2% glycerol as compared to 1%.

Global Natural Product Social (GNPS) molecular networking [43] was employed to detect the

MS/MS structural relatedness among molecules in an automated manner; the software generates a

molecular network wherein molecules with related scaffolds cluster together [43]. A network

representing the ions detected in the crude extract of *Streptomyces sp.* MBT27 grown with 2% glycerol

was constructed, revealing 183 nodes clustered in 19 spectral families (Fig. 3). GNPS dereplication

based on matching with its MS/MS spectral database highlighted some known metabolites. These

included anthranilic acid (**3**), anthranilamide (**4**) [36], actinomycin D (**7**), and actinomycin X2 (**8**) [42].

The annotation of the compounds was supported by comparison of the exact mass, fragmentation

pattern, and UV spectra, with reference data. Moreover, crude extracts of MBT27 possessed

antimicrobial activity against *B. subtilis*, which was most likely due to the expression of actinomycins

(data not shown). Additional metabolites could be dereplicated through manual comparison of their

spectral data against the microbial natural products database Antibase. These previously described

metabolites were 4(3*H*)-quinazolinone (**5**) [22] and 2,2-dimethyl-1,2-dihydroquinazolin-4(3*H*)-one (**6**)

[8].

Besides known molecules, the network also contained many mass features that could not be

assigned to any of the previously identified metabolites. One of these was an ion with an *m/z* value of

281.1151 (**1**), which was very highly increased in intensity (7000 fold, *p*-value = 0.002) when the

glycerol concentration was increased from 1% to 2%. It was closely connected (cosine score 0.91) to

another upregulated ion with an *m/z* value of 209.0932 (**2**), which was also not previously reported.

Both ions were part of the large spectral family comprising the annotated anthranilic acid (**3**),

anthranilamide (**4**), and the 4-quinazolinones **5** and **6**, suggesting that they are structurally related

metabolites. The relationship between the increase in glycerol concentration and the strong increase

in the production of the compounds **1** and **2**, was investigated further by expanding the range of

glycerol concentrations and analyzing the metabolic profiles. For this, *Streptomyces* sp. MBT27 was

cultured in glycerol concentrations ranging from 1-4% (w/v). ANOVA, followed by a *post hoc* Tukey's

HSD test, was performed to trace the variation in the production of **1** and **2** among the different

culturing conditions. A box plot was used to visualize such variation (Fig. S1). As observed earlier, the

production of **1** and **2** was significantly increased when the glycerol concentration was increased from

1% to 2%. However, further increase in glycerol concentration (3% and 4%), did not lead to any

significant increase in the production of **1** and **2**. Accordingly, MM with 2% glycerol was used to culture

the bacteria, for the purpose of purification and identification of the new metabolites **1** and **2**.

Isolation and structure elucidation of novel quinazolinones

To elucidate the structure of **1** and **2**, upscale fermentation was done to obtain larger quantities of the

compounds. For this, *Streptomyces* sp. MBT27 was fermented in a total of 4 L of liquid MM

supplemented with 2% glycerol, and the supernatant was extracted with EtOAc. Following repeated

chromatographic isolation, compounds **1** and **2** were obtained as pure, colorless, amorphous powders.

Both compounds were fluorescent, showing UV absorption maxima at 224 and 322 nm and at 226 and

347 nm, respectively. The final structures of **1** and **2** were determined by the combination of NMR and

high resolution MS (Fig. 4).

The high resolution mass of m/z 281.1151 for an [M + H]$^+$ established a molecular formula of

$C_{13}H_{16}N_2O_5$, with seven degrees of unsaturation, for **1** (yield 0.25 mg/L). The deduced molecular

formula was corroborated by the ^{13}C NMR attached proton test (APT) spectrum that exhibited 13

carbons in total. On the other hand, the ^1H NMR spectrum of **1** (Table 1) presented four coupling

aromatic signals at δ_H 7.88 (dd, J = 7.8, 1.2, H-5), 7.07 (t, J = 7.8, H-6), 7.46 (td, J = 7.8, 1.2, H-7), and

7.23 (brd, J = 7.8, H-8), indicating an o-disubstituted aromatic benzene system. The HMBC correlation

(Fig. 5) from H-5 to C-4 (δ_C 166.1) confirmed one of substituents on the benzene ring to be an

ester/amide carbonyl group, while the downfield chemical shift of C-8a at δ_C 143.7 indicated it was

nitrogenated. Four O-bearing methylene groups at δ_H 4.43 (d, J = 9.0, H-1'a) and 4.24 (d, J = 9.0, H-1'b);

3.59 (d, J = 11.4, H-2'a) and 3.47 (d, J = 11.4, H-2'b); 3.86 (d, J = 12.6, H-1''a) and 3.80 (d, J = 12.6, H-

1''b); and 3.66 (d, J = 3.0, H$_2$-3'') were resolved by HSQC experiments. Three of these methylene groups

were part of a 1,4-dioxepane ring system, which was established based on the HMBC correlations

observed from H-1' to C-2 (δ_C 79.0) and C-2' (δ_C 64.6), from H-2' to C-1'' (δ_C 65.7), and from H-1' to the

hemiacetal C-2'' (δ_C 102.9). The remaining oxymethylene group CH$_2$-3'' was connected to C-2'' based

on the HMBC correlations observed from H$_2$-3'' to C-1'' and C-2''. The two substructures obtained

accounted for all of the oxygens and six out of the seven degrees of unsaturation required by the

molecular formula of **1**. Accordingly, an additional ring including two nitrogen atoms was deduced to

connect the two substructures, forming a 4-quinazolinone ring. Consequently, compound **1** was

identified to be a di-glycerolated 4-quinozolinone, and was named quinazolinone A.

The [M + H]$^+$ molecular ion at m/z 209.0932 in the ESI-HRMS spectrum resulted in a molecular

formula of $C_{10}H_{12}N_2O_3$ for **2** (yield 0.3 mg/L). The ^{13}C NMR spectrum presented 10 signals, which was

consistent with the molecular formula (Table 1). The ^1H NMR spectrum of **2** resembled that of **1** in the

downfield aromatic region, and the major difference is the absence of a set of signals for the additional

glycerol unit. Further HMBC experiment confirmed **2** is a $mono$-glycerolated 4-quinazolinone (Fig. 5).

Compound **2** was named quinazolinone B.

The 4-quinazoline alkaloids represent an important class of nitrogen-containing heterocyclic

compounds [26], which have highly diverse biological activities, such as antitumor, anti-inflammatory,

antihypertensive, antimicrobial, anticonvulsant and antifungal activities [17]. Though the basic bicyclic

core (the fused benzene ring and pyrimidine ring) is normally conserved in most of naturally occurring

4-quinazoline derivatives [26], a variety of structural substitutions have oftentimes been found on the

pyrimidine ring of the 4-quinazolinone. Quinazolinones A and B isolated from *Streptomyces* sp. MBT27

represent the first example that glycerol participates in the construction of the ring system of 4-

quinazolines. In this sense, quinazolinones A and B can be perceived as a new sub-branch in the family

of the quinazolines. Particularly, quinazolinone A contains an exciting seven-membered ring formed

by intermolecular etherification of two units of glycerol, and this ring is further linked with the

quinazolinone backbone by a spiro atom at C-2, which is unprecedented. Antimicrobial activities of

compounds **1** and **2** were tested against *B. subtilis* and *E. coli*. None of the compounds showed

antimicrobial activity against these indictor strains (data not shown). Further bioactivity screening is

required to find out what the biological activity of the two compounds is.

Biosynthesis of quinazolinones A and B

In our study, the production of **1** and **2** was dramatically enhanced in the cultures with 2% glycerol.

Elicitation by glycerol is an indication that glycerol may play a key role during their biosynthesis. The

aromatic ring system of quinazoline alkaloids is known to be derived from anthranilic acid, which in

turn is biosynthesized through the shikimate pathway [12]. In line with this, we propose that the

biosynthesis of the isolated quinazolinones starts from anthranilic acid, which is then converted to

anthranilamide (Fig. 6). Further successive attachments of two molecules of glycerol then results in

the formation of **2**, followed by **1**. However, multiple biosynthetic routes have been proposed for

quinazolinones, because the C-2 residue of the quinazoline ring may originate from various precursors.

For example, the C-2 and the remaining non-aromatic part of the quinazolinone alkaloid chrysogine,

produced by *Penicillium chrysogenum*, are biosynthesized from pyruvic acid via an NRPS system [45],

while the non-aromatic part of the quinazoline alkaloid peganine, produced by the plant *Peganum*

harmala, is derived from ornithine [12]. Furthermore, fumiquinazoline F originates from a fungal

nonribosomal peptide synthetase (TqaA) and is biosynthesized from anthranilic acid, L-tryptophan and

L-alanine [14].

In conclusion, the metabolic potential of *Streptomyces* sp. MBT27 was analyzed under different

nutritional conditions, showing major changes in the metabolome, depending on the carbon source

used. In particular, the relatively unspectacular change from 1% to 2% glycerol resulted in a surprisingly

global change in the secondary metabolome, with some compounds changing by almost four orders

of magnitude. The use of GNPS molecular networking, together with LC-MS based dereplication,

allowed us to annotate a cluster of quinazolinone-family molecules that connects to anthranilic acid

and anthranilamide. Isolation and structure elucidation revealed the novel alkaloid natural products

quinazolinone A and B. Based on the structures, the biosynthesis of the compounds most likely involves

the conversion of anthranilic acid to anthranilamide, which is subsequently attached to two glycerol

units, to produce compounds **1** and **2**. Identification of the biosynthetic gene cluster and subsequent

analysis of the individual enzymatic reactions should reveal the precise biosynthetic pathway for these

exciting novel molecules.

Reference list

1. Baltz RH (2008) Renaissance in antibacterial discovery from actinomycetes. Curr Opin Pharmacol

8:557-563

2. Barka EA, Vatsa P, Sanchez L, Gavaut-Vaillant N, Jacquard C, Meier-Kolthoff J, Klenk HP, Clément C,

Oudouch Y, van Wezel GP (2016) Taxonomy, physiology, and natural products of the

Actinobacteria. Microbiol Mol Biol Rev 80:1-43

3. Bentley SD, Chater KF, Cerdeno-Tarraga AM, Challis GL, Thomson NR, James KD, Harris DE, Quail MA,

Kieser H, Harper D, Bateman A, Brown S, Chandra G, Chen CW, Collins M, Cronin A, Fraser A,

Goble A, Hidalgo J, Hornsby T, Howarth S, Huang CH, Kieser T, Larke L, Murphy L, Oliver K,

O'Neil S, Rabbinowitsch E, Rajandream MA, Rutherford K, Rutter S, Seeger K, Saunders D,

Sharp S, Squares R, Squares S, Taylor K, Warren T, Wietzorrek A, Woodward J, Barrell BG,

Parkhill J, Hopwood DA (2002) Complete genome sequence of the model actinomycete

Streptomyces coelicolor A3(2). Nature 417:141-147

4. Bérdy J (2005) Bioactive microbial metabolites. J Antibiot (Tokyo) 58:1-26

5. Bode HB, Bethe B, Höfs R, Zeeck A (2002) Big Effects from Small Changes: Possible Ways to Explore

Nature's Chemical Diversity. ChemBioChem 3:619–627

6. Chambers MC, Maclean B, Burke R, Amodei D, Ruderman DL, Neumann S, Gatto L, Fischer B, Pratt

B, Egertson J, Hoff K, Kessner D, Tasman N, Shulman N, Frewen B, Baker TA, Brusniak MY,

Paulse C, Creasy D, Flashner L, Kani K, Moulding C, Seymour SL, Nuwaysir LM, Lefebvre B,

Kuhlmann F, Roark J, Rainer P, Detlev S, Hemenway T, Huhmer A, Langridge J, Connolly B,

Chadick T, Holly K, Eckels J, Deutsch EW, Moritz RL, Katz JE, Agus DB, MacCoss M, Tabb DL,

Mallick P (2012) A cross-platform toolkit for mass spectrometry and proteomics. Nat

Biotechnol 30:918–920

7. Chong J, Soufan O, Li C, Caraus I, Li S, Bourque G, Wishart DS, Xia J (2018) MetaboAnalyst 4.0: towards

more transparent and integrative metabolomics analysis. Nucleic Acids Res 46:W486-W494

8. Conti R, Chagas FO, Caraballo-Rodriguez AM, Melo WG, do Nascimento AM, Cavalcanti BC, de

Moraes MO, Pessoa C, Costa-Lotufo LV, Krogh R, Andricopulo AD, Lopes NP, Pupo MT (2016)

Endophytic Actinobacteria from the Brazilian Medicinal Plant Lychnophora ericoides Mart.

and the Biological Potential of Their Secondary Metabolites. Chem Biodivers 13:727–736

9. Cooper MA, Shlaes D (2011) Fix the antibiotics pipeline. Nature 472:32

10. Craney A, Ozimok C, Pimentel-Elardo SM, Capretta A, Nodwell JR (2012) Chemical perturbation of

secondary metabolism demonstrates important links to primary metabolism. Chem Biol

19:1020-1027

11. Cruz-Morales P, Vijgenboom E, Iruegas-Bocardo F, Girard G, Yanez-Guerra LA, Ramos-Aboites HE,

Pernodet JL, Anne J, van Wezel GP, Barona-Gomez F (2013) The genome sequence of

Streptomyces lividans 66 reveals a novel tRNA-dependent peptide biosynthetic system within

a metal-related genomic island. Genome Biol Evol 5:1165-1175

12. Dewick P (2009) Medicinal Natural Products: A Biosynthetic Approach. 3rd edn. ohn Wiley & Sons,

Chichester, UK

13. Du C, van Wezel GP (2018) Mining for Microbial Gems: Integrating Proteomics in the Postgenomic

Natural Product Discovery Pipeline. Proteomics:e1700332

14. Gao X, Chooi YH, Ames BD, Wang P, Walsh CT, Tang Y (2011) Fungal indole alkaloid biosynthesis:

genetic and biochemical investigation of the tryptoquialanine pathway in *Penicillium*

aethiopicum. J Am Chem Soc 133:2729–2741

15. Gaudencio SP, Pereira F (2015) Dereplication: racing to speed up the natural products discovery

process. Nat Prod Rep 32:779–810

16. Hameed A, Al-Rashida M, Uroos M, Ali SA, Arshia, Ishtiaq M, Khan KM (2018) Quinazoline and

quinazolinone as important medicinal scaffolds: a comparative patent review (2011-2016).

Expert Opin Ther Pat 28:281-297

17. He D, Wang M, Zhao S, Shu Y, Zeng H, Xiao C, Lu C, Liu Y (2017) Pharmaceutical prospects of

naturally occurring quinazolinone and its derivatives. Fitoterapia 119:136–149

18. Ikeda H, Ishikawa J, Hanamoto A, Shinose M, Kikuchi H, Shiba T, Sakaki Y, Hattori M, Omura S (2003)

Complete genome sequence and comparative analysis of the industrial microorganism

Streptomyces avermitilis. Nat Biotechnol 21:526-531

19. Jafari E, Khajouei MR, Hassanzadeh F, Hakimelahi GH, Khodarahmi GA (2016) Quinazolinone and

quinazoline derivatives: recent structures with potent antimicrobial and cytotoxic activities.

Res Pharm Sci 11:1-14

20. Kieser T, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA (2000) Practical *Streptomyces* genetics. John

Innes Foundation, Norwich, U.K.

21. Kolter R, van Wezel GP (2016) Goodbye to brute force in antibiotic discovery? Nat Microbiol

1:15020

22. Kornsakulkarn J, Saepua S, Srijomthong K, Rachtawee P, Thongpanchang C (2015) Quinazolinone

alkaloids from actinomycete Streptomyces sp. BCC 21795. Phytochemistry Letters 12:6–8

23. Kshirsagar UA (2015) Recent developments in the chemistry of quinazolinone alkaloids. Org Biomol

Chem 13:9336-9352

24. Laatsch H (2012) Antibase 2012, the Natural Compound Identifier.

25. Martin JF, Sola-Landa A, Santos-Beneit F, Fernandez-Martinez LT, Prieto C, Rodriguez-Garcia A

(2011) Cross-talk of global nutritional regulators in the control of primary and secondary

metabolism in *Streptomyces*. Microb Biotechnol 4:165-174

26. Michael JP (2000) Quinoline, quinazoline and acridone alkaloids. Nat Prod Rep 17: 603–620

27. Nett M, Ikeda H, Moore BS (2009) Genomic basis for natural product biosynthetic diversity in the

actinomycetes. Nat Prod Rep 26:1362-1384

28. Payne DJ, Gwynn MN, Holmes DJ, Pompliano DL (2007) Drugs for bad bugs: confronting the

challenges of antibacterial discovery. Nat Rev Drug Discov 6:29-40

29. Pluskal T, Castillo S, Villar-Briones A, Orešič M (2010) MZmine 2: Modular framework for

processing, visualizing, and analyzing mass spectrometry-based molecular profile data. BMC

Bioinformatics 11:395

30. Rice LB (2008) Federal funding for the study of antimicrobial resistance in nosocomial pathogens:

no ESKAPE. J Infect Dis 197:1079-1081

31. Rigali S, Titgemeyer F, Barends S, Mulder S, Thomae AW, Hopwood DA, van Wezel GP (2008) Feast

or famine: the global regulator DasR links nutrient stress to antibiotic production by

Streptomyces. EMBO Rep 9:670-675

32. Romano S, Jackson SA, Patry S, Dobson ADW (2018) Extending the "One Strain Many Compounds"

(OSMAC) principle to marine microorganisms. Mar Drugs 16:244

33. Rutledge PJ, Challis GL (2015) Discovery of microbial natural products by activation of silent

biosynthetic gene clusters. Nat Rev Microbiol 13:509-523

34. Sanchez S, Chavez A, Forero A, Garcia-Huante Y, Romero A, Sanchez M, Rocha D, Sanchez B, Avalos

M, Guzman-Trampe S, Rodriguez-Sanoja R, Langley E, Ruiz B (2010) Carbon source regulation

of antibiotic production. J Antibiot (Tokyo) 63:442-459

35. Schulze CJ, Bray WM, Woerhmann MH, Stuart J, Lokey RS, Linington RG (2013) "Function-first" lead

discovery: mode of action profiling of natural product libraries using image-based screening.

Chem Biol 20:285–295

36. Shaaban KA, Shepherd MD, Ahmed TA, Nybo SE, Leggas M, Rohr J (2012) Pyramidamycins A-D and

3-hydroxyquinoline-2-carboxamide; cytotoxic benzamides from *Streptomyces* sp. DGC1. J

Antibiot (Tokyo) 65:615–622

37. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T

(2003) Cytoscape: a software environment for integrated models of biomolecular interaction

networks. Genome Res 13:2498–2504

38. Swiatek-Polatynska MA, Bucca G, Laing E, Gubbens J, Titgemeyer F, Smith CP, Rigali S, van Wezel

GP (2015) Genome-wide analysis of *In vivo* binding of the master regulator DasR in

Streptomyces coelicolor identifies novel non-anonical targets. PLoS One 10:e0122479

39. Urem M, Swiatek-Polatynska MA, Rigali S, van Wezel GP (2016) Intertwining nutrient-sensory

networks and the control of antibiotic production in Streptomyces. Mol Microbiol 102:183-

195

40. van der Heul HU, Bilyk BL, McDowall KJ, Seipke RF, van Wezel GP (2018) Regulation of antibiotic

production in Actinobacteria: new perspectives from the post-genomic era. Nat Prod Rep

35:575-604

41. van Wezel GP, McKenzie NL, Nodwell JR (2009) Applying the Genetics of Secondary Metabolism in

Model Actinomycetes to the Discovery of New Antibiotics. In: Complex Enzymes in Microbial

Natural Product Biosynthesis, Part A: Overview Articles and Peptides. Methods in

Enzymology. pp 117–141. doi:https://doi.org/10.1016/s0076-6879(09)04805-8

42. Vater J, Crnovcic I, Semsary S, Keller U (2014) MALDI-TOF mass spectrometry, an efficient technique

for in situ detection and characterization of actinomycins. J Mass Spectrom 49:210–222

43. Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Nguyen DD, Watrous J, Kapono CA,

Luzzatto-Knaan T, Porto C, Bouslimani A, Melnik AV, Meehan MJ, Liu WT, Crusemann M,

Boudreau PD, Esquenazi E, Sandoval-Calderon M, Kersten RD, Pace LA, Quinn RA, Duncan KR,

Hsu CC, Floros DJ, Gavilan RG, Kleigrewe K, Northen T, Dutton RJ, Parrot D, Carlson EE, Aigle

B, Michelsen CF, Jelsbak L, Sohlenkamp C, Pevzner P, Edlund A, McLean J, Piel J, Murphy BT,

Gerwick L, Liaw CC, Yang YL, Humpf HU, Maansson M, Keyzers RA, Sims AC, Johnson AR,

Sidebottom AM, Sedio BE, Klitgaard A, Larson CB, Boya PC, Torres-Mendoza D, Gonzalez DJ,

Silva DB, Marques LM, Demarque DP, Pociute E, O'Neill EC, Briand E, Helfrich EJ, Granatosky

EA, Glukhov E, Ryffel F, Houson H, Mohimani H, Kharbush JJ, Zeng Y, Vorholt JA, Kurita KL,

Charusanti P, McPhail KL, Nielsen KF, Vuong L, Elfeki M, Traxler MF, Engene N, Koyama N,

Vining OB, Baric R, Silva RR, Mascuch SJ, Tomasi S, Jenkins S, Macherla V, Hoffman T, Agarwal

V, Williams PG, Dai J, Neupane R, Gurr J, Rodriguez AM, Lamsa A, Zhang C, Dorrestein K,

Duggan BM, Almaliti J, Allard PM, Phapale P, Nothias LF, Alexandrov T, Litaudon M,

Wolfender JL, Kyle JE, Metz TO, Peryea T, Nguyen DT, VanLeer D, Shinn P, Jadhav A, Muller R,

Waters KM, Shi W, Liu X, Zhang L, Knight R, Jensen PR, Palsson BO, Pogliano K, Linington RG,

Gutierrez M, Lopes NP, Gerwick WH, Moore BS, Dorrestein PC, Bandeira N (2016) Sharing and

community curation of mass spectrometry data with Global Natural Products Social

Molecular Networking. Nat Biotechnol 34:828-837

44. Winnikoff JR, Glukhov E, Watrous J, Dorrestein PC, Gerwick WH (2014) Quantitative molecular

networking to profile marine cyanobacterial metabolomes. J Antibiot (Tokyo) 67:105–112

45. Wollenberg RD, Saei W, Westphal KR, Klitgaard CS, Nielsen KL, Lysoe E, Gardiner DM, Wimmer R,

Sondergaard TE, Sorensen JL (2017) Chrysogine Biosynthesis Is Mediated by a Two-Module

Nonribosomal Peptide Synthetase. J Nat Prod 80:2131–2135

46. Wu C, Kim HK, van Wezel GP, Choi YH (2015) Metabolomics in the natural products field - a gateway

to novel antibiotics. Drug Discov Today Technol 13:11-17

47. Yang JY, Sanchez LM, Rath CM, Liu X, Boudreau PD, Bruns N, Glukhov E, Wodtke A, de Felicio R,

Fenner A, Wong WR, Linington RG, Zhang L, Debonsi HM, Gerwick WH, Dorrestein PC (2013)

Molecular networking as a dereplication strategy. J Nat Prod 76:1686–1699

48. Yoon V, Nodwell JR (2014) Activating secondary metabolism with stress and chemicals. J Ind

Microbiol Biotechnol 41:415-424

49. Zhu H, Sandiford SK, van Wezel GP (2014) Triggers and cues that activate antibiotic production by

actinomycetes. J Ind Microbiol Biotechnol 41:371-386

50. Zhu H, Swierstra J, Wu C, Girard G, Choi YH, van Wamel W, Sandiford SK, van Wezel GP (2014)

Eliciting antibiotics active against the ESKAPE pathogens in a collection of actinomycetes

isolated from mountain soils. Microbiology 160:1714–1725

Table 1. ^{1}H and ^{13}C NMR data for compounds 1 and 2^a

NO.	1		2	
	δ_C	δ_H (mult., J in Hz)	δ_C	δ_H (mult., J in Hz)
2	79.0		73.3	
4	166.1		166.7	
4a	121.8		114.5	
5	128.8	7.88 (dd, J = 7.8, 1.2)	128.5	7.66 (dd, J = 7.8, 1.2)
6	123.0	7.07 (t, J = 7.8)	118.5	6.69 (td, J = 7.8, 1.2)
7	134.6	7.46 (td, J = 7.8, 1.2)	135.3	7.26 (td, J = 7.8, 1.2)
8	122.3	7.23 (brd, J = 7.8)	115.8	6.73 (brd, J = 7.8)
8a	143.7		148.6	
1'	73.9	4.43 (d, J = 9.0); 4.24 (d, J = 9.0)	64.5	3.67 (d, J = 10.8); 3.62 (d, J = 10.8)
2'	64.6	3.59 (d, J = 11.4); 3.47 (d, J = 11.4)	64.5	3.67 (d, J = 10.8); 3.62 (d, J = 10.8)
1"	65.7	3.86 (d, J = 12.6); 3.80 (d, J = 12.6)		
2"	102.9			
3"	61.9	3.66 (d, J = 3.0)		

494 ^{a} **1** and **2** were recorded in CD$_3$OD, at 298 K. All chemical shift assignments were done on the basis of

495 1D and 2D NMR techniques.

Figure Legends

Fig. 1. Analysis of secondary metabolites produced by *Streptomyces* sp. MBT27. The heat map depicts the relative abundance of the metabolites (rows) produced under different growth conditions (columns)

Fig. 2. Secondary metabolite profiles found in 1% and 2% glycerol-grown cultures of *Streptomyces* sp. MBT27. a. Thin layer chromatography of extracts from the cultures grown in MM with either 1% or 2% glycerol as the carbon source. **b.** Volcano plot highlighting the differences in metabolite profiles. Pink circles represent metabolites with an intensity difference of more than 4-fold (p-value ≤ 0.005)

Fig. 3. GNPS molecular network of the ions detected in the crude extract of *Streptomyces* sp. MBT27. Cultured were grown in MM with 2% glycerol, with an enlargement of the clusters including the dereplicated and identified metabolites. Green nodes represent the dereplicated metabolites, while blue nodes represent the masses upregulated when glycerol concentration was increased from 1% to 2%

Fig. 4. Chemical structures of quinazolinones A (1) and B (2).

Fig 5. Key COSY (▬) and HMBC (⌒) correlations for 1 and 2.

Fig. 6 Proposed biosynthetic pathway for quinazolinone A (1) and B (2).

520   **Fig. 1.**

523   **Fig. 2.**

527   **Fig. 3.**

529   **Fig. 4**

Quinazolinone A (**1**)   Quinazolinone B (**2**)

534   **Fig. 5**

1   2

539   **Fig. 6**

3   **4**   **2**   **1**