1 December 2018
Aaron Shafer
Trent University
Peterborough, Ontario Canada
K9J 7B8
aaronshafer@trentu.ca

RH: Breen et al. • Mountain goat genetic management

**Extensive field-sampling reveals the uniqueness of a trophy mountain goat population**

JESSICA BREEN* *Forensics Program, Trent University, K9J 7B8 Peterborough,*

*Canada*

MEGHAN BRITT* *Forensics Program, Trent University, K9J 7B8 Peterborough,*

*Canada*

JUSTIN B. JOHNSON* *Environmental and Life Sciences Graduate Program, Trent*

*University, K9J 7B8 Peterborough, Canada*

DARIA MARTCHENKO *Environmental and Life Sciences Graduate Program, Trent*

*University, K9J 7B8 Peterborough, Canada*

YASAMAN SHAKERI *Division of Wildlife Conservation, Alaska Department of Fish and*

*Game, Juneau, AK, USA*

BOYD PORTER *Division of Wildlife Conservation, Alaska Department of Fish and Game,*

*Ketchikan, AK, USA*

KEVIN S. WHITE *Division of Wildlife Conservation, Alaska Department of Fish and*

*Game, Juneau, AK, USA*

AARON B.A. SHAFER[1] *Forensic Science Program & Environmental and Life Sciences*

*Graduate*

*Program, Trent University, K9J 7B8 Peterborough, Canada*

**ABSTRACT** Collaborations between academic researchers and agencies is crucial for genetic data to have a tangible impact on conservation and wildlife management. Such partnerships are particularly important elusive species where the difficult terrain requires that a significant amount of resources and a combination of methods be used to estimate population parameters needed for conservation. We report and multi-year academic-agency collaboration on the North American mountain goat that used an extensive field sampling of genetic and phenotypic data to determine whether, and to what degree, genetic and phenotypic differences separate an isolated population of mountain goats on the Cleveland Peninsula form those in southeast Alaska. We observed significantly larger horns on the peninsula and the population appears demographically isolated. Isolation-by-distance accompanied by limited migration and low effective population size on the Cleveland Peninsula suggest this population will continue to lose genetic diversity. While the large horns of mountain goats have generated interest in re-opening mountain goat harvest on Cleveland Peninsula, our genetic data suggest this population is vulnerable to demographic and environmental perturbations and is unlikely to support a sustained harvest.

**KEYWORDS** Alaska, horn size, non-invasive sampling, population genetics,

______________________________

*Contributed equally

[1.] *Email* aaronshafer@trentu.ca

For the past decade the relevance of genetic data to conservation and management has been questioned. Vernesi et al. (2008) observed that few researchers used their data or methods to suggest practical solutions, and subsequent essays raised similar concerns and argued for more collaboration between academics and conservation practitioners (Hoban et al. 2013). Meta-analyses have revealed a positive relationship between the inclusion of non-academic co-authors and linking data to policy or management recommendations (Britt et al. 2018), notwithstanding a general long-term decline in implementation of research to management and conservation (Mair et al. 2018). Despite the gap between geneticists and on-the-ground practitioners (Shafer et al. 2015, Haig et al. 2016), headway for genetically-informed management can be achieved via academic partnering with non-academics and non-biologists (Britt et al. 2018, Mair et al. 2018). The benefit presumably stems from the sharing of non-overlapping resources, expertise, and knowledge.

The purchase of hunting licences in the United States grosses close to one billion dollars annually (US Fish And Wildlife Service 2017). The economic benefits derived from hunting often mean significant resources are made available to management and conservation of wildlife that are guided by the principles of the North American Model of Wildlife Conservation (Organ et al. 2012). In Alaska, approximately 500 mountain goats (*Oreamnos americanus*) are harvested each year with the majority of hunting concentrated in coastal areas. The mountain goat is characterized by a thick white coat with relatively short, black horns that have been used by indigenous groups for centuries and are sought after by hunters (Festa-Bianchet and Côté 2007). As such, the sustained persistence of healthy mountain goat populations entails both societal gain and enduring cultural significance.

The mountain goat's extreme habitat and sensitivity to anthropogenic disturbance poses a significant challenge relative to other species in terms of population monitoring (White et al. 2011) and sightability models are often implemented (Poole 2007, Rice et al. 2009). As a result, a combination of observational, GPS tracking, and genetic data have been used to fill in knowledge gaps regarding population patterns and processes (Mainguy et al. 2009, Shafer et al. 2012*a*). Molecular genetics, specifically non-invasive genetic monitoring methods, represents an important tool for obtaining accurate and cost-effective population data that is less invasive for the target species (Taberlet and Luikart 1999). Genetic tools allow for delineating populations boundaries and estimating parameters such as migration rates (i.e. gene flow), heterozygosity, and effective population size ($N_e$), all of which can inform management decisions. This is relevant to Alaska mountain goats as they are managed at the hunt unit level, not at the GMU (see Fig. 1), and hunt units are defined based on expert opinion and all available data.

The Cleveland Peninsula (Fig. 1) is an interesting case study involving mountain goat management as it is separated from the main mountain range by 32 km of low elevation terrain that is generally deemed unsuitable for the species (White et al. 2010). There are additional management and societal interests in the population due to disproprotionatley large horns, with four of the top ten Boone and Crockett records coming from the peninsula (Buckner et al. 2009; Boyd Porter, unpublished data). Based on the strict habitat requirements of mountain goats and the apparent geographic isolation it is currently unclear whether there is any mountain goat movement between the population(s) on the peninsula and those inhabiting the Coast Mountains in southeast Alaska.

In the 1980s there were an estimated 51 individuals across 9 suitable habitat patches found on the lower part of the Cleveland Peninsula (Smith and Raedeke 1982). In 2004, high

harvests and an apparent population decline led managers to close the area to hunting. In 2013-2014, only 4 patches were occupied with an estimated 40 individuals (K. S. White, unpublished data). As little is known about the population(s) on the peninsula, more information is required to guide management decisions, especially as it pertains to harvest. Specifically, knowledge about the degree to which the population(s) are isolated and how much genetic interchange is occurring is needed to assess the resilience of the population to stochastic events and the extent to which genetic rescue could aid the population in the event of natural or anthropogenic mediated declines. Here, we present a multiyear academic-agency collaboration that used an extensive field sampling effort combining genetic and phenotypic data to determine whether, and to what degree, genetic and phenotypic differences separate the population(s) on the Cleveland Peninsula from those found in southeast Alaska.

**METHODS**

**Sample Collection**

A pilot project was initiated in 2009 by the Alaska Department of Fish and Game (ADFG) to obtain precise population estimates for the Lower Cleveland Peninsula (Fig. 1), entailing the capture of 13 mountain goats. Mountain goats were fitted with radio-collars, phenotypic measurements were taken, and ear punches were obtained for DNA analysis. Tissue samples from mountain goats harvested from GMU 01 between 2005-2015 were also used for this analysis.

In the summers of 2014, 2016, 2017, and 2018, helicopter crews identified mountain goats on the Cleveland Peninsula and recorded their locations. Where individuals were sighted, a search was conducted on foot for fecal pellets. Non-invasive pellet samples were directly swabbed and placed in Longmire's solution. Samples from the Cleveland Peninsula were divided

into four sampling regions based on the geographic location of major ridges (Fig. 1); Lower Cleveland Peninsula North (LCP-N), Lower Cleveland Peninsula South (LCP-S), Middle Cleveland Peninsula (MCP), Upper Cleveland Peninsula (UCP). A transition zone Bradfield-Stikine (B-S) between the mainland was sampled, along with neighbouring GMUs 01A, 01B, and 01C.

**Horn Measurements**

Horn measurements from mountain goats from southeastern Alaska between 2005-2017 were collated and compared to captured mountain goats from the LCP. We evaluated horn growth between males in southeast Alaska by first generating length-by-age regression curves. A polynomial regression was fit with horn length as the response variable and location and age as explanatory variables, with a quadratic function applied to the age variable. We compared the annuli length at year 1 and 2 using a Student's t-test (note only the first two years were available for LCP). Analyses were conducted in R v. 3.4.1.

**DNA Extraction and Genotyping**

Genomic DNA was extracted from the tissue and the non-invasive fecal and hair samples with QIAGEN DNeasy Blood and Tissue Kits (Qiagen, Inc., Valencia, CA). For hair samples, a minimum of 10 hairs were placed root down in a 1.5mL tube. Fecal samples were swabbed with a cotton swab that had been wetted with de-ionized water and then placed downward in a 1.5mL tube. Ear punches and swabs that were collected in the field were placed wholly into 1.5mL tubes. All samples were digested overnight (~24 hours) in 180uL of Qiagen's Buffer ATL and 20uL of Pro-K before they were extracted following standard Qiagen protocol. Extractions were quantified using a Qubit 3.0 Fluorometer (ThermoFisher Scientific) and stored at -20° C.

A total of 18 polymorphic microsatellite loci were amplified in 3 PCR pools (Table S1). All non-invasive samples were amplified in triplicate. Each polymerase chain reaction contained approximately 2uL Taq buffer, 1uL BSA, 0.6uL $MgCl_2$, 0.2 dNTPs, <1uL each primer [0.05-0.25 uM], 0.1uL [5 units/uL] Taq polymerase, and 4uL $ddH_2O$. PCR conditions were: 94°C for 5 minutes, 30 cycles of 94°C for 30 seconds, 56°C for 60 seconds, 72°C for 60 seconds, followed by 60°C for 45 minutes. Samples were genotyped on an ABI 3730 DNA Analyzer and scored using the program Geneious R10 (Kearse et al. 2012). Non-invasive sample genotype calls for each locus were compared across three replicates and if there were disagreements (i.e. alleles were not the same), the peaks were compared visually, and the weak ones removed (RFU < 250). If more than one replicate at a marker had discrepancy between heterozygosity or homozygosity after three replicates, the marker was considered missing (scored 0). The resulting Cleveland Peninsula dataset was screened with the R package *Allelematch 2.03* (Galpern et al. 2012) to remove potential duplicate individuals. Individual samples and microsatellite loci with over 50% missing data were removed from analysis.

The number of alleles, heterozygosity, inbreeding and differentiation indices were determined using the program GenAlEx v6.5 (Peakall and Smouse 2006). We estimated $N_m$, or the effective number of migrants per generation between the sampled populations. We compared diversity statistics between the tissue and non-invasive samples from the LCP-S to determine whether sample quality influenced the results. All of our samples were collected within 15 years (1-3 generations) thereby reflecting a snapshot of population history; as such we estimated the effective population size ($N_e$) using a single point in time and linkage disequilibrium method available in NeEstimator V2.1 (Do et al. 2014). Isolation-by-distance was conducted by taking

the natural logarithm of the Euclidean distance between sampling area centroids and comparing this to $F_{ST}$. Significance was assessed using a Mantel test and 1,000 permutations (Mantel 1967).

The Bayesian clustering program STRUCTURE v.2.3.4 was then used to identify subpopulations (Pritchard et al. 2000). Admixture models within STRUCTURE were parallelized with StrAuto for K = 1-10 with a burn-in of $5x10^5$ followed by $1 \times 10^6$ MCMCs with 20 replicate runs for each K. The results of the runs for each K were sorted and averaged with CLUMPP 1.1.2 (Jakobsson and Rosenberg 2007). This process was completed once with the PopFlag parameter set to 1, which defines *a priori* population assignments for known groups, and once with the PopFlag parameter set to 0.  The assignment results were comparable with and without the PopFlag parameter, so we used our results without predefined reference groups to allow the software to find its own population divisions. We also used a principal component analysis (PCA) in the R package *adegenet* (Jombart 2008) to visualize population genetic structure and confirm the previous STRUCTURE results.

**Mitochondrial Sequencing and Analysis**

All samples from the LCP were sequenced at the mtDNA d-loop to confirm species identity as the population is sympatric with Sitka deer. A subset of MCP, UCP, and B-S samples were sequenced to include in the analysis. Samples were amplified following the approach of Shafer et al. (2011) with negative controls, according to the following conditions: 95°C for 5 minutes, 30 cycles of 95°C for 30 seconds, 53°C for 45 minutes, 72°C for 2 minutes, followed by 72°C for 5 minutes. Each polymerase chain reaction contained 10uL Taq buffer, 5uL BSA, 4uL $MgCl_2$, 1uL dNTPs, 0.4uL L15527, 0.4uL H00438, 0.25 uL [5 units/uL] Taq polymerase, and 35.95uL $ddH_2O$. The PCR product was run on a gel to ensure sufficient DNA, and then cleaned via ExoSAP-IT protocol (ThermoFisher Scientific) and ethanol precipitation prior to submission.

Samples were sequenced on an ABI 3730 DNA Analyzer, and edited and aligned with the program BioEdit (Hall 1999). For mtDNA analysis, 24 additional *Oreamnos americanus* d-loop sequences from southeast Alaska were obtained from GenBank to create a spatial outgroup. We quantified the number of unique haplotypes on the Cleveland Peninsula and plotted this using a haplotype network.

**RESULTS**

The regression analysis revealed significantly larger horns on the LCP relative to the mainland, with the polynomial model explaining a moderate amount of the total variation in horn length ($R^2 = 0.45$; Fig. 2). Growth during the first year was higher on the LCP compared to the mainland (18.3 cm $\pm$ 1.5 vs 16.4 cm $\pm$ 1.6; *t*-statistic 2.68, $p = 0.05$). A similar trend was observed in the second annuli (4.4 cm $\pm$ 0.6 vs 3.8 cm $\pm$ 3.2; *t*-statistic 1.46, $p = 0.18$).

A total of 9 days was spent helicopter sampling and a large portion of the peninsula was sampled (Fig. S1). We collected a total of 639 pellet groups and extracted 214 tissue samples from our mountain goat tissue repository. We note that preservation Longmire's solution resulted in more successful amplification compare to ethanol and dry storage. After data filtering (missing data, duplicates) a total of 468 samples were genotyped; 201 from the Mainland (tissue), 15 from the transition area Bradfield-Stikine (pellet), and 252 from the Cleveland Peninsula (239 pellet, 13 tissue). The data set was 84% complete (Dryad Accession no. XXXXX). All 18 loci had less than 50% missing data and were used.

Estimates of $F_{ST}$ between our samples from mainland (GMU 01A, 01B, 01C) and the Cleveland Peninsula, excluding B-S, ranged from 0.02-0.12 with and $N_m$ = approximately 3 migrants per generation (Table S2). Between the four grouped populations on the Peninsula and B-S, pairwise $F_{ST}$ ranged from 0.02-0.11 (Table S3), with B-S being the most differentiated.

Comparing non-invasive to the tissue samples on the LCP revealed similar diversity statistics (Table S4). Observed Heterozygosity ($H_O$) for 01A, B, and C ranged from 0.41 to 0.48 (Table 2). Within the groupings on the Cleveland Peninsula, $H_O$ ranged from 0.23 to 0.34, with B-S at 0.40; thus genetic diversity appears to decrease as mountain goat populations move down the peninsula, with inbreeding coefficients also becoming elevated (Table 2). There was a clear pattern of isolation-by-distance (Mantel $r = 0.60$; p < 0.01; Fig. 3). Our estimates of $N_e$ ranged from 5-15 individuals (Table 2) for the Lower Cleveland Peninsula using the lowest allele frequency. Estimates of $N_e$ for the total Mainland metapopulation ranged from 36-120 for the Mainland population (Table 2). There was no relationship between sample size and estimate of $N_e$ ($\beta = 0.28$; $p = 0.28$; Fig. S2).

We used the algorithm developed by Evanno et al. (Hall 1999) to identify mountain goat group clusters (K), implemented in the program STRUCTUREHARVESTER (Earl and vonHoldt 2012). Choice of appropriate number of clusters was informed by $\Delta(k)$, which identified either $k = 2$ as the optimal number of cluster groups in the genotypic data, which largely divided the Cleveland Peninsula from the mainland (Fig. 4; $k$ of 2-5 are shown). Notably a $k$ of 5 is consistent with the isolation by distance pattern, which is also seen in the PCA (Fig. S3).

We sequenced the mitochondrial DNA of 94 samples from the CP (GenBank Accession no. XXXXX). We add 38 samples from southeast Alaska that were available on GenBank. From our 94 mtDNA sequences from the CP we identified 21 unique haplotypes, with the majority found only on the peninsula and not shared with the mainland (Fig. 5)

**DISCUSSION**

The results of our analysis suggest that the subpopulation(s) on the Cleveland Peninsula are phenotypically and genetically differentiated from the adjacent GMU situated in southeast Alaska. Metrics of diversity suggest the Cleveland Peninsula is isolated with limited gene flow and is thus at risk for inbreeding depression and increased extinction risk (Caballero et al. 2017). Populations on the peninsula do appear interconnected, constituting a meta-population, with a pattern consistent with stepping-stone model exhibited by decreasing diversity and gene flow as you move down the peninsula. Collectively, this suggests the Cleveland Peninsula is largely isolated from the mainland and demographically distinct. How these data and patterns can inform actual management, however, is more nuanced.

**Interpreting genetic parameters for management**

A gap has emerged between genetic studies and on-the-ground conservation and management needs (Shafer et al. 2015, Taylor et al. 2017). While there are a plethora of reasons for this gap (see Taylor et al. 2017, Britt et al. 2018, Mair et al. 2018), part of the problem stems from the abstract and relative nature of many genetic parameters making them difficult to interpret and implement. General convention describes that $F_{ST} < 0.05$ constitutes little genetic difference, $F_{ST} = 0.05\text{-}0.15$ as moderate genetic difference, $F_{ST} = 0.15\text{-}0.25$ as large genetic differentiation, and $F_{ST} > 0.25$ as significant genetic difference (Hartl and Clark 1997). Thus, the Cleveland Peninsula population can be considered moderately differentiated from the mainland. We cannot, however, distinguish whether the population is recently isolated and there is currently no gene flow, meaning signatures of $N_m$ simply stem from recent common ancestry, or the CP mountain goats have been isolated for a long-time with periodic migration. The large number of individuals containing the CP unique mtDNA haplotypes would support the latter, further supporting the demographic uniqueness of this population.

Not only do the data support a demographically distinct population on the Cleveland Peninsula, but one that is vulnerable to environmental and demographic perturbations. The estimate $N_e$ is a key population genetic parameter described by Wright (1931) as the theoretical number of individuals in an ideal population that would lose genetic variation at the same rate as the observed population (i.e. the census size or $N_C$). Interpreting this parameter is difficult, as the approach reflects the parental generation's $N_e$, but can be impacted by changes further in the past (Waples 2005). Efforts to link $N_e$ to $N_C$ are desirable as they allow for going from a genetic parameter to the number of individuals on the landscape; however, $N_e/N_C$ ratios often span multiple orders of magnitude leading to high uncertainty (Palstra and Fraser 2012). Ortego et al. (2011) found the ratio of $N_e/N_C$ between 0.4 and 0.6 in mountain goat population found on Caw Ridge, Alberta. This seems reasonable for the lower Cleveland Peninsula given the $N_e$ of less than 15 and aerial survey estimates from the 1980's of 50-70 individuals, but the key take-home is reduced diversity on the LCP and UCP relative to the mainland.

The small effective population size and high inbreeding coefficient on the lower Cleveland Peninsula is an indication of a lack of genetic variability and may lead to reduced fitness of individuals, an increase of inbreeding, and further reduction in population size if not managed accordingly (Armbruster and Reed 2005). Habitat on the lower Cleveland Peninsula is atypical for mountain goats (Smith and Raedeke 1982), and it is unlikely sufficient suitable habitat and resources exist to accommodate more individuals. Without gene flow to the mainland it is reasonable to predict genetic diversity will continue decrease if population numbers do not change. Low $N_e$ combined with minimal migration to the mainland equates to reduced adaptive potential, making the population more susceptible to environmental and demographic perturbations than their mainland counterparts.

**MANAGEMENT IMPLICATIONS**

Given the geographic isolation, phenotypic differences, and the genetic distinctiveness reported here, this current harvest moratoria and separate hunting is warranted on the LCP (e.g. Moritz 1994, Fraser and Bernatchez 2001). These data suggest a sustainable harvest unlikely on the LCP given the genetic parameters and population numbers; if the population demographically recovers, any future harvest should be limited to adult males (Gonzalez-Voyer et al. 2003) and consideration should be given to alternating harvest years given the high extirpation risk (Hamel et al. 2006). The UCP and MCP are clearly more connected to the mainland, but it too is demographically distinct, and likely warrants a separated hunting unit status from the LCP. Given the inaccessible nature of the MCP and UCP a high impact of harvest seems unlikely, but it too should be regularly monitored. To prevent further loss of genetic variability in the region, translocating individuals from southeast Alaska to the Cleveland Peninsula is a management option. While this could impact the unique and desirable phenotype on the peninsula (Fig. 3), the translocation of even a few individuals carrying novel alleles could be enough to genetically support this population if needed (Hufbauer et al. 2015). That said, given the atypical habitat on the lower Cleveland Peninsula it is not clear that the region could even support the number of individuals that are desirable for a sustained harvest (see Hamel et al. 2006), and thus translocations should be considered a last resort.

**ACKNOWLEDGEMENTS**

We thank Sarah Haworth and Jesse Wolf help with sequencing. JB was supported by an EcoCanada Internship; MB by a NSERC undergraduate student research award and DM by an Ontario Graduate Scholarship. The project was supported by NSERC-DG, CFI-JELF and Compute Canada grants to ABAS; US Federal Aid grants to KSW and ABAS.

## LITERATURE CITED

Armbruster, P., and D. H. Reed. 2005. Inbreeding depression in benign and stressful environments. Heredity 95:235.

Britt, M., S. E. Haworth, J. B. Johnson, D. Martchenko, and A. B. A. Shafer. 2018. The importance of non-academic coauthors in bridging the conservation genetics gap. Biological Conservation 218:118–123.

Buckner, E. L., C. R. Byers, and J. Reneau. 2009. Measuring and scoring North American big game trophies. Boone and Crockett Club, Missoula. Montana, USA.

Caballero, A., I. Bravo, and J. Wang. 2017. Inbreeding load and purging: implications for the short-term survival and the conservation management of small populations. Heredity 118:177–185.

Do, C., R. S. Waples, D. Peel, G. M. Macbeth, B. J. Tillett, and J. R. Ovenden. 2014. NeEstimator v2: re-implementation of software for the estimation of contemporary effective population size (Ne) from genetic data. Molecular Ecology Resources 14:209–214.

Earl, D. A., and B. M. vonHoldt. 2012. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. Conservation Genetics Resources 4:359–361.

Evanno, G., S. Regnaut, and J. Goudet. 2005. Detecting the number of clusters of individuals using the software structure: a simulation study. Molecular Ecology 14:2611–2620.

Festa-Bianchet, M., and S. D. Côté. 2007. Mountain Goats: Ecology, Behavior, and Conservation of an Alpine Ungulate. None edition. Island Press, Washington, DC.

Fraser, D. J., and L. Bernatchez. 2001. Adaptive evolutionary conservation: towards a unified concept for defining conservation units. Molecular Ecology 10:2741–2752.

Galpern, P., M. Manseau, P. Hettinga, K. Smith, and P. Wilson. 2012. Allelematch: an R package for identifying unique multilocus genotypes where genotyping error and missing data may be present. Molecular Ecology Resources 12:771–778.

Haig, S. M., M. P. Miller, R. Bellinger, H. M. Draheim, D. M. Mercer, and T. D. Mullins. 2016. The conservation genetics juggling act: integrating genetics and ecology, science and policy. Evolutionary Applications 9:181–195.

Hall, T. A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Pages 95–98 *in*. Volume 41. [London]: Information Retrieval Ltd., c1979-c2000.

Hamel, S., S. D. Cote, K. G. Smith, and M. Festa-Bianchet. 2006. Population dynamics and harvest potential of mountain goat herds in Alberta. The Journal of wildlife management 70:1044–1053.

Hartl, D. L., and A. Clark. 1997. Principles of polulation genetics.

Hoban, S. M., H. C. Hauffe, Jan W. Arntzen, G. Bertorelle, Josef Bryja, K. Frith, O. E. Gaggiotti, P. Galbusera, J. A. Godoy, A. Rus, H. Richard, A. Nichols, C. R. Primmer, I. Russo, G. Segelbacher, H. R. Siegismund, M. Sihvonen, C. Vernesi, C. Vila, M. W. Bruford, J. W. Arntzen, and J. Bryja. 2013. Bringing genetic diversity to the forefront of conservation policy and management. Volume 5.

Hufbauer, R. A., M. Szűcs, E. Kasyon, C. Youngberg, M. J. Koontz, C. Richards, T. Tuff, and B. A. Melbourne. 2015. Three types of rescue can avert extinction in a changing environment. Proceedings of the National Academy of Sciences 112:10557–10562.

Jakobsson, M., and N. A. Rosenberg. 2007. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. Bioinformatics 23:1801–1806.

Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers. Bioinformatics 24:1403–1405.

Kearse, M., R. Moir, A. Wilson, S. Stones-Havas, M. Cheung, S. Sturrock, S. Buxton, A. Cooper, S. Markowitz, C. Duran, T. Thierer, B. Ashton, P. Meintjes, and A. Drummond. 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28:1647–1649.

Mainguy, J., S. D. Côté, and D. W. Coltman. 2009. Multilocus heterozygosity, parental relatedness and individual fitness components in a wild mountain goat, Oreamnos americanus population. Molecular Ecology 18:2297–2306.

Mair, L., A. C. Mill, P. A. Robertson, S. P. Rushton, M. D. F. Shirley, J. P. Rodriguez, and P. J. K. McGowan. 2018. The contribution of scientific research to conservation planning. Biological Conservation 223:82–96.

Mantel, N. 1967. The detection of disease clustering and a generalized regression approach. Cancer Research 27:209–220.

Moritz, C. 1994. Defining'Evolutionarily Significant Units' for conservation. Trends in ecology & evolution 9:373.

Organ, J., V. Geist, S. Mahoney, S. Williams, P. Krausman, G. Batcheller, T. Decker, R. Carmichael, P. Nanjappa, and R. Regan. 2012. The North American model of wildlife conservation. The Wildlife Society Technical Review 12.

Palstra, F. P., and D. J. Fraser. 2012. Effective/census population size ratio estimation: a compendium and appraisal. Ecology and Evolution 2:2357–2365.

Peakall, R., and P. E. Smouse. 2006. genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. Molecular Ecology Notes 6:288–295.

Poole, K. G. 2007. Does survey effort influence sightability of mountain goats Oreamnos americanus during aerial surveys? Wildlife Biology 13:113–119.

Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of Population Structure Using Multilocus Genotype Data. Genetics 155:945–959.

Rice, C. G., K. J. Jenkins, and W.-Y. Chang. 2009. A Sightability Model for Mountain Goats. The Journal of Wildlife Management 73:468–478.

Shafer, A. B. A., S. D. Côté, and D. W. Coltman. 2011. Hot Spots of Genetic Diversity Descended from Multiple Pleistocene Refugia in an Alpine Ungulate. Evolution 65:125–138.

Shafer, A. B. A., J. M. Northrup, K. S. White, M. S. Boyce, S. D. Côté, and D. W. Coltman. 2012*a*. Habitat selection predicts genetic relatedness in an alpine ungulate. Ecology 93:1317–1329.

Shafer, A. B. A., J. M. Northrup, K. S. White, M. S. Boyce, S. D. Côté, and D. W. Coltman. 2012*b*. Habitat selection predicts genetic relatedness in an alpine ungulate. Ecology 93:1317–1329.

Shafer, A. B. A., J. B. W. Wolf, P. C. Alves, L. Bergström, M. W. Bruford, I. Brännström, G. Colling, L. Dalén, L. De Meester, R. Ekblom, K. D. Fawcett, S. Fior, M. Hajibabaei, J. A. Hill, A. R. Hoezel, J. Höglund, E. L. Jensen, J. Krause, T. N. Kristensen, M. Krützen, J. K. McKay, A. J. Norman, R. Ogden, E. M. Österling, N. J. Ouborg, J. Piccolo, D. Popović, C. R. Primmer, F. A. Reed, M. Roumet, J. Salmona, T. Schenekar, M. K.

Schwartz, G. Segelbacher, H. Senn, J. Thaulow, M. Valtonen, A. Veale, P. Vergeer, N. Vijay, C. Vilà, M. Weissensteiner, L. Wennerström, C. W. Wheat, and P. Zieliński. 2015. Genomics and the challenging translation into conservation practice. Trends in Ecology & Evolution 30:78–87.

Smith, C. A., and K. J. Raedeke. 1982. Group size and movements of a dispersed, low density goat population with comments on inbreeding and human impact. Pages 54–67 *in*. Proceedings of the Biennial Symposium of the Northern Wild Sheep and Goat Council. Volume 3.

Taberlet, P., and G. Luikart. 1999. Non-invasive genetic sampling and individual identification. Biological Journal of the Linnean Society 68:41–55.

Taylor, H., N. Dussex, and Y. van Heezik. 2017. Bridging the conservation genetics gap by identifying barriers to implementation for conservation practitioners. Global Ecology and Conservation 10:231–242.

US Fish And Wildlife Service. 2017. WSFR Historical License Data. <https://wsfrprograms.fws.gov/subpages/licenseinfo/hunting.htm>. Accessed 9 Sep 2018.

Vernesi, C., M. W. Bruford, G. Bertorelle, E. Pecchioli, A. Rizzoli, and H. C. Hauffe. 2008. Where's the Conservation in Conservation Genetics? Conservation Biology 22:802–804.

Voyer, A. G., K. G. Smith, and M. Festa-Bianchet. 2003. Dynamics of hunted and unhunted mountain goat Oreamnos americanus populations. Wildlife Biology 9:213–218.

Waples, R. S. 2005. Genetic estimates of contemporary effective population size: to what time periods do the estimates apply? Molecular Ecology 14:3335–3352.

White, K. S., G. W. Pendleton, D. Crowley, H. J. Griese, K. J. Hundertmark, T. Mcdonough, L. Nichols, M. Robus, C. A. Smith, and J. W. Schoen. 2011. Mountain goat survival in coastal Alaska: Effects of age, sex, and climate. The Journal of Wildlife Management 75:1731–1744.

White, K. S., B. Porter, and S. Bethune. 2010. Cleveland Peninsula mountain goat population monitoring. Annual Progress Report, Alaska Department of Fish and Game.

Wright, S. 1931. Evolution in Mendelian populations. Genetics.

**FIGURES AND TABLES**

Figure 1.  Map showing management unit GMU 01 in southeast Alaska and the Cleveland Peninsula (a); map of the six sub-divisions in the Cleveland Peninsula (CP); Lower Cleveland Peninsula North (LCP-N), Lower Cleveland Peninsula South (LCP-S), Middle Cleveland Peninsula (MCP), Upper Cleveland Peninsula (UCP) and Bradfield-Stikine (BS)

Figure 2. Plot of the polynomial model explaining a total variation in horn length with age between the Cleveland Peninsula and samples from southeast Alaska.

Figure 3. Pattern of isolation-by-distance seen across the study area. Geographic distance is in kilometers and has be log-transformed.

Figure 4. Results of STRUCTURE analysis showing genetic subdivision at $K$ 2 through 5. A map showing population assignments is provided; Lower Cleveland Peninsula North (LCP-N), Lower Cleveland Peninsula South (LCP-S), Middle Cleveland Peninsula (MCP), Upper Cleveland Peninsula (UCP) and Bradfield-Stikine (BS) and management units 01A, 01B and 01C.

Figure 5. Haplotype network showing difference in mitochondrial sequence between the Cleveland Peninsula ($n$ = 95) and samples from the mainland ($n$ = 38). The key unique haplotypes unique to the Cleveland Peninsula are highlighted in red.

Table 1. Sample numbers from the Cleveland Peninsula and the mainland used in the genetic analysis.

| Population | Subdivision | Number of samples |
|---|---|---|
| Mainland | 01A | 60 |
| | 01B | 50 |
| | 01C | 91 |
| | Bradfield Stikine | 15 |
| Cleveland Peninsula | Upper | 36 |
| | Middle | 102 |
| | Lower North | 28 |
| | Lower South | 63 |

Table 2. Diversity statistics for the Mainland the Cleveland Peninsula. The effective population size estimate ($N_e$) reflects the upper and lower estimates.

| Population | Subdivision | No. of Alleles | Observed Heterozygosity | Fixation Index | $N_e$ estimate |
|---|---|---|---|---|---|
| Southeastern Alaska | 01A | 4.8 ± 0.61 | 0.41 ± 0.05 | 0.10 ± 0.04 | 36.0-58.1 |
| | 01B | 5.4 ± 0.58 | 0.46 ± 0.04 | 0.17 ± 0.04 | 81.4-119.4 |
| | 01C | 5.6 ± 0.54 | 0.48 ± 0.0.4 | 0.10 ± 0.03 | 64.3-95.7 |
| | Bradfield-Stikine | 3.1 ± 0.30 | 0.40 ± 0.05 | 0.08 ± 0.08 | 2.0-2.4 |
| Cleveland Peninsula | Upper | 3.9 ± 0.56 | 0.34 ± 0.04 | 0.30 ± 0.06 | 14.7-17.5 |
| | Middle | 4.8 ± 0.64 | 0.31 ± 0.04 | 0.37 ± 0.05 | 27.3-47.7 |
| | Lower North | 4.4 ± 0.46 | 0.23 ± 0.04 | 0.50 ± 0.07 | 7.3-14.9 |
| | Lower South | 4.3 ± 0.44 | 0.28 ± 0.04 | 0.38 ± 0.05 | 4.8-7.9 |

Horn Length between Cleveland and SE Alaska

K2 K3 K4 K5

01C (n = 91)
01B (n = 50)
01A (n = 60)
BS (n = 15)
UCP (n = 36)
MCP (n = 102)
LCP-N (n = 51)
LCP-S (n = 63)

K5 K4 K3 K2