A phosphate starvation response gene (*psr*1-like) is present and expressed in
***Micromonas pusilla* and other marine algae**

Cara L. Fiore[1,*], Harriet Alexander[2], Melissa C. Kido Soule, Elizabeth B. Kujawinski

Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic
Institution, Woods Hole, MA, USA

[1]Current address: Biology Department, Appalachian State University, Boone, NC, USA

[2]Current address: Department of Biology, Woods Hole Oceanographic Institution, Woods
Hole, MA, USA

*Corresponding author: Cara Fiore
572 Rivers St.
Boone, NC 28607, USA
fiorec@appstate.edu
15

Running title: Psr1 transcription factor in marine phytoplankton

Competing interest statement: The authors declare that they have no conflict of interest

Abstract

Phosphorus (P) limits primary production in regions of the surface ocean, and many

plankton species exhibit specific physiological responses to P-deficiency. The metabolic

response of *Micromonas pusilla*, an ecologically relevant marine photoautotroph, to P-

deficiency was investigated using environmental metabolomics and comparative

genomics. The concentrations of some intracellular metabolites were elevated in the P-

deficient cells (e.g., xanthine) and genes involved in the associated metabolic pathways

shared a significant motif in the non-coding regions of the gene. The presence of the

conserved motif suggests that these genes may be co-regulated, and the motif may

constitute a regulatory element for binding a transcription factor (i.e., Psr1). A putative

phosphate starvation response gene (*psr*1-like) was identified in *M. pusilla* with

homology to well characterized *psr*1/*phr*1 genes in freshwater algae and plants. This gene

appears to be present and expressed in other marine algal taxa, such as the abundant

haptophyte *Emiliania huxleyi*, in chronically phosphorus-limited field sites. Results from

the present study have implications for our understanding of taxon-specific roles in

mediating P cycling in the ocean.

Introduction

Phosphorus (P) is found in lipid membranes, genetic material, and energy storage compounds, making it a critical element for life. Because most marine microorganisms preferentially take up P as inorganic phosphate (PO_4^{3-}), concentrations of dissolved PO_4^{3-} are low in most of the surface ocean (<1 µM; ref. 1) thereby limiting phytoplankton productivity (2).

Marine microbes respond to chronic P-deficiency by investing resources in organic and inorganic P uptake (3-7), remodeling cellular metabolism and structures (8-13), and storing P (14,15). Different mechanisms for enacting these responses have evolved in widely distributed and often sympatric microbial taxa. For example, within the diatom, *Phaeodactylum tricornutum*, proteomics indicated broadly depressed metabolic activity as a response to P-starvation, with down-regulated energy metabolism, amino acid metabolism, nucleic acid metabolism, and photosynthesis, and up-regulated protein degradation, lipid accumulation, and photorespiration (16). In contrast, the prymnesiophytes *Prymnesium parvum* and *Emiliania huxleyi,* and the dinoflagellates *Prorocentrum donghaiense* and *Amphidinium carterae*, maintained energy-generating processes (i.e., photosynthesis) and carbon fixation in response to P-deficiency (17-20). Interestingly, *P. donghaiense* increased nitrate assimilation under P-deficiency (20), whereas *E. huxleyi* reduced nitrate uptake with a tight coupling between P and nitrogen (N) pools (19). These varied strategies all minimize non-critical P utilization and maximize P uptake, and thus play important roles in structuring phytoplankton assemblages and P biogeochemical cycling in the oceans (11, 19, 21-23).

Picoeukaryotes, including the abundant and cosmopolitan groups *Micromonas*

and *Ostreococcus* spp., are members of the "green" lineage (Chlorophyta) and are

important marine primary producers (24-26). Recent culture experiments with

Micromonas pusilla indicated that this group is likely to be successful under conditions

of increased acidification and low nutrient (P) availability, yet the mechanisms behind

their success are elusive (27). Similar to other phytoplankton (4), *M. pusilla* may remodel

lipids to increase non-P containing lipids (28) and up-regulate genes for P-transporters

and polyphosphate accumulation (29). However, the broader physiological response of *M.*

pusilla and other prasinophytes to P-deficiency has not been fully explored.

The green algae, Chlorophyta, share a more recent common ancestor with land

plants than other marine algal groups such as diatoms and haptophytes (30,31).

Consequently, their physiological response to P-deficiency may share more traits with

land plants than with other algal taxa. Indeed, P-deficiency is common in lakes and

terrestrial systems (32). The physiological response of model chlorophytes such as the

mustard plant, *Arabidopsis thaliana*, and the freshwater green alga, *Chlamydomonas*

reinhardtii, to P-deficiency led to the identification of a phosphate starvation response

gene described as *psr*1 in *C. reinhardtii* (33) and as *phr*1 in plants (34). This gene

encodes a transcription factor (TF), a protein that binds to specific DNA regions to

activate or repress transcription of one or more genes. The specific region of DNA to

which TFs bind is typically characterized by a short repetitive nucleotide sequence, or

motif, found up- or downstream of a given gene or within introns (35,36). In *A. thaliana,*

regulatory motifs for the Phr1 protein were more abundant in known P-responsive genes

than in the rest of the genome (37), linking Phr1 to the regulation of P-responsive genes.

Phr1/Psr1 TFs have not been described, to our knowledge, in marine algae.

Here, we investigate the impact of P-deficiency on the physiology of the

ecologically relevant picoeukaryote, *Micromonas pusilla* CCMP1545. Many metabolites

are produced by metabolic pathways under genetic regulatory control; thus, our combined

metabolomics and genomics approach provides mechanistic insight into the physiological

response of the organism to P-deficiency (e.g., 38, 39) and its underlying genetic

regulation. We used a targeted metabolomics approach to analyze the suite of molecules

produced by *M. pusilla* within the cells and in the external medium. In combination, we

employed a comparative genetics approach to identify 1) a *psr*1-like gene in *M. pusilla*

and other abundant marine phytoplankton species and 2) a potential regulatory element

that may be the DNA-binding site of the Psr1-like protein in putatively P-responsive

genes. We found evidence for the expression of *psr*1-like genes in *M. pusilla* and other

marine algae under P-deficient conditions in cultures and in the field (i.e., *Tara* Oceans

metatranscriptomes). These results have implications for better understanding and

predicting the metabolic response of diverse phytoplankton groups to P-deficiency.

Materials and Methods

Culture of Micromonas pusilla *CCMP 1545*

All glassware was acid-cleaned and combusted at 450°C for at least 4 h. We grew

an axenic culture of *Micromonas pusilla* CCMP1545 from the National Center for

Marine Algae and Microbiota (Boothbay, ME, USA) in L1-Si media

(https://ncma.bigelow.org/algal-recipes) with 0.2-μm filtered (Omnipore filters; Millipore,

MA, USA) autoclaved seawater from Vineyard Sound (MA, USA). We maintained the

cultures at 22°C under a 12:12 light:dark regime (84 μmol m^{-2} s^{-1}). After at least three

generations, we split the culture into two parallel cultures of L1-Si media amended with

(a) 36 µM phosphate (P-replete) and (b) 0.4 µM phosphate (P-deficient). We completed

three transfers of these parallel cultures prior to this experiment (Fig S1 a,b). For each

media type, we inoculated 9 flasks of media with exponential-phase cells to achieve 320

ml and ~300,000 cells in each flask (33 ml P-replete; 37 ml P-deficient), with three cell-

free control flasks. We grew cultures for two weeks (Fig S2a, b, c) and removed samples

approximately 1 hr into the light cycle at the time of experimental setup (T_0; day 0), in

rapid growth phase (T_1; day 4 P-deficient; day 6 P-replete), and in stationary phase (T_2;

day 5 P-deficient; day 13 P-replete).

At each sampling point, we removed 1 ml for cell counts and chlorophyll

fluorescence, 30 ml for total organic carbon (TOC; Methods S1), and 20 ml of filtrate

(see below) for nutrients. We monitored cell abundance daily by flow cytometry (Guava,

Millipore) and assessed photosystem II efficiency by measuring the variable and

maximum chlorophyll fluorescence (F_v/F_m) using fluorescence induction and relaxation

(FIRe; Satlantic LP, Halifax, Cananda). We used chlorophyll *a* (692 nm) and side scatter

of *M. pusilla* cultures to optimize settings for flow cytometry analysis. We assessed

potential bacterial contamination by viewing DAPI-stained cells of each time point

(Methods S1).

Metabolite extraction and instrument methods

We processed cultures for intracellular and extracellular metabolite extraction as

described previously (ref. 40; Methods S1). For intracellular metabolites, we collected

cells via filtration on 0.2 µm PTFE filters (Omnipore, Millipore) and stored them at -80°C.

Metabolites were extracted in 500 µl of cold 40:40:20 acetonitrile:methanol:water with

0.1 M formic acid. Extracts were reduced to near dryness under vacuum centrifugation

and reconstituted in 500 µl of 95:5 MilliQ water: acetonitrile with deuterated biotin (final

concentration 0.05 µg ml^{-1}) added as a high-performance liquid chromatography (HPLC)

injection standard; 100 µl of the extract was used for targeted metabolomics analysis. We

extracted extracellular metabolites using solid phase extraction (SPE) with 1 g/6 cc PPL

cartridges (BondElut, Agilent, Santa Clara, CA, USA) as described previously (41) and

modified by Longnecker (42) (Methods S1). Targeted metabolomics analysis was

performed as described in Kido Soule *et al*. (43). All metabolomics data are available in

the MetaboLights database (http://www.ebi.ac.uk/metabolights/) with accession number

MTBLS295.

R statistical software (v3.0.2; R Core Team) was used for statistical analyses. We

used a Welch's two sample *t*-test to compare the cell-normalized concentrations of

specific metabolites (log transformed) between P-deficient and P-replete treatments.

*Characterization of the phosphate starvation response (*psr*1) gene and DNA-binding*
motifs

We used the *psr*1 gene sequence from *C. reinhardtii* (ref. 33, NCBI

XM_001700501.1) to query the genome of *M. pusilla* using the BLAST option through

Integrated Microbial Genomes (IMG, blastx, e-value: 1e^{-5}). Conserved domains within

putative sequences were characterized using NCBI conserved domain search (CD-search).

We then used the putative *M. pusilla psr*1-derived amino acid sequence to query several

databases using BLAST (tblastn, e-value: 1e^{-5}), including: a re-assembly of the Marine

Microbial Eukaryote Transcriptional Sequencing Project (MMETSP) (44), NCBI non-

redundant protein sequences (nr), the NCBI metagenome proteins (env nr), the *Tara*

Oceans eukaryotic unigenes (45), and several eukaryotic algal genomes from IMG

(Methods S1). Visualization of the *Tara* Oceans queries were performed using the

basemap library in Python and custom Python scripts (Methods S1). Inspection of

sequences were performed with multiple sequences alignment with *C. reinhardtii psr*1-

derived amino acid sequence. Sequence alignment and phylogenetic analysis was

performed in MEGA 5.2.1 (46) using MUSCLE (47) with default settings in MEGA

(Methods S1).

We searched for a regulatory binding element, or motif, for the Psr1-like protein

within a now-archived gene model of *M. pusilla* CCMP1545

(FrozenGeneCatalog_20080206 (ver 1), Table S1), then again within the updated gene

model from Bachy *et al*. (29) (Table 1). Specifically, we analyzed gene sequences of

interest for a regulatory motif using the motif-based sequence analysis tools Multiple Em

for Motif Elicitation (MEME Suite) 4.10.1 (48). The genes of interest (Table 1) were

selected based on metabolites that were elevated or depressed in concentration in the P-

deficient cultures although not necessarily statistically different between treatments (i.e.,

malate, several amino acids, vitamins, Fig 2) and gene expression information from

Bachy *et al*. (29). These included genes related to central carbon metabolism, lipid

metabolism, vitamin metabolism (pantothenic acid, folic acid), and nucleotide

metabolism (aspartic acid). We also included genes within the carbohydrate and lipid

metabolic pathways based on published work (49, 50) as well as the gene for proline

oxidase (POX), a key enzyme up-regulated in *E. huxleyi* in response to N- and P-

deficiency (19, 51). After initial motif discovery, we optimized our putative motif by

varying discovery parameters such as motif width and number within genes of interest

and within genes that were not expected to be regulated in response to P-deficiency,

including asparagine synthetase (JGI gene ID: 9681548) and acetyltransferase like/FAD

linked oxidase (9687568)). We attempted to identify motifs using the motif comparison

tool (Tomtom; ref. 52) against the *A. thaliana* database (Protein Binding Microarray

(PBM) motifs, (36)) and significant matches were used to guide the optimization of motif

discovery settings and inclusive sequences.

Using the selected sequences and optimized parameters, we searched for motifs

that are overrepresented in the query sequences ("positive") relative to a set of

background sequences ("negative"), the latter we defined using the genome scaffolds.

These positive and negative sequences were used to create a position specific prior (PSP)

distribution file (48). Discriminative motif discovery used the PSP file and the set of *M.*

pusilla gene sequences of interest (parameters: minw = 5, maxw = 10, nmotifs = 3, mod =

oops (one occurrence per sequence)). Each gene sequence consisted of concatenated 500

nucleotides upstream and downstream of the gene as well as untranslated regions (UTRs)

and intronic regions, thereby excluding the coding regions.

Two recent publications provided transcriptomes for testing our hypotheses

regarding expected DE genes in P-deficient and P-replete conditions. Whitney & Lomas

(53) published transcriptomes of *M. commoda* (formerly *M. pusilla*) RCC299, while

Bachy *et al*. (29) published transcriptomes of *M. pusilla* CCMP1545. We utilized these

data to test our hypotheses based on results of our metabolomics analyses with *M. pusilla*

CCMP1545 and *in silico* genomic analysis. We performed the same protocol for motif

discovery and identification on gene sequences from *M. commoda* RCC299 (gene model:

Micromonas pusilla NOUM17/FrozenGeneCatalog_20090404) and CCMP1545 (gene

model: *Micromonas pusilla* CCMP1545/MBARI_models(ver_1)).

Results

Metabolic response of Micromonas pusilla *CCMP1545 to phosphorus deficiency*

M. pusilla grew in P-replete (0.45 day^{-1}) and P-deficient (0.03 day^{-1}) media (Fig

S2). Total organic carbon concentrations mirrored growth curves, increasing

substantially in the P-replete cultures, and increasing slightly in the P-deficient cultures

(Fig S2). Inorganic nutrient concentrations and photochemical efficiency generally

decreased over time in both treatments (Results S2).

Metabolites were generally less abundant under P-deficiency in the intracellular

(Fig 1) and extracellular (Fig S3) fractions, although several intracellular metabolites

were most abundant during rapid growth (T_1) under P-deficiency, including nucleobases,

amino acids, carbohydrates, and vitamins (Fig 1). Only a few intracellular metabolites

showed significantly different concentrations between treatments: (a) the nucleobase

xanthine, which was significantly elevated under P-deficiency (*t*-test, p $= 0.01$), and (b)

the amino acid derivatives, *N*-acetylglutamate and tryptamine, which were significantly

higher in P-replete samples (*t*-test, p $= 0.01$, p $= 0.02$, respectively) (Fig 1). We observed

significant decreases in the intracellular ratio of purine nucleosides to their nucleobases

under P-deficiency at T_1 (Fig 2). Intracellular malate, succinate, and citrate exhibited

divergent responses, although their abundances were not significantly different between

treatments (Fig 1). Malate was more abundant in the P-deficient cells, while succinate

concentrations were similar between treatments and replicates and citrate was detected in

only one of three replicates in each treatment (Fig 1). We noted similarly varied

responses to P-deficiency in the purine nucleosides xanthosine and inosine, with invariant

xanthosine abundances but higher average concentrations and variability of inosine (Fig

1).

Metabolite concentrations are, in part, determined by genetic regulation of the

enzymes producing or degrading them and we hypothesized that the varied metabolite

responses could provide insight into the regulation of these genes. The description of a

phosphate starvation response (*psr*1) gene containing a myb DNA-binding domain, led us

to investigate the presence of *psr*1, and a myb-like motif in genes that could be regulated

by the *psr*1 gene product in *M. pusilla* and other marine algae.

*Characterization of the phosphate starvation response (*psr*1) gene in marine algae*

We found four statistically-significant sequences (Table 2) in response to

querying the now-archived *M. pusilla* CCMP1545 genome with the *psr*1 gene from *C.*

reinhardtii (strain cc125, NCBI accession AF174532). Only one of these sequences (JGI

ID 61323) had the two characteristic myb domains of the *psr*1 gene in *C. reinhardtii*

(myb-like DNA-binding domain, SHAQKYF class (TIGR01557) and myb predicted

coiled-coil type transfactor, LHEQLE motif (pfam PF14379)) (Fig 3a). The glutamine-

rich regions, characteristic of TF activation domains, and putative metal binding domain

originally described for *psr*1 in *C. reinhardtii* (33) were not detected in the putative *M.*

*pusilla psr*1 gene, thus we refer to this as a *psr*1-like gene. Conserved domain analysis

(54) revealed two other domains in the putative *psr*1-like-derived amino acid sequence

from *M. pusilla*: 1) PLN03162 super family (NCBI cl26028), a provisional golden-2 like

transcription factor, and 2) a DUF390 super family domain (NCBI cl25642), comprising

proteins of unknown function.

We identified putative *psr*1-like genes or transcripts in field datasets (Fig 3b, 4, 5,

Table S2), representing diverse taxa such as Dinophyta (e.g., the symbiotic clade C

Symbiodinium sp.), and Haptophyta (e.g., the coccolithophore *E. huxleyi*). These

representative sequences did not necessarily cluster by taxonomic origin (e.g., cluster III,

Fig 4). Transcripts from MMETSP and *Tara* Oceans were generally short sequences and

often contained one of the two characteristic domains for *psr*1-derived amino acid

sequences (Results S2). Field-derived transcripts of the *psr*1-like gene were

geographically dispersed in the *Tara* Oceans dataset with the highest relative expression

in surface samples and in the North Atlantic Ocean and Mediterranean Sea (Fig 5a). The

majority of transcripts from the *Tara* Oceans dataset occurred in samples with low

phosphate concentrations (≤ 0.5 µM) but there was no significant correlation between

*psr*1 expression and phosphate concentration (Fig 5b). We observed a significant positive

relationship between the presence of *Micromonas psr*1-like genes and the relative

expression of one of the TCA cycle genes in *Micromonas,* fumarase, which were up-

regulated in both culture-based studies of *M. pusilla* (29) and *M. commoda* (53) under P-

deficiency (Fig 5c, Discussion S3). We never found genes with homology to the *M.*

pusilla putative *psr*1-like gene, in genomes or transcriptomes within diatoms, the

Heterokonta superphylum or the cryptophytes.

Putative regulatory element in Micromonas pusilla *genes of interest*

MEME analysis yielded a significant motif in *M. pusilla* genes involved in central

carbon metabolism, lipid metabolism, and nucleotide metabolism (Fig S4). The

conserved motif is similar in nucleotide sequence to several TF-binding sites in *A.*

thaliana (Table S3, Fig S5; 36), including myb-like TF-binding motifs. Several iterations

of this analysis using genes not tied to the metabolites of interest yielded either no

significant motif or motif sequences lacking similarity to myb-like domains. We observed

a similar conserved motif in DE transcripts in *M. pusilla* CCMP1545 under P-deficiency

(29) (Table 1, Fig S4). While the actual nucleotide sequence recognized by a TF must be

experimentally determined (36), the presence of a significant motif in each model and

only in the expected genes, indicates that this motif is likely biologically significant in *M.*

pusilla. Specifically, this motif, with similarity to myb-like domains in *A. thaliana*, may

represent a regulatory element where the Psr*1*-like protein binds.

Comparison of metabolomics-based predictions of P-responsive genes in M. pusilla

CCMP1545 to transcriptomics analysis

We queried the DE transcripts in *M. pusilla* CCMP1545 (29) and compared these to our

gene predictions based on our metabolomics data. As Bachy and colleagues (29) used a

more recent gene model than our original analysis, we confirmed the identity of the *psr*1-

like gene in the updated gene model (Fig S5). We analyzed a similar set of genes in

search of a regulatory element where the Psr1-like TF might bind (Fig S4), including DE

genes in the P-deficient transcriptome relative to the replete (Table S1 from 29). Under P-

deficiency, the *psr*1-like gene (JGI gene ID 360) was highly up-regulated (log-2 fold

change = 4.3) and nearly all of the predicted P-responsive genes were DE (Table 1). In

contrast to the *M. commoda* P-deficient transcriptome, *M. pusilla* exhibited many DE

genes in the tricarboxylic acid (TCA) cycle (Results S2) but not pyruvate carboxylase

(gene ID 6984). Additionally, between *M. pusilla* and *M. commoda*, different genes

involved in nucleotide and lipid metabolism were DE and contained a conserved motif

with similarity to myb-like regulatory element in *A. thaliana* (Fig 6). Lastly, the POX

gene observed in *M. commoda* and in the archived gene model for *M. pusilla*, is not

present, or at least not similarly annotated in the updated gene model. Instead, a copper

amine oxidase (gene ID 5863) contained a significant conserved motif (Table 1, Figure

S4) and is DE under P-deficiency.

Comparison to Micromonas commoda *RCC299 gene expression under P-deficiency*

We queried the DE genes in *M. commoda* RCC299 (26) exposed to P-deficient

conditions (53) for the *psr*1-like gene as well as for genes that we expected to be

regulated by the Psr1-like TF based on our *M. pusilla* observations and the literature (e.g.,

19, 55, 56). In *M. commoda*, the *psr*1-like gene (JGI gene ID: 60184) was significantly

up-regulated in the P-deficient treatment (Table S4), and we found a significant

conserved motif in 18 DE genes in *M. commoda*. The significant motif identified in *M.*

commoda differed in sequence from that discovered in *M. pusilla* and occurred in a

different set of genes. However, putative binding motifs for Psr1 are present in the same

pathways in the two *Micromonas* species and are similar to other DNA-binding motifs in

A. thaliana (Fig S6 Results S2; ref. 36). The *psr*1-like genes in RCC299 and CCMP1545

exhibited relatively low homology to each other but were equally similar to the *psr*1-

derived amino acid sequence in *C. reinhardtii* (Table S5).

Discussion

Metabolomics and comparative genomic analysis of the response of *M. pusilla*

CCMP1545 to P-deficiency has led to three conclusions: 1) there is a shift in intracellular

metabolite composition, 2) a *psr*1-like gene is expressed in *M. pusilla* and other marine

phytoplankton, and 3) the genes regulated by the Psr1-like protein appear to differ

amongst algal species.

Micromonas pusilla *exhibits a metabolic shift in response to P-deficiency*

Most intracellular metabolites did not differ significantly in concentration

between treatments. Because these metabolites represent many central metabolic

pathways, it appears that *M. pusilla* does not respond to P-deficiency through a global

decrease in metabolic activity, as was observed for a diatom (16). Interestingly, we found

examples of functionally-related metabolites that behaved differently. For example,

glutamate, proline, and methionine were variable but higher on average under P-

deficiency, while phenylalanine and tryptophan were not. In the TCA cycle, malate was

variable but higher in concentration on average under P-deficiency, while citrate was

lower and succinate was relatively unchanged. The opposing responses of these TCA

intermediates was surprising as others (e.g., 49, 57) have shown that transcripts or

proteins involved in the TCA cycle tend to exhibit a coordinated response to nutrient

limitation. Additional non-uniform metabolite responses occurred, with vitamins and

glucosamine-6-phosphate higher on average under P-deficiency, but other P-containing

compounds (e.g. glycerol-3-phosphate) and sugars were depleted. Previous work has

suggested that organisms placed under nutrient stress are capable of diverting carbon flux

between pathways (reviewed in 39). Changes in carbon flux through the cell could result

in coordinated abundance shifts in seemingly unrelated metabolites, as observed in this

study. If the shunting of carbon is a coordinated response within *M. pusilla*, then we

would expect genes involved in the synthesis and/or catabolism of these compounds to be

co-regulated, and potentially mediated by a specific transcription factor.

Presence and expression of psr*1 in marine algae*

We detected a putative *psr*1-like gene in the genome of *M. pusilla* and in the

genomes and transcriptomes of major algal lineages (e.g., prasinophytes, dinoflagellates).

A gene annotated as *psr*1 is present in *Ostreococcus tauri* (58), although to our

knowledge the potential role of this gene in the metabolism of *O. tauri* or other marine

algae has not been discussed. The *M. pusilla psr*1-like gene contains two myb domains

similar to those in *psr*1 from *C. reinhardtii* (33), and comparable to those in *phr*1 in *A.*

thaliana (34). No *psr*1-like gene was identified in queried diatom transcriptomes or

genomes, suggesting that a phosphate starvation response gene akin to *psr*1 is present in

diverse but phylogenetically constrained phytoplankton groups.

We found significant similarity between the *M. pusilla psr*1-like gene and

transcript sequences from the MMETSP and *Tara* Oceans datasets, indicating that this

gene is expressed by marine algae *in situ*. High-identity hits to sequences in the GOS and

Tara Oceans datasets underscore this gene's prevalence in the oceans, particularly in

regions characterized by chronically low phosphorus concentrations. Although some

MMETSP and *Tara* Oceans sequences contained only one (of two) characteristic Psr1

domains, these short sequences did not include enough of the C-terminal end to capture

the myb coiled-coil domain. Thus, further investigation is required to confirm whether

the identified transcripts are from *psr*1-like genes. Several GOS sequences, previously

described as protein of unknown function, had homology to both characteristic Psr1

domains, suggesting that they should be re-annotated as *psr*1-like genes. The GOS

sequences are likely derived from prasinophytes (i.e., *Micromonas* spp., *Ostreococcus*

spp.) and haptophytes (e.g., *E. huxleyi*), highlighting the need to elucidate the role of *psr*1

in these organisms.

Potential role of Psr1 in the marine algal response to P-deficiency

We observed a conserved and enriched motif that may function as a regulatory

element across genes involved in nucleotide biosynthesis, the TCA cycle and glycolysis,

carbon fixation, fatty acid metabolism, and phosphate transport or salvage. The conserved

motifs discovered in *M. pusilla* and *M. commoda* had unique sequences and were present

in a slightly different set of genes (Fig 6). The motif detected in *M. pusilla* is similar in

nucleotide sequence to a myb DNA-binding domain in *A. thaliana.* This similarity is

potentially significant, because the *psr*1/*phr*1 genes in *C. reinhardtii* and *A. thaliana*

contain a myb DNA-binding domain. Thus, we hypothesize that the conserved motif

detected in the *Micromonas* species represents binding sites for the Psr1-like transcription

factors (TF).

We observed the conserved motif only in DE genes under P-deficiency by each

species (29, 53), suggesting that these genes could be co-regulated. In bacteria, genes

within the well-characterized Pho-regulon, for example, all contain specific sequences

(the PHO box) where the transcription factor binds to activate or repress the gene (59).

Different bacterial species and strains contain distinct PHO box sequences. In *M.*

commoda, the discovered motif was not significantly similar to myb-like DNA-binding

motifs in *A. thaliana* and was not similar to the motif discovered in *M. pusilla*. This was

surprising at first, as we expected that the myb domain of the Psr1-like proteins in the

two species would interact with similar binding regions in the genome. However, *psr*1-

like derived amino acid sequences from *M. pusilla* and *M. commoda* were only 47%

similar. TF protein sequences with up to 79% similarity have been shown to have distinct

DNA-binding motif profiles (36), so these two proteins could reasonably have distinct

DNA-binding motifs. Moreover, as the conserved motif was present in different sets of

genes in each *Micromonas* species, it is likely that the Psr1-like protein regulates

different genes in different taxa and may serve as a niche defining feature between these

taxa. The observed variability in the *psr*1-like gene and the associated binding motifs

may be an adaptation to distinct environments as *M. pusilla* was collected in the

temperate English Channel, while *M. commoda* was isolated from the tropical Atlantic

(26).

The transcriptomes of *M. commoda* (53) and *M. pusilla* (29) provided a means to

test our hypotheses regarding the up- or down-regulation of certain genes in response to

P-deficiency. The *psr*1-like gene in *M. commoda* and *M. pusilla* was one of the highest

DE genes in each experiment (29, 53), suggesting that this potential TF plays a critical

role in the response of *Micromonas* to P-deficiency. Based on previous studies, we

expected genes involved in triacylglycerol (TAG) production to increase under P-

deficiency, but found divergent responses between the two species. Two TAG production

genes were up-regulated in *M. commoda*, including the penultimate step in TAG

production (putative phosphatidic acid phosphatase), which contained a putative

regulatory element where the Psr1-like TF could bind. By contrast, neither of the two

TAG metabolism genes that were DE in *M. commoda* were DE in *M. pusilla*; instead two

different genes involved in fatty acid biosynthesis and metabolism were DE in *M. pusilla*.

Additionally, a starch-binding protein was highly up-regulated in the *M. pusilla*

transcriptome under P-deficiency (gene ID 9633, ref. 29). These observations suggest that

M. pusilla may invest resources in starch storage while *M. commoda* may invest in lipid

storage, a testable hypothesis for future work.

Variable and non-uniform dynamics of several TCA cycle metabolites between

treatments suggested that this pathway may be involved in the metabolic response to P-

deficiency. Indeed, nearly all of the genes involved in the end of glycolysis and in the

TCA cycle were up-regulated in *M. pusilla* and contained the conserved motif that may

function as a regulatory element where the Psr1-like TF could bind. We propose that

carbon flow through the TCA cycle may increase as a means to fuel TAG or starch

production and to pull potentially damaging energy away from the photosystems (60, 61).

Further work is needed to test this hypothesis and to explore the physiological

implications for *Micromonas* spp. *in situ*. In contrast, only a few genes involved in the

TCA cycle were up-regulated in *M. commoda*, including genes involved in malate and

oxaloacetate (OAA) metabolism and a gene for pyruvate carboxylase (PC) which

converts pyruvate to oxaloacetate. The expression of these TCA-cycle genes may reflect

an increase in the malate-aspartate shuttle, a redox reaction that drives the production of

NAD^+ from NADH in the mitochondrial membrane. In *M. commoda*, it is possible that

PC-based conversion of pyruvate to OAA combined with an increase in the malate-

aspartate shuttle function as a stress response that allows continued glycolysis and ATP

production *via* oxidative phosphorylation (62, 63). In this scenario, carbon flow into the

TCA cycle would be limited and the genes controlling malate, aspartate, oxaloacetate and

pyruvate would be co-regulated. This hypothesis is supported by the observation of a

conserved motif that may function as a regulatory element in the relevant genes and by

preliminary gene expression analysis of one of the TCA cycle genes that was up-

regulated in both *M. pusilla* and *M. commoda* transcriptomes under P-deficiency, the

fumarase gene. If fumarase expression is regulated by the Psr1-like protein, expression of

these two genes (fumarase and *psr*1) should be correlated in field populations. We

observed support for this hypothesis within the *Tara* Oceans dataset, although we observe

weaker relationships between Psr1-like gene expression and other genes that were up-

regulated in either *M. pusilla* or *M. commoda* only (Discussion S3).

We observed shifts for several purine nucleosides between treatments for *M.*

pusilla and detected the conserved motif in the genes for a nucleoside phosphatase and

aspartate transcarbamylase. These results suggest increased nucleotide salvage through

purine nucleotides and nucleosides, a mechanism described in *M. commoda* (53) and in

other phytoplankton (13, 64). Interestingly, while a 5'-nucleotidase was observed to be

DE in *M. commoda* (53) under P-deficiency, it was not DE in *M. pusilla* (29). In the *M.*

pusilla transcriptome (29), a nucleotide phosphatase was up-regulated and may function

in a similar capacity in nucleotide salvage.

Another point of contrast between *M. pusilla* and *M. commoda* is the presence and

expression of a proline oxidase (POX) gene. The POX gene was highlighted by Rokitta *et*

al. (19,51) in nutrient limitation experiments (including P-deficiency) with the

haptophyte *E. huxleyi* where this enzyme may have a role in stabilizing the mitochondrial

membrane and in detecting cellular nitrogen levels. The POX gene was up-regulated in *M.*

commoda but not present in *M. pusilla*. By contrast, a copper amine oxidase was up-

regulated in *M. pusilla*. The POX gene in *M. commoda* and the copper amine oxidase

gene in *M. pusilla* contained the conserved motif that may function as a regulatory

element for the Psr1-like TF. The role of these genes in nutrient limitation should be

targets for further investigations (Discussion S3).

The potential impact of Psr1-regulated genes on the ecological roles of

Micromonas spp. is unknown, but our results suggest that the Psr1-like protein

coordinates a metabolic shift in these organisms under P-deficiency, altering the

intracellular flow of carbon and other elements. More comprehensive examination of

these metabolic responses, which likely vary to some extent among these organisms, will

be paramount to improving models of trophic carbon flow. More experiments are needed

to characterize the structure and role of the *psr*1-like gene in *Micromonas* spp. and other

phytoplankton, including: (1) confirming the presence of the *psr*1-like gene with genetic

experiments, (2) determining whether the Psr1-like protein is more abundant under P-

deficiency, (3) identifying the taxon-specific genes affected by the Psr1-like protein, (4)

verifying the interaction of the Psr1-like protein with the hypothesized binding sites, and

(5) comparing the genetic and metabolic responses to P-deficiency between organisms

containing the *psr*1-like and those that do not. Exploring the underlying biology of the

*psr*1 gene will facilitate mechanistic understanding of the complex metabolic response of

these organisms to P-limitation and will enhance our ecological and biogeochemical

predictions.

Acknowledgements

The authors would like to thank Krista Longnecker for performing the TOC analysis and
Matthew Johnson and Elizabeth Harvey for assistance with flow cytometry and FIRe
analysis. This research was funded by the Gordon and Betty Moore Foundation through
Grant GBMF3304 to E. Kujawinski.

References

1. Karl DM. Microbially mediated transformations of phosphorus in the sea: new views
of an old cycle. *Ann Rev Mar Sci.* 2014;6:279-337.

2. Tyrrell T. The relative influences of nitrogen and phosphorus on oceanic primary
production. *Nature* 1999;400:525–531.

3. Björkman KM, Karl DM. Bioavailability of dissolved organic phosphorus in the
euphotic zone at Station ALOHA, North Pacific Subtropical Gyre. *Limnol Oceanogr.*
2003;48:1049–1057.

4. Chung C-C, Hwang S-PL, Chang J. Identification of a high-affinity phosphate
transporter gene in a prasinophyte alga, *Tetraselmis chui*, and its expression under

nutrient limitation. *Appl Environ Microbiol.* 2003;69:754-759.

5. Teeling H, Fuchs BM, Becher D, Klockow C, Gardebrecht A, Bennke CM, Kassabgy
M, Huang S, Mann AJ, Waldmann J, Weber M, Klindworth A, Otto A, Lange J,
Bernhardt J, Reinsch C, Hecker M, Peplies J, Bockelmann FD, Callies U, Gunnar G,
Wichels A, Wiltshire KH, Glöckner FO, Schweder T, Amann R. Substrate-controlled
succession of marine bacterioplankton populations induced by a phytoplankton bloom.
Science 2012;336:608-611.

6. Dyhrman ST. Nutrients and their acquisition: phosphorus physiology in microalgae. In:
Borowitzka MA, Beardall J, and Raven JA (eds). *The Physiology of Microalgae*.
Heidelberg, Germany: Springer Publishing, 2016, pp155-183.

7. Lin S, Litaker RW, Sundra WG. 2016. Phosphorus physiological ecology and
molecular mechanisms in marine phytoplankton. *J Phycol.* 2016;52:10-36.

8. Berdalet E, Latasa M, Estrada M. Effects of nitrogen and phosphorus starvation on
nucleic acid and protein content of Heterocapsa sp. *J Plank Res.* 1994;16: 303–316.

9. Van Mooy BAS, Rocap G, Fredricks HF, Evans CT, Devol AH. Sulfolipids
dramatically decrease phosphorus demand by picocyanobacteria in oligotrophic marine
environments. *Proc Nat Acad Sci.* 2006;103:8607-8612.

10. Van Mooy BAS, Fredricks HF, Pedler BE, Dyhrman ST, Karl DM, Koblížek M, *et al.*
Phytoplankton in the ocean use non-phosphorus lipids in response to phosphorus scarcity.
Nature 2009;457:69–72.

11. Martin P, Van Mooy BA, Heithoff A, Dyhrman ST. Phosphorus supply drives rapid
turnover of membrane phospholipids in the diatom *Thalassiosira pseudonana*. *ISME J.*
2011;5:1057–1060.

12. Shemi A, Schatz D, Fredricks HF, Van Mooy BAS, Porat Z, Vardi A. Phosphorus
starvation induces membrane remodeling and recycling in *Emiliania huxleyi*. *New Phytol.*
2016;211:886-898.

13. Kujawinski EB, Longnecker K, Alexander H, Dyhrman ST, Fiore CL, Haley ST, *et al.*
Phosphorus availability regulates intracellular nucleotides in marine eukaryotic
phytoplankton. *Limnol Oceanogr Lett.* 2017;2:119-129. doi:10.1002/lol2.10043.

14. Orchard ED, Ammerman JW, Lomas MW, Dyhrman ST. Dissolved and organic
phosphorus uptake in Trichodesmium and the microbial community: The importance of
phosphorus ester in the Sargasso Sea. *Limnol Oceanogr.* 2010;55:1390-1399.

15. Martin P, Dyhrman ST, Lomas MW, Poulton NJ, Van Mooy BAS. Accumulation
and enhanced cycling of polyphosphate by Sargasso Sea phytoplankton in response to
low phosphorus. *Proc Nat Acad Sci.* 2014;111:8089-8094.

16. Feng T-Y, Yan Z-K, Zheng J-W, Xie Y, Li D-W, Bala Murugan S, Yang W-D, Liu J-
S, Li H-Y. Examination of metabolic responses to phosphorus limitation via proteomic

analyses in the marine diatom *Phaeodactylum tricornutum. Sci Rep*. 2015;5:10373.

17. Lai J, Yu Z, Song X, Cao X, Han X. Responses of the growth and biochemical
composition of *Prorocentrum donghaiense* to different nitrogen and phosphorus
concentrations. *J Exp Mar Biol Ecol*. 2015;405:6-17.

18. Li M, Shi X, Guo C, Lin S. Phosphorus deficiency inhibits cell division but not
growth in the dinoflagellate *Amphidinium carterae. Front Microbiol.* 2016;7:826.

19. Rokitta SD, von Dassow P, Rost B, John U. P- and N-depletion trigger similar
cellular responses to promote senescence in eukaryotic phytoplankton. *Front Mar Sci*
2016;3:109.

20. Shi X, Lin X, Li L, Palenik B, Lin S. Transcriptomic and microRNAomic profiling
reveals multi-faceted mechanisms to cope with phosphate stress in a dinoflagellate. *ISME*
J. 2017;11:2209-2218.

21. Cavender-Bares KK, Karl DM, Chisholm SW. Nutrient gradients in the western
North Atlantic Ocean: Relationship to microbial community structure and comparison to
patterns in the Pacific Ocean. *Deep Sea Res Part I: Oceanogr Res Pap.* 2001;48:2373–
2395.

22. Dyhrman ST, Ruttenberg KC. Presence and regulation of alkaline phosphatase
activity in eukaryotic phytoplankton from the coastal ocean: Implications for dissolved
organic phosphorus remineralization. *Limnol Oceanogr.* 2006;51:1381–1390.

23. Dyhrman ST, Benitez-Nelson CR, Orchard ED, Haley ST, Pellechia PJ. A microbial
source of phosphonates in oligotrophic marine systems. *Nat Geosci.* 2009;2:696–699.

24. Throndsen J, Kristiansen S. Micromonas pusilla (Prasinophyceae) as part of pico-and
nanoplankton communities of the Barents Sea. *Polar res.* 1991;10:201–208.

25. Not F, Latasa M, Marie D, Cariou T, Vaulot D, Simon N. A single species,
Micromonas pusilla (Prasinophyceae), dominates the eukaryotic picoplankton in the
Western English Channel. *Appl Environ Microbiol.* 2004;70: 4064–4072.

26. Worden AZ. Picoeukaryote diversity in coastal waters of the Pacific Ocean. *Aquat*
Microb Ecol. 2006;43:165–175.

27. Maat DS, Crawfurd KJ, Timmermans KR, Brussaard CPD. Elevated CO_2 and
phosphate limitation favor *Micromonas pusilla* through stimulated growth and reduced
viral impact. *Appl Environ Microbiol.* 2014;80:3119–3127.

28. Maat DS, van Bleijswijk JDL, Witte HJ, Brussaard PD. Virus production in
phosphorus-limited *Micromonas pusilla* stimulated by a supply of naturally low
concentrations of different phosphorus sources, far into the lytic cycle. *FEMS Microbiol*
Ecol. 2016;92:fiw136

29. Bachy C, Charlesworth CJ, Chan AM, Finke JF, Wong C-H, Wei C-L, *et al*.

Transcriptional responses of the marine green alga *Micromonas pusilla* and an infecting
prasinovirus under different phosphate conditions. *Environ Microbiol.* 2018;20:2898-
2912.

30. Falkowski PG, Katz ME, Knoll AH, Quigg A, Raven JA, Schofield O, *et al*. The
evolution of modern eukaryotic phytoplankton. *Science* 2004;16:354-360.

31. Lewis LA, McCourt RM. Green algae and the origin of land plants. *Amer J Bot.*
2004;91:1535–1556.

32. Elser JJ, Bracken MES, Cleland EE, Gruner DS, Harpole WS, Hillebrand H, *et al*.
Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater,
marine and terrestrial ecoystems. *Ecol Lett.* 2007;20:1135-1142.

33. Wykoff DD, Grossman AR, Weeks DP, Usuda H, Shimogawara K. Psr1, a nuclear
localized protein that regulates phosphorus metabolism in *Chlamydomonas. Proc Nat*
Acad Sci. 1999;96:15336–15341.

34. Rubio V, Linhares F, Solano R, Martín AC, Iglesias J, Leyva A, *et al*. A conserved
MYB transcription factor involved in phosphate starvation signaling both in vascular
plants and in unicellular algae. *Genes Devel.* 2001;15: 2122–2133.

35. Barrett LW, Fletcher S, Wilton SD. Regulation of eukaryotic gene expression by the
untranslated gene regions and other non-coding elements. *Cell Molec Life Sci.*
2012;69:3613-3634.

36. Franco-Zorrilla JM, López-Vidriero I, Carrasco JL, Godoy M, Vera P, Solano R.
DNA-binding specificities of plant transcription factors and their potential to define
target genes. *Proc Nat Acad Sci.* 2014;111:2367–2372.

37. Müller R, Morant M, Jarmer H, Nilsson L, Nielsen TH. Genome-wide analysis of the
Arabidopsis leaf transcriptome reveals interaction of phosphate and sugar metabolism.
Plant Physiol. 2006;143:156–171.

38. de Carvalho CCCR, Fernandes P. Production of metabolites as bacterial responses to
the marine environment. *Marine Drugs* 2010;3:705-727.

39. Markou G, Nerantzis E. Microalgae for high-value compounds and biofuels
production: a review with focus on cultivation under stress conditions. *Biotechn Advan.*
2013;31:1532-1542.

40. Fiore CL, Longnecker K, Kido Soule MC, Kujawinski EB. Release of ecologically
relevant metabolites by the cyanobacterium *Synechococcus elongatus* CCMP 1631.
Environ Microbiol. 2015;17:3949–3963.

41. Dittmar T, Koch B, Hertkorn N, Kattner G. A simple and efficient method for solid-
phase extracion of dissolved organic matter (SPE-DOM) from seawater. *Limn. Oceangr.*
Meth. 2008;6: 230–235.

42. Longnecker K. Dissolved organic matter in newly formed sea ice and surface
seawater. *Geochim Cosmochim Acta* 2015;171:39–49.

43. Kido Soule MCK, Longnecker K, Johnson WM, Kujawinski EB. Environmental
Metabolomics: Analytical Strategies. *Mar Chem.* 2015;177:1–62.

44. Johnson LK, Alexander H, Brown CT. Re-assembly, quality
evaluation, and annotation of 678 microbial eukaryotic reference transcriptomes. *Giga*
Sci. bioRxiv p 323576 doi:10.1101/323576 (In review).
604

45. Carradec Q, Pelletier E, Da Silva C, Alberti A, Seeleuthner Y, Blanc-Mathieu R, *et al*.
A global ocean atlas of eukaryotic genes. *Nature Comm.* 2017;9:373.

46. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular
Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and
Maximum Parsimony Methods. *Molec. Biol. Evol.* 2011;28: 2731–2739.

47. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucl Acids Res.* 2004;32:1792–1797.

48. Bailey TL, Elkan C. Fitting a mixture model by expectation maximization to discover
motifs in biopolymers. *Proceedings of the Second International Conference on Intelligent*
Systems for Molecular Biology, AAAI Press, Menlo Park, California, 1994, pp. 28-36,

49. Hockin NL, Mock T, Mulholland F, Kopriva S, Malin G. The response of diatom
central carbon metabolism to nitrogen starvation is different from that of green algae and
higher plants. *Plant Physiol.* 2012;158:299-312.

50. Goncalves EC, Johnson JV, Rathinasabapathi B. Conversion of membrane lipid acyl
groups to triacylglycerol and formation of lipid bodies upon nitrogen starvation in biofuel
green algae *Chlorella* UTEX29. *Planta* 2013;238:895-906.

51. Rokitta SD, von Dassow P, Rost B, John U. *Emiliana huxleyi* endures N-limitation
with an efficient metabolic budgeting and effective ATP synthesis. *BMC Genomics*
2014;15:1051.

52. Gupta S, Stamatoyannopolous JA, Bailey T, Noble WS. Quantifying similarity
between motifs. *Gen Biol.* 2007;8:R24.

53. Whitney LP, Lomas MW. Growth on ATP elicits a P-stress response in the
picoeukaryote *Micrmonas pusilla*. *PLoS ONE* 2016;11:e0155158.

54. Marchler-Bauer A, Bo Y, Han L, Lanczycki CJ, Lu S, Chitsaz F, *et al*.
CDD/SPARCLE: functional classification of proteins via subfamily domain architectures.
Nucl Acids Res. 2017;4:D200-D203.

55. Moseley JL, Chang C-W, Grossman AR. Genome-based approaches to understanding
phosphorus deprivation responses and PSR1 control in *Chlamydomonas reinhardtii*. *Euk*
Cell 2006;5:26–44.

56. Sharma KK, Schuhmann H, Schenk PM. High lipid induction in microalgae for
biodiesel production. *Energies* 2012;5:1532–1553.

57. Mock T, Samanta MP, Iverson V, Berthiaume C, Robison M, Holtermann K, *et al*.
Whole-genome expression profiling of the marine diatom *Thalassiosira pseudonana*
identifies genes involved in silicon bioprocesses. *Proc Nat Acad Sci*. 2008;105:1579-
1584.

58. Derelle E, Ferraz C, Rombauts S, Rouzé P, Worden AZ, Robbens S, *et al.* Genome
analysis of the smallest free-living eukaryote *Ostreococcus tauri* unveils many unique
features. *Proc Nat Acad Sci.* 2006;103:11647–11652.

59. Santos-Beneit F. The Pho regulon: a huge regulatory network in bacteria. *Front
Microbiol.* 2015;6:402.

60. Norici A, Bazonni AM, Pugnetti A, Raven JA, Giordano M. Impact of irradiance on
the C allocation in the coastal marine diatom *Skeletonema marinoi* Sarno and Zingone.
Plant Cell Environ. 2011;34:1666–1677.

61. Klok AJ, Martens DE, Wijffels RH, Lamers PP. Simultaneous growth and neutral
lipid accumulation in microalgae. *Biores technol.* 2013;134:233–243.

62. Yoshida K, Terashima I, Noguchi K. Up-regulation of mitochondrial alternative
oxidase concomitant with chloroplast over-reduction by excess light. *Plant Cell Physiol.*
2007;48:606-614.

63. Easlon E, Tsang F, Skinner C, Wang C, Lin S-J. The malate-aspartate NADH shuttle
components are novel metabolic longevity regulators required for calorie restriction-
mediated life span extension in yeast. *Genes Devel.* 2008;22:931-944.

64. Dyhrman ST, Palenik B. Characterization of ectoenzyme activity and phosphate-
regulated proteins in the coccolithophorid *Emiliania huxleyi*. *J Plank Res.* 2003;25:1215-
1225.

Figure Legends

Figure 1. Average ratio of the P-deficient to P-replete intracellular metabolite
concentrations during rapid growth (T_1) in *Micromonas pusilla* CCMP1545.
Concentrations are normalized to cell number in each treatment and means with one
standard deviation are shown (N=3 unless otherwise noted). Metabolites are listed in
order of descending concentration within three groups; those that were detected in
enough replicates for standard deviation to be calculated, those where only one ratio
could be calculated (black line), and those that were not detected (ND) in any of the three
replicates for one treatment (R = P-replete, D = P-deficient). Metabolite names marked
with an asterisk indicates a significant difference in concentration between treatments (*t*-
test, p < 0.05). Only one replicate in the P-deficient treatment contained a non-zero
concentration for *N*-acetylmuramic acid, thus while there was a significant difference in
concentration for this metabolite, only one ratio could be calculated. (DMSP =
dimethylsulfoniopropionate, NAD = nicotinamide adenine dinucleotide)

Figure 2. The intracellular ratios of two purine nucleosides to their nucleobases for
Micromonas pusilla CCMP1545 in P-deficient (orange) and P-replete (blue) treatments
during rapid growth (T_1). Xanthosine and xanthine, and guanosine and guanine were
quantified in the targeted metabolomics method and normalized to cell abundance. The
average ratio and high and low values are shown based on three replicate cultures. The
asterisk marks a significant difference between the treatments (*t*-test, p = 0.03). The
dashed line represents a ratio of 1. Two samples in the P-replete treatment had guanine
levels below the limit of detection and the one ratio is shown as a line.

Figure 3. Predicted amino acid sequences of the *psr*1-like gene in *Micromonas pusilla*
CCMP1545 and other organisms. The amino acid sequence (with IMG gene ID) is shown
as described in the Joint Genome Institute Integrated Microbial Genomes *M. pusilla*
CCMP1545 database. The myb-like DNA-binding and myb coiled-coiled domains are
highlighted in blue (a). Predicted amino acid sequence alignment of *psr*1 and *psr*1-like
genes in the region of the myb-like DNA-binding domain (SHAQKYF class) and
LHEQL coiled-coil domain (b). Asterisks at the top of each column indicate 100%
conserved residues across species surveyed. Numbers at the end of each row indicate
position of the last shown residue in the amino acid sequence. Accession (MMETSP,
NCBI, *Tara*) or gene identification (IMG) numbers are listed for each sequence.

Figure 4. Occurrence and phylogenetic relationship of *psr*1-like genes in marine
phytoplankton. Maximum likelihood tree based on the derived amino acid sequences for
*psr*1 and *psr*1-like genes from eukaryotic phytoplankton. The alignment for the
phylogenetic tree is based on 44 amino acid positions. Bootstrap values above 50% from
500 replications are shown for each node. Accession or IMG gene identification numbers
are shown with the taxon name and the star indicates *Micromonas pusilla* CCMP1545
cultured in this study. Four clusters are highlighted containing the Chlamydomonadales
(I), prasinophytes (II), dinoflagellates and other taxa (III), and the haptophyte, *Emiliania*
huxleyi and other taxa (IV). The outgroup used to root the tree was *Arabidopsis thaliana*
(NC_194590). The scale indicates the number of amino acid substitutions per site.

Figure 5. Geographic distribution of *Micromonas psr*1-like genes and relationship with
phosphate concentration in the *Tara* Oceans eukaryotic gene atlas (ref 45). The
geographic distribution and relative abundance of *Micromonas psr*1-like genes in surface
samples are depicted. Black dots represent stations where *Micromonas* was detected; the

size of purple circles represents the relative abundance of *psr*1-like gene, where the
FPKM of the psr1-like gene was normalized to the total FPKM of all *Micromonas-*
associated transcripts (a). Relative abundance of *Micromonas* is plotted against phosphate
concentrations for surface stations where *Micromonas* was detected. Grey circles indicate
samples where *psr*1-like genes were not detected, while samples where psr1-like genes
were detected are colored based on the relative expression of *Micromonas psr*1-like gene,
again normalized to total *Micromonas* transcript abundance (b). A split violin plot
depicting the distribution of expression of *Micromonas* fumarase synthase genes as a
function of presence or absence of *psr*1-like gene. Significant difference in the
distribution of expression between the presence and absence is assessed with
Kolmogorov-Smirnov test (p<0.005) (c).

Figure 6. Conceptual model of Psr1-like TF regulation in *Micromonas pusilla* CCMP154
(top half) and *M. commoda* RCC299 (bottom half) in three major metabolic pathways
under phosphorus deficiency. The top half of the figure illustrates the relevant
metabolites quantified in the present study and indicates if they are elevated (orange, ratio
> 1 Figure 2) or depressed (blue, ratio < 1 Figure 2) in concentration in the P-deficient
cells relative to the P-replete cells. Genes that contain the conserved motif with similarity
to the myb-like regulatory element in *Arabidopsis thaliana* are marked with a red star.
The orange and blue text refer to elevated or depressed transcript levels for each gene (ref
29). The same pattern applies to the bottom half of the figure with gene expression data
from Whitney & Lomas (53); metabolite data is not available for this organism.
Descriptions of putative metabolic responses in each pathway are shown and are based on
data in the present study and data from the literature ([1]ref 63; [2]ref 56). Genes encode for:
A) pyruvate kinase, B) pyruvate dehydrogenase, C) citrate synthase, D) aconitase, E) α-
ketoglutarate dehydrogenase, F) succinate dehydrogenase, G) fumarase, H) malate
dehydrogenase, I) pyruvate carboxylase, J) ribulose-1,5-bisphosphate carboxylase, K)
triose sugar transporter, L) nucleotide phosphatase, M) 5'-nucleotidase, N) glycerol
kinase, O) 1-acyl-sn-glycerol-3-phosphate acyltransferase, P) phosphatidate phosphatase,
Q) triacylglycerol lipase, LPAT = 1-acylglycerol-3-phosphate O-acyltransferase

750 **Tables**

751 Table 1. Enzymes that contain a significant motif in the gene sequences of *M. pusilla* CCMP1545 (MBARI_models (ver 1)). Column 2: the
752 Enzyme Commission (E.C.) number or pfam identifier; Column 3: main pathway(s) in which the enzyme is involved; Columns 4-6: fold
753 change and *q*-value of the transcript in P-deficient treatment relative to P-replete conditions from Bachy *et al*. (29) and corresponding gene ID.

Enzyme	E.C. Number	Pathway	Log 2 fold change	*q*-value	JGI gene ID
Glyceraldehyde-3-phosphate dehydrogenase	1.2.1.12	Calvin cycle/glycolysis	2.73	0.000037	4267
Glyceraldehyde-3-phosphate dehydrogenase	1.2.1.12	Calvin cycle/glycolysis	-0.644	0.000037	700
Phosphoglycerate kinase	2.7.2.3	Glycolysis	-0.699	0.000037	7661
Pyruvate kinase	2.7.1.40	Glycolysis	-1.32	0.000037	9370
Pyruvate dehydrogenase	1.2.4.1	TCA[a] cycle/Glycolysis	0.756	0.000037	3535
Citrate synthase	2.3.3.1	TCA cycle	1.68	0.000037	1278
Citrate synthase	2.3.3.1	TCA cycle	1.10	0.000037	3714
Aconitase	4.2.1.3	TCA cycle	1.00	0.000037	2901
Succinate dehydrogenase	1.3.5.1	TCA cycle and ETC[b]	1.16	0.001515	2182
Fumarase	4.2.1.2	TCA cycle	1.68	0.000037	600
Fumarase	4.2.1.2	TCA cycle	1.73	0.000068	6662
Copper amine oxidase	1.4.3.21-22	Arginine and proline metabolism	2.87	0.000037	5863
Aspartate transcarbamylase	2.1.3.2	Pyrimidine biosynthesis	-0.609	0.000037	3615
Nucleoside phosphatase	3.6.1.3	Pyrimidine metabolism	0.711	0.000037	3322
Pantoate-beta-alanine ligase	6.3.2.1	Pantothenate and CoA biosynthesis	-0.616	0.000037	5304
Lysophospholipase	3.1.1.5	Glycerophospholipid metabolism	1.65	0.000037	3652
Acyl-CoA synthetase	6.2.1.3	Fatty acid metabolism	3.20	0.000037	1751
Long-chain acyl-CoA synthetases	6.2.1.3	Fatty acid metabolism	0.736	0.000037	416
Na+/PO$_4$ transporter	PF02690	inorganic nutrient transport	4.54	0.000037	5963

754 [a]Tricarboxylic acid cycle

755 [b]Electron transport chain
756 [c]Triacylglycerol
757

Table 2. Gene matches to *psr*1 in *Micromonas pusilla* CCMP1545 as originally discovered in the Joint Genome Institute (JGI) Integrated
Microbial Genomes (IMG) database and the corresponding gene in the updated gene model for CCMP1545 (MBARI_models (ver 1)). The *M.*
pusilla CCMP 1545 database in IMG contained 10,660 sequences and 4,795,637 letters. The top-scoring gene was considered to be a *psr*1-
like gene and the JGI gene ID is shown in parentheses. The corresponding *psr*1-like gene in the updated gene model is shown for reference,
and includes the JGI gene ID, the gene description, and the E-value from the BLAST search (see Methods) using the top scoring gene
(613233) from CCMP1545.

IMG Gene ID	Locus tag	Bit score	E-value
2615011133 (JGI 61323)	MicpuC2.est_orfs.1_306_4269596:1	90.5	2×10^{-19}
2615008475	MicpuC2.EuGene.0000130210	74.3	4×10^{-14}
2615010887	MicpuC2.est_orfs.10_1861_4270447:1	70.5	3×10^{-13}
2615007841	MicpuC2.EuGene.0000090108	67.8	4×10^{-12}
JGI ID 360	Myb domain-containing protein		8×10^{-156}

A

Micromonas pusilla CCMP1545: 2615011133

MTRMSGIEDDSFSYSFLLISGQLRDPTRLTPRPQTSSAPLGAGGGAAAASGMNTSGTFWPSDLDLDLESSDFL
DSLLLGGEMMTTQQQHHHAGDERLGHGPVPLGARAAGEDHDPRVVLAERDAGDGLLPRALADERREPVAR
AVVASNNNKQRLRWTPELHKMFVDAVKRLGGLDLATPKGIMQLMDVEGMSIQHVKSHLQKYRLQDSGGG
ASEFRVSPDASASGKRPRSEEDDAGGNGKDGNNSAGKTRRRPSAAERSAARLRAAEEKAREERDAARSMAA
AAAAAAAAAVERQNVELALLTGDAAGARDALEMSSHHAASHVGGAYDDVLDLVAGPGESGDGDGAWGD
VLHDHAIVGAAGDDVGLVDDVGSDPEAAAAMLKQLELQKKLHEHLMSQRRLQQQVEAHGVYLETILDQQK
RRRGVE

B

		SHLQKyR		LHEQL	
Arabidopsis thaliana NC_003075.7	P K G V L K I M K V E - G - - - L T I Y H V K S H L Q K Y R	278		L H - E Q L	331
Volvox carteri XP_003054945.1	P K G I L K L M C L E - G - - - L T I Y H I K S H L Q K Y R	326		L H - E Q L	457
Micromonas pusilla CCMP 1545 2615011133 (IMG)	P K G I M Q L M D V E - G - - - M S I Q H V K S H L Q K Y R	205		L H - E H L	396
Ostreococcus lucimarinus 640420317 (IMG)	P K G I M H I M A M S - G - - - M T I Q H I K S H L Q K Y R	170		L H - D Q L	320
Micromonas sp. RCC 299 XP_002504000.1	P K G I M Q L M E V D - G - - - M T I Q H V K S H L Q K Y R	240		L H - A Q L	416
O. tauri 2507574032 (IMG)	P K G I A T L M T T S - G - - - M T L Q H I K S H L Q K Y R	106		L H - D Q L	241
GOS marine metagenome EBR92606.1	P K R I L D L M G V Q - G - - - L T R E N V A S H L Q K Y R	54		P D G G M M	201
GOS marine metagenome EBU22100.1	P K G I M H I M A M S - G - - - M T I Q H I K S H L Q K Y R	67		- - - - - -	84
GOS marine metagenome EBY01923.1	P K G I M S L M T T A - G - - - M T L Q H I K S H L Q K Y R	133		L H S M D A	223
Emiliana huxleyi XP_005765367.1	P Q A I S Q L M N C E - G E G A P T R Q N I K S H L Q K Y R	161		L E - - - -	212
GOS marine metagenome EDD34186.1	P Q A I R Q L M G C K T E E E A P T R Q N I K S H L Q K Y R	228		L H - E Q Q	327
Chlamydomonas reinhardtii NW_001843857.1	P K G I L K L M G V D - G - - - L T I Y H I K S H L Q K Y R	240		L H - E Q L	403
Chlorella variabilis 2507982799 (IMG)	P K R I L D L M N V E - G - - - L T R E N V A S H L Q K Y R	253		- - - - - -	334
MMETSP0052 *Polymella parva*	P K G I V T L M N V R - E - - - I T I Y H V K S H L Q K Y R	239		T A N A T L	425
MMETSP0063 *Chlamydomonas euryale*	P K K I L E I M Q V E - D - - - L T R E N I A S H L Q K Y R	293		G G - - - -	361
MMETSP1127 *Dunaliella tertiolects*	P K G I L K L V N S E - G - - - L T I Y H I K S H L Q K Y R	247		F Q G A G G	335
MMETSP1460 *Bathycoccus prassinos*	P K G V V E L M R V Q - G - - - V T I P H V K S H L Q K Y R	192		L H - E Q L	452
MMETSP1106 Mantoniella antarctica	P K G I M Q L M G V T - G - - - M T I Q H V K S H L Q K Y R	237		S A - A R I	332
MMETSP1403 *M. pusilla* CCMP 1723	P K G I M Q L M E V E - G - - - M T I Q H V K S H L Q K Y R	238		L H - E Q L	413
MMETSP0384 *Alexandrium tamarense*	P K T I L Q L M N V E - G - - - M T R E N V A S H L Q K Y R	97		M G - - - -	176
MMETSP1369 *Symbiodinium* sp. C1	P K N I M Q E M N V E - G - - - L T R E N V A S H L Q K Y R	113		- - - - - -	110
TARA MATOU-v1_62935935	P K G I M H I M A M S - G - - - M T I Q H I K S H L Q K Y R	57		- - - - - -	74
TARA MATOU-v1_45631074	P K G V V E L M R V Q - G - - - V T I P H V K S H L Q K Y R	88		L H - E Q L	348
TARA MATOU-v1_79753402	P K L I L R L L A V P - G - - - M T I Y H V K S H L Q K Y R	146		L Q - E Q L	243
TARA MATOU-v1_83305352	P K F I M R L L A V P - G - - - M T I Y H V K S H L Q K F R	87		L R - E Q L	158

A Normalized *psr1* abundance (proportion of *Micromonas* transcripts)

10^{-5} 10^{-4} 10^{-3}

B Normalized *psr1* abundance

10^{-5} 10^{-4} 10^{-3}

Normalized abundance of *Micromonas* (proportion per sample)

PO_4^{3-} (μM)

C

Gene abundance (proportion *Micromonas* transcripts)

Absent
Present

* $p < 0.005$

Fumarase Synthase

TCA cycle & Carbon fixation

M. pusilla CCMP1545: Targeted metabolomics and transcriptomic analysis

PEP
A ★
pyruvate
B ★
citrate
C ★
malate
D ★
G ★
fumarate
F ★
E ★
succinate
2-oxoglutarate
ribulose-1,5-phosphate
CO_2
J
glycerol-3-phosphate
K
triose sugar/ phosphate transport

Elevated in P-deficient
No significant difference
Depressed in P-deficient
Not quantified
★ Gene contains the conserved motif

Potential P-deficiency strategy: Decreased carbon fixation. Increased carbon flow through the TCA cycle to build metabolites for C, N, and P storage and salvage

M. commoda RCC299: Transcriptomic analysis

PEP
A ★
pyruvate
I ★
oxaloacetate
C
citrate
H ★
malate
G ★
succinate
ribulose-1,5-phosphate
CO_2
J
glycerol-3-phosphate
K
triose sugar/ phosphate transport

Potential P-deficiency strategy: Decreased carbon fixation. Increased carbon flow through the malate-aspartate shuttle with minimal increase in flow through the entire TCA cycle[1]

Purine nucleotide metabolism

ATP
L ★
AMP ⟷ adenosine ⟷ adenine
IMP ⟷ inosine ⟷ hypoxan-thine
XMP ⟷ xanthosine ⟷ xanthine
GMP ⟷ guanosine ⟷ guanine

Shift nucleosides to nucleobases

ATP
M ★
AMP ⟷ adenosine ⟷ adenine
★
IMP ⟷ inosine ⟷ hypoxan-thine
★
XMP ⟷ xanthosine ⟷ xanthine
★
GMP ⟷ guanosine ⟷ guanine

Shift nucleosides to nucleobases

Lipid metabolism

★ Acyl-CoA synthetase
★ Lysophospholipase

glycerol
fatty acid
glycerol-3-phosphate
1-acylglycerol-3-phosphate
monoacylglyceraol
1,2-diacylglycerol-3-phosphate
DAG → TAG

Maintain TAG metabolism and potentially increase some fatty acid and other lipid biosynthesis and salvage

★ LPAT

glycerol
fatty acid
N
glycerol-3-phosphate
Q
1-acylglycerol-3-phosphate
monoacylglycerol
O
1,2-diacylglycerol-3-phosphate
★ P
DAG → TAG

Increase TAG biosynthesis to serve as energy storage molecules[2]