A phosphate starvation response gene (*psr*1-like) is present and expressed in
***Micromonas pusilla* and other marine algae**

Cara L. Fiore[1,*], Harriet Alexander[2], Melissa C. Kido Soule, Elizabeth B. Kujawinski

Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic
Institution, Woods Hole, MA, USA

[1]Current address: Biology Department, Appalachian State University, Boone, NC, USA

[2]Current address: Department of Biology, Woods Hole Oceanographic Institution, Woods
Hole, MA, USA

*Corresponding author: Cara Fiore
572 Rivers St.
Boone, NC 28607, USA
fiorec@appstate.edu

Summary

- Phosphorus (P) limits primary production in regions of the surface ocean, and many plankton species exhibit specific physiological responses to P-deficiency. The metabolic response of *Micromonas pusilla*, an ecologically relevant marine photoautotroph, to P-deficiency was investigated using environmental metabolomics and comparative genomics.

- The concentrations of some intracellular metabolites were elevated in the P-deficient cells (e.g., xanthine, inosine) and genes involved in the associated metabolic pathways shared a predicted conserved amino acid motif in the non-coding regions of each gene. The presence of the conserved motif suggests that these genes may be co-regulated, and the motif may constitute a regulatory element for binding a transcription factor (i.e., Phosphate starvation response, Psr1, first described in the alga, *Chlamydomonas reinhardtii*).

- A putative phosphate starvation response gene (*psr*1-like) was identified in *M. pusilla* with homology to well characterized *psr*1/*phr*1 genes in algae and plants, respectively. This gene appears to be present and expressed in other marine algal taxa (e.g., *Emiliania huxleyi*) in field sites that are chronically phosphorus-limited.

- Results from the present study have implications for understanding phytoplankton taxon-specific roles in mediating P cycling in the ocean.

Introduction

Phosphorus (P) is a critical element for life, found in lipid membranes, genetic material, and energy storage compounds. Because most marine microorganisms preferentially take up P as inorganic phosphate (PO_4^{3-}), concentrations of dissolved PO_4^{3-} are generally below 1 µM in the surface ocean, (Karl, 2014) limiting phytoplankton productivity (Tyrell, 1999).

Marine phytoplankton respond to chronic P-deficiency by investing resources in P uptake via inorganic and organic compounds (Chung *et al*., 2003; Lin *et al*., 2016), remodeling cellular metabolism and structures (Berdalet *et al*., 1994; Van Mooy *et al*., 2009; Martin *et al*., 2011; Shemi *et al*., 2016; Kujawinski *et al*., 2017), and storing P (Dyhrman *et al*., 2012; Martin *et al*., 2014). Different mechanisms for enacting these responses have evolved in widely distributed and often sympatric phytoplankton taxa. For example, within the diatom, *Phaeodactylum tricornutum*, proteomics indicated broadly depressed metabolic activity as a response to P-starvation, with down-regulated energy metabolism, amino acid metabolism, nucleic acid metabolism, and photosynthesis, and up-regulated protein degradation, lipid accumulation, and photorespiration (Feng *et al*., 2016). In contrast, the prymnesiophytes *Prymnesium parvum* and *Emiliania huxleyi,* and the dinoflagellates *Prorocentrum donghaiense* and *Amphidinium carterae*, maintained energy-generating processes (i.e., photosynthesis) and carbon fixation in response to P-deficiency (Lai *et al*., 2015; Li *et al*., 2016; Rokitta *et al*., 2016; Shi *et al*., 2017). Interestingly, *P. donghaiense* increased nitrate assimilation under P-deficiency (Shi *et al*., 2017), whereas *E. huxleyi* reduced nitrate uptake with a tight coupling between P and nitrogen (N) pools (Rokitta *et al*., 2016). These varied strategies all minimize non-critical

P utilization and maximize P uptake, and thus play important roles in structuring

phytoplankton assemblages in the oceans (Dyhrman *et al*., 2006; Dyhrman *et al*., 2009;

Martin *et al*., 2011; Rokitta *et al*., 2016; Guo *et al*., 2018).

The cosmopolitan picoeukaryotic groups *Micromonas* and *Ostreococcus* spp. are

members of the "green" lineage (Chlorophyta), a monophyletic group including land

plants, and are important marine primary producers (Li,1994). Recent culture

experiments with *Micromonas pusilla* (Hoppe *et al*., 2018) and *M. pusilla* Lac38 (Maat *et*

al., 2014) indicated that these strains are likely to be successful under conditions of

increased acidification and low nutrient (P) availability. This success may in part, be due

to higher efficiency of carbon sequestration for growth under high pCO$_2$ conditions

and/or the ability to remodel photosystem proteins under nutrient-limited conditions

(Maat *et al*., 2014; Hoppe *et al*., 2018, Guo *et al*., 2018). Additionally, similar to other

phytoplankton (Chung *et al*., 2003), *M. pusilla* may shift lipid composition to increase

non-P containing lipids (Maat *et al*., 2016, Guo *et al*., 2018), and up-regulate genes and

proteins for P-transporters (Whitney & Lomas, 2016; Guo *et al*., 2018) and genes for

polyphosphate accumulation (Bachy *et al*., 2018). However, some of the broader

physiological response mechanisms of *M. pusilla* and other prasinophytes to P-deficiency

have yet to be fully explored.

The Chlorophyta share common ancestry with land plants, stemming from a

single endosymbiotic event, in contrast to more distantly related phytoplankton groups

such as diatoms and haptophytes, which are hypothesized to have undergone multiple

endosymbioses (Falkowski *et al*., 2004; Lewis & McCourt, 2004). Consequently, their

physiological response to nutritional stressors, like P-deficiency, may share more traits

with land plants than with other algal taxa. Indeed, P-deficiency is common in lakes and

terrestrial systems (Elser *et al*., 2007). The physiological response of model chlorophytes

such as the mustard plant, *Arabidopsis thaliana*, and the freshwater green alga,

Chlamydomonas reinhardtii, to P-deficiency led to the identification of a phosphate

starvation response gene described as *psr*1 in *C. reinhardtii* (Wykoff *et al*., 2009) and as

*phr*1 in plants (Rubio *et al*., 2001). This gene encodes for a transcription factor (TF), a

protein that binds to specific DNA regions to activate or repress transcription of one or

more genes. The specific region of DNA to which TFs bind is typically characterized by

a short repetitive nucleotide sequence, or motif, found up- or downstream of a given gene

or within introns (Barrett *et al*., 2012; Franco-Zorrilla *et al*., 2014). In *A. thaliana,*

regulatory motifs for the Phr1 protein were more abundant in known P-responsive genes

than in the rest of the genome (Müller *et al*., 2006), linking Phr1 to the regulation of P-

responsive genes. Phr1/Psr1 TFs have not been described, to our knowledge, in marine

algae.

Here, we use a combined metabolomic and comparative genomic approach to

investigate the impact of P-deficiency on the physiology of *Micromonas pusilla*

CCMP1545. Many metabolites are produced by metabolic pathways under genetic

regulatory control (e.g., de Carvalho & Fernandes, 2010; Markou & Nerantzis, 2013);

thus, our approach complements recent physiological, transcriptional, and proteomic

work (Hoppe *et al*., 2018; Guo *et al*., 2018) and provides mechanistic insight into the

physiological response to P-deficiency and its underlying genetic regulation. We used a

targeted metabolomics approach to analyze the suite of intracellular and extracellular

molecules produced by *M. pusilla*. In combination, we employed a comparative genetics

approach to identify 1) a *psr*1-like gene in *M. pusilla* and other prevalent marine

phytoplankton species and 2) a potential DNA-binding site of the Psr1-like protein in

putatively P-responsive genes. We found evidence for the expression of *psr*1-like genes

in *M. pusilla* and other marine algae under P-deficient conditions in cultures and in the

field. These results have implications for better understanding and predicting the

metabolic response of diverse phytoplankton groups to P-deficiency.

Materials and Methods

Culture of Micromonas pusilla *CCMP 1545*

118   All glassware was acid-cleaned and combusted at 450°C for at least 4 h. We grew

an axenic culture of *Micromonas pusilla* CCMP1545 from the National Center for

Marine Algae and Microbiota (Boothbay, ME, USA) in L1-Si media

(https://ncma.bigelow.org/algal-recipes) with 0.2-μm filtered (Omnipore filters; Millipore,

MA, USA) autoclaved seawater from Vineyard Sound (MA, USA). We maintained the

cultures at 22°C under a 12:12 light:dark regime (84 μmol m^{-2} s^{-1}). After acclimation to

these culture conditions, we split the culture into two parallel cultures of L1-Si media

amended with (a) 36 μM phosphate (P-replete) or (b) 0.4 μM phosphate (P-deficient). We

maintained these parallel cultures for three generations prior to this experiment (Fig. S1

a,b). At the start of the experiment, we inoculated 9 flasks of each media type with

exponential-phase cells to achieve 320 ml and ~300,000 cells in each flask (33 ml P-

replete; 37 ml P-deficient), with three cell-free control flasks. We grew cultures for two

weeks (Fig. S2a, b, c) and removed samples approximately 1 hr into the light cycle at

experiment initiation (T_0; day 0), in late exponential growth phase (T_1; day 4 P-deficient;

day 6 P-replete), and in stationary phase (T_2; day 5 P-deficient; day 13 P-replete).

At each sampling point, we removed 1 ml for cell counts and chlorophyll

fluorescence, 30 ml for total organic carbon (TOC; Methods S1), and 20 ml of filtrate

(see below) for nutrients. We monitored cell abundance daily by flow cytometry (Guava,

Millipore) and assessed photosystem II efficiency by measuring the variable and

maximum chlorophyll fluorescence (F_v/F_m) using fluorescence induction and relaxation

(FIRe; Satlantic LP, Halifax, Cananda). We used chlorophyll *a* (692 nm) and side scatter

of *M. pusilla* cultures to optimize flow cytometry settings. We assessed potential bacterial

contamination by viewing DAPI-stained cells at each time point (Methods S1).

Metabolite extraction and instrument methods

Metabolite extraction and analytical methods are optimized for semi-polar

compounds with a size range of 100-1000 Da, capturing many primary metabolites such

as organic acids, vitamins, and nucleosides. We processed cultures for intracellular and

extracellular metabolite extraction as described previously (Fiore *et al*., 2015; Methods

S1). For intracellular metabolites, we collected cells via filtration on 0.2 μm PTFE filters

(Omnipore, Millipore) and stored them at -80°C. Metabolites were extracted in 500 μl of

cold 40:40:20 acetonitrile:methanol:water with 0.1 M formic acid. Extracts were reduced

to near dryness under vacuum centrifugation and reconstituted in 500 μl of 95:5 MilliQ

water: acetonitrile with deuterated biotin (d2-biotin, final concentration 0.05 μg ml^{-1})

added as a high-performance liquid chromatography (HPLC) injection standard; 100 μl of

the extract was used for targeted metabolomics analysis. We extracted extracellular

metabolites using solid phase extraction (SPE) with 1 g/6 cc PPL cartridges (BondElut,

Agilent, Santa Clara, CA, USA) as described previously (Dittmar *et al*., 2008;

Longnecker, 2015).

Intracellular and extracellular extracts were used in targeted metabolomics

analysis with HPLC (Thermo PAL autosampler and Accela pump) coupled to tandem

mass spectrometry (TSQ Vantage, Thermo Fisher Scientific, MA, USA) via a heated

electrospray ionization (H-ESI) source operated in both positive and negative ion modes.

All 62 standards were analyzed using optimized selected reaction monitoring (SRM)

conditions (XCalibur 2.0) in the intracellular samples (Kido Soule et al., 2015), while 23

metabolites with high SPE recovery (Johnson *et al*., 2017) were analyzed in the

extracellular samples. Concentrations of metabolites were calculated based on a five-to-

seven point manually-curated external calibration curve (0.5-500 ng ml^{-1}). A pooled

sample for quality control consisted of 10 parts intracellular extract and one part

extracellular extract of each experimental sample and was analyzed every ten injections

(Methods S1). A laboratory-fortified pooled sample containing a standard mix was

analyzed at the start, middle, and end of the analytical run to assess matrix effects and

analytical quality. Chromatographic peaks derived from calibration standards, pooled

samples, and experimental samples were manually assessed for quality and metabolite

concentrations were exported to Excel. To facilitate comparison with other studies, we

also report cell-normalized ratios of the P-deficient to P-replete metabolite concentrations.

All metabolomics data are available in MetaboLights

(http://www.ebi.ac.uk/metabolights/) with accession number MTBLS295.

While both intracellular and extracellular metabolites were quantified in our

experiment, the results and discussion will focus primarily on changes in the intracellular

metabolite profile in response to P-deficiency, highlighting compounds that were most

relevant to the discovery of the *psr*1-like gene. We further focus the intracellular

metabolite comparison on the exponential growth phase (T_1) to capture clear changes in

metabolite profiles between treatments and facilitate comparison to previous studies

(Whitney & Lomas, 2016; Bachy *et al*., 2018). R statistical software (v3.0.2; R Core

Team) was used for statistical analyses. We used a Welch's two sample *t*-test to compare

the cell-normalized concentrations of specific metabolites (log transformed to achieve a

normal distribution) between P-deficient and P-replete treatments.

*Characterization of the phosphate starvation response (*psr*1) gene and putative DNA-*
binding motifs

We used the *psr*1 gene sequence from *C. reinhardtii* (Wykoff *et al*., 1999, NCBI

XM_001700501.1) to query the genome of *M. pusilla* using the BLAST option through

Integrated Microbial Genomes (IMG, blastx, e-value: $1e^{-5}$). Conserved domains within

putative sequences were characterized using NCBI conserved domain search (CD-search).

We then used the putative *M. pusilla psr*1-derived amino acid sequence to query several

databases using BLAST (tblastn, e-value: $1e^{-5}$), including: a re-assembly of the Marine

Microbial Eukaryote Transcriptional Sequencing Project (MMETSP) (Johnson *et al*.,

2018), NCBI non-redundant protein sequences (nr), the NCBI metagenome proteins (env

nr), the *Tara* Oceans eukaryotic unigenes (Carradec, Pelletier, *et al*., 2017), and several

eukaryotic algal genomes from IMG (Methods S1). For the *Tara* Oceans data, we

selected only the *psr*1-like unigenes from the metatranscriptome assemblies that were

assigned the taxonomic classification of *Micromonas*. Within each sample, we

normalized the expression of the *psr*1 unigenes to the total abundance of all unigenes

taxonomically classified as *Micromonas* (Methods S1). Visualization of the *Tara* Oceans

queries were performed using the basemap library in Python and custom Python scripts

(Methods S1). For several genes of interest, we compared the distribution of gene

expression to the expression of *Micromonas psr*1-like genes in the *Tara* Oceans

metatranscriptome using a Kolmogorov-Smirnov test. We tested a relationship between

phosphate concentration and normalized expression of *Micromonas psr*1-like genes from

Tara Oceans metatranscriptomes by comparing the abundance of *Micromonas psr*1-like

transcripts above and below the mean concentration of phosphate using a Student's *t*-test.

We manually checked significant matches for all nucleotide sequences that were

collected from database-based (NCBI, IMG, MMETSP, *Tara* Oceans) BLAST searches

using the *psr*1-like gene from *M. pusilla*. First, we used conserved domain analysis

(Marchler-Bauer *et al*., 2017) to assess domain similarity to *psr*1 and second, we used the

sequences with both characteristic domains of the *C. reinhardtii psr*1-derived amino acid

sequence in a multiple sequence alignment to the *C. reinhardtii psr*1-derived amino acid

sequence. Sequences that did not align well, and specifically did not align with the myb

coiled-coil domain, were removed from further consideration. Sequence alignment and

phylogenetic analysis was performed in MEGA 5.2.1 (Tamura *et al*., 2011) using

MUSCLE (Edgar, 2004) with default settings in MEGA. Phylogenetic reconstruction of

the *psr*1 gene was performed in MEGA using neighbor-joining, maximum parsimony,

and maximum likelihood algorithms and bootstrap replication 500 times. For visual

clarity, a subset of significant sequence matches was used to create the trees, including

representative sequences from each major taxonomic group (full tree in Supplementary

Information). The tree topology for the different phylogenetic reconstructions was similar

for nodes with high bootstrap values ($> 50\%$, data not shown). We present the maximum

likelihood tree because this algorithm (Jones-Taylor-Thornton model, default maximum

likelihood settings in MEGA) uses the most information in its alignments and typically

has high accuracy (Felsenstein, 1981; Kishino *et al*., 1990).

We searched genes of *M. pusilla* CCMP1545 for a regulatory binding element, or

motif, where the Psr1-like protein would bind to activate or repress gene transcription.

We first searched within a now-archived gene model of *M. pusilla* CCMP1545

(FrozenGeneCatalog_20080206 (ver_1), Table S1), then within the updated gene model

from Bachy *et al*. (2018) (*Micromonas pusilla* CCMP1545/MBARI_models(ver_1),

Table 1). Specifically, we analyzed gene sequences of interest for a regulatory element,

or motif, using the motif-based sequence analysis tools Multiple Em for Motif Elicitation

(MEME Suite) 4.10.1 (Bailey *et al*., 1994). The genes of interest (Table 1) were selected

based on metabolites that were elevated or depressed in concentration in the P-deficient

cultures though not necessarily statistically different between treatments (i.e., malate,

several amino acids, vitamins, Fig. 1), as well as gene expression information from

Bachy *et al*. (2018). These included genes related to central carbon metabolism, lipid

metabolism, vitamin metabolism (pantothenate, folate), and nucleotide metabolism

(aspartate). We included genes within the carbohydrate and lipid metabolic pathways

based on published work (Hockin *et al*., 2012; Goncalves *et al*., 2013) as well as the gene

for proline oxidase (POX), a key enzyme up-regulated in *E. huxleyi* in response to N- and

P-deficiency (Rokitta *et al*., 2014; Rokitta *et al*., 2016). After initial motif discovery, we

optimized our putative motif by varying motif width and number within genes of interest

and within genes that were not expected to be regulated under P-deficiency, including

asparagine synthetase (JGI gene ID: 9681548) and acetyltransferase like/FAD linked

oxidase (9687568)). We attempted to identify motifs using the motif comparison tool

(Tomtom; Gupta *et al*., 2007) against the *A. thaliana* database (Protein Binding

Microarray (PBM) motifs, (Franco-Zorrilla *et al*., 2014)) and significant matches were

used to guide the optimization of motif discovery settings and inclusive sequences.

Using the selected sequences and optimized parameters, we searched for motifs

that are overrepresented in the query sequences ("positive") relative to a set of

background sequences ("negative"), the latter defined using the genome scaffolds. We

used positive and negative sequences to create a position specific prior (PSP) distribution

file (Bailey *et al*., 1994), for further discriminative motif discovery within *M. pusilla*

gene sequences of interest (parameters: minw = 5, maxw = 10, nmotifs = 3, mod = oops

(one occurrence per sequence)). Each gene sequence consisted of concatenated 500

nucleotides upstream and downstream of the gene as well as untranslated regions (UTRs)

and intronic regions, thereby excluding the coding regions.

Two recent publications provided transcriptomes for testing our hypotheses

regarding expected differentially expressed (DE) genes in P-deficient and P-replete

conditions. Whitney & Lomas (2016) published transcriptomes of *M. commoda* (formerly

M. pusilla) RCC299, while Bachy *et al*. (2018) published transcriptomes of *M. pusilla*

CCMP1545, under P-deficient or limiting (respectively) and P-replete conditions. Genes

discussed as DE here were defined by each study (Whitney & Lomas, 2016; Bachy *et al*.,

2018). We note that the each *Micromonas* experiment used slightly different growth

conditions and were sampled at different stages of growth. Thus, we focus our

comparisons on work in the same strain (*M. pusilla* CCMP1545, Bachy *et al*., 2018), with

brief complementary discussions about *M. commoda* (see Notes S1 for further details).

Bachy *et al.* (2018) used semi-continuous cultures grown under continuous light and

were grown with a moderately higher concentration of P (2.5 µM) than in our experiment

(0.4 µM P). Despite these differences, the transcriptional data of *M. pusilla* under P-

limitation provided a preliminary test of our hypotheses derived from the metabolomics

analyses with *M. pusilla* CCMP1545 and *in silico* genomic analysis. We further reasoned

that a signal observed across species and experiments would likely represent a conserved

and biologically meaningful response to P-deficiency.

Results

Metabolic response of Micromonas pusilla *CCMP1545 to phosphorus deficiency*

280   Metabolite concentrations were generally lower under P-deficiency in the

extracellular (Fig. S3) fraction relative to the P-replete cultures, while there was a mixed

response in metabolite concentrations in the intracellular fraction (Fig. 1). The

extracellular metabolite concentrations represent cumulative changes in the media over

time. In stationary phase, nearly all of the quantified extracellular metabolites were less

abundant in the P-deficient samples than the P-replete samples (Fig. S3). In contrast,

several of the intracellular metabolites were more abundant during exponential growth

(T_1) under P-deficiency, including nucleobases (xanthine, adenine), the nucleoside

inosine, amino acids (glutamine, aspartate), the dicarboxylic acid malate, and vitamins

(thiamin, pantothenate, folate) (Fig. 1). None of the extracellular metabolites and only a

few intracellular metabolites showed statistically significant differences between

treatments, including: (a) the nucleobase xanthine, significantly elevated under P-

deficiency (*t*-test, p = 0.01), and (b) the amino acid derivatives, *N*-acetylglutamate and

tryptamine, significantly higher in P-replete samples (*t*-test, p = 0.01, p = 0.02,

respectively) (Fig. 1). We observed decreases in the intracellular ratios of purine

nucleosides (guanosine, xanthosine) to their nucleobases (guanine, xanthine) under P-

deficiency at T_1 (Fig. 2). The decrease in the xanthosine:xanthine ratio was statistically

significant (*t*-test, p = 0.03). The change in the guanosine:guanine ratio could not be

statistically tested as guanine was below detection in two of three P-replete cultures (Fig.

2). Intracellular malate, succinate, and citrate exhibited divergent responses, although

their abundances were not significantly different between treatments (Fig. 1). Malate was

more abundant in P-deficient cells, while succinate concentrations were similar between

treatments and replicates and citrate was detected in only one of three replicates in each

treatment (Fig. 1). We noted similarly varied responses to P-deficiency in the purine

nucleosides xanthosine and inosine, with invariant xanthosine abundances but higher

average concentrations and variability of inosine (Figs. 1,2).

306   Metabolite concentrations are, in part, determined by genetic regulation of the

enzymes producing or degrading them and we hypothesized that the varied metabolite

responses could provide insight into the regulation of these genes. The description of a

phosphate starvation response (*psr*1) gene containing a myb DNA-binding domain

(Wykoff *et al*., 1999), motivated us to investigate the presence of *psr*1 and a myb-like

motif in genes that could be regulated by the *psr*1 gene product in *M. pusilla* and other

marine algae.

*Characterization of the phosphate starvation response (*psr*1) gene in marine algae*

314   We found four statistically-significant sequences (Table 2) in response to

querying the now-archived *M. pusilla* CCMP1545 genome with the *psr*1 gene from *C.*

reinhardtii (strain cc125, NCBI accession AF174532). Only one of these sequences (JGI

ID 61323) had the two characteristic myb domains of the *psr*1 gene in *C. reinhardtii*

(myb-like DNA-binding domain, SHAQKYF class (TIGR01557) and myb predicted

coiled-coil type transfactor, LHEQLE motif (pfam PF14379)) (Fig. 3a). The glutamine-

rich regions, characteristic of TF activation domains, and putative metal binding domain

originally described for *psr*1 in *C. reinhardtii* (Wykoff *et al*., 1999) were not detected in

the putative *M. pusilla psr*1 gene, thus we refer to this as a *psr*1-like gene. Conserved

domain analysis (Marchler-Bauer *et al*., 2017) revealed two other domains in the putative

*psr*1-like-derived amino acid sequence from *M. pusilla*: 1) PLN03162 super family

(NCBI cl26028), a provisional golden-2 like transcription factor, and 2) a DUF390 super

family domain (NCBI cl25642), comprising proteins of unknown function. A comparison

of the psr1-like gene in *M. pusilla*, *M. commoda*, *C. reinhardtii*, and *A. thaliana* is

provided in Table S2.

We identified putative *psr*1-like genes or transcripts in cultured isolates as well as

field datasets (Figs. 3b, 4, 5, Table S3), representing diverse taxa such as Dinophyta (e.g.,

the symbiotic clade C *Symbiodinium* sp.), and Haptophyta (e.g., the coccolithophore *E.

huxleyi)*. In our phylogenetic analysis (Fig. 4; see Fig. S3 for full tree and Notes S1),

representative sequences clustered by taxonomic origin only to some extent. For example,

the prasinophytes, including *M. pusilla* CCMP1545, generally clustered together, as did

the haptophytes, including *E. huxleyi*. However, other clusters included a mix of

taxonomic groups such as the dinoflagellate *Alexandrium tamarense*, the glaucophyte

Gloeochaete wittrockiana, and the green alga *Chlorella vulgaris* (Fig. 4).

Transcripts from MMETSP and *Tara* Oceans were generally short sequences and

often contained one of the two characteristic domains for *psr*1-derived amino acid

sequences (Notes S1). Field-derived transcripts of the *psr*1-like gene were geographically

dispersed in the *Tara* Oceans dataset with the highest relative expression in surface

samples and in the North Atlantic Ocean and Mediterranean Sea (Fig 5a). The majority of

these transcripts from the *Tara* Oceans dataset occurred in samples with low phosphate

concentrations (≤ 0.5 μM), but there was no significant relationship between *psr*1

expression and phosphate concentration (*t*-test, $p < 0.05$; Fig. 5b). We observed a

significant positive relationship (Kolmogorov-Smirnov, $p < 0.05$) between the presence

of *Micromonas psr*1-like genes and the relative expression of fumarase, a TCA cycle

gene in *Micromonas*, which was up-regulated in cultures of *M. pusilla* (Bachy *et al*.,

2018) and *M. commoda* (Whitney & Lomas, 2016) under P-deficiency (Fig. 5c, Notes

S2). We did not find genes with significant homology to the *M. pusilla* putative *psr*1-like

gene, in genomes or transcriptomes, within diatoms, the Heterokonta superphylum or the

cryptophytes.

Putative regulatory element in Micromonas pusilla *genes of interest*

MEME analysis yielded a significant motif in *M. pusilla* genes involved in central

carbon metabolism, lipid metabolism, and nucleotide metabolism (Fig. S5). The

conserved motif is similar in nucleotide sequence to several TF-binding sites in *A.*

thaliana (Table S4, Fig. S6; Franco-Zorrilla *et al*., 2014), including myb-like TF-binding

motifs. The presence of a myb-like TF-binding motif is notable as the Psr1 protein

contains a myb-like DNA binding domain (Wykoff *et al*., 1999). Several iterations of this

analysis using genes not tied to the metabolites of interest yielded either no significant

motif or motif sequences lacking similarity to myb-like domains. We observed a similar

conserved motif in author-defined DE transcripts (Bachy *et al*., 2018) in *M. pusilla*

CCMP1545 under P-deficiency (Table 1, Fig. S5). While the actual nucleotide sequence

recognized by a TF must be experimentally determined (Franco-Zorrilla *et al*., 2014), the

presence of a significant motif in each model and only in the expected genes, indicates

that this conserved motif is likely biologically significant in *M. pusilla*. Specifically, this

motif, with similarity to myb-like domains in *A. thaliana*, may represent a regulatory

element where the Psr1-like protein binds.

Comparison of metabolomics-based predictions of P-responsive genes in M. pusilla

CCMP1545 to transcriptomics analysis

We compared the author-defined DE transcripts in *M. pusilla* CCMP1545 to gene

predictions based on our metabolomics data. As Bachy and colleagues (2018) used a

more recent gene model than our original analysis, we confirmed the identity of the *psr*1-

like gene in the updated model (CCMP1545/MBARI_models(ver_1), Table 2). We then

analyzed a similar set of genes for a regulatory element where the Psr1-like TF might

bind (Fig. S5), including genes that were DE in the P-deficient transcriptome relative to

the replete (Table S1 from Bachy *et al.*, 2018). Under P-deficiency, the *psr*1-like gene

(JGI gene ID 360) was highly up-regulated (log-2 fold-change = 4.3, q = 0.00037) and

nearly all of the predicted P-responsive genes were DE (Table 1). For example, *M.*

pusilla up-regulated many genes in the TCA cycle but not pyruvate carboxylase (gene ID

6984). Genes involved in nucleotide and lipid metabolism (e.g., nucleoside phosphatase,

long-chain acyl-coA synthetase) were DE and contained a conserved motif with

similarity to myb-like regulatory element in *A. thaliana* (Table 1, Fig. S5). Lastly, the

POX gene described in *E. huxleyi* (Rokitta *et al.*, 2016) and observed in the archived gene

model for *M. pusilla*, was absent, or not similarly annotated in the updated gene model.

Instead, a copper amine oxidase (gene ID 5863) contained a significant conserved motif

(Table 1, Fig. S5) and was DE under P-deficiency (Bachy *et al.* 2018).

We conducted a similar analysis with published transcriptome data from *M.*

commoda (Whitney & Lomas, 2016). The *psr*1-like gene (JGI gene ID 60184) in *M.*

commoda was significantly up-regulated in the P-deficient treatment (defined by Whitney

& Lomas, 2016; Table S5), and we found a significant conserved motif in 18 DE genes.

The significant motif identified in *M. commoda* differed in sequence from that discovered

in *M. pusilla* and occurred in a different set of genes (Notes S1).

Discussion

Our combined metabolomics and comparative genomic analyses of the response

of *M. pusilla* CCMP1545 to P-deficiency has led to three conclusions: 1) there is a shift

in intracellular metabolite composition, 2) a *psr*1-like gene is expressed in *M. pusilla* and

other marine phytoplankton, and 3) the genes regulated by the Psr1-like protein may

differ amongst algal species.

Micromonas pusilla *exhibits a metabolic shift in response to P-deficiency*

The general decrease in concentration of extracellular metabolites of P-deficient

cells indicates reduced release of these central metabolites under P-deficiency. This is

expected as nutrient-limited microalgae may have lower photosynthetic activity (Loebl *et*

al., 2010; Halsey *et al*., 2014; Guo *et al*., 2018) and hence, produce less metabolic

overflow. In contrast, most intracellular metabolites did not differ significantly in

concentration between treatments. Because these metabolites represent many central

metabolic pathways, it appears that *M. pusilla* does not respond to P-deficiency through a

global decrease in metabolic activity, as was observed for the diatom, *Phaeodactylum*

tricornutum (Feng *et al*., 2015). Interestingly, we found examples of functionally-related

metabolites that behaved differently. For example, glutamate, proline, and methionine

were higher on average, although variable, under P-deficiency, while phenylalanine and

tryptophan were similar. In the TCA cycle, malate was variable but higher in

concentration on average under P-deficiency, while citrate was lower on average and

succinate was relatively unchanged. The opposing responses of these TCA intermediates

was surprising given that others (e.g., Hockin *et al*., 2012; Goncalves *et al*., 2013) have

observed a coordinated response to nutrient limitation in transcripts or proteins involved

in the TCA cycle.

We observed additional non-uniform metabolite responses, with vitamins and

glucosamine-6-phosphate higher on average under P-deficiency, but other P-containing

compounds (e.g. glycerol-3-phosphate) and sugars were depleted. Previous work has

suggested that organisms placed under nutrient stress are capable of diverting carbon flux

between pathways (reviewed in Markou & Nerantzis, 2013). Changes in carbon flux

through the cell could result in coordinated abundance shifts in seemingly unrelated

metabolites, as observed in this study. If the shunting of carbon is a coordinated response

within *M. pusilla*, then we would expect genes involved in the synthesis and/or

catabolism of these compounds to be co-regulated, and potentially mediated by a specific

transcription factor. Indeed, in *M. pusilla,* we discovered a gene with significant

homology to the transcription factor phosphorus starvation response gene (*psr*1)

originally described in the freshwater alga *Chlamydomonas reinhardtii* (Wykoff et al.,

1999).

Presence and expression of the transcription factor gene, psr*1, in marine algae*

In *C. reinhardtii* and in many plants, a *psr*1 gene encodes for a transcription factor

(Psr1) that coordinates the metabolic response to P-deficiency in these organisms

(Wykoff *et al*., 1999; Mosely *et al*., 2006). We detected a putative *psr*1-like gene in the

M. pusilla genome and in the genomes and transcriptomes of major algal lineages (e.g.,

prasinophytes, dinoflagellates). The *psr*1-like genes that we discovered were generally

not annotated as *psr*1 with the exception of *Ostreococcus tauri* (Derelle et al., 2006),

although to our knowledge the potential role of this gene in the metabolism of *O. tauri* or

other marine algae has not been discussed. The *M. pusilla psr*1-like gene contains two

myb domains similar to those in *psr*1 from *C. reinhardtii* (Wykoff *et al.*, 1999), and

comparable to those in *phr*1 in *A. thaliana* (Rubio *et al.*, 2001). No *psr*1-like gene was

identified by our queries of diatom transcriptomes or genomes, suggesting that a

phosphate starvation response gene akin to *psr*1 is present in diverse but phylogenetically

constrained phytoplankton groups.

We found significant similarity between the *M. pusilla psr*1-like gene and

transcript sequences from the MMETSP and *Tara* Oceans datasets, indicating that this

gene is expressed by marine algae in cultures and in the environment. High-identity hits

to sequences in the global ocean survey (GOS, Rusch *et al.* 2007) and *Tara* Oceans

datasets underscore this gene's prevalence in the oceans, particularly in regions

characterized by chronically low phosphorus concentrations. Short sequences in some

MMETSP and *Tara* Oceans sequences did not include enough of the C-terminal end to

capture the myb coiled-coil domain. Thus, further investigation is required to confirm

whether the identified transcripts are from *psr*1-like genes. Several GOS sequences,

previously described as protein of unknown function, had homology to both characteristic

Psr1 domains, suggesting that they could be re-annotated as *psr*1-like genes. The GOS

sequences are likely derived from prasinophytes (i.e., *Micromonas* spp., *Ostreococcus*

spp.) and haptophytes (e.g., *E. huxleyi*), highlighting the need to elucidate the role of *psr*1

in these organisms.

Potential role of Psr1 in the marine algal response to P-deficiency

463   Transcription factors typically recognize a specific motif in DNA where the

protein binds to regulate each gene. If such a motif is observed, one can hypothesize that

a gene containing the motif is regulated by the protein. We observed a conserved and

enriched motif across genes involved in nucleotide biosynthesis, the TCA cycle and

glycolysis, carbon fixation, fatty acid metabolism, and phosphate transport or salvage

(Fig. 6). The conserved motifs discovered in both *M. pusilla* and *M. commoda* had unique

sequences and were present in a slightly different set of genes (Notes S2). The motif

detected in *M. pusilla* is similar in nucleotide sequence to a myb DNA-binding domain in

A. thaliana. This similarity is potentially significant, because the *psr*1/*phr*1 genes in *C.*

reinhardtii and *A. thaliana* contain a myb DNA-binding domain. Thus, we hypothesize

that the conserved motif detected in the *Micromonas* species represents binding sites for

the Psr1-like transcription factors (TF).

475   The conserved motif was observed only in DE genes under P-deficiency by each

Micromonas species (Bachy *et al*., 2018; Whitney & Lomas, 2016), suggesting that these

genes could be co-regulated. In bacteria, for example, genes within the well-characterized

Pho-regulon all contain specific sequences (the PHO box) where the transcription factor

binds to activate or repress the gene (Santos-Beneit, 2015). These PHO box sequences

are distinct among different bacterial species and strains. In *M. commoda*, the discovered

motif was not significantly similar to myb-like DNA-binding motifs in *A. thaliana* and

was not similar to the motif discovered in *M. pusilla*. Although we expected that the myb

domain of the Psr1-like proteins in the two species would interact with similar binding

regions in the genome, *psr*1-like derived amino acid sequences from *M. pusilla* and *M.*

commoda were only 47% similar. TF protein sequences with up to 79% similarity have

been shown to have distinct DNA-binding motif profiles (Franco-Zorrilla et al., 2014), so

the two proteins in different species could reasonably have distinct DNA-binding motifs.

If the Psr1-like TF regulates a different set of genes in the two *Micromonas* species, this

may serve as a niche defining feature between these taxa. As further support for this

hypothesis, the conserved motif was only observed in a subset of genes that were

differentially expressed under P-limitation or deficiency (Whitney & Lomas, 2016;

Bachy *et al.*, 2018). The observed differences in the *psr*1-like gene and differences in

associated binding motifs may represent adaptation to distinct environments as *M. pusilla*

was collected in the temperate English Channel, while *M. commoda* was isolated from

the tropics (Worden, 2006).

496   The *psr*1-like gene in *M. commoda* and *M. pusilla* was one of the highest DE

genes in each experiment (Whitney & Lomas, 2016; Bachy *et al.*, 2018), suggesting that

this potential TF plays a critical role in the P-deficiency response of *Micromonas*.

Moreover, the *psr*1-like gene was highly expressed in two culture experiments with

different growth conditions and sampled at different growth stages, suggesting a broad

physiological relevance for the Psr1-like TF (Fig. 6). In contrast to the transcriptome

results, a recent study describing proteome changes in *M. pusilla* CCMP1545 grown

under P-limited conditions in a bioreactor (Guo *et al.*, 2018), did not observe significant

regulation of the Psr1-like protein (significance defined by Guo *et al.*, 2018). The

discordant results may be due to known differences in regulation between genes and

proteins (Guo *et al.*, 2018) or to different Psr1-like protein behavior under continuous

culturing conditions. Regardless, these results warrant further investigation into the

physiological role of the Psr1-like protein in *M. pusilla*.

Variable and non-uniform dynamics of several TCA cycle metabolites between

treatments suggested that this pathway may be involved in the metabolic response to P-

deficiency in *M. pusilla*. Indeed, nearly all of the genes involved in the end of glycolysis

and in the TCA cycle were up-regulated in *M. pusilla* (Bachy *et al*., 2018) and these

genes contained the conserved motif that may function as a regulatory element where the

Psr1-like TF could bind. Fumarase, in particular, was upregulated in both *M. pusilla* and

M. commoda transcriptomes under P-limitation or deficiency. If fumarase expression is

regulated by the Psr1-like protein, expression of these two genes (fumarase and *psr*1)

should be correlated in field populations. We observed support for this hypothesis within

the *Tara* Oceans dataset, although we observe weaker relationships between Psr1-like

gene expression and other up-regulated genes in either *M. pusilla* or *M. commoda* (Notes

S2). In *M. pusilla*, carbon flow through the TCA cycle may increase as a means to fuel

TAG or starch production and to pull potentially damaging energy away from the

photosystems (Norici *et al*., 2011; Klok *et al*., 2013). Similarly, several proteins involved

in glycolysis (e.g., pyruvate dehydrogenase) and the TCA cycle (e.g., succinate

dehydrogenase) were found to be up-regulated in another study with *M. pusilla* under P-

limitation (Guo *et al*., 2018), lending some support to this hypothesis. Insight from the

latter study underscores potential differences in regulation of genes and enzymes

involved in glycolysis and the TCA cycle (Guo *et al*., 2018). For example, we found

congruency in up- or down-regulation of some proteins from Guo *et al*. (2018) that we

would expect in *M. pusilla* under P-deficiency based on our combined comparative

genomics and metabolomics approach (i.e., pyruvate kinase, pyruvate dehydrogenase,

oxoglutarate dehydrogenase, and succinate dehydrogenase) but not others (i.e., fumarase,

aconitase, citrate synthase). Analysis of transcript, protein, and metabolite responses in *M.*

pusilla under the same growth conditions will be required to elucidate the role of Psr1 in

this and other taxa.

In addition to TCA cycle intermediates, we observed shifts for several purine

nucleosides between treatments and detected the conserved motif in genes for a

nucleoside phosphatase and aspartate transcarbamylase. These results suggest increased

nucleotide salvage through purine nucleotides and nucleosides, a mechanism described in

M. commoda (Whitney & Lomas, 2016) and in other phytoplankton (Dyhrman & Palenik,

2003; Kujawinski *et al*., 2017). Previous phytoplankton work has also highlighted

expression of proline oxidase (POX) (Rokitta *et al*., 2014; Rokitta *et al*., 2015) in nutrient

limitation experiments (including P-deficiency) with the haptophyte *E. huxleyi,* where

this enzyme may have a role in stabilizing the mitochondrial membrane and in detecting

cellular nitrogen levels. The POX gene was not present in *M. pusilla* but was up-

regulated in *M. commoda* (Whitney & Lomas, 2016; Table S4). In *M. pusilla*, a copper

amine oxidase was up-regulated (Bachy *et al*., 2018) and contained the conserved motif

that may function as a regulatory element for the Psr1-like TF. Thus, POX and copper

amine oxidase represent gene targets for further investigation.

Maat *et al*. (2014) noted that both *E. huxleyi* (Borchard *et al*., 2011) and *M.*

pusilla (Maat *et al*., 2014) exhibited a similar metabolic response to elevated pCO$_2$ levels

and P-deficient conditions. While there may be multiple factors underlying the response

of these organisms, both contain the *psr*1-like gene which could direct part of a shared

metabolic response to elevated pCO$_2$ and nutrient-limited conditions. While

transcriptomic analysis of *E. huxleyi* was not performed in the study by Borchard *et al*.

(2011), Rokitta and colleagues (2014; 2016) performed microarray transcriptome analysis

of *E. huxleyi* cells under separate nitrogen- and P-deficient conditions (normal pCO$_2$).

The transcripts were only partial sequences; however, two up-regulated genes detected by

Rokitta *et al*. (2016), contained myb-like DNA binding domains (gene IDs: GJ02872,

GJ03978). Targeted experimental work is necessary to elucidate the presence and role of

a *psr*1-like gene in *E. huxleyi*. If there is a common role for the *psr*1-like gene in *E.*

huxleyi and *M. pusilla* in their response to nutrient limitation and elevated pCO_2, this

could have important implications for predicting the physiological response of the wide

range of organisms that contain this *psr*1-like gene to changing environmental conditions.

The potential impact of Psr1-regulated genes on the ecological roles of

Micromonas spp. is unknown, but our results suggest that the Psr1-like protein

coordinates a metabolic shift in these organisms under P-deficiency, altering the

intracellular flow of carbon and other elements. More comprehensive examination of

these metabolic responses, which likely vary to some extent among organisms, will be

paramount to improving models of trophic carbon flow. More experiments are needed to

characterize the structure and role of the *psr*1-like gene in *Micromonas* spp. and other

phytoplankton, including: (1) confirming the presence of the *psr*1-like gene with genetic

experiments, (2) determining the conditions under which the Psr1-like protein is abundant,

(3) identifying the taxon-specific genes affected by the Psr1-like protein, (4) verifying the

interaction of the Psr1-like protein with the hypothesized binding sites, and (5)

comparing the genetic and metabolic responses to P-deficiency between organisms

containing the *psr*1-like and those that do not. Exploring the underlying biology of the

*psr*1 gene will facilitate mechanistic understanding of the complex response of these

organisms to P-limitation and will enhance our ecological and biogeochemical

predictions.

Acknowledgements

The authors would like to thank Krista Longnecker for performing the TOC analysis and Matthew Johnson and Elizabeth Harvey for assistance with flow cytometry and FIRe analysis. This research was funded by the Gordon and Betty Moore Foundation through Grant GBMF3304 to E. Kujawinski.

Author contributions

CLF, MCKS, and EBK conceived and designed the experiments; CLF performed the experimental work; CLF and HA conducted data analyses; EBK supervised the study; CLF drafted the manuscript and HA, MCKS, and EBK provided input and helped revise the manuscript.

References

Bachy C, Charlesworth CJ, Chan AM, Finke JF, Wong C-H, Wei C-L, Sudek S, Coleman ML, Suttle CA, Worden AZ. 2018. Transcriptional responses of the marine green alga *Micromonas pusilla* and an infecting prasinovirus under different phosphate conditions. *Environ Microbiol.* **20**:2898-2912.

Bailey TL, Elkan C. 1994. Fitting a mixture model by expectation maximization to discover motifs in biopolymers. *Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology*, AAAI Press, Menlo Park, California, pp. 28-36,

Barrett LW, Fletcher S, Wilton SD. **2012.** Regulation of eukaryotic gene expression by the untranslated gene regions and other non-coding elements. *Cell Molec Life Sci.* **69**:3613-3634.

Berdalet E, Latasa M, Estrada M. 1994. Effects of nitrogen and phosphorus starvation on nucleic acid and protein content of Heterocapsa sp. *J Plank Res.* **16**: 303–316.

Borchard C, Borges AV, Haendel N, Engel A. 2011. Biogeochemical response of *Emiliania huxleyi* (PML B92/11) to elevated CO_2 and temperature under phosphorous limitation: a chemostat study. *J. Exp. Mar. Biol. Ecol.* **410**:61–71.

Carradec Q, Pelletier E, Da Silva C, Alberti A, Seeleuthner Y, Blanc-Mathieu R, Lima-Mendez G, Rocha F, Tirichine L, Labadie K, Kirilovsky A, et al. 2017. A global ocean atlas of eukaryotic genes. *Nature Comm.* **9**:373.

Cavender-Bares KK, Karl DM, Chisholm SW. 2001. Nutrient gradients in the western North Atlantic Ocean: Relationship to microbial community structure and comparison to

patterns in the Pacific Ocean. *Deep Sea Res Part I: Oceanogr Res Pap.* **48**:2373–2395.

Chung C-C, Hwang S-PL, Chang J. 2003. Identification of a high-affinity phosphate
transporter gene in a prasinophyte alga, *Tetraselmis chui*, and its expression under
nutrient limitation. *Appl Environ Microbiol.* **69**:754-759.

de Carvalho CCCR, Fernandes P. 2010. Production of metabolites as bacterial
responses to the marine environment. *Marine Drugs* **3**:705-727.

Derelle E, Ferraz C, Rombauts S, Rouzé P, Worden AZ, Robbens S, Partensky F,
Degroeve S, Echeynié S, Cooke R, et al. 2006. Genome analysis of the smallest free-
living eukaryote *Ostreococcus tauri* unveils many unique features. *Proc Nat Acad Sci.*
103:11647–11652.

Dittmar T, Koch B, Hertkorn N, Kattner G. 2008. A simple and efficient method for
solid-phase extracion of dissolved organic matter (SPE-DOM) from seawater. *Limn.*
Oceangr. Meth. **6**: 230–235.

Dyhrman ST, Ruttenberg KC. 2006. Presence and regulation of alkaline phosphatase
activity in eukaryotic phytoplankton from the coastal ocean: Implications for dissolved
organic phosphorus remineralization. *Limnol Oceanogr.* **51**:1381–1390.

Dyhrman ST, Benitez-Nelson CR, Orchard ED, Haley ST, Pellechia PJ. 2009. A
microbial source of phosphonates in oligotrophic marine systems. *Nat Geosci.* **2**:696–699.

Dyhrman ST, Jenkins DB, Rynearson TA, Saito MA, Mercier ML, Alexander H,
Whitney LP, Drzewianowski A, Bulygin VV, Bertrand EM, et al. 2012. The
transcriptome and proteome of the diatom *Thalassiosira psuedonana* reveal a diverse
phosphorus stress response. PLoS ONE **7**:e33768.

Easlon E, Tsang F, Skinner C, Wang C, Lin S-J. 2008. The malate-aspartate NADH
shuttle components are novel metabolic longevity regulators required for calorie
restriction-mediated life span extension in yeast. *Genes Devel.* **22**:931-944.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucl Acids Res.* **32**:1792–1797.

Elser JJ, Bracken MES, Cleland EE, Gruner DS, Harpole WS, Hillebrand H, Ngai
JT, Seabloom EW, Shurin JB, Smith JE. 2007. Global analysis of nitrogen and
phosphorus limitation of primary producers in freshwater, marine and terrestrial
ecoystems. *Ecol Lett.* 20:1135-1142.

Falkowski PG, Katz ME, Knoll AH, Quigg A, Raven JA, Schofield O, *et al*. 2004.
The evolution of modern eukaryotic phytoplankton. *Science* **16**:354-360.

Franco-Zorrilla JM, López-Vidriero I, Carrasco JL, Godoy M, Vera P, Solano R.
2014. DNA-binding specificities of plant transcription factors and their potential to define
target genes. *Proc Nat Acad Sci.* **111**:2367–2372.

Felsenstein J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood
approach. *J molec evol.* **17**:368–376.

Feng T-Y, Yan Z-K, Zheng J-W, Xie Y, Li D-W, Bala Murugan S, *et al*. 2015.
Examination of metabolic responses to phosphorus limitation via proteomic analyses in
the marine diatom *Phaeodactylum tricornutum. Sci Rep.* **5**:10373.

Fiore CL, Longnecker K, Kido Soule MC, Kujawinski EB. 2015. Release of
ecologically relevant metabolites by the cyanobacterium *Synechococcus elongatus*
CCMP 1631. *Environ Microbiol.* **17**:3949–3963.

Goncalves EC, Johnson JV, Rathinasabapathi B. 2013. Conversion of membrane lipid
acyl groups to triacylglycerol and formation of lipid bodies upon nitrogen starvation in
biofuel green algae *Chlorella* UTEX29. *Planta* **238**:895-906.

Guo J, Wilken S, Jimenez V, Choi CJ, Ansong C, Dannebaum R, Sudek L, Milner
DS, Bachy C, Reistetter EN, et al. 2018. Specialized proteomic responses and an
ancient photoprotection mechanism sustain marine green algal growth during phosphate
limitation. *Nat Microbiol Lett.* 3:781-790.

Gupta S, Stamatoyannopolous JA, Bailey T, Noble WS. 2007. Quantifying similarity
between motifs. *Gen Biol.* **8**:R24.

Halsey KH, Milligan AJ, Behrenfeld MJ. 2014. Contrasting strategies of
photosynthetic energy utilization drive lifestyle strategies in ecologically important
picoeukaryotes. *Metabolites* **4**: 260-280.

Hockin NL, Mock T, Mulholland F, Kopriva S, Malin G. 2012. The response of
diatom central carbon metabolism to nitrogen starvation is different from that of green
algae and higher plants. *Plant Physiol.* **158**:299-312.

Hoppe CJM, Flintrop CM, Rost B. 2018. The Arctic picoeukaryote *Micromonas*
pusilla benefits synergistically from warming and ocean acidification. *Biogeosciences*
15:4553-4365.

Johnson WM, Kido Soule MC, Kujawinski EB. 2017. Extraction efficiency and
quantification of dissolved metabolites in targeted marine metabolomics. *Limnol*
Oceanogr Meth. **15**: 417-428
682
683 **Johnson LK, Alexander H, Brown CT. 2018.** Re-assembly, quality
684 evaluation, and annotation of 678 microbial eukaryotic reference transcriptomes. *Giga*
685 *Sci.* **8**:giy158
686
687 **Jones DT, Taylor WR, Thornton J.M. 1992.** The rapid generation of mutation data
688 matrices from protein sequences. *Computer Appl Biosci.* **8**:275-282.
689
Karl DM. 2014. Microbially mediated transformations of phosphorus in the sea: new
views of an old cycle. *Ann Rev Mar Sci.* **6**:279-337.

Kido Soule MCK, Longnecker K, Johnson WM, Kujawinski EB. 2015.
Environmental Metabolomics: Analytical Strategies. *Mar Chem.* **177**:1–62.

Kishino H, Miyata T, Hasegawa M. 1990. Maximum likelihood inference of protein
phylogeny and the origin of chloroplasts. *J Molec Evol.* 31:151–160.

Klok AJ, Martens DE, Wijffels RH, Lamers PP. 2013. Simultaneous growth and
neutral lipid accumulation in microalgae. *Biores Technol.* **134**:233–243.

Kujawinski EB, Longnecker K, Alexander H, Dyhrman ST, Fiore CL, Haley ST,
Johnson WM. 2017. Phosphorus availability regulates intracellular nucleotides in marine
eukaryotic phytoplankton. *Limnol Oceanogr Lett.* **2**:119-129. doi:10.1002/lol2.10043.

Kumar S, Stecher G, and Tamura K. 2016. MEGA7: Molecular Evolutionary Genetics
Analysis version 7.0 for bigger datasets. *Molec Biol Evol.* **33**:1870-1874.

Lewis LA, McCourt RM. 2004. Green algae and the origin of land plants. *Amer J Bot.*
91:1535–1556.

Lai J, Yu Z, Song X, Cao X, Han X. 2015. Responses of the growth and biochemical
composition of *Prorocentrum donghaiense* to different nitrogen and phosphorus
concentrations. *J Exp Mar Biol Ecol.* **405**:6-17.

Li WKW. 1994. Primary production of prochlorophytes, cyanobacteria, and eukaryotic
ultraphytoplankton: Measurements from flow cytometric sorting. *Limnol Oceanogr*
39:169-175.

Li M, Shi X, Guo C, Lin S. 2016. Phosphorus deficiency inhibits cell division but not
growth in the dinoflagellate *Amphidinium carterae*. *Front Microbiol.* **7**:826.

Lin S, Litaker RW, Sundra WG. 2016. Phosphorus physiological ecology and
molecular mechanisms in marine phytoplankton. *J Phycol.* **52**:10-36.

Loebl M, Cockshutt AM, Campbell D, Finkel ZV. 2010. Physological basis for high
resistance to photoinhibition under nitrogen depletion in *Emiliania huxleyi*. *Limnol*
Oceanogr 55:2150-2160.

Longnecker K. 2015. Dissolved organic matter in newly formed sea ice and surface
seawater. *Geochim Cosmochim Acta* **171**:39–49.

Maat DS, Crawfurd KJ, Timmermans KR, Brussaard CPD. 2014. Elevated CO_2 and
phosphate limitation favor *Micromonas pusilla* through stimulated growth and reduced
viral impact. *Appl Environ Microbiol.* **80**:3119–3127.

Maat DS, van Bleijswijk JDL, Witte HJ, Brussaard PD. 2016. Virus production in
phosphorus-limited *Micromonas pusilla* stimulated by a supply of naturally low
concentrations of different phosphorus sources, far into the lytic cycle. *FEMS Microbiol*
Ecol. **92**:fiw136

Marchler-Bauer A, Bo Y, Han L, Lanczycki CJ, Lu S, Chitsaz F, Derbyshire MK,
Geer RC, Gonzales NR, Gwadz M, et al. 2017. CDD/SPARCLE: functional
classification of proteins via subfamily domain architectures. *Nucl Acids Res.* **4**:D200-
D203.

Markou G, Nerantzis E. 2013. Microalgae for high-value compounds and biofuels
production: a review with focus on cultivation under stress conditions. *Biotechn Advan.*
31:1532-1542.

Martin P, Van Mooy BA, Heithoff A, Dyhrman ST. 2011. Phosphorus supply drives
rapid turnover of membrane phospholipids in the diatom *Thalassiosira pseudonana*.
ISME J. **25**:1057–1060.

Martin P, Dyhrman ST, Lomas MW, Poulton NJ, Van Mooy BAS. 2014.
Accumulation and enhanced cycling of polyphosphate by Sargasso Sea phytoplankton in
response to low phosphorus. *Proc Nat Acad Sci.* **111**:8089-8094.

Mock T, Samanta MP, Iverson V, Berthiaume C, Robison M, Holtermann K, *et al*.
2008. Whole-genome expression profiling of the marine diatom *Thalassiosira*
pseudonana identifies genes involved in silicon bioprocesses. *Proc Nat Acad Sci.*
105:1579-1584.

Moseley JL, Chang C-W, Grossman AR. 2006. Genome-based approaches to
understanding phosphorus deprivation responses and PSR1 control in *Chlamydomonas*
reinhardtii. *Euk Cell* **5**:26–44.

Müller R, Morant M, Jarmer H, Nilsson L, Nielsen TH. 2006. Genome-wide analysis
of the *Arabidopsis* leaf transcriptome reveals interaction of phosphate and sugar
metabolism. *Plant Physiol.* **143**:156–171.

Norici A, Bazonni AM, Pugnetti A, Raven JA, Giordano M. 2011. Impact of
irradiance on the C allocation in the coastal marine diatom *Skeletonema marinoi* Sarno
and Zingone. *Plant Cell Environ.* **34**:1666–1677.

Rokitta SD, von Dassow P, Rost B, John U. 2014. *Emiliana huxleyi* endures N-
limitation with an efficient metabolic budgeting and effective ATP synthesis. *BMC*
Genomics **15**:1051.

Rokitta SD, von Dassow P, Rost B, John U. 2016. P- and N-depletion trigger similar
cellular responses to promote senescence in eukaryotic phytoplankton. *Front Mar Sci*
3:109.

Rubio V, Linhares F, Solano R, Martín AC, Iglesias J, Leyva A, Paz-Ares J. 2001. A
conserved MYB transcription factor involved in phosphate starvation signaling both in
vascular plants and in unicellular algae. *Genes Devel.* **15**: 2122–2133.

Rusch DB, Halpern AL, Sutton G, Heidelberg KB, Williamson S, Yooseph S, Wu D,
Eisen JA, Hoffman JM, Remington K, et al. 2007. The Sorcerer II global ocean
sampling expedition: northwest Atlantic through eastern tropical Pacific. *PLoS Biol.*

5:e77.

Santos-Beneit F. 2015. The Pho regulon: a huge regulatory network in bacteria. *Front*
Microbiol. **6**:402.

Sharma KK, Schuhmann H, Schenk PM. 2012. High lipid induction in microalgae for
biodiesel production. *Energies* **5**:1532–1553.

Shemi A, Schatz D, Fredricks HF, Van Mooy BAS, Porat Z, Vardi A. 2016.
Phosphorus starvation induces membrane remodeling and recycling in *Emiliania huxleyi.*
New Phytol. **211**:886-898.

Shi X, Lin X, Li L, Palenik B, Lin S. 2017. Transcriptomic and microRNAomic
profiling reveals multi-faceted mechanisms to cope with phosphate stress in a
dinoflagellate. *ISME J.* **11**:2209-2218.

Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5:
Molecular Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary
Distance, and Maximum Parsimony Methods. *Molec. Biol. Evol.* **28**: 2731–2739.

Tyrrell T. 1999. The relative influences of nitrogen and phosphorus on oceanic primary
production. *Nature* **400**:525–531.

Van Mooy BAS, Fredricks HF, Pedler BE, Dyhrman ST, Karl DM, Koblížek M,
Lomas MW, Mincer TJ, Moore LR, Moutin T, et al. 2009. Phytoplankton in the ocean
use non-phosphorus lipids in response to phosphorus scarcity. *Nature* **457**:69–72.

Whitney LP, Lomas MW. 2016. Growth on ATP elicits a P-stress response in the
picoeukaryote *Micrmonas pusilla. PLoS ONE* **11**:e0155158.

Worden AZ. 2006. Picoeukaryote diversity in coastal waters of the Pacific Ocean. *Aquat*
Microb Ecol. **43**:165–175.

Wykoff DD, Grossman AR, Weeks DP, Usuda H, Shimogawara K. 1999. Psr1, a
nuclear localized protein that regulates phosphorus metabolism in *Chlamydomonas. Proc*
Nat Acad Sci. **96**:15336–15341.

Yoshida K, Terashima I, Noguchi K. 2007. Up-regulation of mitochondrial alternative
oxidase concomitant with chloroplast over-reduction by excess light. *Plant Cell Physiol.*
48:606-614.

Figure Legends

Figure 1. Average ratio of the P-deficient to P-replete intracellular metabolite
concentrations during exponential growth (T_1) in *Micromonas pusilla* CCMP1545.
Concentrations are normalized to cell number in each treatment and means with one
standard deviation are shown (N=3 unless otherwise noted). Metabolites are listed in
order of descending concentration within three groups; those that were detected in
enough replicates for standard deviation to be calculated, those where only one ratio
could be calculated (black line), and those that were not detected (ND) in any of the three
replicates for one treatment (R = P-replete, D = P-deficient). Metabolite names marked
with an asterisk indicates a significant difference in concentration between treatments (*t*-
test, p < 0.05). Only one replicate in the P-deficient treatment contained a non-zero
concentration for *N*-acetylmuramic acid, thus while there was a significant difference in
concentration for this metabolite, only one ratio could be calculated. (DMSP =
dimethylsulfoniopropionate, NAD = nicotinamide adenine dinucleotide)

Figure 2. The intracellular ratios of two purine nucleosides to their nucleobases for
Micromonas pusilla CCMP1545 in P-deficient (orange) and P-replete (blue) treatments
during exponential growth (T_1). Xanthosine and xanthine, and guanosine and guanine
were quantified in the targeted metabolomics method and normalized to cell abundance.
The average ratio and high and low values are shown based on three replicate cultures.
The asterisk marks a significant difference between the treatments (*t*-test, p = 0.03). The
dashed line represents a ratio of 1. Two samples in the P-replete treatment had guanine
levels below the limit of detection and the one ratio is shown as a line.

Figure 3. Predicted amino acid sequences of the *psr*1-like gene in *Micromonas pusilla*
CCMP1545 and a subset of other organisms. The amino acid sequence (with IMG gene
ID) is shown as described in the Joint Genome Institute Integrated Microbial Genomes *M.*
pusilla CCMP1545 database. The myb-like DNA-binding and myb coiled-coiled domains
are highlighted in blue (a). Predicted amino acid sequence alignment of *psr*1 and *psr*1-
like genes in the region of the myb-like DNA-binding domain (SHAQKYF class) and
LHEQL coiled-coil domain (b). Asterisks at the top of each column indicate 100%
conserved residues across species surveyed. Numbers at the end of each row indicate
position of the last shown residue in the amino acid sequence. Accession (MMETSP,
NCBI, *Tara*) or gene identification (IMG) numbers are listed for each sequence.

Figure 4. Occurrence and phylogenetic relationship of *psr*1-like genes in marine
phytoplankton. Maximum likelihood method based on the JTT matrix-based model
(Jones *et al*., 1992). The tree with the highest log likelihood (-2442.94) is shown and is
based on the derived amino acid sequences for *psr*1 and *psr*1-like genes from eukaryotic
phytoplankton. The analysis involved 55 amino acid sequences. All positions containing

gaps and missing data were eliminated yielding a total of 43 positions in the final dataset.
Evolutionary analyses were conducted in MEGA7 (Kumar *et al*., 2016) Accession or
IMG gene identification numbers are shown with the taxon name and the star indicates
Micromonas pusilla CCMP1545 cultured in this study. The tree is unrooted and the scale
indicates the number of amino acid substitutions per site.

Figure 5. Geographic distribution of *Micromonas psr*1-like genes and relationship with
phosphate concentration in the *Tara* Oceans eukaryotic gene atlas (Carradec, Pelletier, *et*
al., 2017). The geographic distribution and relative abundance of *Micromonas psr*1-like
genes in surface samples are depicted. Black dots represent stations where *Micromonas*
was detected; the size of purple circles represents the relative abundance of *psr*1-like
gene, where the FPKM of the psr1-like gene was normalized to the total FPKM of all
Micromonas-associated transcripts (a). Relative abundance of *Micromonas* is plotted
against phosphate concentrations for surface stations where *Micromonas* was detected.
Grey circles indicate samples where *psr*1-like genes were not detected, while samples
where psr1-like genes were detected are colored based on the relative expression of
*Micromonas psr*1-like gene, again normalized to total *Micromonas* transcript abundance.
No significant relationship was observed between the phosphate concentration and
abundance of *Micromonas psr*1-like transcripts (*t*-test, $p > 0.05$) (b). A split violin plot
depicting the distribution of expression of *Micromonas* fumarase synthase genes as a
function of presence or absence of *psr*1-like gene. The larger dashed line represents the
distribution mean and the smaller dashed lines represent the 25% and 75% quartiles.
Significant difference in the distribution of expression between the presence and absence
is assessed with Kolmogorov-Smirnov test ($p < 0.005$) (c).

Figure 6. Conceptual model of Psr1-like TF regulation in *Micromonas pusilla* CCMP154
in three major metabolic pathways under phosphorus deficiency. The figure illustrates the
relevant metabolites quantified in the present study and indicates if they are elevated
(orange, ratio > 1 Figure 2), depressed (blue, ratio < 1 Figure 2) in concentration in the P-
deficient cells relative to the P-replete cells. Some metabolite concentrations were equal
in each treatment (black) or were not quantified (gray). Genes that contain the conserved
motif with similarity to the myb-like regulatory element in *Arabidopsis thaliana* are
marked with a red star. The orange and blue letters refer to elevated or depressed
transcript levels for each gene ([1]data from Bachy *et al*., 2018). Descriptions of putative
metabolic responses in each pathway are shown. Genes encode for: A) pyruvate kinase,
B) pyruvate dehydrogenase, C) citrate synthase, D) aconitase, E) α-ketoglutarate
dehydrogenase, F) succinate dehydrogenase, G) fumarase, H) ribulose-1,5-bisphosphate
carboxylase, I) triose sugar transporter, J) nucleotide phosphatase DAG = diacylglycerol,
TAG = triacylglycerol

884   **Tables**

885 Table 1. Enzymes that contain a significant motif in the gene sequences of *M. pusilla* CCMP1545 (MBARI_models (ver 1)). Column 2: the
886 Enzyme Commission (E.C.) number or pfam identifier; Column 3: main pathway(s) in which the enzyme is involved; Columns 4-6: fold
887 change and *q*-value of the transcript in P-deficient treatment relative to P-replete conditions are data from Bachy *et al*. (2018) and
888 corresponding gene ID.

Enzyme	E.C. Number	Pathway	Log 2 fold change*	*q*-value*	JGI gene ID*
Glyceraldehyde-3-phosphate dehydrogenase	1.2.1.12	Calvin cycle/glycolysis	2.73	0.000037	4267
Glyceraldehyde-3-phosphate dehydrogenase	1.2.1.12	Calvin cycle/glycolysis	-0.644	0.000037	700
Phosphoglycerate kinase	2.7.2.3	Glycolysis	-0.699	0.000037	7661
Pyruvate kinase	2.7.1.40	Glycolysis	-1.32	0.000037	9370
Pyruvate dehydrogenase	1.2.4.1	TCA[a] cycle/Glycolysis	0.756	0.000037	3535
Citrate synthase	2.3.3.1	TCA cycle	1.68	0.000037	1278
Citrate synthase	2.3.3.1	TCA cycle	1.10	0.000037	3714
Aconitase	4.2.1.3	TCA cycle	1.00	0.000037	2901
Succinate dehydrogenase	1.3.5.1	TCA cycle and ETC[b]	1.16	0.001515	2182
Fumarase	4.2.1.2	TCA cycle	1.68	0.000037	600
Fumarase	4.2.1.2	TCA cycle	1.73	0.000068	6662
Copper amine oxidase	1.4.3.21-22	Arginine and proline metabolism	2.87	0.000037	5863
Aspartate transcarbamylase	2.1.3.2	Pyrimidine biosynthesis	-0.609	0.000037	3615
Nucleoside phosphatase	3.6.1.3	Pyrimidine metabolism	0.711	0.000037	3322
Pantoate-beta-alanine ligase	6.3.2.1	Pantothenate and CoA biosynthesis	-0.616	0.000037	5304
Lysophospholipase	3.1.1.5	Glycerophospholipid metabolism	1.65	0.000037	3652
Acyl-CoA synthetase	6.2.1.3	Fatty acid metabolism	3.20	0.000037	1751
Long-chain acyl-CoA synthetases	6.2.1.3	Fatty acid metabolism	0.736	0.000037	416
Na+/PO$_4$ transporter	PF02690	inorganic nutrient transport	4.54	0.000037	5963

889
890 *From Bachy *et al*. (2018)
891 [a]Tricarboxylic acid cycle
892 [b]Electron transport chain
893 [c]Triacylglycerol
894

Table 2. Gene matches to *psr*1 in *Micromonas pusilla* CCMP1545 as originally discovered in the Joint Genome Institute (JGI) Integrated
Microbial Genomes (IMG) database and the corresponding gene in the updated gene model for CCMP1545 (MBARI_models (ver 1)). The *M.*
pusilla CCMP 1545 database in IMG contained 10,660 sequences and 4,795,637 letters. The top-scoring gene was considered to be a *psr*1-
like gene and the JGI gene ID is shown in parentheses. The corresponding *psr*1-like gene in the updated gene model is shown for reference,
and includes the JGI gene ID, the gene description, and the E-value from the BLAST search (see Methods) using the top scoring gene
(613233) from CCMP1545.

IMG Gene ID	Locus tag	Bit score	E-value
2615011133 (JGI 61323)	MicpuC2.est_orfs.1_306_4269596:1	90.5	2×10^{-19}
2615008475	MicpuC2.EuGene.0000130210	74.3	4×10^{-14}
2615010887	MicpuC2.est_orfs.10_1861_4270447:1	70.5	3×10^{-13}
2615007841	MicpuC2.EuGene.0000090108	67.8	4×10^{-12}
JGI ID 360	Myb domain-containing protein		8×10^{-156}

Ratio of cell normalized metabolite concentrations
(P-deficient:P-replete)

(a)

Micromonas pusilla CCMP1545: 2615011133

MTRMSGIEDDSFSYSFLLISGQLRDPTRLTPRPQTSSAPLGAGGGAAAASGMNTSGTFWPSDLDLDLESSDFLD
SLLLGGEMMTTQQQHHHAGDERLGHGPVPLGARAAGEDHDPRVVLAERDAGDGLLPRALADERREPVARA
VVASNNN**KQRLRWTPELHKMFVDAVKRLGGLDLATPKGIMQLMDVEGMSIQHVKSHLQKYRLQDSGGGAS**
EFRVSPDASASGKRPRSEEDDAGGNGKDGNNSAGKTRRRPSAAERSAARLRAAEEKAREERDAARSMAAAA
AAAAAAAVERQNVELALLTGDAAGARDALEMSSHHAASHVGGAYDDVLDLVAGPGESGDGDGAWGDVLH
DHAIVGAAGDDVGLVDDVGSDPEAAAA**MLKQLELQKKLHEHLMSQRRLQQQVEAHGVYLETILDQQK**RRR
GVE

(b)

Figure showing multiple sequence alignment with columns labeled SHLQKyR and LHEQL.

Species	Alignment (SHLQKyR region)	Pos	LHEQL	Pos
Arabidopsis thaliana NC_003075.7	PKGVLKIMKVE-G---LTIYHVKSHLQKYR	278	LH-EQL	331
Volvox carteri XP_003054945.1	PKGILKLMCLE-G---LTIYHIKSHLQKYR	326	LH-EQL	457
Micromonas pusilla CCMP 1545 2615011133 (IMG)	PKGIMQLMDVE-G---MSIQHVKSHLQKYR	205	LH-EHL	396
Ostreococcus lucimarinus 640420317 (IMG)	PKGIMHIMAMS-G---MTIQHIKSHLQKYR	170	LH-DQL	320
Micromonas sp. RCC 299 XP_002504000.1	PKGIMQLMEVD-G---MTIQHVKSHLQKYR	240	LH-AQL	416
O. tauri 2507574032 (IMG)	PKGIATLMTTS-G---MTLQHIKSHLQKYR	106	LH-DQL	241
GOS marine metagenome EBR92606.1	PKRILDLMGVQ-G---LTRENVASHLQKYR	54	PDGGMM	201
GOS marine metagenome EBU22100.1	PKGIMHIMAMS-G---MTIQHIKSHLQKYR	67	- - - - - -	84
GOS marine metagenome EBY01923.1	PKGIMSLMTTA-G---MTLQHIKSHLQKYR	133	LHSMDA	223
Emiliana huxleyi XP_005765367.1	PQAISQLMNCE-GEGAPTRQNIKSHLQKYR	161	LE- - - -	212
GOS marine metagenome EDD34186.1	PQAIRQLMGCKTEEEAPTRQNIKSHLQKYR	228	LH-EQQ	327
Chlamydomonas reinhardtii NW_001843857.1	PKGILKLMGVD-G---LTIYHIKSHLQKYR	240	LH-EQL	403
Chlorella variabilis 2507982799 (IMG)	PKRILDLMNVE-G---LTRENVASHLQKYR	253	- - - - - -	334
MMETSP0052 *Polymella parva*	PKGIVTLMNVR-E---ITIYHVKSHLQKYR	239	TANATL	425
MMETSP0063 *Chlamydomonas euryale*	PKKILEIMQVE-D---LTRENIASHLQKYR	293	GG- - - -	361
MMETSP1127 *Dunaliella tertiolects*	PKGILKLVNSE-G---LTIYHIKSHLQKYR	247	FQGAGG	335
MMETSP1460 *Bathycoccus prassinos*	PKGVVELMRVQ-G---VTIPHVKSHLQKYR	192	LH-EQL	452
MMETSP1106 Mantoniella antarctica	PKGIMQLMGVT-G---MTIQHVKSHLQKYR	237	SA-ARI	332
MMETSP1403 *M. pusilla* CCMP 1723	PKGIMQLMEVE-G---MTIQHVKSHLQKYR	238	LH-EQL	413
MMETSP0384 *Alexandrium tamarense*	PKTILQLMNVE-G---MTRENVASHLQKYR	97	MG- - - -	176
MMETSP1369 *Symbiodinium* sp. C1	PKNIMQEMNVE-G---LTRENVASHLQKYR	113	- - - - - -	110
TARA MATOU-v1_62935935	PKGIMHIMAMS-G---MATIQHIKSHLQKYR	57	- - - - - -	74
TARA MATOU-v1_45631074	PKGVVELMRVQ-G---VTIPHVKSHLQKYR	88	LH-EQL	348
TARA MATOU-v1_79753402	PKLILRLLAVP-G---MTIYHVKSHLQKYR	146	LQ-EQL	243
TARA MATOU-v1_83305352	PKFIMRLLAVP-G---MTIYHVKSHLQKFR	87	LR-EQL	158

(a) Normalized *psr1* abundance (proportion of *Micromonas* transcripts)

10^{-5} 10^{-4} 10^{-3}

(b) Normalized *psr1* abundance

10^{-5} 10^{-4} 10^{-3}

Normalized abundance of *Micromonas* (proportion per sample)

PO_4^{3-} (µM)

(c) Gene abundance (proportion *Micromonas* transcripts)

Absent
Present

* p<0.005

Fumarase Synthase