A phosphate starvation response gene (*psr*1-like) is present and expressed in
***Micromonas pusilla* and other marine algae**

Cara L. Fiore[1,*], Harriet Alexander[2], Melissa C. Kido Soule, Elizabeth B. Kujawinski

Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic
Institution, Woods Hole, MA, USA

[1]Current address: Biology Department, Appalachian State University, Boone, NC, USA

[2]Current address: Department of Biology, Woods Hole Oceanographic Institution, Woods
Hole, MA, USA

*Corresponding author: Cara Fiore
572 Rivers St.
Boone, NC 28607, USA
fiorec@appstate.edu
15

16 Running page head: psr1 in marine phytoplankton

Abstract

Phosphorus (P) limits primary production in regions of the surface ocean, and many

plankton species exhibit specific physiological responses to P-deficiency. The metabolic

response of *Micromonas pusilla*, an ecologically relevant marine photoautotroph, to P-

deficiency was investigated using environmental metabolomics and comparative

genomics. The concentrations of some intracellular metabolites were elevated in the P-

deficient cells (e.g., xanthine, inosine) and genes involved in the associated metabolic

pathways shared a predicted conserved amino acid motif in the non-coding regions of

each gene. The presence of the conserved motif suggests that these genes may be co-

regulated, and the motif may constitute a regulatory element for binding a transcription

factor, specifically that of Psr1 (Phosphate starvation response,), first described in the

alga, *Chlamydomonas reinhardtii*. A putative phosphate starvation response gene (*psr*1-

like) was identified in *M. pusilla* with homology to well characterized *psr*1/*phr*1 genes in

algae and plants, respectively. This gene appears to be present and expressed in other

marine algal taxa (e.g., *Emiliania huxleyi*) in field sites that are chronically phosphorus-

limited. Results from the present study have implications for understanding

phytoplankton taxon-specific roles in mediating P cycling in the ocean.

Key words: *Micromonas pusilla*, phosphate stress response, marine algae, metabolomics,
dissolved organic matter, biological oceanography

1. Introduction

Phosphorus (P) is a critical element for life, found in lipid membranes, genetic material, and energy storage compounds. Because most marine microorganisms preferentially take up P as inorganic phosphate (PO_4^{3-}), concentrations of dissolved PO_4^{3-} are generally below 1 µM in the surface ocean, (Karl 2014) limiting phytoplankton productivity (Tyrell 1999).

Marine phytoplankton respond to chronic P-deficiency by investing resources in P uptake via inorganic and organic compounds (Chung et al. 2003, Lin et al. 2016), remodeling cellular metabolism and structures (Berdalet et al. 1994, Van Mooy et al. 2009, Martin et al. 2011, Shemi et al. 2016, Kujawinski et al. 2017), and storing P (Dyhrman et al. 2012, Martin et al. 2014). Different mechanisms for enacting these responses have evolved in widely distributed and often sympatric phytoplankton taxa. For example, within the diatom, *Phaeodactylum tricornutum*, proteomics indicated broadly depressed metabolic activity as a response to P-starvation, with down-regulated energy metabolism, amino acid metabolism, nucleic acid metabolism, and photosynthesis, and up-regulated protein degradation, lipid accumulation, and photorespiration (Feng et al. 2016). In contrast, the prymnesiophytes *Prymnesium parvum* and *Emiliania huxleyi,* and the dinoflagellates *Prorocentrum donghaiense* and *Amphidinium carterae*, maintained energy-generating processes (i.e., photosynthesis) and carbon fixation in response to P-deficiency (Lai et al. 2015, Li et al. 2016, Rokitta et al. 2016, Shi et al. 2017). Interestingly, *P. donghaiense* increased nitrate assimilation under P-deficiency (Shi et al. 2017), whereas *E. huxleyi* reduced nitrate uptake with a tight coupling between P and nitrogen (N) pools (Rokitta et al. 2016). These varied strategies all minimize non-critical

P utilization and maximize P uptake, and thus play important roles in structuring

phytoplankton assemblages in the oceans (Dyhrman et al. 2006, Dyhrman et al. 2009,

Martin et al. 2011, Rokitta et al. 2016, Guo et al. 2018).

The cosmopolitan picoeukaryotic groups *Micromonas* and *Ostreococcus* spp. are

members of the "green" lineage (Chlorophyta), a monophyletic group including land

plants, and are important marine primary producers (Li 1994). Recent culture

experiments with *Micromonas pusilla* (Hoppe et al. 2018) and *M. pusilla* Lac38 (Maat et

al. 2014) indicated that these strains are likely to be successful under conditions of

increased acidification and low nutrient (P) availability. This success may in part, be due

to higher efficiency of carbon sequestration for growth under high pCO$_2$ conditions

and/or the ability to remodel photosystem proteins under nutrient-limited conditions

(Maat *et al.*, 2014; Hoppe *et al.*, 2018, Guo *et al.*, 2018). Additionally, similar to other

phytoplankton (Chung et al. 2003), *M. pusilla* may shift lipid composition to increase

non-P containing lipids (Maat et al. 2016, Guo et al. 2018), and up-regulate genes and

proteins for P-transporters (Whitney & Lomas 2016, Guo et al. 2018) and genes for

polyphosphate accumulation (Bachy et al. 2018). However, some of the broader

physiological response mechanisms of *M. pusilla* and other prasinophytes to P-deficiency

have yet to be fully explored.

The Chlorophyta share common ancestry with land plants, stemming from a

single endosymbiotic event, in contrast to more distantly related phytoplankton groups

such as diatoms and haptophytes, which are hypothesized to have undergone multiple

endosymbioses (Falkowski et al. 2004; Lewis & McCourt 2004). Consequently, their

physiological response to nutritional stressors, like P-deficiency, may share more traits

with land plants than with other algal taxa. Indeed, P-deficiency is common in lakes and

terrestrial systems (Elser et al. 2007). The physiological response of model chlorophytes

such as the mustard plant, *Arabidopsis thaliana*, and the freshwater green alga,

Chlamydomonas reinhardtii, to P-deficiency led to the identification of a phosphate

starvation response gene described as *psr*1 in *C. reinhardtii* (Wykoff et al. 2009) and as

*phr*1 in plants (Rubio et al. 2001). This gene encodes for a transcription factor (TF), a

protein that binds to specific DNA regions to activate or repress transcription of one or

more genes. The specific region of DNA to which TFs bind is typically characterized by

a short repetitive nucleotide sequence, or motif, found up- or downstream of a given gene

or within introns (Barrett et al. 2012; Franco-Zorrilla et al. 2014). In *A. thaliana,*

regulatory motifs for the Phr1 protein were more abundant in known P-responsive genes

than in the rest of the genome (Müller et al. 2006), linking Phr1 to the regulation of P-

responsive genes. Phr1/Psr1 TFs have not been described, to our knowledge, in marine

algae.

Here, we use a combined metabolomic and comparative genomic approach to

investigate the impact of P-deficiency on the physiology of *Micromonas pusilla*

CCMP1545. Many metabolites are produced by metabolic pathways under genetic

regulatory control (e.g., de Carvalho & Fernandes 2010, Markou & Nerantzis 2013); thus,

our approach complements recent physiological, transcriptional, and proteomic work

(Hoppe et al. 2018, Guo et al. 2018) and provides mechanistic insight into the

physiological response to P-deficiency and its underlying genetic regulation. We used a

targeted metabolomics approach to analyze the suite of intracellular and extracellular

molecules produced by *M. pusilla*. In combination, we employed a comparative genetics

approach to identify 1) a *psr*1-like gene in *M. pusilla* and other prevalent marine

phytoplankton species and 2) a potential DNA-binding site of the Psr1-like protein in

putatively P-responsive genes. We found evidence for the expression of *psr*1-like genes

in *M. pusilla* and other marine algae under P-deficient conditions in cultures and in the

field. These results have implications for better understanding and predicting the

metabolic response of diverse phytoplankton groups to P-deficiency.

2. Materials and Methods

2.1 Culture of Micromonas pusilla *CCMP 1545*

All glassware was acid-cleaned and combusted at 450°C for at least 4 h. We grew

an axenic culture of *Micromonas pusilla* CCMP1545 from the National Center for

Marine Algae and Microbiota (Boothbay, ME, USA) in L1-Si media

(https://ncma.bigelow.org/algal-recipes) with 0.2-μm filtered (Omnipore filters; Millipore,

MA, USA) autoclaved seawater from Vineyard Sound (MA, USA). We maintained the

cultures at 22°C under a 12:12 light:dark regime (84 μmol m^{-2} s^{-1}). After acclimation to

these culture conditions, we split the culture into two parallel cultures of L1-Si media

amended with (a) 36 μM phosphate (P-replete) or (b) 0.4 μM phosphate (P-deficient). We

maintained these parallel cultures for three generations prior to this experiment (Fig. S1

a,b). At the start of the experiment, we inoculated 9 flasks of each media type with

exponential-phase cells to achieve 320 ml and ~300,000 cells in each flask (33 ml P-

replete; 37 ml P-deficient), with three cell-free control flasks. We grew cultures for two

weeks (Fig. S2a, b, c) and removed samples approximately 1 hr into the light cycle at

experiment initiation (T_0; day 0), in late exponential growth phase (T_1; day 4 P-deficient;

day 6 P-replete), and in stationary phase (T_2; day 5 P-deficient; day 13 P-replete).

At each sampling point, we removed 1 ml for cell counts and chlorophyll

fluorescence, 30 ml for total organic carbon (TOC; see Methods in the supplement), and

20 ml of filtrate (see below) for nutrients. We monitored cell abundance daily by flow

cytometry (Guava, Millipore) and assessed photosystem II efficiency by measuring the

variable and maximum chlorophyll fluorescence (F_v/F_m) using fluorescence induction and

relaxation (FIRe; Satlantic LP, Halifax, Cananda). We used chlorophyll *a* (692 nm) and

side scatter of *M. pusilla* cultures to optimize flow cytometry settings. We assessed

potential bacterial contamination by viewing DAPI-stained cells at each time point (see

extended Methods in the supplement).

2.2 Metabolite extraction and instrument methods

Metabolite extraction and analytical methods are optimized for semi-polar

compounds with a size range of 100-1000 Da, capturing many primary metabolites such

as organic acids, vitamins, and nucleosides. We processed cultures for intracellular and

extracellular metabolite extraction as described previously (Fiore et al. 2015, extended

Methods in the supplement). For intracellular metabolites, we collected cells via

filtration on 0.2 μm PTFE filters (Omnipore, Millipore) and stored them at -80°C.

Metabolites were extracted in 500 μl of cold 40:40:20 acetonitrile:methanol:water with

0.1 M formic acid. Extracts were reduced to near dryness under vacuum centrifugation

and reconstituted in 500 μl of 95:5 MilliQ water: acetonitrile with deuterated biotin (d2-

biotin, final concentration 0.05 μg ml^{-1}) added as a high-performance liquid

chromatography (HPLC) injection standard; 100 μl of the extract was used for targeted

metabolomics analysis. We extracted extracellular metabolites using solid phase

extraction (SPE) with 1 g/6 cc PPL cartridges (BondElut, Agilent, Santa Clara, CA,

USA) as described previously (Dittmar et al. 2008, Longnecker 2015).

Intracellular and extracellular extracts were used in targeted metabolomics

analysis with HPLC (Thermo PAL autosampler and Accela pump) coupled to tandem

mass spectrometry (TSQ Vantage, Thermo Fisher Scientific, MA, USA) via a heated

electrospray ionization (H-ESI) source operated in both positive and negative ion modes.

All 62 standards were analyzed using optimized selected reaction monitoring (SRM)

conditions (XCalibur 2.0) in the intracellular samples (Kido Soule et al. 2015), while 23

metabolites with high SPE recovery (Johnson et al. 2017) were analyzed in the

extracellular samples. Concentrations of metabolites were calculated based on a five-to-

seven point manually-curated external calibration curve (0.5-500 ng ml^{-1}). A pooled

sample for quality control consisted of 10 parts intracellular extract and one part

extracellular extract of each experimental sample and was analyzed every ten injections

(see extended Methods in the supplement). A laboratory-fortified pooled sample

containing a standard mix was analyzed at the start, middle, and end of the analytical run

to assess matrix effects and analytical quality. Chromatographic peaks derived from

calibration standards, pooled samples, and experimental samples were manually assessed

for quality and metabolite concentrations were exported to Excel. To facilitate

comparison with other studies, we also report cell-normalized ratios of the P-deficient to

P-replete metabolite concentrations. All metabolomics data are available in MetaboLights

(http://www.ebi.ac.uk/metabolights/) with accession number MTBLS295.

While both intracellular and extracellular metabolites were quantified in our

experiment, the results and discussion will focus primarily on changes in the intracellular

metabolite profile in response to P-deficiency, highlighting compounds that were most

relevant to the discovery of the *psr*1-like gene. We further focus the intracellular

metabolite comparison on the exponential growth phase (T_1) to capture clear changes in

metabolite profiles between treatments and facilitate comparison to previous studies

(Whitney & Lomas, 2016, Bachy et al. 2018). R statistical software (v3.0.2; R Core

Team) was used for statistical analyses. We used a Welch's two sample *t*-test to compare

the cell-normalized concentrations of specific metabolites (log transformed to achieve a

normal distribution) between P-deficient and P-replete treatments.

*2.3 Characterization of the phosphate starvation response (*psr*1) gene and putative DNA-*
binding motifs

We used the *psr*1 gene sequence from *C. reinhardtii* (Wykoff et al. 1999, NCBI

XM_001700501.1) to query the genome of *M. pusilla* using the BLAST option through

Integrated Microbial Genomes (IMG, blastx, e-value: $1e^{-5}$). Conserved domains within

putative sequences were characterized using NCBI conserved domain search (CD-search).

We then used the putative *M. pusilla psr*1-derived amino acid sequence to query several

databases using BLAST (tblastn, e-value: $1e^{-5}$), including: a re-assembly of the Marine

Microbial Eukaryote Transcriptional Sequencing Project (MMETSP) (Johnson *et al*.,

2018), NCBI non-redundant protein sequences (nr), the NCBI metagenome proteins (env

nr), the *Tara* Oceans eukaryotic unigenes (Carradec, Pelletier et al. 2017), and several

eukaryotic algal genomes from IMG (Methods in the supplement). For the *Tara* Oceans

data, we selected only the *psr*1-like unigenes from the metatranscriptome assemblies that

were assigned the taxonomic classification of *Micromonas*. Within each sample, we

normalized the expression of the *psr*1 unigenes to the total abundance of all unigenes

taxonomically classified as *Micromonas* (Methods in the supplement). Visualization of

the *Tara* Oceans queries were performed using the basemap library in Python and custom

Python scripts (see extended Methods in the supplement). For several genes of interest,

we compared the distribution of gene expression to the expression of *Micromonas psr*1-

like genes in the *Tara* Oceans metatranscriptome using a Kolmogorov-Smirnov test. We

tested a relationship between phosphate concentration and normalized expression of

*Micromonas psr*1-like genes from *Tara* Oceans metatranscriptomes by comparing the

abundance of *Micromonas psr*1-like transcripts above and below the mean concentration

of phosphate using a Student's *t*-test.

We manually checked significant matches for all nucleotide sequences that were

collected from database-based (NCBI, IMG, MMETSP, *Tara* Oceans) BLAST searches

using the *psr*1-like gene from *M. pusilla*. First, we used conserved domain analysis

(Marchler-Bauer et al. 2017) to assess domain similarity to *psr*1 and second, we used the

sequences with both characteristic domains of the *C. reinhardtii psr*1-derived amino acid

sequence in a multiple sequence alignment to the *C. reinhardtii psr*1-derived amino acid

sequence. Sequences that did not align well, and specifically did not align with the myb

coiled-coil domain, were removed from further consideration. Sequence alignment and

phylogenetic analysis was performed in MEGA 5.2.1 (Tamura et al. 2011) using

MUSCLE (Edgar 2004) with default settings in MEGA. Phylogenetic reconstruction of

the *psr*1 gene was performed in MEGA using neighbor-joining, maximum parsimony,

and maximum likelihood algorithms and bootstrap replication 500 times. For visual

clarity, a subset of significant sequence matches was used to create the trees, including

representative sequences from each major taxonomic group (see Fig. S4 for the full tree).

The tree topology for the different phylogenetic reconstructions was similar for nodes

with high bootstrap values ($> 50\%$, data not shown). We present the maximum likelihood

tree because this algorithm (Jones-Taylor-Thornton model, default maximum likelihood

settings in MEGA) uses the most information in its alignments and typically has high

accuracy (Felsenstein 1981, Kishino et al. 1990).

We searched genes of *M. pusilla* CCMP1545 for a regulatory binding element, or

motif, where the Psr1-like protein would bind to activate or repress gene transcription.

We first searched within a now-archived gene model of *M. pusilla* CCMP1545

(FrozenGeneCatalog_20080206 (ver_1), Table S1), then within the updated gene model

from Bachy et al. (2018) (*Micromonas pusilla* CCMP1545/MBARI_models(ver_1),

Table 1). Specifically, we analyzed gene sequences of interest for a regulatory element,

or motif, using the motif-based sequence analysis tools Multiple Em for Motif Elicitation

(MEME Suite) 4.10.1 (Bailey et al. 1994). The genes of interest (Table 1) were selected

based on metabolites that were elevated or depressed in concentration in the P-deficient

cultures though not necessarily statistically different between treatments (i.e., malate,

several amino acids, vitamins, Fig. 1), as well as gene expression information from

Bachy et al. (2018). These included genes related to central carbon metabolism, lipid

metabolism, vitamin metabolism (pantothenate, folate), and nucleotide metabolism

(aspartate). We included genes within the carbohydrate and lipid metabolic pathways

based on published work (Hockin et al. 2012, Goncalves et al. 2013) as well as the gene

for proline oxidase (POX), a key enzyme up-regulated in *E. huxleyi* in response to N- and

P-deficiency (Rokitta et al. 2014, Rokitta et al. 2016). After initial motif discovery, we

optimized our putative motif by varying motif width and number within genes of interest

and within genes that were not expected to be regulated under P-deficiency, including

asparagine synthetase (JGI gene ID: 9681548) and acetyltransferase like/FAD linked

oxidase (9687568)). We attempted to identify motifs using the motif comparison tool

(Tomtom; Gupta et al. 2007) against the *A. thaliana* database (Protein Binding

Microarray (PBM) motifs, (Franco-Zorrilla et al. 2014)) and significant matches were

used to guide the optimization of motif discovery settings and inclusive sequences.

Using the selected sequences and optimized parameters, we searched for motifs

that are overrepresented in the query sequences ("positive") relative to a set of

background sequences ("negative"), the latter defined using the genome scaffolds. We

used positive and negative sequences to create a position specific prior (PSP) distribution

file (Bailey et al. 1994), for further discriminative motif discovery within *M. pusilla* gene

sequences of interest (parameters: minw = 5, maxw = 10, nmotifs = 3, mod = oops (one

occurrence per sequence)). Each gene sequence consisted of concatenated 500

nucleotides upstream and downstream of the gene as well as untranslated regions (UTRs)

and intronic regions, thereby excluding the coding regions.

Two recent publications provided transcriptomes for testing our hypotheses

regarding expected differentially expressed (DE) genes in P-deficient and P-replete

conditions. Whitney & Lomas (2016) published transcriptomes of *M. commoda* (formerly

M. pusilla) RCC299, while Bachy et al. (2018) published transcriptomes of *M. pusilla*

CCMP1545, under P-deficient or limiting (respectively) and P-replete conditions. Genes

discussed as DE here were defined by each study (Whitney & Lomas 2016, Bachy et al.

2018). We note that the each *Micromonas* experiment used slightly different growth

conditions and were sampled at different stages of growth. Thus, we focus our

comparisons on work in the same strain (*M. pusilla* CCMP1545, Bachy et al. 2018), with

brief complementary discussions about *M. commoda* (see extended Results in the

supplement). Bachy et al. (2018) used semi-continuous cultures grown under continuous

light and were grown with a moderately higher concentration of P (2.5 µM) than in our

experiment (0.4 µM P). Despite these differences, the transcriptional data of *M. pusilla*

under P-limitation provided a preliminary test of our hypotheses derived from the

metabolomics analyses with *M. pusilla* CCMP1545 and *in silico* genomic analysis. We

further reasoned that a signal observed across species and experiments would likely

represent a conserved and biologically meaningful response to P-deficiency.

3. Results

3.1 Metabolic response of Micromonas pusilla *CCMP1545 to phosphorus deficiency*

Metabolite concentrations were generally lower under P-deficiency in the

extracellular (Fig. S3) fraction relative to the P-replete cultures, while there was a mixed

response in metabolite concentrations in the intracellular fraction (Fig. 1). The

extracellular metabolite concentrations represent cumulative changes in the media over

time. In stationary phase, nearly all of the quantified extracellular metabolites were less

abundant in the P-deficient samples than the P-replete samples (Fig. S3). In contrast,

several of the intracellular metabolites were more abundant during exponential growth

(T_1) under P-deficiency, including nucleobases (xanthine, adenine), the nucleoside

inosine, amino acids (glutamine, aspartate), the dicarboxylic acid malate, and vitamins

(thiamin, pantothenate, folate) (Fig. 1). None of the extracellular metabolites and only a

few intracellular metabolites showed statistically significant differences between

treatments, including: (a) the nucleobase xanthine, significantly elevated under P-

deficiency (*t*-test, p = 0.01), and (b) the amino acid derivatives, *N*-acetylglutamate and

tryptamine, significantly higher in P-replete samples (*t*-test, p = 0.01, p = 0.02,

respectively) (Fig. 1). We observed decreases in the intracellular ratios of purine

nucleosides (guanosine, xanthosine) to their nucleobases (guanine, xanthine) under P-

deficiency at T_1 (Fig. 2). The decrease in the xanthosine:xanthine ratio was statistically

significant (*t*-test, p = 0.03). The change in the guanosine:guanine ratio could not be

statistically tested as guanine was below detection in two of three P-replete cultures (Fig.

2). Intracellular malate, succinate, and citrate exhibited divergent responses, although

their abundances were not significantly different between treatments (Fig. 1). Malate was

more abundant in P-deficient cells, while succinate concentrations were similar between

treatments and replicates and citrate was detected in only one of three replicates in each

treatment (Fig. 1). We noted similarly varied responses to P-deficiency in the purine

nucleosides xanthosine and inosine, with invariant xanthosine abundances but higher

average concentrations and variability of inosine (Figs. 1,2).

Metabolite concentrations are, in part, determined by genetic regulation of the

enzymes producing or degrading them and we hypothesized that the varied metabolite

responses could provide insight into the regulation of these genes. The description of a

phosphate starvation response (*psr*1) gene containing a myb DNA-binding domain

(Wykoff et al. 1999), motivated us to investigate the presence of *psr*1 and a myb-like

motif in genes that could be regulated by the *psr*1 gene product in *M. pusilla* and other

marine algae.

*3.2 Characterization of the phosphate starvation response (*psr*1) gene in marine algae*

We found four statistically-significant sequences (Table 2) in response to

querying the now-archived *M. pusilla* CCMP1545 genome with the *psr*1 gene from *C.*

reinhardtii (strain cc125, NCBI accession AF174532). Only one of these sequences (JGI

ID 61323) had the two characteristic myb domains of the *psr*1 gene in *C. reinhardtii*

(myb-like DNA-binding domain, SHAQKYF class (TIGR01557) and myb predicted

coiled-coil type transfactor, LHEQLE motif (pfam PF14379)) (Fig. 3a). The glutamine-

rich regions, characteristic of TF activation domains, and putative metal binding domain

originally described for *psr*1 in *C. reinhardtii* (Wykoff et al. 1999) were not detected in

the putative *M. pusilla psr*1 gene, thus we refer to this as a *psr*1-like gene. Conserved

domain analysis (Marchler-Bauer et al. 2017) revealed two other domains in the putative

*psr*1-like-derived amino acid sequence from *M. pusilla*: 1) PLN03162 super family

(NCBI cl26028), a provisional golden-2 like transcription factor, and 2) a DUF390 super

family domain (NCBI cl25642), comprising proteins of unknown function. A comparison

of the psr1-like gene in *M. pusilla*, *M. commoda*, *C. reinhardtii*, and *A. thaliana* is

provided in Table S2.

We identified putative *psr*1-like genes or transcripts in cultured isolates as well as

field datasets (Figs. 3b, 4, 5, Table S3), representing diverse taxa such as Dinophyta (e.g.,

the symbiotic clade C *Symbiodinium* sp.), and Haptophyta (e.g., the coccolithophore *E.*

huxleyi). In our phylogenetic analysis (Fig. 4, Fig. S4 for full tree and extended Results

in the supplement), representative sequences clustered by taxonomic origin only to some

extent. For example, the prasinophytes, including *M. pusilla* CCMP1545, generally

clustered together, as did the haptophytes, including *E. huxleyi*. However, other clusters

included a mix of taxonomic groups such as the dinoflagellate *Alexandrium tamarense*,

the glaucophyte *Gloeochaete wittrockiana*, and the green alga *Chlorella vulgaris* (Fig. 4).

Transcripts from MMETSP and *Tara* Oceans were generally short sequences and

often contained one of the two characteristic domains for *psr*1-derived amino acid

sequences (extended Results in the supplement). Field-derived transcripts of the *psr*1-like

gene were geographically dispersed in the *Tara* Oceans dataset with the highest relative

expression in surface samples and in the North Atlantic Ocean and Mediterranean Sea

(Fig 5a). The majority of these transcripts from the *Tara* Oceans dataset occurred in

samples with low phosphate concentrations (≤ 0.5 µM), but there was no significant

relationship between *psr*1 expression and phosphate concentration (*t*-test, $p < 0.05$; Fig.

5b). We observed a significant positive relationship (Kolmogorov-Smirnov, $p < 0.05$)

between the presence of *Micromonas psr*1-like genes and the relative expression of

fumarase, a TCA cycle gene in *Micromonas*, which was up-regulated in cultures of *M.*

pusilla (Bachy et al. 2018) and *M. commoda* (Whitney & Lomas 2016) under P-

deficiency (Fig. 5c, see extended Discussion in the supplement). We did not find genes

with significant homology to the *M. pusilla* putative *psr*1-like gene, in genomes or

transcriptomes, within diatoms, the Heterokonta superphylum or the cryptophytes.

3.3 Putative regulatory element in Micromonas pusilla *genes of interest*

MEME analysis yielded a significant motif in *M. pusilla* genes involved in central

carbon metabolism, lipid metabolism, and nucleotide metabolism (Fig. S5). The

conserved motif is similar in nucleotide sequence to several TF-binding sites in *A.*

thaliana (Table S4, Fig. S6; Franco-Zorrilla et al. 2014), including myb-like TF-binding

motifs. The presence of a myb-like TF-binding motif is notable as the Psr1 protein

contains a myb-like DNA binding domain (Wykoff et al. 1999). Several iterations of this

analysis using genes not tied to the metabolites of interest yielded either no significant

motif or motif sequences lacking similarity to myb-like domains. We observed a similar

conserved motif in author-defined DE transcripts (Bachy et al. 2018) in *M. pusilla*

CCMP1545 under P-deficiency (Table 1, Fig. S5). While the actual nucleotide sequence

recognized by a TF must be experimentally determined (Franco-Zorrilla et al. 2014), the

presence of a significant motif in each model and only in the expected genes, indicates

that this conserved motif is likely biologically significant in *M. pusilla*. Specifically, this

motif, with similarity to myb-like domains in *A. thaliana*, may represent a regulatory

element where the Psr1-like protein binds.

3.4 Comparison of metabolomics-based predictions of P-responsive genes in M. pusilla

CCMP1545 to transcriptomics analysis

We compared the author-defined DE transcripts in *M. pusilla* CCMP1545 to gene

predictions based on our metabolomics data. As Bachy and colleagues (2018) used a

more recent gene model than our original analysis, we confirmed the identity of the *psr*1-

like gene in the updated model (CCMP1545/MBARI_models(ver_1), Table 2). We then

analyzed a similar set of genes for a regulatory element where the Psr1-like TF might

bind (Fig. S5), including genes that were DE in the P-deficient transcriptome relative to

the replete (Table S1 from Bachy et al. 2018). Under P-deficiency, the *psr*1-like gene

(JGI gene ID 360) was highly up-regulated (log-2 fold-change = 4.3, $q = 0.00037$) and

nearly all of the predicted P-responsive genes were DE (Table 1). For example, *M.*

pusilla up-regulated many genes in the TCA cycle but not pyruvate carboxylase (gene ID

6984). Genes involved in nucleotide and lipid metabolism (e.g., nucleoside phosphatase,

long-chain acyl-coA synthetase) were DE and contained a conserved motif with

similarity to myb-like regulatory element in *A. thaliana* (Table 1, Fig. S5). Lastly, the

POX gene described in *E. huxleyi* (Rokitta et al. 2016) and observed in the archived gene

model for *M. pusilla*, was absent, or not similarly annotated in the updated gene model.

Instead, a copper amine oxidase (gene ID 5863) contained a significant conserved motif

(Table 1, Fig. S5) and was DE under P-deficiency (Bachy et al. 2018).

We conducted a similar analysis with published transcriptome data from *M.*

commoda (Whitney & Lomas 2016). The *psr*1-like gene (JGI gene ID 60184) in *M.*

commoda was significantly up-regulated in the P-deficient treatment (defined by Whitney

& Lomas 2016, Table S5), and we found a significant conserved motif in 18 DE genes.

The significant motif identified in *M. commoda* differed in sequence from that discovered

in *M. pusilla* and occurred in a different set of genes (see extended Results in the

supplement).

4. Discussion

Our combined metabolomics and comparative genomic analyses of the response

of *M. pusilla* CCMP1545 to P-deficiency has led to three conclusions: 1) there is a shift

in intracellular metabolite composition, 2) a *psr*1-like gene is expressed in *M. pusilla* and

other marine phytoplankton, and 3) the genes regulated by the Psr1-like protein may

differ amongst algal species.

4.1 Micromonas pusilla *exhibits a metabolic shift in response to P-deficiency*

The general decrease in concentration of extracellular metabolites of P-deficient

cells indicates reduced release of these central metabolites under P-deficiency. This is

expected as nutrient-limited microalgae may have lower photosynthetic activity (Loebl et

al. 2010, Halsey et al. 2014, Guo et al. 2018) and hence, produce less metabolic overflow.

In contrast, most intracellular metabolites did not differ significantly in concentration

between treatments. Because these metabolites represent many central metabolic

pathways, it appears that *M. pusilla* does not respond to P-deficiency through a global

decrease in metabolic activity, as was observed for the diatom, *Phaeodactylum*

tricornutum (Feng et al. 2015). Interestingly, we found examples of functionally-related

metabolites that behaved differently. For example, glutamate, proline, and methionine

were higher on average, although variable, under P-deficiency, while phenylalanine and

tryptophan were similar. In the TCA cycle, malate was variable but higher in

concentration on average under P-deficiency, while citrate was lower on average and

succinate was relatively unchanged. The opposing responses of these TCA intermediates

was surprising given that others (e.g., Hockin et al. 2012, Goncalves et al. 2013) have

observed a coordinated response to nutrient limitation in transcripts or proteins involved

in the TCA cycle.

We observed additional non-uniform metabolite responses, with vitamins and

glucosamine-6-phosphate higher on average under P-deficiency, but other P-containing

compounds (e.g. glycerol-3-phosphate) and sugars were depleted. Previous work has

suggested that organisms placed under nutrient stress are capable of diverting carbon flux

between pathways (reviewed in Markou & Nerantzis 2013). Changes in carbon flux

through the cell could result in coordinated abundance shifts in seemingly unrelated

metabolites, as observed in this study. If the shunting of carbon is a coordinated response

within *M. pusilla*, then we would expect genes involved in the synthesis and/or

catabolism of these compounds to be co-regulated, and potentially mediated by a specific

transcription factor. Indeed, in *M. pusilla,* we discovered a gene with significant

homology to the transcription factor phosphorus starvation response gene (*psr*1)

originally described in the freshwater alga *Chlamydomonas reinhardtii* (Wykoff et al.,

1999).

4.2 Presence and expression of the transcription factor gene, psr*1, in marine algae*

In *C. reinhardtii* and in many plants, a *psr*1 gene encodes for a transcription factor

(Psr1) that coordinates the metabolic response to P-deficiency in these organisms

(Wykoff et al. 1999; Mosely et al. 2006). We detected a putative *psr*1-like gene in the *M.*

pusilla genome and in the genomes and transcriptomes of major algal lineages (e.g.,

prasinophytes, dinoflagellates). The *psr*1-like genes that we discovered were generally

not annotated as *psr*1 with the exception of *Ostreococcus tauri* (Derelle et al. 2006),

although to our knowledge the potential role of this gene in the metabolism of *O. tauri* or

other marine algae has not been discussed. The *M. pusilla psr*1-like gene contains two

myb domains similar to those in *psr*1 from *C. reinhardtii* (Wykoff et al. 1999), and

comparable to those in *phr*1 in *A. thaliana* (Rubio et al. 2001). No *psr*1-like gene was

identified by our queries of diatom transcriptomes or genomes, suggesting that a

phosphate starvation response gene akin to *psr*1 is present in diverse but phylogenetically

constrained phytoplankton groups.

We found significant similarity between the *M. pusilla psr*1-like gene and

transcript sequences from the MMETSP and *Tara* Oceans datasets, indicating that this

gene is expressed by marine algae in cultures and in the environment. High-identity hits

to sequences in the global ocean survey (GOS, Rusch et al. 2007) and *Tara* Oceans

datasets underscore this gene's prevalence in the oceans, particularly in regions

characterized by chronically low phosphorus concentrations. Short sequences in some

MMETSP and *Tara* Oceans sequences did not include enough of the C-terminal end to

capture the myb coiled-coil domain. Thus, further investigation is required to confirm

whether the identified transcripts are from *psr*1-like genes. Several GOS sequences,

previously described as protein of unknown function, had homology to both characteristic

Psr1 domains, suggesting that they could be re-annotated as *psr*1-like genes. The GOS

sequences are likely derived from prasinophytes (i.e., *Micromonas* spp., *Ostreococcus*

spp.) and haptophytes (e.g., *E. huxleyi*), highlighting the need to elucidate the role of *psr*1

in these organisms.

4.3 Potential role of Psr1 in the marine algal response to P-deficiency

6   Transcription factors typically recognize a specific motif in DNA where the

protein binds to regulate each gene. If such a motif is observed, one can hypothesize that

a gene containing the motif is regulated by the protein. We observed a conserved and

enriched motif across genes involved in nucleotide biosynthesis, the TCA cycle and

glycolysis, carbon fixation, fatty acid metabolism, and phosphate transport or salvage

(Fig. 6). The conserved motifs discovered in both *M. pusilla* and *M. commoda* had unique

sequences and were present in a slightly different set of genes (see extended Discussion

in the supplement). The motif detected in *M. pusilla* is similar in nucleotide sequence to a

myb DNA-binding domain in *A. thaliana*. This similarity is potentially significant,

because the *psr*1/*phr*1 genes in *C. reinhardtii* and *A. thaliana* contain a myb DNA-

binding domain. Thus, we hypothesize that the conserved motif detected in the

Micromonas species represents binding sites for the Psr1-like transcription factors (TF).

18   The conserved motif was observed only in DE genes under P-deficiency by each

Micromonas species (Bachy et al. 2018, Whitney & Lomas 2016), suggesting that these

genes could be co-regulated. In bacteria, for example, genes within the well-characterized

Pho-regulon all contain specific sequences (the PHO box) where the transcription factor

binds to activate or repress the gene (Santos-Beneit 2015). These PHO box sequences are

distinct among different bacterial species and strains. In *M. commoda*, the discovered

motif was not significantly similar to myb-like DNA-binding motifs in *A. thaliana* and

was not similar to the motif discovered in *M. pusilla*. Although we expected that the myb

domain of the Psr1-like proteins in the two species would interact with similar binding

regions in the genome, *psr*1-like derived amino acid sequences from *M. pusilla* and *M.*

commoda were only 47% similar. TF protein sequences with up to 79% similarity have

been shown to have distinct DNA-binding motif profiles (Franco-Zorrilla et al. 2014), so

the two proteins in different species could reasonably have distinct DNA-binding motifs.

If the Psr1-like TF regulates a different set of genes in the two *Micromonas* species, this

may serve as a niche defining feature between these taxa. As further support for this

hypothesis, the conserved motif was only observed in a subset of genes that were

differentially expressed under P-limitation or deficiency (Whitney & Lomas 2016, Bachy

et al. 2018). The observed differences in the *psr*1-like gene and differences in associated

binding motifs may represent adaptation to distinct environments as *M. pusilla* was

collected in the temperate English Channel, while *M. commoda* was isolated from the

tropics (Worden 2006).

15   The *psr*1-like gene in *M. commoda* and *M. pusilla* was one of the highest DE

genes in each experiment (Whitney & Lomas 2016, Bachy et al. 2018), suggesting that

this potential TF plays a critical role in the P-deficiency response of *Micromonas*.

Moreover, the *psr*1-like gene was highly expressed in two culture experiments with

different growth conditions and sampled at different growth stages, suggesting a broad

physiological relevance for the Psr1-like TF (Fig. 6). In contrast to the transcriptome

results, a recent study describing proteome changes in *M. pusilla* CCMP1545 grown

under P-limited conditions in a bioreactor (Guo et al. 2018), did not observe significant

regulation of the Psr1-like protein (significance defined by Guo et al. 2018). The

discordant results may be due to known differences in regulation between genes and

proteins (Guo et al. 2018) or to different Psr1-like protein behavior under continuous

culturing conditions. Regardless, these results warrant further investigation into the

physiological role of the Psr1-like protein in *M. pusilla*.

Variable and non-uniform dynamics of several TCA cycle metabolites between

treatments suggested that this pathway may be involved in the metabolic response to P-

deficiency in *M. pusilla*. Indeed, nearly all of the genes involved in the end of glycolysis

and in the TCA cycle were up-regulated in *M. pusilla* (Bachy et al. 2018) and these genes

contained the conserved motif that may function as a regulatory element where the Psr1-

like TF could bind. Fumarase, in particular, was upregulated in both *M. pusilla* and *M.*

commoda transcriptomes under P-limitation or deficiency. If fumarase expression is

regulated by the Psr1-like protein, expression of these two genes (fumarase and *psr*1)

should be correlated in field populations. We observed support for this hypothesis within

the *Tara* Oceans dataset, although we observe weaker relationships between Psr1-like

gene expression and other up-regulated genes in either *M. pusilla* or *M. commoda* (see

extended Discusssion in the supplement). In *M. pusilla*, carbon flow through the TCA

cycle may increase as a means to fuel TAG or starch production and to pull potentially

damaging energy away from the photosystems (Norici et al. 2011, Klok et al. 2013).

Similarly, several proteins involved in glycolysis (e.g., pyruvate dehydrogenase) and the

TCA cycle (e.g., succinate dehydrogenase) were found to be up-regulated in another

study with *M. pusilla* under P-limitation (Guo et al. 2018), lending some support to this

hypothesis. Insight from the latter study underscores potential differences in regulation of

genes and enzymes involved in glycolysis and the TCA cycle (Guo et al. 2018). For

example, we found congruency in up- or down-regulation of some proteins from Guo et

al. (2018) that we would expect in *M. pusilla* under P-deficiency based on our combined

comparative genomics and metabolomics approach (i.e., pyruvate kinase, pyruvate

dehydrogenase, oxoglutarate dehydrogenase, and succinate dehydrogenase) but not others

(i.e., fumarase, aconitase, citrate synthase). Analysis of transcript, protein, and metabolite

responses in *M. pusilla* under the same growth conditions will be required to elucidate the

role of Psr1 in this and other taxa.

In addition to TCA cycle intermediates, we observed shifts for several purine

nucleosides between treatments and detected the conserved motif in genes for a

nucleoside phosphatase and aspartate transcarbamylase. These results suggest increased

nucleotide salvage through purine nucleotides and nucleosides, a mechanism described in

M. commoda (Whitney & Lomas 2016) and in other phytoplankton (Dyhrman & Palenik

2003, Kujawinski et al. 2017). Previous phytoplankton work has also highlighted

expression of proline oxidase (POX) (Rokitta et al. 2014, Rokitta et al. 2015) in nutrient

limitation experiments (including P-deficiency) with the haptophyte *E. huxleyi,* where

this enzyme may have a role in stabilizing the mitochondrial membrane and in detecting

cellular nitrogen levels. The POX gene was not present in *M. pusilla* but was up-

regulated in *M. commoda* (Whitney & Lomas 2016, Table S4). In *M. pusilla*, a copper

amine oxidase was up-regulated (Bachy et al. 2018) and contained the conserved motif

that may function as a regulatory element for the Psr1-like TF. Thus, POX and copper

amine oxidase represent gene targets for further investigation.

Maat et al. (2014) noted that both *E. huxleyi* (Borchard et al. 2011) and *M. pusilla*

(Maat et al. 2014) exhibited a similar metabolic response to elevated pCO$_2$ levels and P-

deficient conditions. While there may be multiple factors underlying the response of

these organisms, both contain the *psr*1-like gene which could direct part of a shared

metabolic response to elevated pCO$_2$ and nutrient-limited conditions. While

transcriptomic analysis of *E. huxleyi* was not performed in the study by Borchard et al.

(2011), Rokitta and colleagues (2014, 2016) performed microarray transcriptome analysis

of *E. huxleyi* cells under separate nitrogen- and P-deficient conditions (normal pCO_2).

The transcripts were only partial sequences; however, two up-regulated genes detected by

Rokitta et al. (2016), contained myb-like DNA binding domains (gene IDs: GJ02872,

GJ03978). Targeted experimental work is necessary to elucidate the presence and role of

a *psr*1-like gene in *E. huxleyi*. If there is a common role for the *psr*1-like gene in *E.*

huxleyi and *M. pusilla* in their response to nutrient limitation and elevated pCO_2, this

could have important implications for predicting the physiological response of the wide

range of organisms that contain this *psr*1-like gene to changing environmental conditions.

The potential impact of Psr1-regulated genes on the ecological roles of

Micromonas spp. is unknown, but our results suggest that the Psr1-like protein

coordinates a metabolic shift in these organisms under P-deficiency, altering the

intracellular flow of carbon and other elements. More comprehensive examination of

these metabolic responses, which likely vary to some extent among organisms, will be

paramount to improving models of trophic carbon flow. More experiments are needed to

characterize the structure and role of the *psr*1-like gene in *Micromonas* spp. and other

phytoplankton, including: (1) confirming the presence of the *psr*1-like gene with genetic

experiments, (2) determining the conditions under which the Psr1-like protein is abundant,

(3) identifying the taxon-specific genes affected by the Psr1-like protein, (4) verifying the

interaction of the Psr1-like protein with the hypothesized binding sites, and (5)

comparing the genetic and metabolic responses to P-deficiency between organisms

containing the *psr*1-like and those that do not. Exploring the underlying biology of the

*psr*1 gene will facilitate mechanistic understanding of the complex response of these

organisms to P-limitation and will enhance our ecological and biogeochemical

predictions.

Acknowledgements

The authors would like to thank Krista Longnecker for performing the TOC analysis and
Matthew Johnson and Elizabeth Harvey for assistance with flow cytometry and FIRe
analysis. This research was funded by the Gordon and Betty Moore Foundation through
Grant GBMF3304 to E. Kujawinski.

Literature Cited

Bachy C, Charlesworth CJ, Chan AM, Finke JF, Wong C-H, Wei C-L, Sudek S, Coleman
ML, Suttle CA, Worden AZ (2018) Transcriptional responses of the marine green alga
Micromonas pusilla and an infecting prasinovirus under different phosphate conditions.
Environ Microbiol 20:2898-2912.

Bailey TL, Elkan C (1994) Fitting a mixture model by expectation maximization to
discover motifs in biopolymers. Proc 2nd Int Conf Intelligent Sys Molec Biol, AAAI
Press, Menlo Park, California, p 28-36.

Barrett LW, Fletcher S, Wilton SD (2012) Regulation of eukaryotic gene expression by
the untranslated gene regions and other non-coding elements. Cell Molec Life Sci
69:3613-3634.

Berdalet E, Latasa M, Estrada M (1994) Effects of nitrogen and phosphorus starvation on
nucleic acid and protein content of *Heterocapsa* sp. J Plank Res 16: 303–316.

Borchard C, Borges AV, Haendel N, Engel A (2011) Biogeochemical response of
Emiliania huxleyi (PML B92/11) to elevated CO2 and temperature under phosphorous
limitation: a chemostat study. J Exp Mar Biol Ecol 410:61–71.
25
Carradec Q, Pelletier E, Da Silva C, Alberti A, Seeleuthner Y, Blanc-Mathieu R, Lima-
Mendez G, Rocha F, Tirichine L, Labadie K, Kirilovsky A, et al (2017) A global ocean
atlas of eukaryotic genes. Nature Comm 9:373.

Cavender-Bares KK, Karl DM, Chisholm SW (2001) Nutrient gradients in the western
North Atlantic Ocean: relationship to microbial community structure and comparison to
patterns in the Pacific Ocean. Deep Sea Res Part I: Oceanogr Res Pap 48:2373–2395.

Chung C-C, Hwang S-PL, Chang J (2003) Identification of a high-affinity phosphate
transporter gene in a prasinophyte alga, *Tetraselmis chui*, and its expression under
nutrient limitation. Appl Environ Microbiol 69:754-759.

de Carvalho CCCR, Fernandes P (2010) Production of metabolites as bacterial responses
to the marine environment. Marine Drugs 3:705-727.

Derelle E, Ferraz C, Rombauts S, Rouzé P, Worden AZ, Robbens S, Partensky F, Degroeve S, Echeynié S, Cooke R, et al (2006) Genome analysis of the smallest free-living eukaryote *Ostreococcus tauri* unveils many unique features. Proc Nat Acad Sci 103:11647–11652.

Dittmar T, Koch B, Hertkorn N, Kattner G (2008) A simple and efficient method for solid-phase extracion of dissolved organic matter (SPE-DOM) from seawater. Limn Oceangr Meth 6:230–235.

Dyhrman ST, Ruttenberg KC (2006) Presence and regulation of alkaline phosphatase activity in eukaryotic phytoplankton from the coastal ocean: Implications for dissolved organic phosphorus remineralization. Limnol Oceanogr 51:1381–1390.

Dyhrman ST, Benitez-Nelson CR, Orchard ED, Haley ST, Pellechia PJ (2009) A microbial source of phosphonates in oligotrophic marine systems. Nat Geosci 2:696–699.

Dyhrman ST, Jenkins DB, Rynearson TA, Saito MA, Mercier ML, Alexander H, Whitney LP, Drzewianowski A, Bulygin VV, Bertrand EM, et al (2012) The transcriptome and proteome of the diatom *Thalassiosira psuedonana* reveal a diverse phosphorus stress response. PLoS ONE 7:e33768.

Easlon E, Tsang F, Skinner C, Wang C, Lin S-J (2008) The malate-aspartate NADH shuttle components are novel metabolic longevity regulators required for calorie restriction-mediated life span extension in yeast. Genes Devel 22:931-944.

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucl Acids Res 32:1792–1797.

Elser JJ, Bracken MES, Cleland EE, Gruner DS, Harpole WS, Hillebrand H, Ngai JT, Seabloom EW, Shurin JB, Smith JE (2007) Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecoystems. Ecol Lett 20:1135-1142.

Falkowski PG, Katz ME, Knoll AH, Quigg A, Raven JA, Schofield O, et al (2004) The evolution of modern eukaryotic phytoplankton. Science 16:354-360.

Franco-Zorrilla JM, López-Vidriero I, Carrasco JL, Godoy M, Vera P, Solano R (2014) DNA-binding specificities of plant transcription factors and their potential to define target genes. Proc Nat Acad Sci 111:2367–2372.

Felsenstein J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood approach. J molec evol 17:368–376.

Feng T-Y, Yan Z-K, Zheng J-W, Xie Y, Li D-W, Bala Murugan S, *et al*. 2015. Examination of metabolic responses to phosphorus limitation via proteomic analyses in the marine diatom *Phaeodactylum tricornutum*. Sci Rep 5:10373.

Fiore CL, Longnecker K, Kido Soule MC, Kujawinski EB (2015) Release of ecologically relevant metabolites by the cyanobacterium *Synechococcus elongatus* CCMP 1631.

Environ Microbiol 17:3949–3963.

Goncalves EC, Johnson JV, Rathinasabapathi B (2013) Conversion of membrane lipid acyl groups to triacylglycerol and formation of lipid bodies upon nitrogen starvation in biofuel green algae *Chlorella* UTEX29. Planta 238:895-906.

Guo J, Wilken S, Jimenez V, Choi CJ, Ansong C, Dannebaum R, Sudek L, Milner DS, Bachy C, Reistetter EN, et al (2018) Specialized proteomic responses and an ancient photoprotection mechanism sustain marine green algal growth during phosphate limitation. Nat Microbiol Lett 3:781-790.

Gupta S, Stamatoyannopolous JA, Bailey T, Noble WS (2007) Quantifying similarity between motifs. Gen Biol 8:R24.

Halsey KH, Milligan AJ, Behrenfeld MJ (2014) Contrasting strategies of photosynthetic energy utilization drive lifestyle strategies in ecologically important picoeukaryotes. Metabolite*s* 4: 260-280.

Hockin NL, Mock T, Mulholland F, Kopriva S, Malin G (2012) The response of diatom central carbon metabolism to nitrogen starvation is different from that of green algae and higher plants. Plant Physiol 158:299-312.

Hoppe CJM, Flintrop CM, Rost B (2018) The Arctic picoeukaryote *Micromonas pusilla* benefits synergistically from warming and ocean acidification. Biogeosciences 15:4553-4365.

Johnson WM, Kido Soule MC, Kujawinski EB (2017) Extraction efficiency and quantification of dissolved metabolites in targeted marine metabolomics. Limnol Oceanogr Meth 15: 417-428

Johnson LK, Alexander H, Brown CT (2018) Re-assembly, quality evaluation, and annotation of 678 microbial eukaryotic reference transcriptomes. Giga Sci 8:giy158.

Jones DT, Taylor WR, Thornton JM (1992) The rapid generation of mutation data matrices from protein sequences. Computer Appl Biosci 8:275-282.

Karl DM (2014) Microbially mediated transformations of phosphorus in the sea: new views of an old cycle. Ann Rev Mar Sci 6:279-337.

Kido Soule MCK, Longnecker K, Johnson WM, Kujawinski EB (2015) Environmental Metabolomics: Analytical Strategies. Mar Chem 177:1–62.

Kishino H, Miyata T, Hasegawa M (1990) Maximum likelihood inference of protein phylogeny and the origin of chloroplasts. J Molec Evol 31:151–160.

Klok AJ, Martens DE, Wijffels RH, Lamers PP (2013) Simultaneous growth and neutral lipid accumulation in microalgae. Biores Technol 134:233–243.

Kujawinski EB, Longnecker K, Alexander H, Dyhrman ST, Fiore CL, Haley ST, Johnson WM (2017) Phosphorus availability regulates intracellular nucleotides in marine eukaryotic phytoplankton. Limnol Oceanogr Lett 2:119-129. doi:10.1002/lol2.10043.

Kumar S, Stecher G, and Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. Molec Biol Evol 33:1870-1874.

Lewis LA, McCourt RM (2004) Green algae and the origin of land plants. Amer J Bot 91:1535–1556.

Lai J, Yu Z, Song X, Cao X, Han X (2015) Responses of the growth and biochemical composition of *Prorocentrum donghaiense* to different nitrogen and phosphorus concentrations. J Exp Mar Biol Ecol 405:6-17.

Li WKW (1994) Primary production of prochlorophytes, cyanobacteria, and eukaryotic ultraphytoplankton: Measurements from flow cytometric sorting. Limnol Oceanogr 39:169-175.

Li M, Shi X, Guo C, Lin S (2016) Phosphorus deficiency inhibits cell division but not growth in the dinoflagellate *Amphidinium carterae*. Front Microbiol 7:826.

Lin S, Litaker RW, Sundra WG (2016) Phosphorus physiological ecology and molecular mechanisms in marine phytoplankton. J Phycol 52:10-36.

Loebl M, Cockshutt AM, Campbell D, Finkel ZV (2010) Physological basis for high resistance to photoinhibition under nitrogen depletion in *Emiliania huxleyi*. Limnol Oceanogr 55:2150-2160.

Longnecker K (2015) Dissolved organic matter in newly formed sea ice and surface seawater. Geochim Cosmochim Acta 171:39–49.

Maat DS, Crawfurd KJ, Timmermans KR, Brussaard CPD (2014) Elevated CO_2 and phosphate limitation favor *Micromonas pusilla* through stimulated growth and reduced viral impact. *Appl Environ Microbiol.* 80:3119–3127.

Maat DS, van Bleijswijk JDL, Witte HJ, Brussaard PD (2016) Virus production in phosphorus-limited *Micromonas pusilla* stimulated by a supply of naturally low concentrations of different phosphorus sources, far into the lytic cycle. FEMS Microbiol Ecol 92:fiw136.

Marchler-Bauer A, Bo Y, Han L, Lanczycki CJ, Lu S, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, Gwadz M, et al (2017) CDD/SPARCLE: functional classification of proteins via subfamily domain architectures. Nucl Acids Res 4:D200-D203.

Markou G, Nerantzis E (2013) Microalgae for high-value compounds and biofuels production: a review with focus on cultivation under stress conditions. Biotechn Advan 31:1532-1542.

Martin P, Van Mooy BA, Heithoff A, Dyhrman ST (2011) Phosphorus supply drives

rapid turnover of membrane phospholipids in the diatom *Thalassiosira pseudonana*. ISME J 25:1057–1060.

Martin P, Dyhrman ST, Lomas MW, Poulton NJ, Van Mooy BAS (2014) Accumulation and enhanced cycling of polyphosphate by Sargasso Sea phytoplankton in response to low phosphorus. Proc Nat Acad Sci 111:8089-8094.

Mock T, Samanta MP, Iverson V, Berthiaume C, Robison M, Holtermann K, et al. (2008) Whole-genome expression profiling of the marine diatom *Thalassiosira pseudonana* identifies genes involved in silicon bioprocesses. Proc Nat Acad Sci 105:1579-1584.

Moseley JL, Chang C-W, Grossman AR (2006) Genome-based approaches to understanding phosphorus deprivation responses and PSR1 control in *Chlamydomonas reinhardtii*. Euk Cell 5:26–44.

Müller R, Morant M, Jarmer H, Nilsson L, Nielsen TH (2006) Genome-wide analysis of the *Arabidopsis* leaf transcriptome reveals interaction of phosphate and sugar metabolism. Plant Physiol 143:156–171.

Norici A, Bazonni AM, Pugnetti A, Raven JA, Giordano M (2011) Impact of irradiance on the C allocation in the coastal marine diatom *Skeletonema marinoi* Sarno and Zingone. Plant Cell Environ 34:1666–1677.

Rokitta SD, von Dassow P, Rost B, John U (2014) *Emiliana huxleyi* endures N-limitation with an efficient metabolic budgeting and effective ATP synthesis. BMC Genomics 15:1051.

Rokitta SD, von Dassow P, Rost B, John U (2016) P- and N-depletion trigger similar cellular responses to promote senescence in eukaryotic phytoplankton. Front Mar Sci 3:109.

Rubio V, Linhares F, Solano R, Martín AC, Iglesias J, Leyva A, Paz-Ares J (2001) A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. Genes Devel 15: 2122–2133.

Rusch DB, Halpern AL, Sutton G, Heidelberg KB, Williamson S, Yooseph S, Wu D, Eisen JA, Hoffman JM, Remington K, et al (2007) The Sorcerer II global ocean sampling expedition: northwest Atlantic through eastern tropical Pacific. PLoS Biol 5:e77.

Santos-Beneit F (2015) The Pho regulon: a huge regulatory network in bacteria. Front Microbiol 6:402.

Sharma KK, Schuhmann H, Schenk PM (2012) High lipid induction in microalgae for biodiesel production. Energies 5:1532–1553.

Shemi A, Schatz D, Fredricks HF, Van Mooy BAS, Porat Z, Vardi A (2016) Phosphorus starvation induces membrane remodeling and recycling in *Emiliania huxleyi*. New Phytol 211:886-898.

Shi X, Lin X, Li L, Palenik B, Lin S (2017) Transcriptomic and microRNAomic profiling reveals multi-faceted mechanisms to cope with phosphate stress in a dinoflagellate. ISME J 11:2209-2218.

Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular Evolutionary Genetics Analysis Using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. Molec Biol Evol 28: 2731–2739.

Tyrrell T. 1999. The relative influences of nitrogen and phosphorus on oceanic primary production. Nature 400:525–531.

Van Mooy BAS, Fredricks HF, Pedler BE, Dyhrman ST, Karl DM, Koblížek M, Lomas MW, Mincer TJ, Moore LR, Moutin T, et al (2009) Phytoplankton in the ocean use non-phosphorus lipids in response to phosphorus scarcity. Nature 457:69–72.

Whitney LP, Lomas MW (2016) Growth on ATP elicits a P-stress response in the picoeukaryote *Micrmonas pusilla*. PLoS ONE 11:e0155158.

Worden AZ (2006) Picoeukaryote diversity in coastal waters of the Pacific Ocean. Aquat Microb Ecol 43:165–175.

Wykoff DD, Grossman AR, Weeks DP, Usuda H, Shimogawara K (1999) Psr1, a nuclear localized protein that regulates phosphorus metabolism in *Chlamydomonas*. Proc Nat Acad Sci 96:15336–15341.

Yoshida K, Terashima I, Noguchi K (2007) Up-regulation of mitochondrial alternative oxidase concomitant with chloroplast over-reduction by excess light. Plant Cell Physiol 48:606-614.

8   Table 1. Enzymes that contain a significant motif in the gene sequences of *M. pusilla* CCMP1545 (MBARI_models (ver 1)). Column 2: the
9   Enzyme Commission (E.C.) number or pfam identifier; Column 3: main pathway(s) in which the enzyme is involved; Columns 4-6: fold
10   change and *q*-value of the transcript in P-deficient treatment relative to P-replete conditions are data from Bachy et al. (2018) and
11   corresponding gene ID.

Enzyme	E.C. Number	Pathway	Log 2 fold change*	*q*-value*	JGI gene ID*
Glyceraldehyde-3-phosphate dehydrogenase	1.2.1.12	Calvin cycle/glycolysis	2.73	0.000037	4267
Glyceraldehyde-3-phosphate dehydrogenase	1.2.1.12	Calvin cycle/glycolysis	-0.644	0.000037	700
Phosphoglycerate kinase	2.7.2.3	Glycolysis	-0.699	0.000037	7661
Pyruvate kinase	2.7.1.40	Glycolysis	-1.32	0.000037	9370
Pyruvate dehydrogenase	1.2.4.1	TCA[a] cycle/Glycolysis	0.756	0.000037	3535
Citrate synthase	2.3.3.1	TCA cycle	1.68	0.000037	1278
Citrate synthase	2.3.3.1	TCA cycle	1.10	0.000037	3714
Aconitase	4.2.1.3	TCA cycle	1.00	0.000037	2901
Succinate dehydrogenase	1.3.5.1	TCA cycle and ETC[b]	1.16	0.001515	2182
Fumarase	4.2.1.2	TCA cycle	1.68	0.000037	600
Fumarase	4.2.1.2	TCA cycle	1.73	0.000068	6662
Copper amine oxidase	1.4.3.21-22	Arginine and proline metabolism	2.87	0.000037	5863
Aspartate transcarbamylase	2.1.3.2	Pyrimidine biosynthesis	-0.609	0.000037	3615
Nucleoside phosphatase	3.6.1.3	Pyrimidine metabolism	0.711	0.000037	3322
Pantoate-beta-alanine ligase	6.3.2.1	Pantothenate and CoA biosynthesis	-0.616	0.000037	5304
Lysophospholipase	3.1.1.5	Glycerophospholipid metabolism	1.65	0.000037	3652
Acyl-CoA synthetase	6.2.1.3	Fatty acid metabolism	3.20	0.000037	1751
Long-chain acyl-CoA synthetases	6.2.1.3	Fatty acid metabolism	0.736	0.000037	416
Na+/PO$_4$ transporter	PF02690	inorganic nutrient transport	4.54	0.000037	5963

Table 2. Gene matches to *psr*1 in *Micromonas pusilla* CCMP1545 as originally discovered in the Joint Genome Institute (JGI) Integrated
Microbial Genomes (IMG) database and the corresponding gene in the updated gene model for CCMP1545 (MBARI_models (ver 1)). The *M.*
pusilla CCMP 1545 database in IMG contained 10,660 sequences and 4,795,637 letters. The top-scoring gene was considered to be a *psr*1-
like gene and the JGI gene ID is shown in parentheses. The corresponding *psr*1-like gene in the updated gene model is shown for reference,
and includes the JGI gene ID, the gene description, and the E-value from the BLAST search (see Methods) using the top scoring gene
(613233) from CCMP1545.

IMG Gene ID	Locus tag	Bit score	E-value
2615011133 (JGI 61323)	MicpuC2.est_orfs.1_306_4269596:1	90.5	2×10^{-19}
2615008475	MicpuC2.EuGene.0000130210	74.3	4×10^{-14}
2615010887	MicpuC2.est_orfs.10_1861_4270447:1	70.5	3×10^{-13}
2615007841	MicpuC2.EuGene.0000090108	67.8	4×10^{-12}
JGI ID 360	Myb domain-containing protein		8×10^{-156}

Fig. 1. Average ratio of the P-deficient to P-replete intracellular metabolite
concentrations during exponential growth (T_1) in *Micromonas pusilla* CCMP1545.
Concentrations are normalized to cell number in each treatment and means with one
standard deviation are shown (N=3 unless otherwise noted). Metabolites are listed in
order of descending concentration within three groups; those that were detected in
enough replicates for standard deviation to be calculated, those where only one ratio
could be calculated (black line), and those that were not detected (ND) in any of the three
replicates for one treatment (R = P-replete, D = P-deficient). Metabolite names marked
with an asterisk indicates a significant difference in concentration between treatments (*t*-
test, p < 0.05). Only one replicate in the P-deficient treatment contained a non-zero
concentration for *N*-acetylmuramic acid, thus while there was a significant difference in
concentration for this metabolite, only one ratio could be calculated. (DMSP =
dimethylsulfoniopropionate, NAD = nicotinamide adenine dinucleotide)

Fig. 2. The intracellular ratios of two purine nucleosides to their nucleobases for
Micromonas pusilla CCMP1545 in P-deficient (orange) and P-replete (blue) treatments
during exponential growth (T_1). Xanthosine and xanthine, and guanosine and guanine
were quantified in the targeted metabolomics method and normalized to cell abundance.
The average ratio and high and low values are shown based on three replicate cultures.
The asterisk marks a significant difference between the treatments (*t*-test, p = 0.03). The
dashed line represents a ratio of 1. Two samples in the P-replete treatment had guanine
levels below the limit of detection and the one ratio is shown as a line.

A) *Micromonas pusilla* CCMP1545: 2615011133

MTRMSGIEDDSFSYSFLLISGQLRDPTRLTPRPQTSSAPLGAGGGAAAASGMNTSGTFWPSDLDLDLESSDFL
DSLLLGGEMMTTQQQHHHAGDERLGHGPVPLGARAAGEDHDPRVVLAERDAGDGLLPRALADERREPVAR
AVVASNNNKQRLRWTPELIIKMFVDAVKRLGGLDLATPKGIMQLMDVEGMSIQIIVKSIILQKYRLQDSGGG
ASEFRVSPDASASGKRPRSEEDDAGGNGKDGNNSAGKTRRRPSAAERSAARLRAAEEKAREERDAARSMAA
AAAAAAAAAVERQNVELALLTGDAAGARDALEMSSHHAASHVGGAYDDVLDLVAGPGESGDGDGAWGD
VLHDHAIVGAAGDDVGLVDDVGSDPEAAAAMLKQLELQKKLHEHLMSQRRLQQQVEAHGVYLETILDQQK
RRRGVE

B)

Fig. 3. Predicted amino acid sequences of the *psr*1-like gene in *Micromonas pusilla*
CCMP1545 and a subset of other organisms. The amino acid sequence (with IMG gene
ID) is shown as described in the Joint Genome Institute Integrated Microbial Genomes *M.*
pusilla CCMP1545 database. The myb-like DNA-binding and myb coiled-coiled domains
are highlighted in blue (A). Predicted amino acid sequence alignment of *psr*1 and *psr*1-
like genes in the region of the myb-like DNA-binding domain (SHAQKYF class) and
LHEQL coiled-coil domain (B). Asterisks at the top of each column indicate 100%
conserved residues across species surveyed. Numbers at the end of each row indicate
position of the last shown residue in the amino acid sequence. Accession (MMETSP,
NCBI, *Tara*) or gene identification (IMG) numbers are listed for each sequence.

Fig. 4. Occurrence and phylogenetic relationship of *psr*1-like genes in marine
phytoplankton. Maximum likelihood method based on the JTT matrix-based model
(Jones *et al*., 1992). The tree with the highest log likelihood (-2442.94) is shown and is
based on the derived amino acid sequences for *psr*1 and *psr*1-like genes from eukaryotic
phytoplankton. The analysis involved 55 amino acid sequences. All positions containing
gaps and missing data were eliminated yielding a total of 43 positions in the final dataset.
Evolutionary analyses were conducted in MEGA7 (Kumar *et al*., 2016) Accession or
IMG gene identification numbers are shown with the taxon name and the star indicates
Micromonas pusilla CCMP1545 cultured in this study. The tree is unrooted and the scale
indicates the number of amino acid substitutions per site.

Fig. 5. Geographic distribution of *Micromonas psr*1-like genes and relationship with
phosphate concentration in the *Tara* Oceans eukaryotic gene atlas (Carradec, Pelletier, *et*
al., 2017). The geographic distribution and relative abundance of *Micromonas psr*1-like
genes in surface samples are depicted. Black dots represent stations where *Micromonas*
was detected; the size of purple circles represents the relative abundance of *psr*1-like
gene, where the FPKM of the psr1-like gene was normalized to the total FPKM of all
Micromonas-associated transcripts (A). Relative abundance of *Micromonas* is plotted
against phosphate concentrations for surface stations where *Micromonas* was detected.
Grey circles indicate samples where *psr*1-like genes were not detected, while samples
where psr1-like genes were detected are colored based on the relative expression of
*Micromonas psr*1-like gene, again normalized to total *Micromonas* transcript abundance.
No significant relationship was observed between the phosphate concentration and
abundance of *Micromonas psr*1-like transcripts (*t*-test, $p > 0.05$) (B). A split violin plot
depicting the distribution of expression of *Micromonas* fumarase synthase genes as a
function of presence or absence of *psr*1-like gene. The larger dashed line represents the
distribution mean and the smaller dashed lines represent the 25% and 75% quartiles.
Significant difference in the distribution of expression between the presence and absence
is assessed with Kolmogorov-Smirnov test ($p < 0.005$) (C).

Fig. 6. Conceptual model of Psr1-like TF regulation in *Micromonas pusilla* CCMP154 in
three major metabolic pathways under phosphorus deficiency. The figure illustrates the
relevant metabolites quantified in the present study and indicates if they are elevated
(orange, ratio > 1 Figure 2), depressed (blue, ratio < 1 Figure 2) in concentration in the P-
deficient cells relative to the P-replete cells. Some metabolite concentrations were equal
in each treatment (black) or were not quantified (gray). Genes that contain the conserved
motif with similarity to the myb-like regulatory element in *Arabidopsis thaliana* are
marked with a red star. The orange and blue letters refer to elevated or depressed
transcript levels for each gene ([1]data from Bachy *et al*., 2018). Descriptions of putative
metabolic responses in each pathway are shown. Genes encode for: A) pyruvate kinase,
B) pyruvate dehydrogenase, C) citrate synthase, D) aconitase, E) α-ketoglutarate
dehydrogenase, F) succinate dehydrogenase, G) fumarase, H) ribulose-1,5-bisphosphate
carboxylase, I) triose sugar transporter, J) nucleotide phosphatase DAG = diacylglycerol,
TAG = triacylglycerol