Title: Effects of male age and female presence on male associations in a large, polygynous

mammal in southern India

Authors: P. Keerthipriya, S. Nandini[†], T.N.C. Vidya[*]

Author affiliation (for all authors): Evolutionary and Organismal Biology Unit,

Jawaharlal Nehru Centre for Advanced Scientific Research (JNCASR), Bengaluru 560064,

India.

[†] Present address: Azim Premji Foundation, Doddakannelli, Sarjapur Road, Bengaluru.

Email: P. Keerthipriya: keerthipriya@jncasr.ac.in; S. Nandini:

nandini.shetty@azimpremjifoundation.org; T.N.C. Vidya: tncvidya@jncasr.ac.in; ORCID

iD for T.N.C. Vidya: 0000-0002-7143-9008.

[*] Corresponding author.

Address for correspondence:

T.N.C. Vidya

Jawaharlal Nehru Centre for Advanced Scientific Research

Jakkur

Bengaluru – 560 064, India.

Email: tncvidya@jncasr.ac.in

Phone: +91-80-22082968

Fax: +91-80-22082766

Abstract (<=250 words)

We present a detailed study of male associations in a roving species, the Asian elephant,

using six years of data on identified, nonmusth males. Adult males spent greater proportions

of their time solitarily than in mixed-sex or in all-male groups. Old (over 30 years) males

were sighted more frequently with their age-peers and less frequently with young (15-30

years) males than expected at random in all-male groups. Young males were not sighted

more frequently with old males than with young males, and did not disproportionately

initiate associations with old males. These results suggest that male associations, in the

absence of females, primarily allow for old nonmusth males to test strengths against age-

peers. Social learning from older individuals did not seem to be important in male

associations, unlike that observed in African savannah elephants. We also found a constraint

on the sizes of all-male groups, similar to that seen in female groups in our study

population, and with male group sizes being smaller than that of African savannah

elephants. However, most males had a significant top associate in female absence. In mixed-

sex groups, male associations occurred at random, suggesting that males were tracking

female groups independently. Thus, we find some differences in male social organisation

compared to the phylogenetically related African savannah elephant that occupies a similar

niche, and suggest that ecological factors might have shaped the differences in these male

societies.

Keywords

Asian elephant, male associations, association networks, Kabini Elephant Project, testing-

strengths hypothesis, social learning hypothesis, female presence/absence, male age.

Introduction

Adult males and females of many large mammals exhibit sexual dimorphism and strikingly different lifestyles, with female philopatry and male dispersal (see Greenwood 1980, Ruckstuhl and Neuhaus 2000). Following dispersal from their natal groups, males may spend some time alone or in all-male groups, and then form a) relatively stable bonds with female groups (for example, Hrdy 1977 in Hanuman langurs, Packer and Pusey 1987 in lions, Alberts and Altmann 1995 in baboons), b) form mixed-species groups during the breeding season and all-male groups outside of the breeding season (for example, Clutton-Brock *et al.* 1987 in red deer, Villaret and Bon 1995 in Alpine ibex, Mooring *et al.* 2003 in desert bighorn sheep), or c) rove between different female groups and form only temporary associations with female groups throughout the year (for example, Best 1979 in sperm whales, Poole 1982 in African savannah elephants, Desai and Johnsingh 1995 in Asian elephants, Baird and Whitehead 2000 in killer whales). The interactions between males themselves in polygynous species with these different kinds of lifestyles are expected to be competitive rather than affiliative, with males competing with one another for access to receptive females (van Hooff and van Schaik 1994). Therefore, strong associations are not expected between males in species with female-philopatry and may occur primarily in the context of coalitions to defend or to contest access to females (for example, Saayman 1971 in baboons, Connor *et al.* 1992, Gerber *et al.* 2020 in bottlenose dolphins, van Hooff and van Schaik 1994 in non-human primates, Wagner *et al.* 2008 in hyaenas). However, male-male interactions in all-male groups may be less competitive or aggressive than those within mixed-sex groups (Pusey and Packer 1987 in primates, Robbins 1996 in mountain gorillas). All-male groups may provide an opportunity to test strength against and assess competitors in a more relaxed setting (Bon *et al.* 1993 in mouflon sheep, Chiyo *et al.* 2011 in African savannah elephants). Male associations in all-male groups may also be motivated by the

opportunities available for social learning from older, more experienced males (Evans and

Harris 2008, Chiyo *et al*. 2011, 2012 in African savannah elephants, Bercovitch and Berry

2014 in giraffes). Increased efficiency in obtaining food resources (river otters, Blundell *et*

al. 2002) and improved defense against predators (sperm whales, Curé *et al*. 2013) are also

possible benefits from associating with other males, despite possible intrasexual

competition. There has been little study on male association patterns in mammals overall,

especially on those species that rove between female groups and do not form stable

multimale-multifemale groups or large multimale groups in the non-breeding season. We,

therefore, wanted to examine the extent and nature of male associations in such a roving

species that is large, polygynous, and faces potentially high male-male competition and

potential group size restriction, but is also phylogenetically related to a species with

complex male association patterns.

Asian elephants (*Elephas maximus*) are polygynous, with males and females exhibiting

different morphologies and adult lifestyles. Female society in this species is organised into

clans that show fission-fusion dynamics (de Silva *et al*. 2011, Nandini *et al*. 2017, 2018),

while pubertal males disperse from their natal groups and only temporarily associate with

other males and with female groups thereafter (McKay 1973, Desai and Johnsingh 1995).

There are often no clearly defined bull areas. Males are not known to form coalitions to

defend females. Males can breed throughout the year, but females are sexually receptive

only for a few days (Eisenberg *et al*. 1971) every four to five years, making receptive

females a rare resource, for which males are expected to compete intensely. Male-male

dominance interactions have been observed, indicating contest competition (McKay 1973,

Daniel *et al*. 1987, Chelliah and Sukumar 2013, Keerthipriya 2018).

Male Asian and African elephants are thought to be more reproductively active during

musth (a rut-like state); however, nonmusth males also acquire some mating and

reproductive success (Hollister-Smith *et al*. 2007, Rasmussen *et al*. 2007, Chelliah and

Sukumar 2015, Kabini Elephant Project, unpublished data). Therefore, competition for

females may also exist among nonmusth males, and male-male affiliative associations are

expected to be weak. However, males in the African savannah elephant have been shown to

have complex relationships, with males preferring to associate with age-peers (Chiyo *et al*.

2011, Goldenberg *et al*. 2014 – in the case of sexually inactive males) and related males

(Chiyo *et al*. 2011), and there is some evidence for older males being preferred associates or

being more central to male society than young males (Evans and Harris 2008, Chiyo *et al*.

2011, Murphy *et al*. 2020). Male associations were also shown to facilitate social learning:

bulls who had an older crop raider as a top associate were more likely to raid themselves

(Chiyo *et al*. 2012). Thus, temporary all-male groups seem to provide an opportunity to spar

and test strengths, and also possibly for younger males to learn from knowledgeable, older

males in African savannah elephants.

Although Asian and African elephants were previously assumed to have similar societies,

we now know that there are some differences between the female Asian elephant and

African savannah elephant societies (de Silva and Wittemyer 2012, Nandini *et al*. 2018),

probably because of a constraint on female group sizes due to ecology in the Asian elephant

(Nandini *et al*. 2017, 2018). Since males, being larger and continuing to grow in size as they

age (Sukumar *et al*. 1988), are likely to require more food than females, such a constraint

might also exist in male Asian elephants and lead to differences in male societies across

species, despite their phylogenetic closeness. Moreover, male African savannah elephants

were known to return to the same bull areas (areas frequented by males and not by many

female groups) when sexually inactive (Poole 1982), providing an opportunity for repeated

associations with specific individuals (although the selection of a bull area may also

possibly be a decision to associate with other males in the area; see Lee *et al*. 2011). An

absence of separate bull areas in Asian elephant populations might reduce the frequency of

males meeting each other, and hinder the formation of strong male associations. We,

therefore, wanted to examine associations among adult male Asian elephants to find out

whether possible ecological differences correlated with a different male social structure than

in the African savannah elephant, despite the phylogenetic similarity between species.

We analysed associations of nonmusth adult males in this paper because musth males spend

only a small proportion of their time in all-male groups (Keerthipriya *et al*. 2020). We

aimed to examine the prevalence, strength, and stability of associations among nonmusth

Asian elephant males and the factors (male age, presence of female groups, restriction on

group size) that might affect male associations. We hypothesized that male (adult,

nonmusth, male Asian elephants in this paper, unless specified otherwise) associations

might be based on opportunities available for a) social learning from older individuals

and/or b) testing strengths. Increased efficiency in obtaining food resources was not likely to

be a factor leading to adult male groupings in elephants because individuals require large

amounts of food and grouping is expected to create food competition. Defense against

predators was also not likely to be important because adult male elephants do not have any

natural predators and our work was carried out in Protected Areas. We did not examine

genetic relatedness as a cause for associations.

We set out to address the following specific questions:

1) *What are the proportions of time that nonmusth males spend in all-male groups and*

mixed-sex groups and how are they related to male age? While examining male

associations regardless of the males' musth status, we had found that age did not affect

the time spent in mixed-sex or all-male groups (Keerthipriya *et al*. 2018). We had also

found that old males increased their association with females when they entered musth,

while young males decreased their association with females when they were in musth,

compared to their nonmusth associations (Keerthipriya *et al*. 2020). Therefore, we

expected that nonmusth older males would spend less time with female groups than

nonmusth younger males. Although older males were seen more often than younger

males in all-male groups of the African savannah elephant (see Chiyo *et al*. 2011,

Goldenberg *et al*. 2014), that pattern might be reversed among nonmusth males in

Kabini if there was a constraint on group size.

2) *How does male age and the presence or absence of females in the vicinity affect patterns*

of associations between nonmusth males? We expected male age to affect associations

differently depending on whether males associated primarily for social learning or for

testing strength. We expected males to use their time in all-male groups, rather than in

mixed-sex groups, to test strength. However, social learning from older males could

occur in female absence (learning related to resources) and/or female presence (learning

related to reproduction). If male associations were primarily based on social learning

from older individuals, younger males would seek out older males more often than

expected by chance alone. If male associations were primarily driven by opportunities

for social learning, but older males were restricted in the amount of time they spent with

other males (possibly due to group size constraints), older males might not spend more

time in all-male groups, but the proportion of sightings of young males in which young

males associated with older males would be higher than the proportion of sightings of

young males in which young males associated with other young males. Older males

would also be better connected in male networks if they were preferred as associates by

young and old males, although better connectedness of old males could also be observed

if old males had stronger age-peer associations than younger males. If the primary

purpose of male associations was to test strengths, males would be expected to associate

(relative to population age-structure) to a greater extent with age-peers than with much

younger or older individuals, whose relative strengths are easily assessed by size

differences. We did not have *a priori* expectations about whether old or young males

would be more likely to associate amongst themselves if there were preferential

associations with age-peers. It was possible that young males might show greater

preferential association with age-peers than old males since old males might better know

their strengths (through experience) than young males. However, it was also possible

that old males might show greater preferential association with age-peers because, being

more competitive breeders than young males, resolving dominance relationships might

be more crucial for them.

Since competition for females could play a major role in how males associated, we

examined male associations in the immediate presence and absence of females. Unlike

the case of the African savannah elephants, no clear indicators of active and inactive

sexual states outside of musth have been recognised in Asian elephants. We expected

the average number of other males that males met to be lower, and associations to be

weaker, in the presence of females than in their absence. It should be noted that if males

spent a greater amount of time in the absence of females than in their presence, males

would likely also meet other males more frequently in female absence, resulting in the

male association network in female absence being denser and better connected than that

in female presence. However, this difference should not be significant after controlling

for the difference in the number of sightings, if the greater number of associations was

merely an effect of the greater amount of time spent. Due to the expected competition

among males, we also predicted smaller experienced group sizes of the males in female

presence than absence. It was also possible that males approached female groups by

tracking them independent of other males, in which case, we expected males to associate

with each other at random. In such a case, we had no *a priori* expectation about whether

the group sizes of males would be higher in female presence or in female absence.

3) Is there a restriction on male group size? If there was a constraint on the group sizes of

all-male groups, we expected a trade-off between the number of associates of a male and

how often the focal male was sighted with those associates. We also expected older

males to spend less time in all-male groups or to form smaller all-male groups than

younger males because of greater food competition (due to larger body size). However,

a similar pattern could also be observed if young males had stronger age-peer

associations than old males. On the other hand, if there were stronger age-peer

associations among old than young males, we would expect old males to spend more

time in all-male groups or to form larger all-male groups than young males, although

group-size restriction might reduce the difference.

4) *Are there preferential associations between nonmusth males and, if so, are they stable*

over time? We did not have any *a priori* expectation about whether preferred, stable

associations should be present or not, but, if they occurred, we expected them to be less

frequent than that in the African savannah elephant due to possible group size

constraints.

Methods

Field data collection

The field study was carried out in Nagarahole and Bandipur National Parks and Tiger

Reserves (Nagarahole: 11.85304°-12.26089° N, 76.00075°-76.27996° E, 644 km^2;

Bandipur: 11.59234°-11.94884° N, 76.20850°-76.86904° E, 872 km^2) in southern India

from March 2009 to July 2014. Nagarahole and Bandipur National Parks are separated by

the Kabini reservoir, and we refer to the elephants in the two parks as the Kabini population.

Because of the high density of elephants around the reservoir and better visibility for

behavioural observations, our sampling was centred around the reservoir, and extended to

the forests in either direction with lower frequency of sampling (see Nandini *et al*. 2017).

We sampled pre-selected forest routes (see Nandini *et al*. 2017 for details) in the study area

from early morning to late evening (~6:30 AM am to 6:00-6:45 PM depending on field

permits and light conditions). There were no distinct bull areas in our study area. The adult

sex ratio in the study area was 1 male : 4-5 females (Gupta *et al*. 2016).

We tried to sex, age, and identify all the elephants sighted. Asian elephants are sexually

dimorphic, with males being taller and bulkier than females apart from differences in

genitalia. Females do not possess tusks, although some males (called makhnas) are also

tuskless. We estimated age based on shoulder height, body length, skull size, and skin folds

(see Vidya *et al*. 2014), with semi-captive elephants in the same area serving as a reference

for ageing older animals. We placed males into the following age categories, based on their

estimated ages: calves (<1 year), juveniles (1-<5 years), sub-adults (5-<15 years), young

adults (15-<30 years), and old adults (>=30 years). Young adult males were likely to have

completely dispersed from their natal herds but were possibly less reproductively

competitive than old adult males, based on studies of African savannah elephants (Poole

1982, Poole *et al*. 2011). The ages used for classifying males into these categories were

those calculated at the mid-point of the study period (November 2012). We identified

individuals based on a combination of ear, back, tail, tusk, and body characteristics (detailed

in Vidya *et al*. 2014). We recorded group size, GPS location, time of sighting, and whether

adult males were in the presence or absence of females. Females were classified as adults

when they were 10 years old (see Nandini *et al*. 2018). Adult males were said to associate

with a female group (one or more adult females and their young that were in close proximity

and showed coordinated movement; see Nandini *et al*. 2018) if they fed within 10 m (easy

physical reach) of a group member or interacted with any group member. When two males

associated with the same female group at the same time, they were said to be associating

with each other in female presence. Rarely (only three different sightings), males were seen

to associate with subadult females (5-10 years old) in the absence of an adult female and

this was also considered to be association in female presence. Males were said to associate

with each other in female absence if they fed within about 50 m of each other and there

were no females in the vicinity. At this distance, the males would be able to display or react

to visual signals, apart from sensing one another through sound or smell. Males could

indulge in sparring during their associations, but if males, upon encountering each other,

displayed only aggressive interactions and moved away, they were not considered to be

associating.

Data Analysis

Data analysis was carried out using only those sightings in which all adult males were aged

and identified and female group compositions (if applicable) were known. Of the 878 days

of field work between 2009 and 2014, elephants were sighted on 853 days, identified adult

males were sighted on 718 days, and nonmusth adult males on 681 days. In many of the

analyses mentioned below, only males who were sighted at least 10 times in that particular

category (such as group composition type or female presence) while not in musth were used,

as associations of males seen rarely are unlikely to represent their actual association patterns

and may bias the results. Similarly, if there was a comparison between different categories

(such as associations in female presence and absence), common males sighted at least 10

times (when not in musth) in each of the categories were used, unless otherwise mentioned.

ANOVAs, GLMs/GLMEs (General Linear Models/Generalised Linear Mixed-Effects

Models), and non-parametric tests were performed using Statistica 7 (StatSoft, Inc. 2004)

and randomisations (as randomisations of observational data can account for potential

sampling bias in social network analyses; see Farine 2017) were carried out using

MATLAB (MATLAB R2011a, MathWorks, Inc, 1984-2011, www.mathworks.com) unless

specified otherwise.

Proportions of time spent in all-male and mixed-sex groups and their relationship with male

age

We calculated the number of minutes individual males (that were seen on at least 5 different

days when not in musth) were seen in the following group types and calculated the

proportions of each individual's time spent in such groups: 1) solitary, 2) all-male groups

with only one adult male (but including subadult or juvenile males and, therefore, not

solitary), 3) all-male groups with more than one adult male, and 4) mixed-sex groups. Since

the four proportions add up to one and are, therefore, not independent, and the number of

males seen in group type 2 was small, we compared only two of the four categories – all-

male groups with more than one adult male and mixed-sex groups. Males were categorized

into old and young adults as explained in the *Field data collection* section above. We

carried out Wilcoxon's matched-pairs tests, separately on old and young males, to test

whether the proportions of time spent in the different group types differed significantly.

Since the proportions were calculated for each male, each male was represented only once

in this analysis, effectively making male identity random. In order to find out whether old

males spent a smaller proportion of their time in mixed-sex groups than young males did,

we used a Mann-Whitney *U* test.

Changes in male associations and time to sighting independence

Using data from the last two years of observation (2013-2014), we examined continuous

observations of individual males and recorded the number of changes in that male's

associations in five-minute bins (all observations that lasted less than five minutes were

removed for this calculation). If there were changes in the identities of the adult males in the

focal male's group or if the focal male's association changed from female presence to

absence (or vice versa), they were counted as changes in the focal male's associations. The

time interval at which there was roughly equal probability of a male's association changing

or not would indicate the time interval at which sightings could be recorded as independent

of each other (in order to avoid pseudoreplication of data). We found this time interval to be

about 80 minutes (see Supplementary Material 1). Therefore, for all subsequent analyses of

sighting data, we only included independent sightings based on this time interval.

Effect of male age and the presence or absence of females on male association patterns

<u>Male age and social learning</u>

We examined the initiation of associations and the pattern of associations between males to

find out whether social learning might be a reason for associating. As mentioned in the

Introduction, if associations were primarily based on social learning, younger males would

be expected to seek out older males more often than expected by random chance. We,

therefore, examined all the instances in which two males (dyadic combinations in the small

number of cases where there were more than two males) of different ages were in close

proximity and one approached the other and associated with it. It was possible under such

circumstances for either of the males to approach the other. We randomly chose one of the

males as the focal male and used a generalized linear mixed-effects model (GLME) with a

binomial dependent variable (0 if the focal male approached the associate and 1 if the focal

male was approached by the associate), age difference (age of the focal male minus the age

of the associate) as a continuous predictor variable, and dyad identity as a random factor. If

younger males approached older males more often than expected, a significant positive

effect of the age difference on the dependent variable would be found. We used the *fitglme*

function in MATLAB R2011a, with a logit link function and Laplace estimation method,

and the model was fit using maximum likelihood. For this analysis, we used data only from

the years 2011-2014, during which detailed behavioural observations were available, and

did not include dyads that were already present when we began the observation. We only

used instances (of all males) when observers were present during the beginning of the

association and one of the males clearly moved towards the other.

We had expected that if social learning was the primary reason for the formation of all-male

groups, but older males were restricted in the amount of time they spent with other males,

the proportion of young males' time (sightings) that was spent with older males would still

be higher than the proportion of young males' sightings spent with other young males.

Therefore, we compared these proportions, in female presence and absence, using

Wilcoxon's matched-pairs tests. (Young males would spend a greater proportion of their

sightings with other young males than with old males if the testing-strength hypothesis was

correct.)

We had also expected older males to be better connected in male networks if they were

preferred as associates by both young and old males (or disproportionately by old males).

We examined this by analysing social networks of males. We first calculated the association

index (AI; Ginsberg and Young 1992) between pairs of identified males as the number of

times (independent sightings) two males were sighted together (N_{AB}) divided by the total

number of (independent) sightings of the two males. AI between each pair of males was

calculated separately in female presence and female absence (for instance,

$AI_{AB(F\_abs)}=N_{AB(F\_abs)}/(N_{A(F\_abs)}+N_{B(F\_abs)}-N_{AB(F\_abs)})$, where $N_{A(F\_abs)}$ is the total number of

sightings of A in female absence and $N_{B(F\_abs)}$ is the total number of sightings of B in female

absence). We used these AIs to calculate association strength (or weighted degree) for each

individual male, as the sum of the AI values between a male and all his associates. We

calculated the difference in association strength between the two age-classes (mean

association strength of old males – mean association strength of young males) in the

observed dataset. We then obtained the same value from each of 5000 permuted datasets

(see the section below for details of permutations) and compared the difference in the

observed dataset with those from the permuted datasets (see section below; $P<0.025$ for

statistical significance as we had no prior expectation about whether the observed values

would be lower or higher than the permuted values).

<u>Male age and testing strengths</u>

In order to find out whether males preferentially associated with age-peers (as predicted by

the testing-strengths hypothesis) more often than expected by chance, we used a

permutation procedure following Whitehead (2008, pg. 124). This procedure was carried

out on the datasets of males in female absence. We generated 5000 *males permuted* datasets,

by flipping (switching) adult males across independent sightings, while keeping the group

size and the number of sightings for each male constant, in each permuted dataset. The

number of flips performed in each permutation was five times the number of sightings in

that dataset. We calculated the number of times old ($>=30$ years) and young (15-30 years)

males were sighted with other males of the same or different age class in the observed

dataset and compared these observed values with the values from the permuted datasets. We

used the distribution of values from the permuted datasets to determine whether the

observed value was greater or lower than that expected by chance (P = proportion of

permuted values that were higher – if the observed value was higher than more than half of

the permuted values – or lower – if the observed value was lower than more than half of the

permuted values – than the observed value; $P<0.025$ for statistical significance).

We also compared the proportion of their sightings that old males spent associating with

other old males with the proportion of their sightings that they spent associating with young

males using Wilcoxon's matched-pairs tests. A higher proportion of sightings with other old

males than young males would be seen if the testing-strengths hypothesis was correct.

Male associations in the presence and absence of females

We performed a similar permutation procedure as described in the section above on the

female presence dataset, comparing the observed numbers of sightings in which males

associated with their age-peers and the other age-class. If association between males was

influenced by competition for females, we expected old males to associate less frequently

than that expected by random chance. We had also expected the male association network to

be denser and better connected in female absence than in female presence because of

potential competition between males in the presence of females. We, therefore, compared

the following network statistics between male association networks in female presence and

absence: mean *degree* (number of associates of a focal male), mean *clustering coefficient*

(the proportion of the total possible connections between a male's associates that exist),

mean *path length* (the smallest number of connections between two individuals in the

network), and *network density* (the proportion of all possible connections that exist in the

network) (Latapy 2008; see Wasserman and Faust 1994). We compared these network

statistics and mean AIs between female presence and absence using a sampled

randomization test (Sokal and Rohlf 1981, pp. 791-794). We created 5000 permuted

datasets (permuted by randomly assigning sightings to female presence or absence, while

conserving the sample sizes for both the categories and ensuring that at least ten sightings of

each male were assigned to each category in each permutation). We then compared the

observed differences in network statistics and AI between the original female presence and

female absence datasets with the differences in statistics between the permuted 'female

presence' and 'female absence' datasets. The probability of a significant difference between

the observed values was calculated as the proportion of permutations in which the difference

in statistic between the permuted datasets was greater than or equal to the difference in

statistic between the observed datasets. Since the total numbers of sightings in female

presence and female absence were conserved, a significant difference in this test would

indicate a difference in network statistic that could not be attributed to the difference in the

numbers of sightings.

We compared the degree distributions of association networks in female presence and

absence with their Poisson expectations (expected for a Erdös-Rényi random network;

Erdös and Rényi 1960) to test whether males were associating with each other at random.

We examined the effect of female presence on male group size by performing a GLM on

experienced group sizes (experienced by individual adult males; counted as the number of

adult males in each sighting of the focal male, including the focal male). We used female

presence/absence as a fixed factor and male identity as a random factor.

Restriction on all-male group size

If there was a constraint on group size in all-male groups, a negative relationship would be

seen between the number of associates of a male and how frequently the focal male was

sighted with those associates. We, therefore, calculated the average of the proportions of

sightings of a focal male (out of all the sightings of that male, including solitary sightings)

that the male spent with his different associates. This average proportion of the focal male's

sightings spent with his associates was correlated with the number of associates of that male

(in female absence) using a Spearman's rank-order correlation. We also examined whether

old males spent less time than young males in all-male groups by using the data on

proportions of time spent in different types of groups (section towards the beginning of the

Methods) and performing a Mann-Whitney U test. We also examined the relationship

between the mean experienced group size of a male and his age using Spearman's rank-

order correlation.

Preferred male associations and stability of associations

We tested for preferred associations or avoidance amongst identified males across sampling

periods smaller than the entire dataset using SOCPROG 2.6 (Whitehead 2009). We used a

sampling period of 14 days and 10,000 permutations with 10,000 flips for each permutation.

We used the '*permute associations within samples*' method, which tests for long-term

(across sampling period) preferences and avoidances (Whitehead 2009). The presence of

long-term preference/avoidance is indicated by significantly (>95% of the values of the

randomised datasets) higher SD and CV of AI values from the real dataset when compared

to the randomised datasets. We used sightings of all males for this analysis.

We additionally determined a top associate (based on AI value) for males, separately in

female presence and in female absence. Directed networks of males and their top associates

were constructed using Gephi 0.8.2 (Bastian *et al.* 2009). In order to examine whether the

identity and strength (AI) of a male's association with his top associate was different from

random, we again used the procedure for permuting associations (Whitehead 2008, pg. 124)

explained in a section above. We permuted associations between males separately in the

female presence and female absence datasets (5000 permutations on each dataset, with the

number of flips in each permutation being five times the number of sightings in that

dataset). We compared the AI of a male and his observed top associate with AI values for

the same dyad from the permuted dataset, and considered the association to be significant if

the observed AI was greater than 95% of the permuted values. In each permutation, for each

male, we also recorded the identity of his permuted top associate. We examined how often

the male had the same top associate in the permuted datasets, and considered the identity of

his top associate to be significantly different from random if the observed top associate was

not the top associate in greater than 95% of the permuted datasets.

In order to find out whether male associations were stable across years, we compared AI

matrices between consecutive years, using those males that were common to and sighted at

least five times in both years, by performing Mantel tests of matrix correlation (Mantel

1967), with 5000 permutations, in MATLAB (MATLAB R2011a, MathWorks, Inc, 1984-

2011, www.mathworks.com). Due to limitations of the number of sightings, only data from

2011-2014 and only in female absence were used for this analysis.

Results

Proportions of time spent in all-male and mixed-sex groups and their relationship with male

age

We sighted 83 nonmusth males in all (see Supplementary Material 2), of which 44 were

seen in the presence of females and 81 in the absence of females. Based on the set of

nonmusth males seen on at least five different days (*N*=38), we found that males spent an

average (± SD) of 12.3% (± 11.79%) of their time in all-male groups and 29.6% (± 22.73%)

of their time in mixed-sex groups (Figure 1a). Old males spent similar proportions of their

time in all-male and mixed-sex groups (Wilcoxon's matched-pairs test: $T=63.00$, $Z=0.259$,

$N=16$, $P=0.796$), whereas young males spent a significantly higher proportion of their time

in mixed-sex groups than in all-male groups (Wilcoxon's matched-pairs test: $T=9.00$,

$Z=3.815$, $N=22$, $P<0.001$, see Figure 1b). Old males spent a significantly smaller proportion

of their time in mixed-sex groups than young males did (Mann-Whitney U test: $U=79.50$,

$Z_{adj}=-2.853$, $P=0.004$, see Figure 1b).

Effect of male age and the presence or absence of females on male association patterns

Based on the 80-minute cutoff, there were 2466 independent sightings of adult nonmusth

males (374 sightings in female presence and 2092 sightings in female absence) in the study

period. Fourteen males (4 old and 10 young males) were sighted at least 10 times in female

presence, and 32 males (13 old males and 19 young males) were sighted at least 10 times in

female absence. Only 12 males were sighted at least 10 times each in the presence and

absence of females.

<u>Male age and social learning</u>

The age difference between the focal male and the other male did not significantly affect

who approached whom either in female presence (GLME: *Estimated coefficient*$_{Age\ difference}$= -

0.035, t_{stat}= -0.821, $N=19$ approaches, $df=17$, $P=0.423$) or in female absence (GLME:

Estimated coefficient$_{Age\ difference}$= -0.026, t_{stat}= -0.772, $N=48$ approaches, $df=46$, $P=0.444$)

(Figure 2). Thus, old and young males were equally likely to approach each other to initiate

association. We also found no significant difference in the proportion of its sightings that a

young male spent with old males (Female presence: mean $\pm$ SD 0.058 ± 0.035; Female

absence: 0.073 ± 0.086) and the proportion of its sightings a young male spent with other

young males in female presence or absence (Female presence: mean $\pm$ SD: 0.111 ± 0.090;

Wilcoxon's matched-pairs test: T=6.50, Z=1.610, N=10, P=0.107; Female absence: 0.065 ±

0.045; Wilcoxon's matched-pairs test: T=58.00, Z=0.114, N=19, P=0.910).

Examining the effect of male age-class on association strength, we found old males to have

only a slightly higher association strength (mean=0.087) than young males (mean=0.052) in

the absence of females (difference in mean association strength (old-young): observed

dataset: 0.035; permuted datasets: mean (2.5 and 97.5 percentiles): 0.003 (-0.015, 0.020),

P<0.001, Table 1). The observed difference in mean association strengths between the age-

classes was not different from the permuted values in female presence (see Table 1).

<u>Male age and testing strengths</u>

In female absence, old males were sighted together more frequently than expected by

chance (permutation test: observed: 42 sightings, *males permuted* mean (2.5, 97.5

percentiles): 25.5 (18, 34) sightings, P<0.001; Figure 3a). Old and young males were

sighted together less frequently than expected by chance (observed: 51 sightings, permuted:

69.4 (58, 80) sightings, P=0.001) and young males were sighted together as expected by

chance (observed: 38 sightings, permuted: 42.9 (33, 53) sightings, P=0.178; Figure 3a).

Although these tests were based on small numbers of males, the results remained unchanged

when we repeated the analysis using all the identified males (Supplementary Material 3).

The proportion of their sightings that old males spent with other old males (mean ± SD:

0.141 ± 0.116) was higher than the proportion that they spent with young males (mean ±

SD: 0.085 ± 0.069; Wilcoxon's matched-pairs test: T=5.00, Z=2.489, N=13, P=0.013), in

female absence.

Male associations in the presence and absence of females

We found that old males were sighted together as expected by chance in the presence of

females (permutation test: observed: 4 sightings, *males permuted* mean (2.5 and 97.5

percentiles): 1.6 (0, 4) sightings, $P=0.069$). The numbers of sightings were not different

from those expected by chance in the case of old and young males seen together (observed:

12 sightings, permuted: 11.1 (7, 16) sightings, $P=0.439$) and young males seen together

either (observed: 13 sightings, permuted: 14.7 (10, 19) sightings, $P=0.289$) (Figure 3b).

These results remained unchanged when we repeated the analyses using all identified males

without a sighting cutoff (Supplementary Material 3). All three age-class combinations of

males were sighted together more often in female absence than in female presence (Figure

3). However, while the observed male association network appeared more connected and

denser (higher mean degree, mean clustering coefficient, and density, and lower mean path

length) in female absence than in female presence, none of these statistics was significantly

different based on sampled randomisation tests (all $P>0.05$, 12 males seen at least 10 times

each in female presence (249 sightings) and in female absence (855 sightings), Table 2).

There was also no significant difference in mean AI between males in the presence and

absence of females (Table 2, see Supplementary Material 4 for distributions). Thus, the

seemingly more connected network in female absence was probably a result of greater time

spent by males in female absence.

The association network of adult males in female presence was not significantly different

from a random network ($\chi^2=6.122$, $df=8$, $P=0.633$), but the network in female absence was

significantly different from random ($\chi^2=184.647$, $df=15$, $P<0.001$) (see Supplementary

Material 5).

Group sizes (number of nonmusth adult males in the sighting) were small in general, with

the modal group size of all groups that had an adult male being 1 and the modal group size

of all multi-male groups being 2 (Figure 4a). The experienced group sizes of common males

(N=12, seen 273 times in all in the presence of females and 1304 times in all the absence of

females) were larger in female presence (mean ± SD: 1.6 ± 0.80) than in female absence

(mean ± SD: 1.2 ± 0.52; $F_{1,1553}$=48.520, P<0.001). There was a significant main effect of

male identity ($F_{11,1553}$=2.918, P=0.045) and a significant interaction effect between male

identity and female presence ($F_{11,1553}$=2.157, P=0.014; Figure 4b).

Restriction on all-male group size

The mean proportions of sightings that a male spent with an associate was significantly

negatively correlated with the number of associates of that male (Spearman's rank-order

correlation: N=31 males, each with at least one associate and seen at least ten times in

female absence, r_s= -0.582, P<0.001; Figure 5). The relationship was even more negative

when the mean proportion of all-male group sightings (rather than all sightings) spent with

an associate was correlated with the number of associates of those males (r_s= -0.984,

P<0.001; Figure 5). Thus, in female absence, males that associated with a greater number of

males spent a smaller proportion of their sightings on average with each of their associates.

However, contrary to expectation, old and young males did not spend significantly different

proportions of their time in all-male groups (Mann-Whitney U test: U=157.50, Z_{adj}=0.547,

N_{Old}=16, N_{Young}=22, P=0.584), and there was no significant relationship between male age

and mean experienced group size (Spearman's rank-order correlation: N=31 males, each

with at least one associate and seen at least ten times in female absence, r_s=0.213, P>0.05).

Preferred male associations and stability of associations

We found no evidence of preferred male associations across 14-day sampling periods in

female presence or in female absence (details in Supplementary Material 6). Only 14.3% (2

out of 14 males; 2 out of 15 dyads as one male had two top associates) of the males had a

significant association with their top associates in female presence, whereas 61.3% (19 of 31

males; one male did not associate with any of the other males) of the males had a significant

association with their top associate in female absence (mean significant AI with top

associate in female absence=0.032, see Supplementary Material 4 for AI distribution;

networks shown in Figure 6). We found that the identity of the focal male-top associate

dyad was significantly different from random in 13.3% (2 out of 15) of the dyads in female

presence and in 83.9% (26 out of 31) of the dyads in female absence.

Mantel tests showed no significant correlation between association matrices across years, in

two out of the three comparisons (Supplementary Material 7).

Discussion

This is the first detailed study of nonmusth adult male associations in Asian elephants in a

relatively undisturbed natural habitat. We found male-male affiliative associations to be

weak as expected, and in contrast to that found in the African savannah elephant.

Proportions of time spent in all-male and mixed-sex groups and their relationship with male

age

Nonmusth adult males spent only ~12% of their time in all-male groups in Kabini, in

contrast to ~63% after adjusting for age-effects (Chiyo *et al*. 2011) and 30.5% of all bull

sightings (Lee *et al*. 2011) in the Amboseli African savannah elephant population. Unlike

Kabini, Amboseli has distinct "bull areas" (Poole 1982) that 'sexually inactive' adult male

elephants inhabit but females seldom do, and it is possible that adult males may be more

likely to encounter and form strong associations with one another in such bull areas.

However, Chiyo *et al*. (2011) used observations of all-male groups of males in an area used

by both males and females; therefore, the difference cannot be attributed to bull areas in that

case. We found evidence for a restriction of all-male group sizes in Kabini (see below), and

this restriction may contribute to the smaller time spent in all-male groups. We expected and

found old nonmusth males to spend less time with female groups than young nonmusth

males. Whereas old males spent a similar proportion of their time in all-male and mixed-sex

groups, young males spent more of their time in mixed-sex than all-male groups. As young

males invest less in a competitive breeding strategy like musth (see Keerthipriya *et al*.

2020), their greater associations with females might facilitate learning about female groups

in areas they have dispersed to and possible opportunistic matings (see Chelliah and

Sukumar 2015, Kabini Elephant Project, unpublished data), or reflect a reduced cost of

feeding alongside females compared to that faced by old males. Further data are required to

examine these possibilities.

Effect of male age on association patterns: Reasons for adult male associations

As adult Asian elephants do not face any major risks of predation within protected areas and

are not known to form coalitions, we had hypothesised that all-male groups could provide

an opportunity for younger males to learn from older males or for males to test strengths

against age-peers (learning about their relative dominance status, instead of resources). Our

results on male associations favour the testing-strengths rather than the social learning

hypothesis. Similar-aged males have been shown to preferentially associate with each other

and test strengths in all-male groups of other species (Villaret and Bon 1995 - Alpine ibex,

Cransac *et al*. 1998 - mouflon sheep) and all-male groups in the Kabini elephant population

may similarly provide a relaxed setting for old nonmusth males to test strengths with age-

peers. However, young males in Kabini were sighted together as expected by chance both in

female presence and absence. Similar to our results, old (30+ years) males associated more

with age-peers, while the within age-class associations were as expected by chance in

younger age-classes in all-male groups in the Amboseli population (Chiyo *et al*. 2011). As

old males regularly enter musth, it may be more important for them than for young males to

clarify their relationships when they are not in musth. It would be interesting to find out if

dominance relationships formed when not in musth have a bearing on males entering into

musth. Old males have also spent more time in their post-dispersal home ranges and might

have formed significant associations with familiar age-peers. Male African savannah

elephants in Amboseli picked familiar age-peers as sparring partners (Chiyo *et al*. 2011),

and, in other species, interactions with familiar opponents tended to be less aggressive than

those with unfamiliar opponents (Smith *et al*. 1999, López and Martin 2001, Wich and

Sterck 2007). It would be interesting to examine the identities of sparring partners amongst

males and whether familiarity plays a role in the Kabini population. Murphy *et al*. (2020)

suggested that, perhaps, younger (20-30-year-old African savannah male elephants in their

study) males had not yet established a consistent nonmusth home range, leading to lowered

stability in association strength compared to older males. However, we found no difference

between the turnover rates of young and old nonmusth males (see Supplementary Material

2); therefore, this is unlikely to be the reason for the difference observed in our six-year

dataset.

We did not find any evidence for the social learning hypothesis, which suggests that the

superior knowledge of experienced, older males would attract younger males to associate

with them in order to learn about the location of food resources in all-male groups or about

interactions with females in mixed-sex groups. Contrary to the expectations based on the

social learning hypothesis, we found that young males were not sighted with old males to a

greater extent than with other young males, either in female presence or absence. During the

small amount of time that old and young males spent together, males were equally likely to

approach the other to associate regardless of their relative ages, both in female presence and

absence. Therefore, young males did not seek out old males. Although old males had higher

association strengths than young males in female absence, this was due to the stronger old

male-old male associations, rather than old male-young male associations.

In African savannah elephants, older males were found to be important in male networks.

Older males spent more time with other males than younger males did (Poole 1982, Chiyo

et al. 2011, Lee *et al*. 2011) and had a greater number of associates in all-male groups

(Chiyo et al. 2011) in the Amboseli population. In the Samburu population, sexually

inactive older males showed significant affiliation with a higher proportion of available

dyads compared to sexually inactive younger males (Goldenberg et al. 2014). Old males had

higher centrality in association networks based on all-male groups in Amboseli (Chiyo *et al*.

2011), although when males were classified based on their sexual state (sexually active and

inactive) in the Samburu population, there was no correlation between centrality and age in

sexually inactive networks and a negative correlation between centrality and age in sexually

active networks (Goldenberg *et al*. 2014). In a recent study in South Africa that examined

nonmusth male associations, older males showed higher stability in association strength

than younger males across sampling periods; however, age of the male did not affect the

stability of the identity of top ranked associates across sampling periods, eigenvector

centrality, or association strength of individual males in the network (Murphy *et al*. 2020).

Further, younger males preferentially associated with older males, unlike what we found in

Kabini. In Amboseli, males of the 20-29 year age-class associated with older (30+) males

more than expected by chance (Chiyo *et al*. 2011), and male associations seemed to

facilitate social learning; males who had an older crop raider as a top associate were more

likely to raid themselves (Chiyo *et al*. 2012). Older African savannah elephant males were

also preferred as neighbours while associating by males of all ages, based on observations in

a bull area in Okavango Delta, Botswana (Evans and Harris 2008), and have been

considered analogous to the knowledgeable matriarchs of female groups in the species

(McComb et al. 2001, Evans and Harris 2008). More generally, preferred association with,

and social learning from older, more experienced individuals has been observed in the

context of mating behaviour (Pereira 1988, Beecher et al. 1994), and foraging strategies

(Rajpurohit et al. 1995, Biro et al. 2003) in other species.

The greater social role of older males in the African savannah elephant (with younger males

preferentially associating with older males) compared to the Asian elephant in female

absence may stem from differences in the habitats they occupy. Asian elephants occupy

moister, more forested habitats, in which food is possibly more dispersed and unpredictable

in space at a very local scale (but possibly more predictable at a larger spatiotemporal scale

due to clear seasonality). This local heterogeneity, along with possibly limited food in a

feeding patch, and greater predictability at a larger scale might make social learning as

adults sub-optimal, as the habitat experienced by individual males might be different (see

Boyd and Richardson 1988), and/or superfluous. We also did not find any evidence for

social learning in female presence. As young male Asian elephants have been observed to

obtain some mating success through opportunistic matings (see Chelliah and Sukumar 2015,

Kabini Elephant Project, unpublished data), it is possible that learning in the context of

reproductive behaviours occurs at a younger age. It would be interesting to examine whether

there is evidence for subadult male Asian elephants socially learning from adults in mixed-

sex groups. In our study population, matriarchs of female clans were not the most central

individuals (Shetty 2016), which has also been suggested in Uda Walawe in Sri Lanka (de

Silva et al. 2011). However, while female group size constraints were found to result in

clans being split among small groups (Nandini et al. 2017), which might result in the

matriarch not being central to the clan, group size constraint alone may not lead to the

pattern we see amongst males. Although males also associated in small group sizes, young

males did not prefer older age-class males as associates more than their age-peers, and

young males did not preferentially approach old males when associations did occur.

Therefore, it appears that social learning is not the primary reason for adult male

associations, even accounting for limitations on group size.

Male associations in the presence and absence of females

We found that the association network of males was non-random in female absence but

random in female presence. There were no significant differences in network statistics and

AI values between female presence and absence, after accounting for the different number

of sightings in the two categories. Thus, the seemingly better connected network in female

absence was due to the greater amount of time that males spent in female absence.

Contrastingly, in African savannah elephants in Samburu, association networks of sexually

inactive males were significantly denser and more clustered than those of sexually active

males (Goldenberg *et al.* 2014). However, sexually active males included musth males and

it is not clear to what extent this might contribute to the significant difference between the

networks in Samburu.

If males were approaching female groups solely due to the presence of females and

independent of the presence of other males, we had predicted that male-male associations in

female presence would be random, both at an individual and at an age-class level. Our

results (network degree distribution being random and all age-classes meeting each other as

expected by chance) agree with these predictions. Moss and Poole (1983) observed that

males in Amboseli also associated at random when in mixed-sex groups. Males in multi-

male mixed-sex groups of other species have also been found to invest in affiliations with

females rather than with each other (for example, mountain gorillas, see Sicotte 1994). The

slightly larger male group size in female presence than in female absence could be a result

of multiple males tracking the same resource (female groups).

Restriction on all-male group size

We found a negative relationship between how many associates a male had and how much

time he spent with them, indicating that all-male groups have a constraint on their group

size. Males of both age-classes spent similar proportions of time in all-male groups and

experienced similar all-male group sizes, which was expected if there was age-peer

association among old but not young males, and a restriction on group size. Female group

sizes had also been found to be constrained in the Kabini population (Nandini *et al*. 2017)

and smaller compared to an African savannah elephant population (Nandini *et al*. 2018).

The group size of independent sightings of all-male groups was lower in Kabini than in the

Amboseli population of African savannah elephants (mean ± SD group size of all-male

groups with more than one male: Kabini: 2.159 ± 0.502, Amboseli: 3.325 ± 1.995;

Supplementary Material 8, see Chiyo *et al*. 2011). Low male group sizes have also been

observed in other Asian elephant populations. The average group size of adult males

(including solitary males) was 1.1 in Mudumalai Wildlife Sanctuary, southern India (Daniel

et al. 1987). The maximum all-male group size was 2 in Mudumalai (Daniel *et al*. 1987), 6

in Kabini (although observed in only one sighting), and 5 in Gal Oya in Sri Lanka (McKay

1973). In contrast, the maximum group size in Amboseli was 40 (Lee *et al*. 2011). Lee *et al*.

(2011) classified gregarious males (who spend most of their nonmusth time in all-male

groups) as promiscuous (males who spent a lot of time with many associates) or

discriminating (males who associated with fewer males for a lot of time). The presence of

promiscuous males, along with much larger group sizes, suggest that there is no strong

negative relationship between how many associates a male has and how strong his

associations are with them in Amboseli.

There are methodological differences between the current study and the one in Amboseli.

We considered 50-m as the radius for male association while Chiyo *et al.* (2011) used a 100-

m radius. However, we did not find a large number of instances of males within a 100-m

radius but outside a 50-m radius. Chiyo *et al.* (2011) also included males regardless of

musth status/sexual state (which may artificially reduce strength of association; see

Goldenberg *et al.* 2014). However, since musth males in Amboseli associated more with

female groups than in all-male groups, and the reverse was true for nonmusth males (Poole

1987), it is unlikely that the removal of musth males from the data will lead to a substantial

reduction in the all-male group sizes reported in Chiyo *et al.* (2011).

We expected a group size constraint to affect old males to a greater extent than young males

due to greater food requirements. However, we found no effect of male age on the time

spent in all-male groups. We also found no relationship between male age and the mean

experienced group size, but the mean experienced group sizes were close to 1.2, precluding

much further reduction in group sizes. Larger Asian elephant male groups are reported

elsewhere when they raid crops in risky, but resource-rich environments (see Srinivasaiah *et*

al. 2012), which would relax any constraints on group size. However, anthropogenic threat

that could result in larger group sizes is a confounding factor. Intra-group feeding

competition has been discussed as a constraint on group sizes in primates (see Chapman *et*

al. 1995), including in all-male groups (Rajpurohit 1995, Steenbeek *et al.* 2000). Future

studies on foraging in male elephants are required to find out the extent to which foraging

constraints exist and affect group sizes.

Preferred male associations and stability of associations

We did not find high correlations between associations across years, in female absence, or

evidence for preferred associations among males across two week intervals in either female

absence or presence. However, when we examined the males' top associates, we found, in

female absence, that the identity of the top associate was different from random for most

(84%) males and a majority of the pairs (61%) had a higher AI than expected by chance,

although the absolute values of these AIs were quite low (Supplementary Material 4). Thus,

there were males who showed preference in their choice of top associate but did not or

could not spend more time with those associates, and group size restriction may possibly

play a role. Such nonrandom top associates were hardly observed in the presence of

females. In all-male groups in Amboseli, most of the associations were not significant, with

less than 10% of all the observed AI values being greater than that (AI=0.1) predicted under

a model of random associations (Chiyo *et al*. 2011). However, similar to our findings in

Kabini, older (>20 years old) adult males in Amboseli also had at least one significant top

associate (Lee *et al*. 2011). Stable and significant affiliation among adult males have been

observed in several other species (Packer and Pusey 1982 - lions, Connor *et al*. 2001 -

bottlenose dolphins, Mitani 2009 - chimpanzees, Berghänel *et al*. 2011 - barbary macaques)

but as mentioned in the Introduction, significant relationships among adult males are often

thought to be a means to form coalitions to defend females. Adult male coalitions have not

been observed in Asian elephants and are unlikely, given the low probability of finding a

receptive female and the small sizes of female groups (Nandini *et al*. 2017). It will be

interesting to explore other possible reasons for the significant top male associates we find

in Kabini. It is possible that significant affiliations occur between males who are related (see

Vidya and Sukumar 2005).

In summary, we show that associations among nonmusth adult male Asian elephants,

although temporary, were affected by male age and the immediate presence of females. We

found associations with age-peers among old males in the absence of females, likely

allowing for testing strengths. There was no evidence for social learning driving male

associations. There was a constraint on all-male group size, which probably contributed to

the small proportion of time males spent in all-male groups. Males associated with each

other at random in the presence of females. When we compared our results with those

observed in African savannah elephants, we found that Kabini males spent a much smaller

proportion of their time in all-male groups of smaller sizes, possibly due to the group size

constraints found in Kabini. We also found no evidence for young males preferentially

associating with older males in Kabini, unlike the case of African savannah elephants. We

posit that the difference in the role of older males is due to the difference in the dispersion

of food resources in habitats they occupy, making social learning about resources less

valuable in adult male Asian elephants. Overall, while there were some similarities in male

associations, ecological differences possibly result in the differences in male social structure

between the two species, despite phylogenetic similarity. The relative extents to which the

presence/absence of bull areas and differences in feeding competition in non-bull areas

explain differences in group size and association time between adult males in the African

savannah and Kabini would be interesting to examine. Differences in the nature of male

associations between phylogenetically related species have been observed in other taxa:

chimpanzees and bonobos show different levels of agonism (Furuichi and Ihobe 1994),

macaque populations/species differ in the frequencies of male affiliations depending on

group sizes and group sex ratios (Hill 1994). However, while differences between related

species in overall social organisation (dolphin species: Parra *et al*. 2011, Grevy's zebra and

onager: Rubenstein *et al*. 2015) or female social organisation (colobine species: Korstjens *et*

al. 2002, African savannah and Asian elephants: Nandini *et al*. 2018) that are consistent

with resource distributions are known, there is little information on food resource

distributions differently affecting male social organisation in related species of large

mammals. Studies on the foraging ecology of male elephants are required in the future to

further understand the differences in social organisation between species.

Acknowledgments

This work was supported by the Department of Science and Technology's (Government of

India) Ramanujan Fellowship (to TNCV) under Grant No. SR/S2/RJN-25/2007 (dated

09/06/2008), Council of Scientific and Industrial Research, Government of India, under

Grant No. 37(1375)/09/EMR-II and No. 37(1613)/13/EMR-II, National Geographic Society,

USA, under Grant #8719-09 and #9378-13, and JNCASR. PK was supported as a Ph.D.

student by the Council of Scientific and Industrial Research (No. 09/733(0152)/2011-EMR-

I). This work is part of PK's Ph.D. thesis. JNCASR also provided logistic support. The

funders had no role in study design, data collection and analysis, preparation of the

manuscript, or decision to publish.

We thank the offices of the PCCF, Karnataka Forest Department, and of the Conservators of

Forests of Nagarahole and Bandipur National Parks and Tiger Reserves for field permits.

We also thank various officials, from various PCCFs and APCCFs, to the Conservators of

Forests and Range Forest Officers, to the staff of Nagarahole and Bandipur National Parks

for their support across the years. We thank Krishna, Althaf, Ranga, Shankar, Gunda,

Rajesh, Binu, and others for field assistance. We thank Deepika Prasad and Arjun Ghosh for

help with the initial field data collection. We thank Ajay Desai for useful discussions.

References

1. Alberts SC, Altmann J. 1995. Balancing costs and opportunities: dispersal in male

baboons. Am Nat. 145:279-306.

2. Baird RW, Whitehead H. 2000. Social organization of mammal-eating killer whales:

group stability and dispersal patterns. Can J Zool. 78:2096-2105.

3. Bastian M, Heymann S, Jacomy M. 2009. Gephi: an open source software for exploring

and manipulating networks. Proc Third Internat ICWSM Conf. 20098:361–362.

4. Beecher MD, Campbell SE, Stoddard PK. 1994. Correlation of song learning and

territory establishment strategies in the Song Sparrow. Proc Natl Acad Sci. 91:1450-

1454.

5. Bercovitch FB, Berry PS. 2015. The composition and function of all‐male herds of T

hornicroft's giraffe, *Giraffa camelopardalis thornicrofti*, in Zambia. Afr J Ecol. 53:167-

174.

6. Berghänel A, Ostner J, Schröder U, Schülke O. 2011. Social bonds predict future

cooperation in male Barbary macaques, *Macaca sylvanus*. Anim Behav. 81:1109-1116.

7. Best PB. 1979. Social organisation in sperm whales, *Physeter macrocephalus*. In: Winn

HE, Olla BL, editors. Behavior of marine mammals, Volume 3: Cetaceans. New York:

Plenum press. p. 227-289.

8. Biro D, Inoue-Nakamura N, Tonooka R, Yamakoshi G, Sousa C, Matsuzawa T. 2003.

Cultural innovation and transmission of tool use in wild chimpanzees: evidence from

field experiments. Anim Cogn. 6:213-223.

9. Blundell GM, Ben-David M, Bowyer RT. 2002. Sociality in river otters: cooperative

foraging or reproductive strategies? Behav Ecol. 13:134-141.

10. Bon R, Dubois M, Maublanc ML. 1993. Does age influence between-rams

companionship in mouflon (*Ovis gmelini*)? Rev Ecol-Terre Vie. 48:57-64.

11. Boyd R, Richerson PJ. 1988. An Evolutionary Model of Social Learning: The Effects of

Spatial and Temporal Variation. In: Zentall T and Galef BG, editors. Social Learning:

Psychological and Biological Perspectives. Hillsdale, New Jersey: Lawrence Erlbaum

Associates. p. 29-48.

12. Chapman CA, Chapman LJ, Wrangham RW. 1995. Ecological constraints on group

size: an analysis of spider monkey and chimpanzee subgroups. Behav Ecol Sociobiol.

36:59-70.

13. Chelliah K, Sukumar R. 2013. The role of tusks, musth and body size in male–male

competition among Asian elephants, *Elephas maximus*. Anim Behav. 86:1207-1214.

14. Chelliah K, Sukumar R. 2015. Interplay of male traits, male mating strategies and

female mate choice in the Asian elephant, *Elephas maximus*. Behaviour. 152:1113-1144.

15. Chiyo PI, Archie EA, Hollister-Smith JA, Lee PC, Poole JH, Moss CJ, Alberts SC.

2011. Association patterns of African elephants in all-male groups: the role of age and

genetic relatedness. Anim Behav. 81:1093-1099.

16. Chiyo PI, Moss CJ, Alberts SC. 2012. The influence of life history milestones and

association networks on crop-raiding behavior in male African elephants. PLoS

One. 7:e31382.

17. Clutton-Brock TH, Iason GR, Guinness FE. 1987. Sexual segregation and

density-related changes in habitat use in male and female Red deer (*Cervus elaphus*). J

Zool. 211:275-289.

18. Connor RC, Smolker RA, Richards AF. 1992. Two levels of alliance formation among

male bottlenose dolphins (*Tursiops* sp.). Proc Natl Acad Sci. 89:987-990.

19. Connor RC, Heithaus MR, Barre LM. 2001. Complex social structure, alliance stability

and mating access in a bottlenose dolphin 'super-alliance.' Proc R Soc Lond B Biol

Sci. 268:263-267.

20. Cransac N, Gerard JF, Maublanc ML, Pépin D. 1998. An example of segregation

between age and sex classes only weakly related to habitat use in mouflon sheep (*Ovis

gmelini*). J Zool. 244:371-378.

21. Curé C, Antunes R, Alves AC, Visser F, Kvadsheim PH, Miller PJ. 2013. Responses of

male sperm whales (*Physeter macrocephalus*) to killer whale sounds: implications for

anti-predator strategies. Sci Rep. 3:1579.

22. Daniel JC, Desai AA, Sivaganesan N, Datye HS, Rameshkumar S, Baskaran N,

Balasubramanian M, Swaminathan S. 1987. Ecology of the Asian elephant (Final

report). Bombay: Bombay Natural History Society.

23. de Silva S, Ranjeewa AD, Kryazhimskiy S. 2011. The dynamics of social networks

among female Asian elephants. BMC Ecol. 11:17.

24. de Silva S, Wittemyer G. 2012. A comparison of social organization in Asian elephants

and African savannah elephants. Int J Primatol. 33:1125–1141.

25. Desai AA, Johnsingh AJT. 1995. Social organization and reproductive strategy of the

male Asian elephant (*Elephas maximus*). In: Daniel JC and Datye H, editors. A week

with elephants. Bombay Natural History Society: Oxford University Press. p. 532-532.

26. Eisenberg JF, McKay GM, Jainudeen MR. 1971. Reproductive behavior of the Asiatic

elephant. Behaviour. 38:193-224.

27. Erdös P, Rényi A. 1960. On the evolution of random graphs. Publ Math Inst Hungarian

Acad Sci. 5:17–61.

28. Evans KE, Harris S. 2008. Adolescence in male African elephants, *Loxodonta africana*,

and the importance of sociality. Anim Behav. 76:779-787.

29. Farine DR. 2017. A guide to null models for social network analysis. Methods Ecol

Evol. 8:1309-1320.

30. Furuichi T, Ihobe H. 1994. Variation in male relationships in bonobos and chimpanzees.

Behaviour. 130:211-228.

31. Gerber L, Connor RC, King SL, Allen SJ, Wittwer S, Bizzozzero MR, Friedman WR,

Kalberer S, Sherwin WB, Wild S, Willems EP, Krützen M. (2020). Affiliation history

and age similarity predict alliance formation in adult male bottlenose dolphins. Behav

Ecol. 31: 361-370.

32. Ginsberg JR, Young TP. 1992. Measuring association between individuals or groups in

behavioural studies. Anim Behav. 44:377-379.

33. Goldenberg SZ, de Silva S, Rasmussen HB, Douglas-Hamilton I, Wittemyer G. 2014.

Controlling for behavioural state reveals social dynamics among male African elephants,

Loxodonta africana. Anim Behav. 95:111-119.

34. Greenwood PJ. 1980. Mating systems, philopatry and dispersal in birds and

mammals. Anim Behav. 28:1140-1162.

35. Gupta M, Ravindranath S, Prasad D, Vidya TNC. 2016. Short-term variation in sex ratio

estimates of Asian elephants due to space use differences between the sexes. Gajah

44:5-15.

36. Hill DA. 1994. Affiliative behaviour between adult males of the genus

Macaca. Behaviour. 130:293-308.

37. Hollister-Smith JA, Poole JH, Archie EA, Vance EA, Georgiadis NJ, Moss CJ, Alberts

SC. 2007. Age, musth and paternity success in wild male African elephants, *Loxodonta*

africana. Anim Behav. 74:287-296.

38. Hrdy SB. 1977. The langurs of Abu- Female and Male Strategies of Reproduction.

Cambridge: Harvard University Press.

39. Keerthipriya P. 2018. Associations, dominance interactions, and musth in male Asian

elephants in Nagarahole and Bandipur national parks, southern India. Doctoral

dissertation. JNCASR.

40. Keerthipriya P, Nandini S, Gautam H, Revathe T, Vidya TNC. 2020. Musth and its

effects on male-male and male-female associations in Asian elephants in Nagarahole-

Bandipur, southern India. J Mammal. 101:259-270.

41. Keerthipriya P, Nandini S, Vidya TNC. 2018. Effects of male age and female presence

on male associations in a large, polygynous mammal in southern India. bioRxiv.

485144.

42. Korstjens AH, Sterck EH, Noë R. 2002. How adaptive or phylogenetically inert is

primate social behaviour? A test with two sympatric colobines. Behaviour. 139:203-226.

43. Latapy M. 2008. Main-memory triangle computations for very large (sparse (power-

law)) graphs. Theor Comput Sci. 407:458–473.

44. Lee PC, Poole JH, Njiraini N, Sayialel CN, Moss CJ. 2011. Male social dynamics:

Independence and beyond. In: Moss CJ, Croze H, Lee PC, editors. The Amboseli

elephants: a long-term perspective on a long-lived mammal. Chicago: University of

Chicago Press. p. 224-237.

45. López P, Martín J. 2001. Fighting rules and rival recognition reduce costs of aggression

in male lizards, *Podarcis hispanica*. Behav Ecol Sociobiol. 49:111-116.

46. Mantel N. 1967. The detection of disease clustering and a generalized regression

approach. Cancer Res. 27:209-220.

47. McComb K, Moss C, Durant SM, Baker L, Sayialel S. 2001. Matriarchs as repositories

of social knowledge in African elephants. Science. 292:491-494.

48. McKay GM. 1973. Behavior and ecology of the Asiatic elephant in southeastern Ceylon.

Smithson Contrib Zool. 125:1–113.

49. Mitani JC. 2009. Male chimpanzees form enduring and equitable social bonds. Anim

Behav. 77:633-640.

50. Mooring MS, Fitzpatrick TA, Benjamin JE, Fraser IC, Nishihira TT, Reisig DD,

Rominger EM. 2003. Sexual segregation in desert bighorn sheep (*Ovis canadensis*

mexicana). Behaviour. 140:183-207.

51. Moss, CJ, Poole JH. 1983. Relationships and social structure of African elephants. In:

Hinde RA, editor. Primate social relationships: An integrated approach. London:

Blackwell Scientific Publications. p. 315-325.

52. Murphy D, Mumby HS, Henley MD. 2020. Age differences in the temporal stability of a

male African elephant (*Loxodonta africana*) social network. Behav Ecol. 31:21-31.

53. Nandini S, Keerthipriya P, Vidya TNC. 2017. Seasonal variation in female Asian

elephant social structure in Nagarahole-Bandipur, southern India. Anim

Behav. 134:135-145.

54. Nandini S, Keerthipriya P, Vidya TNC. 2018. Group size differences may mask

underlying similarities in social structure: a comparison of female elephant

societies. Behav Ecol. 29:145-159.

55. Packer C, Pusey AE. 1982. Cooperation and competition within coalitions of male lions:

kin selection or game theory? Nature. 296:740-742.

56. Packer C, Pusey AE. 1987. The evolution of sex-biased dispersal in lions.

Behaviour. 101:275-310.

57. Parra GJ, Corkeron PJ, Arnold P. 2011. Grouping and fission–fusion dynamics in

Australian snubfin and Indo-Pacific humpback dolphins. Anim Behav. 82:1423-1433.

58. Pereira ME. 1988. Effects of age and sex on intra-group spacing behaviour in juvenile

savannah baboons, *Papio cynocephalus cynocephalus*. Anim Behav. 36:184-204.

59. Poole JH. 1982. Musth and male-male competition in the African elephant. Doctoral

dissertation. University of Cambridge.

60. Poole JH. 1987. Rutting behavior in African elephants: the phenomenon of musth.

Behaviour. 102:283-316.

61. Poole JH, Lee PC, Njiraini N, Moss CJ. 2011. Longevity, competition and musth: a

long-term perspective on male reproductive strategies. In: Moss CJ, Croze H, Lee PC,

editors. The Amboseli elephants: a long-term perspective on a long-lived mammal.

Chicago: University of Chicago Press. p. 272-290.

62. Pusey AE, Packer C. 1987. Dispersal and Philopatry. In: Smuts BB, Cheney DL,

Seyfarth RM, Wrangham RW, Struhsaker TT, editors. Primate societies. Chicago:

University of Chicago Press. p. 250-266.

63. Rajpurohit LS. 1995. Temporary splitting or subgrouping in male bands of Hanuman

langurs, *Presbytis entellus*, around Jodhpur, western India. Mammalia. 59:3-8.

64. Rajpurohit LS, Mohnot SM, Sommer V. 1995. Wanderers Between Harems and

Bachelor Bands: Male Hanuman Langurs (*Presbytis entellus*) At Jodhpur in Rajasthan.

Behaviour. 132:255–299.

65. Rasmussen HB, Okello JBA, Wittemyer G, Siegismund HR, Arctander P, Vollrath F,

Douglas-Hamilton I. 2007. Age-and tactic-related paternity success in male African

elephants. Behav Ecol. 19:9-15.

66. Robbins MM. 1996. Male‐male Interactions in heterosexual and all‐male wild mountain

gorilla groups. Ethology. 102:942-965.

67. Rubenstein DI, Sundaresan SR, Fischhoff IR, Tantipathananandh C, Berger-Wolf TY.

2015. Similar but different: dynamic social network analysis highlights fundamental

differences between the fission-fusion societies of two equid species, the onager and

Grevy's zebra. PLoS One. 10:e0138645.

68. Ruckstuhl KE, Neuhaus P. 2000. Sexual segregation in ungulates: a new

approach. Behaviour. 137:361-377.

69. Saayman GS. 1971. Behaviour of the adult males in a troop of free-ranging chacma

baboons (*Papio ursinus*). Folia Primatol. 15:36-57.

70. Shetty N. 2016. Social Structure, Genetic Relatedness, and Dominance Relationships in

Female Asian Elephants in Nagarahole and Bandipur National Parks, Southern India.

Doctoral dissertation. JNCASR.

71. Sicotte P. 1994. Effect of male competition on male-female relationships in bi-male

groups of mountain gorillas. Ethology. 97:47-64.

72. Smith LK, Fantella SL, Pellis SM. 1999. Playful defensive responses in adult male rats

depend on the status of the unfamiliar opponent. Aggress Behav. 25:141-152.

73. Sokal RR, Rohlf FJ. 1981. Biometry (second edition). New York: WH Feeman and

Company.

74. Srinivasaiah NM, Anand VD, Vaidyanathan S, Sinha A. 2012. Usual populations,

unusual individuals: insights into the behavior and management of Asian elephants in

fragmented landscapes. PLoS One. 7:e42571.

75. StatSoft, Inc. 2004. STATISTICA (data analysis software system), version 7.0. StatSoft,

Inc., Tulsa. www.statsoft.com.

76. Steenbeek R, Sterck EHM, deVries H, van Hooff JARAM. 2000. Costs and benefits of

the one-male, age-graded, and all-male phases in wild Thomas's langur groups. In:

Kappeler PM, editor. Primate males. Cambridge: Cambridge University Press. p 130–

145.

77. Sukumar R, Joshi NV, Krishnamurthy V. 1988. Growth in the Asian elephant. Proc:

Anim Sci. 97:561-571.

78. The MathWorks, Inc. 2004. MATLAB 7 (R14). The MathWorks, Inc., Natick.

www.mathworks.com.

79. Van Hooff JA, Van Schaik CP. 1994. Male bonds: afilliative relationships among

nonhuman primate males. Behaviour. 130:309-337.

80. Vidya TNC, Prasad D, Ghosh A. 2014. Individual identification in Asian elephants.

Gajah. 40:3–17.

81. Vidya TNC, Sukumar R. 2005. Social organization of the Asian elephant (*Elephas*

maximus) in southern India inferred from microsatellite DNA. J Ethol. 23:205–210.

82. Villaret JC, Bon R. 1995. Social and spatial segregation in Alpine ibex (*Capra ibex*) in

Bargy, French Alps. Ethology. 101:291-300.

83. Wagner AP, Frank LG, Creel S. 2008. Spatial grouping in behaviourally solitary striped

hyaenas, *Hyaena hyaena*. Anim Behav. 75:1131-1142.

84. Wasserman S, Faust K. 1994. Social Network Analysis: Methods and Applications.

Cambridge and New York: Cambridge University Press.

85. Whitehead H. 2008. Analyzing Animal Societies: Quantitative Methods for Vertebrate

Social Analysis. Chicago: University of Chicago Press.

86. Whitehead H. 2009. SOCPROG programs: analyzing animal social structures. Behav

Ecol Sociobiol. 63:765-778.

87. Wich S, Sterck E. 2007. Familiarity and threat of opponents determine variation in

Thomas langur (*Presbytis thomasi*) male behaviour during between-group encounters.

Behaviour. 144:1583-1598.

1096

1097

1098

 Figure legends

 Figure 1. The proportion of time that a) males (sighted on five days or more) spent in

 different group types and b) old and young males (sighted on five days or more) spent in

 mixed-sex and adult all-male groups. Pairwise comparisons with significant results are

 marked with an asterix, and those with non-significant results are marked as *NS* (Not

 Significant). Error bars are standard deviations.

 Figure 2. The proportion of instances when the younger male approached the older male,

 and the proportion when the older male approached the younger male, to initiate association,

 in female presence and absence. The proportions were not different from one another in

 female presence and absence and hence, pairwise comparisons results are marked as *NS*

 (Not Significant).

 Figure 3. Permuted and observed numbers of times adult males (sighted 10 times or more in

 that category) of the same and different age-classes were sighted together in a) female

 absence and b) female presence. Comparisons where the observed value was significantly

 different from those of the permuted datasets are marked with an asterix. Old males are

 >=30 years and young males are 15-30 years old. Please note that the Y axis is on different

 scales in the two panels.

 Figure 4. a) Proportion of sightings of all groups containing at least one male, of different

 group sizes (different numbers of adult males) and b) the experienced group sizes of

 common males in female presence and absence. Error bars are 95% CI.

1124 Figure 5. The mean proportion of a) sightings (out of the total sightings of the male) and b)

1125 all-male group sightings (out of the total all-male group sightings of the male), that a male

1126 spent with an associate plotted against the number of associates of that male. Results from

1127 Spearman's rank-order correlations are written in the plot.

1128

1129 Figure 6. Networks of males and their top associates in a) female presence and b) female

1130 absence. Nodes representing old males are coloured dark blue and those representing young

1131 males are coloured light blue.

1132

1133

1134 **Tables**

1135

1136 Table 1. Difference in mean association strength between males of the two age-classes in

1137 observed and permuted datasets. The associated probability values (P=number of

1138 permutations whose difference in mean association strength is higher than that based on the

1139 observed dataset / number of permutations) are provided. Significant comparisons

1140 (P<0.025) are marked in bold.

1141

Category	Observed difference in mean association strength (Old-Young)	Difference in mean association strength (Old-Young) in permuted datasets Mean (2.5 and 97.5 percentiles)	P
Female presence Age-class (Old)	0.061	-0.003 (-0.0625, 0.0621)	0.027
Female absence Age-class (Old)	**0.035**	**0.003 (-0.0147, 0.0199)**	**<0.001**

1142

1143

1144

1145 Table 2. Network statistics based on observed and permuted male associations in female

1146 presence and female absence. P=(number of times difference$_{random}$ ≥ difference$_{observed}$) /

1147 number of randomisations (5000). None of the comparisons was significant.

1148

Category	Mean AI	Mean degree	Mean clustering coefficient	Mean path length	Density
Female presence observed	0.005	2.167	0.300	2.345	0.197
Female absence observed	0.003	4.500	0.577	1.697	0.409
Female presence permuted mean (SD)	0.004 (0.0011)	1.463 (0.4246)	0.224 (0.2263)	2.054 (0.4013)	0.133 (0.0386)
Female absence permuted mean (SD)	0.003 (0.0002)	4.887 (0.3184)	0.592 (0.0723)	1.623 (0.0621)	0.444 (0.0289)
P value	0.1126	0.9948	0.6316	0.2784	0.9448

1149

1150

1151

1152   **Figures**

1153

1154   Figure 1.

1155

a)

b)

1156

1157

1158

1159   Figure 2.

1160

1161

1162

1163

1164

1165   Figure 3.

1166

a) Female absence
b) Female presence

1167

1168

1169

1170   Figure 4.

1171

a)

b)

1172

1173

1174

1175   Figure 5.

1176

1177

1178

1179

1180

1181   Figure 6.

1182

a) Female presence   b) Female absence

1183

1184