Title: Co-opted transposons help perpetuate conserved higher-order chromosomal structures

Short title: Co-opted TEs maintain conserved 3D genome

One-sentence summary: Co-option of transposable elements maintains conserved 3D

genome structures via CTCF binding site turnover in human and mouse.

Authors: Mayank NK Choudhary[1], Ryan Z Friedman[1], Julia T Wang[1], Hyo Sik Jang[1], Xiaoyu

Zhuo PhD[1], Ting Wang PhD[1,*]

Affiliations: [1]The Edison Family Center for Genome Sciences & Systems Biology, Department

of Genetics, Washington University, St. Louis, MO

[*]Corresponding author:

Ting Wang, PhD, Department of Genetics,

Washington University School of Medicine, 4515 McKinley Avenue

Campus Box 8510

St. Louis, MO 63110, USA.

Phone: (314) 286-0865

E-mail address: twang@genetics.wustl.edu

3782 Words

0 Table

4 Figures

ABSTRACT (227 words)

Transposable elements (TEs) make up half of mammalian genomes and shape genome

regulation by harboring binding sites for regulatory factors. These include architectural

proteins—such as CTCF, RAD21 and SMC3—that are involved in tethering chromatin loops and

marking domain boundaries. The 3D organization of the mammalian genome is intimately

linked to its function and is remarkably conserved. However, the mechanisms by which these

structural intricacies emerge and evolve have not been thoroughly probed. Here we show that

TEs contribute extensively to both the formation of species-specific loops in humans and mice

via deposition of novel anchoring motifs, as well as to the maintenance of conserved loops

across both species via CTCF binding site turnover. The latter function demonstrates the ability

of TEs to contribute to genome plasticity and reinforce conserved genome architecture as

redundant loop anchors. Deleting such candidate TEs in human cells leads to a collapse of such

conserved loop and domain structures. These TEs are also marked by reduced DNA

methylation and bear mutational signatures of hypomethylation through evolutionary time.

TEs have long been considered a source of genetic innovation; by examining their contribution

to genome topology, we show that TEs can contribute to regulatory plasticity by inducing

redundancy and potentiating genetic drift locally while conserving genome architecture

globally, revealing a paradigm for defining regulatory conservation in the noncoding genome

beyond classic sequence-level conservation.

Keywords: 3D genome, loops, evolution, conservation, transposable elements, binding

site turnover

BACKGROUND

The 3D organization of various genomes has been mapped at high resolution using a variety of methods (1-5). While genome folding is largely conserved in mammals (1,4), the genetic forces shaping its emergence and evolution remain poorly understood. Two distinct yet mutually non-exclusive models (6) have recently gained much traction: that of phase separation (7) and of loop extrusion (8,9) by factors such as CTCF. In relation to the latter, TEs are known to contain and disseminate functional regulatory sequences (10-13) including that of CTCF. In contrast to relying on point mutations to evolve a functional CTCF binding site, TE transposition presents an attractive model for rapid regulatory sequence dissemination and regime building (14-17). Hence, we hypothesized that TEs have been a rich source of sequence for the assembly and tinkering of higher-order chromosomal structures. We studied the influence of all repetitive elements (REs) in establishing higher-order chromosomal structures and, more specifically, the role of TEs in the evolution of these higher-order chromosomal structures in humans and mice.

RESULTS

We examined REs' contribution to loop anchor CTCF sites using published genome-wide chromosomal conformation capture data from assays including ChIA-PET (2) and Hi-C in human (GM12878, HeLa, HMEC, IMR90, K562, NHEK) and mouse (ESCs, NSCs, CH12-LX) cell lines (1). We determined that 398 out of 3159 (12.6%) unique loop anchor CTCF sites were derived from REs in the mouse lymphoblastoid cell line. These RE-derived CTCF sites help establish 451 out of 2718 (16.6%) loops with discernible, unique CTCF loop anchors (Fig 1A, B). In the corresponding human lymphoblastoid cell line, REs contributed 935 out of 8324 (11.2%) unique loop anchor CTCF sites that help establish 1244 out of 8007 (15.6%) loops. Overall, REs contributed 9-15% of the anchor CTCF sites that result in 12-18% loops in humans and 12-23% of the anchor CTCF sites that result in 15-27% loops in mouse, across a variety of cell lines (Fig 1A, B).

In both species, RE-derived loop anchor CTCF sites were largely derived from TEs (>95%) and their class of origin (SINE, LINE, LTR, DNA) showed a species-biased distribution

(Fig 1C). Using the highest resolution *in-situ* HiC maps in matched lymphoblastoid cell types in

mice (CH12-LX) and humans (GM12878), we compared the composition of the RE-derived loop

anchor CTCF sites. While the mouse lineage was profoundly shaped by the SINEs (70%, 4x

enrichment over background), the human lineage was overrepresented by retroviral LTR

elements and DNA transposons (36% and 22%, 2x and 3x enriched over the background

respectively) (Fig 1D). At the family level, the B2 SINEs in mice were 13-fold enriched over

background and contributed 65% of TE-derived loop anchor CTCF sites. In humans, the hAT-

Charlie family of DNA transposons contributed 13% of TE-derived loop anchor CTCF sites, a 4-

fold enrichment over background (Fig 1E). These contributions are underestimates as we have

yet to (i) uniquely identify all loop anchor CTCF sites (especially in repetitive regions), and (ii)

annotate all repetitive elements, especially ancient TEs that have diverged far from their

identity (*18*). Further, we looked at the cell-type specificity of these loop anchor CTCF sites in

humans and see that 1334 out of 2017 (66%) RE-derived loop anchor CTCF sites were found in

only one cell type (Supplementary Figure 1A). However, we did not find any specific TE family

that enriches for cell-type specific loop anchor CTCF sites in the cell lines profiled

(Supplementary Figure 1B).

To study the evolution of chromatin loops, we compared their conservation

(Supplementary Methods) in matched human and mouse cell-types. Briefly, we used the

liftOver tool (*19*) to compare loops across species and required exactly one reciprocal match

(reciprocal best hit) to designate conserved loops. We found that 48% of all mouse loops (1596

out of 3331) had a loop call in the corresponding syntenic region in humans (Table S1.1). Our

observation is in close agreement with prior studies (*1,4*) that show about half of all higher-

order chromosomal structures to be conserved. We then sought to characterize the

contribution of TEs to various classes of loops based on their orthology.

We compared the origin of loop anchor CTCF sites of orthologous loops in mouse and

human. We found that out of 1596 orthologous loops, 142 (8.9%) in mouse and 108 (6.7%) in

human had at least one TE-derived loop anchor CTCF site (Fig 2A). In addition to orthologous

loops, TE-derived loop anchor CTCF sites also gave rise to 24% (409 out of 1735) and 15% (1136

out of 7852) non-orthologous (species-specific) loops in mouse and humans, respectively (Fig

2A), consistent with the appreciable role of TEs in genome innovation (*14-16,20,21*). Overall,

the majority of TE-derived loop anchors in mouse were established by a handful of young TE

subfamilies (B3, B2_Mm2, B3A, B2_Mm1t) that expanded in the rodent lineage (*22*) (Fig 2B). In

contrast, multiple TE subfamilies of varying evolutionary ages contributed diffusely to CTCF

loop anchors in humans (Fig 2C). Altogether, TEs in humans contributed to fewer orthologous

loops and distributed over more TE subfamilies than in mouse.

Intriguingly, 123/142 (87%) TE-derived orthologous loops in mouse were discordant for

TEs in humans (Table S1.2). In the sense: while the loops in humans were anchored at the

putative ancestral CTCF binding sites, the syntenic ancestral CTCF motifs were largely

degraded or deleted in mouse and the loops were now anchored at CTCF sites derived from

nearby, co-opted TEs instead. One such example is an orthologous loop at the 5′ end of the

Akap8l gene (Fig 2D) maintained in mouse by a MER20 element transposed ~1.5kb upstream

of the degraded ancestral motif which was well conserved in most non-rodent mammals

(Supplementary Figure 2). The degradation of the ancestral CTCF motif derived from an

ancient MIR3 element that is over 147 million-years-old (see *Methods*) incapacitates CTCF

binding as evidenced by the CTCF-ChIP track (Fig 2E). In contrast, the younger MER20 element

that inserted ~90 million-years ago harbored strong CTCF binding, providing an anchor site to

maintain the conserved loop in mouse. Similarly, we find that 89/108 (82%) TE-derived

orthologous loops in human GM12878 cells were discordant for TEs in mouse (Table S1.3). We

hypothesized that TEs provide redundant CTCF sites and mediated binding site turnover for

CTCF contributing to conserved genome folding events between human and mouse.

Moreover, the 123 turned-over loops in mouse represent 127 turnover events (4 loops

had both loop anchors turned-over) mediated by 124 unique loop anchors (3 turned-over loop

anchors tethered 2 loops each). Out of the 124 unique loop anchors, 61 events represent

turnover of the left loop anchor and 63 events represent turnover of the right loop. In terms of

CTCF motif orientation—for the 61 left loop anchor turnover events, 53 were positive and 8

were negative; and for the 63 right loop anchor turnover events, 45 were negative and 18 were

positive (Chi-square test, p-value=5.3×10^{-11}). Similarly, in humans the 89 turned-over loops

represent 93 turnover events (4 loops had both loop anchors turned-over) were mediated by 84

unique loop anchors (1 turned-over loop anchor tethered 3 loops, and 7 loop anchors tethered 2

loops each). Out of the 84 unique loop anchors, 43 events represent turnover of the left loop

anchor (43 positive orientation CTCF motif and 0 negative orientation CTCF motif), and 41

events represent turnover of the right loop (40 positive orientation CTCF motif and 1 negative

orientation CTCF motif) (Chi-square test, p-value=3.6×10^{-19}). These results further lend

credence to the loop extrusion model (8) and suggest that TE exaptation is more likely when

the orientation of the inserted TE (and the underlying CTCF motif provided) is compatible with

the local loop structure.

mm9 CH12-LX (n=124)	Left Loop Anchor	Right Loop Anchor
+ve CTCF motif	53	18
-ve CTCF motif	8	45

hg19 GM12878 (n=84)	Left Loop Anchor	Right Loop Anchor
+ve CTCF motif	43	1
-ve CTCF motif	0	40

Since the mouse genome is replete with repeat-derived CTCF sites (22) that could

interfere with the targeted study of specific TE candidates, we decided to validate these

hypotheses in human cell lines.

Here we examine two candidate TEs that maintain conserved higher-order

chromosomal structures in humans: one belonging to the L1M3f subfamily of LINEs, and the

other belonging to the LTR41 subfamily of endogenous-retrovirus-derived long terminal

repeat (LTR). The former TE replaces the function of a lost ancestral CTCF site (Supplementary

Figure 3), while the latter is functionally redundant for an ancestral CTCF site still present in

humans (Supplementary Figure 4). These two TEs were specifically chosen as they could be

unambiguously attributed to the genome folding function (no other CTCF/Cohesin binding site

in the vicinity). Using CRISPR-Cas9, we obtained clones of GM12878 cells bearing homozygous

deletions of the L1M3f and LTR41 elements, respectively (Supplementary Figure 5, Table S2.4).

We then performed HYbrid-Capture on the *in situ* Hi-C library (Hi-C^2) to examine the effect of

the TE deletion on the local 3D structure (*8*) (Table S2.1, S2.2, S2.3).

The L1M3f-derived CTCF site was positioned at a conserved domain border and

anchored three chromatin loops (Supplementary Figure 3). Upon deletion of this L1M3f, the

conserved local chromosomal structure collapsed as evidenced by (i) the loss of focal

enrichment in the homozygous TE knockout (KO) contact map in comparison to the wild-type

(WT) contact map, and (ii) the fusion of two neighboring domains (Hi-C^2 results: Fig 3A, Hi-C

results: Supplementary Figure 6). The Virtual 4C plot anchored at the region surrounding the

L1M3f element showed three distinct peaks (corresponding to the three loops in the WT cell

line), which were lost in the KO (ΔL1M3f) cell line. We also found that cross-domain

interactions significantly increased from 8% in WT to 19% in KO cell lines (~2.4x, Welch's t-test

p-value<1.5x10^{-16}, Table S2.5) across the L1M3f-established domain boundary, a change

specific to the targeted domain and not seen in a control domain from a nearby region (Fig 3C).

Thus, the L1M3f element is necessary for maintaining the conserved loops and domain

boundary in humans. It represents a novel class of binding site turnover (*23-26*) for CTCF

leading to conservation in terms of function via establishment of long-range interactions and

potentially the underlying gene regulation, but not in primary local sequence.

Our second candidate was a species-specific LTR41-derived CTCF site ("c" in Fig 3D, E)

that replaced an ancestral CTCF site derived from a much older TE ("d" in Fig 3D, E) of the

MER82 subfamily that is conserved in humans and mouse. The ancestral MER82-derived CTCF

site was "decommissioned" as the LTR41 insertion (after the primate-rodent split) provided a

negative orientation CTCF motif upstream of the MER82 element. Based on the loop extrusion

model, the LTR41-derived CTCF motif would be encountered before the MER82-derived CTCF

site and hence the ancestral site is mostly decommissioned in present-day human genome as

evidenced by the drastically reduced CTCF binding (Supplementary Figure 4B). In the WT

contact map, we observed a bright focal enrichment corresponding to CTCF binding sites a-c

suggesting a looping interaction. In contrast, there was little focal enrichment corresponding

to a-d (Fig 3D, top row). Additionally, in the WT Virtual 4C track anchored on "a", we observed

a clear peak corresponding to LTR41 ("c") suggesting an a-c loop (Fig 3E). Upon deletion of

LTR41, the conserved loop's anchor is offset to the MER82-derived CTCF site ("d") downstream

of the LTR41 as evidenced by the shift in the focal enrichment in the KO contact map (Fig 3D,

bottom row) and an increase in the KO Virtual 4C peak corresponding to the MER82-derived

CTCF site (i.e., a-d loop) (Fig 3E, Supplementary Figure 7). Upon anchoring the Virtual 4C on a

5-kb window containing LTR41 (c), we observed a peak loss at "a" corresponding to the loss of

the a-c loop in the KO, an interaction that existed in the WT cells (Fig 3F). With the ~39kb shift

of the anchor site, the half-megabase scale chromosomal structure around the anchor region

remained largely preserved (Supplementary Figure 4C). Upon deletion of this TE candidate,

the local sequence configuration probably resembled that of the pre TE-insertion, ancestral

genome. This example therefore illustrates a potential path by which the local 3D genome

evolved upon insertion of the LTR41 element as well as the plasticity TEs, like LTR41 and

MER82 in this case, can encode in their host genomes by providing redundant CTCF binding

sites.

These results support the hypothesis that TEs are able to contribute regulatory

robustness and strengthen conserved regulatory architecture as redundant or "shadow" loop

anchors. The mouse genome that underwent a lineage-specific expansion of SINE B2s (22),

which carry a CTCF binding motif, is saturated with such events.

TEs are typically silenced by host repressive machineries including DNA and histone

methylation (27-29). However, a small fraction of TEs escape epigenetic silencing and provide

functional regulatory elements for the host in a process termed exaptation (30-33). Since CTCF

is a methylation sensitive chromatin factor and only binds to unmethylated DNA (34,35), we

examined the DNA methylation levels of loop anchor CTCF sites of orthologous loops

(Supplementary Methods). We found that TE-derived CTCF sites were marked by reduced

DNA methylation, similar to their non-TE derived genomic counterparts (Fig 4A). To

understand the DNA methylation dynamics through evolution, we took advantage of the

differential mutation rate of 5-methylcytosine (5mC) to Thymine (T) (36). Unmethylated

cytosines (C) mutate to T at a lower rate than 5mC; thus, methylated DNA exhibits higher

frequency of C to T mutations (37). We found that TEs involved in turnover events had a

significantly lower frequency of methylation-associated C-to-T and G-to-A mutations

compared to an identically sampled background of TEs not involved in looping (1000

simulations), but no difference in all other combined substitutions (summarized human results:

Fig 4B; full human and mouse results: Supplementary Figure 8, 9, Table S3). These results

suggest that TEs providing CTCF turnover were hypomethylated over evolutionary time to

maintain their functional role, compared to other TE copies (Fig 4C).

DISCUSSION

TEs have substantially contributed to higher order chromatin structures by serving as

chromatin loop anchors—a large fraction of which were found to be species-specific,

confirming TEs' role in genome innovation. Pioneering work in the last decade has extensively

outlined this contribution of TEs in shaping gene regulatory networks by depositing TF binding

sites in host genomes, leading to the origins of novel phenotypes like innate immunity and

pregnancy in mammals. Herein, lies the catch: research to date showcases the role of TEs in

bringing novelty and new regulatory functions to the host genome. Hence, TEs have long been

considered a source of genetic innovation. However, by comparing topologies instead of raw

DNA sequences in this study, for the first time, we have been able to reveal the role of TEs in

3D genome conservation. This seemingly counter-intuitive role of species-specific parasitic

sequences in helping maintain ancestral genome architecture is fundamentally different from

all current and previous work regarding TEs' role in gene regulation. This role is mediated by a

long-postulated, classic genetic phenomena of binding site turnover—for CTCF in this case.

Redundant TE-derived CTCF sites in the vicinity of conserved chromatin anchor/ boundary can

sometimes take over from the conserved anchor/boundary element, thus slightly shifting the

anchor/boundary site while largely maintaining the 3D structure. Certain TE subfamilies like

mouse SINE B2s contain pre-existing CTCF motifs within them, while others like mouse

RLTR30 provide sequence fodder which upon a couple of specific point mutations can acquire

CTCF binding and potentiate this binding site turnover.

In this study, 123 turnover events were observed in mouse on the basis of 3331

annotated loops (3.7%) whereas in humans 89 turnover events were observed out of 9448

loops (0.94%). This 4-fold higher rate of turnover events in mouse highlights differences in

between species and the turnover phenomenon being investigated. The higher rate of loop
anchor CTCF turnover in the mouse genome was amplified by the arrival of CTCF-motif
containing B2 elements. The genome is replete with such events and we have for the first time
functionally dissected and validated them in the context of 3D genome conservation, opening
the doors up for such investigations in the field for enhancer or promoter turnover events.

The *fons et origo* of CTCF motifs in B2 SINEs has been extensively researched. B2 SINEs
are derived from tRNA genes. Mouse tRNA genes have been shown to possess classical
insulator activity and the potential to function as boundary elements (*38*). Moreover, CTCF-
binding enrichment in B2 SINEs and repeat-driven dispersal of CTCF-binding has been shown
to be a fundamental, ancient, and still highly active mechanism of genome evolution in
mammalian lineages (*22*).

Similarly, the role CTCF motifs in viral genome regulation has been a topic of
tremendous interest and investigation. In EBV, this control involves direct binding of CTCF
across the viral genome and the formation of three-dimensional loops between virus
promoters and enhancers (*39*). CTCF has been shown to be important in the regulation of gene
expression of a number of human DNA viruses (*40*). CTCF also plays a critical role in epigenetic
regulation of viral gene expression to establish and/or maintain a form of latent infection that
can reactivate efficiently (*41*). Recent evidence has also shown that HTLV-1 inserts an ectopic
CTCF binding site forming loops between the provirus and host genome, altering expression of
proviral and host genes (*42*). CTCF has also been shown to promote HSV-1 lytic transcription
by facilitating the elongation of RNA Pol II and preventing silenced chromatin on the viral
genome (*43*). Moreover, one can speculate that having a CTCF motif can not only help in
maintaining viral genome confirmation but can also help insulate the chromatin activity of the
neighborhood wherein the virus inserts into the host genome. It may also increase the chances
of long-range interactions taking place which can sometimes bring in other TFs and/or
polymerase, leading to enhanced transcription at the site of viral integration.

Our in-depth analysis of 3D genome structures upon genetic manipulation of candidate
TEs revealed principles of how 3D genome evolves. In one example, a human TE provided a
conserved chromatin boundary and loop anchor, whereas the ancestral CTCF site had decayed.

Upon deletion, the chromatin domains collapsed, and loops eliminated, underscoring the
importance of the TE in maintaining the local 3D genome structure.

In another case where a human TE provided a similarly conserved boundary and loop
anchor, the ancestral CTCF site was still recognizable but was decommissioned. Deletion of the
TE resulted in reinstallation of the ancestral CTCF site to form a slightly shifted boundary and
loop anchor, and the local chromatin domains were largely preserved. In this second case that
we validated, we undid the events that took place during the course of (tens of millions of
years) evolution by removing a young TE (LTR41) and having the ancestral "decommissioned"
TE (MER82) re-uptake its function, thereby "reversing" the path of evolution in a dish (in days).
Thus, experimentally demonstrating the evolutionary impact of a TE-derived CTCF site.
Moreover, the concept of such shadow loop anchors residing in TEs that can be activated upon
escape from epigenetic silencing is extremely crucial to take into account for studies pertaining
to diseases of the epigenome like certain cancers, their treatment and therapy. This study also
underscores the redundancy that exists in the genome when it comes to CTCF binding sites
and can potentially explain why we may not always see a change in 3D genome structure upon
deleting CTCF binding sites.

It is important to remember that the contribution outlined in this manuscript are
underestimates as we have yet to (i) uniquely identify all loop anchor CTCF sites (especially in
highly repetitive regions), (ii) annotate all repetitive elements, especially ancient TEs that have
diverged far from their identity (18), and (iii) identify other architectural proteins and expand
this framework beyond just CTCF-derived loop anchors.

While most studies highlight TEs' role in innovating new functions by providing novel
regulatory elements such as enhancers and promoters, we implicate the role of TEs in
functional conservation inviting us to reexamine this unconventional role—perhaps many novel
regulatory elements derived from TEs are not creating new functions, but rather providing
redundant genetic material thus contributing to the robustness of gene regulatory networks.
These findings will undoubtedly stimulate investigations to explore the multitude modes of
regulatory evolution mediated by TEs. Indeed, recent evidence has linked the transcriptional

activation of retrotransposons to restructuring of genome architecture during human
cardiomyocyte development (44).

A major caveat of the analysis presented in this study is that the *in situ* Hi-C maps (re-
analyzed in this study) of the 9 cell lines were sequenced to varying depths, and thus differ in
their resolution and "completeness" of loop annotations. Hence, due to this limitation of
publicly available high-resolution HiC data, our findings likely represent a lower bound of TE's
involvement in shaping both the conserved and species-specific 3D genome. These analyses
need to be revisited as and when higher-resolution datasets are available.

Lastly, our study opens the doors for population-scale genetic variation studies that
identify polymorphic TE insertions to be reconciled with population-scale 3D genome and
regulatory variation. These future explorations will present yet another vignette of
transposable elements and their very many roles in accelerating adaptive evolution.

CONCLUSIONS

Taken together, our findings reveal a formerly uncharacterized role that TEs have
played in the evolution of higher-order chromosomal structures in mammals. TEs have
contributed a substantial number of loop anchors in mouse and human 3D genomes, a fraction
of which were co-opted to help maintain conserved higher-order chromosomal structures. TE
transposition provides redundant CTCF motifs and a novel method for CTCF binding site
turnover to maintain regulatory conservation (defined here as the preservation of long-range
chromosomal interactions, loop and boundary formation), by compensating for the loss of
local primary sequence—local sequence that would have otherwise allowed the assessment of
purifying selection. Deletion of these TEs in human cell lines eliminated the chromatin loops
that they anchor and resulted in collapse of conserved chromatin structure, as expected by our
hypothesis. More strikingly, we demonstrate that in another case the loop anchor shifted to an
alternative TE-derived CTCF site nearby, resulting in largely unchanged chromatin structure,
underscoring the dynamic nature and robustness of the 3D genome upon TE infiltration. These
TEs that maintain conserved chromatin loops via turnover are hypomethylated through deep
time, an observation that highlights the intimate interplay between genome, epigenome, and

3D genome in evolution. This research provides a foundation to study the impact of TEs and
expand our understanding of chromosomal folding—its emergence, maintenance and
transformation—in the context of evolving genomes. Ultimately, our study reveals how selfish
genetic elements, regardless of their origins, can be repurposed to provide redundant TF
motifs, maintain latent genome sanctity and regulatory fidelity by conserving 3D structure.

FIGURE LEGENDS
Figure 1: Contribution of repetitive elements (REs) to chromatin loops in humans and
mouse. (**A**) Pie charts representing percentage of loops and (**B**) unique loop anchor CTCF sites
derived from REs in a variety of human and mouse cell types. (**C**) Stacked bar plots showing the
distribution of RE-derived anchor CTCF across major RE classes in the various human and
mouse cell types. Stacked bar plots showcasing the distribution of RE-derived anchor CTCF vs.
background and CTCF ChIP peaks across (**D**) major RE classes and (**E**) major RE families in
matched blood lymphoblastoid cell line (mouse = CH12-LX; human = GM12878).

Figure 2: Contribution of TEs in the conservation landscape of human and mouse loops. (**A**)
Venn diagram representing the various classes of chromatin loops based on their orthology
and bar plots showing the contribution of REs to anchor CTCFs of each class of loops. (**B**) Age
distribution and age of individual TEs that contribute loop anchor CTCF sites (black dots for
orthologous loops; gold dots for non-orthologous loops) (left), total contribution to loop
anchor CTCF sites (middle), distribution of orthologous and non-orthologous loops (right)
derived from the top 13 TE subfamilies in mouse and (**C**) humans. Estimated primate/rodent
divergence time (82 million years ago) is from (Meredith et al, 2011). (**D**) Contact maps
representing a conserved chromatin loop in a syntenic region between human and mouse (**E**) A
MER20 transposon insertion provides a redundant CTCF motif that helps in maintaining the
conserved 3D structure via CTCF binding site turnover with remnants of the ancestral CTCF
motif, well conserved in most non-rodent mammals (Supplementary Figure 2), still seen in the
mouse genome.

Figure 3: TEs are necessary for maintaining conserved higher-order chromosomal structures in humans. (**A**) Results of a CRISPR/Cas9-based deletion of an L1M3f element at chr10:26–28 Mb in GM187278 cells. Mega-contact maps (details in Methods) generated using Hi-C² technology for the (top) WT locus and (bottom) KO (ΔL1M3f) locus. (**B**) Virtual 4C plot displaying total percent interactions emanating from an anchor on a 5kb-window containing the L1M3f element. (**C**) Boxplot measuring the percent inter-domain interactions (Table S2.5) across the targeted domain and a control domain (boundaries unaffected by CRISPR edits) using subsampled contact maps (details in Methods). (**D**) Results of CRISPR/Cas9-based deletion of an LTR41 element at chr8:70.3–71.8 Mb in GM187278 cells. Mega-contact maps generated in Hi-C² experiments for the (top) WT locus and (bottom) KO (ΔLTR41) locus. (**E**) Virtual 4C plot displaying total percent interactions emanating from an anchor on a 5kb-window containing the left anchor CTCF of the conserved loop, and (**F**) the LTR41 element.

Figure 4: Turnover TEs are hypomethylated through evolutionary time. (**A**) Methylation signature ±2kb around CTCF sites that help maintain orthologous loops segmented by the origin of the anchor CTCF site (**B**) Methylation-associated and non-methylation mutational signature of individual TEs relative to its ancestral sequence in humans (mouse TE data available in Supplementary Figure 8). Alignments were performed using crossmatch (shown here) and Needle (details in Methods, results in Supplementary Figure 9). Error bars show one standard deviation of the means from 1000 simulations. (**C**) Schematic depicting the framework of TE-mediated CTCF binding site turnover that highlights the intimate reciprocity between the TE, genome and epigenome, to help maintain conserved 3D genome.

REFERENCES
1. SSP Rao et al., *Cell* **159**, 1665-1680 (2014).
2. Z Tang et al., *Cell* **163**, 1611-1627 (2015).
3. T Sexton et al., *Cell* **148**, 458-472 (2012).
4. JR Dixon et al., *Nature*. **485**, 376-380 (2012).
5. DU Gorkin et al., *Cell Stem Cell* **14**, 771-775 (2014).
6. W Schwarzer et al., *Nature* **551**, 51–56 (2017).
7. AR Strom et al., *Nature* **547**, 241–245 (2017).
8. AL Sanborn et al., *Proc Natl Acad Sci.* **112**, E6456-E6465. (2015).
9. G Fudenberg et al., *Cell Rep.* **15**, 2038–2049 (2016).
10. V Sundaram et al., *Genome Res.* **24**, 1963-1976 (2014).

11.   G Bourque et al., *Genome Res*. **18**, 1752-1762 (2008).

12.   G Kunarso et al. *Nat Genet*. **42**, 631-634 (2010).

13.   PÉ Jacques et al., *PLoS Genet*. **9**(5) (2013).

14.   T Wang et al., *Proc Natl Acad Sci*. **104**, 18613-18618 (2007).

15.   EB Chuong et al., *Science* **351**, 1083-1087 (2016).

16.   VJ Lynch et al., *Nat Genet* **43**, 1154-1159 (2011).

17.   BJ Britten, EH Davidson. *Q Rev Biol*. **46**, 111-138 (1971).

18.   APJ de Koning et al., *PLoS Genet*. **7** (2011).

19.   AS Hinrichs, *Nucleic Acids Res*. **34**, 590-598 (2006).

20.   C Feschotte, EJ Pritham. *Annu Rev Genet*. **41**, 331-368 (2007).

21.   V Sundaram, T Wang. *BioEssays*. **40**, 1700155 (2018).

22.   D Schmidt et al., *Cell* **148**, 335-348 (2012).

23.   MZ Ludwig et al., *Development* **12**, 3325–3330 (1998).

24.   AM Moses et al., *PLoS Comput Biol*. **2**, 130 (2006).

25.   S Venkataram, JC Fay. *Genome Biol Evol*. **2**, 851–858 (2010).

26.   D Villar et al., *Nat Rev Genet*. **15**, 221-233 (2014).

27.   MA Matzke et al., *Plant Mol Biol*. **43**, 401-415 (2000).

28.   JA Yoder et al., *Trends Genet*. **13**, 335-340 (1997).

29.   RK Slotkin et al., *Nat Rev Genet*. **8**, 272-285 (2007).

30.   A Huda et al., *Mob DNA* **1**, 2-12 (2010).

31.   CB Lowe, D Haussler. *PLoS One* **7**, 43128 (2012).

32.   G Bejerano et al., *Nature* **441**, 87–90 (2006).

33.   MG Kidwell, DR Lisch. *Trends Ecol Evol*. **15**, 95-99 (2000).

34.   C Kanduri et al., *Curr Biol*. **10**, 853-856 (2000).

35.   S Kurukuti et al., *Proc Natl Acad Sci*. **103**, 10684-10689 (2006).

36.   JC Shen et al., *Nucleic Acids Res*. **22**, 972-976 (1994).

37.   AP Bird. *Nucleic Acids Res*. **8**, 1499–1504 (1980).

38.   T Ebersole et al. *Cell Cycle*. **10**, 2779–2791 (2011).

39.   I Tempera et al. *PLoS Pathog* **7**:e1002180 (2011).

40.   I Pentland et al. *Viruses* **7**:3574-85 (2015).

41.   JS Lee et al. *mBio* **9**:e02372-17 (2018).

42.   Y Satou et al. *Proc Natl Acad Sci*. **113**(11):3054-3059 (2016).

43.   F Lang et al. *Sci. Rep.* **7**, 39861 (2017).

44.   Y Zhang, T Li, S Preissl et al., *biorXiv*. (2018).

45.   RW Meredith et al., *Science* **334**, 521-524 (2011).

46.   RH Waterston et al., *Nature* **420**, 520-562 (2002).

47.   M Haeussler et al., *Genome Biol*. **17**, 148 (2016).

48.   MA Moreno-Mateos et al., *Nature Methods* **12**, 982-988 (2015).

49.   W Bao et al., *Mob DNA* **6**, 11 (2015).

50.   A Smit et al., RepeatMasker Open-4.0.6 2013-2015. (2017).

51.   P Rice et al., *Trends Genet*. **16**, 276-277 (2000).

METHODS

Dataset GEO accession numbers:

The genomic data analyzed in this study were obtained from publicly available
datasets. HiC datasets were obtained from GSE63525 (mouse: CH12; humans: GM12878, HeLa,
HMEC, IMR90, K562, NHEK). GM12878 ChIA-PET dataset was obtained from GSE72816.
GM12878 CTCF ChIP-seq datasets were obtained from ENCODE (ENCSR000AKB and
ENCSR000DZN). CH12 CTCF ChIP-seq datasets were obtained from Mouse ENCODE
(ENCSR000ERM and ENCSR000DIU). WGBS methylation dataset for GM12878 was also
obtained from ENCODE, GEO: GSE86765 (ENCSR890UQO). Mouse ESC and NSC HiC data was
obtained from PMID: 30414923.

Loop anchor CTCF–RE intersection:

We generated a list of unique anchor CTCF sites using the HiCCUPS output[1] for various
mentioned cell lines. We then overlapped loop anchor CTCF motifs identified using HiCCUPS
(1) with *RepeatMasker* (RMSK v4.0.7, for hg19 and mm9) and required at least 10bp of the core
CTCF motif to intersect with a repetitive element (RE) to call it a RE-derived loop anchor CTCF
site. Further, only loops with (i) at least one known RE-derived anchor CTCF site, or (ii) two
non-RE derived anchor CTCF sites were taken into consideration for analysis of RE-derived
loop counts, because we can definitively say whether the loops and their loop anchor CTCF
sites were derived from REs or not. Loops with both unidentified loop anchor CTCF sites, or
one unidentified and one non-RE derived anchor CTCF site were not considered as there is the
possibility of having at least one of the other anchor CTCF sites derived from a RE. We followed
the same methodology when considering ChIA-PET loops.

TE class and family distribution

We ran RepeatMasker v4.0.7 with the -s slow search parameter on the hg19 and mm9
genomes to obtain a comprehensive list of REs in the genome and their corresponding
subfamily, family and class annotations. We used RE counts (generated as previously outlined)
to characterize their distribution to loop anchor CTCF sites. For characterizing RE-derived
CTCF binding peaks, we repurposed a previously used strategy (10). Briefly, we required that
the centers of the MACS-called peaks of ENCODE-generated CTCF ChIP datasets overlapped
with RE fragments. We used the length distribution of various RE family and classes in the
entire genome as the background distribution.

Loop orthology check:

We used liftOver (19) to convert CH12 loop annotations from mm9 mouse genome
coordinates to hg19 human genome coordinates. We used various sequence match rates
(minMatch = 0.05, 0.1..., 1) to convert CH12 mouse peaks from mm9 genome coordinates to
hg19 genome coordinates. To optimize for the minMatch parameter, we generated ten

shuffled (randomized) peak annotations by using bedtools shuffle –chrom command to
permute their location on the chromosome of origin. minMatch parameter of 0.1 was chosen
for liftOver analyses henceforth, as it resulted in the greatest number of features being lifted
over (on average) and lower coefficient of variation across the 10 simulated sets. We lifted over
3245 out of 3331 mouse peaks from mm9 to hg19, using the minMatch 0.1, to facilitate cross-
species peak annotations comparison. To call a mouse feature conserved in humans, we
required that the loop anchor pairs individually lie within a min(half of loop length, vicinity
threshold) window of an existing loop anchor pair. The vicinity threshold was put in place to
account for cross-species liftOver errors and facilitate comparison of higher-order
chromosomal features that vary from 120Kb to 125Mb in length (in mouse). We tested multiple
vicinity thresholds ranging from 500bp to 100Mb and identified false discovery rates using
simulated sets of mouse features and comparing them to the orthology observed between the
real CH12 (mouse) and GM12878 (human) features. We decided to use 50kb as the vicinity
threshold as it corresponded to a false discovery less than 0.1. We found that 1688 CH12
mouse peaks overlapped at least one corresponding peak in GM12878 human lymphoblastoid
cells. We performed a similar analysis to compare 'muranized' human features (liftOver from
GM12878) to actual mouse features (CH12). We found that 1900 GM12878 human peaks
overlapped at least one corresponding peak in CH12 mouse lymphoblastoid cells. We then
filtered for features that displayed reciprocal matches (reciprocal best hits) in the two
comparisons (mouse-to-human and human-to-mouse) as stated above. Finally, we curated the
list by considering genic, epigenomic and transcriptomic synteny to pick exactly one
orthologous human loop to a corresponding mouse loop, to enlist 1596 high-confidence
orthologous peak calls (Table S1.1). A brief flowchart of the pipeline is shown below:

494
495

 <u>TE age estimation:</u>

         Species divergence times were based on (*45*). Repeat ages were estimated by dividing
 the percent divergence of extant copies from the consensus sequence by the species neutral
 substitution rate. Substitution rates (mutations/yr) used were as follows: humans: 2.2×10^{-9};
 mouse: 4.5×10^{-9}, from (*46*). Jukes-Cantor and Kimura distances were calculated by aligning
 each TE to its consensus sequence and counting all possible mutations (see below). Single
 nucleotide substitution counts were normalized by the length of the genomic TE minus the
 number of insertions (gaps in the consensus). These mutation rates were then used to calculate
 the Jukes-Cantor and Kimura distances for each genomic TE.

 <u>Candidate selection and filtering:</u>

         After manually curating the list of conserved loops, we looked for TE-derived
 orthologous loops in humans that were discordant for TEs in mouse. After identifying the list
 of TE-derived CTCF turnover events in humans, we comprehensively surveyed the local CTCF
 binding landscape (CTCF ChIP-seq peaks) to ensure (i) there weren't other CTCF binding sites
 in the vicinity that could function as loop anchors in humans (in the first case); and (ii) there
 was only one other unique CTCF binding site, i.e. the ancestral CTCF motif (in the second case).
 We also ensured that the TE insertion from which the loop anchor CTCF site was derived was
 human-specific and not present in mouse (Table S1.2). We repeated this analysis to identify
 TE-mediated turnover in mouse as well (Table S1.3). We also identified events wherein TEs
 mediated turnover events both in mouse as well as human (Table S1.4). One possible
 explanation for this observation is that similar selective pressures (i.e. the need to maintain
 higher-order chromosomal structure) led to the convergent co-option of species-specific TEs
 at syntenic locus, independently in both the genomes.

 <u>Cell culture methods:</u>

         GM12878 cell lines were grown between 200K-800K cells/ml in 10ml cultures in T-25
 flasks, in a humidified incubator with 95% CO_2 at 37°C in RPMI1640 media (Gibco, 1187-085)
 supplemented with 15% fetal bovine serum (Corning, 35-011-CV) and 100U/ml penicillin-
 streptomycin (Gibco, 15140-122) as per the ENCODE standards.

 <u>CRISPR-Cas9 mediated genome engineering:</u>

         Our CRISPR workflow consisted of the following steps: We identified turned over
 chromatin loops that are maintained by TEs, with unique, correctly oriented TE-derived CTCF
 motifs within loop anchors (*1*). We used two independent CRISPR sgRNA design engines
 CRISPOR (*47*) and CRISPRScan (*48*) to rationally design multiple pairs of sgRNAs that have
 high cutting efficiency and minimizing off-target effects. We used pU6-(BbsI)_CBh-Cas9-T2A-
 BFP plasmid (Addgene, 64323) and pU6-(BbsI)_CBh-Cas9-T2A-mCherry plasmid (Addgene,

64324) as the CRISPR delivery vectors. For each sgRNA, we designed and annealed two single-
stranded oligos with compatible overhangs that can be cloned into BbsI-digested BFP and
mCherry CRISPR vectors through standard ligation techniques. For every pair of sgRNAs, we
constructed BFP-CRISPR vectors and mCherry-CRISPR vectors that express sgRNAs targeting
upstream and downstream of the candidate TEs, respectively. BFP-CRISPR vectors and
mCherry-CRISPR vectors each were co-transfected into GM12878 cells in antibiotic-free media
using the Neon transfection system. After 24 hours of incubation, the transfected cells were
analyzed by flow-cytometry (Beckman Coulter MoFlo) for BFP-positive and mCherry-positive
subpopulations. Transfection efficiencies were usually between 3-5%. We single-cell sorted
these double-positive fluorescent cells into 96-well plates for clone expansion and allowed to
grow for 21-28 days. After that, 20-48 clones were screened per transfection. Genomic DNA
from CRISPR clones was extracted using *Quick-DNA* Miniprep kit for genotyping and validated
with Sanger sequencing. Details of sequences used to generate clones used in this study are
listed in Table S2.4. We then performed *in situ* Hi-C on the selected mutated cell lines and
performed hybrid selection on the in situ Hi-C libraries for a region around the targeted TE to
generate Hi-C^2 libraries that can easily and cheaply be sequenced to read off the effects of our
TE deletions on local genome folding.
551
Hi-C^2 probe design:
To design probes targeting the two regions for HYbrid Capture Hi-C (Hi-C^2), we
followed a similar approach as (8). In short, we (i) identified all MboI restriction sites within the
target region, (ii) we designed our bait probe sequences to target sequences within a certain
distance of the MboI restriction sites as Hi-C ligation junctions occur between them, (iii) we
followed a similar three-pass probe design strategy sequentially increasing various parameters
like the distance of the probe from the MboI restriction site, the number of repetitive bases,
the GC content, probe density in gaps with relaxed probe design quality filters. We then
removed overlapping probes or probes with identical sequences. After all three passes, we
identified 2741 unique probes covering region 1 (chr10:26-28Mb; 1.37 probes/kb) and 1856
probes covering region 2 (chr8:70.3-71.8Mb; 1.24 probes/kb). 15bp primer sequences (unique
for each region, details in Table S2.3) were then appended to both ends of the 120bp probe
sequence to facilitate single oligo pool synthesis and subsequent amplification of region-
specific sub-pools. Probe construction and hybrid selection was then followed with sequences
specific to this study using the same strategy detailed in (8).
567
Hi-C experiments:
The Hi-C datasets used in our analyses were generated using the *in situ* Hi-C protocol
standardized by the 4DN consortia. In brief, the in situ Hi-C protocol involves crosslinking cells
with 1% formaldehyde for 10 minutes, permeabilizing them with nuclei intact, digesting the

DNA with MboI (4-cutter restriction enzyme), filling the 5'-overhangs while incorporating
biotin-14-dATP (a biotinylated nucleotide), followed by ligating the resulting blunt-end
fragments, shearing the DNA to a 400-700bp fragment size, capturing the biotinylated ligation
junctions with streptavidin beads, building an Illumina library with 10-12 rounds of PCR
amplification, and finally analyzing the resulting fragments with paired-end sequencing. The
resulting library was always shallow sequenced to 500K-4M reads to check for library build
quality looking at key statistics such as complexity, number of Hi-C contacts, inter vs.
intrachromosomal interactions, and long-range vs/ short-range intrachromosomal
interactions. Libraries that passed the quality check were either sequenced deeper and/or used
as pools for subsequent Hi-C^2 experiments.

For our genome engineering experiments, we generated 14 in situ Hi-C libraries (Table
S2.1) from GM12878 cells. We also generated 16 *in situ* Hi-C^2 libraries from various genome-
engineered GM12878 cell lines on which we performed hybrid selection. All *in situ* Hi-C libraries
generated as part of this study are detailed in Table S2.2. All the Hi-C data was processed using
the computational pipeline described in full detail in (1). Hi-C libraries were sequenced to a
depth of between 624K-333M reads (on average, 63.8M reads). Hi-C^2 libraries were sequenced
to a depth of between 6.7M-168M reads (on average, 35.8M reads). All data was initially
processed using the pipeline published in (1) and visualized on the desktop and web version of
Juicebox. We combined Hi-C and Hi-C^2 contact maps corresponding to the same genotype and
the same locus using the Juicer's mega.sh script as these are in essence "biological" replicates,
to generate higher resolution megamaps.

<u>Analysis of cross-domain interactions:</u>

We subsampled the Hi-C^2 corresponding to the R1-WT megamap (containing 46M
reads) and R1-KO (containing 56M reads) for 5M reads, 10 times to create 10 independent R1-
WT and R1-KO mini-maps. For each of these HiC maps, we used the Juicer Tools dump
command to extract the raw contact matrix. Intradomain interactions were defined as
interaction that (i) originate and terminate in domain 1, or (ii) originate and terminate in
domain 2. Interdomain interactions were defined as interactions that originate in domain 1 and
terminate in domain 2. We then calculate percentage of cross-domain interactions for each of
the mini-maps using the formula: (number of intradomain-interactions)*100 / ((number of
intradomain-interactions) + (number of interdomain-interactions)). The percentage of cross-
domain interactions were calculated for the target domain as well as a control domain. The
distribution of cross-domain interactions across the targeted domain was found to be
significantly different in the KO vs. the WT (t-Test: Two-Sample Assuming Unequal Variances,
p-value = 1.40668×10^{-16}). The distribution of cross-domain interactions across a nearby control
domain however was not found to be significantly different in the KO vs. the WT (t-Test: Two-

Sample Assuming Unequal Variances, p-value = 0.013254165). Raw simulation data and
statistics are provided in Table S2.5.
For Figure S3C, we used the Hi-C megamap corresponding to R2-WT and R2-KO to
retrieve raw interaction counts at a 100kb resolution. Percent cross-domain interactions was
calculated using the formula stated above. We calculated the enrichment of cross-domain
interactions in the LTR41-DKO w.r.t. the WT across the targeted domain as well as a nearby
control domain.
616
617 <u>DNA Methylation analysis:</u>
618   We generated a methylation metaplot representing the mean CpG methylation value
619 from WGBS data (ENCODE dataset: ENCFF835NTC) of 20bp sliding windows, centered on
620 CTCF motifs (and ±2kb around it) segmented by their origin/TE-derivation status.
621
<u>Analysis of TE Mutational Profile:</u>
1. <u>TE consensus construction</u>
For most of the TE subfamilies, we retrieved the consensus sequences from the
RepBase library (RepBase 22.02, RepeatMaskerEdition20170127) (49). However, LINE
elements are fragmented to 5' end, ORF2 and 3' end regions in RepBase library. To reconstruct
full-length LINE consensus, we identified TE fragments in human and mouse genome using
RepeatMasker and compared the standard output (.out file) with the alignment output (.align
file) from the same RepeatMasker run (50). For each LINE element in the standard output, we
summarized which 5' end, ORF2, and 3' end fragments have been used most to construct the
full-length element. Then we use EMBOSS Water local alignment algorithm to align the three
pieces together and generated the full-length LINE consensus sequences (51).
2. <u>Crossmatch alignments</u>
We ran RepeatMasker 4.0.7 on the mm9 and hg19 genomes using crossmatch as the
search engine. We then parsed the alignment file to determine the substitution rates between
the ancestral sequence and the genomic element. For each genomic element, we counted the
number of A-to-C, A-to-G, A-to-T, C-to-A, C-to-G, C-to-T, G-to-A, G-to-C, G-to-T, T-to-A, T-
to-C, and T-to-G substitutions (single nucleotide substitutions), where the first nucleotide
indicates the ancestral sequence and the second nucleotide indicates the genomic sequence.
We ignored any substitutions that involved ambiguous nucleotides. We also counted the
number of insertions and deletions. All substitution frequencies were normalized by the length
of the genomic sequence to estimate the substitution rates in each TE. Any genomic TE with a
length less than 20% of the ancestral sequence was filtered out. For each single nucleotide
substitution, we calculated the average substitution rate in two subsets of TEs (details below).
We also calculated the combined C-to-T and G-to-A substitution rate (methylation-associated
substitutions) and the combined rate of all other substitutions (non-methylation-associated

substitutions) to compare the rate of DNA methylation-induced mutations to other mutations.
The methylation substitution rate was computed by taking the average of the C-to-T and G-to-
A rates for each TE and then averaging over turnover events. The non-methylation substitution
rate was computed by taking the average of all other (ten) single nucleotide substitutions for
each TE and then averaging over turnover events.

We generated a background distribution by repeating this analysis on 1000
permutations of all genomic TEs. We first calculated the frequency of each TE subfamily in the
set of turnover events. For each permutation, we randomly selected genomic TEs (not involved
in anchoring loops) from each subfamily to reflect their frequency in turnover events. The
single nucleotide substitution rate, methylation-associated substitution rate, and non-
methylation-associated substitution rate were calculated as described above. The distribution
of all substitution rates from the permutations follow a normal distribution (KS test, P >
0.0036, Bonferroni correction alpha = 0.05 for N = 14 hypotheses, Supplemental Table S3.1).
The background distribution was then used to perform a left-tailed z-test. We did not compute
a two-tailed p-value because our null hypothesis is that the observed mutation rates are
greater than or equal to the background distribution mean. For the 12 single nucleotide
substitutions, we used Bonferroni correction to account for multiple hypotheses.

3. Needle realignments
RepeatMasker performs post-processing after running crossmatch, so coordinates and
TE subfamily assignments in the .out file do not always reflect the contents of the .align file. To
improve our estimates of mutation rates, we realigned each TE to its matched consensus
sequences. We extracted the genomic and subfamily consensus sequence using the
coordinates reported in the .out file. We then performed a global alignment using EMBOSS
Needle v6.6.0.0 using a gap open penalty of 10, a gap extension penalty of 0.5, and the
EDNAFULL scoring matrix. We used the alignment to recompute single nucleotide
substitutions for each TE and then repeated the same analysis we used for crossmatch
alignments. We did not filter out TEs with a length less than 20% of the ancestral sequence
because this filter was originally put in place to account for discrepancies between the .align
and .out files. As before, the distribution of all substitution rates from the permutations follow
a normal distribution (KS test, P > 0.0036, Bonferroni correction alpha = 0.05 for N = 14
hypotheses, Supplemental Table S3.2).

678
679 **DECLARATIONS**
680
681 **Ethics approval and consent to participate** Not applicable
682
683 **Consent for publication** Not applicable.
684

Availability of data and material The data sets generated and analyzed in this current study will be uploaded to GEO upon acceptance

Competing Interests Authors declare no competing interests.

Funding M.N.K.C. was partly supported by the Precision Medicine Pathway, Washington University; H.S.J. was partly supported by NIH grant T32 GM007067; X.Z. was partly supported by R25DA027995; T.W. is supported by R01HG007175, U01CA200060, U24ES026699, U01HG009391, U41HG010972 and American Cancer Society RSG-14-049-01-DMC.

Authors' Contributions M.N.K.C. and T.W. conceived and designed this study; M.N.K.C. analyzed the data, performed experiments, generated sequencing libraries, and wrote the manuscript with inputs from T.W.; R.Z.F., J.T.W., and X.Z. contributed text and revised the manuscript; H.S.J. contributed reagents and resources; R.Z.F performed mutation frequency simulations along with M.N.K.C.; X.Z. generated TE ancestral sequences and TE alignments; T.W. supervised the project. All authors subsequently edited and approved the final manuscript.

Acknowledgements We thank members of the Wang Lab for helpful discussions related to the project; Jessica Hoisington-López and Maria Lynn Jaeger from The Edison Family Center for Genome Sciences & Systems Biology for assistance with sequencing; Matthew Patana & Daniel Schweppe from the Siteman Flow Cytometry core for FACS expertise.

Corresponding Author Correspondence and requests for materials should be addressed to Ting Wang.

Supplementary Materials
Table S1 – S3
Supplementary Figure 1 – 9

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4