3D Chromatin Architecture Remodeling during Human Cardiomyocyte Differentiation Reveals A Novel Role Of HERV-H In Demarcating Chromatin Domains

Authors:

Yanxiao Zhang[1*], Ting Li[2*], Sebastian Preissl[1,3*], Jonathan Grinstein[2], Elie N. Farah[2], Eugin Destici[2], Ah Young Lee[1], Sora Chee[1], Yunjiang Qiu[1], Kaiyue Ma[1], Zhen Ye[1], Quan Zhu[1], Hui Huang[1], Rong Hu[1], Rongxin Fang[1], Sylvia M. Evans[2,4], Neil C. Chi[2,5,7], Bing Ren[1,3,5,6,7]

*these authors contributed equally

Affiliations:

[1]Ludwig Institute for Cancer Research, La Jolla, CA 92093, USA.

[2] Department of Medicine, Division of Cardiology, University of California, San Diego, La Jolla, California 92093, USA.

[3] Department of Cellular and Molecular Medicine, Center for Epigenomics, University of California, San Diego, School of Medicine, La Jolla, CA 92093, USA.

[4]Department of Pharmacology and Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California, San Diego, La Jolla, CA 92093, USA.

[5]School of Medicine, Institute of Genomic Medicine, University of California, San Diego, La Jolla, CA 92093, USA

[6]School of Medicine, Moores Cancer Center, University of California, San Diego, La Jolla, CA 92093, USA.

[7]Correspondence: Neil Chi (nchi@ucsd.edu) and Bing Ren (biren@ucsd.edu)

Abstract/Summary

Restructuring of chromatin architecture is an essential process for establishing cell type-specific gene regulatory programs in eukaryotic cells including cardiomyocytes[1-3]. Supporting its importance, recent studies have reported that a substantial number of mutations discovered in congenital heart disease (CHD) patients reside in genes encoding chromatin remodeling factors[4-6]; yet, how chromatin structure reorganizes to assemble gene regulatory networks crucial for controlling human cardiomyocyte development remains to be elucidated. Here, through comprehensively analyzing high-resolution genomic maps that detail the dynamic changes of chromatin architecture, chromatin accessibility and modifications, and gene expression during human pluripotent stem cell (PSC) cardiomyocyte differentiation, we reveal novel molecular insights into how human PSC chromatin architecture is iteratively remodeled to build gene regulatory networks directing cardiac lineage specification. Specifically, we uncover a new class of human PSC-specific topologically associating domain (TAD) that is created by the active transcription of the primate-specific endogenous retrotransposon HERV-H. Silencing of specific HERV-Hs during the initial stages of human PSC differentiation or by genome-editing results in the elimination of corresponding TAD boundaries and reduced transcription of genes upstream of HERV-Hs. Supporting their role in maintaining pluripotency, we discovered that deletion of specific HERV-Hs leads to more efficient human PSC cardiomyocyte differentiation. Using chromatin interaction maps from these analyses, we also assigned potential target genes to distal regulatory elements involved in cardiac differentiation. Genome-editing of enhancers harboring cardiac-disease risk loci associated with congenital and adult heart diseases further confirmed that these loci regulate predicted target genes. Our results highlight a novel role for HERV-Hs in establishing human-specific PSC chromatin architecture, delineate the dynamic gene regulatory networks during cardiomyocyte development and inform how non-coding genetic variants contribute to congenital and adult heart diseases.

MAIN TEXT

The three-dimensional organization of chromosomes enables long-range communications between enhancers and promoters that are critical for building complex gene regulatory networks in multicellular species[1,3]. In somatic cells, interphase chromosomes occupy separate nuclear spaces known as chromosome territories[7]. Each chromosome is folded into a dynamic but non-random hierarchical structure characterized by stretches of transcriptionally active, megabase-long compartments that are interspersed with stretches of transcriptionally inactive compartments[8]. These compartments can be further partitioned into topologically associating domains (TADs), which exhibit high levels of intra-domain interactions and relatively low levels of inter-domain interactions[9-11]. The boundaries of TADs are generally conserved between cell types and closely related species[9,12], and are necessary to insulate enhancer/promoter interactions[13,14]. However, recent studies have shown that chromatin architecture is reconfigured in a cell-type specific manner during stem cell differentiation and somatic cell reprogramming[12,15-18].

Chromatin remodelers and transcription factors compose a large portion of genes disrupted in CHD[4-6], one of the leading causes of birth defects affecting 1-3% of newborns[19]. Thus, we investigated how chromatin architecture is dynamically remodeled to coordinate the silencing and activation of key transcriptional programs during differentiation of human PSCs to cardiomyocytes. Utilizing a transgenic human embryonic stem cell (hESC) H9 *MYL2:H2B-GFP* ventricular cardiomyocyte reporter line, which can be used to purify ventricular cardiomyocytes[20] (**Extended Data Fig. 1a**), we generated and collected samples in biological duplicates at the following six critical time points of cardiomyocyte differentiation: hESCs (Day 0), mesodermal cells (Day 2), cardiac mesodermal cells (Day 5), cardiac progenitors (Day 7), primitive cardiomyocytes (Day 15) and ventricular cardiomyocytes (Day 80) (**Fig. 1a**). Flow cytometry confirmed that differentiation efficiency for D2 – D15 time point samples was at least ~80-90%, and *MYL2:H2B-GFP+* ventricular cardiomyocytes were sorted at day 80 (**Extended Data Fig. 1b**). *In situ* Hi-C[8] experiments were performed, and on average three billion raw read pairs and one billion unique long-range cis contacts (minimum distance 10 kb,

Extended Data Table 1) were generated for each time point. Close correlation of Hi-C

maps between biological replicates verified that these maps were highly reproducible at

multiple scales (**Extended Data Fig. 2)**. Complementing these Hi-C contact maps,

ATAC-seq, ChIP-seq and RNA-seq experiments were performed on these samples to

interrogate chromatin accessibility, locate CTCF occupancy and key histone

modifications, and profile transcriptomes, respectively (**Extended Data Fig. 3)**.

Analyses of these genome-wide datasets revealed extensive reorganization of

chromatin architecture at all structural levels during hESC differentiation, which was

also accompanied by corresponding epigenomic changes (**Fig. 1a, b, c**). Across the

entire genome, the abundance of short-range chromatin interactions decreased while

the number of long-range chromatin interactions (above 500 kilobases) increased

(**Extended Data Fig. 4a**). Identification of active (A) and inactive (B) compartments at

all stages using principal component analysis showed that 81.6% of the genome

maintained the same compartment state throughout cardiomyocyte differentiation (**Fig.**

1d). Among the 18.4% of the genome that switched compartments, a similar proportion

of the genome switched from A to B and from B to A at each stage transition (**Extended**

Data Fig. 4b), and the majority (73%) only switched once, suggesting linear nuclear

architectural changes during lineage commitment (**Fig. 1d**). Hierarchical clustering of

genomic bins that switched compartments showed stage specific dynamics (**Fig. 1d**),

which positively correlated with gene expression (**Extended Data Fig. 4c**) as reported

previously[12]. For instance, gene loci of pluripotency factors including *SOX2* moved to

the inactive compartment when hESCs differentiated into mesodermal cells, whereas

genes involved in cardiomyocyte specification such as the transcription factor *HAND2*

and the cardiomyocyte-specific calcium channel *RYR2* displayed B to A transitions at

day 5 and day 15, respectively (**Fig. 1d; Extended Data Fig. 4d**).

Consistent with recent reports describing loss of TADs during the differentiation of

embryonic stem cells[15,21], we also observed that the number of TADs decreased as

hESCs differentiated into ventricular cardiomyocytes using multiple TAD-calling

algorithms[9,13,22] (**Extended Data Fig. 5a, b, c**). Clustering of stage-specific TAD

boundaries using Directionality Index (DI) delta scores, which estimates the strength of

the TAD boundaries, further showed that the majority of stage-specific TAD boundaries

were lost or weakened during differentiation and the effect was most pronounced at

early mesoderm specification (D0-D2), and ventricular cardiomyocyte

differentiation/maturation (D15-D80) (**Fig. 1e**). To investigate the mechanism of TAD

dynamics during cardiomyocyte differentiation, we examined the sequence features of

TAD boundaries that were lost in the course of differentiation with a particular focus on

D0 hESC-specific TAD boundaries [ESC(+)] and the boundaries lost during D15-D80

ventricular cardiomyocyte transition [vCM(-)] (**Extended Data Table 2**). While CTCF

has been shown to play a critical role in defining TAD boundaries[9,13,23], we observed no

significant difference in CTCF occupancy across all stages examined at ESC(+), vCM(-)

and stable TAD boundaries (**Extended Data Fig. 5d**). Thus, we explored whether other

potential mechanisms might be involved in the formation of these TAD boundaries.

Interestingly, we discovered that eight classes of repeat elements were over-

represented specifically at ESC(+) TAD boundaries when compared to stable TAD

boundaries (p-value < 0.01 and fold change > 2), whereas no enrichment of these

repeats was observed in vCM(-) TAD boundaries (**Extended Data Fig. 6a**).

Additionally, the ESC(+) TAD boundaries were associated with hESC-specific H3K27ac

signal and mRNA transcription (**Extended Data Fig. 5e, f**), which was particularly

enriched for HERV-H transcription (**Fig. 2a and Extended Data Fig. 6b**).

HERV-H is a class of primate-specific endogenous retrotransposons. It is

transcriptionally active in both human preimplantation embryos and pluripotent stem

cells[24,25], and plays a critical role in pluripotency, stem cell maintenance and somatic

cell reprograming[26-30]. Among the ~1200 HERV-H sequences in the human genome,

only a small number are transcriptionally active in hESCs (**Fig. 2b**)[26]. We found that the

top 50 highly transcribed HERV-H loci were predominantly located at TAD boundaries in

hESCs (**Fig. 2c, Extended Data. Fig 6c**). However, transcriptional silencing of these

HERV-Hs occurred promptly by day 2 in mesodermal cells and coincided with the loss

of the corresponding TAD boundaries (**Fig. 2d, e, f and Extended Data Fig 6d, e, f**).

As controls, genes with similar transcriptional levels to these top 50 HERV-H loci did not

show any boundary activity as evidenced by the DI scores (**Extended Data Fig. 7a**). This correlation was not specific to only the H9 hESC line used in this study, as we also observed HERV-H associated TAD boundaries in other hESCs (H1)[12,31] and human induced PSCs[32] (**Extended Data Fig. 7b**). Consistent with these findings, we notably discovered that in human mesendoderm cells (an early human embryonic state that gives rise to mesoderm and endoderm cells), both the expression of these HERV-Hs and their corresponding TAD boundary strength were approximately half the levels compared to those observed in hESCs (**Extended Data Fig. 7b**).

Because HERV-Hs might contribute to TAD boundary formation by recruiting DNA binding proteins, we examined publicly available ChIP-seq data of histone modifications and transcription factors (TFs) in hESCs. We identified several modifications and factors enriched at sites for the top 50 transcribed HERV-Hs when compared to those for the lower-ranking HERV-Hs. They included POL2 transcriptional machinery (POLR2A and POLR2AphosphoS5), active histone marks (H3K4me3, H3K4me2 and H3K27ac), ubiquitously active TFs (JUND and SP1), pluripotency factor NANOG[28] and all three subunits of the cohesin complex (**Extended Data Fig. 7c**). To further investigate the role of cohesin in HERV-H-mediated TAD boundary formation, we performed ChIP-seq in hESCs for the cohesin complex subunit SMC3. Analyzing this ChIP-seq data along with RNA-seq as well as CTCF and POLR2A ChIP-seq data, revealed broad enrichment of cohesin, CTCF and POLR2A specifically at the 3' end of these highly transcribed HERV-H sequences (**Fig. 2g**). In contrast to these findings, the enrichment of these factors was not similarly observed at sites of lower ranked HERV-Hs (**Fig. 2g**), suggesting that these factors may be involved in mediating the formation of TAD boundaries by HERV-Hs. However, we cannot completely rule out that CTCF may also participate in defining these HERV-H-associated TAD boundaries as weak but reproducible enrichment of CTCF occupancy was observed in these regions despite the absence of CTCF binding motifs (**Fig. 2g**). Overall, these data support that HERV-Hs form TAD boundaries by accumulating cohesin complexes at its 3' end, and cohesin is likely positioned by the transcribing POL2 complex[33,34].

To determine whether HERV-H sequences are indeed required for TAD boundary formation, we deleted two HERV-H elements located at different hESC TAD boundaries, using CRISPR-Cas9 genome editing tools (termed as HERV-H1-KO and HERV-H2-KO, respectively). In both cases, deletion of these HERV-Hs resulted in elimination of TAD boundaries, as evidenced by the merging of the two TADs bordering each HERV-H element (**Fig. 3a**), and reduced expression of genes in the TAD domain immediately 5' upstream of the HERV-H sequence (**Fig. 3a**). These findings thus reveal that HERV-H sequences are indispensable for the TAD boundaries formed at these regions and furthermore the presence of HERV-H increases the overall transcriptional activity of the neighboring genes in the same TAD domain, supporting that HERV-Hs may serve as enhancers in hESCs[35,36]. Notably, this activating effect is only seen in the TAD domain on the 5' end of HERV-H and not the 3' end, suggesting that boundaries present on the 3' end of HERV-Hs, may insulate the effect of enhancers. Supporting this possibility, additional analyses of all genes located within 500 kb of boundary-associated HERV-Hs showed that only the genes in regions 5' of HERV-H were up-regulated in hESCs compared to differentiated cells (**Fig. 3b**).

Further examination of the transcriptional profiles of both HERV-H knockouts revealed that expression of 353 and 144 genes were significantly changed (**Extended Data Fig. 8a, b, Extended Data Table 3**). The fold changes of HERV-H1-KO and HERV-H2-KO were highly concordant (**Extended Data Fig. 8c**, Pearson r=0.5, p-value< 1e-15). Out of the 74 genes that were significantly altered in these knockouts, 69 genes (92%) were altered in the same direction with similar fold changes (**Extended Data Fig. 8c**). Of the 43 genes down-regulated in both, 10 (23%) were located within 20 kb of other HERV-H sequences (**Extended Data Fig. 8d**), including two well studied chimeric transcripts *SCGB3A2*[37] and *LINC00458*[38], which are known to be regulated by HERV-Hs (**Extended Data Fig. 8e, f**). Thus, these data showed that deletion of individual HERV-H sequences could lead to down-regulation of HERV-H related transcription at other loci. In addition to these findings, we also discovered that the human-specific long-coding RNA (lncRNA), *Heart Brake* (*HBL1*), which suppresses cardiomyocyte differentiation[39], was also significantly reduced in the HERV-H1-KO (**Extended Data**

Fig. 8f). To further explore whether this decrease in *HBL1* lncRNA in the HERV-H1-KO

promotes cardiomyocyte differentiation, we examined the cardiomyocyte differentiation

efficiency of HERV-H1-KO hESCs compared to control hESCs under suboptimal

cardiomyocyte differentiation conditions. While these conditions led to ~40%

cardiomyocyte differentiation efficiency at day 15 in control hESCs, we discovered that

HERV-H1-KO hESCs exhibited >80% cardiomyocyte differentiation efficiency (**Fig. 3c**),

which is consistent with the increased cardiomyocyte differentiation efficiency observed

in *HBL1* knockout hPSCs[39]. Overall, these findings suggest that HERV-Hs may globally

maintain pluripotency in hPSCs through a common regulatory network; however,

individual HERV-Hs may also regulate specific genes involved in the differentiation of

distinct lineages, such as *HBL1* and the cardiac lineage.

Because HERV-H sequences are primate specific, we predicted that the TAD

boundaries established by HERV-Hs would be absent in ESCs from non-primate

mammals. To test this prediction, we examined Hi-C profiles from mouse ESC and

observed no TAD boundaries within regions syntenic with the HERV-H-associated

human TAD boundaries (**Fig. 3d, e**). Next, we calculated the fraction of the human TAD

boundaries that overlapped or were within 25 kb of TAD boundaries in mouse ESCs

and discovered that HERV-H-associated TAD boundaries were less likely to be present

in mouse ESC than other types of TAD boundaries (**Extended Data Fig. 9a**). This

result is consistent with the model that HERV-H transcription is involved in forming ESC

specific TAD boundaries in the primate lineages. It is estimated that HERV-H integrated

into the genome 30–40 million years ago (MYA) at the time of divergence of Old and

New World monkeys[40,41], with the largest expansion (LTR7 and LTR7B) occurring at

about 25 MYA[40], and a second expansion (LTR7Y) occurring between 10 and 17

MYA[40]. We found that the top 50 most highly expressed HERV-Hs, which have the

potential to form TAD boundaries, were exclusively flanked by LTR7 at both ends

(**Extended Data Fig. 9b**). In addition, these LTRs belonged predominantly to a 450 bp

LTR7 subtype (**Extended Data Fig. 9c**). Finally, these TAD-boundary HERV-Hs

showed less sequence divergence between their 5’ LTRs and 3’ LTRs (**Extended Data

Fig. 9d**), suggesting that they likely were inserted more recently than other HERV-Hs

that do not form TAD boundaries, or were impacted by gene conversion or selection[42].

From our extensive interrogation of the genome during hESC cardiomyocyte

differentiation, we also identified 14,216 chromatin loops spanning a median genomic

distance of 180 kb at 10 kb resolution (**Extended Data Table 4**). More than 70% of the

loops corresponded to either promoter-to-promoter or promoter-to-enhancer interactions

(**Extended Data Fig. 10a**). Consistent with chromatin loops detected in different cell

types or model systems[13,14], the loop anchors were enriched for accessible chromatin,

histone marks, transcriptional start sites and CTCF binding sites (**Extended Data Fig.**

10b). More importantly, 23% (3,251) of these chromatin loops exhibited significant

changes in interaction strength during cardiomyocyte differentiation, which illustrates a

dramatic reconfiguration of the gene regulatory network (**Fig. 4a, Extended Data Fig.**

10c). Gene transcription and chromatin marks at anchors of stage-specific chromatin

loops were correlated to a significantly higher degree than that of anchors of static

chromatin loops and distance-matched control (**Fig. 4e**). Using K-means clustering, we

defined five distinct clusters of stage-specific chromatin loops (**Fig. 4b**), each

accompanied by corresponding epigenomic changes (**Extended Data Figure 10d**).

Remarkably, the gene sets enriched for each chromatin loop cluster recapitulated the

developmental processes from early differentiation, to heart development, and

culminated in cardiac muscle development (**Fig. 4c**). Chromatin loops gained during

cardiomyocyte differentiation included promoters of many well studied heart

transcription factors, such as *GATA6*, *TBX5*, *NKX2-5*, *TBX3* and *MEF2A*[43]. Analysis of

ATAC-seq peaks at loop anchors also revealed distinct patterns of transcription factor

binding throughout differentiation (**Fig. 4d**). For instance, enrichment for GATA motifs

emerged at day 2, and T-BOX motifs at day 5, whereas MEF2 motifs were only highly

enriched at day 80. These findings suggest a process involving pioneering factor

activity of GATA family members such as GATA4 for the initial rewiring of chromatin

loops to activate relevant early cardiac developmental gene programs, whereas MEF

family members such as MEF2C may play a role in the maturation of cardiomyocytes.

Finally, loops formed at later stages tended to span longer genomic distances, to have a

higher percentage of convergent CTCF-CTCF motifs, and were more likely to occur at

TAD boundaries (**Extended Data Fig. 10e, f, g**). These observations further support

the global trend of increased long-distance interactions during cardiomyocyte

differentiation (**Extended Data Fig. 4a**).

Integrative analysis of the chromatin architecture and ChIP-seq data revealed two major

types of loops: 1) loops enriched for the polycomb mark H3K27me3 (**Extended Data**

Fig 10d, 11a); and 2) loops enriched for H3K27ac (**Extended Data Fig 10d, 11b**).

H3K27me3-associated, polycomb dependent loop interactions, which have been

described in *Drosophila* and mouse ESC[15,44,45], were prominent in hESCs, spanned

very long genomic distances and were abruptly lost at the mesoderm stage (**Extended**

Data Fig 10d, 11a). Notably, such loops frequently involved promoters of transcription

factors, which were not expressed at the hESC stage (Fold enrichment=9.7, adjusted p-

value=1e-70, hypergeometric test), including the early cardiac transcription factor

HAND2. Further analyses of the *HAND2* locus revealed a multistep assembly of loops

formed between the *HAND2* promoter and several distal elements (**Fig. 1b**). In hESCs

and mesodermal cells, which do not express *HAND2*, the *HAND2* promoter formed an

H3K27me3 associated loop with the promoter region of the lncRNA *LINC02268*.

However, by day 5, this loop was eliminated and two new loop interactions were

established between the *HAND2* promoter and two distal regions marked by H3K27ac

and H3K4me1. This chromatin remodeling coincided with the onset of *HAND2*

expression in cardiac mesodermal cells (D5) and a subsequent 4-fold increase in

HAND2 expression in primitive cardiomyocytes (D15) and ventricular cardiomyocytes

(D80); however, lncRNA *LINC02268* remained transcriptionally repressed (**Fig. 1b, c**).

Based on this multi-loop chromatin remodeling at the *HAND2* locus, we further

examined whether multi-loop formation was a general feature to tightly control

expression levels of transcription factors whose function may impact cell fate more than

that of other genes. To this end, we calculated the degree of interaction for each loop

anchor (**Fig. 4f**), and observed that 45% of the loop anchors interacted with at least two

other anchors (**Fig. 4g**), suggesting that chromatin loops frequently form network hubs.

In addition to *HAND2*, many other well-known cardiac TFs, including *TBX5*, *GATA4*,

TBX20, *MEIS1/2*, also interacted with four or more *cis*-elements and formed similar

network hubs (**Extended Data Fig 11c, d, e,** and **Extended Data Table. 5**). Supporting

these findings, the network hubs with more chromatin interactions were more likely to

include transcription start sites (TSSs) of TFs (**Fig. 4h**), and less likely to involve TSSs

of house-keeping genes (**Fig. 4i**). Finally, those network hubs with more interactions

frequently contained more CTCF peaks (**Extended Data. Fig. 11f**), suggesting that

these interactions were likely facilitated by CTCF.

We next used the chromatin interaction maps to link non-coding genetic variants from

genome-wide association studies (GWAS) to potential target genes[46-49], and found a

strong enrichment of single nucleotide polymorphisms (SNPs) associated with cardiac

traits in the chromatin loop anchors (**Fig. 5a**). The list of predicted candidate genes for

these SNPs predominantly included genes involved in cardiac function (**Table 1** and

Extended Data Table 6) and cardiac ion channel genes (**Fig. 5b**). One notable cardiac

GWAS example was rs6781009, which was found in strong linkage disequilibrium (LD)

with lead SNPs of four cardiac electrophysiologic GWAS traits (QRS complex, QRS

duration, QT interval and electrocardiographic traits). Using our chromatin interaction

maps, we found that this SNP was located in a stage-specific D80 enhancer that

interacted with the promoter of the cardiac sodium channel gene *SCN5A* (**Extended**

Data Fig. 12a), supporting similar findings recently observed by circular chromosome

conformation capture (4C) studies in human cardiac tissue and functional studies in

mouse hearts[50]. Another cardiac GWAS example was rs17608766, the lead SNP for

six cardiac traits including aortic root size, blood pressure, pulse pressure, QRS

complex, QRS duration and systolic blood pressure. Our chromatin interaction maps

revealed that this SNP overlapped a D2 mesoderm specific enhancer that interacted

~110 kb away with the promoter of *WNT3* (**Fig. 5c**), which is known to be involved in

cardiac mesoderm induction[51,52] and thus may explain the broad cardiac traits

potentially affected by this SNP. Because this SNP is within a highly conserved

genomic region (**Fig. 5d**) and predicted to alter a binding motif for the KLF4 transcription

factor (**Extended Data Fig. 12b),** which has been shown to regulate cardiac function in

mice[53], we deleted the putative enhancer containing this SNP in hESCs using CRISPR-

mediated genome editing. This deletion consequently resulted in a ~20-50% reduction

in *WNT3* expression in mesodermal cells at day 2 (**Fig. 5e**), but had no effect on

expression of the nearest gene *GOSR2* (**Extended Data Fig. 12c**), which has

previously been annotated to rs17608766. Finally, in addition to investigating SNPs

associated with cardiac functional traits, we also examined long-range acting SNPs for

cardiac developmental traits (i.e. congenital heart disease – CHD). Interrogating

rs870142, a SNP associated with atrial septal defect[54]*,* revealed that it was located in an

active D2 mesodermal enhancer within the intron of *STX18-AS1*, which formed dynamic

long-range interactions (~200 kb) with the promoter of *MSX1* (**Fig. 5f, g**), a

developmental TF reported to interact with *TBX5* during atrial septal development[55,56].

Deletion of this enhancer containing rs870142 using CRISPR/Cas9-mediated genome

editing resulted in a 50% reduction in *MSX1* expression (**Fig. 5h**), whereas the

expression of the nearest gene *STX18* remained unchanged (**Extended Data Fig. 12d**).

Thus, these findings support that our chromatin interaction maps can be used to

discover candidate genes affected by genetic disease or trait variants located in distal

regulatory elements controlling gene expression.

Evolution of multi-cellular organisms is driven, in large part, by the invention of new cell-

type specific gene regulatory circuits responsible for the fitness traits of each species.

Our interrogation of the dynamic re-organization of 3D chromatin architecture during

cardiomyocyte differentiation reveals novel insights into not only how these gene

regulatory networks are rewired to specifically control cardiac development but also

more broadly how the primate specific retrotransposon, HERV-H, creates new human

PSC-specific TADs during primate evolution. Applying these findings to identifying

chromatin loops during cardiomyocyte differentiation moreover aided in predicting

potential target genes for cardiac GWAS variants, which were further validated by

CRISPR-mediated deletion studies in both congenital and adult heart disease/trait

examples. Although previous research suggests that HERV-Hs integrated into the

human genome during primate evolution to regulate human-specific pluripotency

through creating novel chimeric transcripts (*ESRG*[26], *linc-ROR*[30,57] and *LINC00458*[38])

and providing potential binding sites to recruit pluripotency factors (*NANOG*, *SOX2* and

POU5F1)[27], our findings show that HERV-H sequences may also create human PSC

specific TAD boundaries to build human-specific pluripotency gene regulatory networks,

thus supporting a long-standing theory that retroviruses may play an important role in

modifying gene regulatory logic during evolution[58,59]. Additionally, we provide evidence

that HERV-H transcription may achieve TAD boundary formation through accumulating

cohesin complexes at the 3' terminal of these actively transcribed retrotransposons.

Consistent with this possibility, we observed that silencing of HERV-H transcription

during differentiation or CRISPR-mediated removal of HERV-H sequences results in

loss of corresponding TAD boundaries. In addition to affecting the transcription of

genes within the TADs that are upstream of the 5' end of HERV-H*s*, this transcriptional

silencing is closely linked to hPSC differentiation, and in one particular case, deletion of

a specific HERV-H that appears to control the cardiac regulatory lncRNA *HBL1*, leads to

increased differentiation of hPSCs into cardiomyocytes. Overall, these findings not only

highlight a novel role for endogenous retroviruses in constructing species-specific PSC

chromatin architecture but also provide a genomic blueprint for understanding how gene

regulatory networks are rewired during cardiomyocyte development and disrupted by

non-coding genetic variants to lead to congenital and adult heart diseases.

Acknowledgements

We thank Samantha Kuan and Dr. Bin Li for sequencing and bioinformatics support.

We thank Eric van Nostrand for RNA extraction. This project is supported by funding

from the Ludwig Institute for Cancer Research (to B.R.), the NIH (1UM1HL128773 to

S.M.E., N.C.C., E.D. and B.R.). S.P. was supported by a postdoctoral fellowship from

the Deutsche Forschungsgemeinschaft (DFG, PR 1668/1-1). E.N.F. was supported by

NIH pre-doctoral training grant (5T32HL007444-35).

Author contributions

N.C.C. and B.R. designed and supervised the experiments, analysis, and data

interpretation. Y.Z. implemented the analysis pipeline and analyzed all sequencing

datasets, interpreted the results, designed the experiments for HERV-H and chromatin

loop functional studies. T.L. generated the CRISPR-cas9 edited cell lines for HERV-H

and chromatin loop functional studies, performed differentiation and qPCR of the

corresponding cell lines. S.P. initiated the project, performed the Hi-C and ATAC-seq

experiments, and helped with interpretation of the results. J.G. and E.N.F performed cell

culture, differentiation and collected cells for Hi-C, ChIP-seq, ATAC-seq and RNA-seq

assays. E.D. contributed to analysis and interpretation of the ChIP-seq data. ChIP-seq

experiments were performed by A.Y.L. (histone modifications), S.C. (CTCF), Q.Z. and

H.H. (SMC3). Y.Q. and R.F. helped with the analysis of Hi-C datasets. K.M. helped with

the genome-editing experiments. Z.Y. performed the RNA-seq experiments. R.H.

performed the Hi-C experiments for HERV-H KO cell lines. S.M.E. helped with

interpretation of the results. Y.Z., T.L., S.P., N.C.C., and B.R. wrote the manuscript with

input from all authors.

Author Information

Sequencing data from this study have been deposited in the Gene Expression Omnibus

under the accession number GSE116862. Reprints and permissions information is

available at www.nature.com/reprints. The authors declare no competing financial

interests. Readers are welcome to comment on the online version of the paper.

Correspondence and requests for materials should be addressed to N.C.C.

(nchi@ucsd.edu) and B.R. (biren@ucsd.edu).

Figure Legends

Figure 1. Global chromatin architecture reorganization during human ventricular cardiomyocyte differentiation. (**a & b**) Hi-C contact matrices for each stage of cardiomyocyte differentiation at mega-base resolution and at higher resolution at the *HAND2* locus with corresponding epigenomic profiles. (**c**) Schematic illustration of the multi-step decommissioning of inhibitory loop and assembly of subsequent activating loops anchored at *HAND2* promoter. (**d**) A heatmap showing hierarchical clustering of dynamic chromatin compartments during cardiomyocyte differentiation. The pseudo color reflects the PC1 values (compartment A/B) of compartment bins. Negative PC1 value stands for compartment B and positive for compartment A. Representative genes located in corresponding compartment bins are annotated to the right of the heatmap. (**e**) A heatmap showing the DI delta scores for the stage-specific TAD boundaries, ordered by the presence of TADs at six stages. vCM(-) stands for TAD boundaries lost in purified ventricular cardiomyocytes at D80, ESC(+) for hESC-specific TAD boundaries.

Figure 2. Transcriptionally active HERV-H forms human ESC-specific TAD boundaries. (**a**) Aggregate RNA-seq expression profile (RPKM normalized) at ESC(+) TAD boundaries that overlap HERV-H element. (**b**) Scatterplot shows the expression levels (RPKM) across different HERV-H loci (ordered by expression levels from high to low in ESC) at D0, D2 and D5 stages of differentiation. (**c**) Heatmap of aggregate Hi-C contact matrix [log2(observed/expected)] within 200 kb of the top 50, 51-100 and 101-150 ranked HERV-Hs, at D0. (**d**) Heatmap of the aggregated Hi-C matrix [log2(observed/expected)] within 200 kb of the top 50 HERV-Hs, at D0, D2 and D5. (**e,f**) Representative Hi-C interaction matrices of two HERV-H loci located at ESC(+) TAD boundaries at D0, D2 and D5 (top) are shown as heatmaps along with genome browser tracks of POLR2A, SMC3, CTCF, H3K27ac ChIP-seq and RNA-seq data of the expanded genomic region containing the TAD boundary (arrow). (**g**) Aggregated genomic profiles of RNA-seq, POLR2A, SMC3 and CTCF ChIP-seq around top 50

HERV-Hs located on the ESC(+) TAD boundaries (red) and lower ranked HERV-Hs

(grey).

Figure 3. Deletion of two HERV-H sequences leads to merging of TADs and

dysregulation of gene expression programs in hESC. (**a**) Hi-C interaction matrices

of the wild-type (WT) and mutant hESC lines (HERV-H1-KO and HERV-H2-KO*)* are

shown, along with DI scores, and fold changes of gene expression at the HERV-H1 and

HERV-H2 loci. The loss of TAD boundary in the mutant cells is accompanied with

decrease of RNA expression 5' terminus to the HERV-H sequences. (**b**) Boxplots show

expression levels (RPKMs) of genes whose TSSs are located from -500 kb to the 5'

LTR and from 3' LTR to +500 kb of HERV-Hs of boundary-associated HERV-Hs. (**c**)

Line chart (mean ± standard error, N=3 cardiomyocyte differentiations) shows

percentage of TNNT2 positive cells during cardiomyocyte differentiation of WT and

HERV-H1-KO hESC lines (two HERV-H1-KO clones analyzed). HERV-H1-KO hESCs

display increased cardiomyocyte differentiation efficiency compared to WT hESCs. (**d**)

Hi-C contact matrix and DI scores of the mouse syntenic loci to the human loci

examined in (**a**). The arrows indicate the boundaries observed in human ESCs. (**e**)

Heatmap of the aggregated Hi-C contact matrix [log2(observed/expected)] of the lifted-

over hESC TAD boundaries associated with HERV-H loci.

Figure 4. Dynamics of chromatin loops during human cardiomyocyte

differentiation. (**a**) Bar-chart shows the percentages of static or dynamic chromatin

loops during cardiomyocyte differentiation. (**b**) Heatmap shows K-means clustering of

the stage-specific chromatin loops. Pseudo color reflects the normalized contact

frequencies between the loop anchors for each stage-specific loop. (**c**) Enriched Gene

Ontology (GO) terms for genes located at the loop anchors of each cluster in (**b**) along

with their adjusted p-values. (**d**) Over-represented transcription factor binding motifs for

distal open chromatin regions located at the loop anchors of each cluster in (**b**) and their

p-values. CTCF motif enrichment was not shown but observed in all clusters. (**e**)

Boxplots of Pearson correlation coefficients of histone marks of distal enhancers and

expression levels of their target genes predicted based on the chromatin loops.

Distance-matched controls are "mirror" loops that connect to genomic regions from the

opposite direction with the same distances. P-values are from Wilcoxon Rank Sum test.

(**f**) Diagram illustrates how the degree of chromatin interactions for each loop anchor is

determined. (**g**) Histogram of degrees of network connections for all loop anchors. (**h**)

Fraction of chromatin interaction network hubs containing at least one transcription start

site (TSS) of a transcription factor (TF) plotted against the degree of connectivity. (**i**)

Fraction of network hubs containing at least one TSS of a house-keeping gene (HKG)

plotted against the degree of connectivity.

Figure 5. Identification of target gene for heart-related GWAS SNPs using

chromatin loops detected during cardiomyocyte differentiation. (**a**) Bar graph

shows the enrichment of GWAS SNPs associated with cardiac traits at the chromatin

loop anchors detected during the cardiomyocyte differentiation. (**b**) Enriched GO terms

for the genes predicted to be linked to SNPs from cardiac GWAS studies along with P

values. Red dashed line indicates P value cutoff of 0.05. (**c**) Hi-C contact matrices, and

genome browser tracks of various histone modification ChIP-seq and RNA-seq data at

the *WNT3* locus. A cardiac GWAS SNP rs17608766 is marked by a red line. (**d**)

Expanded view of the genomic region containing rs17608766. The putative enhancer

containing rs17608766, which located in an open chromatin region flanked by H3K27ac

and H3K4me1 peaks, is targeted for CRISPR/Cas9-mediated deletion. (**e**) Bar charts

show the normalized expression levels of *WNT3* in three CRISPR genome edited

clones at D2 (mean ± standard error, N =5). Deleting the putative enhancer containing

rs17608766 specifically reduces *WNT3* expression. P-values are from one-sided paired

t-test. (**f**) Hi-C contact matrices, epigenomic profiles at *MSX1* locus. A cardiac GWAS

SNP rs870142 is marked by red line. (**g**) Expanded view of the genomic region

containing rs870142. CRISPR mediated deletion of the putative enhancer is marked.

(**h**) Bar charts show that deleting the putative enhancer containing rs870142 specifically

reduces *MSX1* expression in three independent clones at D2 (mean ± standard error, N

= 9). P-values are from one-sided paired t-test.

Table 1. List of target genes assigned to cardiac GWAS SNPs with chromatin

loops. The first three columns show the ATAC-seq peaks that the SNPs overlapped.

Last column "tier" was based on our knowledge of the gene's known or likely

importance in cardiac differentiation and function.

Extended Data Figure Legends.

Extended Data Figure 1. Imaging and Flow cytometry data for cardiomyocyte differentiation. (**a**) Immunostaining for MYL2 (white) shows exclusive expression in *MYL2*-H2B-GFP + hESC-derived cardiomyocytes after 80 days of cardiac differentiation. *MYL2*-H2B-GFP + cells were positive for cardiac troponin T (cTNT) (red), but not all cTnT+ cells were MYL2+ cardiomyocytes. DNA was stained with DAPI (blue). Images are representative of a minimum of three independent experiments. (**b**) Flow cytometric quantification of distinct cell-surface markers for D2-D15 time points: T (D2), KDR/PDGFRα (D5), TNNT2 (D15). H2B-GFP + ventricular cardiomyocytes were sorted at day 80. Numbers represent the respective percentage of cells.

Extended Data Figure2. The Hi-C data are highly reproducible. (a) Smoothed scatterplot of the Hi-C contacts between biological replicates for each stage. bin size = 100 kb. (**b**) Hierarchical clustering of Hi-C contact matrices based on 1-PCC (Pearson correlation coefficients) of the contacts between all samples, bin size = 100 kb. (**c**) Hierarchical clustering of the compartment A/B scores (PC1 values) between samples. (**d**) Hierarchical clustering of the insulation scores between samples.

Extended Data Figure 3. Quality metrics of ChIP-seq, RNA-seq and ATAC-seq data used in the present study. (**a**) Boxplot showing the number of reads for each sample. (**b**) Boxplot showing the percentage of alignments to human genome for each sample. (**c**) Boxplot showing the percentage of potential duplicated reads for each sample. (**d**) Boxplot showing the number of peaks called for each ChIP-seq and ATAC-seq sample. (**e**) Barplots showing the relative expression levels (by RNA-seq) of representative genes during cardiomyocyte differentiation.

Extended Data Figure 4. Global changes in chromatin organization during

cardiomyocyte differentiation. (**a**) Probability of Hi-C interaction versus interacting

distance (log2) during cardiomyocyte differentiation. (**b**) Barplot shows the percentages

of compartment switches for each stage transition. (**c**) Histogram of the Pearson

correlation coefficients between gene expression and compartment A/B (PC1) value. (**d**)

Genome browser view of *SOX2*, *HAND2* and *RYR2* loci, showing PC1 (Blue/red), RNA-

seq (Black) and H3K27ac (Blue) signals.

Extended Data Figure 5. Dynamics of TADs during cardiomyocyte differentiation.

(**a**) Number of TADs as defined by a domain caller algorithm first reported by Dixon et

al. 2012[9]. (**b**) Number of TADs (or loop domains) found by Arrowhead algorithm[13]. (**c**)

The number of TADs found by Insulation score[22]. (**d**) Fraction of TAD boundaries that

contain CTCF ChIP-seq peaks at each stage of cardiomyocyte differentiation. "stable"

group stands for TADs that are present at all stages. (**e**) Boxplot showing the dynamics

of H3K27ac signal at ESC(+) TADs. (**f**) Boxplot showing the dynamics of RNA-seq

signal at ESC(+) TADs.

Extended Data Figure 6. HERV-H elements are enriched at ESC(+) TAD

boundaries. (a) Scatter plot shows the fold enrichment and -log10(p-values) for various

repeat element classes at ESC(+) and vCM(-) TAD boundaries relative to the rest of the

TAD boundaries. P-values are from two-sided proportion test. (**b**) Aggregate RNA-seq

expression profile (RPKM normalized) at ESC(+) TAD boundaries that overlap each of

the eight repeat elements (HERVH-int and LTR7 are combined as they both belong to

HERV-H). (**c**) Heatmap of DI scores within 70 kb of the top 50 most highly-expressed

HERV-Hs. (**d,e,f**) Hi-C heatmap, DI, and epigenomic profiles for three ESC(+) TAD

boundaries harboring HERV-H/LTR7 sequences.

Extended Data Figure 7. Further characterization of the HERV-H-associated TAD

boundaries. (**a**) DI score profiles around the TSSs of genes expressed at similar levels

to the top 50 transcribed HERV-Hs. (**b**) Aggregated DI score profiles of the TAD-

boundary associated with HERV-Hs in multiple H1 ESC derived lineages and iPSCs.

ESC: embryonic stem cell; MES: mesendoderm. MSC: mesenchymal stem cell; NPC:

neural stem cell; TRO: trophoblast-like cell; iPSC: induced pluripotent stem cell. (**c**)

Barplot shows the ChIP-seq signal fold enrichment of the top 50 HERV-Hs comparing to

HERV-Hs ranking 51-300.

Extended Data Figure 8. Changes in gene expression in HERV-H knockout

hESCs. (**a**) MA plot showing average gene expression levels and fold changes of each

gene in HERV-H1-KO and wild-type (WT). (**b**) Same as (**a**) but for HERV-H2-KO. (**c**)

Scatterplot shows the changes in gene expression in HERV-H1-KO and HERV-H2-KO

cells over WT cells. The red dots are genes significantly changed in both mutant cell

lines. The numbers at the corners show the number of significantly changed genes in

each Quadrant. (**d**) Barplot showing the number of significantly changed genes located

within 20 kb of the HERV-H sequences. Genes down-regulated in both HERV-H

knockouts were more likely to be within 20 kb of HERV-H sequences. (**e**, **f**) RNA-seq

profile of wild-type (*WT*) and HERV-H1-KO and HERV-H2-KO lines at the *SCGB3A2*

and *LINC00458/HBL1* gene loci.

Extended Data Figure 9. Evolutionary analysis of HERV-H sequences and

associated TAD boundaries. (**a**) Comparison between hESC and mESC TAD

boundaries reveals that a lower portion of HERV-H associated boundaries is present in

mESCs. P-value of the difference between groups was computed using two-sided

proportion test. (**b**) Bar graph shows the percentage of HERV-Hs flanked by various

types of LTRs. HERV-Hs are ranked by their expression levels in the hESCs, and

grouped by bins of 50. (**c**) Violin plot shows the length of the flanking LTRs. HERV-Hs

ranked and binned as described in (**b**). (**d**) Boxplot shows the sequence divergence of

the 5' LTR and 3' LTR for each bin of HERV-Hs. HERV-Hs ranked and binned same as

(**b**).

Extended Data Figure 10. Relationships between chromatin loops and histone

modification during cardiomyocyte differentiation. (**a**) Barplot showing fraction of

chromatin loops in each category of interactions. E-E: enhancer to enhancer; E-P:

enhancer to promoter; P-P: promoter to promoter. (**b**) Barplot showing fraction of

chromatin loop anchors or genome-wide background containing each of the epigenomic

element. (**c**) Representative data for Hi-C contact matrices of stage-specific and static

chromatin loop interactions (**d**) Heatmap shows the clustering of stage-specific

chromatin loops and histone modification states at the corresponding loop anchors. (**e**)

Violin plot of the genomic distances for each class of chromatin loops in log scale.

Symbol S: static loop. 1-5: Loop clusters 1-5 in **Fig.4b** and **Ext. Fig. 10d**. (**f**) Barplot for

fraction of loops harboring different type of CTCF orientations. No-CTCF: at least one

anchor does not contain CTCF peak; unknown: at least one anchor does not contain a

matched CTCF motif; incorrect: CTCF motifs at both anchors do not face inward;

inward: CTCF motifs at both anchors face inward. (**g**) Barplot for fraction of loops in

each category related to TAD boundaries. Inter-TAD: loop is outside of all TADs; TAD-

corner: loop is within 20 kb of any TAD corner; intra-TAD: loop is inside of TAD.

Extended Data Figure 11. Chromatin interaction hubs around genes encoding

cardiac transcription factors. (**a**) Example of a H3K27me3-associated inhibitory loop

connecting *MSX1* and *HMX1* promoters. (**b**) Example of several stage-specific and

H3K27ac associated activating loops at *HAND1* locus. (**c,d,e**) Hi-C contact matrices

(top), loop arcs (middle), and epigenomic profiles (bottom) for *GATA4* (**c**), *MEIS1* (**d**)

and *TBX5* (**e**). (**f**) Fraction of network hubs containing CTCF peaks plotted against the

degree of connectivity as illustrated by the number of CTCF peaks at each anchor

connectivity.

Extended Data Figure 12. Validation of target gene predictions for heart-related

GWAS SNPs. (**a**) Hi-C contact matrices, epigenomic profiles (ATAC-seq, H3K4me1,

H3K27ac and RNA-seq) and cardiac GWAS SNP rs6781009 at *SCN5A* locus. (**b**)

rs17608766 is predicted to alter KLF4 motif in the putative enhancer. (**c**) Normalized

expression values of *GOSR2* in three CRISPR-genome edited clones shows that

deleting the putative enhancer containing rs17608766 does not change *GOSR2*

expression (mean ± standard error, N=5). (**d**) Normalized expression values of *STX18*

in three CRISPR-genome edited clones shows that deleting the putative enhancer

containing rs870142 does not change *STX18* expression (mean ± standard error, N=7).

Extended Data Table 1. Summary statistics for Hi-C data

Extended Data Table 2. List of stage-specific TAD boundaries.

Extended Data Table 3. List of differentially expressed genes in HERV-H1-KO and

HERV-H2-KO.

Extended Data Table 4. List of chromatin loops.

Extended Data Table 5. List of genes located on the network hubs.

Extended Data Table 6. List of additional target genes assigned to cardiac GWAS

SNPs with chromatin loops.

Extended Data Table 7. List of antibodies used for ChIP-seq.

Extended Data Table 8. List of primers and cell lines used in this study.

Supplementary Materials:

Materials and Methods

Figures S1-S12

Tables S1-S7

References

1   Dekker, J. & Mirny, L. The 3D Genome as Moderator of Chromosomal Communication. *Cell* **164**, 1110-
1121, doi:10.1016/j.cell.2016.02.007 (2016).
2   Dixon, J. R., Gorkin, D. U. & Ren, B. Chromatin Domains: The Unit of Chromosome Organization. *Mol*
Cell **62**, 668-680, doi:10.1016/j.molcel.2016.05.018 (2016).
3   Yu, M. & Ren, B. The Three-Dimensional Organization of Mammalian Genomes. *Annu Rev Cell Dev Biol*
33, 265-289, doi:10.1146/annurev-cellbio-100616-060531 (2017).
4   Zaidi, S. *et al.* De novo mutations in histone-modifying genes in congenital heart disease. *Nature* **498**, 220-
223, doi:10.1038/nature12141 (2013).
5   Homsy, J. *et al.* De novo mutations in congenital heart disease with neurodevelopmental and other
congenital anomalies. *Science* **350**, 1262-1266, doi:10.1126/science.aac9396 (2015).

667	6	Jin, S. C. *et al.* Contribution of rare inherited and de novo variants in 2,871 congenital heart disease
668		probands. *Nat Genet* **49**, 1593-1601, doi:10.1038/ng.3970 (2017).
669	7	Meaburn, K. J. & Misteli, T. Cell biology: chromosome territories. *Nature* **445**, 379-781,
670		doi:10.1038/445379a (2007).
671	8	Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range interactions reveals folding principles of
672		the human genome. *Science* **326**, 289-293, doi:10.1126/science.1181369 (2009).
673	9	Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin
674		interactions. *Nature* **485**, 376-380, doi:10.1038/nature11082 (2012).
675	10	Sexton, T. *et al.* Three-Dimensional Folding and Functional Organization Principles of the Drosophila
676		Genome. *Cell* **148**, 458-472, doi:10.1016/j.cell.2012.01.010 (2012).
677	11	Nora, E. P. *et al.* Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature* **485**,
678		381-385, doi:10.1038/nature11049 (2012).
679	12	Dixon, J. R. *et al.* Chromatin architecture reorganization during stem cell differentiation. *Nature* **518**, 331-
680		336, doi:10.1038/nature14222 (2015).
681	13	Rao, S. S. *et al.* A 3D map of the human genome at kilobase resolution reveals principles of chromatin
682		looping. *Cell* **159**, 1665-1680, doi:10.1016/j.cell.2014.11.021 (2014).
683	14	Jin, F. *et al.* A high-resolution map of the three-dimensional chromatin interactome in human cells. *Nature*
684		**503**, 290-294, doi:10.1038/nature12644 (2013).
685	15	Bonev, B. *et al.* Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell* **171**, 557-572
686		e524, doi:10.1016/j.cell.2017.09.043 (2017).
687	16	Stadhouders, R. *et al.* Transcription factors orchestrate dynamic interplay between genome topology and
688		gene regulation during cell reprogramming. *Nat Genet* **50**, 238-249, doi:10.1038/s41588-017-0030-7
689		(2018).
690	17	Krijger, P. H. *et al.* Cell-of-Origin-Specific 3D Genome Structure Acquired during Somatic Cell
691		Reprogramming. *Cell Stem Cell* **18**, 597-610, doi:10.1016/j.stem.2016.01.007 (2016).
692	18	Beagan, J. A. *et al.* Local Genome Topology Can Exhibit an Incompletely Rewired 3D-Folding State
693		during Somatic Cell Reprogramming. *Cell Stem Cell* **18**, 611-624, doi:10.1016/j.stem.2016.04.004 (2016).
694	19	Gelb, B. D. Genetic basis of congenital heart disease. *Curr Opin Cardiol* **19**, 110-115 (2004).
695	20	Veevers, J. *et al.* Cell-Surface Marker Signature for Enrichment of Ventricular Cardiomyocytes Derived
696		from Human Embryonic Stem Cells. *Stem cell reports*, doi:10.1016/j.stemcr.2018.07.007 (2018).
697	21	Fields, P. A. *et al.* Dynamic reorganization of nuclear architecture during human cardiogenesis. *bioRxiv*
698		(2017).
699	22	Crane, E. *et al.* Condensin-driven remodelling of X chromosome topology during dosage compensation.
700		*Nature* **523**, 240-244, doi:10.1038/nature14450 (2015).
701	23	Nora, E. P. *et al.* Targeted Degradation of CTCF Decouples Local Insulation of Chromosome Domains
702		from Genomic Compartmentalization. *Cell* **169**, 930-944 e922, doi:10.1016/j.cell.2017.05.004 (2017).
703	24	Grow, E. J. *et al.* Intrinsic retroviral reactivation in human preimplantation embryos and pluripotent cells.
704		*Nature* **522**, 221-225, doi:10.1038/nature14308 (2015).
705	25	Goke, J. *et al.* Dynamic transcription of distinct classes of endogenous retroviral elements marks specific
706		populations of early human embryonic cells. *Cell Stem Cell* **16**, 135-141, doi:10.1016/j.stem.2015.01.005
707		(2015).
708	26	Lu, X. *et al.* The retrovirus HERVH is a long noncoding RNA required for human embryonic stem cell
709		identity. *Nat Struct Mol Biol* **21**, 423-425, doi:10.1038/nsmb.2799 (2014).
710	27	Romer, C., Singh, M., Hurst, L. D. & Izsvak, Z. How to tame an endogenous retrovirus: HERVH and the
711		evolution of human pluripotency. *Curr Opin Virol* **25**, 49-58, doi:10.1016/j.coviro.2017.07.001 (2017).
712	28	Santoni, F. A., Guerra, J. & Luban, J. HERV-H RNA is abundant in human embryonic stem cells and a
713		precise marker for pluripotency. *Retrovirology* **9**, 111, doi:10.1186/1742-4690-9-111 (2012).
714	29	Ohnuki, M. *et al.* Dynamic regulation of human endogenous retroviruses mediates factor-induced
715		reprogramming and differentiation potential. *Proc Natl Acad Sci U S A* **111**, 12426-12431,
716		doi:10.1073/pnas.1413299111 (2014).
717	30	Loewer, S. *et al.* Large intergenic non-coding RNA-RoR modulates reprogramming of human induced
718		pluripotent stem cells. *Nat Genet* **42**, 1113-1117, doi:10.1038/ng.710 (2010).
719	31	Xie, W. *et al.* Epigenomic analysis of multilineage differentiation of human embryonic stem cells. *Cell*
720		**153**, 1134-1148, doi:10.1016/j.cell.2013.04.022 (2013).
721	32	Greenwald, W. W. *et al.* Integration of phased Hi-C and molecular phenotype data to study genetic and
722		epigenetic effects on chromatin looping. *bioRxiv* (2018).

33   Lengronne, A. *et al.* Cohesin relocation from sites of chromosomal loading to places of convergent
transcription. *Nature* **430**, 573-578, doi:10.1038/nature02742 (2004).
34   Busslinger, G. A. *et al.* Cohesin is positioned in mammalian genomes by transcription, CTCF and Wapl.
Nature **544**, 503-507, doi:10.1038/nature22063 (2017).
35   Jacques, P. E., Jeyakani, J. & Bourque, G. The majority of primate-specific regulatory sequences are
derived from transposable elements. *PLoS Genet* **9**, e1003504, doi:10.1371/journal.pgen.1003504 (2013).
36   Barakat, T. S. *et al.* Functional Dissection of the Enhancer Repertoire in Human Embryonic Stem Cells.
Cell Stem Cell **23**, 276-288 e278, doi:10.1016/j.stem.2018.06.014 (2018).
37   Wang, J. *et al.* Primate-specific endogenous retrovirus-driven transcription defines naive-like stem cells.
Nature **516**, 405-409, doi:10.1038/nature13804 (2014).
38   Ng, S. Y., Johnson, R. & Stanton, L. W. Human long non-coding RNAs promote pluripotency and neuronal
differentiation by association with chromatin modifiers and transcription factors. *EMBO J* **31**, 522-533,
doi:10.1038/emboj.2011.459 (2012).
39   Liu, J., Li, Y., Lin, B., Sheng, Y. & Yang, L. HBL1 Is a Human Long Noncoding RNA that Modulates
Cardiomyocyte Development from Pluripotent Stem Cells by Counteracting MIR1. *Dev Cell* **42**, 333-348
e335, doi:10.1016/j.devcel.2017.07.023 (2017).
40   Goodchild, N. L., Wilkinson, D. A. & Mager, D. L. Recent evolutionary expansion of a subfamily of
RTVL-H human endogenous retrovirus-like elements. *Virology* **196**, 778-788, doi:10.1006/viro.1993.1535
(1993).
41   Mager, D. L. & Freeman, J. D. HERV-H endogenous retroviruses: presence in the New World branch but
amplification in the Old World primate lineage. *Virology* **213**, 395-404, doi:10.1006/viro.1995.0012
(1995).
42   Kijima, T. E. & Innan, H. On the estimation of the insertion time of LTR retrotransposable elements. *Mol
Biol Evol* **27**, 896-904, doi:10.1093/molbev/msp295 (2010).
43   McCulley, D. J. & Black, B. L. Transcription factor pathways and congenital heart disease. *Curr Top Dev
Biol* **100**, 253-277, doi:10.1016/B978-0-12-387786-4.00008-7 (2012).
44   Bantignies, F. *et al.* Polycomb-dependent regulatory contacts between distant Hox loci in Drosophila. *Cell*
144, 214-226, doi:10.1016/j.cell.2010.12.026 (2011).
45   Schoenfelder, S. *et al.* Polycomb repressive complex PRC1 spatially constrains the mouse embryonic stem
cell genome. *Nat Genet* **47**, 1179-1186, doi:10.1038/ng.3393 (2015).
46   Javierre, B. M. *et al.* Lineage-Specific Genome Architecture Links Enhancers and Non-coding Disease
Variants to Target Gene Promoters. *Cell* **167**, 1369-1384 e1319, doi:10.1016/j.cell.2016.09.037 (2016).
47   Won, H. *et al.* Chromosome conformation elucidates regulatory relationships in developing human brain.
Nature **538**, 523-527, doi:10.1038/nature19847 (2016).
48   Montefiori, L. E. *et al.* A promoter interaction map for cardiovascular disease genetics. *Elife* **7**,
doi:10.7554/eLife.35788 (2018).
49   Choy, M. K. *et al.* Promoter interactome of human embryonic stem cell-derived cardiomyocytes connects
GWAS regions to cardiac gene networks. *Nat Commun* **9**, 2526, doi:10.1038/s41467-018-04931-0 (2018).
50   van der Harst, P. *et al.* 52 Genetic Loci Influencing Myocardial Mass. *J Am Coll Cardiol* **68**, 1435-1448,
doi:10.1016/j.jacc.2016.07.729 (2016).
51   Pfeiffer, M. J. *et al.* Cardiogenic programming of human pluripotent stem cells by dose-controlled
activation of EOMES. *Nat Commun* **9**, 440, doi:10.1038/s41467-017-02812-6 (2018).
52   Cohen, E. D., Tian, Y. & Morrisey, E. E. Wnt signaling: an essential regulator of cardiovascular
differentiation, morphogenesis and progenitor self-renewal. *Development* **135**, 789-798,
doi:10.1242/dev.016865 (2008).
53   Liao, X. *et al.* Kruppel-like factor 4 is critical for transcriptional control of cardiac mitochondrial
homeostasis. *J Clin Invest* **125**, 3461-3476, doi:10.1172/JCI79964 (2015).
54   Cordell, H. J. *et al.* Genome-wide association study of multiple congenital heart disease phenotypes
identifies a susceptibility locus for atrial septal defect at chromosome 4p16. *Nat Genet* **45**, 822-824,
doi:10.1038/ng.2637 (2013).
55   Boogerd, K. J. *et al.* Msx1 and Msx2 are functional interacting partners of T-box factors in the regulation
of Connexin43. *Cardiovasc Res* **78**, 485-493, doi:10.1093/cvr/cvn049 (2008).
56   Li, Q. Y. *et al.* Holt-Oram syndrome is caused by mutations in TBX5, a member of the Brachyury (T) gene
family. *Nat Genet* **15**, 21-29, doi:10.1038/ng0197-21 (1997).
57   Wang, Y. *et al.* Endogenous miRNA sponge lincRNA-RoR regulates Oct4, Nanog, and Sox2 in human
embryonic stem cell self-renewal. *Dev Cell* **25**, 69-80, doi:10.1016/j.devcel.2013.03.002 (2013).

58   Britten, R. J. & Davidson, E. H. Repetitive and non-repetitive DNA sequences and a speculation on the
780   origins of evolutionary novelty. *Q Rev Biol* **46**, 111-138 (1971).
781   59   King, M. C. & Wilson, A. C. Evolution at two levels in humans and chimpanzees. *Science* **188**, 107-116
782   (1975).
783

Materials and Methods

Table of Contents

1 Experimental approaches

1.1 Cell culture, differentiation, and collections for large-scale cardiac molecular studies.

For the large-scale genomic and transcriptomic human embryonic stem cell (hESC) cardiac studies (i.e. ATAC-seq, ChIP-seq, RNA-seq and HiC experiments), an engineered H9 hESC *MLC2v:H2B-GFP* reporter transgenic line, which specifically expresses H2B-GFP in differentiated ventricular cardiomyocytes (CMs), was used[1]. This H9 hESC *MLC2v:H2B-GFP* line was differentiated into cardiomyocytes and their intermediate developmental stages utilizing a well-established Wnt-based cardiomyocyte differentiation protocol[2] that we have previously used to efficiently create hESC-CMs[3]. Specifically, the following developmental cardiac cell populations were generated and collected: Day (D)0 (hESC), D2 (mesoderm), D5 (cardiac mesoderm), D7 (cardiac progenitors), D15 (primitive CMs) and D80 (ventricular CMs). Differentiation efficiency was assessed at specific developmental cardiac stages by flow cytometry using antibodies against POU5F1 (Cell Signaling C30A3) - D0/hESC, BRACHYURY/T (R&D Systems AF2085) - D2/mesoderm, KDR/PDGFRA (R&D Systems AF357P and AF1264A, respectively) - D5/cardiac mesoderm and D7/cardiac progenitors, and cTNNT (ThermoFisher MA5-12960) - D7/cardiac progenitors, D15/primitive CMs, D80/ventricular CMs. Differentiation success for collected D2-D7 samples was further confirmed by re-plating a portion of the cells dissociated for collection and then maintaining them until ~D15 when CM differentiation could be assessed using TNNT2 flow analysis. Collected samples that resulted in high differentiation efficiency (>90%) were then further processed and analyzed for molecular studies (see below). For D80 ventricular CMs, we additionally sorted and purified these cells based on *MLC2v*:H2B-GFP prior to their processing and analyses (**Extended Data Fig.1**). For RNA-seq studies, ~2M dissociated cells were washed in PBS, lysed in Trizol and then stored at -80°C until further processing. For ChIP-seq studies, dissociated cells were washed in PBS, fixed in 1% formaldehyde for 10 min, quenched with 0.125 M Glycine for 5 min at room temperature (RT), then 10 min on ice and finally washed twice with PBS. These fixed cell pellets were then snap-frozen and stored at -80°C until further processing. For ATAC-seq, dissociated cells were washed with PBS and cryopreserved in culture medium + 10% DMSO at -80°C overnight in Mr.

1 Frosty freezing containers (Thermo Scientific) and then stored in liquid N[2] until further

2 processing as previously described[4].

1.2 In-situ Hi-C

In situ HiC was performed as described previously with minor modifications[5]. Cells were

harvested and pelleted by centrifugation at 300 x g for 5 min. Cells were resuspended in

fresh media at 1 million cells/mL and fixed with 2% formaldehyde for 10min at RT. The

reaction was quenched by incubating with 200 mM glycine for 5 min at RT before

transferring onto ice. Cells were pelleted (300 x g for 5 min at 4°C) and subsequently

washed with ice cold PBS. Following supernatant removal, the pellet was flash frozen in

liquid nitrogen and stored at -80°C. Nuclei were isolated using lysis buffer (10mM Tris-

HCl (pH 8.0), 10mM NaCl, 0.2% Igepal CA630) and incubation on ice for 15 min.

Pelleted nuclei were permeabilized using 0.5% SDS and incubation for 10 min at 62°C.

SDS was quenched by adding Triton-X100. Chromatin was digested using the four base

cutter restriction enzyme MboI (200 units, NEB) at 37 °C overnight. Digestion efficiency

was checked by gel electrophoresis. After inactivation of MboI for 20 min at 62°C, single

strand overhangs were filled with biotinylated-14-ATP (Life Technologies) using Klenow

DNA polymerase (40 units, NEB) for 90 min at 37°C. Next, DNA was ligated for 4 h at

16 °C using T4 ligase (2000 units, NEB). Reverse crosslinking and protein degradation

was performed using proteinase K (NEB) and incubation at 55 °C for 30 min. Finally,

NaCl (500mM) was added and the reaction was completed at 68 °C overnight. DNA

was purified using ethanol precipitation and sonicated using an ultra sonicator (Covaris

M220; duty cycle: 10; intensity: 4; cycles/burst: 220; duration: 55s) to 300-700 bp small

fragments. After size selection and purification using SPRI beads (Beckman Coulter),

biotinylated DNA was isolated using Dynabeads MyOne T1 Streptavidin beads (Life

Technologies). Sequencing libraries were prepared on magnetic beads and final PCR

amplification was carried out for 6 cycles. Size distribution of bead purified libraries was

checked using a High Sensitivity D1000 ScreenTape on a Tapestation (Agilent) and

quantified using Qubit (Life Technologies). Libraries were sequenced on Hiseq 4000

(100 bp PE, Illumina).

1.3 ChIP-seq

ChIP-seq experiments for all histone marks at all time points were performed as previously described[6,7]. The experiments were conducted according to experimental guidelines and read depth standards of the ENCODE consortium. Harvested cells were washed in PBS and fixed in 1% formaldehyde for 10 min at room temperature. Reaction was quenched with 125 mM Glycine for 5 min at room temperature followed by 10 min incubation on ice. Next, the cells were washed twice with PBS. Crosslinked cells were lysed and chromatin sheared using a Branson Sonifier 450 (20 cycles, 15 sec ON, 45 sec OFF at power 3). Detailed protocols can be found on the ENCODE homepage for immunoprecipitation (https://www.encodeproject.org/documents/89795b31-e65a-42ca-9d7bd75196f6f4b3/@@download/attachment/Ren%20Lab%20ENCODE%20Chromatin%20Immunoprecipitation%20Protocol_V2.pdf)

and library preparation (https://www.encodeproject.org/documents/4f73fbc3-956e-47ae-aa2d41a7df552c81/@@download/attachment/Ren_ChIP_Library_Preparation_v060614.pdf).

ChIP-seq experiments for CTCF and SMC3 were carried out on two and ten million crosslinked cells respectively, as previously described[8,9]. Nuclei were isolated using hypotonic buffer and subsequecntly lysed in RIPA buffer (140 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% SDS, 0.1% sodium deoxycholate, 10 mM Tris-HCl pH 8.0 supplemented with protease inhibitor (Roche)) and chromatin was sheared using a Covaris M220. Cell lysates were precleared using Protein G beads (Sigma) and rotation for 3h at 4°C. For immunoprecipitation, chromatin was incubated with antibodies against CTCF overnight at 4°C. Antibody chromatin complexes were bound by Protein G beads for 2h at 4°C. After eight rounds of washing using combinations of high and low salt buffers, crosslinks were reversed with Proteinase K (NEB) at 65°C overnight. DNA was purified by phenol /chloroform/isoamylalcohol extraction and ethanol precipitation. Sequencing libraries were generated using the Accel-NGS® 2S Plus DNA Library Kit (Swift).

All ChIP-seq libraries were sequenced on HiSeq2500/4000 (50 bp SE, Illumina).

A list of antibodies used in this study can be found in **Extended Data Table 7**.

1.4 RNA-seq

Total RNA from cells was extracted using Trizol (Thermo) and purified with RNA Clean & Concentrator™-5 (R1013, Zymo Research). RNA sequencing libraries were prepared from 1-2 µg total RNA using the Illumina TruSeq Stranded mRNA Library Prep Kit Set A (Illumina, RS-122-2101) or Set B (Illumina, RS-122-2102). Sequencing was carried out on a HiSeq4000 (100 bp PE, Illumina).

1.5 ATAC-seq

ATAC-seq was performed on 50,000 cells as described previously with minor modifications[10]. Cells were thawed at 37°C and pelleted for 5 min at 500 x g at 4 °C. Cell pellets were resuspended in 250 µl permeabilization buffer (5 % BSA, 0.2 % IGEPAL-CA630, 1mM DTT and cOmplete™, EDTA-free protease inhibitor cocktail (Roche) in PBS) and incubated on an overhead shaker for 5 min at 4°C, followed by centrifugation for 5 min at 500 x g. Pellet was resupended in 25 µl tagmentation buffer (36.3 mM Tris-acetate (pH = 7.8), 72.6 mM K-acetate, 11 mM Mg-acetate, 17.6 % DMF) and counted on a hemocytometer. 40,000 nuclei were used for tagmentation and the reaction volume was adjusted to 19 µl using tagmentation buffer. After addition of 1 µl Tn5 transposomes (Illumina) tagmentation was performed at 37°C for 60 min with shaking (500 rpm). Next, samples were purified using MinElute columns (Qiagen), PCR-amplified for 9 cycles with NEBNext® High-Fidelity 2X PCR Master Mix (NEB, 72°C 5 min, 98°C 30 s,[98°C 10 s, 63°C 30 s, 72°C 60 s] x cycles, 72°C forever). Amplified libraries were purified using MinElute columns (Qiagen) and Ampure XP Bead (Beckmann Coulter). Sequencing was carried out on a HiSeq2500 or 4000 (50 bp PE, Illumina).

1.6 Cardiac chromatin loop functional studies

To further validate identified chromatin loops regulating predicted enhancer-promoter interactions, selected enhancers were genetically deleted from hESC lines using CRISPR-Cas9 genome-editing strategies. To this end, specific gRNA pairs with binding sites surrounding the target region were designed[11] using a web-based software tool CRISPOR[12], which generates gRNA sequences for optimal targeting of region of interest while minimizing potential off-target effects. These identified gRNAs were then

synthesized *in vitro* using the GeneArt Precision gRNA Synthesis kit (Invitrogen). To

better monitor the differentiation efficiency, we generated both eWNT3 and eMSX1

knockout line in H9 *TNNT2*:GFP hESCs, which enables us to dynamically track the

cardiomyocyte generation by GFP[13]. One day before transfection, 1×10^5 H9

TNNT2:GFP hESCs were seeded in 12-well plates. When cells were 30-60% confluent,

a pair of RNP complexes containing 2 µg of Cas9 protein (NEB) and 400 ng of *in vitro*

transcribed gRNA were transfected using Lipofectamine Stem Transfection Reagent

(Invitrogen)[14,15]. Three days after transfection, cells were diluted and clonally expanded

for genotyping and functional assays[16]. Clonal mutant lines were genotyped using PCR

and Sanger sequencing (**Extended Data Table 8**). For heterozygous knock-out alleles,

a second round of targeting was performed to create homozygous knock-out lines.

After confirmation of genome-edited clones, at least three successfully targeted clones

per deletion were then selected for further functional analyses. Using the Wnt-based

CM differentiation protocol[2] described above, enhancer deletion clones (i.e. eWNT3[-/-]

and eMSX1[-/-]) were differentiated to respective stages as to when enhancer-promoter

loops appear. Differentiation efficiency was assessed as described above by flow

cytometry and replating studies if applicable. Samples that resulted in high efficiency

differentiation (>90%) were then further processed and analyzed by qPCR for the

expression of cognate genes that are predicted to be regulated by studied enhancer. As

a negative control, expression of the nearest genes of the studied enhancer was also

examined. Gene expression levels were normalized to the expression of TBP of each

sample. For WNT3 locus, four differentiations were performed; the first three

differentiations had one replicate, the 4[th] differentiation had two replicates. For MSX1

locus, three differentiations were performed, each with three replicates.

1.7 HERV-H functional studies

To interrogate the functional significance of HERV-Hs, selected HERV-H sequences

were genetically deleted in H9 *MLC2v*:H2B-GFP hESC lines using CRISPR-Cas9

genome-editing strategies as described above for cardiac chromatin loop functional

studies. After confirmation of genome-edited clones by Sanger sequencing, at least two

successfully targeted clones per deletion were selected for functional analysis. To

examine the effects of deleting HERV-Hs on chromatin boundary formation and gene

expression, *in-situ* Hi-C and RNA-seq studies were respectively performed as described

above. Because of the potential role of HERV-H1 in cardiac differentiation, HERV-H1

deleted H9 *MLC2v*:H2B-GFP hESC clones were additionally examined for their ability to

differentiate into cardiomyocytes. To this end, hESC cell seeding density and canonical

Wnt signaling levels were titrated to initially find suboptimal CM differentiation conditions

that result in 50% CM differentiation by Day 15 in control H9 *MLC2v*:H2B-GFP hESCs.

These suboptimal CM differentiation conditions were then applied to both control and

HERV-H1 deleted H9 *MLC2v*:H2B-GFP hESC clones to further assess their ability to

differentiate into CMs from Day 7-15 as detected by TNNT2 FACS analyses. The

differentiation experiments were repeated three times.

2 Data Analysis

2.1 ChIP-seq data processing:

2.1.1 Read alignment

Reads were aligned to hg19 (GRC37) following standards of ENCODE ChIP-seq

pipeline[17]. The full pipeline is available at https://github.com/ren-lab/chip-seq-pipeline.

Briefly, the reads were mapped with bwa[18]. PCR duplicates were removed using

Picard[19]. Only reads with mapping quality > 30 were kept.

2.1.2 ChIP-seq peak calling

For each stage, ChIP-seq peaks were called using MACS2[20] for each biological

replicate and pooled sample, using the pooled input sample as control. MACS2 was run

with the default setting with "--nomodel –extsize 180" parameter. Peaks called in the

pooled sample that also intersected with peaks in both replicates by at least 50% bases

were defined as replicated peaks and were used as the final peak list for that stage.

Peaks from all stages were merged into a union peak list using bedtools mergeBed[21].

2.1.3 ChIP-seq signal visualization

ChIP-seq bigWig files were generated using deeptools[22] bamCoverage function with the

RPKM (Reads Per Kilobase of transcript per Million mapped reads) normalization

option. Aggregated signal profiles were generated with the deeptools computeMatrix

and plotProfile functions and further modified in R.

2.2 ATAC-seq data processing:

2.2.1 Read alignment

Reads were aligned to hg19 (GRC37) following standards of ENCODE ATAC-dnase-

seq pipeline[23]. The full pipeline (including peak-calling) is available at

https://github.com/ren-lab/atac-seq-pipeline. Briefly, the reads were mapped with

bowtie2[24]. PCR duplicates were removed using Picard. Only reads with mapping

quality > 30 were kept.

2.2.2 ATAC-seq peak calling

For each stage, ATAC-seq peaks were called using MACS2 for each biological replicate

and pooled sample, using the pooled input sample as control. MACS2 was run with the

default setting with "--nomodel –extsize 73 –p 0.01" parameter. Peaks called in the

pooled sample that also intersected with peaks in both replicates by at least 50% were

defined as replicated peaks and were used as the final peak list for that stage. Peaks

from all stages were merged into a union peak list using bedtools mergeBed. The peaks

were further separated into promoter-distal and promoter-proximal by whether the

peaks intersected with GENCODE v19 transcription start sites (TSSs).

2.3 RNA-seq data processing:

2.3.1 Read alignment and quantification

RNA-seq alignment and quantification pipeline is available at https://github.com/ren-

lab/rnaseq-pipeline. Briefly, reads were aligned to hg19 (GRC37) and GENCODE GTF

v19 with rnaSTAR[25] following "ENCODE" options outlined in the rnaSTAR manual

(http://labshare.cshl.edu/shares/gingeraslab/www-

data/dobin/STAR/STAR.posix/doc/STARmanual.pdf). PCR duplicates were removed

using Picard. Raw reads for each gene were quantified using featureCounts[26]. Read per

million reads and kilobases (RPKM) were calculated for each gene.

 2.3.2 Differentiation gene expression analysis

2 Differentially gene expression was performed using edgeR[27] with the default

3 normalization method, tagwise dispersion estimation function and glmLRT (negative

4 binomial generalized log-linear model).

6 2.4 Hi-C data processing:

The HiC data processing pipeline is available at https://github.com/ren-lab/hic-pipeline.

Hi-C reads were aligned on hg19(GRC37) using BWA-MEM[18] for each read separately,

and then paired. For chimeric reads, only 5' end mapped locations were kept.

Duplicated read pairs mapped to the same location were removed to leave only one

unique read pair. The output bam files were transformed into juicer file format and tag

directories for further analysis in Juicebox [28]and HOMER[29], respectively. Unless

specified otherwise, we used the contact matrix in 10kb resolution and normalized using

KR matrix balancing method.

2.4.1 Compartments A/B

Principal component analysis was performed using HOMER[29] with resolution of 50kb.

DNA segments with positive PC1 value in both biological replicates were defined as

compartment A, whereas negative value for compartment B. Inconsistent PC1 signs

between replicates were defined as inconsistent. For dynamic and static compartment

analysis (**Fig 1.d**), only DNA segments with consistent compartment state at each of the

six stages (which was 85% of all genomic bins) were included. A DNA segment was

defined as static if it stayed compartment A or B at all six stages; Otherwise it was

named dynamic. Compartment switch types were defined as "BA" if it switched from B

to A, and *vice versa*. If the compartment bin switched from A to B, and then switched

back to A, it will be defined as "ABA", and the same rule applies to "BAB". Bins with

more than 2 switches were very rare and left out in the analysis/plot. Hierarchical

clustering was performed on the PC1 values of all dynamic bins, by both rows

(segments) and columns (sample replicates), using (1- Pearson correlation coefficient)

as distance and average linkage method to merge branches.

1. *2.4.2 Insulation score*

2. To quantify insulation based on KR normalized contact frequencies we used the

3. insulation score defined in Crane et al. 2015[30], using 500kb as the square size, and

4. 200kb as the delta size. Insulation TADs were called using the perl script at

5. https://github.com/dekkerlab/crane-nature-2015.

6. *2.4.3 Directionality Index*

7. We calculated the Directionality Index (DI) as described in Dixon et al[31]. The KR

8. normalized contact matrices were used as input to this transformation. We generated

9. the DI score at two different resolutions: 10kb and 40kb. In both cases, we limited the

10. scope of interaction to 2MB upstream or downstream.

11.

12. *2.4.4 Topological associated domains (TADs)*

13. *2.4.4.1 TAD identification.*

14. We called TADs using three different algorithms: domain caller[31], Arrowhead[5] and

15. insulation score[30]. All methods were run with the default parameters recommended in

16. the user manual and with the resolution at 10kb. They produced different number of

17. TADs because of inherent difference in their algorithms and choice of default

18. parameters. Nevertheless, the same trend of TAD loss was observed (**Extended Data**

19. **Fig. 5a,b,c**).

20. *2.4.4.2 Identification of non-redundant TAD boundaries*

21. TAD boundaries were previously found to be conserved between cell types[32]. Based on

22. this assumption, we should be able to find a set of non-redundant TAD boundaries

23. across different cell types. To identify non-redundant boundaries across all samples, we

24. collapsed spatially identical or close TADs across all time points and replicates. To do

25. this, we first quantile normalized the Directionality Index (DI) scores genome-wide

26. between all twelve samples. Then we calculated the DI delta scores for each TAD

27. boundary. DI delta score is the average difference in DI between the 4 bins (10kb per

28. bin) downstream and 4 bins upstream of the boundary. We concatenated TAD

29. boundaries from all samples (twelve) and sorted them by their DI delta scores in

30. descending order. Then we picked one TAD boundary from the top of the list and

31. remove any remaining boundary within 50Kb of the top TAD boundary. The number as

well as the stages of removed boundaries are recorded in the kept TAD boundaries as two additional columns. Then we picked the next TAD boundary on the list and repeat the same process, until the entire list was traversed. To remove unwanted spurious calls, any TAD boundary without a redundant (duplicate) call is discarded. For each non-redundant TAD boundary, it is denoted as a boundary (TRUE) in a stage if the DI delta score is greater than 200 in both replicates in that stage, otherwise is FALSE.

2.4.4.3 Dynamic TAD boundaries across stages

To determine dynamic TAD boundaries across stages, we first obtained the DI delta scores for each non-redundant TAD boundary. Then we log transformed the score and used the values as input to LIMMA[33]. LIMMA package was used to call differential boundary strength between each adjacent stage comparisons (e.g. D00 vs D02, D02 vs D05, etc). P-values were calculated with the eBayes function within LIMMA with trend parameter disabled, and were adjusted using Benjamini-Hochberg method. A TAD boundary was called dynamic if its adjusted p-value was less than 0.01 in any adjacent stage comparisons. Otherwise, it was called a stable TAD boundary. However, if the delta scores of that boundary were consistently greater than 200 in all samples, regardless of its state called in LIMMA, it was denoted as stable TAD boundary. Within dynamic TAD boundaries, we denoted the ones that were only present (TRUE) at D00 as ESC(+) TAD boundaries, those that were only FALSE at D80 as vCM(-) TAD boundaries (**Fig. 1e**).

2.4.4.4 Overlap of TAD and TAD boundaries with CTCF and repeat elements

Overlap of CTCF ChIP-seq peaks and repeat elements to TAD and/or TAD boundaries were performed using BEDtools intersect[21]. When performing intersections, the TAD boundaries were increased to ±25kb from the original TAD boundaries to account for algorithmic inaccuracies in calling boundaries. Repeat elements were RepeatMasker annotation downloaded directly from UCSC[34]. To avoid spurious results introduced by repeats with few copies in the genome, only repeats with more than 1000 copies genome-wide were tested here. The intersections were performed against each repeat name category separately, and for ESC(+) TADs, vCM(-) TADs and all stable TADs. For each repeat element, we obtained the fraction of TAD boundaries that contained that repeat. Then we calculated the fold enrichment of each repeat element at the TAD

boundaries of ESC(+) (or vCM(-)) TADs over stable TAD boundaries. The significance

of the enrichment was computed using two-sided proportion test function in R. The

repeat elements with fold enrichment greater than two and p-value less than 0.01 were

reported as significantly enriched.

2.4.4.5 Gene expression profile at the TAD boundaries containing repeat elements

*(relating to **Fig. 2a** and **Extended Fig. 6b**)*

For each of the eight repeat elements found in **Extended Fig.6a**, we kept the DNA

sequences that intersected with ESC(+) TAD boundaries. Using these sequences as

reference point, we generated the average RPKM normalized signal over ±50kb of the

reference point with deeptools computeMatrix function[22]. Only LTR7 and HERVH-int

sequences displayed high expression in the hESC.

*Mouse ES cell TAD boundaries and overlap with human TAD boundaries (**Fig.3d,e,***

***Extended Fig.9a**)*

Mouse ESC Hi-C data from *Bonev et al*[35] was downloaded from SRA and aligned to

mm10 with the same procedure as described above. The TAD boundary coordinates

were converted to hg19 using UCSC liftover tool with minimum ratio of base remap of

0.1. Human TAD boundaries overlap with lifted-over mouse TAD boundaries were

performed using bedtools intersect. The significance of the difference in fraction of

overlap was computed using two-sided proportion test function in R.

2.4.5 Chromatin Loops

2.4.5.1 Identification of chromatin loops

Chromatin loops were called using an adapted algorithm from HICCUPS[5] at 10kb

resolution, with the major difference being that only the "donut region" around the pixel

were used to model the expected count.

The input for the method were raw and KR-normalized contact matrices of 10kb

resolution. In preparation, distance-based mean interaction frequency was calculated

from KR-normalized matrix, as well as the raw-to-KR normalization factor matrix. For

each bin (B_i) in the matrix, we isolated the surrounding bins (B_s) within 40kb (both

horizontal and vertical) of that bin. For each surrounding bin, we corrected its contact

frequency with respect to B_i by distance-based curve, i.e.

$$B'_s = B_s \times F_i / F_s$$

F_i and F_s are the genome-wide average interaction frequencies for the distance of the test bin and surrounding bin, respectively. B'_s is the corrected frequency. Then we used the mean of the corrected frequencies from all surrounding bins as the expected contact (E_i) for the test bin. Because the expected contact was calculated using the balanced matrix, we transformed it to the raw expected contact by multiplying it with the raw-to-KR normalization factor. Finally, we calculated the probability of seeing the raw observed contact based on Poisson distribution with lambda equals expected raw contact.

In the first pass of post-processing, bins with p-value less than 0.01 and distance greater than 10kb were marked as candidate bins and kept for further analysis. Candidate bins often form clusters, because if a genomic locus is interacting with another, their nearby bins should also be physically close. Therefore, if a candidate bin is not adjacent to any other candidate bins, it is removed because it is very likely a false positive. Finally, we collapsed candidate bins within 20kb of each other into the bin with most significant p-value. The collapsed bins with p-value less than 1e-5 was designated as significant chromatin loops.

2.4.5.2 Merge redundant loops across stages

Non-redundant loops across all stages and replicates (total of 12 samples) were identified using a similar algorithm as that for non-redundant TAD boundaries in the previous section. To begin with, loop calls from all samples were concatenated and sorted by the p-values in ascending order. Loop from the top of the list was selected and any loops within 30kb of that top loop were collapsed into the top loop. This process was repeated for the next top loop until the whole list was traversed. To remove unwanted spurious calls, any loop without a redundant (replicated) loop call was discarded. For each non-redundant loop, it is denoted as TRUE in a stage if it is called as loop in that stage, otherwise FALSE.

2.4.5.3 Identification of stage-specific loops

For each non-redundant loop, Bioconductor package edgeR[27] was utilized to evaluate the dynamicity of the interaction frequencies over the stages. Raw interaction

frequencies were provided as an input to edgeR. The contrast matrix was constructed to

test the difference between every adjacent stages, e.g. D02.vs.D00, D05.vs.D07, etc.

The statistical tests were performed using the glmLRT function per the pacakge's

instruction. The resulting p-values were adjusted using Bonferroni correction. A loop

was defined as stage-specific if its adjusted p-value was less than 0.01.

2.4.5.4 K-means clustering

In search of over-represented patterns of stage-specific loop interactions over

differentiation, we performed K-means clustering. The input matrix for K-means was

number of contacts per million read pairs. Each row (loop) was further divided by a

common factor to become a unit vector. K-means clustering was performed using K

from two to ten. When K was greater than five, there was no apparent gain of new

patterns. Therefore, we chose K=5 to split the clusters.

2.4.5.5 Gene set and motif enrichment testing at the loop anchors

For each loop cluster, we extended the loop anchors to 15kb from the center of the

anchor to account for potential errors in finding the exact pixel of the loop. Loop anchors

were overlapped with transcription start sites of genes and merged ATAC-seq peaks.

The genes whose TSSs overlapped anchors from each loop cluster were provided as

input to DAVID[36] to perform gene set enrichment analysis. On the other hand, The

ATAC-seq peaks were analyzed for transcription factor motif enrichment using

HOMER[29]. The default parameters were chosen according to the user manual.

2.4.5.6 Multi-loop anchors and overlap with transcription factors, house-keeping gene

TSSs, and CTCF peaks

We took all loop anchors (In total 19938 bins, each with 10kb resolution) from the 14215

chromatin loops we identified and extend the anchors to 15kb around their centers. We

calculated the number of loops that interacted with each bin as the degree of

connectivity for that bin (**Fig. 4f**). Then we overlapped the anchors with the TSSs of

1999 transcription factors[37], 3804 house-keeping genes[38], and merged CTCF ChIP-seq

peaks from our data. The fraction of anchors overlapping these features were plotted for

anchors with different degree of connectivity (**Fig. 4g,h,i**).

2.4.5.7 CTCF motif and orientation

We scanned the CTCF motif (MA0139.1) from JASPAR database[39] and kept track of

their orientation in merged CTCF ChIP-seq peaks with FIMO[40]. We overlapped the

chromatin loops with the CTCF ChIP-seq peaks, and separated the loops into four

categories with respect to their CTCF motif orientation located on the loop anchors. The

four categories are: (1) No-CTCF: at least one anchor does not contain any CTCF

ChIP-seq peak; (2) Unkown: Both anchors overlapped with CTCF peaks. At least one

anchor does not contain a matched CTCF motif; (3) Incorrect: both anchors overlapped

with CTCF peaks. CTCF motifs at loop anchors do not face inward; (4) Inward: both

anchors overlapped with CTCF peaks. CTCF motifs at loop anchors face inward.

2.4.5.8 loop location relative to TADs

We assigned chromatin loops into three categories with respect to their location to TAD

boundaries. (1) TAD-corner: a loop is located within 20kb of any TAD corner. (2) Intra-

TAD: a loop is inside of any TAD. (3) inter-TAD: a loop that is outside of all TADs. When

there are conflicts, the priority of assignment is (1)>(2)>(3).

2.5 HERV-H related analysis

2.5.1 Directionality index profile on HERV-Hs transcribed at different levels

To investigate the boundary activity of HERV-Hs that transcribed at different levels, we

visualized the Directionality Index (DI) scores at 10kb resolution. We sorted the HERV-

H loci by their expression RPKMs in the hESC (day 0) in descending order and assign

every 50 loci to a bin. For each bin, we generated the aggregated median DI score

profile 70kb surrounding the HERV-H loci. The same DI scores were computed at day 2

and day 5. As comparison, we also plotted DI score profile for genes that expressed

with similar RPKMs as the top-tier HERV-Hs in the hESC, and down-regulated at least 2

fold after differentiation (**Extended Data. Fig7. a**).

2.5.2 Directionality index profile on HERV-Hs in iPSCs and H1-derived cells (relating to

Extended Data. Fig7. b).

The Hi-C data for human iPSCs were each from Greenwald *et al*[41]. The Hi-C data from

H1 ESC and its derived cells were from Dixon *et al*[31,32]. The data were downloaded and

processed the same way as described above. Because DI scores are dependent on the

sequencing depth, the H1ESC DI profile may look less pronounced comparing to the

iPSC.

2.5.3 ChIP-seq enrichment analysis at TAD boundary-associating HERV-H loci (relating

*to **Extended Data. Fig. 7c**)*

To investigate which protein(s) may be responsible for TAD boundary formation at top

50 transcribing HERV-H loci, we downloaded hESC ChIP-seq bigWig files from

ENCODE (primary) and other sources in the public domain (prefixed with the SRA ID if

downloaded from other than ENCODE)[42]. For each ChIP-seq data, we computed the

average bigWig signal on the top 50 HERV-Hs with *bigWigAverageOverBed* command

from UCSC, and the average bigWig signal on HERV-Hs ranking from 51 to 300. Then

we calculated the fold change with these two averages.

*2.5.4 HERV-H LTR type and divergence analysis (relating to **Extended Data. Fig. 9**)*

To investigate the sequence features of the LTRs flanking TAD boundary associated

HERV-Hs and non-boundary associated HERV-Hs, we examined their presence, length

and sequence divergence. The LTRs for HERV-H include LTR7, LTR7A, LTR7B,

LTR7C and LTR7Y. We downloaded the annotations for those LTRs from repeatMasker

and assigned them to each HERV-H-int sequence by base overlap. The HERV-H loci

were ordered by their expression levels and binned as described above (the top 50

were enriched at TAD boundaries).

Sequence divergence between the 5' LTR and 3'LTR sequences was used to measure

the insertion time for retrotransposons. Based on the retrotransposon insertion

mechanism, at the time of insertion, 5' and 3' LTR sequences should be identical; the

LTR sequences will each accumulate mutations over time, and thus their sequence

differences could be used as a proxy to their ages. We followed the steps outlined in

https://github.com/SIWLab/Lab_Info/wiki/Ageing-LTR-insertions to align each LTR pairs

and calculated the sequence divergences[43,44].

2.6 GWAS analysis:

2.6.1 GWAS lead SNP and SNPs in linkage disequilibrium

Human GWAS association data was downloaded from human GWAS catalog
(https://www.ebi.ac.uk/gwas/) as of Sept 28th, 2017. Heart-related traits were selected
for analysis. These traits are: "Atrial fibrillation", "Cardiac repolarization", "Coronary
heart disease", "Conotruncal heart defects", "Congenital left-sided heart lesions",
"Congenital heart disease", "Congenital heart malformation", "Heart failure", "Mortality in
heart failure", "PR interval", "QRS duration", "QT interval", and "Sudden cardiac arrest".
For all lead SNPs, we obtained the SNPs in linkage disequilibrium (LD) using SNAP[45]
with default parameters (SNP data set was set to "1000 Genome Pilot1", Population
panel set to "CEU", r square threshold > 0.8 and distance limit equals 500). The output
from SNAP was in hg18 coordinates, and therefore was lifted over to hg19 with UCSC
liftover tool.

2.6.2 Binomial test for significance of enrichment of heart-traits GWAS SNPs within
chromatin loops
The LD SNPs were overlapped with merged ATAC-seq peaks within chromatin loop
anchors using bedtools. The chromatin loops used here were loops present in
cardiomyocyte differentiation day 2 – day 80. For each GWAS trait, the significance of
enrichment was computed using a binomial test as described in Maurano *et al*[46]. The
total number of tests was total number of lead SNPs for each trait. The number of
successes was the number of lead SNPs that were overlapped with open chromatin
region within the loop anchor. For each lead SNP, if itself or any of the LD SNPs were
overlapped with an open chromatin region within chromatin loop anchors, that lead SNP
was considered a positive hit. The hypothesized probability of success is the width of
ATAC-seq peaks within chromatin loop anchors divided by the length of the genome.

Gene Ontology Enrichment testing of the genes.
We collected all of the SNPs overlapped with chromatin loop anchors and ATAC-seq
peaks in the previous section, and identified the candidate target gene at the other end
of the chromatin loop. Using this list of gene as input, we performed GO enrichment
analysis using PANTHER[47].

2.7 General data processing and plots

Most of the described data processing steps (statistical tests, clustering, plotting, etc) were performed in python 3.4.5 (www.python.org) and statistical computing environment R 3.4.3 (www.r-project.org). Boxplots were made with ggplot2 (https://cran.r-project.org/web/packages/ggplot2). The elements of the boxplot are: center line, median; box limit, upper and lower quartiles; whiskers, 1.5x interquartile range; points, outliers.

3 Data and script availability:

All sequencing datasets have been deposited on GEO with the accession number

GSE116862.

Scripts are available at https://github.com/shawnzhangyx/cvdc_scripts.

Additional References:

Veevers, J. *et al.* Cell-Surface Marker Signature for Enrichment of Ventricular Cardiomyocytes Derived from Human Embryonic Stem Cells. *Stem cell reports*, doi:10.1016/j.stemcr.2018.07.007 (2018).
Lian, X. *et al.* Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/beta-catenin signaling under fully defined conditions. *Nature protocols* **8**, 162-175, doi:10.1038/nprot.2012.150 (2013).
Hashem, S. I. *et al.* Brief Report: Oxidative Stress Mediates Cardiomyocyte Apoptosis in a Human Model of Danon Disease and Heart Failure. *Stem cells* **33**, 2343-2350, doi:10.1002/stem.2015 (2015).
Milani, P. *et al.* Cell freezing protocol suitable for ATAC-Seq on motor neurons derived from human induced pluripotent stem cells. *Scientific reports* **6**, 25474, doi:10.1038/srep25474 (2016).
Rao, S. S. *et al.* A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* **159**, 1665-1680, doi:10.1016/j.cell.2014.11.021 (2014).
Shen, Y. *et al.* A map of the cis-regulatory sequences in the mouse genome. *Nature* **488**, 116-120, doi:10.1038/nature11243 (2012).
Gorkin, D. *et al.* Systematic mapping of chromatin state landscapes during mouse development. *bioRxiv* (2017).
Jolma, A. *et al.* DNA-binding specificities of human transcription factors. *Cell* **152**, 327-339, doi:10.1016/j.cell.2012.12.009 (2013).
Yan, J. *et al.* Transcription factor binding in human cells occurs in dense clusters formed around cohesin anchor sites. *Cell* **154**, 801-813, doi:10.1016/j.cell.2013.07.034 (2013).

Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nature methods* **10**, 1213-1218, doi:10.1038/nmeth.2688 (2013).

Liu, Z. *et al.* Efficient CRISPR/Cas9-Mediated Versatile, Predictable, and Donor-Free Gene Knockout in Human Pluripotent Stem Cells. *Stem cell reports* **7**, 496-507, doi:10.1016/j.stemcr.2016.07.021 (2016).

Haeussler, M. *et al.* Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome biology* **17**, 148, doi:10.1186/s13059-016-1012-2 (2016).

Wrighton, P. J. *et al.* Signals from the surface modulate differentiation of human pluripotent stem cells through glycosaminoglycans and integrins. *Proc Natl Acad Sci U S A* **111**, 18126-18131, doi:10.1073/pnas.1409525111 (2014).

Kim, S., Kim, D., Cho, S. W., Kim, J. & Kim, J. S. Highly efficient RNA-guided genome editing in human cells via delivery of purified Cas9 ribonucleoproteins. *Genome research* **24**, 1012-1019, doi:10.1101/gr.171322.113 (2014).

Zuris, J. A. *et al.* Cationic lipid-mediated delivery of proteins enables efficient protein-based genome editing in vitro and in vivo. *Nature biotechnology* **33**, 73-80, doi:10.1038/nbt.3081 (2015).

Santos, D. P., Kiskinis, E., Eggan, K. & Merkle, F. T. Comprehensive Protocols for CRISPR/Cas9-based Gene Editing in Human Pluripotent Stem Cells. *Current protocols in stem cell biology* **38**, 5B 6 1-5B 6 60, doi:10.1002/cpsc.15 (2016).

https://github.com/ENCODE-DCC/chip-seq-pipeline.

Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).

http://broadinstitute.github.io/picard/.

Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome biology* **9**, R137, doi:10.1186/gb-2008-9-9-r137 (2008).

Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842, doi:10.1093/bioinformatics/btq033 (2010).

Ramirez, F., Dundar, F., Diehl, S., Gruning, B. A. & Manke, T. deepTools: a flexible platform for exploring deep-sequencing data. *Nucleic Acids Res* **42**, W187-191, doi:10.1093/nar/gku365 (2014).

https://github.com/kundajelab/atac_dnase_pipelines.

Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature methods* **9**, 357-359, doi:10.1038/nmeth.1923 (2012).

Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21, doi:10.1093/bioinformatics/bts635 (2013).

Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930, doi:10.1093/bioinformatics/btt656 (2014).

Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140, doi:10.1093/bioinformatics/btp616 (2010).

Durand, N. C. *et al.* Juicebox Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. *Cell Syst* **3**, 99-101, doi:10.1016/j.cels.2015.07.012 (2016).

Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589, doi:10.1016/j.molcel.2010.05.004 (2010).

Crane, E. *et al.* Condensin-driven remodelling of X chromosome topology during dosage compensation. *Nature* **523**, 240-244, doi:10.1038/nature14450 (2015).

Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**, 376-380, doi:10.1038/nature11082 (2012).

Dixon, J. R. *et al.* Chromatin architecture reorganization during stem cell differentiation. *Nature* **518**, 331-336, doi:10.1038/nature14222 (2015).

Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47, doi:10.1093/nar/gkv007 (2015).

Casper, J. *et al.* The UCSC Genome Browser database: 2018 update. *Nucleic Acids Res* **46**, D762-D769, doi:10.1093/nar/gkx1020 (2018).

Bonev, B. *et al.* Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell* **171**, 557-572 e524, doi:10.1016/j.cell.2017.09.043 (2017).

Huang da, W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**, 1-13, doi:10.1093/nar/gkn923 (2009).

Vaquerizas, J. M., Kummerfeld, S. K., Teichmann, S. A. & Luscombe, N. M. A census of human transcription factors: function, expression and evolution. *Nat Rev Genet* **10**, 252-263, doi:10.1038/nrg2538 (2009).

Eisenberg, E. & Levanon, E. Y. Human housekeeping genes, revisited. *Trends Genet* **29**, 569-574, doi:10.1016/j.tig.2013.05.010 (2013).

Sandelin, A., Alkema, W., Engstrom, P., Wasserman, W. W. & Lenhard, B. JASPAR: an open-access database for eukaryotic transcription factor binding profiles. *Nucleic Acids Res* **32**, D91-94, doi:10.1093/nar/gkh012 (2004).

Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif. *Bioinformatics* **27**, 1017-1018, doi:10.1093/bioinformatics/btr064 (2011).

Greenwald, W. W. *et al.* Integration of phased Hi-C and molecular phenotype data to study genetic and epigenetic effects on chromatin looping. *bioRxiv* (2018).

Xu, C. & Corces, V. G. Nascent DNA methylome mapping reveals inheritance of hemimethylation at CTCF/cohesin sites. *Science* **359**, 1166-1170, doi:10.1126/science.aan5480 (2018).

Hu, T. T. *et al.* The Arabidopsis lyrata genome sequence and the basis of rapid genome size change. *Nat Genet* **43**, 476-481, doi:10.1038/ng.807 (2011).

Slotte, T. *et al.* The Capsella rubella genome and the genomic consequences of rapid mating system evolution. *Nat Genet* **45**, 831-835, doi:10.1038/ng.2669 (2013).

Johnson, A. D. *et al.* SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* **24**, 2938-2939, doi:10.1093/bioinformatics/btn564 (2008).

Maurano, M. T. *et al.* Systematic localization of common disease-associated variation in regulatory DNA. *Science* **337**, 1190-1195, doi:10.1126/science.1222794 (2012).

Mi, H. *et al.* PANTHER version 11: expanded annotation data from Gene Ontology and Reactome pathways, and data analysis tool enhancements. *Nucleic Acids Res* **45**, D183-D189, doi:10.1093/nar/gkw1138 (2017).

Figure 1

a

D00 hESCs → D02 mesodermal cells → D05 cardiac mesodermal cells → D07 cardiac progenitors → D15 primitive cardiomyocytes → D80 ventricular cardiomyocytes

chr4:167M-182M
chr4:167M-182M
3.3Mb
100

b

chr4:174.35-175.25MB
HAND2
LINC02268
200kb
50

CTCF — 15
ATAC-seq — 100
H3K27ac — 50
H3K27me3 — 40
H3K4me1 — 12
H3K4me3 — 40
mRNA — 500

HAND2   LINC02268

c

D00
HAND2

D05
-D07
HAND2

D15
-D80
HAND2

LINC02268 promoter
putative enhancer

d

Fraction of compartment types
1.00
0.75
0.50
0.25
0.00

stage-specific (8,683)
static (38,585)

compartment types
BAB   BA   B
ABA   AB   A

SOX2
HAND2
TTN
NFIB
RYR2

D00 D02 D05 D07 D15 D80
stages of differentiation
PC1 value (Compartment A/B)
-2   0   2

e

Fraction of TAD boundary types
1.00
0.68
0.00

stage-specific (1,233)
static (2,622)

TAD boundaries

vCM(-) (329)
ESC(+) (198)

D00 D02 D05 D07 D15 D80
stages of differentiation
DI delta
0 100 200 300 400 500

Figure 2

a HERVH-int — D00, D02, D05, D07, D15, D80
Expression (RPKM) vs TAD boundary (-50KB, ESC(+), +50KB)

b HERV-H expression (RPKM) vs Rank of HERV-H expression at D00 — D00, D02, D05; Top% of all genes at D00 (1%, 5%, 10%, 25%)

c D00 — Rank 1-50, Rank 51-100, Rank 101-150; -200kb, HERV-H, 200kb

d Rank 1-50 — D00, D02, D05; -200kb, HERV-H, 200kb; log2(obs/exp)

e HERV-H1 locus — chr4:91M-95M; contacts 30; D00, D02, D05; CCSER1, LNCPRESS2, GRID2, ATOH1; chr4:93,170,027-93,202,961; POLR2A, SMC3, CTCF, H3K27ac, RNA-seq; LNCPRESS2, HERVH-int 3'LTR7 5'LTR7

f HERV-H2 locus — chr13:55M-57.7M; contacts 30; D00, D02, D05; MIR5007; chr13:56,131,656-56,166,400; HERVH-int 3'LTR7 5'LTR7

g HERV-H rank 001-050, 051-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400; RNA-seq, POLR2A, SMC3, CTCF; -30KB, 5'LTR, 3'LTR, +30KB

Figure 3

Figure 4

Figure 5

Extended Data Figure 1

a

b

Extended Data Figure 2

Extended Data Figure 3

Extended Data Figure 4

Extended Data Figure 5

Extended Data Figure 6

Extended Data Figure 7

Extended Data Figure 8

Extended Data Figure 9

a

b

c

d

Extended Data Figure 10

Extended Data Figure 11

Extended Data Figure 12

a

SCN5A Locus

b

MA0039.3:KLF4(JASPAR)

rs17608766

c

d