Assessment of Environmental and Occupational Exposure while Working with *Candida*

***auris* a Multidrug Resistant (MDR) Fungus in the Animal Facility.**

Keywords: *Candida auris,* Biocontainment, Real-time PCR, Animal Facility, Biosafety level 2.

Word Count:

Steven R. Torres[1,2+], Heather C. Kim[1,2+], Lynn Leach[1], Sudha Chaturvedi[1], Corey Bennett[1],

David Hill[1] and Magdia De Jesus[1,2*]

[1]Division of Infectious Diseases, Wadsworth Center, New York State Department of Health,

Albany NY USA. [2] Department of Biomedical Sciences, University at Albany, School of Public

Health, Albany, New York, USA

+ Both authors contributed equally to this publication

Address correspondence to:

Magdia De Jesus, mdejesus2@albany.edu.

*Present address: 120 New Scotland Avenue, Albany, NY 12208

ABSTRACT

In less than a decade since its identification in 2009, the emerging fungal pathogen

Candida auris has become a major public health threat due to its multidrug resistant (MDR)

phenotype, high transmissibility, and high mortality. Unlike any other *Candida* species, *C. auris*

has acquired high levels of resistance to an already limited arsenal of antifungals. As an

emerging pathogen, there are currently a limited number of documented murine models of *C.*

auris infection. These animal models use inoculums as high as 10^7-10^8 cells per mouse, and the

environmental and occupational exposure of working with these models has not been clearly

defined. Using an intravenous model of *C. auris* infection, this study determined that shedding of

the organism is dose-dependent. *C. auris* was detected in the cage bedding when mice were

infected with 10^7 and 10^8 cells, but not with doses of 10^5 and 10^6 cells. Potential for exposure to

C. auris during necropsies and when working with infected tissues was also demonstrated. To

mitigate these potential exposures, a rigorous "buddy system" workflow, biomonitoring and

disinfection procedures were developed that can be used to prevent accidental exposures when

using small animal models of *C. auris* infection.

INTRODUCTION

Since its identification in 2009, the emerging fungal pathogen *Candida auris* has become a major public health threat, due to its status as a multidrug resistant (MDR) fungus with high levels of resistance to an already limited arsenal of antifungals, such as the azoles and polyenes.[1-4] Minimal inhibitory concentrations (MICs) for antifungals used against *C. auris* have been reported to be as high as 256 mg/L for fluconazole and > 2 mg/L for amphotericin B.[1-5] The Centers for Disease Control (CDC) has documented that *C. auris* behaves more like transmissible bacterial multidrug resistant organisms (MDROs) such as methicillin resistant *Staphylococcus aureus* (MRSA) than any other *Candida* species. [6] Of particular concern is that the unpredictable antifungal resistance profile of *C. auris* is also showing evidence of reduced susceptibility to the echinocandins, a third class of antifungals.[4] *C. auris* causes invasive bloodstream infections with reported mortality rates as high as 30-60% in Venezuela and India, and most recently, a reported mortality rate of up to 78% in Panama.[7-9] Although the origin of *C. auris* is unknown, infections with this organism have been documented on five continents in at least 18 countries, including the United States.[7, 8, 10-18]

C. auris has a predilection for the skin, and recent studies have suggested that transmission and colonization can occur rapidly after a few days or even a few hours of exposure.[6] *C. auris* is also alarmingly persistent in the environment and can spread rapidly in hospital and nursing home settings.[19, 20] *C. auris* is often misidentified as *Candida haemulonii*, and without specialized laboratory methodologies such as Matrix-Assisted Laser Desorption Ionization Time of Flight Spectrometry (MALDI-TOF), diagnosis is delayed, leading to mortality rates as high as 35%.[16, 20, 21]

The current animal models of disseminated *C. auris* infection use concentrations that

range from 10^6-10^8, along with the immunosuppressive agent cyclophosphamide. These animal

models are being used to test new alternative therapeutics, including a small molecule antifungal

agent called APX001A derived from the pro-drug APX001 that inhibits the fungal protein Gwt1

(glycosylphosphatidylinositol-anchored wall transfer protein 1). A novel echinocandin called

rezafungin is also being tested in these animal models as a potential therapeutic to combat *C.*

auris.[22-25] Because of its multidrug resistance, environmental persistence, and the large

inoculum used when working with animal models infected with *C. auris*, the risk of an

occupational exposure during experiments for researchers and animal care staff is still unclear.

In this study, immune-competent and neutrophil-depleted mice were infected with the *C.*

auris strain M5658 that belongs to the South Asia clade. The concentrations of *C. auris* used to

infect mice ranged from 10^5-10^8 inoculum per animal to determine potential for exposure.[26] The

animal holding room and procedure rooms were monitored to determine whether there was a risk

for exposure during cage changes animal infections and animal procedures such as necropsies.

As part of this study, a rigorous "buddy system" workflow and disinfection protocol was

developed as part of safety procedures to prevent accidental exposures.

METHODS

Animal Use Ethics Statement

Animal experiments described in this study were approved by the Wadsworth Center's

Institutional Animal Care and Use Committee (IACUC) under protocol #18-450. The

Wadsworth Center complies with the Public Health Service Policy on Humane Care and Use of

Laboratory Animals and was issued assurance number A3183-01. The Wadsworth Center is fully

accredited by the Association for Assessment and Accreditation of Laboratory Animal Care

(AAALAC).

The "Buddy System" Workflow: Cage Handling and Cleaning in Animal Holding Room

C. auris is recommended by the Centers for Disease Control and Prevention (CDC) to be

handled at Biosafety Level 2 (ABSL-2). However, due to the limited knowledge of its

environmental exposure risk in an animal research setting, the organism was handled at Animal

Biosafety Level 2 (ABSL-2) with Animal Biosafety Level 3 (ABSL-3) containment practices

that included enhanced Personal Protective Equipment (PPE) and a strict no open handling

policy. Staff entering the animal facility used standard PPE that included booties, gloves,

bouffant, face mask, and a gown. Before entering the animal holding room, additional PPE was

donned which included a second pair of booties, gloves and a snap front gown (Valumax, New

York, NY).

Mice infected with *C. auris* were housed in locked micro-isolator cages (Tecniplast, West

Chester, PA) with a negative air flow set to 51%, RH% 47, 75 ACH. Micro-isolator cages also

provided a way to autoclave bedding and water bottles as a single unit after use, thus minimizing

exposure to animal caretakers. A "buddy system" was implemented such that one person could

hand the clean caging and other supplies to a second person performing the cage changes inside

of the biosafety cabinet **(Figure 1).** This allowed for a "clean" to "dirty" workflow with minimal

disruption of the laminar airflow of the biosafety cabinet. When there was a need to leave the

biosafety cabinet, the exterior gloves were doffed inside of the cabinet and a new pair was

donned. Blue pads placed inside of a biosafety cabinet (Kendall Healthcare Products, Mansfield,

MA) were sprayed with a 10% bleach solution. It is important to note that the 10% bleach

solution was made fresh daily as recommended by the CDC, since quaternary ammonia products

routinely used for disinfection in animal research settings are not effective against *C. auris*.[4, 27]

To change infected animals from cages, the in-use micro-isolator cage and an empty

clean cage were placed inside of the biosafety cabinet on top of the blue pads. Mice were

carefully transferred to the new cage. Cage bedding that fell into the biosafety cabinet was

collected by the blue pads sprayed with 10% bleach. Food was added to the new cages from a

stock strictly kept inside of the biosafety cabinet. Before removal from the biosafety cabinet,

cages were wiped down with 10% bleach by the individual inside the cabinet, followed by a

wipe down of 70% ethanol by the individual outside the cabinet. At the end of cage changes, the

blue pads were rolled up and placed into a biohazard waste bag. The biosafety cabinet and

contents were sprayed down with a 10% bleach solution and allowed to sit for 5 minutes,

followed by wipe down with 70% ethanol **(Figure 2)**. Before leaving the room, the exterior

booties, gloves and gown were doffed. Dirty micro-isolator cages were sent to be autoclaved as a

single sealed unit with bedding and water bottle still inside to reduce potential occupational

exposures to animal care takers during the transport of cages to the autoclave. While changing

animals from their cages, opened Sabouraud Dextrose Agar plates with antibiotics (40 µg/ml

streptomycin; 20 U/ml penicillin; 25 µg/ml chloramphenicol; 40 µg/ml gentamicin) were

positioned inside of the biosafety cabinet, as well as on the floor directly outside of the cabinet.

At the completion of the work flow, these plates were incubated at 37℃ for up to 2 weeks and

checked daily to monitor growth of *C. auris.*

Transferring Cages from Holding to Procedure Room

Cages with infected animals were transferred on to a laboratory cart that was placed

outside of the holding room. The cart contained dry blue pads that were not sprayed with 10%

bleach; the cages were placed on each shelf of the cart. Before leaving the holding room, exterior

gloves, booties and gowns were removed and placed in a biohazard waste bin. The cages were

then transferred from the holding room to the procedure room. When entering the procedure

room, clean outer booties, outer gloves, and snap-front gown (Valumax, New York, NY) were

donned. The cart with the isolator cages as well as any supplies was kept on the "clean" side of

the room, indicated by a line on the floor.

Animal Handling during Weight Measurements

Dry blue pads were placed inside of a biosafety cabinet to cover the surfaces and cages

were placed on top. To monitor daily weights throughout the experimental trial, mice were

placed into a sterilized container that contained airholes and weighed using a laboratory balance.

After the last mouse in each cage was weighed, the cage was sealed and wiped down with 10%

bleach by the person inside of the cabinet, followed by a wipe with 70% ethanol by the person

outside of the cabinet, before returning cages to the cart. This was repeated until all mice had

been weighed. At the end of workflow, if no other procedures were done, the blue pads in the

biosafety cabinet were rolled up and placed in a biohazard waste bag. The biosafety cabinet,

scale, weigh chambers, and all contents were sprayed with a 10% bleach solution. After 5

minutes, the surfaces were wiped and then cleaned with a 70% ethanol solution **(Figure 3).** As in

the holding room, open Sabouraud Dextrose Agar plates with antibiotics were positioned inside

of the biosafety cabinet, as well as one on the floor directly outside of the cabinet. The plates

were incubated at 37℃ for up to 2 weeks and checked daily to monitor growth of *C. auris.*

Animal Infections and Necropsies

Prior to *C. auris* infection, 100 µg of the monoclonal antibody RB6-5C8 that depletes Ly6G neutrophils (Thermo Fisher, San Diego, CA) was administered by intraperitoneal (i.p.) injection using a 25G needle (BD, Franklin Lakes, NJ). Twenty-four hours later, 100 µl of *C. auris* was administered via tail vein (i.v.) injection at concentrations of that range from 10^5-10^8 cells. To do the infections, mice were initially anesthetized in an isoflurane chamber located outside of the biosafety cabinet. Once the mouse was anesthetized, the individual outside the biosafety cabinet placed it inside the cabinet. The individual with the hands inside the cabinet then placed the mouse into an illuminated mouse restrainer (Braintree Scientific, Braintree, MA) and placed a nose cone on the mouse. After the injection, the mice were transferred back into the cage located inside of the biosafety cabinet. The injection syringe was immediately placed in a sharps container to prevent needlestick accidents. When all mice in the group had been injected, the cage was wiped with 10% bleach by the individual inside of the cabinet, followed by a wipe with 70% ethanol by the individual outside of the cabinet. This procedure was repeated until all groups had been injected. To collect organs by necropsy, mice were euthanized in a CO_2 chamber located inside of the biosafety cabinet and death was confirmed by cervical dislocation. Blood was collected via cardiac puncture using a 26G needle, and spleen, liver, kidney, stomach, heart, brain and lungs were collected and placed in labeled 5 ml tubes (CELLTREAT, Pepperell, MA) containing 2 ml of HBSS with no dye on ice. Organs were ground through a 70 µm cell strainer (CELLTREAT, Pepperell, MA) using a pestle (CELLTREAT, Pepperell, MA) and twenty microliters (20 µl) of the resulting homogenate were plated on Sabouraud Dextrose Agar plates with antibiotics in triplicate and incubated at 37℃ for 24 hrs for colony forming units (CFU). When there was a need to leave the biosafety cabinet, the exterior gloves were doffed and a new

pair donned before continuing to work. Glove changes were also done in between handling of

different treatment groups to avoid cross-contamination. Surgical tools were sanitized with 10%

bleach, 70% ethanol, and water between mice. The carcasses were placed into a sealable

biohazard bag. At the end of the procedure, blue pads were rolled up and placed into a biohazard

bag. All contents of the biosafety cabinet including the ice bucket, tubes where organs were

collected, and surgical tools were sprayed with 10% bleach, allowed to sit for 5 minutes, and

then wiped down with 70% ethanol. The biohazard bag containing the mouse carcasses was

placed into a second sealable biohazard bag and then frozen prior to final disposal by

incineration. The biohazard bags containing blue pads and any waste were placed into a

biohazard waste bin. The biosafety cabinet was wiped down with 10% bleach and then cleaned

with 70% ethanol.

Transfer of Animals and Tissues for Processing

The collected and ground organ samples were moved from the ice bucket and placed

inside of a specimen transport box (Nalgene, Rochester NY). The transport box was also sprayed

with 10% bleach for five minutes followed by 70% ethanol, wiped and placed on the clean side

of the room to await transfer. When mice were returned to the holding room, the blue pads from

the cart were disposed of in a biohazard bag and the cart wiped with 10% bleach followed by

70% ethanol.

Environmental Monitoring of *C. auris* in Holding and Procedure Rooms by qPCR

In addition to biomonitoring the rooms by culture using the Sabouraud Dextrose Agar

plates, surfaces in both the holding and procedure rooms were tested after disinfection with 10%

bleach followed by 70% ethanol. Surfaces were swabbed using a sponge-stick (3M Health Care,

St. Paul, MN), the sponges placed back into their respective bags, and the handles removed. The

bag was folded 4-5 times and sealed, taking care that only the sponge touched the inside of the

bag during sampling. Surfaces were again wiped with 10% bleach followed by 70% ethanol to

remove residue left by the sponge-stick. The sponges were processed immediately but could be

stored at 4°C in the sealed bag for no more than a week. The protocol for sponge processing was

done as previously described by Leach et al., who developed the method for rapid identification

of *C. auris* by PCR.[26] Briefly, in a Whirl-Pak bag (Nasco, Fort Atkinson, WI), 45 ml of 1x PBS

with 0.02% of Tween 80 was added to the swabbing sponge and placed in a stomacher 400

circulator (Laboratory Supply Network, Inc., Atkinson, NH) for 1 min at 260 rpm. The liquid

was centrifuged at 4,000 rpm for 5 min and the supernatant was decanted, leaving approximately

2 ml of liquid in the bottom of the tube. One milliliter of sample was used for plating and the

other 1ml was used for DNA extraction. The sample was centrifuged at 13,000 rpm for 5 min

and the collected pellet was washed twice then resuspended in 48 µl of PBS-BSA. The 48 µl was

then transferred to a sterile screw cap tube (Heathrow Scientific, Vernon Hills, IL) containing

two 3 mm glass beads and 2 µl of bicoid inhibition control plasmid DNA (5×10^{-3} ng/µl).[26] The

tubes were then homogenized using a Vortex Genie in 10 min intervals followed by a 2-min

incubation on ice for a total of 50 min of homogenization. Samples were either run on qPCR

immediately, or stored at -20℃. The qPCR targets the internal transcribed spacer 2 (*ITS*2) gene

of *C. auris* as previously described.[26] The PCR also contains an internal bicoid inhibition

control.[26] Any samples that provided a Ct value were deemed positive. Samples that gave no Ct

value were found to be negative.

Culturing Samples Directly from Mice

Anal and oral swabs were taken from neutrophil-depleted mice infected with *C. auris* and

immune-competent control mice using Q-tips (Puritan, Guilford, ME). The Q-tips were placed in

labeled 15 mL tubes, and each was submerged in 1 mL 0.02% PBS-Tween-80 and vortexed for

10 seconds. The Q-tip was pressed against the sides of the tube as it was lifted out to remove as

much moisture as possible. The Q-tip was discarded in a biohazard bag. The liquid collected was

transferred to a labeled 2 ml screwcap tube. These samples were stored at $4\,°C$ for no longer than

one week before processing was continued. The collected liquid was cultured in three ways: (1)

50 µl of the liquid was plated on Sabouraud Dextrose Agar plates with antibiotics, (2) 50 µl of

the liquid was plated on Sabouraud Dulcitol Agar plates with 10% NaCl and antibiotics, and (3)

200 µl was inoculated in a Sabouraud Dulcitol Agar, 10% NaCl and antibiotics broth. Plates

were incubated at $40\,°C$ for 2 weeks. Broth cultures are incubated in a shaker at $40\,°C$ for 2 weeks.

Cultures that were positive were verified by PCR as previously described by Leach et al.[26]

RESULTS

Biomonitoring of *C. auris* in Animal Holding and Procedure Rooms

Due to the persistence of *C. auris* in hospitals and nursing home settings, coupled with its

resistance to antifungals, it was important to identify potential sources of exposure while

working with intravenously (i.v.) infected mice. Mice were infected with *C. auris* at a range of

10^5-10^8 cells. At concentrations of 10^8, *C. auris* was fulminant to neutrophil-depleted mice, as

these animals succumbed to infection in less than 24 hrs. At the lowest inoculum of 10^5 cells, *C.*

auris targets multiple organs such as the spleen, liver, kidney, stomach, heart, lungs, and brain.

Immune-competent mice exhibited 3.65×10^3 cells per organ for spleen, 7.85×10^1 cells per organ

for liver, 2.80×10^2 cells per organ for kidney, 1.07×10^3 cells per organ for heart, and 4.30×10^1

cells per organ for brain. No *C. auris* was detectable in the lungs and stomach. Neutrophil-

depleted mice exhibited 1.81×10^4 cells per organ for spleen, 1.12×10^2 cells per organ for liver,

5.78×10^2 cells per organ for kidney, 4.22×10^1 cells per organ for stomach, 1.56×10^3 cells per

organ for heart, and 1.70×10^3 cells per organ for lungs. No *C. auris* was detected in the brain

(Figure 4).

To assess the potential sources of *C. auris* contamination and exposure during the study,

both the animal holding and procedure rooms were monitored on a daily basis for five days.

Surfaces monitored for *C. auris* exposure were chosen based on their direct or indirect contact

with the infected mice and hands of lab personnel. Prior to doing any work with *C. auris,*

baseline environmental swabs were taken on selected surfaces **(Tables 1 and 2).** In the animal

holding rooms where routine cage changes were done, environmental monitoring was done by

swabbing the air flow nozzle of the micro-isolator cages, the floor near the biosafety cabinet, the

interior door handle, the exterior door handle, and the cart used to transport mice between the

holding and procedure rooms. The air flow nozzle of the mouse rack served as an indicator of

whether *C. auris* could be horizontally transmitted in the animal room, as has been described for

C. albicans.[28] The floor of the room, door knobs and transport cart were monitored to identify

potential cross-contamination that might have arisen from improper PPE use or clean up

procedure. After collection, swabs were analyzed via qPCR designed specifically for the

detection of *C. auris* DNA.[26] The results summarized in **Table 1** reveal that there was no *C.*

auris DNA detected at any time point in the holding room. To eliminate false negative results,

during the qPCR a bicoid plasmid was added to the samples as positive internal control.

As in the animal holding room, environmental monitoring of the procedure room was

also assessed for five days. Baseline swabs were also taken before any work with *C. auris* was

done in the room. Surfaces such as the isoflurane nose cone, floor near the biosafety cabinets,

chair backs, interior and exterior of the sink faucets and spigots, and interior and exterior door

handles were tested. The results, summarized in **Table 2,** reveal that there was no *C. auris* in the

animal holding room itself. However, samples of the cage bedding that held infected mice, as

well as the interior of the biosafety cabinet and exterior gloves right after animal necropsies all

tested positive at certain *C. auris* concentrations (**Table 3; Figure 5**). The results reveal that at

10^7, the mouse bedding and the exterior gloves were positive with Ct values of 29.52 and 28.36,

respectively. A 10^8 inoculum revealed that the mouse bedding, exterior gloves, and interior of the

biosafety cabinet were positive with Ct values of 28.90, 29.01, and 22.93, respectively,

suggesting that at these concentrations, mice begin to shed *C. auris* into the bedding. No

shedding into the bedding was found at inoculums of 10^5 and 10^6, [as] the PCR was negative.

Positive tests of the bedding at 10^7 and 10^8 inoculums provided the opportunity to test

whether autoclaving and disinfection were efficient. To remediate *C. auris* in the bedding, the

closed micro-isolator cages with water bottle attached were autoclaved at 121°C for 60 minutes.

Both culture and PCR results were negative, revealing that autoclaving is essential for cage

disinfection. At 10^7 and 10^8 inoculums, oral and anal swabs were taken from mice to determine if

these were the source of *C auris*. Swabs were cultured in three different media, including (1)

Sabouraud Dextrose Agar plates with antibiotics, (2) Sabouraud Dulcitol Agar plates with 10%

NaCl and antibiotics, and (3) Sabouraud Dulcitol, 10% NaCl and antibiotics broth. The cultures

did not show any growth at these concentrations after 2 weeks; therefore, no PCR was performed

on these swabs (data not shown).

During necropsies, the interior of the biosafety cabinet was only positive for *C. auris*

inoculums of 10^8 cells. The disinfection protocol of 10% bleach for five minutes followed by

70% ethanol was found to be effective, as the interior of the biosafety cabinet tested negative for

C. auris by PCR and by culture following this procedure. Glove exteriors during necropsies were

positive at 10^6, 10^7 and 10^8 concentrations, with Ct counts of 27.65, 28.36 and 29.01,

respectively, indicating a potential way to contaminate the procedure room. To easily remediate

this potential source of contamination, exterior gloves were removed right after the full necropsy

of each mouse was done and before the workers' hands touched anything else within the

biosafety cabinet. As an important reminder, the "buddy" always kept a watch for the removal of

exterior gloves.

Post-Six Month Biomonitoring using *C. auris* Inoculums Ranging from 10^5-10^8 Cells

To assess the effectiveness of the workflow and disinfection protocol, the procedure

room was assessed 6 months after initial studies began. Throughout this time, at least twenty

experiments were done at inoculums ranging from 10^5-10^8. Areas swabbed included the

isoflurane nose cone, sharps container, the floor under the biosafety cabinet, the back of the

chairs, the interior and exterior of the sink, the facets and spigots of the sink, the stockpot lid, and

the interior and exterior door handles, as well as the mouse restrainer. The results showed all

negative Ct values for all of the areas sampled, providing a strong indicator that the workflow

and disinfection protocol was effective. The results are summarized in **Table 4.**

DISCUSSION

As an MDR emerging pathogen, recent work in the *C. auris* field has focused on several

areas such as the development of rapid identification methods to prevent delayed treatment,[21, 26]

the epidemiology and genomic analysis of *C. auris*,[1] the testing of disinfectants and best

practices to reduce *C. auris* persistence, especially in healthcare settings,[10] and the testing of

small molecule antifungal drugs such as APX001 and the echinocandin rezafungin as alternative

therapeutics.[22, 23] Because *C. auris* is a new organism, there are a limited number of animal

models. The current animal models use immunosuppressive agents such as cyclophosphamide to

establish a *C. auris* infection.[22] However, to date no studies have evaluated the risk of *C. auris*

as an occupational exposure during animal experiments for researchers and animal care staff.[22]

As part of this study, rigorous safety procedures were developed, and *C. auris* was handled at

Animal Biosafety Level 2 (ABSL-2) with Animal Biosafety Level 3 (ABSL-3) containment

practices that included enhanced Personal Protective Equipment (PPE) and a strict no open

handling policy. A "buddy system" was also implemented in the workflow to prevent accidental

contamination of the room, as well as allowing researchers to maintain clearly defined clean and

dirty work areas. The "buddy system" was also a way for workers to monitor each other and

ensure that exterior PPE was changed during cage changes, infections and necropsies. Since

exterior gloves tested positive at inoculums of 10^6-10^8 cells, they were considered to be

important sources of contamination, emphasizing the importance of changing exterior gloves

even when moving within the biosafety cabinet. Another important part of the developed

procedures included keeping mice in micro-isolator cages, as *C. auris* was detected as positive at

10^7 and 10^8 inoculums. The closed micro-isolator cages were effective at keeping dirty bedding

isolated and contained water bottles that could be autoclaved as a single unit. This system

prevented exposure of animal caretakers to *C. auris* positive bedding when handling cages or if

cages fell to the floor prior to autoclaving. Autoclaving was also found to be highly effective in

the elimination of *C. auris* in the bedding. This study and others also found the use of 10%

bleach followed by 70% ethanol to be an effective disinfectant regimen against *C. auris*,[4] as

mice were clearly infected, yet all surfaces were qPCR-negative for *C. auris*.

CONCLUSIONS

Because there are currently a limited number of documented murine models of *C. auris*

infection, occupational exposure of working with these models has not been clearly defined. This

study tested both the animal holding room and procedure room and evaluated potential sources

of contamination and exposure during animal infections. Using an intravenous model of *C. auris*

infection, shedding of the organism was determined to be dose-dependent. *C. auris* was

detectable in the cage bedding when mice were infected with 10^7 and 10^8 cells, but not with

doses of 10^5 and 10^6 cells. Testing by anal and oral swabs on mice was unable to identify the

source of shedding. Results indicated potential for exposure with *C. auris* during necropsies and

when working with infected tissues.

To mitigate these potential exposures, laboratory personnel used two layers of PPE and

the lab developed a rigorous "buddy system" workflow. The "buddy's" role was to remind the

person working inside of the biosafety cabinet to change their exterior gloves, ensure that there

were no needlestick accidents during infection, and enable the person to move around the

procedure room and hand over cages and materials to the person working inside of the biosafety

cabinet. The rigorous use of 10% bleach and 70% ethanol and the use of closed micro-isolator

cages and cage autoclaving were all determined to be essential. Assessment of the procedure

room after six months of experiments using this workflow and disinfection protocol found

negative Ct values, indicating that these methods were successful. In conclusion, this study

showed that it is possible to work with large inoculums of *C. auris* in an animal facility, and that

by following these recommendations, it can be done safely without exposing the researchers and

animal caretakers to infection.

RECOMMENDATIONS

For any study using a *C. auris* infection models, it is recommended that investigators

handle the organism at Animal Biosafety Level 2 (ABSL-2) with Animal Biosafety Level 3

(ABSL-3) containment practices. In addition, researchers should wear two layers of protective

PPE, such as double gloves, booties, and lab coats. Also, the "buddy system" and a directional

workflow should be used, and micro-isolator cages should be used and autoclaved. To maintain a

clean animal holding and procedure rooms, surfaces should be decontaminated with 10% bleach

for five minutes followed by 70% ethanol. Following these recommendations will ensure a safe

laboratory environment for both the researchers and animal care takers.

FUNDING

M.D.J. was supported by the University at Albany and Wadsworth Center start-up funds. The

funders had no role in study design, data collection and interpretation, or the decision to submit

the work for publication. The authors declared no potential conflicts of interest with respect to

the research, authorship, and/or publication of this article.

ACKNOWLEDGMENTS

We thank Dr. Kimberly McClive-Reed for the critical reading of this manuscript. We also thank

Dr. Frank Blaisdell, Dr. Robert Keefe and Mrs. Deirdre Torrisi at the Wadsworth Center's

Veterinary Sciences Facility for their suggestions and special accommodations that allowed this

work to be possible.

REFERENCES

1   **Jeffery-Smith, A., S.K. Taori, S. Schelenz, K. Jeffery, E.M. Johnson, A. Borman et**

al.: Candida auris: a Review of the Literature. *Clin Microbiol Rev* 31(1)(2018).

2   **Satoh, K., K. Makimura, Y. Hasumi, Y. Nishiyama, K. Uchida, and H. Yamaguchi:**

Candida auris sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an

inpatient in a Japanese hospital. *Microbiol Immunol* 53(1): 41-44 (2009).

3   **Sears, D., and B.S. Schwartz:** Candida auris: An emerging multidrug-resistant

pathogen. *Int J Infect Dis* 63: 95-98 (2017).

4   **Ku, T.S.N., C.J. Walraven, and S.A. Lee:** Candida auris: Disinfectants and

Implications for Infection Control. *Front Microbiol* 9: 726 (2018).

5   **Berkow, E.L., and S.R. Lockhart:** Activity of novel antifungal compound APX001A

against a large collection of Candida auris. *J Antimicrob Chemother* (2018).

6   **Forsberg, K., K. Woodworth, M. Walters, E.L. Berkow, B. Jackson, T. Chiller et al.:**

Candida auris: The recent emergence of a multidrug-resistant fungal pathogen. *Med Mycol*

(2018).

7   **Calvo, B., A.S. Melo, A. Perozo-Mena, M. Hernandez, E.C. Francisco, F. Hagen et**

al.: First report of Candida auris in America: Clinical and microbiological aspects of 18 episodes

of candidemia. *J Infect* 73(4): 369-374 (2016).

8   **Chowdhary, A., C. Sharma, S. Duggal, K. Agarwal, A. Prakash, P.K. Singh et al.:**

New clonal strain of Candida auris, Delhi, India. *Emerg Infect Dis* 19(10): 1670-1673 (2013).

9   **Arauz, A.B., D.H. Caceres, E. Santiago, P. Armstrong, S. Arosemena, C. Ramos et**

al.: Isolation of Candida auris from 9 patients in Central America: Importance of accurate

diagnosis and susceptibility testing. *Mycoses* 61(1): 44-47 (2018).

10 **Lee, W.G., J.H. Shin, Y. Uh, M.G. Kang, S.H. Kim, K.H. Park et al.:** First three

reported cases of nosocomial fungemia caused by Candida auris. *J Clin Microbiol* 49(9): 3139-

3142 (2011).

11 **Magobo, R.E., C. Corcoran, S. Seetharam, and N.P. Govender:** Candida auris-

associated candidemia, South Africa. *Emerg Infect Dis* 20(7): 1250-1251 (2014).

12 **Emara, M., S. Ahmad, Z. Khan, L. Joseph, I. Al-Obaid, P. Purohit et al.:** Candida

auris candidemia in Kuwait, 2014. *Emerg Infect Dis* 21(6): 1091-1092 (2015).

13 **Schelenz, S., F. Hagen, J.L. Rhodes, A. Abdolrasouli, A. Chowdhary, A. Hall et al.:**

First hospital outbreak of the globally emerging Candida auris in a European hospital.

Antimicrob Resist Infect Control 5: 35 (2016).

14 **Vallabhaneni, S., A. Kallen, S. Tsay, N. Chow, R. Welsh, J. Kerins et al.:**

Investigation of the First Seven Reported Cases of Candida auris, a Globally Emerging Invasive,

Multidrug-Resistant Fungus - United States, May 2013-August 2016. *MMWR Morb Mortal Wkly*

Rep 65(44): 1234-1237 (2016).

15 **Ben-Ami, R., J. Berman, A. Novikov, E. Bash, Y. Shachor-Meyouhas, S. Zakin et**

al.: Multidrug-Resistant Candida haemulonii and C. auris, Tel Aviv, Israel. *Emerg Infect Dis*

23(1)(2017).

16 **Morales-Lopez, S.E., C.M. Parra-Giraldo, A. Ceballos-Garzon, H.P. Martinez, G.J.**

Rodriguez, C.A. Alvarez-Moreno et al.: Invasive Infections with Multidrug-Resistant Yeast

Candida auris, Colombia. *Emerg Infect Dis* 23(1): 162-164 (2017).

17 **Ruiz Gaitan, A.C., A. Moret, J.L. Lopez Hontangas, J.M. Molina, A.I. Aleixandre**

Lopez, A.H. Cabezas et al.: Nosocomial fungemia by Candida auris: First four reported cases in

continental Europe. *Rev Iberoam Micol* 34(1): 23-27 (2017).

18   **Wang, X., J. Bing, Q. Zheng, F. Zhang, J. Liu, H. Yue et al.:** The first isolate of

Candida auris in China: clinical and biological aspects. *Emerg Microbes Infect* 7(1): 93 (2018).

19   **Durante, A.J., M.H. Maloney, V.H. Leung, J.H. Razeq, and D.B. Banach:** Challenges

in identifying Candida auris in hospital clinical laboratories: a need for hospital and public health

laboratory collaboration in rapid identification of an emerging pathogen. *Infect Control Hosp*

Epidemiol: 1-2 (2018).

20   **Chowdhary, A., C. Sharma, and J.F. Meis:** Candida auris: A rapidly emerging cause of

hospital-acquired multidrug-resistant fungal infections globally. *PLoS Pathog* 13(5): e1006290

(2017).

21   **Kathuria, S., P.K. Singh, C. Sharma, A. Prakash, A. Masih, A. Kumar et al.:**

Multidrug-Resistant Candida auris Misidentified as Candida haemulonii: Characterization by

Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass Spectrometry and DNA

Sequencing and Its Antifungal Susceptibility Profile Variability by Vitek 2, CLSI Broth

Microdilution, and Etest Method. *J Clin Microbiol* 53(6): 1823-1830 (2015).

22   **Hager, C.L., E.L. Larkin, L. Long, F. Zohra Abidi, K.J. Shaw, and M.A.**

Ghannoum: In Vitro and In Vivo Evaluation of the Antifungal Activity of APX001A/APX001

against Candida auris. *Antimicrob Agents Chemother* 62(3)(2018).

23   **Larkin, E., C. Hager, J. Chandra, P.K. Mukherjee, M. Retuerto, I. Salem et al.:** The

Emerging Pathogen Candida auris: Growth Phenotype, Virulence Factors, Activity of

Antifungals, and Effect of SCY-078, a Novel Glucan Synthesis Inhibitor, on Growth

Morphology and Biofilm Formation. *Antimicrob Agents Chemother* 61(5)(2017).

24 **Hager, C.L., E.L. Larkin, L.A. Long, and M.A. Ghannoum:** Evaluation of the

efficacy of rezafungin, a novel echinocandin, in the treatment of disseminated Candida auris

infection using an immunocompromised mouse model. *J Antimicrob Chemother* (2018).

25 **Lepak, A.J., M. Zhao, E.L. Berkow, S.R. Lockhart, and D.R. Andes:**

Pharmacodynamic Optimization for Treatment of Invasive Candida auris Infection. *Antimicrob*

Agents Chemother 61(8)(2017).

26 **Leach, L., Y. Zhu, and S. Chaturvedi:** Development and Validation of a Real-Time

PCR Assay for Rapid Detection of Candida auris from Surveillance Samples. *J Clin Microbiol*

56(2)(2018).

27 **(CDC), C.D.C.:** "Environmental Disinfection." [Online] Available at

https://www.cdc.gov/fungal/candida-auris/c-auris-infection-control.html#disinfection, 2018).

28 **Cutler, J.E., M. Corti, P. Lambert, M. Ferris, and H. Xin:** Horizontal transmission of

Candida albicans and evidence of a vaccine response in mice colonized with the fungus. *PLoS*

One 6(7): e22030 (2011).

FIGURES

Figure 1: The "buddy system" work flow. A "buddy system" was implemented, such that one
person hands clean caging and other supplies to a second person performing the procedures
inside of the biosafety cabinet. This allows for a "clean" to "dirty" workflow with minimal
disruption of the laminar airflow of the biosafety cabinet.

Figure 2: Workflow for cage changes during *C. auris* experiments. The in-use micro-isolator
cage and an empty clean cage were placed inside of the biosafety cabinet atop the blue pads that
were sprayed with 10% bleach. Mice were carefully transferred to the new cage and food was
added to the new cages. The food stock was kept strictly kept inside of the biosafety cabinet.
Before removal from the biosafety cabinet, cages were wiped down with 10% bleach by the
individual inside of the cabinet, followed by 70% ethanol by the individual outside of the
cabinet. At the end of cage changes, the blue pads were rolled up and placed into a biohazard
waste bag. The biosafety cabinet and contents were sprayed down with a 10% bleach solution
and allowed to sit for 5 minutes and wiped. Additionally, the cabinet was cleaned with a 70%
ethanol solution.

Figure 3: Biosafety cabinet setup within the procedure room. (1) On the right-hand side of
the biosafety cabinet, mice were weighed within the chamber on the scale. Daily weights were
taken for all mice in the study before moving on to the next step. (2) Mice were euthanized in the
CO_2 chamber on the left-hand side, as the CO_2 source was only available on that side of the
biosafety cabinet. (3) Cardiac punctures were performed directly after CO_2 euthanasia. Procedure
needles were disposed of in the sharps container. Cervical dislocation was done as a secondary
method of animal euthanasia. (4) The mice were transferred to the folded blue mat on the right.
70% ethanol was sprayed on the mouse's abdomen prior to making a midline incision for organ
collection. Organs were collected and placed inside labeled tubes containing 2mL HBSS on the
organ tube rack. When necropsy was finished, the carcass was placed into a sealable biohazard
bag. Tools were rinsed in the following order: 10% bleach, water, then 70% ethanol between
each mouse. The folded blue mat was switched to a clean side between mice. Any waste
generated throughout necropsies was put into the biohazard bag in the back of the biosafety
cabinet. At the end of workflow, the blue pads were rolled up and placed in the biohazard waste
bag. The cabinet and all contents were sprayed with 10% bleach and allowed to sit for 5 minutes.
After 5 minutes all contents and the cabinet were wiped and then cleaned using 70% ethanol.

Figure 4. Organ fungal burdens in *C. auris* mouse model of infection. Organ fungal burden in
mice infected with 10^5 *C. auris* by tail vein injection. Neutrophil-depleted and control mice were
assessed 5 days post infection. Neutrophil-depleted mice were found to have additional fungal
burden in the lungs (black bar). Data represent mean cells per organ; n = 4 mice per group.

Figure 5. Autoclaving is an important step in disinfecting cage bedding. Neutrophil-depleted mice were infected with 10^8 *C. auris* by tail vein injection. (A) Before autoclaving, swabbed bedding is shown to have *C. auris* growth on Sabouraud dextrose agar plates, confirmed by PCR. (B) Bedding autoclaved in micro-isolator cages as a closed unit displays no *C. auris* growth, confirmed by PCR.

544 **TABLES**

	Target	Baseline	Day 1	Day 2	Day 3	Day 4	Day 5
Mouse Rack Nozzle	*C. auris*	(-)	ND	(-)	(-)	(-)	(-)
	Bicoid Plasmid	26.87	ND	28.16	28.46	26.20	26.51
Floor near BSC	*C. auris*	(-)	ND	(-)	(-)	(-)	(-)
	Bicoid Plasmid	28.03	ND	28.07	29.83	26.04	27.09
Interior Door handle	*C. auris*	(-)	(-)	ND	(-)	(-)	(-)
	Bicoid Plasmid	29.38	28.30	ND	29.54	29.07	27.74
Exterior Door handle	*C. auris*	(-)	(-)	ND	(-)	(-)	(-)
	Bicoid Plasmid	28.63	27.57	ND	27.88	29.23	27.88
Lab Cart 3-tier	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	27.07	27.58	28.93	28.37	29.26	29.01

546 **Table 1. Monitoring of Animal Holding Room by qPCR**

548 **Table 1.** Monitoring of the animal facility holding room where daily cage changes are done. Ct values
549 show (-) negative detection of *C. auris* DNA by qPCR. Bicoid positive internal control shows positive Ct
550 values. N.D. denotes not determined.

562 **Table 2. Monitoring of Animal Procedure Room by qPCR**

	Target	Baseline	Day 1	Day 2	Day 3	Day 4	Day 5
Isoflurane Nose Cone	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	26.49	29.07	28.78	28.27	28.53	29.20
Sharps Container	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	26.94	28.21	30.43	28.27	28.46	29.37
Floor near BSC	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	27.06	29.00	27.69	28.14	31.09	28.00
Chair Backs	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	26.96	29.27	27.51	27.89	30.70	27.33
Interior of Sink	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	28.56	29.10	28.05	28.87	30.56	26.18
Exterior of Sink	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	27.96	28.25	28.09	29.40	29.33	27.01
Faucets and Spigots	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	28.27	27.11	29.11	28.78	28.83	27.46
Stockpot Lid	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	28.31	26.79	28.40	28.12	29.03	28.63
Interior Door handle	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	26.18	27.82	27.67	27.89	29.60	28.50
Exterior Door handle	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	26.74	29.02	27.77	28.29	30.85	27.57
Mouse Water Bottle Tip	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	27.38	28.87	29.72	28.29	28.82	31.12
Exterior of Mouse Cage	*C. auris*	(-)	(-)	(-)	(-)	(-)	(-)
	Bicoid Plasmid	27.05	28.87	28.17	27.78	29.72	28.86

564 **Table 2.** Monitoring of the animal facility procedure room post necropsy shows (-) negative Ct values for
565 *C. auris* DNA by qPCR. Bicoid positive internal control shows positive Ct values. N.D. denotes not
566 determined.

572 **Table 3. Monitoring inside the biosafety cabinet at different inoculums by qPCR**

	Target	1×10^5	1×10^6	1×10^7	1×10^8
Mouse Bedding (Before Autoclave)	*C. auris*	(-)	(-)	(+) 29.52	(+) 28.90
	Bicoid Plasmid	28.38	27.66	26.8	27.42
Mouse Bedding (After Autoclave)	*C. auris*	ND	ND	ND	(-)
	Bicoid Plasmid	ND	ND	ND	24.93
Exterior Gloves (Before Disinfection)	*C. auris*	(-)	(+) 27.65	(+) 28.36	(+) 29.01
	Bicoid Plasmid	28.21	28.34	28.34	27.58
Exterior Gloves (After Disinfection)	*C. auris*	(-)	(-)	(-)	(-)
	Bicoid Plasmid	27.65	28.74	28.74	28.96
Interior of BSC (Before Disinfection)	*C. auris*	(-)	(-)	(-)	(+) 22.93
	Bicoid Plasmid	26.90	28.20	28.20	34.10
Interior of BSC (After Disinfection)	*C. auris*	(-)	(-)	(-)	(-)
	Bicoid Plasmid	28.90	27.93	27.93	27.36

574 **Table 3.** Monitoring within the biosafety cabinet within the procedure room post necropsy across several
575 concentrations shows (+) positive Ct values for *C. auris* DNA by qPCR. Monitoring within the biosafety
576 cabinet post-disinfection and post-autoclaving samples shows (-) negative detection of *C. auris* DNA by
577 qPCR. Bicoid positive internal control shows positive Ct values. N.D. denotes not determined.

Table 4: Monitoring of the procedure room six months later

	Target	Ct Value
Isoflurane Nose Cone	*C. auris*	-
	Bicoid Plasmid	27.89
Sharps Container	*C. auris*	-
	Bicoid Plasmid	27.8
Floor near BSC	*C. auris*	-
	Bicoid Plasmid	28.32
Chair Backs	*C. auris*	-
	Bicoid Plasmid	28.08
Interior of Sink	*C. auris*	-
	Bicoid Plasmid	29.88
Exterior of Sink	*C. auris*	-
	Bicoid Plasmid	27.41
Faucets and Spigots	*C. auris*	-
	Bicoid Plasmid	28.04
Stockpot Lid	*C. auris*	-
	Bicoid Plasmid	27.96
Interior Door handle	*C. auris*	-
	Bicoid Plasmid	28.64
Exterior Door handle	*C. auris*	-
	Bicoid Plasmid	27.34
Mouse Restrainer	*C. auris*	-
	Bicoid Plasmid	27.52

Table 4. Monitoring of procedure room six months after working with *C. auris* inoculums that ranged from 10^5-10^8 cells. All areas show (-) negative Ct values for *C. auris* DNA by qPCR. Bicoid positive internal control shows positive Ct values.