Title:

Mitogenomes illuminate the origin and migration patterns of the indigenous people of the Canary Islands

Authors:

Rosa Fregel[1,2], Alejandra C. Ordóñez[3], Jonathan Santana-Cabrera[4], Vicente M. Cabrera[2], Javier Velasco-Vazquez[5], Verónica Alberto[6], Marco A. Moreno-Benítez[6], Teresa Delgado-Darias[7], Amelia del Carmen Rodríguez-Rodríguez[5], Juan Carlos Hernández[8], Jorge Pais[9], Rafaela González-Montelongo[10], José M. Lorenzo-Salazar[10], Carlos Flores[10,11,12], M. Carmen Cruz de Mercadal[7], Nuria Álvarez-Rodríguez[8], Beth Shapiro[13,14], Matilde Arnay[3*] and Carlos D. Bustamante[1*]

*These authors contributed equally

Affiliations:

1. Department of Genetics, Stanford University, Stanford, California, United States of America.
2. Department of Biochemistry, Microbiology, Cell Biology and Genetics, Universidad de La Laguna, San Cristóbal de La Laguna, Spain.
3. Department of Prehistory, Anthropology and Ancient History, Universidad de La Laguna, San Cristóbal de La Laguna, Spain.
4. Department of Archaeology, Durham University, Durham, United Kingdom.
5. Department of Historical Sciences, Universidad de Las Palmas de Gran Canaria, Las Palmas de Gran Canaria, Spain.
6. Tibicena Arqueología y Patrimonio, Las Palmas de Gran Canaria, Spain.
7. Museo Canario, Las Palmas de Gran Canaria, Spain.
8. Museo Arqueológico de La Gomera, San Sebastián de La Gomera, Spain.
9. Museo Arqueológico Benahoarita, Los Llanos de Aridane, Spain.
10. Genomics Division, Instituto Tecnológico y de Energías Renovables, Granadilla, Spain.
11. Research Unit, Hospital Universitario N. S. de Candelaria, Santa Cruz de Tenerife, Spain.
12. CIBER de Enfermedades Respiratorias, Instituto de Salud Carlos III, Madrid, Spain

34   13. Department of Ecology and Evolutionary Biology, University of California Santa
35   Cruz, Santa Cruz, California, United States of America.
36   14. Howard Hughes Medical Institute, University of California Santa Cruz, Santa Cruz,
37   California, United States of America

Abstract

The Canary Islands' indigenous people have been the subject of substantial archaeological, anthropological, linguistic and genetic research pointing to a most probable North African Berber source. However, neither agreement about the exact point of origin nor a model for the indigenous colonization of the islands has been established. To shed light on these questions, we analyzed 48 ancient mitogenomes from 25 archaeological sites from the seven main islands. Most lineages observed in the ancient samples have a Mediterranean distribution, and belong to lineages associated with the Neolithic expansion in the Near East and Europe (T2c, J2a, X3a…). This phylogeographic analysis of Canarian indigenous mitogenomes, the first of its kind, shows that some lineages are restricted to Central North Africa (H1cf, J2a2d and T2c1d3), while others have a wider distribution, including both West and Central North Africa, and, in some cases, Europe and the Near East (U6a1a1, U6a7a1, U6b, X3a, U6c1). In addition, we identify four new Canarian-specific lineages (H1e1a9, H4a1e, J2a2d1a and L3b1a12) whose coalescence dates correlate with the estimated time for the colonization of the islands (1st millennia CE). Additionally, we observe an asymmetrical distribution of mtDNA haplogroups in the ancient population, with certain haplogroups appearing more frequently in the islands closer to the continent. This reinforces results based on modern mtDNA and Y-chromosome data, and archaeological evidence suggesting the existence of two distinct migrations. Comparisons between insular populations show that some populations had high genetic diversity, while others were probably affected by genetic drift and/or bottlenecks. In spite of observing interinsular differences in the survival of indigenous lineages, modern populations, with the sole exception of La Gomera, are homogenous across the islands, supporting the theory of extensive human mobility after the European conquest.

Introduction

The Canaries archipelago is located off the southern coast of Morocco (Figure 1). Due to their oceanic volcanic origin, they have probably never been connected to the continent. Mediterranean sailors discovered several groups of islands in the Atlantic Ocean in the 13th century, but only the Canary Islands were found to be inhabited by an indigenous population [1]. European chroniclers recorded that different islands were inhabited by populations exhibiting different ways of life and speaking distinct dialects of what they believed to be a Berber language. Ethno-historical sources provided ethnonyms for the native population of each island (e.g. Guanches for Tenerife, Benehaoritas for La Palma, and Bimbapes for El Hierro). However, for clarity, we will refer to them in general terms, as the Canarian indigenous or native population.

Chroniclers were amazed to discover that the Canarian natives were unaware of navigational methods and had remained isolated from the African continent [2,3]. During the 15th century, the Spanish kingdom of Castile gradually conquered all of the Canary Islands, after previous European attempts. In most of the islands, the indigenous people resisted the European conquest [4]. The crushing of the resistance, and subsequent European colonization, had a great impact on the indigenous people [5]. In spite of the abolishment of slavery on the Islands in 1498, a large number of natives were deported during and after the conquest [6]. Those that survived and stayed progressively mixed with the European colonizers, leading to the loss of indigenous culture and language.

The geographic origin of the Canarian indigenous people was initially inferred from both the interpretation of historical written sources and the analysis of archaeological evidence. Most archaeological and anthropological data support a North African origin for the Canarian indigenous people, relating to the Berber populations [7,8]. Key evidence supporting a Berber origin includes inscriptions belonging to the Libyco-Berber and Lybico-Canarian alphabets [9,10], pottery [11], communal granaries [12], and domestic species [13-15]. Non-metric dental traits [16-18] and morphological analyses of cranial and long bones [19,20] also show similarities between current inhabitants of Northwest Africa and the Canarian indigenous people.

In regards to the time of the arrival of the first population groups, some authors have proposed the first millennium BCE as the upper bound for human presence in the

archipelago [21], based on radiocarbon dating of charcoal and sediment samples. In

addition, there is evidence of a Roman short-stay settlement in Lobos islet dated during or

before the first centuries of the present era [22], which did not, according to the

archaeological data, involve attempting to colonize the Canaries. Recently, there has been

an effort to review and contextualize radiocarbon dates in the Canary Islands to better

assess the time of the archipelago's indigenous colonization. Accelerator mass

spectrometry (AMS) analyses support a later colonisation of the Canary Islands dating to

the outset of the first millennium AD. If only AMS analyses performed on short-lived

samples are considered [23], the earliest dates from the eastern islands of Lanzarote and

Fuerteventura range between 100-300 cal AD [21,22], whereas those from the central

island of Gran Canaria range between 400-500 AD [3]. The oldest AMS dates from

Tenerife are around 660-880 cal AD [24], while the western islands of La Palma, El

Hierro and La Gomera yield AMS dates ranging respectively between 260-450 cal AD

[24], 420-610 cal AD [24], and 120-330 cal AD [25]. On the other hand, older

radiocarbon dates that place the arrival of human populations before the 1st century BCE

were obtained from sediment, wood and charcoal samples that could be older than the

archaeological site where they were excavated.

Mitochondrial DNA (mtDNA) is a powerful tool for inferring the geographic

origin of populations [26]. MtDNA is maternally inherited, does not undergo

recombination and its different lineages are geographically structured in human

populations. For those reasons, mtDNA has been widely applied in phylogeographic

studies. The analysis of current Canary Islands samples using mtDNA has provided

support for a North African origin for the indigenous people, based on the presence of the

mtDNA U6 haplogroup [27], which has a clear Berber ascription [28,29]. Within the U6

lineages observed in the current Canary Islanders, it is worth mentioning U6b1a, a

haplogroup that is not present today in North Africa and which is considered a Canarian

autochthonous lineage [30]. Interestingly, U6b1a's coalescence age (3,600 years ago)

predates the proposed time of arrival of the first inhabitants of the islands, suggesting an

origin in North Africa [30]. Other haplogroups observed in the current Canarian people

have Eurasian (H, T, J…), sub-Saharan African (L1, L2 and L3) and Amerindian (A2 and

C1) affiliations [31]. These results highlighted the multiethnic nature of the modern

population of the Canary Islands, correlating with historical events, such as the
implementation of a slave workforce for the sugar cane plantations, or the commercial
connection with the Americas in the colonial period [32]. The detailed analysis of current
mtDNA of the modern Canary islanders has also suggested possible origins for the
indigenous population, including Morocco, Tunisia, Algeria or Sahara, but an overall
agreement has not yet been reached [31,33].

Regarding the colonization model, linguistic research has pointed to at least two
migration waves from North Africa [10,34]. Also, the observation of different cultural
backgrounds affecting the island of La Palma has been interpreted as evidence of
consecutive migrations. The specific timing for those migrations is still unclear, except
for La Palma, where the second wave of migration has been proposed to have taken place
around the 10^{th} century [7]. This idea has also been supported by asymmetrical
distribution of both mtDNA [31] and Y-chromosome lineages [35] in the modern
Canarian population. The first colonization wave may have affected the entire
archipelago, creating the substrate population and bringing mtDNA and Y-chromosome
haplogroups observed today in most of the islands, including the mtDNA lineages U6b1a
or H1cf. The second colonization would have brought new migrants to certain islands and
created an asymmetrical distribution of haplogroups, such as T2c1 and U6c1.

The direct analysis of ancient remains from the Canary Islands, using mtDNA by
means of PCR techniques, confirmed the presence of North African markers in the
indigenous people, including the U6b1a haplogroup, as well as some of the Eurasian
lineages observed in the modern population [36]. Admixture analysis based on mtDNA
data, using the natives as parental population, determined that 42% of modern Canarian
mtDNA lineages have an indigenous origin [36]. Ancient mtDNA results from four of the
seven islands found high diversities for Tenerife and La Palma [33,36,37], and the partial
and complete fixation of certain haplogroups in La Gomera [38] and El Hierro [39],
suggesting that the colonization of the archipelago was a heterogeneous process and that
different islands could have had different evolutionary histories.

Although previous ancient DNA (aDNA) studies have been fundamental to
understanding the origin and evolution of the Canarian population, most of the ancient
mtDNA data produced so far has been obtained using PCR amplification. This classical

aDNA technique has provided valuable information, but results have always been
hindered by the risk of sample contamination. This is due to the fact that aDNA from
warm climates is often extremely degraded and the PCR technique is highly sensitive,
thus minute amounts of modern contaminant DNA can be preferentially amplified [40].
Additionally, because the molecules are short and degraded, aDNA analyses based on
PCR amplification have tended to isolate small, but informative, regions of the
mitochondrial genome, such as the hypervariable region (HVR). This partial information
does not allow for refined classification within haplogroups, which is needed to
discriminate between close geographical regions. This is especially true within
haplogroup H, which comprises ~40% of the ancient Canarian mtDNA lineages. The
advent of next-generation-sequencing (NGS) has greatly expanded the capacity of aDNA
research. NGS allows damage patterns that are unique to aDNA, such as short fragment
size and post-mortem damage, to be detected easily, thus authenticating mtDNA results.
NGS also has the advantage of providing complete mtDNA genomes to allow a better
geographic assignment, compared to those obtained from partial HVR sequences.

A recent NGS study of the Canarian indigenous people presented the first
complete mtDNA genomes and low-coverage full genomes from this population, and,
more specifically, from the central islands of Tenerife and Gran Canaria [41]. However,
previous aDNA data [36-39] suggested that the indigenous populations from different
islands might have experienced different demographic processes. The inclusion of data
from all seven islands is therefore of paramount importance to accurately characterizing
the archipelago's indigenous population. Additionally, to fully benefit from the potential
of ancient mtDNA data, a more detailed phylogeographic analysis is required.

In order to obtain a comprehensive mtDNA perspective on the origin of the
indigenous people of the Canary Islands, we have applied aDNA protocols and NGS to
assemble ancient mtDNA genomes from all seven sub-populations. Since human remains
from warm regions like the Canary Islands are expected to have low endogenous DNA
content, we applied an enrichment technique [42] to improve mtDNA coverage and
reduce sequencing costs.

Methods

Sample collection

Samples were collected in collaboration with both Canarian universities, La Laguna (Tenerife) and Las Palmas de Gran Canaria (Gran Canaria), as well as the insular museums of Gran Canaria (El Museo Canario), La Palma (Museo Arqueológico Benahorita) and La Gomera (Museo Arqueológico de La Gomera). A total of 25 archaeological sites were selected for this project (Figure 1). Radiocarbon calibrated dates are available for several sites (Figure S1): El Agujero (1030 - 1440 cal AD), La Angostura (1318 - 1394 cal AD), Las Arenas (540 - 650 cal AD), El Capricho (400 - 480 cal AD), Cascajo (1640 - 1700 cal AD), Cuermeja (1270 - 1316 cal AD), La Fortaleza (599 – 633 cal AD), Guayadeque (540 - 737 cal AD), El Hormiguero (1020 - 1160 cal AD), Huriamen (1015 - 1050 cal AD; 1080 - 1150 cal AD), Lomo Galeón (1260 - 1290 cal AD), Montaña Mina (1313 - 1365 cal AD), El Pescante (150 - 350 cal AD), Portillo (1500 - 1580 cal AD), Puente de La Calzada (1265 - 1312 cal AD; 1358 - 1388 cal AD), Punta Azul (1015 - 1155 cal AD) and El Salitre (1060 - 1179 cal AD). For those sites with no available calibrated dates (Antoncojo, Barranco Majona, El Cedro, Cueva del Agua, El Espigón, Huerto de Los Morales, Los Pasitos and Salto del Casimiro), their assignation to the indigenous population was based on general context, the archaeological remains themselves and the presence of specific funerary practices. Sample CAN.005 is a tooth sample that was taken from a private collection of ancient human remains donated to El Museo Canario (Gran Canaria, Spain). Although this sample is not associated with any specific archaeological site, its calibrated radiocarbon date (1265 - 1312 cal AD) is in agreement with a pre-Hispanic origin. It is also worth mentioning that some archaeological sites from Tenerife (Cascajo and Portillo) are from the post-conquest period [43], but they are associated with the so-called "Alzados", indigenous people that rebelled against the European colonizers and retired to the mountains, leaving all contact with the Europeans behind [44].

DNA extraction and library preparation

Best-conserved samples were selected for DNA extraction. Although the petrous
bone is considered the best source for aDNA [45], we used teeth and small bones (e.g.
phalanx) to avoid destroying valuable archaeological material.

Required precautions were taken during the handling of samples, and all
experiments that included aDNA were carried out in dedicated, clean lab facilities at the
Paleogenomics Lab, University of California Santa Cruz, to avoid contamination. DNA
extraction was performed following Dabney et al. [46]. Bone samples were sanded to
remove the external surface, and then one bone piece was cut with a Dremel tool and
pulverized using a bone mill. The surface of tooth samples was decontaminated using a
bleach solution, and then the teeth were cut down the midline and the cementum drilled
using a Dremel tool and a metallic bit. Pulverized bone and tooth samples were incubated
overnight, using a proteinase K/EDTA solution, and DNA extracted using a silica-based
and guanidine method. Ancient DNA was then built into double-stranded libraries, with
7-bp single-index barcoding to allow for multiplexing sequencing, following Meyer and
Kircher [47]. Libraries were sequenced for an initial screening on an Illumina NextSeq
500 apparatus for obtaining paired-end shotgun data (~1 M reads per library) with a
sequencing read length of 2 x 75 bp.

Enrichment

After the screening of shotgun libraries, those samples with an endogenous DNA
content lower than 10% were enriched using whole-genome in solution capture [42].
Briefly, aDNA libraries were captured in singleplex reactions using human genomic
RNA baits, with the aim of increasing endogenous DNA rates and reducing sequencing
costs. Although this method is directed at capturing the whole genome, multicopy regions
of the mtDNA become particularly enriched. Post-capture libraries were sequenced as
indicated before, to obtain at least ~5 M reads per post-capture library.

HVR analysis

In order to perform population-based analyses, we included in our study
previously published [36-39] and unpublished HVR data from the seven islands. Newly

reported HVR data from the islands of Gran Canaria (n = 75), Lanzarote (n = 8) and

Fuerteventura (n = 10) was obtained following the methodology described by Maca-

Meyer et al. [36] and Ordóñez et al. [39]. Briefly, after external decontamination, tooth

samples were extracted by means of a GuSCN-silica based protocol. MtDNA

quantification was performed on a 7500 Real Time PCR system (Applied Biosystem,

Foster City, CA, USA), using a human-specific mtDNA fluorescent probe [48], and

~3,000 copies were submitted to PCR with the aim of reducing the effects of DNA

damage. The mtDNA HVRI (from positions 16,000 to 16,400) was amplified using seven

overlapping fragments, with sizes ranging from 82 to 124 bp, to improve the

amplification of endogenous DNA. All the sequencing reactions were prepared with the

BigDye v3.1 Terminator Cycle Sequencing kit (Applied Biosystems) and run on an ABI

PRISM 3130xl Genetic Analyzer (Applied Biosystems). Standard contamination

prevention and monitoring were conducted as described earlier [39].

262 Modern mtDNA genomes

We included in this study several current Canary Islands mtDNA genomes,

analyzed using both whole-genome and Sanger sequencing. Complete genomes were

obtained in Instituto Tecnológico y de Energías Renovables (ITER) by whole-genome

sequencing from a set of 18 unrelated Canarians. Briefly, DNA samples were processed

with a Nextera DNA Prep kit, with dual indexes following the manufacturer's

recommendations (Illumina Inc., San Diego, CA). Library sizes were checked on a

TapeStation 4200 (Agilent Technologies, Santa Clara, CA) and their concentration

determined by the Qubit dsDNA HS Assay (Thermo Fisher, Waltham, MA). Samples

were sequenced to a depth of 30X on a HiSeq 4000 instrument (Illumina) with paired-end

150-base reads. Sanger sequencing mtDNA genomes were obtained at University of La

Laguna following previously published methodologies [49], for samples classified as

T2c1 (determined by HVRI analysis). These samples were selected because of their

potential to define new sub-lineages within T2c1. Institutional review board approval for

the analysis of human subjects was obtained from Stanford University.

278 Data analysis

Mapping and filtering of ancient mtDNA reads

Shotgun sequencing reads were trimmed and adapters removed using AdapterRemoval version 1.5.4 [50]. Specifically, the paired-end reads were merged, and low-quality bases (BASEQ < 20) and short reads (< 30 bp) removed. Merged trimmed reads were then mapped to the human reference genome (hg19) using BWA version 0.7.12 [51], while unmerged reads were discarded. Unmapped, low-quality (MAPQ<30) and duplicate reads were removed using SAMtools version 0.1.19 [52]. The percentage of endogenous DNA was calculated by dividing the number of reads remaining after filtering by the total number of trimmed reads.

Authentication

Damage patterns were assessed using MapDamage v2.0 [53]. Insert size of libraries was obtained with SAMtools mpileup, and plotted using R software v.3.2.0 [54]. Contamination rates of libraries were calculated using contamMix v.1.0-10 [55] and Schmutzi [56].

Analysis of complete mtDNA genomes

MtDNA reads were directly mapped to the revised Cambridge Reference Sequence (rCRS) [57] and filtered as described before. MapDamage was used to rescale the quality of bases likely affected by post-mortem damage. Indel Realigner from the GATK pipeline version 2.5.2 was also used for improving alignment quality around indels [58]. MtDNA consensus sequences were obtained using SAMtools and BCFtools version 0.1.19 [52]. A list of variants was then obtained using SAMtools mpileup, with a minimum depth of 5. Haplogroups were determined with HaploGrep version 2.0 [59], using PhyloTree build 17 version (http://www.phylotree.org) [60]. MtDNA haplotypes were manually curated by visual inspection, using Tablet v.1.17.08.17 [61]. Modern DNA sequencing data was analyzed following the same protocol used for ancient samples, except for the MapDamage rescaling step. After retrieving all available mtDNA genomes belonging to the haplogroups of interest from NCBI (http://www.ncbi.nlm.nih.gov), phylogenetic trees were built using median-joining networks [62]. Indels around nucleotides 309, 522, 573 and 16193, and hotspot

mutations (e.g. 16519) were excluded from phylogenetic analysis. For estimating

coalescence ages for specific clades, we used the ρ statistic [63]. We established a

mutation rate for the complete mtDNA sequence of one substitution in every 3,624 years,

correcting for purifying selection as in Soares et al. [64]. Accompanying standard errors

were calculated as per Saillard et al. [65]. For highly frequent haplogroups, such as H1cf

and T2cd3, we only kept one sample per site, to avoid relatedness interfering with

coalescence age estimations.

Analysis of HVRI data

Newly reported HVR sequences were analyzed using BioEdit software v.7.0.9.0

[66], and haplotypes were obtained by means of HaploSearch software [67] and further

confirmed by manually inspecting the electropherograms. Haplogroup nomenclature was

assessed following the most updated mtDNA phylotree (Build 17) [60].

Genome-wide data was combined with previous HVRI sequencing data to

perform population-based analysis. Published samples used for comparisons are detailed

in Table S1. As we do not know if samples in the same burial can be related, when

several samples with the same haplogroup were observed from the same archaeological

site, only one was included in the analysis. Two-tailed Fisher's exact test was used to

assess differences in mtDNA haplogroup frequencies between eastern and western

islands. Gene diversity was calculated according to Nei [68]. Distances between

populations were estimated using haplogroup frequency-based linearized F_{ST} [69] as in

Arlequin v.3.5 [70]. Multidimensional Scaling (MDS) was performed using R software

and the "smacof" package [71]. Admixture estimates were calculated with the

WLSAdmix program [72], which was kindly provided by Dr Jeffrey Long.

Results and Discussion

The average endogenous DNA content for the Canarian indigenous samples is

7.92%, a relatively high value considering the warm and humid environmental conditions

of the archipelago (Table S2). However, endogenous DNA values varied within and

between archaeological sites, ranging between 0.02% and 39.0% (IQR= 0.67% - 11.5%).

All samples meet the standard aDNA authentication criteria, including observation of
DNA fragmentation and damage patterns at both ends of molecules, and low modern
DNA contamination rates (Figure 2). Those contamination rates calculated with
contamMix are larger than those produced with Schmutzi. One possible reason is that
contamMix estimations are more sensitive to low coverage values (Table S2). For
example, sample CAN.033, with a 7.9X mtDNA coverage, has a contamination rate of
10.2% based on contamMix and 1.0% on schmutzi. Schmutzi has been reported to be
able to obtain accurate contamination rates for coverage down to ~5X [56]. However, in
other cases, variable contamination estimations do not seem to be related to low
coverage, and other factors may be interfering.

After capture, we obtained complete mtDNA genomes from 48 ancient human
remains sampled in 25 different archaeological sites (Table S2). Our sample set covers
the entire archipelago and a time span of 1,200 years (Figure S1). The average mtDNA
depth is ~140X, with a minimum value of 8X (Table S2). Observed haplogroups agree
with previous studies [33,36-39,41], indicating the presence of North African (U6),
Eurasian (H, J2, T2 and X) and sub-Saharan African lineages (L1 and L3) in the Canarian
indigenous population (Figure S2). As delineated before [36], the majority of
haplogroups observed are of Eurasian origin, most with a Mediterranean distribution.
This result is expected, as recent aDNA data from North Africa has indicated the
presence of Neolithic European lineages as early as the Late Neolithic period (~5,000
BP) [73].

We also obtained complete mtDNA genomes from a set of 18 modern Canarians
(Figure S3). More than 50% of the samples belong to haplogroup H, with a higher
diversity of sub-haplogroups than the one observed in the indigenous population. In
addition to H1cf and H1e1a, we observe other H1 sub-lineages and other branches, such
as H6a1, H3c2 or H43, which are most likely of European origin. Other haplogroups
present in the indigenous people are also observed in the modern population, including
J2a2d, U6b1a and X3a. In line with previous analyses [27,31], a sub-Saharan African
(L3d1b3a) [74] and an Amerindian lineage (A2) [75] are observed in the current
population of the Canary Islands. Assuming that our set of 48 ancient genomes is
representative of the native population, we performed a rough admixture estimate of

27.8% of maternal lineages in members of the present-day population possessing

indigenous origins, while 61.1% would be of European ascription (Figure S3).

Population-based analysis

In order to compare our samples to previously published data, we combined the

newly generated mtDNA genomes with HVRI data from the Canarian indigenous

population (Table S1) [33,36-39]. Given that sample sizes for Lanzarote and

Fuerteventura are small and their indigenous populations are considered to be similar

based on archaeological data [76], these data sets were pooled together. It is worth

mentioning that those samples for which mitochondrial data were generated, using both

classical techniques and NGS sequencing, produced identical HVRI haplotypes, proving

our PCR-based approach generates authentic results.

As previously observed, the indigenous populations of the Canary Islands in the

past were not homogenous (Table 1; Figure 3). The islands of La Palma and Tenerife

show a relatively diverse mtDNA composition (>70%) [33,36,37], while the others show

signs of genetic drift and/or diversity reduction events, such as a bottleneck or a founder

effect. In La Gomera, mtDNA diversity was 54.2%, due to the high frequency of

haplogroup U6b1a [38], while in El Hierro, this value was 2.9%, with the almost

complete fixation of H1cf haplogroup in the Punta Azul site [39]. With new data on the

indigenous population of Gran Canaria, Lanzarote and Fuerteventura (Table S3), we

show that Gran Canaria had high mtDNA diversity, similar to Tenerife and La Palma,

while Lanzarote and Fuerteventura had low diversity (51.1%) because of the high

frequency of H*(xH1cf, H4a1a) lineages. These findings emphasize that results obtained

from the larger islands of Tenerife and Gran Canaria should not be extrapolated to the

entire archipelago. Estimations of population sizes during pre-colonial times based on

archaeological evidence agree with mtDNA results. Populations in Gran Canaria,

Tenerife and, to a lesser degree, La Palma, were large and able to sustain relatively high

diversity, while Lanzarote, Fuerteventura and El Hierro were almost depopulated at the

time of the conquest [77]. In the case of La Gomera, the population size was also reported

to be small [78].

By directly comparing the mtDNA types found in the indigenous population of
each island, we observe that H4a1e, L3b1a, U5 and U6c haplogroups are present only in
the eastern islands (Gran Canaria, Lanzarote and Fuerteventura). Differences between
eastern and western islands were shown to be significant for the four haplogroups, when
all the ancient samples were considered: H4a1 (p=0.0127), L3b1a (p=0.0037), U5
(p=0.0114) and U6c (p=0.0012). Though also present in the western islands, haplogroups
T2c1 (p=0.0164) and U6a (p=0.0028) appeared more frequently in the eastern islands.
However, these results can be artifacts caused by the high frequency of H1cf in El Hierro
and U6b1a in La Gomera. After removing these two populations from the western group,
only differences in the distribution of U6c remained significant (p=0.0328).

In contrast with the heterogeneity we observe in pre-Hispanic times, mtDNA
haplogroup frequencies in modern populations of the Canary Islands are homogenous
(Figure 3), with the sole exception of La Gomera [27]. The high frequency of haplogroup
U6b1a observed in the indigenous population of La Gomera is also detected in its
present-day population [38]. However, the same pattern is not observed for El Hierro. In
pre-colonial times, H1cf was almost fixated in El Hierro [39], while the frequency of this
haplogroup today is 4.6%, not significantly different from the average 2.4% observed in
the entire archipelago (p=0.2364).

In order to determine the admixture pattern at an insular level, we compared
modern Canarian samples with their principal parental populations: indigenous people,
Iberians, and sub-Saharan Africans (Table S1). Global admixture estimations using the
new mtDNA dataset (Table 2) confirm previous results on the survival of native lineages
in the modern population (55.9%). However, we observed that results within islands are
variable. When the miscellaneous ancient sample is used as one of the parental
populations, indigenous contribution to the modern population ranges from 30.8% in
Gran Canaria to 71.4% in La Gomera. However, this approach is not correct, as we know
that the indigenous population of the archipelago was heterogeneous and mtDNA
frequencies were variable. With our new data, we were able to estimate admixture, using
aDNA sampled directly from each island. Indigenous mtDNA contribution estimates are
lower when a direct comparison is performed, with values ranging from 0% in El Hierro
to 55.5% in La Gomera (Table 2). The extreme result observed in El Hierro is evidently

due to the marked difference between the ancient and current people. It is interesting that,
when the miscellaneous sample is used, the indigenous contribution increases to 36.2%.
This result is reasonable, given that the present-day sample from El Hierro is not
significantly different from other islands. This can be explained if we consider that El
Hierro was almost depopulated at the time of the European conquest [79]. In fact, it was
recounted in the chronicles that the indigenous population of El Hierro was decimated
due to razzias (raids for the purpose of capturing slaves) at the time of the Spanish
conquest, and was later repopulated with indigenous populations from other islands and
European colonizers [80,81].

To determine if a more specific origin for the Canarian indigenous population
could be ascertained, the ancient mtDNA sample was combined with a reference modern
DNA database containing samples from the Canary Islands, Europe, North Africa, Sub-
Saharan Africa and the Near East (Table S1). In the MDS analysis (Figure 4), the
indigenous sample from El Hierro and the indigenous and modern samples from La
Gomera act as outliers, due to the high frequency of H1cf and U6b1a, respectively.
When the two outliers were removed and all the remaining ancient samples were pooled
together, the first dimension differentiates sub-Saharan populations from Eurasian
populations, including North Africa and the Canary Islands. The second dimension places
Canarian and European/Near Eastern populations on both ends, with North Africans in an
intermediate position. The closest North African sample to the Canarian indigenous
population in the second dimension is West Sahara. However, the ancient sample is
differentiated from all current North African populations and placed closer to modern
Canarians. This is due to the fact that haplogroups occurring frequently in the Canarian
indigenous and current samples (e.g. U6b1a) are not present or appear in low frequencies
within the reference populations. This result concurs with later demographic processes
reshaping the mtDNA landscape of North Africa, and/or founder effects and isolation in
the Canary Islands. It is interesting that, compared to the other islands, the modern
populations of Tenerife and Gran Canaria are closer to the European populations. This
result is expected, because they each have capital cities of the two Canarian provinces
and, thus, have received substantial historical migration from the mainland.

Phylogeographic analysis of mitogenomes

The HVRI has been proven to be of limited value in providing a clear picture of
the origin of the indigenous people of the Canary Islands. In order to conduct a better
assignment of the geographic origin of the maternal Canarian indigenous lineages, we
performed detailed phylogeographic analysis of all the lineages observed in the aDNA
dataset (Figure S2), including those from Rodriguez-Varela et al. [41]. For detailed
information on our phylogenetic analysis, see Supplementary Text.
469

We observe five different H sub-lineages in the indigenous people of the Canary
Islands: H1cf, H1e1a9, H2, H3 and H4a1e. H1cf (Figure S4) seems to be restricted to
both the Canary Islands and Central North Africa, and shows a coalescence age (~3,400
years ago) that is in agreement with a continental origin before the colonization of the
islands (Figure S5). Newly defined haplogroups H1e1a9 (Figure S6) and H4a1e (Figure
S7) are both restricted to the Canary Islands, with a distribution similar to that observed
for U6b1a. However, in this case, H1e1a9 and H4a1e coalescence ages overlap with the
human occupation period (Figure S5) and are compatible with an origin in the islands.
The presence of lineages derived of H1e1a and H4a1 in both European Neolithic and the
Canary Islands indigenous samples corresponds with Eurasian prehistoric intrusions in
North Africa (Fregel et al. 2018). Two samples were classified within basal H2 and H3
haplogroups, preventing further phylogenetic analysis.

Two sublineages of haplogroup J are observed in the indigenous population of the
Canary Islands: J1c3 and the newly defined J2a2d1a1. J1c3 is present in Europe, North
Africa and the Near East, and more interestingly, in ancient Neolithic samples from Spain
and Sardinia (Figure S8). Although J2a2d1a* has been spotted in Central North Africa,
subhaplogroup J2a2d1a1 is exclusive to the Canary Islands and Brazil, the latter
representing an area with known historical migrations from the islands (Figure S9).
Accordingly, this new autochthonous Canarian lineage has a coalescence age that
overlaps with the indigenous occupation of the islands (Figure S5).

Phylogenetic analysis of the Canarian T2 sequences places them within T2b and
T2c1d, two haplogroups thoroughly observed in Neolithic and Bronze Age sites from
Europe. The inclusion of ancient and modern Canarian samples allows us to define four
new T2c1d subhaplogroups (Figure S10). T2c1d3 haplogroup is present in both Tunisia

and the Canary Islands. T2c1d1c1 and its two subclades (T2c1d1c1a and T2c1d1c1b) are

present in both North Africa and the current population of the eastern Canary Islands.

This distribution could be explained by an asymmetrical migration pattern, or, given its

absence in the indigenous people, by a higher impact of Moorish slave trade in the

eastern islands (Supplementary Text).

We identify several indigenous samples within macrohaplogroup L, belonging to

L1b1a and the newly defined L3b1a12. Although Later Stone Age [82], and Early and

Late Neolithic [73] samples from North Africa did not show any mtDNA lineage of sub-

Saharan origin, our results imply the presence of L1b and L3b1a in North Africa at the

time of the colonization of the Canary Islands. Regarding L3b1a12 (Figure S11), this

lineage can also be considered autochthonous of the Canary Islands, with a coalescence

age posterior to the proposed colonization date (Figure S5). Interestingly, this lineage was

only present in the eastern islands in ancient times, but has a wider distribution at the

present time, suggesting extensive movement of native people after the conquest.

Canarian indigenous sequences belonging to X haplogroup are classified within

the X3a clade (Figure S12). This lineage is present both in Europe, the Near East and

northeast Africa, as well as in the ancient and current populations of the Canary Islands.

Finally, several U6 sublineages are observed in the indigenous population of the

Canary Islands, including U6a1a1 (Figure S13), U6a7a1 (Figure S14), U6b1a (Figure

S16) and U6c1 (Figure S16). U6a1a1, U6a7a1 and U6c1 are present in the Maghreb,

southern Europe and the Canary Islands, and are most probably related to prehistoric

Mediterranean expansions (Figure S13, Figure S14 and Figure S16). As reported before,

the Canarian autochthonous U6b1a is also present in regions with recent Canarian

migration, including mainland Spain and Cuba (Figure S15). Given its coalescence age

and the oldest calibrated radiocarbon dates from human remains from the Canary Islands

(Figure S5), U6b1a most probably originated in North Africa and later migrated to the

Canaries. However, to date, this lineage has not been observed in the continent,

indicating the migrations occurred after the colonization of the Canary Islands reshaped

the North African mtDNA landscape.

Discussion

Our mtDNA results on the indigenous people of the Canary Islands shed light on the prehistory of North Africa. Our data are in agreement with recent aDNA data from Morocco [73] and further evidence of a complex pattern of Mediterranean migrations in North Africa. Archaeological records in the Maghreb support this result, and also suggest further European intrusions during the Chalcolithic and Bronze Age eras [83,84]. Additionally, Phoenicians, Carthaginians and Romans arrived in the North African region in historical times [85-88]. The presence of haplogroups of Mediterranean distribution in the indigenous people of the Canaries demonstrates the impact of these prehistoric and historical migrations in the Berbers and that they were already an admixed population at the time of the indigenous colonization of the islands [89].

In our phylogeographic analysis of complete mtDNA sequences from the Canarian indigenous population, we found lineages that are only observed in Central North Africa and the Canary Islands (H1cf, J2a2d and T2c1d3), while others have a wider distribution including both West and Central North Africa, and, in some cases, Europe and the Near East (U6a1a1, U6a7a1, U6b, X3a, U6c1). These results point to a complex scenario, where different migration waves from a dynamic and evolving North African population reached the islands over time. Every island experienced their own evolutionary path, determined by the environmental conditions and limitations of insularity. Those islands with the capability of sustaining large populations retained variability, while others with more restricted means (La Gomera and probably El Hierro) had to develop cultural practices to avoid inbreeding, like mandatory exogamic practices [78,90].

Although the North African Berber origin is the most widely accepted hypothesis, other lines of research have proposed that certain funerary practices and religious beliefs observed in the indigenous population of the Canary Islands could be linked to Punic-Phoenician influence [91], thus proposing the colonization of the Canary Islands as the result of Phoenicians expanding their control to the Atlantic Ocean. Based on the limits of the territorial occupation of the Atlantic West Africa by Phoenicians, Carthaginians and Romans, most researchers consider it unlikely that there was a political occupation or economic exploitation of the archipelago [92-94]. However, the islands were not

unknown to Mediterranean cultures, and Romans possessed the seafaring skills needed to
travel to the islands [22]. Some authors think Phoenicians also had the navigational
technology required to reach the Canary Islands [95,96], although this idea has been
challenged [97]. The first Phoenician aDNA sample published was a complete mtDNA
sequence of a child from Carthage dated to the 6[th] century BC [98]. This Carthaginian
sample was classified within U5b2c1 haplogroup. This result is interesting, given that U5
was more frequent in the indigenous population of the eastern islands, including the
island of Lanzarote, where a Punic-Phoenician influence has been proposed. As U5
haplogroup was not uncommon in Neolithic European samples, and its presence in North
Africa might be due to prehistoric migrations, an alterative explanation would be that
haplogroup U5 was incorporated into the Berber mtDNA pool before the Carthaginians
were established in Tunisia. Recently, Matisoo-Smith et al. [99] published thirteen
complete mitogenomes from Punic-Phoenician samples from Lebanon and Sardinia. The
only haplogroups in common with the indigenous population of the Canary Islands are
H3 and H1e1a, although, in this case, the Phoenician H1e1a sample is classified within
the sub-lineage H1e1a10. The lack of overlap between the mtDNA composition of
Phoenicians and the Canarian indigenous people disagrees with either a Punic-Phoenician
origin for the ancient islanders or sustained contact between the two populations.

Previous genetic analyses of the modern Canarian population detected an
asymmetrical distribution of maternal and paternal lineages in the archipelago [31,35].
Our aDNA results confirm the existence of asymmetrical distribution of mtDNA
haplogroups in pre-colonial times, with the presence of haplogroups H1e1a9, H4a1e,
L3b1a12 and U6c1 only in the eastern islands. However, it is worth mentioning that La
Palma, the island with the most anthropological evidence of two migrations waves, does
not show any of these lineages. If we consider the presence of H1e1a9, H4a1e, L3b1a12
and U6c1 haplogroups to be the result of further population movements from North
Africa to the eastern islands, we could approximate the date based on radiocarbon dates
of the sites where the sample was taken. Most sites where these lineages have been
observed have radiocarbon dates placed around the 13[th] century, and all except one are
from after the 10[th] century. The only site with an older date is Guayadeque; however, we
have to take into account that this is a large site, with evidence of human occupation

extending until the 14[th] centuries AD [100], and the dating was not performed directly on
the analyzed sample.

Archaeological data has evidenced significant changes in the productive strategies
of some islands around the 11[th] - 12[th] centuries [12,76,101-103]. In fact, recent data
indicates probable population growth in Gran Canaria at that time, suggesting the
appearance of new settlements associated with an exploitation model that intensified the
use of marine resources, the increase in the size of settlements linked to agricultural
nuclei, and changes in the production of some craftsmanships [12,104,105]. These
changes have been interpreted as part of an endogenous process, as it has been
determined that this population growth involved neither significant changes in the
structure of human settlements or burials, nor introduced differences in land management
or the types of domestic species that were exploited. However, it is also possible to
explain those changes as the result of the arrival of new migrants to the island of Gran
Canaria. Although it is still under study, there is evidence for transformations in the
configuration of some settlements in Lanzarote, between the 8[th] and 13[th] centuries [106].
Again, these modifications could be reflecting changes in the conception of domestic
space due to an endogenous process, or associated with the arrival of new colonizers.
Archaeological information from Fuerteventura is not abundant enough to determine
population size changes that could be related to the arrival of new migrants. Nevertheless,
it is clear from the archaeological record that Fuerteventura and Lanzarote maintained
frequent contact and shared both cultural and economic elements [76,107]. Future
paleogenomic efforts to obtain high-coverage genomes from all seven islands, in
combination with proper archaeological contextualization of the genetic data and detailed
radiocarbon dating, will be essential for improving our knowledge of the origins and
evolution of the indigenous population of the Canary Islands.
612

613 **DATA AVAILABILITY:**
614

615   Mitochondrial DNA sequence data are available through the European Nucleotide
616 Archive (PRJEB29569). Consensus mtDNA sequences are available at the National
617 Center of Biotechnology Information (Accession Numbers MK139577 - MK139649).

618 Requests for additional materials should be addressed to R.F. (e-mail:
619 rfregel@ull.edu.es).
620
621 **ACKNOWLEDGMENTS:**
622

C.D.B. and R.F. were funded by a grant from the National Science Foundation (1201234)
and the Chan Zuckerberg Biohub Investigator Award; R.F. was funded by a Fundación
Canaria Dr. Manuel Morales Fellowship and by a grant from Dirección General de
Patrimonio Cultural del Gobierno de Canarias (MITOCAN); B.S. was funded by a grant
from the Gordon and Betty Moore Foundation (GBMF-3804); R.G.M., J.M.L.S and C.F.
were funded by grants from the Spanish Ministry of Science and Innovation (RTC-2017-
6471-1) and Área Tenerife 2030 from Cabildo de Tenerife (CGIEU0000219140), and
from the agreement OA17/008 with Instituto Tecnológico y de Energías Renovables
(ITER). Finally, C.F. wants to acknowledge technical assistance from Ana Díaz-de-
Usera.

REFERENCES

1. Farrujia de la Rosa J (2014) An archaeology of the margins: colonialism, amazighity and heritage management in the Canary Islands. New York: Springer.

2. Navarro JF (2005) Todo sobre Canarias: Los aborigenes. Santa Cruz de Tenerife: Centro de la Cultura Popular Canaria.

3. Velasco-Vázquez J (2015) Más allá del horizonte: Una "perspectiva humana" del poblamiento de Canarias. In: Farrujía de la Rosa J, editor. Orígenes Enfoques interdisciplinares sobre el poblamiento indígena de Canarias. Santa Cruz de Tenerife: Thesaurus Arqueología, Ediciones Idea. pp. 23-89.

4. Rumeu de Armas A (2006) La conquista de Tenerife (1494-1496). La Laguna: Instituto de Estudios Canarios.

5. Crosby AW (1988) Imperialismo ecologico. La expansion biologica de Europa 900-1900. Barcelona: Ed. Critica.

6. Suarez JJ, Rodriguez F, Quintero CL (1988) Historia popular de Canarias. Vol. 2. Conquista y Colonizacion: Centro de la Cultura Popular Canaria.

7. Navarro JF (1997) Arqueologia de las Islas Canarias. Espacio, tiempo y forma Serie I, Prehistoria y Arqueologia 10: 447-478.

8. Tejera-Gaspar A (2006) Canarias y el África Antigua. San Cristóbal de La Laguna: Centro de la Cultura Popular de Canarias.

9. Springer R (2014) Die libysch-berberischen Inschriften der Kanarischen Inseln in ihrem Felsbildkontext; Stroomer H, editor. Koln: Rudiger Koppe Verlag.

10. Tejera-Gaspar A, Perera-Betancort MA (2011) Las supuestas inscripciones púnicas y neopúnicas de las Islas Canarias. Spal Revista de Prehistoria y Arqueología 20: 175-184.

11. Navarro-Mederos J (1999) «El viaje de las loceras»: la transmisión de tradiciones cerámicas prehistóricas e históricas de África a Canarias y su reproducción en las islas. Anuario de Estudios Atlánticos 45: 61-118.

12. Morales J, Rodriguez-Rodriguez A, González-Marrero MC, Martín-Rodríguez E, Henríquez-Valido P, del Pino M (2014) The archaeobotany of long-term crop storage in northwest African communal granaries: a case study from pre-Hispanic

Gran Canaria (cal. ad 1000–1500). Vegetation history and archaeobotany 23: 789-
804.

13. Hagenblad J, Morales J, Leino M, Rodriguez-Rodriguez A (2017) Farmer fidelity in
the Canary Islands revealed by ancient DNA from prehistoric seeds. Journal of
Archaeological Science 78: 78 - 87.

14. Ferrando A, Manunza A, Jordana J, Capote J, Pons A, Pais J, et al. (2015) A
mitochondrial analysis reveals distinct founder effect signatures in Canarian and
Balearic goats. Anim Genet 46: 452-456.

15. Olalde I, Capote J, Del-Arco MC, Atoche P, Delgado T, Gonzalez-Anton R, et al.
(2015) Ancient DNA sheds light on the ancestry of pre-hispanic Canarian pigs.
Genet Sel Evol 47: 40.

16. Guatelli-Steinberg D, Irish JD, Lukacs JR (2001) Canary islands-north African
population affinities: measures of divergence based on dental morphology. Homo
52: 173-188.

17. Bermudez De Castro JM (1989) The Carabelli trait in human prehistoric populations
of the Canary Islands. Hum Biol 61: 117-131.

18. Bermudez de Castro JM (1989) Third molar agenesis in human prehistoric
populations of the Canary Islands. Am J Phys Anthropol 79: 207-215.

19. Billy G (1982) Le peuplement prehistorique de l'Archipel Canarien. El Museo
Canario 41: 59-74.

20. Santana-Cabrera J, Velasco-Vázquez J, Rodríguez-Rodríguez A (2012) Patrón
cotidiano de actividad física y organización social del trabajo en la Gran Canaria
prehispánica (siglos XI-XV): la aportación de los marcadores óseos de actividad
física. Tabona: Revista de Prehistoria y de Arqueología 19: 125-163.

21. Atoche P, Ramirez-Rodriguez MA (2011) Nuevas dataciones radiocarbonicas para la
Protohistoria canaria: el yacimiento de Buenavista (Lanzarote). Anuario de
Estudios Atlanticos 57: 139-170.

22. Arco-Aguilar MC, Arco-Aguilar MM, Benito-Mateo C, Rosario-Adrian MC (2017)
Un taller romano de púrpura en los limites de la Ecumene: Lobos 1. Santa Cruz de
Tenerife: Cabildo de Tenerife.

23. Bronk-Ramsay C (2009) Bayesian analysis of radiocarbon dates. Radiocarbon 51:
337-360.

24. Morales-Mateos J, Rodríguez-Rodríguez A, Herbig A (2017) Agricultura y
recolección vegetal en la arqueología prehispánica de las Islas Canarias (siglos
XI-XV dC): la contribución de los estudios carpológicos. In: Fernández-Eraso J,
Mujika-Alustiza JA, Arrizabalaga-Valbuena A, editors. Miscelánea en homenaje
a Lydia Zapata Peña (1965-2015). Vitoria: Universidad del País Vasco/Euskal
Herriko Unibertsitatea. pp. 189-218.

25. Hernández-Marrero J-C, Navarro-Mederos J-F, Trujillo-Mora J-M, Cancel S,
Machado-Yanes C, Pais J, et al. (2016) An approach to prehistoric shepherding in
La Gomera (Canary Islands) through the study of domestic spaces. Quaternary
International 414: 337-349.

26. Kivisild T (2015) Maternal ancestry and population history from whole mitochondrial
genomes. Investig Genet 6: 3.

27. Rando JC, Cabrera VM, Larruga JM, Hernandez M, Gonzalez AM, Pinto F, et al.
(1999) Phylogeographic patterns of mtDNA reflecting the colonization of the
Canary Islands. Ann Hum Genet 63: 413-428.

28. Bosch E, Calafell F, Comas D, Oefner PJ, Underhill PA, Bertranpetit J (2001) High-
resolution analysis of human Y-chromosome variation shows a sharp
discontinuity and limited gene flow between northwestern Africa and the Iberian
Peninsula. Am J Hum Genet 68: 1019-1029.

29. Rando JC, Pinto F, Gonzalez AM, Hernandez M, Larruga JM, Cabrera VM, et al.
(1998) Mitochondrial DNA analysis of northwest African populations reveals
genetic exchanges with European, near-eastern, and sub-Saharan populations.
Ann Hum Genet 62: 531-550.

30. Secher B, Fregel R, Larruga JM, Cabrera VM, Endicott P, Pestano JJ, et al. (2014)
The history of the North African mitochondrial DNA haplogroup U6 gene flow
into the African, Eurasian and American continents. BMC evolutionary biology
14: 109.

31. Santos C, Fregel R, Cabrera VM, Gonzalez AM, Larruga JM, Lima M (2010)
Mitochondrial DNA patterns in the Macaronesia islands: Variation within and
among archipelagos. Am J Phys Anthropol 141: 610-619.

32. Santana J, Fregel R, Lightfoot E, Morales J, Alamon M, Guillen J, et al. (2016) The
early colonial atlantic world: New insights on the African Diaspora from isotopic
and ancient DNA analyses of a multiethnic 15th-17th century burial population
from the Canary Islands, Spain. Am J Phys Anthropol 159: 300-312.

33. Fregel R (2010) La evolucion genetica de las poblaciones humanas canarias:
determinacion mediante marcadores polimorficos y uniparentales. La Laguna:
Servicio de Publicaciones de la Universidad de La Laguna.

34. Farrujía de la Rosa AJ, Pichler W, Rodrigue A, Garcia-Marin S (2010) The Libyco–
Berber and Latino–Canarian Scripts and the Colonization of the Canary Islands.
African Archaeology Review 27: 13-41.

35. Flores C, Maca-Meyer N, Perez JA, Gonzalez AM, Larruga JM, Cabrera VM (2003)
A predominant European ancestry of paternal lineages from Canary Islanders.
Ann Hum Genet 67: 138-152.

36. Maca-Meyer N, Arnay M, Rando JC, Flores C, Gonzalez AM, Cabrera VM, et al.
(2004) Ancient mtDNA analysis and the origin of the Guanches. Eur J Hum
Genet 12: 155-162.

37. Fregel R, Pestano J, Arnay M, Cabrera VM, Larruga JM, Gonzalez AM (2009) The
maternal aborigine colonization of La Palma (Canary Islands). Eur J Hum Genet
17: 1314-1324.

38. Fregel R, Cabrera VM, Larruga JM, Hernandez JC, Gamez A, Pestano JJ, et al.
(2015) Isolation and prominent aboriginal maternal legacy in the present-day
population of La Gomera (Canary Islands). Eur J Hum Genet 23: 1236-1243.

39. Ordóñez AC, Fregel R, Trujillo-Mederos A, Hervella M, de-la-Rúa C, Arnay-de-la-
Rosa M (2017) Genetic studies on the prehispanic population buried in Punta
Azul cave (El Hierro, Canary Islands). Journal of Archaeological Science 78: 20-
28.

40. O'Rourke DH, Hayes MG, Carlyle SW (2000) Ancient DNA studies in physical
anthropology. Annual Review of Anthropology 29: 217-242.

41. Rodriguez-Varela R, Gunther T, Krzewinska M, Stora J, Gillingwater TH,
MacCallum M, et al. (2017) Genomic Analyses of Pre-European Conquest
Human Remains from the Canary Islands Reveal Close Affinity to Modern North
Africans. Curr Biol 27: 3396-3402 e3395.

42. Carpenter ML, Buenrostro JD, Valdiosera C, Schroeder H, Allentoft ME, Sikora M,
et al. (2013) Pulling out the 1%: Whole-Genome Capture for the Targeted
Enrichment of Ancient DNA Sequencing Libraries. American Journal of Human
Genetics 93: 852-864.

43. Arnay-De-La-Rosa M, González-Reimers E, Yanes Y, Romanek CS, Noakes JE,
Galindo-Martín L (2011) Paleonutritional and paleodietary survey on prehistoric
humans from Las Cañadas del Teide (Tenerife, Canary Islands) based on
chemical and histological analysis of bone. Journal of Archaeological Science 38:
884-895.

44. Espinosa A (1980) Del origen y milagros de la Santa Imagen de Nuestra Señora de
Candelaria que apareció en la Isla de Tenerife con la descripción de esta Isla.
Santa Cruz de Tenerife: Ediciones Goya.

45. Pinhasi R, Fernandes D, Sirak K, Novak M, Connell S, Alpaslan-Roodenberg S, et al.
(2015) Optimal Ancient DNA Yields from the Inner Ear Part of the Human
Petrous Bone. PLoS One 10: e0129102.

46. Dabney J, Knapp M, Glocke I, Gansauge MT, Weihmann A, Nickel B, et al. (2013)
Complete mitochondrial genome sequence of a Middle Pleistocene cave bear
reconstructed from ultrashort DNA fragments. Proc Natl Acad Sci U S A 110:
15758-15763.

47. Meyer M, Kircher M (2010) Illumina sequencing library preparation for highly
multiplexed target capture and sequencing. Cold Spring Harb Protoc 2010: pdb
prot5448.

48. Fregel R, Almeida M, Betancor E, Suarez NM, Pestano J (2011) Reliable nuclear and
mitochondrial DNA quantification for low copy number and degraded forensic
samples. Forensic Science International: Genetics Supplement Series 3: e303-
e304.

49. Gonzalez AM, Cabrera VM, Larruga JM, Tounkara A, Noumsi G, Thomas BN, et al.
(2006) Mitochondrial DNA variation in Mauritania and Mali and their genetic
relationship to other Western Africa populations. Ann Hum Genet 70: 631-657.

50. Lindgreen S (2012) AdapterRemoval: easy cleaning of next-generation sequencing
reads. BMC Res Notes 5: 337.

51. Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25: 1754-1760.

52. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. (2009) The
Sequence Alignment/Map format and SAMtools. Bioinformatics 25: 2078-2079.

53. Ginolhac A, Rasmussen M, Gilbert MT, Willerslev E, Orlando L (2011)
mapDamage: testing for damage patterns in ancient DNA sequences.
Bioinformatics 27: 2153-2155.

54. R_Development_Core_Team (2008) A language and environment for statistical
computing. R Foundation for Statistical Computing. Vienna, Austria. ISBN 3-
900051-07-0, URL http://www.R-project.org.

55. Fu Q, Mittnik A, Johnson PL, Bos K, Lari M, Bollongino R, et al. (2013) A revised
timescale for human evolution based on ancient mitochondrial genomes. Current
biology : CB 23: 553-559.

56. Renaud G, Slon V, Duggan AT, Kelso J (2015) Schmutzi: estimation of
contamination and endogenous mitochondrial consensus calling for ancient DNA.
Genome Biol 16: 224.

57. Andrews RM, Kubacka I, Chinnery PF, Lightowlers RN, Turnbull DM, Howell N
(1999) Reanalysis and revision of the Cambridge reference sequence for human
mitochondrial DNA. Nat Genet 23: 147.

58. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. (2011) A
framework for variation discovery and genotyping using next-generation DNA
sequencing data. Nat Genet 43: 491-498.

59. Kloss-Brandstatter A, Pacher D, Schonherr S, Weissensteiner H, Binna R, Specht G,
et al. (2011) HaploGrep: a fast and reliable algorithm for automatic classification
of mitochondrial DNA haplogroups. Hum Mutat 32: 25-32.

60. van Oven M, Kayser M (2009) Updated comprehensive phylogenetic tree of global
815   human mitochondrial DNA variation. Hum Mutat 30: E386-394.

816   61. Milne I, Stephen G, Bayer M, Cock PJA, Pritchard L, Cardle L, et al. (2013) Using
817   Tablet for visual exploration of second-generation sequencing data. Briefings in
818   Bioinformatics 14: 193-202.

819   62. Bandelt HJ, Forster P, Rohl A (1999) Median-joining networks for inferring
820   intraspecific phylogenies. Mol Biol Evol 16: 37-48.

821   63. Forster P, Harding R, Torroni A, Bandelt HJ (1996) Origin and evolution of Native
822   American mtDNA variation: a reappraisal. Am J Hum Genet 59: 935-945.

823   64. Soares P, Ermini L, Thomson N, Mormina M, Rito T, Rohl A, et al. (2009)
824   Correcting for purifying selection: an improved human mitochondrial molecular
825   clock. Am J Hum Genet 84: 740-759.

826   65. Saillard J, Forster P, Lynnerup N, Bandelt HJ, Norby S (2000) mtDNA variation
827   among Greenland Eskimos: the edge of the Beringian expansion. Am J Hum
828   Genet 67: 718-726.

829   66. Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and
830   analysis program for Windows 95/98/NT. Nucleic acids symposium series 41: 95-
831   98.

832   67. Fregel R, Delgado S (2011) HaploSearch: a tool for haplotype-sequence two-way
833   transformation. Mitochondrion 11: 366-367.

834   68. Nei N (1987) Molecular evolutionary genetics. New York: Columbia University
835   Press.

836   69. Slatkin M (1995) A measure of population subdivision based on microsatellite allele
837   frequencies. Genetics 139: 457-462.

838   70. Excoffier L, Lischer HEL (2010) Arlequin suite ver 3.5: A new series of programs to
839   perform population genetics analyses under Linux and Windows. Mol Ecol
840   Resour 10: 564-567.

841   71. de Leeuw J, Mair P (2009) Multidimensional Scaling Using Majorization: SMACOF
842   in R. Journal of Statistical Software 31: 1-30.

843   72. Long JC (1991) The genetic structure of admixed populations. Genetics 127: 417-
844   428.

73. Fregel R, Mendez FL, Bokbot Y, Martin-Socas D, Camalich-Massieu MD, Santana J,

et al. (2018) Ancient genomes from North Africa evidence prehistoric migrations

to the Maghreb from both the Levant and Europe. Proc Natl Acad Sci U S A in

press.

74. Heinz T, Cardenas JM, Alvarez-Iglesias V, Pardo-Seco J, Gomez-Carballa A, Santos

C, et al. (2015) The Genomic Legacy of the Transatlantic Slave Trade in the

Yungas Valley of Bolivia. PLoS One 10: e0134129.

75. Bandelt HJ, Herrnstadt C, Yao YG, Kong QP, Kivisild T, Rengo C, et al. (2003)

Identification of Native American Founder mtDNAs Through the Analysis of

Complete mtDNA Sequences: Some Caveats. Annals of Human Genetics 67:

512--524.

76. Cabrera-Pérez J, Perera-Betancort MA, Tejera-Gaspar A (1999) Majos La primitiva

población de Lanzarote. Islas Canarias. Teguise: Fundación César Manrique.

77. Macias-Hernandez AM (1988) Fuentes y principales problemas metodológicos de la

demografía histórica de Canarias. Anuario de Estudios Atlánticos 34: 51-157.

78. Navarro JF (1992) La Prehistoria de Canarias. Vol. 5. La Gomera y Los Gomeros.

Santa Cruz de Tenerife: Centro de la Cultura Popular Canaria.

79. Ruiz-Gonzalez MT (2008) El territorio de los antiguos bimbaches / Algunas

cuestiones en torno al territorio de Esero. VI Congreso de Patrimonio Histórico:

'Arqueología en Canarias, Territorio y Sociedad'. Lanzarote: Cabildo Insular de

Lanzarote.

80. Jimenez JJ (1992) La Prehistoria de Canarias. Vol. 6. El Hierro y los Bimbaches.

Santa Cruz de Tenerife: Centro de la Cultura Popular Canaria.

81. Pico B, Aznar E, Corbella D (2003) Le Canarien. Manuscritos, trancripción y

traducción, La Laguna. Santa Cruz de Tenerife: Instituto de Estudios Canarios.

82. van de Loosdrecht M, Bouzouggar A, Humphrey L, Posth C, Barton N, Aximu-Petri

A, et al. (2018) Pleistocene North African genomes link Near Eastern and sub-

Saharan African human populations. Science.

83. Souville G (1998) Contacts et échanges entre la péninsule Ibérique et le Nord-Ouest

de l'Afrique durant les temps préhistoriques et protohistoriques. Comptes rendus

des séances de l'Académie des Inscriptions et Belles-Lettres. pp. 163-177.

84. Camps G (1974) La civilisations préhistoriques de l'Afrique du Nord et du Sahara.
Paris: Doin.

85. Clark JD (1997) The Cambridge History of Africa,1. From the earliest times to c.
500BC. Cambridge: Cambridge University Press.

86. Aubet ME (1994) Tiro y las colonias fenicias de Occidente. Barcelona: Crítica.

87. Lopez-Pardo F (1993) La colonización fenicio-púnica en el África noroccidental.
Hispania Antiqua 17: 435-449.

88. Le Bohec Y (2005) Histoire de l'Afrique romaine (146 avant J.C. - 439 après J. C.).
Paris: Picard.

89. Camps G (1980) Berberes: aux marges de l´histoire. Paris: Hesperides.

90. Tejera-Gaspar A (1989) Parentesco, familia y matrimonio en las culturas
prehistóricas de las Islas Canarias. In: Bermejo JC, editor. Parentesco, familia y
matrimonio en la historia de Galicia. Santiago de Compostela: Torculo Ediciones.

91. Atoche P, Paz JA, Ramirez MA, Ortiz ME (1995) Evidencias arqueologicas del
mundo romano en Lanzarote. Arrecife: Servicio de publicaciones del Excmo.
Cabildo Insular de Lanzarote.

92. Delgado-Delgado JA (2001) Las islas de Juno: ¿hitos de la navegación fenicia en el
Atlántico en época arcaica? The Ancient History Bulletin 15: 29-43.

93. Delgado-Delgado JA (2012) Canarias en la antigüedad como problema histórico.
Tabona: Revista de Prehistoria y de Arqueología 19: 9-13.

94. Raven S (2003) Rome in Africa. London: Routledge.

95. Gonzalez-Anton R, del Arco-Aguilar MC, Berhmann RB, Bueno-Ramírez P (1998)
El poblamiento de un archipiélago Atlántico: Canarias en el proceso colonizador
del primer milenio a. C. Eres 8: 43-100.

96. González-Antón R, Del Arco-Aguilar MC (2009) Navegaciones exploratorias en
Canarias a finales del II milenio AC e inicios del primero: El cordón litoral de La
Graciosa (Lanzarote). Canarias Arqueológica: Arqueología-Bioantropología 17:
9-80.

97. Guerrero-Ayuso VM (2008) Canarias en el contexto de las navegaciones atlánticas de
la Antigüedad. SAGVNTVM Papeles del Laboratorio de Arqueología 40: 37-62.

98. Matisoo-Smith EA, Gosling AL, Boocock J, Kardailsky O, Kurumilian Y, Roudesli-
Chebbi S, et al. (2016) A European Mitochondrial Haplotype Identified in
Ancient Phoenician Remains from Carthage, North Africa. PLoS One 11:
e0155046.

99. Matisoo-Smith E, Gosling AL, Platt D, Kardailsky O, Prost S, Cameron-Christie S, et
al. (2018) Ancient mitogenomes of Phoenicians from Sardinia and Lebanon: A
story of settlement, integration, and female mobility. PLoS One 13: e0190169.

100. Delgado-Darias T, Alberto-Barroso V, Velasco-Vazquez J (2018) Violence in
paradise: Cranial trauma in the prehispanic population of Gran Canaria (Canary
Islands). Am J Phys Anthropol.

101. Delgado T (2009) La Historia en los dientes. Una aproximación a la Prehistoria de
Gran Canaria desde la Antropología Dental. Las Palmas: Cabildo Insular de Gran
Canaria.

102. Rodriguez-Rodriguez A, Morales-Mateos J, Pino-Curbelo MD, Naranjo-Mayor Y,
Martín-Rodriguez E, González-Marrero MC (2012) Espacios de producción
especializada, excedentes y estratificación social en la Gran Canaria pre-europea.
Tabona: Revista de Prehistoria y de Arqueología 19: 101-123.

103. Morales J, Rodriguez-Rodriguez A, Marrero A (2013) Prehistoric Plant Use on La
Palma Island (Canary Islands, Spain): an example of disappearance of agriculture
in an isolated environment. In: Stevens GJ, Nixon S, Murray MA, Fuller D,
editors. Archaeology of African Plant Use. California: Institute of Archaeology
Publications, UCL. pp. 195-204.

104. Moreno-Benítez M (2014) Tiempo y espacio: La cronología de Gran Canaria y el
territorio. Boletín electrónico de Patrimonio Histórico 2.

105. del Pino-Curbelo M, Rodriguez-Rodriguez A, Buxeda-Garrigos J, Mangas-Viñuela
J, González-Quintero P, Moreno-Benítez MA (2016) Las cerámicas aborígenes de
Gran Canaria (Islas Canarias) a través del yacimiento de La Cerera: materias
primas, tecnología y función. Trabajos de Prehistoria 73: 90-114.

106. Santana-Cabrera J, Moreno-Benítez M, Suarez-Medina I, Mendoza-Medina F,
Alberto-Barroso V (2018) Zonzamas: un yacimiento singular en la isla de
Lanzarote. Nuevos datos arqueológicos. Actas del XXII Coloquio de Historia

Canario Americana. Las Palmas de Gran Canaria: Cabildo de Gran Canaria. pp.

1-14.

107. Cabrera-Pérez JC (1996) La Prehistoria de Fuerteventura. Un modelo insular de

adaptación. Las Palmas de Gran Canaria: Cabildo Insular de Gran Canaria y

Fuerteventura.

TABLES:

Table 1 – MtDNA haplogroup absolute frequencies for the indigenous population of

the Canary Islands. Haplogroup frequencies and diversity were calculated using HVRI

sequence data from this study and previously published data. 1: This study; 2: Ordoñez et

al. 2017; 3: Fregel et al. 2009; 4: Maca-Meyer et al. 2004; 5: Fregel et al. 2014; 6:

Rodríguez-Varela et al. 2017.

Haplogroup	El Hierro[1,2]	La Palma[1,3]	Tenerife[1,4,6]	La Gomera[1,5]	Gran Canaria[1,6]	Lanzarote & Fuerteventura[1]	Total
H*	12	10	15	2	33	13	85
H1cf	57	8	6	2	1	-	74
H4a1e	-	-	-	-	3	1	4
HV0	-	-	1	-	-	-	1
J	-	3	4	5	2	-	14
K	-	1	1	1	2	-	5
L1/L2	-	2	2	1	1	-	6
L3	-	-	2	4	1	-	7
L3b1a12	-	-	-	-	5	-	5
HV0	-	-	-	-	1	-	1
Other T	-	1	-	-	3	-	4
T2c1	-	3	12	-	15	2	32
U5	-	-	-	-	3	1	4
U6a	-	-	2	-	6	2	10
U6b	-	2	8	38	4	-	52
U6c	-	-	-	-	5	1	6
U7	1	-	-	-	-	-	1
W1e1	-	1	-	-	-	-	1
X3a	-	4	-	4	1	-	9
Sample size	70	35	53	57	86	20	321
Haplogroup diversity	2.86% ± 2.76%	72.10% ± 7.63%	77.43% ± 4.02%	54.20% ± 7.50%	77.10% ± 3.78%	51.05% ± 12.84%	69.69% ± 2.37%

Table 2 – Admixture results based on mtDNA haplogroup frequencies. Admixture results for the modern population of the Canary Islands using the three main parental populations: Iberian Peninsula (IBP), sub-Saharan Africa (SSA) and the Canarian indigenous population (CIP). Admixture calculations were performed using two approximations: A) we used the whole ancient dataset (combining the ancient samples from all the seven islands) as CIP for calculating admixture estimates for all islands; B) we used each indigenous sample to calculate the admixture of its respective island (e.g. to calculate admixture in the modern population of Gran Canaria we exclusively used the ancient samples from Gran Canaria as CIP). Results are shown for: the whole Canary Islands population (CAN) and the seven individual islands (FUE=Fuerteventura; GCA=Gran Canaria; GOM=La Gomera; HIE=El Hierro; LAN=Lanzarote; PAL=La Palma; TFE=Tenerife).

A: Whole indigenous sample									
Component	IBP			SSA			CIP		
FUE	0.4108	±	0.0071	0.0199	±	0.0015	0.5692	±	0.0070
GCA	0.6486	±	0.0039	0.0438	±	0.0012	0.3076	±	0.0038
GOM	0.2170	±	0.0181	0.0691	±	0.0075	0.7139	±	0.0186
HIE	0.6379	±	0.0122	0.0000	±	0.0000	0.3621	±	0.0122
LAN	0.3303	±	0.0084	0.0448	±	0.0026	0.6248	±	0.0084
PAL	0.5599	±	0.0106	0.0000	±	0.0001	0.4401	±	0.0107
TFE	0.5989	±	0.0041	0.0452	±	0.0012	0.3559	±	0.0040
CAN	0.3978	±	0.0100	0.0432	±	0.0029	0.5589	±	0.0100
B: Indigenous sample from each island									
Component	IBP			SSA			CIP		
FUE	0.6458	±	0.0095	0.0203	±	0.0020	0.3339	±	0.0093
GCA	0.6944	±	0.0040	0.0593	±	0.0015	0.2463	±	0.0038
GOM	0.3768	±	0.0049	0.0682	±	0.0024	0.5550	±	0.0049
HIE	1.0000	±	0.0087	0.0000	±	0.0000	0.0000	±	0.0087
LAN	0.7202	±	0.0116	0.0239	±	0.0027	0.2559	±	0.0113
PAL	0.5896	±	0.0134	0.0000	±	0.0116	0.4104	±	0.0118
TFE	0.7306	±	0.0030	0.0495	±	0.0012	0.2199	±	0.0029

 TITLES AND LEGENDS TO FIGURES:

Figure 1 – Map of the Canary Islands showing the geographical location of the

archaeological sites included in this study. Codes are as follows: 1 – Cueva del Agua; 2

– Huerto de los Morales; 3 – Salto del Casimiro; 4 – El Espigón; 5 – Los Pasitos; 6 –

Punta Azul; 7 – Barranco de Majona; 8 – El Pescante; 9 – Antoncojo; 10 – Las Arenas;

11 – El Cedro; 12 – El Salitre; 13 – El Portillo; 14 – La Angostura; 15 – El Cascajo; 16 –

El Capricho; 17 – El Agujero; 18 – El Hormiguero; 19 – Guayadeque; 20 – La Fortaleza;

21 – Cuermeja; 22 – Lomo Galeón; 23 – Puente de la Calzada; 24 – El Huriamen; 25 –

Montaña Mina.

Figure 2 – DNA authentication results for all the samples included in this study. A)

Insert size density plot. B) Contamination rates estimated using contamMix and

schmutzi. C) Damage patterns.

Figure 3 – MtDNA haplogroup frequencies for ancient and current populations of

the Canary Islands.

Figure 4 – MDS plot based on haplogroup frequency distances. A) MDS analysis

comparing the individual ancient populations (FUI=Fuerteventura; GCI=Gran Canaria;

GOI=La Gomera; HII=El Hierro; LAI=Lanzarote; PAI=La Palma; TFI=Tenerife), with

modern Canarian (codes as in Table 2), Caucasus (CAU), North African (codes as in

Table S1), Sub-Saharan African (SSA), European (codes as in Table S1) and Near

Eastern populations (codes as in Table S2). B) MDS analysis as in Figure 4A, but

removing outliers (HIE, HII and GOM) and pooling all the remaining indigenous samples

together (CIP).

TITLES AND LEGENDS TO SUPPLEMENTARY MATERIAL:

Table S1 –Populations used for comparisons in this study.

Table S2 – Summary of mtDNA results for all aDNA samples.

Table S3 – HVRI data used for this study, including new results on the islands of Gran Canaria, Lanzarote and Fuerteventura.

Figure S1 – Combined calibrated radiocarbon per archaeological site (A) and per mtDNA lineage (B).

Figure S2 – Phylogenetic tree of complete ancient Canarian mtDNA sequences. Number along links refers to nucleotide changes, whereas "@", "d" and "i" indicates back mutations, deletions and insertions, respectively. Recurrent mutations, such as 309iC, 315iC and 16519, have not been taken into account.

Figure S3 – Phylogenetic tree of complete modern Canary Islands sequences. The most probable geographic origin of the sequences is indicated.

Figure S4 – Phylogenetic tree of complete haplogroup H1cf sequences. GenBank accessions and geographic origin are indicated for each complete sequence taken from the bibliography.

Figure S5 – Coalescence ages for mtDNA haplogroups observed in the indigenous population of the Canary Islands. All the coalescence ages have been calculated in this study, except for H2a, H3 and T2b, whose ages have been obtained from previous results (Behar et al. 2008).

Figure S6 – Phylogenetic tree of complete haplogroup H1e1a sequences. GenBank accessions and geographic origin are indicated for each complete sequence taken from the bibliography. Sub-haplogroups in dark grey and white fonts indicate newly defined branches.

Figure S7 – Phylogenetic tree of complete haplogroup H4a1 sequences. GenBank accessions and geographic origin are indicated for each complete sequence taken from the bibliography. Sub-haplogroups in dark grey and white fonts indicate newly defined branches.

Figure S8 – Phylogenetic tree of complete haplogroup J1c3 sequences. GenBank accessions and geographic origin are indicated for each complete sequence taken from the bibliography.

Figure S9 – Phylogenetic tree of complete haplogroup J2a2d sequences. GenBank

accessions and geographic origin are indicated for each complete sequence taken from

the bibliography. Sub-haplogroups in dark grey and white fonts indicate newly defined

branches.

Figure S10 – Phylogenetic tree of complete haplogroup T2c1d sequences. GenBank

accessions and geographic origin are indicated for each complete sequence taken from

the bibliography. Sub-haplogroups in dark grey and white fonts indicate newly defined

branches.

Figure S11 – Phylogenetic tree of complete haplogroup L3b1a sequences. GenBank

accessions and geographic origin are indicated for each complete sequence taken from

the bibliography. Sub-haplogroups in dark grey and white fonts indicate newly defined

branches.

Figure S12 – Phylogenetic tree of complete haplogroup X3a sequences. GenBank

accessions and geographic origin are indicated for each complete sequence taken from

the bibliography.

Figure S13 – Phylogenetic tree of complete haplogroup U6a1a1 sequences. GenBank

accessions and geographic origin are indicated for each complete sequence taken from

the bibliography.

Figure S14 – Phylogenetic tree of complete haplogroup U6a7a1 sequences. GenBank

accessions and geographic origin are indicated for each complete sequence taken from

the bibliography.

Figure S15 – Phylogenetic tree of complete haplogroup U6b1a sequences. GenBank

accessions and geographic origin are indicated for each complete sequence taken from

the bibliography.

Figure S16 – Phylogenetic tree of complete haplogroup U6c sequences. GenBank

accessions and geographic origin are indicated for each complete sequence taken from

the bibliography.

1050